Review 1

Abstract Please use abbreviations in case they are introduced! Page 1 / Line 12: the-polar stratospheric clouds -> PSCs Corrected Page 1 / Line 14: Polar stratospheric ice clouds -> ice PSCs Corrected

Page 1 / Line 17: decrease of the water vapor

Corrected

1 Introduction

Page 2 / Line 51: Please add a citation for the increase in radiative cooling Citation of Shindell et al., (1998) have added. Page 2 / Line 52: Rex et al. (2006) found an increase in the PSC volume trend Done Page 2 / Line 54: (see also Dameris et al., 2014) Done Page 2 / Line 54: Please reformulate the sentence. For example: However, the relationship between increasing greenhouse gas concentrations and increasing PSC occurrences remains a controversial issue. Thanks for reformulation, we corrected the text. Page 2 / Line 57: Overall The result of a climate model study ... Done Page 2 / Line 58: would lead to an increase in the total ozone column ... Done Please check carefully the use of articles in the entire manuscript! I will not comment on this further. Tried to do Page 4 / Line 108: Solomon et al. (2010) Done Page 4 / Line 109: Hegglin et al. (2014) Done Page 4 / Line 112: Recently, Urban et al. (2014) reported ... Done

2.1.1 Water vapour and PSC measurements

The numbering of the subsections were wrong, 2.1.1 were changed to 2.2 You introduced already satellite names, PSCs etc. Extra introductions removed

3 Water vapour distribution

Page 9 / Line 1: MLS observations calculated over winters 2004 – 2014. In Figure 2, the period 2005 – 2014 is mentioned. Please check!

MLS water vapour data is available from August 2004. Winter 2004 there were no MLS data. Year corrected to the text.

4 The origin and long term variability of water vapour

Page 10 / Line 327: values for the years period 1990 – 2014 Done

Page 10 / Line 329: positive trends

Done

Caption Figure 5: (b-d) Same as panel (a), but levels 10, 56 and <u>100</u> hPa. -> Sentence not necessary anymore since you replaced the panels labelling by levels.

Corrected anyhow

page 10 / Line 334: "At 10 hPa the increase stopped by 2014 but at 1 hPa it did not stop until the end of the time series." -> I am confused because your time series ends 2014!?

The text is corrected and is now:"At 10 hPa the increase stopped

by the end of 2013 but at 1 hPa it did not stop until the end of the time series"

Page 10 / Line 336: lower and upper stratosphere, which suggests that ...

Done

Page 10 / Line 343: QBO index (QBO, equatorial winds at 50 hPa) Done

Table 2: The meaning and units of the numbers are missing. I guess the first line are water vapour values in ppm. I don't understand the meaning of the second and third line.

We apology for missing these explanations from the table. The description of the table is improved in the revised version. And the new caption is:

"Table 2. Monthly mean ice PSC areas (10⁶ km²) in FinROSE (left) and CALIPSO (right) at XX pressure level as a function of 56 hPa monthly mean water vapour concentration (ppm) averaged north of 60N and the area of temperatures below 188K (10⁶ km²). The number of months in each group are shown in parentheses."

6 Case study: Winter 2009/2010

Page 13 / Line 425: Temperatures were / The temperature was Done

Page 13 / Line 426: In the lower stratosphere temperatures dropped below 195 K Done

Page 14 / Line 453 – 458 and Figure 9: On 17 Jan, areas with low water vapor coincide with areas where temperatures are coldest. The water vapor is frozen and with proceeding time, ice particles grow and sediment. The sedimentation of ice particles results in a redistribution of water vapour. 20 - 23 Jan: You can nicely see in your FinROSE simulation how the dehydrated air masses (no ice present) move around the pole. Here, minimum water vapour values do not coincide with coldest temperatures!

Thansk for the comment. We addes following to the text:

" Areas with very cold temperatures correlates with very low water vapour content areas because of the ice PSC formation. On 17th January the driest areas coincide both in space and time with the coldest temperatures. The water vapour is frozen and condensed into ice particles and with proceeding time, ice particles grow and sediment downwards resulting in dehydrated air masses. The sedimentation of ice particles results in a redistribution of water vapour. On 20th-23rd January the dehydrated air masses, where no ice presents move around the pole, but the minimum water vapour values do not coincide with coldest temperatures."

Figure 9, CALIPSO – FinROSE comparison: I am still not convinced if this comparison is fair. CALIPSO sees massive ice PSCs on e.g. 17 Jan and your plot looks like only sparse PSC coverage has been observed by CALIPSO. This is of course due to the fact that CALIPSO measures only single orbits. However, you may find a way to improve this comparison.

We gridded the CALIPSO data again to larger grid boxes and drowe the figure again. Now the CALIPSO area is larger than the area in FinROSE simulation. The text changed as follow:

"To facilitate the comparison, CALIPSO observations have been gridded to the grid of 5 x 10 and

the grid box is marked as having ice PSC if there were at least one ice PSC observation within the grid box area.Note that each day CALIPSO orbits cover only a part of the globe; therefore observations appear sparse. Nevertheless it can be seen that the grid boxes where CALIPSO observed ice PSC are scattered over larger areas in comparison to FinROSE simulations." Page 14 / Line 476 + Page 15 / Line 500 and Line 504: ICE PSC -> ice PSC Done

7 Conclusions

Page 14 / Line 481: "The full chemistry in FinROSE can add the water vapour to the ECMWF ERA-Interim water vapour." -> I don't understand the meaning of this sentence. Corrected text:

"The full chemistry in FinROSE produces more water than simplified parameterisation of ECMWF ERA-Interim, alleviates the dry bias of ERA-Interim and improves the comparison with observations."

Page 15 / Line 500: "10 out of 25 simulated winters" -> I thought that 12 out of 25 winters show ice PSCs (see page 12, line 386) Corrected to be 12 Page 15, Line 511: reported by Khaykin et al. (2013) Done

Acknowledgements Page 15 / Line 517: Sodankylä Corrected

I would highly recommend proofreading the whole article by a native speaker before final publication.

The language have been checked

Review 2

Comments on the revised version of the ACPD manuscript entitled "Variability of water vapour in the Arctic stratosphere" by Laura Thölix, Leif Backman, Rigel Kivi and Alexey Karpechko.

The revised is clearly improved compared to the submitted version. The main problems I have concern the regression analysis, I think there is room for clarifications and improvements. Please find my comments below:

Comments:

▶ page 2, line 39: "... the abundance of other PSCs ..." may be changed to "... the abundance of other PSC types ..." to make a distinction to ice PSCs more directly.

Done

▶ page 2 and 3, line 58 to 60: The are some problems with that sentence here. "caused" should read "causing a" and "... and to an increase in ..." may read "... and an increase of ...". Done

• page 4, lines 99 to 107: It is probably fair to separate SMR and MIPAS from the other instruments that employ/employed the occultation technique, which limits the temporal and spatial coverage in general. SCIAMACHY is kind of an hybrid that provides some limited coverage of the Arctic in its solar occultation mode. Its limb mode however is in the same class as MLS, MIPAS and SMR providing global coverage on a daily basis.

Done

▶ page 5, line 138 and 139: Please add "lower" before "stratosphere" in the first line. In the upper stratosphere the trends reported by Hegglin et al. (2014) are positive. In the second the phrase "not good" is probably a bit to strong. "not optimal" might be a better choice. Done

▶ page 6, line 170: Please remove the brackets around the Hanson and Mauersberger reference. Done

▶ page 7, line 211: There is something strange with "First larger set …" sentence.

The text has changed as follow: "First set of sounding observations large enough to derive some statistics of water vapour vertical distribution was obtained during the..."

▶ page 7, lines 225 to 227: Why are the CFH measurements from the first LAPBIAT not used, only the NOAA FPH?

There was a mistake in the previous version. We have included 9 NOAA soundings and 4 CFH soundings. In total we have 13 soundings from the first campaign. We have corrected the text. Now it reads

"In this study we have used in total 13 NOAA and CFH frostpoint hygrometer profiles obtained during the first campaign and 13 CFH ..."

▶ page 8, line 268: "... is about 0.3 hPa moister ...". I guess ppmv is meant here. Changed to ppm

▶ page 8, lines 268 and 269: The sentence "Too moist air …" does not really fit here as the previous sentence focused on the upper part of the stratosphere where no PSCs exist. 1 of 3 Sentence taken off

▶ page 8, line 269 to 273: The information that ERA-interim shows lower water vapour volume mixing ratios compared to MLS in the upper stratosphere is doubled. Please remember that ERA-interim is setup to relax to 6 ppmv at the stratopause, which largely explains the discrepancies. The doubled information is taken off, and the text has modified as:

"...which is likely due to a relaxation of water vapour towards an equivalent value applied in ERA-Interim at the stratopause (Dethof et al., 2002)."

▶ page 10, line 321: I think the word "could" is not appropriate here. At least I assumed you did it that way.

Taken off

▶ page 10, line 328 to 330: I am not sure if that sentence is really needed. Information in that direction has been conveyed earlier on one side and on the other side it seems obvious. Taken off

▶ page 11, line 365 to 367: I wonder if it is a better choice to use the reference period from 2004 to 2014 even for the de-seasonalisation of the simulations. There is a clear offset between MLS and the simulations especially lower down. This may disappear and result result in nicer comparisons. The temporal variations seems to be consistent between the individual data sets, though.

The reviewer is right that the offset between MSL and FinROSE depends on the choice of the reference period. Choosing 2004-2014 period as a reference would reduce the positive FinROSE offset around 2010-2012 but it would create the negative FinROSE offset around 2004-2008 which is missing in the current version. After playing with different reference periods we have decided not to make any changes in the manuscript. We believe that our choice is correct because it highlights the divergence of the timeseries after 2008, which is not a result of the offset but the result of different trends during this period.

▶ page 12, line 385: This fits also your artificial tracer results, which attribute most of the water vapour changes to dynamical processes.

Thanks for comment. We changed the sentence and add the comment to the text:

"Since the air in the upper polar stratosphere is older than that in the lower stratosphere (Stiller et al. 2012) the delay between water vapour changes in the lower and upper stratosphere suggests that these changes are driven by transport processes, consistent with our tracer results, which attribute most of the water vapour changes to dynamical processes."

• page 12, line 397 to 399: To this date it is not hundred percent clear what this 500 hPa tropical temperature term really considers. Dessler et al. (2013) used it mainly to derive their radiative feedback. There are are certainly some contributions of ENSO and climate change in this terms, but it is not as obvious like for the QBO or a solar proxy. The cold point temperature is likely more covered by the BDC proxy than the 500 hPa temperature overall. Also, the Dessler approach is optimised for the tropics. So, there seems to be some plausible explanations for your problems. Thank you for the comment.

page 12, line 409 to 418: Do the correlation coefficients consider the time lags that have been mentioned earlier? Then I miss the regression coefficients. I would prefer those more than correlations for the discussion of the importance of the individual processes contributing to the water vapour distribution. It would be also interesting to see the regression fits and/or the residuals to identify missing processes.

Thank you for this comment. Yes, the correlation coefficients are at the lags mentioned earlier. Following reviewer's suggestion we also add the regression coefficients to the table. The text is modified as follows:

'The individual correlation and regression coefficients with our proxies taken at the lags mentioned earlier are shown in the Table 1.'

Please note that comparison of correlation coefficients often gives better information about the importance of individual processes then regression coefficients. The latter depend on the units used and it is not always straightforward to compare values if different units are used.

We add figure showing the contribution of individual processes to water vapour variability and add the discussion of the figure as follows:

"The multiple regression coefficients are 0.57 and 0.51 at 82 hPa and at 56 hPa correspondingly, showing that our models only explain 25–30 % of the variability. This is considerably less than that of Dessler et al. (2014) suggesting that different processes contribute to the polar water vapour variability in comparison to those in the tropics. The modelled timeseries together with contributions of individual processes and residuals are shown in Figure 6. It is seen that the cold

point variability and QBO make comparable contributions to the water vapour variability at both levels. Note that the regression somewhat explains the increase of the water vapour from 2005 to 2010, which is more clear at 82 hPa. However the peak of the water 365 vapor during 2011–2013 is not explained by these proxies.'

▶ page 13, line 424: I think "Fig. 1" should be "Fig. 3".

Yes, right, corrected.

▶ page 14, line 472: "Figure 7 shows …" should reference to the right panel of Fig. 7. Corrected

▶ page 14, line 484: You mean the 95% significance level here? 2 of 3

Here we refer to the significance level used to reject the null hypothesis. Sorry for confusing the p-value (0.05) with the significance level (5%), which indicate the probability of wrong rejection of the null hypothesis. We rewrite the text as:

'..., both statistically significant at the 5% level.'

▶ page 15, line 495 to 500: I think this part needs some rewriting as it is raising more questions than it answers. Of course if there is a major warming then the vortex is not destroyed directly. For one cells can remain and also the definition of the warming central date depends on the altitude and place you look at. Also the vortex reformation after the warming comes into play here. Maybe you can spare many parts of this and simply say you focus on the coldest week while after that temperatures were too high.

We agree with the reviewer that an SSW does not necessarily mean that temperatures must rise everywhere. The fact we referred to is simply that after the SSW in January 2010 there were no temperatures cold enough to support PSC formation. To avoid possible confusion we have rewritten this part as follows:

'Cold conditions favouring PSC formations lasted until the end of January 2010. A major sudden stratospheric warming (SSW) took place in the end of January with the central date around 24 January (e.g., Khaykin et al., 2013; Dörnbrack et al., 2012; Pitts et al., 2011). Following the SSW stratospheric temperatures rose above the PSC formation threshold so that no more PSC were observed that winter.'

▶ page 15, line 508 to 511: Is Fig. 8 (the old Fig.7) now considering the whole Arctic instead of Sodankylä?

The Sodankylä temperature have now changed to the area of temperatures colder than 188K. Al the panels are now areas calculated over the whole Arctic.

▶ page 24, table 1: The caption sounds strange.

Caption corrected.

▶ page 24, table 2: You may add that the unit is 1e6 km^2.

Unitst have inserted.

▶ page 32, figure 9: There is a blank missing between "occurrence" and "from" in the second line of the caption.

Corrected

References:

Dessler, A., et al., "Stratospheric water vapor feedback", Proceedings of the National Academy of Science, 110, 18087 - 18091, doi:10.1073/pnas.1310344110, 2013.
Hegglin, M. I., et al., "Vertical structure of stratospheric water vapour trends derived from merged satellite data", Nature Geoscience, 7, 768 - 776, doi:10.1038/ngeo2236, 2014.

Manuscript prepared for Atmos. Chem. Phys. with version 2014/07/29 7.12 Copernicus papers of the $\[Mathebase]$ Class copernicus.cls. Date: 10 March 2016

Variability of water vapour in the Arctic stratosphere

L. Thölix¹, L. Backman¹, R. Kivi², and A. Karpechko³

¹Climate Research, Finnish Meteorological Institute, Helsinki, Finland ²Arctic Research, Finnish Meteorological Institute, Sodankylä, Finland ³Arctic Research, Finnish Meteorological Institute, Helsinki, Finland

Correspondence to: L. Thölix (laura.tholix@fmi.fi)

Abstract.

This study evaluates the stratospheric water vapour distribution and variability in the Arctic. A Fin-ROSE chemistry transport model simulation covering years 1990–2014 is compared to observations (satellite and frostpoint hygrometer soundings) and the sources of stratospheric water vapour are

- 5 studied. In the simulations, the Arctic water vapour shows decadal variability with a magnitude of 0.8 ppm. Both observations and the simulations show an increase in the water vapour concentration in the Arctic stratosphere after year 2006, but around 2012 the concentration started to decrease. Model calculations suggest that this increase in water vapour is mostly explained by transport related processes, while the photochemically produced water vapour plays a relatively smaller role.
- 10 The increase of water vapour in the presence of the low winter temperatures in the Arctic stratosphere led to more frequent occurrence of ice polar stratospheric clouds (PSCs) in the Arctic vortex. The We perform a case study of ice PSC formation focusing on January 2010 when the polar vortex was unusually cold in early 2010 and allowed large scale formation of the polar stratospheric clouds. The cold pool in the stratosphere over the Northern polar latitudes was large and stable and
- 15 <u>PSCs. At the same time</u> a large scale persistent dehydration was observed. Polar stratospheric ice clouds Ice PSCs and dehydration observed at Sodankylä with accurate water vapour soundings in January and February 2010 during the LAPBIAT (Lapland Atmosphere-Biosphere facility) atmospheric measurement campaign were well reproduced by the model. In particular, both the observed and simulated decrease of the water vapour in the dehydration layer was up to 1.5 ppm.

20 1 Introduction

Water vapour is the most important natural greenhouse gas in the atmosphere accounting for about half of the current greenhouse effect (Schmidt et al., 2010). Although the majority of water vapour

resides in the troposphere, it has been highlighted that stratospheric water vapour variations may play an important role in the decadal scale variability of the climate (Solomon et al., 2010). Recently the

- 25 existence of a positive stratospheric water vapour feedback was shown based on observations, i.e. stratospheric water vapour increases with tropospheric temperature, which contributes to the climate sensitivity (e.g., Dessler et al., 2013). Therefore, investigating the changes in stratospheric water vapour abundance is helpful in veluable for the detection and attribution of the ongoing climate change.
- 30 Water vapour is also an important constituent in the stratospheric chemistry. It intensifies ozone destruction both by producing odd-hydrogen species, which can destroy odd-oxygen, and by formation of polar stratospheric clouds (PSC), which enable efficient conversion of halogen reservoir species to halogen radicals (e.g., Solomon et al., 1986). In the winter polar vortex water vapour can condense to form ice PSCs, i.e. type II PSCs. Increased water vapour may also affect the abun-
- 35 dance of other PSCs PSC types as the formation of both NAT (nitric acid trihydrate, type Ia PSC) (Hanson and Mauersberger, 1988) and STS (super cooled ternary solution, type Ib PSC) (Carslaw, 1995) are dependent on nitric acid and water vapour concentrations. Furthermore, the reaction rates on/in STS depend on the composition of the particles, which is a function of water vapour (Sander et al., 2011, Section 5).
- 40 Ice PSCs contribute only to a minor part of the chlorine activation. According to a model study by Kirner et al. (2015) 90 % of the ozone depletion in the Antarctic spring is caused by halogen activation on liquid particles. In a model study of the Arctic winter 2009/2010 Wohltmann et al. (2013) showed that chlorine activation on liquid aerosols alone explained the observed changes in the ozone column to within 10 %. The additional chlorine activation caused by ice PSCs is mod-
- 45 est. However, when ice PSC particles sediment to lower altitudes, a reduction of water vapour, i.e. dehydration, occurs (Kelly et al., 1989). Sedimenting ice PSC particles also contribute to the denitrification (Hintsa et al., 1998), which can prolong the ozone depletion in the spring due to a slower conversion of active chlorine back to the reservoir species ClONO₂.
- The Arctic polar vortex is often less stable and maintains higher temperatures than its Antarctic counterpart, and thus ice PSCs and dehydration are seldom observed (Solomon, 1999). However, stratospheric water vapour is expected to increase due to elimate change caused warming of the tropical tropopause (Gettelman et al., 2009) and to an increase in an increase of the atmospheric methane concentration, both consequences of climate change. This, in addition to the increased radiative cooling in the stratosphere due to the increase of CO₂ and water vapour (Shindell et al., 1998), might
- 55 lead to enhanced PSC formation. For example Rex et al. (2006) found increased an increase in PSC volume trend in the lower Arctic stratosphere during dynamically quiescent winters since 1960s (see also Dameris et al. (2014))(see also Dameris et al., 2014). However the potential of greenhouse gas increases to increase PSC occurrence relationship between increasing greenhouse gas concentrations

and increasing PSC occurrences remains controversial issue (Rieder et al., 2013; Langematz et al., 2014).

The combined effect of temperature and water vapour on ozone, through homogeneous and heterogeneous chemistry and dynamics, is complex. Overall the The result of climate model study by Tian et al. (2009) indicated that the increased stratospheric water vapour would lead to increase an increase in the total column of ozone in the Arctic in the future despite increased active chlorine in the polar spring regions, while in the Antarctic the ozone recovery would be delayed. The evolution

of stratospheric water vapour and its effect on the formation of PSCs are therefore of interest.

65

The stratospheric water vapour concentration is controlled by atmospheric dynamics and photochemistry. Its main sources are intrusion from the troposphere via the tropical tropopause (Brewer, 1949) and production through oxidation of methane and also molecular hydrogen (Bates and Nicolet,

- 70 1950; Le Texier et al., 1988). When rising air masses pass through the cold tropical tropopause region, moisture is removed due to freezing and sedimentation of particles (Brewer, 1949). The variability in the entry of water vapour into the stratosphere is largely controlled by the variability in the tropical cold point temperature. The oxidation of methane leads to formation of water through a series of reactions. The reaction with OH is the dominating methane loss reaction through most
- of the stratosphere, while the reaction with excited oxygen becomes increasingly important above 30 km, and photolysis is dominant above 65 km (Le Texier et al., 1988). In addition, the reaction with atomic chlorine has some significance as a sink for methane, but also as a termination reaction of ozone depleting cycles especially in the Antarctic vortex where denitrification reduces the importance of the reaction between ClO and NO₂ (Fahey et al., 1990).
- Due to the cold tropical tropopause, only a small fraction of tropospheric water vapour propagates to the stratosphere. As a result, the stratosphere is very dry, but it exhibits considerable variability both in space and time. The exceptional dryness of the stratosphere makes observation of stratospheric water vapour challenging. Long-term time series of stratospheric water vapour are rare, which complicates the study of concentration trends. Frostpoint hygrometer soundings have
- 85 been performed in Boulder, Colorado, since 1980 and for shorter periods of time also in other locations, including Sodankylä, Finland (Oltmans et al., 2000). Additionally, global data is available from satellite instruments, but only for a limited time span, for example from Microwave Limb Sounder (MLS) on board the Upper Atmosphere Research Satellite (UARS) (1991–1993) and the Earth Observing System on Aura (EOS-Aura) (2004–today) (e.g., Lambert et al., 2007). The
- 90 Odin Sub-Millimetre Radiometer (Odin/SMR) and the Envisat Michelson Interferometer for Passive Soundings (Envisat/MIPAS) provides also global coverage on a daily basis. Also the Stratospheric Aerosol and Gas Experiment II and III (ERBS/SAGE-II, Meteor-3M/SAGE-III), Polar Ozone and Aerosol Measurement (SPOT-4/POAM III), the UARS Halogen Occultation Experiment (UARS/HALOE), the Odin Sub-Millimetre Radiometer (Odin/SMR), the SCanning Imaging Absorption spectroMe-
- 95 ter for Atmospheric ChartographY (Envisat/SCIAMACHY), the Envisat Michelson Interferometer

for Passive Soundings (Envisat/MIPAS), Solar Occultation for Ice Experiment (AIM/SOFIE) and the SCISAT Atmospheric Chemistry Experiment Fourier Transform Spectrometer (SCISAT-1/ACE-FTS) have measured stratospheric water vapour in the Northern high latitudes, but the spatial and temporal coverage is limited.

- Several studies have used these available measurements to look into water vapour trends, especially in the mid-latitudes. Oltmans et al. (2000) analysed frostpoint hygrometer measurements above Boulder Colorado and reported a trend of about $+0.048 \pm 0.001$ ppm yr⁻¹ between 1980 and 2000 at the level of 18–20 km. Randel et al. (2004) compared the Boulder data to the HALOE measurements and reported differences between the Boulder data set and HALOE water vapour
- 105 data. The seasonal and interannual changes were comparable, but the long term increase observed in soundings were not seen in HALOE data. Later Scherer et al. (2008) did corrections for the instrumental bias of the Boulder frostpoint hygrometer data, updated the Boulder trend and reported a trend of +0.03–0.04 ppm yr⁻¹ between 1980 and 2000 at the same altitude, but noted a sudden drop in the stratospheric water vapour beginning in 2001. Hurst et al. (2011) presented a new
- 110 trend analysis of the 30 year record of Boulder stratospheric water vapour measurements (1980–2010) and found a +1.0 ppm increase over that time period at 16–26 km altitude, with significant short term variability. However, Hegglin et al. (2014) suggested, based on merged satellite data set, that the Boulder time series is not globally representative and instead reported negative trends in mid- and high latitudes at 16 km altitude between the end of 1980s and 2010. Based on satel-
- 115 lite and sounding measurements Solomon et al. (e.g., 2010) Solomon et al. (2010) reported negative trends in Boulder and generally in the mid-latitudes at 18 km altitude between 2000 and 2009 while Hegglin et al. (e.g., 2014) Hegglin et al. (2014) showed that these negative trends are mainly related to the sudden drop in the water vapour in 2000 and that after a few years with very low water vapour mixing ratios a recovery started in 2005. Recently (Urban et al., 2014) Urban et al. (2014) reported
- 120 another drop in the tropical water vapour during 2011–2012.

On the other hand, due to the lack of long term time series, there have been very few studies of stratospheric water vapour trends in the Arctic, where variations in the water content can have large effects on spring-time ozone depletion. Recently-Hegglin et al. (2013) have compared water vapour climatologies from 13 satellite products within the SPARC data initiative and analysed also the

- 125 anomalies in the Northern extratropics water vapour. They found that the uncertainty in water vapour increases toward the polar regions, the mesosphere and the upper troposphere-lower stratosphere (UTLS) region. Hegglin et al. (2014) showed water vapour trends up to 80° N latitude for the time period between the late 1980s and 2010 and reported negative trends in the <u>lower</u> stratosphere. On the other hand one should keep in mind that the coverage before 1998, which was based on HALOE
- 130 and SAGE II, was not good optimal, which warrants some caution to the results.

In this study, we use the FinROSE chemistry transport model (FinROSE-CTM) (Damski et al., 2007; Thölix et al., 2010) to investigate the stratospheric water vapour in the Arctic for the period

1990–2014. The model is described in Section 2. In Sect. 3 we describe the water vapour distribution in FinROSE simulations and observations. The modelled water vapour is evaluated against soundings

- 135 at Sodankylä, Finland (67.4° N, 26.6° E) and MLS satellite observations. The studied period includes the exceptionally cold January 2010 Arctic vortex with large scale ice PSC formation, which was observed also by the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP), the primary instrument onboard Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO) space-borne lidar (Pitts et al., 2011). Section 4 shows the long term variations of water vapour and
- 140 its sources. Section 5 describes the dehydration frequency in the Arctic stratosphere from 1990 to 2014. Section 6 deals with the Arctic winter 2010, including results from the Lapland Atmosphere-Biosphere Facility (LAPBIAT-2) measurement campaign in January–March 2010.

2 Modelling and data

2.1 FinROSE

- 145 The FinROSE-CTM (Damski et al., 2007) is a global off-line chemistry-transport model describing the stratosphere and mesosphere. The model produces the distribution of 36 species and the chemistry scheme consists of 110 gas phase reactions and 37 photodissociation processes. Water vapour is produced from oxidation of methane and molecular hydrogen. The PSC scheme includes liquid binary aerosols (LBA), super-cooled ternary solutions (STS, type Ib) and solid nitric acid trihydrate
- 150 (NAT, type Ia) and ice (ice, type II) PSCs. The model chemistry includes altogether 30 heterogeneous reactions on/in liquid binary aerosols and type Ia, Ib and II PSCs. Particle sedimentation, leading to dehydration and denitrification of the stratosphere, is also included in the model. The heterogeneous chemistry scheme in FinROSE is based on the calculation of the composition and volume of sulphate aerosols and PSCs and the partitioning of species between gas phase and condensed phase.
- The composition of LBA and STS are calculated using the method by Carslaw (1995). The STS are not considered below the ice PSC formation temperature. The number density profile for LBA and STS is estimated from McLinden et al. (1999) and the sulphuric acid distribution $[\mu m^2 cm^{-3}]$ is based on 2-D model data (Bekki and Pyle, 1992). NAT formation is based on the thermodynamic equilibrium equations by (Hanson and Mauersberger, 1988)Hanson and Mauersberger (1988). The
- 160 model includes an option to include a supersaturation requirement for NAT and ice formation, but this option was not used in the simulations reported in this paper. The choice was made due to the relative modest resolution of the model. Co-existence of NAT and STS is allowed. A scheme for growth of NAT particles is included based on Fahey et al. (2001). The number density of NAT particles is initially assumed to be 1 cm^{-3} (Krämer et al., 2003). For large NAT particles the number
- 165 density is reduced. The temperature threshold for ice particle formation is based on expressions by Marti and Mauersberger (1993). The equilibrium pressure of nitric acid above ice is calculated ac-

cording to Hanson and Mauersberger (1988). The ice number density is assumed to be $0.04 \,\mathrm{cm}^{-3}$, as estimated from synoptic scale PSCs (Dye et al., 1992).

The chemical kinetics used in this work follow the recommendations by Sander et al. (2011) and 170 Atkinson et al. (2007). Photodissociation coefficients were calculated using the PHODIS radiative transfer model (Kylling et al., 1997) and were used in the model through look-up tables. The model transport is calculated using a flux-form semi-lagrangian transport scheme (Lin and Rood, 1996).

The tropopause height is calculated at every time step using potential vorticity as defining parameter. Model levels below ± 2 PVU are considered to be in the troposphere. The 380 K potential

- 175 temperature level is used to define the tropopause height between 20° S and 20° N. The tropopause is thus changing with time depending on meteorological conditions. The tropospheric concentrations of the chemical species are not calculated in the model but prescribed via model boundary conditions. Tropospheric water vapour and ozone were obtained from the ECMWF ERA-Interim reanalysis (Simmons et al., 2007; Dee et al., 2011). Tropospheric methane (CH₄) is from Global view-data
- 180 (http://www.esrl.noaa.gov/gmd/ccgg/globalview/ch4), nitrous oxide (N₂O) from Advanced Global Atmospheric Gases Experiment (AGAGE) data (Prinn et al., 2000),and halogens (Cly and Bry) are from Montzka et al. (2009) updated data. Carbon dioxide (CO₂) is based on global annual mean trend data (ftp://aftp.cmdl.noaa.gov/products/trends/co2). At the upper model boundary (0.1 hPa), climatological values averaged over 2005–2014 from MLS data were used for water vapour and
- 185 ozone. The model has also a tracer aimed for water vapour studies; a chemically passive tracer for describing the amount of water vapour entering through the tropopause.

In this study, the model was run with a horizontal resolution of $6^{\circ} \times 3^{\circ}$ (longitude × latitude) at 35 hybrid-sigma levels, from the surface up to 0.1 hPa (about 65 km). The wind, temperature and surface pressure fields were obtained from the European Centre for Medium-range Weather Forecasts (ECMWF) ERA-Interim reanalysis (Dee et al., 2011).

2.1.1 Water vapour and PSC measurements

190

2.2 Water vapour and PSC measurements

High resolution soundings of stratospheric water vapour from northern high latitudes are rare. However, such measurements have been been made at Sodankylä (67.4° N, 26.6° E), northern Finland
since early 2000 (Vömel et al., 2007a, c). Sodankylä site is representative of high latitude conditions in the northern Europe, and the upper air soundings in winter and spring sample air both inside and outside the polar stratospheric vortex. Here we have used stratospheric water vapour measurements from two atmospheric sounding campaigns and some additional soundings obtained outside the major campaigns. First larger set of observations set of sounding observations large enough

200 to derive some statistics of water vapour vertical distribution was obtained during the Lapland Atmosphere-Biosphere Facility (LAPBIAT) LAPBIAT Upper Tropospheric Lower Stratospheric Water Vapour Validation Project (LAUTLOS-WAVVAP) campaign in early 2004 (e.g., Deuber et al., 2005; Vömel et al., 2007c; Karpechko et al., 2007; Suortti et al., 2008). The second campaign (the LAPBIAT-2 Atmospheric Sounding Campaign) took place in January–March 2010 (Kivi et al., 2010;

- 205 Khaykin et al., 2013; Engel et al., 2014; Grooß et al., 2014). During these campaigns two types of frost point hygrometers were flown. The NOAA frostpoint instrument (Oltmans, 1985; Vömel et al., 1995) was flown during the first campaign, while the Cryogenic Frostpoint Hygrometer (CFH) was deployed during both campaigns. CFH is a well characterised instrument capable of accurate water vapour measurements in the lower stratosphere typically up to the altitude of 25–28 km
- 210 (Vömel et al., 2007a, b). In this study we have used in total 13 NOAA and CFH frostpoint hygrometer profiles obtained during the first campaign and 13 CFH soundings obtained during the second atmospheric sounding campaign. In addition we included 8 NOAA or CFH soundings outside the main campaign periods, in order to improve temporal coverage. These additional soundings were made during January-February in year 2003, 2006, 2008, 2013 and 2014.
- 215 In addition to the balloon soundings, observations from the Microwave Limb Sounder (MLS) MLS on board the Aura satellite provide global profile measurements of H₂O, temperature and several trace gases (Lambert et al., 2007). In this study, we use the MLS version 3.3 Level 2 data, which are available from August 2004 to present. The data is published in EOS MLS Science team (2011) and it is accessed at http://disc.sci.gsfc.nasa.gov/datacollection/ML2H2O_V003.html. The
- 220 Level 2 data are produced on pressure surfaces from 316 to 0.1 hPa with a vertical resolution of about 3 km. Each day about 3500 vertical profiles are measured along a sun-synchronous suborbital track. For Sodankylä we used MLS overpass-data from Aura validation data center. All profiles within 300 km from Sodankylä were averaged for getting daily profiles of water vapour. We also used polar stratospheric cloud observations provided by the Cloud-Aerosol Lidar and Infrared Pathfinder
- 225 Satellite Observation (CALIPSO) CALIPSO space-borne lidar (Pitts et al., 2007). The CALIPSO PSC algorithm classifies PSCs by composition. Six different classes are defined: supercooled ternary solution (STS), two classes of liquid/NAT mixtures and mix 2 enhanced and water ice (including synoptic-scale ice and wave ice) (Pitts et al., 2011). We used these CALIPSO PSC composition classes for calculating the areas where PSCs were observed. The area is calculated separately for ice
- 230 and NAT. Both water ice and wave ice are included in the ice area and all the NAT mix classes to the NAT area. CALIPSO data is available from June 2006 to present.

3 Water vapour distribution

The FinROSE-CTM has been run using ERA-Interim meteorology and ERA-Interim water vapour data as tropospheric boundary condition. Thus, the evolution of water vapour in the FinROSE model

235 is strongly constrained by the water vapour at the ERA-Interim tropopause. Kunz et al. (2014) recently compared ERA-Interim water vapour in UTLS against independent sounding observations and found that while in the majority of the cases the agreement is satisfactory, in some cases the discrepancies between ERA-Interim and observations are large. Thus one can expect that these biases would affect FinROSE simulations in the stratosphere. Nevertheless, since the description of strato-

spheric H_2O in the ECMWF model is simplified (Monge-Sanz et al., 2013), the chemistry scheme in FinROSE produces a more realistic water vapour distribution, as we show in the manuscript.

First, we evaluated the simulated stratospheric water vapour distribution from FinROSE against measurements above Sodankylä. Figure 1 shows simulated and measured climatologies of water vapour distribution over Sodankylä between 2004 and 2014. Overall, the FinROSE (top panel) is

- 245 capable of reproducing the MLS observations (middle panel) of water vapour concentration and its vertical and temporal distributions. The maximum values of water vapour are located at the same altitude in both data sets. The largest differences are between 10 and 1 hPa: in winter/spring the concentration in FinROSE is about 1 ppm higher compared to MLS but in summer MLS is about 0.3 ppm moister than FinROSE. Too moist air in FinROSE in spring can lead to overestimated
- 250 PSC formation. In comparison, the ECMWF ERA-Interim reanalysis (bottom panel) clearly underestimates the observed water vapour concentrations in the upper stratosphere , while the by 1 ppm, which is likely due to a relaxation of water vapour towards an equivalent value applied in ERA-Interim at the stratopause citepdet02. The lower stratosphere compares well with MLS. The largest discrepancies between ERA-Interim and MLS are seen in the upper stratosphere where
- 255 ERA-Interim underestimates water vapour by 1. The dryness in the reanalysis data is likely a consequence of cold bias in the tropics in the ERA-Interim data (Schoeberl et al., 2012). Also the methane parameterisation in the ECMWF model leads to <u>a</u> too dry air (Dethof, 2003). It is also possible that too fast general circulation previously identified in the ECMWF model causes reduced moisture in the polar regions (Simmons et al., 1999; Schoeberl et al., 2012; Monge-Sanz et al., 2013). Note that
- 260 these problems with the general circulation affect FinROSE simulations because ERA-Interim meteorology is used. However, the full chemistry of the FinROSE improves the water vapour distribution of the model. During winter and spring the very top levels of ERA-Interim are too moist compared to observations. This is probably due to a too low model upper boundary; the ECMWF model does not extend to the upper mesospheric altitudes where photochemical processes destroy water vapour
- 265 causing the observed dry upper stratosphere. This process is also missing from the FinROSE; however, it has been indirectly included by using a water vapour climatology calculated from MLS data as an upper boundary condition. Overall, Fig. 1 shows that FinROSE is capable of simulating the distribution and magnitude of stratospheric water vapour in the high northern latitudes, which gives us confidence in its applicability for a more detailed study of water vapour distribution, sources, and
- 270 long-term variability.

For a more detailed comparison of model results with observations at northern high latitudes, we calculated the average mixing ratios and standard deviations of water vapour profiles above Sodankylä from the FinROSE model, ERA-Interim and observations. We chose January–February, because of the availability of balloon soundings during this time. In this comparison ERA-Interim and

- 275 FinROSE data are available daily from all the January–February months between 2004 and 2013, MLS almost daily, but soundings are available less frequently (altogether 34 profiles). From Fin-ROSE and ERA-Interim the gridpoint closest to Sodankylä has been chosen (Lon=30° E, lat=69° N). From MLS all the profiles measured within 300 km from the Sodankylä gridpoint and flagged as good quality are used. Standard deviation is calculated across the individual profiles for each data
- 280 set and thus represents uncertainty due to natural variability and random measurements errors. Figure 2 compares January–February-mean water vapour mixing ratios and standard deviations above Sodankylä from the FinROSE simulation (black), ECMWF ERA-Interim reanalysis (green), MLS satellite measurements (blue), and frostpoint hygrometer soundings (red). The data are shown for seven different pressure levels; 100, 56, 30, 21, 10, 3 and 1 hPa. Left panel shows the mixing ratios
- 285 in winter 2010 and right panel shows the relative differences compared to MLS observations calculated over winters 2004–20142005–2014. The winter 2010 was chosen for comparison because of the largest amount of soundings.

The modelled water vapour concentration profile agrees well with MLS measurements in winter 2010 (shown in Fig. 2 left panel). The model data is within 0–0.5 ppm of the MLS data, except at 3 hPa where the model gives 0.8 ppm more water vapour. Compared to the Sodankylä sound-ings, FinROSE has about 0.7 ppm more water vapour at 100 hPa, but the difference decreases with

- altitude, except at 30 hPa altitude where the model is again about 0.7 ppm moister. At the levels between 100 and 21 hPa the soundings fit to the range of variation of MLS. The difference is less than 0.5 ppm. Balloon sounding data are not available above 20 hPa. ERA-Interim is generally drier
- 295 compared to MLS and soundings throughout the studied altitude range. ERA-Interim water vapour concentration is also always about 0.7 ppm lower than the FinROSE's which is consistent with Fig. 1.

The right panel shows the differences between models and observations averaged over several winters. The differences have been calculated using all the available data pairs during Januaries and Februaries between 2004 and 2014 and then averaged. For FinROSE and ECMWF all the MLS profiles could have been used, but for sounding only the coincident MLS profiles were used. The differences between FinROSE and MLS, ERA-Interim and MLS and soundings and MLS remain smaller than 10 % at all altitudes. ERA-Interim is drier than MLS also in this climatology but Fin-ROSE is moister than MLS except at the 21 hPa level. Soundings are also drier compared to MLS,

305 but the difference is smaller than the difference between ERA-Interim and MLS. Sounding versus model comparisons are complicated, because firstly the number of soundings is limited and secondly, some of the soundings are obtained in the vicinity of the stratospheric vortex where the spatial water vapour gradients are large. Model to satellite differences have been calculated using all available data, both vortex conditions and non-vortex conditions included.

310 4 The origin and long term variability of water vapour

The sources of stratospheric water vapour are transport from the tropical troposphere and chemical production mainly from methane oxidation. Water vapour enters the stratosphere through the tropical tropopause and propagates then to the upper altitudes and higher latitudes. The FinROSE model has a tracer for studying these two water vapour sources. A passive H_2O -tracer, that is not affected by

- 315 chemistry, represents the transported water vapour. The difference between H_2O -tracer and H_2O represents the amount of water vapour produced by chemistry, i.e. mainly through oxidation of methane, but also hydrogen. Figure 3 shows the fractions of water vapour due to transport (upper panel) and chemistry (lower panel) according to simulations. Transport from the troposphere covers more than a half of the water vapour. At lower altitudes the transported part is clearly the most
- 320 important one. The chemically produced water vapour becomes more important at higher altitudes with a maximum between 1 and 3 hPa. In the summer and autumn, the fraction of chemistry part reaches almost 50 % there.

The water vapour variability and trends above Sodankylä, was investigated from a FinROSE model simulation covering the years 1986–2014. The first four years were discarded as spin-up

- 325 and the period 1990–2014 is analysed below. Figure 4 compares monthly-mean water vapour mixing ratios at 56 hPa above Sodankylä from the FinROSE simulation, ECMWF ERA-Interim reanalysis and MLS satellite measurements. The red dots denote individual sounding measurements. Throughout the investigated period, the water vapour amount in FinROSE and the observations varies within about 4–5.5 ppm. The modelled water vapour concentration agrees well with MLS
- 330 measurements from 2005–2008; however, after that FinROSE shows higher concentrations than observed by MLS and thus reveals a stronger increasing trend in these last four years. The growth rate is about 1 ppm decade⁻¹ in FinROSE but only 0.6 ppm decade⁻¹ in MLS. As can be expected based on Fig. 1, the ERA-Interim water vapour concentration is about 0.5 ppm lower than in FinROSE. It is, however, noteworthy that the difference remains approximately the same throughout the study
- 335 period. The increasing trend from 2007 to 2012 is very similar in FinROSE and ERA-Interim data, resulting from the use of ERA-Interim meteorology as driver data in the FinROSE simulation.

Figure 5 shows the anomaly of FinROSE water vapour and the sources of it between latitudes $70-90^{\circ}$ N. All the anomalies in the figure are calculated with respect to the mean values for the years period 1990-2014 for FinROSE and ERA-Interim and for the period 2004–2014 for MLS anoma-

- 340 lies. The altitudes of the panels are 1, 10, 56 and 100 hPa. At all the levels there are small positive trend-trends in the water vapour from the beginning of the timeseries until years 1994–1995. In the lower stratosphere (100–56 hPa) the anomalies decrease until 1998 and then stay constant until 2007. Thereafter a strong increase lasted until 2012 followed by another decrease, in agreement with observations by Urban et al. (2014) in the tropics. In the upper stratosphere (10–1 hPa) the
- water vapour decreased from 1995 until about 2004 before starting to increase around 2007. At 10 hPa the increase stopped by 2014 the end of 2013 but at 1 hPa it did not stop until the end of

the time series. Since the air in the upper polar stratosphere is older than that in the lower stratosphere (Stiller et al., 2012) the delay of between water vapour changes between in the lower and upper stratosphere suggests that these changes are driven by transport precesses,

350 consistent with our tracer results, which attribute most of the water vapour changes to dynamical processes.

In the FinROSE model results for the high northern latitudes the FinROSE simulations the long term change of stratospheric water vapour in the high northern latitudes is positive. The simulated water vapour is positive in the stratosphere. The water vapour trends for mid-latitudes and tropics

355 in FinROSE develop similarly as in are similar to those in higher latitudes (not shown). The anomalies seen in FinROSE also agree with the results by Dessler et al. (2013) for tropical water vapour between 2005 and 2013.

The Arctic water vapour concentration in FinROSE increases by about 0.8 ppm at altitude 56 hPa from year 2004 until year 2012. This increase corresponds to ca 1 K increase in the frostpoint

- 360 temperature (Marti and Mauersberger, 1993). However, the concentration during recent years (2012 to 2014) has decreased more than 0.5 ppm. The FinROSE water vapour anomaly and the passive tracer anomaly have nearly the same changes as the water vapour, in line with the results presented in Fig. 3. The evolution of the FinROSE water vapour anomaly is similar to the ERA-Interim anomaly (green line), which is expected as the FinROSE is driven by the ERA-Interim data. The chemical
- 365 part (purple line) which is mainly due to the contribution of methane oxidation, has only a small positive trend consistent with previous studies, because the stratospheric methane concentration was nearly stable in the analysed time period.

In order to attribute water vapour changes to physical processes we performed regression analysis following Dessler et al. (2014). We used three proxies: <u>qbo_QBO</u> index (QBO, equatorial winds at

- 50 hPa), Brewer Dobson circulation index (BD, residual vertical winds at 70 hPa averaged from 30° S to 30° N), and cold point temperature (CPT). Unlike Dessler et al. (2014) we found that the use of tropical temperatures at 500 hPa was not enough to explain the variability of the cold point temperature, and therefore used the cold point temperature as one of proxies. Although, there is some correlation between CPT and QBO (0.32) QBO also affect the transport of the water vapour
- 375 not directly influenced by CPT; therefore the use of both proxies is justified. We apply multiple regression analysis with all three proxies to water vapour time series averaged north of 70° N and at 82 hPa and 56 hPa. Cross-correlation analysis shows broad peaks at lags 6–12 months for the proxies. The maximum of the correlations of QBO and CPT with water vapour at 56 hPa is at about 10 months lag, and with 82 hPa is at 8–9 month lags, suggesting that propagation of the tropical
- 380 anomalies in the lower stratosphere is faster than that in the middle stratosphere, likely due to more efficient mixing. We use 10 month lag for all proxies for regression at 56 hPa and 9 month lag for the regression at 82 hPa.

The individual correlation and regression coefficients with our proxies taken at the lags mentioned earlier are shown in the Table 1. The main contribution to the polar water vapour variability is

- 385 CPT, followed by QBO. We found very weak contribution of BD proxy to the variability of the water vapour. One reason is that the effect of BD contribution is accumulated over time and this is not well represented by the monthly mean proxy. The multiple regression coefficients are 0.57 and 0.51 at 82 hPa and at 56 hPa correspondingly, showing that our models only explain 25–30 % of the variability. This is considerably less than that of Dessler et al. (2014) suggesting that different
- 390 processes contribute to the polar water vapour variability in comparison to those in the tropics. The modelled timeseries together with contributions of individual processes and residuals are shown in Figure 6. It is seen that the cold point variability and QBO make comparable contributions to the water vapour variability at both levels. Note that the regression somewhat explains the increase of the water vapour from 2005 to 2010, which is more clear at 82-hPa hPa. However the peak of the 395 water vapor during 2011–2013 is not explained by these proxies.

The Arctic water vapour concentration in FinROSE increases by about 0.8 at altitude 56 from year 2004 until year 2012. This increase corresponds to ca-1 increase in the frostpoint temperature (Marti and Mauersberger, 1993). However, the concentration during recent years (2012 to 2014) has decreased more than 0.5. The FinROSE water vapour anomaly and the passive tracer anomaly have

- 400 nearly the same changes as the water vapour, in line with the results presented in Fig. 1. The evolution of the FinROSE water vapour anomaly is similar to the ERA-Interim anomaly (green line), which is expected as the FinROSE is driven by the ERA-Interim data. The chemical part (purple line) which is mainly due to the contribution of methane oxidation, has only a small positive trend consistent with previous studies, because the stratospheric methane concentration was nearly stable in the analysed
 405 time period
- 405 time period.

5 Arctic dehydration frequency

In the polar vortex areas the concentration of water vapour can be changed also by dehydration. Water freezes into ice particles, sediments to lower altitudes and sublimates. Dehydration is frequently observed in the Antarctic polar vortex and there the magnitude of decrease of the water vapour by

- 410 dehydration is several ppms. The water vapour mixing ratio can be reduced to 1.5 ppm in the cold stable Antarctic vortex (Vömel et al., 1995). In the Arctic vortex the dehydration is rare, because the temperatures are higher. However, ice PSCs are formed also in the Arctic vortex and it is possible for dehydration to occur (Khaykin et al., 2013).
- Ice PSC formation is controlled by temperature and water vapour concentration, which both ex-415 hibit considerable variability. FinROSE simulates significant ice PSC conditions in the Arctic in 12 out of 25 winters (1990, 1993, 1995, 1996, 2000, 2005, 2008, 2010, 2011, 2012, 2013 and 2014). That means that nearly 50 % of winters offer conditions, which allow the formation of ice PSCs.

However, they cover only a small fraction of the vortex. The largest extent of simulated ice PSCs was seen in 1990, with an area of 3×10^6 km² at 56 hPa. In 1993, 2005, 2011, 2012 and 2014 an

- 420 area of around 2.5×10^6 km² were reached. Figure 7 shows the area of the gridpoints where ice PSC (black) form in the FinROSE model and the minimum temperature of the area between 50° N and 90° N at 56 hPa level. From 2007 onwards the ice PSC areas from CALIPSO lidar observations are shown as comparison (red). Based on the timeseries in Fig. 7 it seems that there is an increase in ice PSCs in the recent winters. Between 2007 and 2014 there are 6 ice PSC winters in FinROSE and 7
- 425 winters in CALIPSO, which leads to frequency 0.75 (0.87 for CALIPSO), considerably higher than the mean frequency for the whole studied period. This is likely a combined effect of cold conditions and the increase in the water vapour concentration. Another period with frequent ice PSC occurrence was between 1990 and 1996 when ice PSCs occurred in 4 out of 7 winters. The temperature conditions were cold in the beginning of the 90's, but water vapour concentrations were lower than
- 430 in the 2000's. The PSC areas in FinROSE agrees well with the ones calculated from CALIPSO data. However in some cases, e.g. beginning of winters, CALIPSO detects PSCs that are not simulated by FinROSE. This may indicate a warm bias in ERA-Interim in that period. Also in 2007 CALIPSO detects ice PSCs, which are not seen in the model. This might be due to the resolution of the model $(3^{\circ} \times 6^{\circ}, latitude \times longitude)$, as the areas are quite small.
- Figure 8 left panel shows the relation of the area of simulated and observed ice PSCs and the area with air colder than 188 K temperatures in December–February at the level of 56 hPa. One point or X denotes one winter day between 2007 and 2014 in FinROSE simulation or CALIPSO observation. The colour of the marks show the water vapour concentration averaged in the vortex. Figure 8 right panel shows the dependence of ice PSC on water vapour. It can be seen that while temperature is
- 440 the main factor controlling the ice PSC formation, the formation also noticeably depends on water vapour concentration. Higher water vapour concentrations produce larger areas of ice PSC but in dry vortex no ice PSCs form even when the temperature is low enough. The effect of both temperature and water vapour on ice PSC area is also seen in Table 2: for similar temperature conditions the ice PSC area increases when water vapour concentration increases. This holds for cases when areas
- with temperature below 188 K are small or large, and is seen in both FinROSE and CALIPSO. Consistent with expectations, the correlations between cold temperature area and ice PSC area are 0.89 for FinROSE and 0.64 for CALIPSO. The correlations between water vapour and ice PSC is 0.30 in FinROSE and 0.35 in CALIPSO, both statistically significant at 0.05 significance 5% level. CALIPSO detects small ice PSC areas with small cold temperature areas, but in case of FinROSE
- 450 the ice PSCs are not always created although the cold temperature area is large. The increase of water vapour in the vortex area have been about 0.8 ppm after 2007. That would have increased the ice PSC areas even if the temperatures have been the same, consistent with earlier estimations by (Kirk-Davidoff et al., 1999).

6 Case study: Winter 2009/2010

- 455 The winter 2009/2010 was unusually cold in the Arctic polar region. The temperature were was below 190 K in a large area of the polar vortex. In the lower stratosphere temperatures dropped below 195 K were simulated observed even south of 60° N. PSCs were formed and even persistent dehydration was observed over Northern polar latitudes for the first time (Khaykin et al., 2013). The vortex lasted until the beginning of April, and it was located above Sodankylä until 11th of February. Cold
- 460 conditions favouring PSC formations lasted until the end of January -2010 but a major stratospheric warming 2010. A major sudden stratospheric warming (SSW) took place in the end of January with the central date around 24 January rise temperatures and precluded further PSC formation (e.g., Khaykin et al., 2013; Dörnbrack et al., 2012; Pitts et al., 2011)(e.g., Khaykin et al., 2013; Dörnbrack et al., 2012; Pitts et al., 2 Following the SSW stratospheric temperatures rose above the PSC formation threshold so that no
- 465 more PSC were observed during that winter.

The FinROSE simulation was studied more thoroughly for the We use winter 2009/2010. The results were also compared to observations. Cold temperatures occurred between 20 and 28 in the ERA-Interim data and cold temperatures can be seen also in measured temperatures. Also a reduction 2010 as a case study and compare FinROSE simulations of water vapour at around 56, seen both in

- 470 the simulations and in observations, can be attributed to the formation of ice PSCs. Indeed, also ice PSC particles were simulated at the level 22–24 by the model and also seen in the balloon sounding data from the LAPBIAT campaign (Khaykin et al., 2013)and ice PSC to observations (see Figs. 9 and 10).
- Figure 9 shows the area of ice PSCs from CALIPSO (top panel) and from FinROSE (middle panel) and the area of temperatures colder than 188 K (lower panel) in the Arctic winter 2009/2010. Ice PSCs occur at the coldest dates and the same time both in FinROSE simulation and CALIPSO observations. The ice PSC areas in the beginning of January 2010 are smaller in FinROSE than in CALIPSO, but after mid January the areas are larger in FinROSE than in CALIPSO. However, the clouds appear at the same altitudes in the model as in observations. The timing is also comparable.
- 480 The differences might be attributed to the model coarse resolution and the simplicity of the PSC parameterization. The area of cold temperatures in the lowest panel is larger than the modelled and observed ice PSC areas.

Figure 10 shows maps of temperature from ERA-Interim, water vapour mixing ration from Fin-ROSE and ice PSC from FinROSE and CALIPSO from northern hemisphere vortex area at the

- 485 sounding dates between 17th January and 23th January 2010 at the level 35 hPa (24 km). The level 35 hPa was chosen because it was the coldest level with large ice PSCs. The minimum temperatures in the vortex (first row) are very low in this time period, even below 188 K. The water vapour mixing ratio from FinROSE (second row) is the highest at the vortex boundary and the lowest in the middle of the vortex. The driest areas coincide both in space and time with the coldest temperatures. Areas
- 490 with very cold temperatures correlates with very low water vapour content areas because of the ice

PSC formation. On 17th January the driest areas coincide both in space and time with the coldest temperatures. The water vapour is frozen and condensed into ice particles , which are sedimented and with proceeding time, ice particles grow and sediment downwards resulting in dehydrated air masses. The sedimentation of ice particles results in a redistribution of water vapour. On 20th–23rd

495 January the dehydrated air masses, where no ice presents move around the pole, but the minimum water vapour values do not coincide with coldest temperatures.

Figure 10 show that at 35 hPa ice PSCs were simulated by FinROSE in the same areas as they were observed by CALIPSO. To facilitate the comparison, CALIPSO observations have been gridded to the same grid than FinROSEgrid of $5 \times 10^{\circ}$, and the grid box is marked as having ice PSC if there

The bottom row in Fig. 10 shows modelled frostpoint temperature profiles above Sodankylä from

- 500 were at least one ice PSC observation within the grid box area. <u>CALIPSO measures only single</u> orbits and the gridded data appears sparce. Note that each day CALIPSO orbits cover only a part of the globe; therefore observations appear sparse. Nevertheless it can be seen that the grid boxes where CALIPSO observed ice PSC are scattered over larger areas in comparison to FinROSE simulations
- 505 FinROSE and temperature profiles from ECMWF ERA-Interim analysis, frostpoint temperature calculated from MLS satellite water vapour and frostpoint temperature from Sodankylä soundings. The FinROSE output at lon 30° E, lat 69° N located next to Sodankylä (26.6° E, 67.4° N) is shown. Overall FinROSE can simulate the frostpoint temperature quite well. The ECMWF ERA-Interim temperature reaches or almost reaches the frostpoint temperature during the analysed time period.
- 510 Then the formation of ice PSC is possible also in the FinROSE. The coldest date in Sodankylä is 17th of January. Very low frostpoint temperature values can be seen above 40 hPa altitudes also after 17th is likely a result of dehydration. This can be seen both in observations and in FinROSE. The water vapour concentration decreases about 1 to 1.5 ppm from the median values. A small increase of water vapour observed below 40 hPa on 23rd of January is interpreted as rehydration. It is not
- 515 simulated by FinROSE at the grid point shown, but is simulated at other coldest grid points near Sodankylä (not shown). In summary, FinROSE was able to reproduce well stratospheric water vapour and ICE ice PSC evolution during the record coldest period in winter 2010.

7 Conclusions

- FinROSE has been shown to be capable of simulating the water vapour distribution and evolution in
 the Northern high latitude stratosphere. The representation of water vapour in FinROSE is improved compared to the ERA-Interim data, even though the ERA-Interim data is used as tropospheric boundary condition. The full chemistry in FinROSE can add the water vapour to the produces more water than simplified parameterisation of ECMWF ERA-Interimwater vapour, alleviates the dry bias of ERA-Interim and improves the comparison with observations. The model gives results comparable
- to the MLS satellite measurements. However, some discrepancies compared to MLS remain.

The concentration of stratospheric water vapour in FinROSE is too high, especially in the summer time. Compared to the Sodankylä frost point hygrometer the model is too moist. However, the number of the comparisons is limited. In addition, some of the soundings have been made in the vicinity of the polar vortex, which further complicates comparison with the model data.

- 530 The main sources of the stratospheric water vapour are transport from the tropical troposphere and methane oxidation. A passive tracer was used in the FinROSE model for investigating the relative importance of the different sources of water vapour. The chemically produced fraction shows a maximum at altitudes between 6 hPa and 0.3 hPa. At these altitudes in the summertime the photochemical part is nearly as big as the transported part.
- The considerable decadal variability can be seen in water vapour below 10 hPa; water vapour increases between 2007 and 2012 by about 0.8 ppm in 5 years, which has been followed by a decrease. In the upper stratosphere the increase is smaller than in the lower stratosphere. In the MLS data the increase is smaller than seen in FinROSE. The increase can be attributed to water vapour transported trough the tropical tropopause while the contribution of the photochemically produced part of water
- 540 vapour to the increase is neglible, due to comparably smaller changes in the stratospheric methane concentration. A regression analysis suggests that cold point temperature variability and QBO make comparable contribution to the Arctic water vapour variability in the lower stratosphere; however they only explain 25-30% of the monthly variability.

In the FinROSE simulation ICE-ice PSCs occurred in Northern high latitudes in 10-12 out of 25 simulated winters. Comparison with CALIPSO instrument, which measured PSCs since 2006, shows that FinROSE simulates PSCs generally at the same time as observed by CALIPSO. Also the area of ice PSCs occurrence is comparable to the CALIPSO observations.

ICE Ice PSC have become more frequent in the recent years of the simulation. While cold temperatures observed during these years favoured ice PSC formations, our results suggest that increased

550 stratospheric water vapour concentration in high latitudes may have increased the ice PSC occurrence after year 2006.

Finally, as a test for faithfulness of FinROSE simulations, we perform a case study of the extremely cold winter 2009/2010. FinROSE is able to reproduce ice PSC extent and associated dehydration and rehydration at lower altitudes in good agreement with observations reported by (Khaykin et al., 2013)Khaykin et al. (2013)

555 FinROSE is also able to reproduce the observed magnitude of the dehydration associated with ice PSC formation, which is about 1 ppm. These results add credibility to FinROSE's ability to reproduce stratospheric water vapour changes.

Acknowledgements. We want to thank the UARS reference atmosphere project and MLS/Aura teams for water vapour data and the Atmospheric Science Data Center for the CALIPSO data. The MLS data were ob-

560 tained through the Aura MLS website (http://mirador.gsfc.nasa.gov/). CALIPSO data were obtained from the NASA Langley Research Center Atmospheric Science Data Center. The Sodankyl ("aä water vapour sounding providers are also thanked. The funding from the Academy of Finland through the UTLS project (140408) is gratefully acknowledged.

References

565 Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume III – gas phase reactions of inorganic halogens, Atmos. Chem. Phys., 7, 981–1191, doi:10.5194/acp-7-981-2007, 2007.

Bates, D. R. and Nicolet, M.: The photochemistry of water vapor, J. Geophys. Res., 55, 301-327, 1950.

- 570 Bekki, S. and Pyle, J.: Two-dimensional assessment of the impact of aircraft sulphur emissions on the stratospheric sulphate aerosol layer, J. Geophys. Res., 9, 15839–15847, 1992.
 - Brewer, A. W.: Evidence for a world circulation provided by the measurements of helium and water vapor distribution in the stratosphere, Q. J. Roy. Meteor. Soc., 75, 351–363, 1949.
 - Carslaw, K. S., Luo, B. and Peter, T.: An analytic expression for the composition of aqueous HNO3 -H2SO4
- 575 stratospheric aerosols including gas phase removal of HNO3. Geophys. Res. Lett., 22, 14, 1877–1880, 1995. Dameris, M., and Godin-Beekmann, S. (Lead Authors), Alexander, S., Braesicke, P., Chipperfield, M., de Laat, A. T. J., Orsolini, Y., Rex, M. and Santee, M. L.: Update on Polar ozone: Past, present, and future, Chapter 3 in Scientific Assessment of Ozone Depletion: 2014, Global Ozone Research and Monitoring Project – Report No. 55, World Meteorological Organization, Geneva, Switzerland, 2014.
- 580 Damski, J., Thölix, L., Backman, L., Taalas, P., and Kulmala, M.: FinROSE middle atmospheric chemistry and transport model, Boreal Environ. Res., 12, 535–550, 2007.
 - Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Holm, E. V., Isaksen,
- L., Kallberg, P., Kohler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park,
 B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thepaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, Q. J. Roy. Meteorol. Soc., 137, 553–597, 2011.

Dethof, A.: Aspects of modelling and assimilation for the stratosphere at ECMWF, SPARC Newslett., 2, 11-14,

590 2003.

- Deuber, B., Haefele, A., Feist, D. G., Martin, L., Kampfer, N., Nedoluha, G. E., Yushkov, V., Khaykin, S., Kivi, R., and Vömel, H.: Middle Atmospheric Watre Vapor Radiometer (MIAWARA): validation and first results of the LAPBIAT Upper Tropospheric Lower Stratospheric Water Vapor Validation Project (LAUTLOS-WAVVAP) campaign, J. Geophys. Res., 110, D13306, doi:10.1029/2004JD005543, 2005.
- 595 Dessler, A. E., Schoeberl, M. R., Wang, T., Davis, S. M., and Rosenlof, K. H.: Stratospheric water vapor feedback, P. Natl. Acad. Sci. USA, 45, 18087–18091, doi:10.1073/pnas.1310344110, 2013.
 - Dessler, A. E., Schoeberl, M. R., Wang, T., Davis, S. M., Rosenlof, K. H., Vernier, J.-P.: Variations of stratospheric water vapor over the past three decades. J. Geophys. Res. 119, 12588–12598. doi:10.1002/2014JD021712, 2014.
- 600 Dörnbrack, A., Pitts, M. C., Poole, L. R., Orsolini, Y. J., Nishii, K. and Nakamura, H.: The 2009–2010 Arctic stratospheric winter – general evolution, mountain waves and predictability of an operational weather forecast model, Atmos. Chem. Phys., 12, 3659–3675, doi:10.5194/acp-12-3659-2012, 2012.

Dye, J. E., Baumgardner, D., Gandrud, B. W., Kawa, S. R., Kelly, K. K., Loewenstein, M., Ferry, G.V., Chan, K. R. and Gary, B. L.: Particle size distributions in arctic polar stratospheric clouds, growth and freezing of sulfuric acid droplets, and implications for cloud formation. J. Geophys. Res. 97: 8015–8034, 1992.

Engel, I., Luo, B. P., Khaykin, S. M., Wienhold, F. G., Vömel, H., Kivi, R., Hoyle, C. R., Grooß, J.-U., Pitts, M. C., and Peter, T.: Arctic stratospheric dehydration – Part 2: Microphysical modeling, Atmos. Chem. Phys., 14, 3231–3246, doi:10.5194/acp-14-3231-2014, 2014.

605

- EOS MLS Science Team: MLS/Aura Level 2 Water Vapor (H2O) Mixing Ratio,version 003,
 Greenbelt, MD, USA:NASA Goddard Earth Science Data and Information Services Center. http://disc.sci.gsfc.nasa.gov/datacollection/ML2H2O_V003.html, 2011.
 - Fahey, D. W., Kelly, K. K., Kawa, S. R., Tuck, A. F., Loewenstein, M., Chan, K. R., and Heidt, L. E.: Observations of denitrification and dehydration in the winter polar stratospheres, Nature, 344, 321–324, doi:10.1038/344321a0, 1990.
- 615 Fahey, D.W., Gao, R.S., Carslaw, K.S., Kettleborough, J., Popp, P.J., Northway, M.J., Holecek, J.C., Ciciora, S.C., McLaughlin, R.J., Thompson, T.L., Winkler, R.H., Baumgardner, D.G., Gandrud, B., Wennberg, P.O., Dhaniyala, S., McKinney, K., Peter, Th., Salawitch, R.J., Bui, T.P., Elkins, J.W., Webster, C.R., Atlas, E.L., Jost, H., Wilson, J.C., Herman, R.L., Kleinböhl, A. and von König, M.: The detection of large nitric-acid particles in the winter Arctic stratosphere, Science 291: 1026–1031, 2001.
- 620 Gettelman, A., Birner, T., Eyring, V., Akiyoshi, H., Bekki, S., Brühl, C., Dameris, M., Kinnison, D. E., Lefevre, F., Lott, F., Mancini, E., Pitari, G., Plummer, D. A., Rozanov, E., Shibata, K., Stenke, A., Struthers, H. and Tian, W.: The Tropical Tropopause Layer 1960–2100, Atmos. Chem. Phys., 9, 1621–1637, doi:10.5194/acp-9-1621-2009, 2009.
 - Grooß, J.-U., Engel, I., Borrmann, S., Frey, W., Günther, G., Hoyle, C. R., Kivi, R., Luo, B. P., Molleker, S.,
- 625 Peter, T., Pitts, M. C., Schlager, H., Stiller, G., Vömel, H., Walker, K. A., and Müller, R.: Nitric acid trihydrate nucleation and denitrification in the Arctic stratosphere, Atmos. Chem. Phys., 14, 1055–1073, doi:10.5194/acp-14-1055-2014, 2014.
 - Hanson, D. and Mauersberger, K.: Vapor pressures of HNO₃/H₂O solutions at low temperatures, J. Phys. Chem., 92, 6167–6170, 1988.
- 630 Hegglin, M., Tegtmeier, S., Anderson, J., Froidevaux, L., Fuller, R., Funke, B., Jones, A., Lingenfelser, G., Lumpe, J., Pendlebury, D., Remsberg, E., Rozanov, A., Toohey, M., Urban, J., von Clarmann, T., Walker, K. A., Wang, R., and Weigel, K.: SPARC Data Initiative: comparison of water vapor climatologies from international satellite limb sounders, J. Geophys. Res., 118, 11824–11846, doi:10.1002/jgrd.50752, 2013.
- 635 Hegglin, M. I., Plummer, D. A., Shepherd, T. G., Scinocca, J. F., Anderson, J., Froidevaux, L., Funke, B., Hurst, D., Rozanov, A., Urban, J., von Clarmann, T., Walker, K. A., Wang, H. J., Tegtmeier, S., and Weigel, K.: Vertical structure of stratospheric water vapour trends derived from merged satellite data, Nat. Geosci., 7, 768–776, doi:10.1038/ngeo2236, 2014.
 - Hintsa, E. J., Newman, P. A., Jonsson, H. H., Webster, C. R., May, R. D., Herman, R. L., Lait, L. R., Schoeberl,
- 640 M. R., Elkins, J. W., Wamsley, P. R., Dutton, G. S., Bui, T. P., Kohn, D. W. and Anderson, J. G.: Dehydration and denitrification in the Arctic polar vortex during the 1995–1996 winter, Geophys. Res. Let., 25, 501–504, 1998.

Hurst, D. F., Oltmans, S. J., Vömel, H., Rosenlof, K. H., Davis, S. M., Ray, E. A., Hall, E. G., and Jordan, A. F.: Stratospheric water vapor trends over Boulder, Colorado: analysis of the 30 year Boulder record, J. Geophys.

- 645 Res., 116, D02306, doi:10.1029/2010JD015065, 2011.
- Karpechko, A., Lukyanov, A., Kyrö, E., Khaikin, S., Korshunov, L., Kivi, R., and Vömel, H.: The water vapour distribution in the Arctic lowermost stratosphere during the LAUTLOS campaign and related transport processes including stratosphere-troposphere exchange, Atmos. Chem. Phys., 7, 107–119, doi:10.5194/acp-7-107-2007, 2007.
- 650 Kelly, K. K., Tuck, A. F., Murphy, D. M., Proffitt, M. H., Fahey, D. W., Jones, R. L., Mckenna, D. S., Loewenstein, M., Podolske, J. R., Strahan, S. E., Ferry, G. V., Chan, K. R., Vedder, J. F., Gregory, G. L., Hypes, W. D., Mccormick, M. P., Browell, E. V., and Heidt, L. E.: Dehydration in the lower Antarctic stratosphere during late winter and early spring, 1987, J. Geophys. Res, 94, 11317–11357, doi:10.1029/JD094iD09p11317, 1989.
- 655 Meier, S., Naebert, T., Pitts, M. C., Santee, M. L., Spelten, N., Wienhold, F. G., Yushkov, V. A., and Peter, T.: Arctic stratospheric dehydration – Part 1: Unprecedented observation of vertical redistribution of water, Atmos. Chem. Phys., 13, 11503–11517, doi:10.5194/acp-13-11503-2013, 2013.

Kirk-Davidoff, D. B., Anderson, J. G., Hintsa, E. J., and Keith, D. W.: The effect of climate change on ozone depletion through changes in stratospheric water vapour, Nature, 402, 399–401, 1999.

Khaykin, S. M., Engel, I., Vömel, H., Formanyuk, I. M., Kivi, R., Korshunov, L. I., Krämer, M., Lykov, A. D.,

- 660 Kirner, O., Müller, R., Ruhnke, R., and Fischer, H.: Contribution of liquid, NAT and ice particles to chlorine activation and ozone depletion in Antarctic winter and spring, Atmos. Chem. Phys., 15, 2019–2030, doi:10.5194/acp-15-2019-2015, 2015.
 - Kivi, R., Vömel, H., Immler, F., Lehtola, T., Kämpfer, N., Straub, C., Yushkov, V., Khaykin, S., Christensen, T., and Wienhold, F. G.: LAPBIAT atmospheric sounding campaign in 2010, upper-air and remote sensing
- 665 observations of water vapour, in: WMO Technical Conference on Meteorological and Environmental Instruments and Methods of Observation (TECO-2010), Helsinki, Finland, 30 August–1 September 2010, Instruments and Observing System Methods Report No. 104, WMO/TD-No. 1546, 2010.

Krämer, M., Müller, R., Bovensmann, H., Burrows, J., Brinkmann, J., Röth, E.-P., Grooß, J.-U., Müller, R., Woyke, T., Ruhnke, R., Günther, G., Hendricks, J., Lippert, E., Carslaw, K. S., Peter, T., Zieger, A., Brühl,

- 670 C., Steil, B., Lehmann, R. and McKenna, D.S.: Intercomparison of stratospheric chemistry models under polar vortex conditions. J. Atmos. Chem., 45, 51–77, 2003.
 - Kunz, A., Spelten, N., Konopka, P., Müller, R., Forbes, R. M. and Wernli, H.: Comparison of Fast In situ Stratospheric Hygrometer (FISH) measurements of water vapor in the upper troposphere and lower stratosphere (UTLS) with ECMWF (re)analysis data. Atmos. Chem. Phys., 14, 10803–10822, doi:10.5194/acp-14-10803-2014. 2014.
- **675** 2014, 2014.
 - Kylling, A., Albold, A., and Seckmeyer, G.: Transmittance of a cloud is wavelength dependent in the UVrange: physical interpretation, Geophys. Res. Lett., 24, 397–400, doi:10.1029/97GL00111, 1997.
 - Lambert, A., Read, W. G., Livesey, N. J., Santee, M. L., Manney, G. L., Froidevaux, L., Wu, D. L., Schwartz, M. J., Pumphrey, H. C., Jimenez, C., Nedoluha, G. E., Cofield, R. E., Cuddy, D. T., Daffer, W. H.,
- 680 Drouin, B. J., Fuller, R. A., Jarnot, R. F., Knosp, B. W., Pickett, H. M., Perun, V. S., Snyder, W. V., Stek, P. C., Thurstans, R. P., Wagner, P. A., Waters, J. W., Jucks, K. W., Toon, G. C., Stachnik, R. A., Bernath, P. F., Boone, C. D., Walker, K. A., Urban, J., Murtagh, D., Elkins, J. W., and Atlas, E.: Validation of the Aura Mi-

crowave Limb Sounder middle atmosphere water vapor and nitrous oxide measurements, J. Geophys. Res., 112, D24S36, doi:10.1029/2007JD008724, 2007.

- 685 Langematz, U., Meul, S., Grunow, K., Romanowsky, E., Oberläder, S., Abalichin, J. and Kubin A.: Future Arctic temperature and ozone: The role of stratospheric composition changes, J. Geophys. Res. Atmos., 119, 2092–2112, doi:10.1002/2013JD021100, 2014.
 - Le Texier, H., Solomon, S., and Garcia, R. R.: The role of molecular hydrogen and methane oxidation in the water vapour budget of the stratosphere, Q. J. Roy. Meteorol. Soc., 114, 281–295, 1988.
- 690 Lin, S.-J. and Rood, R. B.: Multidimensional flux-form semi-lagrangian transport schemes, Mon. Weather Rev., 124, 2046–2070, 1996.
 - Marti, J. and Mauersberger, K.: A survey and new measurements of ice vapor pressure at temperatures between 170 and 250 K, Geophys. Res. Lett., 20, 363–366, doi:10.1029/93GL00105, 1993.
- McLinden, C. A., McConnell, J. C., McElroy, C. T. and Griffioen, E.: Observations of stratospheric aerosol using CPFM polarized limb radiances, JAS, 56, 233–240, 1999.
- Monge-Sanz, B. M., Chipperfield, M. P., Untch, A., Morcrette, J.-J., Rap, A., and Simmons, A. J.: On the uses of a new linear scheme for stratospheric methane in global models: water source, transport tracer and radiative forcing, Atmos. Chem. Phys., 13, 9641–9660, doi:10.5194/acp-13-9641-2013, 2013.
- Montzka, S. A., Butler, J. H., Elkins, J. W., Thompson, T. M., Clarke, A. D., and Lock, L. T.: Present and future
 trends in the atmospheric burden of ozone-depleting halogens Nature, 398, 690–694, 1999.
- Oltmans, S. J.: Measurements of water vapor in the stratosphere with a frost point hygrometer, measurement and control in science and industry, in: Proceedings of the 1985 International Symposium on Moisture and Humidity, Instrument Society of America, Washington, DC, 251–258, 1985.
- Oltmans, S. J., Vömel, H., Hofmann, D. J., Rosenlof, K. H., and Kley, D.: The increase in stratospheric water
- 705 vapor from balloonborne, frostpoint hygrometer measurements at Washington, D. C., and Boulder, Colorado, Geophys. Res. Lett., 27, 3453–3456, doi:10.1029/2000GL012133, 2000.
 - Pitts, M. C., Thomason, L. W., Poole, L. R., and Winker, D. M.: Characterization of Polar Stratospheric Clouds with spaceborne lidar: CALIPSO and the 2006 Antarctic season, Atmos. Chem. Phys., 7, 5207–5228, doi:10.5194/acp-7-5207-2007, 2007.
- Pitts, M. C., Poole, L. R., Dörnbrack, A., and Thomason, L. W.: The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective, Atmos. Chem. Phys., 11, 2161–2177, doi:10.5194/acp-11-2161-2011, 2011.
 - Prinn, R. G., Weiss, R. F., Fraser, P. J., Simmonds, P. G., Cunnold, D. M., Alyea, F. N., O'Doherty, S., Salameh, P., Miller, B. R., Huang, J., Wang, R. H. J., Hartley, D. E., Harth, C., Steele, L. P., Sturrock, G.,
- 715 Midgley, P. M., and McCulloch, A.: A history of chemically and radiatively important gases in air deduced from ALE/GAGE/AGAGE, J. Geophys. Res., 105, 17751–17792, 2000.
 - Randel, W., Wu, F., Oltmans, S., Rosenlof, K., and Nedoluha, G.: Interannual changes in stratospheric water vapor and correlations with tropical tropopause temperatures, J. Atmos. Sci., 61, 2133–2148, 2004.
- Randel, W., Wu, F., Vömel, H., Nedoluha, G. E. and Forster, P.: Decreases in stratospheric water vapor after
 2001: Links to changes in the tropical tropopause and the Brewer-Dobson circulation, J. Geophys. Res., 111,

D12312, doi:10.1029/2005JD006744, 2006.

- Rex, M., Salawitch, R. J., Deckelmann, H., von der Gathen, P., Harris, N. R. P., Chipperfield, M. P., Naujokat,
 B., Reimer, E., Allaart, M., Andersen, S. B., Bevilacqua, R., Braathen, G. O., Claude, H., Davies, J., De
 Backer, H., Dier, H., Dorokhov, V., Fast, H., Gerding, M., Godin-Beekmann, S., Hoppel, K., Johnson, B.,
- 725 Kyrö, E., Litynska, Z., Moore, D., Nakane, H., Parrondo, M. C., Risley, A. D., Skrivankova, P., Stübi, R., Viatte, P., Yushkov, V., and Zerefos, C.: Arctic winter 2005: Implications for stratospheric ozone loss and climate change, Geophys. Res. Lett., 33, 123808, 2006.
 - Rieder, H. E. and Polvani, L. M.: Are recent Arctic ozone losses caused by increasing greenhouse gases?, Geophys. Res. Lett., 40, 4437–4441, 2013.
- 730 Sander, S. P., Abbatt, J., Barker, J. R., Burkholder, J. B., Friedl, R. R., Golden, D. M., Huie, R. E., Kolb, C. E., Kurylo, M. J., Moortgat, G. K., Orkin, V. L., and Wine, P. H.: Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies, Evaluation No. 17, JPL Publication 10-6, Jet Propulsion Laboratory, Pasadena, USA, 2011.
 - Scherer, M., Vömel, H., Fueglistaler, S., Oltmans, S. J., and Staehelin, J.: Trends and variability of midlatitude
- stratospheric water vapour deduced from the re-evaluated Boulder balloon series and HALOE, Atmos. Chem.
 Phys., 8, 1391–1402, doi:10.5194/acp-8-1391-2008, 2008.
 - Schmidt, G. A., Ruedy, R. A., Miller, R. L., and Lacis, A. A.: Attribution of the present-day total greenhouse effect, J. Geophys. Res., 115, D20106, doi:10.1029/2010JD014287, 2010.
- Schoeberl, M. R., Dessler, A. E., and Wang, T.: Simulation of stratospheric water vapor and trends using three
 reanalyses, Atmos. Chem. Phys., 12, 6475–6487, doi:10.5194/acp-12-6475-2012, 2012.
- Shindell, D. T., Rind, D. end Lonergan, P.: Increased polar stratospheric ozone losses and delayed eventual recovery owing to increasing greenhouse-gas concentrations, Nature 392, 589–592, 1998.
 - Simmons, A. J., Untch, A., Jakob, C., Kållberg, P., and Undén, P.: Stratospheric water vapour and tropical tropopause temperatures in ECMWF analyses and multi-year simulations, Q. J. Roy. Meteorol. Soc., 125,
- 745 353–386, 1999.
 - Simmons, A. J., Uppala, S. M., Dee, D., and Kobayashi, S.: ERA-Interim: new ECMWF reanalysis products from 1989 onwards, ECMWF Newslett., 110, 25–35, 2007.
 - Solomon, S.: Stratospheric ozone depletion: a review of concepts and history, Rev. Geophys., 37, 275–316, 1999.
- 750 Solomon, S., Garcia, R. R., Rowland, F. S., and Wuebbles, D. J.: On the depletion of Antarctic ozone, Nature, 321, 755–758, 1986.
 - Solomon, S., Rosenlof, K. H., Portmann, R. W., Daniel, J. S., Davis, S. M., Sanford, T. J., and Plattner, G.-K.: Contributions of stratospheric water vapor to decadal changes in the rate of global warming, Science, 327, 5970, 1219–1223, doi:10.1126/science.1182488, 2010.
- 755 Stiller, G. P., von Clarmann, T., Haenel, F., Funke, B., Glatthor, N., Grabowski, U., Kellmann, S., Kiefer, M., Linden, A., Lossow, S., and López-Puertas, M.: Observed temporal evolution of global mean age of stratospheric air for the 2002 to 2010 period, Atmos. Chem. Phys., 12, 3311–3331, doi:10.5194/acp-12-3311-2012, 2012.

<sup>Suortti, T. M., Kats, A., Kivi, R., Kämpfer, N., Leiterer, U., Miloshevich, L. M., Neuber, R., Paukkunen, A.,
Ruppert, P., Vömel, H., and Yushkov, V.: Tropospheric comparisons of Vaisala radiosondes and balloon-</sup>

Ruppert, I., voniei, II., and Tushkov, v.. Hopospheric comparisons of varsata radiosondes and ba

borne frost-point and lyman-alpha hygrometers during the LAUTLOS-WAVVAP experiment, J. Atmos. Ocean. Tech., 25, 149-166, 2008.

- Thölix, L., Backman, L., Ojanen, S.-M.: The effects of driver data on the performance of the FinROSE chemistry transport model, IJRS, 31, 6401-6408, 2010.
- 765 Tian, W. S., Chipperfield, M. P., and Lü, D. R.: Impact of increasing stratospheric water vapor on ozone depletion and temperature change, Adv. Atmos. Sci., 26, 423-437, doi:10.1007/s00376-009-0423-3, 2009.
 - Urban, J., Lossow, S., Stiller, G. and Read, W.: Another drop in water vapor, Eos Trans. AGU, 94, 245-246, doi:10.1002/2014EO270001, 2014.
- Vömel, H., Oltmans, S. J., Hofmann, D. J., Deshler, T., and Rosen, J. M.: The evolution of the dehydration in 770 the Antarctic stratospheric vortex, J. Geophys. Res., 100, 13919-13926, 1995.
- Vömel, H., Barnes, J., Forno, R., Fujiwara, M., Hasebe, F., Iwasaki, S., Kivi, R., Komala, N., Kyrö, E., Leblanc, T., Morel, B., Ogino, S.-Y., Read, W., Ryan, S. C., Saraspriya, S., Selkirk, H., Shiotani, M., Valverde-Canossa, J., Whiteman, D.: Validation of Aura/MLS water vapor by balloon borne cryogenic frostpoint hygrometer measurements, J. Geophys. Res., 112, D24S37, doi:10.1029/2007JD008698, 2007a.
- 775 Vömel, H., David, D. E., and Smith, K.: Accuracy of tropospheric and stratospheric water vapor measurements by the cryogenic frost point hygrometer: instrumental details and observations, J. Geophys. Res., 112, D08305, doi:10.1029/2006JD007224, 2007b.
 - Vömel, H., Yushkov, V., Khaykin, S., Korshunov, L., Kyrö, E., and Kivi, R.: Intercomparisons of Stratospheric Water Vapor Sensors: FLASH-B and NOAA/CMDL Frost-Point Hygrometer, J. Atmos. Ocean. Tech., 24, 941-952, 2007c.
- 780
 - Wohltmann, I., Wegner, T., Müller, R., Lehmann, R., Rex, M., Manney, G. L., Santee, M. L., Bernath, P., Sumińska-Ebersoldt, O., Stroh, F., von Hobe, M., Volk, C. M., Hösen, E., Ravegnani, F., Ulanovsky, A., and Yushkov, V.: Uncertainties in modelling heterogeneous chemistry and Arctic ozone depletion in the winter 2009/2010, Atmos. Chem. Phys., 13, 3909-3929, doi:10.5194/acp-13-3909-2013, 2013.

Table 1. Correlation coefficients Coefficients of correlation and regression. The units for the regression coefficients are ppmv K^{-1} for CPT, ppmv $m^{-1} s^{-1}$ for QBO, and ppmv mm⁻¹ s⁻¹. Regression coefficients are shown together with the 95% confidence intervals.

		СРТ	QBO	BD	Multiple
$56\mathrm{hPa}$		0.45	0.32	-0.21	0.51
	Rgression	<u>0.12±0.03</u>	0.003±0.002	<u>-1.51±0.83</u>	~
82 hPa	Correlation	0.52	0.40	-0.18	0.57
	Regression	$\underbrace{0.13 \pm 0.03}_{0.03}$	$\underbrace{0.004 \pm 0.002}_{0.004 \pm 0.002}$	<u>-1.15±0.80</u>	$\overline{\sim}$



Figure 1. Climatology (2004-2014) of the water vapour distribution from FinROSE, MLS v3.3 data and ECMWF ERA-Interim above Sodankylä.

Table 2. Lee Winter mean ice PSC areas (10^{6} km^{2}) in FinROSE (left) and CALIPSO (right) at 56 hPa pressure level as a function of 56 hPa monthly mean water vapour concentration (ppm) averaged north of 60° N and the area of temperatures below 188K (10^{6} km^{2}) . The number of hits months in each group are shown in parentheses.

Water vapour (ppm)									
$\frac{\text{Area T} < 188 \text{ K}}{(10^6 \text{ km}^2)}$		FinROSE			CALIPSO				
	Wet (>5.4)	5.2–5.4	Dry (<5.2)	Wet (>5.4)	5.2–5.4	Dry (<5.2)			
Warm (0.5–1.5)	1.61 (10)	1.09 (10)	0.31 (4)	0.70 (4)	0.46 (9)	0.18 (2)			
Cold (1.5–3.0)	2.49 (10)	2.48 (10)	1.28 (4)	0.96 (4)	1.20 (9)	0.24 (2)			



Figure 2. (a) 2010 mean January and February water vapour mixing ratio and standard deviation above Sodankylä as function of pressure. (b) Difference of the modelled or observed winter water vapour mixing ratio and MLS calculated over years 2005 to 2014 above Sodankylä as function of pressure. FinROSE (black), MLS (blue), ECMWF ERA-Interim (green) and soundings (red).



Figure 3. Monthly mean of (**a**) fraction of transported water vapour and (**b**) fraction of chemically produced water vapour (%) for 2010 calculated over 70–90° N.



Figure 4. Sodankylä monthly mean water vapour mixing ratio from FinROSE (black), MLS (blue), ECMWF ERA-Interim (green) and soundings (red dots) at 56 hPa between 1990–2014.



Figure 5. (a) Anomalies of water vapour (black), tracer describing transported water vapour (orange), tracer describing water produced by methane oxidation (purple) from FinROSE and water vapour anomaly from ECMWF ERA-Interim (green) as ppm in 1990–2014 and, water vapour anomaly from MLS (blue) in 2004–2014. Anomalies are calculated at latitudes between $70–90^{\circ}$ N at the level of 1 hPa. (b-d) Same as panel (a), but levels 10, 56 and $\frac{1000}{100}$ hPa.



Figure 6. The contribution of individual processes to water vapour variability.



Figure 7. The area where ice PSCs formed in the FinROSE simulation (black) and was observed by CALIPSO (red) and the minimum temperature between $50-90^{\circ}$ N from ERA-Interim (green, right y axis) for winters between 1990 and 2014 at 56 hPa.



Figure 8. (a) Scatter plot of December–February ice PSC area versus the area of colder than 188 K in the northern hemisphere from FinROSE and CALIPSO at 56 hPa. The colour denotes the vortex mean water vapour content (ppm). (b) Scatter plot of the December–February ice PSC area versus the vortex mean water vapour content (ppm) from FinROSE and CALIPSO at 56 hPa. The colour denotes the vortex average temperature (K). FinROSE is shown with dots and CALIPSO is shown with crosses.



Figure 9. (a) Ice PSC area from CALIPSO, (b) Ice PSC area from FinROSE and (c) area of colder than 188 K from ECMWF ERA-Interim in winter 2009/2010.



Figure 10. Upper four rows: temperature, water vapour (ppm) and ice PSC occurrence from FinROSE and ice PSC occurrence from CALIPSO during the extreme cold period of winter 2010. The black contour marks the border of the vortex defined as modified PV > 36. All maps are at the 35 hPa pressure level. Lowest row: ERA-Interim temperature (green), frostpoint temperature from Sodankylä soundings (red dots), from MLS (blue) and from FinROSE (black). Purple dots shows the altitudes where FinROSE simulated ice PSC.