

Responses to the comments of referee#2

Interactive comment on “Fine particulate matter associated with monsoonal effect and the responses of biomass fire hotspots in the tropical environment” by M. F. Khan et al.

Anonymous Referee #2

Received and published: 13 September 2015

The manuscript of Khan et al. focuses on the health implications of fine particulate matter (PM_{2.5}) and its main constituents in the western part of Malaysia. The study period covers two monsoonal regimes, one coming from the south-west and one from the north-east and levels of PM, heavy metals and main ions are determined for both regimes. Overall it has been found that 48% of the samples were 4 times or more than guideline values, with the north-east monsoon regime exhibiting somewhat higher values. Based on the enrichment factor of metals it occurred that the majority of heavy metals were attributed to anthropogenic sources, while arsenic was identified as a potentially significant health concern as its concentrations were nearly equal to the WHO and US EPA guideline values, posing an exposure risk for inhabitants in the specific location. Using a mass closure model a mineral component was found to comprise 35% of the PM_{2.5} concentration, followed by 11% by secondary inorganic aerosol, while 45% was unidentified, possibly a large part of it being the organic aerosol fraction. Using positive matrix factorization (PMF) analysis as a source apportionment tool, motor vehicle and biomass burning emissions were the dominant sources, followed by marine and sulfate aerosol, coal burning, nitrate aerosol and mineral dust. Finally, based on the health risk assessment and the PMF-identified sources, the non- carcinogenic risk posed by exposure of PM was at a safer level than the respective one in the South and East Asian region while the cancer risk posed by the exposure to toxic metals was 3 to 4 in 1 million people in the specific location. Motor vehicle emissions and trans-boundary pollution were the major identified reasons for change in the chemical composition of PM in tropical Peninsular Malaysia.

The manuscript is well written and interesting, with an added value of the presented results being from an area of the globe that is not very well documented. It is a pity that analysis of the organic fraction is not available, as it is expected that a large portion of the unidentified mass from the mass closure model will be the organic component. There are several details missing and more thorough discussion should be made in specific sections. Other than that the paper can be recommended for publication after addressing the issues listed below.

- 1) One of the features of the manuscript that the authors draw our attention to is the comparison

between the $PM_{2.5}$ from the PMF and from the high volume samplers (HVS) which was found to have a slope of 0.91 and a very good correlation. It appears somewhat strange that even though 45% of the $PM_{2.5}$ mass remains unidentified, the comparison of the mass concentrations from the PMF and the HVS results in a slope of 0.91. Please comment. Also, do the authors believe that the remaining 45% of the unidentified mass, with the organics possibly being a large part of this mass, will be part of the already identified components? Please clarify in the text that the PMF analysis results are based on the obtained filter analyses results.

Authors' response

We understand the reviewer's concern and thus add the following explanation in the text as follows (Page 22232, lines 22-27; Page 22233; lines 1-2):

“The reported PMF analysis is based on the chemical components of filter samples. As described in the mass closure, a large portion of the $PM_{2.5}$ mass fraction (about 45%) was not apportioned. In the PMF 5.0 procedure, the contributions of five factors were estimated and then the integrated contribution of the five factors was regressed over the measured $PM_{2.5}$ (HVS). The regression fit line was forced through the origin. Thus, our regression of the $PM_{2.5}$ (PMF) and $PM_{2.5}$ (HVS) showed that the $PM_{2.5}$ had been reproduced by PMF 5.0 with an error of less than 10% and the correlation of $PM_{2.5}$ (PMF) and $PM_{2.5}$ (HVS) showed a strong and significant correlation (slope = 0.91, $r^2 = 0.88$, $p < 0.01$) (Fig. 6b). To evaluate the results of the PMF model, the regression between predicted and observed data for each variable is shown during the operation. A linear correlation between the predicted and measured mass represents the goodness-of-fit of linear regression. Our values strongly suggested that the five identified sources could be readily interpreted.”

Comment to “slope of 0.91 regression result of PMF-HVS”

We acknowledge the reviewer's concern regarding the PMF-MLR results. However, we would like to inform that we did follow all the crucial and necessary steps in running the PMF tools as per mentioned in section 2.7. The mass contribution was calculated following procedure as per mentioned in Ke et al.

(2008) where first, regressing the factor scores obtained from the PMF using multi linear regression against the $PM_{2.5}$ measured mass concentration of each sample; and secondly, the regression coefficients were used to convert the contributions of each factor score into a physical mass.

- 2) Authors mention that the K^+ ion has been widely used in the literature as an excellent tracer representing wood burning, and authors examined the correlation between K^+ concentrations and MODIS fire counts. To my opinion, and correlation coefficient of 0.36 does not suggest that K^+ can be used as a biomass tracer. I would propose to differentiate between nss- K^+ and total K^+ and check if the correlation is stronger. ss- K^+ can be calculated using Na as a reference and the K/Na ratio in bulk seawater.

Authors' response

We agree with the reviewer's suggestions. We have revised Fig. 5, included Fig. 5b and 5c and annotated in the text as below (Page 22235; lines 19-26):




Figure 5 (a) Enrichment factor (EF) of heavy metals in PM_{2.5}, (b) correlation plot of nss-K⁺ and total-K⁺ ($r^2 = 0.95$), (c) correlation plot of K⁺ and Na⁺ and (d) reconstructed mass concentration of PM_{2.5} by mass closure (MC) model and (e) correlation plot of PM_{2.5} (μg m⁻³) HVS and PM_{2.5} (μg m⁻³) MC

“Further, a comparison of nss-K⁺ with the respective total K⁺ is shown in Fig. 5b. The correlation of nss-K⁺ as a function of total K⁺ showed a strong correlation coefficient ($r^2 = 0.95$) which suggests that K⁺ can be used as a biomass tracer. K⁺ may also be emitted from local fire sources. Additionally, the molar equivalent of K⁺ and Na⁺, as shown in Fig 5c, demonstrated significant correlation ($r^2 = 0.70$) with a slope value of 0.34 which is much higher compared to 0.0225-0.230 and 0.0218, reported by Wilson (1975) and Hara et al. (2012), respectively. The higher molar ratio of K⁺ and Na⁺ indicates that at the current location, Na⁺ depletion was high and the K⁺ might also release from other dominant sources. Additional significant sources of K⁺ which may attribute to the mass are soil dust, sea salt,

vegetation and meat cooking (Zhang et al., 2010).”

- 3) Based on the presented biomass fire hotspots, it is clear that from December 2013 to March 2014, the hotspots are a lot more compared to the respective ones from June to September 2013. Maybe authors should consider adding a comment of why this is the case. Is December-March the dry season? Is this period of intense biomass fire hotspots present every year or was the presented year uncharacteristic?

Authors' response

We have revised our text as the reviewer suggested (Page 22229; lines 1-3):

“The back trajectory plots showed that there were high numbers of biomass fire hotspots during both seasons but from different regions (Fig. 3). The period of June to September is the dry season each year in Malaysia and Sumatra of Indonesia. During this dry season, biomass fire hotspots are densely located in this area due to the burning of agricultural waste and forest fires. Several other researchers also reported the high number of biomass fire - related hotspots to these regions (Khan et al., 2015; Sahani et al., 2014). On the other hand, December to March is usually the wet season in Malaysia. However, the backward trajectories showed that air masses were transported from Mainland China and neighbouring regions. In Mainland China and neighbouring regions, this is dry season. During the dry season in this region there are a lot of fires, as reported by Zhang et al. (2015) and Ho et al. (2014), and this influences the pollution of air masses transported to the present location. This scenario of biomass fire hotspots is commonly noticed in these two seasons. In past years Malaysia and Singapore have experienced intensified haze episodes in this particular season, e.g. 1997, 2005, 2013 and 2015.”

- 4) In the enrichment factor section (2.5) in the whole paragraph EFs of two, five or even 10 are cited from the bibliography as thresholds for anthropogenic sources. Nevertheless authors choose a cut-off of EF=1. Please comment.

Authors' response

We understand and accept the reviewer's concern. Therefore, we have revised the cut-off point with explanation pertaining to that as follows (Page 22223; lines 11-24):

“Chester et al. (2000); Cheung et al. (2012); Khan et al. (2010); Mohd Tahir et al. (2013); Torfs and Van Grieken (1997) proposed a EF cut-off of ten to differentiate between crustal and natural and

anthropogenic origins of heavy metals. Thus, we consider $EF = 10$ as the cut-off point. Therefore, a good number of the metals (Zn, Cr, Rb, Be, V, Fe, Ca, Co, Sr, Pb, As and Bi) in $PM_{2.5}$ in this study can be assumed to originate from anthropogenic sources; Fig. 5a. These heavy metals were not natural or from the Earth's crust. No seasonal differences were observed in the EF of the heavy metals.”

Technical corrections:

P22217, L15-18: Not clear, especially the part “. . .with respect to inhalation follows the order. . .”. Also in PMF 5.0 delete the “>”.

Authors' response

We agree with the suggestion. We have revised the sentences as below (Page 22217; lines 14-17):

“The hazard quotient (HQ) for four selected metals (Pb, As, Cd and Ni) in $PM_{2.5}$ mass was highest in $PM_{2.5}$ mass from the coal burning source and least in $PM_{2.5}$ mass originating from the mineral/road dust source. The main carcinogenic heavy metal of concern to health at the current location was As; the other heavy metals (Ni, Pb, and Cd) did not pose a significant cancer risk in $PM_{2.5}$ mass concentration.”

P22219, L17: . . .a period from July to September

Authors' response

We have revised the sentence as below (Page 22219; lines 17-18):

Sampling was carried out on a 24 h basis for a period from July to September 2013 and January to February 2014 for a total of 27 samples.

P22220, L27: . . .ionic composition was determined

Authors' response

We have corrected the error (Page 22220, line 27):

“the water-soluble ionic composition (Na^+ , NH_4^+ , K^+ , Ca^{2+} , Mg^{2+} , Cl^- , NO_3^- and SO_4^{2-}) was determined”

P22221, L18: Replace “fresh” for the calibration curves. Maybe use “updated” instead.

Authors' response

We have revised the sentence as below (Page 22221, line 18):

“During the trace element analysis by ICP-MS, two modes of analysis were applied with updated calibration curves each time.”

P22223, L14: . . .metals in PM_{2.5} originated from. . . (delete “had”)

Authors' response

We have updated the sentence as below (Page 22223, line 14):

“Therefore, a good number of the metals (Zn, Cr, Rb, Be, V, Fe, Ca, Co, Sr, Pb, As and Bi) in PM_{2.5} in this study can be assumed to originate from anthropogenic sources; Fig. 5a”

P22227, L18: . . .classified by the International Agency. . .

Authors' response

We have corrected the sentence as below (Page 22227, line 18):

“Further, we examined the non-carcinogenic risk (presented by the hazard quotient (HQ)) and lifetime carcinogenic risk (LCR) of selected heavy metals as classified by the International Agency for Research on Cancer (IARC).”

P22230, L28: Boreddy et al. (2014) also notice a chlorine depletion due to atmospheric processing in the western north Pacific.

Authors' response

We have added the new reference suggested by the reviewer as below (Page 22230, line 28):

“Boreddy et al. (2014) also noticed a chlorine depletion due to atmospheric processing in the western north Pacific.”

P22233, L1: Based on Figure 6a the slope is 0.91 and R²=0.88 and not the other way round.

Authors' response

We have corrected the typo errors as below (Page 22233, Line 1):

“the correlation of $PM_{2.5}$ (PMF) and $PM_{2.5}$ (HVS) showed a strong and significant correlation (slope = 0.91, $r^2 = 0.88$, $p < 0.01$) (Fig. 6b)”

P22235, L8: Zhang et al. (2015) demonstrate that during the dry season there is an important biomass burning activity in the Pearl River Delta (China), which can result in trans-border transport and a regional scale character of biomass burning. Therefore under the north-east monsoonal regime it is possible that outflow from that area can maybe influence the specific area.

Authors' response

We are thankful to the reviewer suggesting the reference as below (Page 22235, line 8):

“However, the backward trajectories showed that air masses were transported from Mainland China and neighbouring regions. In Mainland China and neighbouring regions, this is dry season. During the dry season in this region there are a lot of fires, as reported by Zhang et al. (2015) and Ho et al. (2014), and this influences the pollution of air masses transported to the present location.”

P22235, L22: An $r=0.36$ does not suggest that K^+ can be used as a biomass tracer.

Authors' response

We have revised the text as the reviewer suggested. Please find the revised text as shown below (Page 22235, line 22):

“Further, a comparison of $nss-K^+$ with the respective total K^+ is shown in Fig. 5b. The correlation of $nss-K^+$ as a function of total K^+ showed a strong correlation coefficient ($r^2 = 0.95$) which suggests that K^+ can be used as a biomass tracer. K^+ may also be emitted from local fire sources. Additionally, the molar equivalent of K^+ and Na^+ , as shown in Fig 5c, demonstrated significant correlation ($r^2 = 0.70$) with a slope value of 0.34 which is much higher compared to 0.0225-0.230 and 0.0218, reported by Wilson (1975) and Hara et al. (2012), respectively. The higher molar ratio of K^+ and Na^+ indicates that at the current location, Na^+ depletion was high and the K^+ might also release from other dominant sources. Additional significant sources of K^+ which may attribute to the mass are soil dust, sea salt, vegetation and meat cooking (Zhang et al., 2010).”

P22238, L17: . . . follows the following order: $As > Ni$. . . Also “follows” and “following” is a repetition, maybe replace one of the two

Authors' response

We have corrected and re-worded the sentences as shown below (Page 22238, line 17):

“the lifetime carcinogenic risk level is estimated for PM_{2.5} mass concentration and may be contributed to by several heavy metals from different sources: mineral/road dust, motor vehicle emissions/biomass burning and coal combustion.”

L18: . . .PM_{2.5} concentration and As > Pb. . . (delete “;”)

Authors' response

We have re-worded the sentences as below (Page 22238, line 18):

“The main carcinogenic heavy metal of concern to the health of people at the current location is As; the other heavy metals (Ni, Pb, and Cd) did not pose a significant cancer risk.”

References

- Boreddy, S. K. R., Kawamura, K., and Jung, J.: Hygroscopic properties of particles nebulized from water extracts of aerosols collected at Chichijima Island in the western North Pacific: An outflow region of Asian dust, *Journal of Geophysical Research: Atmospheres*, 119, 2013JD020626, 10.1002/2013JD020626, 2014.
- Chester, R., Nimmo, M., Fones, G. R., Keyse, S., and Zhang, Z.: Trace metal chemistry of particulate aerosols from the UK mainland coastal rim of the NE Irish sea, *Atmos. Environ.*, 34, 949-958, [http://dx.doi.org/10.1016/S1352-2310\(99\)00234-4](http://dx.doi.org/10.1016/S1352-2310(99)00234-4), 2000.
- Cheung, K., Shafer, M. M., Schauer, J. J., and Sioutas, C.: Historical trends in the mass and chemical species concentrations of coarse particulate matter in the Los Angeles Basin and relation to sources and air quality regulations, *J. Air Waste Manage. Assoc.*, 62, 541-556, 10.1080/10962247.2012.661382, 2012.
- Hara, K., Osada, K., Yabuki, M., and Yamanouchi, T.: Seasonal variation of fractionated sea-salt particles on the Antarctic coast, *Geophysical Research Letters*, 39, L18801, 10.1029/2012GL052761, 2012.
- Ho, K. F., Engling, G., Ho, S. S. H., Huang, R., Lai, S., Cao, J., and Lee, S. C.: Seasonal variations of anhydrosugars in PM_{2.5} in the Pearl River Delta Region, China, 2014, 10.3402/tellusb.v66.22577, 2014.
- Ke, L., Liu, W., Wang, Y., Russell, A. G., Edgerton, E. S., and Zheng, M.: Comparison of PM_{2.5} source apportionment using positive matrix factorization and molecular marker-based chemical mass balance, *Sci. Total. Environ.*, 394, 290-302, <http://dx.doi.org/10.1016/j.scitotenv.2008.01.030>, 2008.
- Khan, M. F., Shirasuna, Y., Hirano, K., and Masunaga, S.: Urban and suburban aerosol in Yokohama, Japan: a comprehensive chemical characterization, *Environ. Monit. Assess.*, 171, 441-456, 10.1007/s10661-009-1290-1, 2010.
- Khan, M. F., Latif, M. T., Lim, C. H., Amil, N., Jaafar, S. A., Dominick, D., Nadzir, M. S. M., Sahani, M., and Tahir, N. M.: Seasonal effect and source apportionment of polycyclic aromatic hydrocarbons in PM_{2.5}, *Atmos. Environ.*, 106, 178-190, 10.1016/j.atmosenv.2015.01.077, 2015.

- Mohd Tahir, N., Suratman, S., Fong, F. T., Hamzah, M. S., and Latif, M. T.: Temporal distribution and chemical characterization of atmospheric particulate matter in the eastern coast of Peninsular Malaysia, *Aerosol Air Qual. Res.*, 13, 584-595, 10.4209/aaqr.2012.08.0216, 2013.
- Sahani, M., Zainon, N. A., Mahiyuddin, W. R. W., Latif, M. T., Hod, R., Khan, M. F., Tahir, N. M., and Chan, C.-C.: A case-crossover analysis of forest fire haze events and mortality in Malaysia, *Atmos. Environ.*, 96, 257-265, 10.1016/j.atmosenv.2014.07.043, 2014.
- Torfs, K., and Van Grieken, R.: Chemical relations between atmospheric aerosols, deposition and stone decay layers on historic buildings at the mediterranean coast, *Atmos. Environ.*, 31, 2179-2192, [http://dx.doi.org/10.1016/S1352-2310\(97\)00038-1](http://dx.doi.org/10.1016/S1352-2310(97)00038-1), 1997.
- Wilson, T. R.: Salinity and the major elements of sea-water, in *Chemical Oceanography*, edited by J. P. Riley and G. Skirrow, pp. 365–413, Academic, San Diego, Calif., 1975.
- Zhang, X., Hecobian, A., Zheng, M., Frank, N. H., and Weber, R. J.: Biomass burning impact on PM 2.5 over the southeastern US during 2007: integrating chemically speciated FRM filter measurements, MODIS fire counts and PMF analysis, *Atmos. Chem. Phys.*, 10, 6839-6853, 10.5194/acp-10-6839-2010, 2010.
- Zhang, Z., Gao, J., Engling, G., Tao, J., Chai, F., Zhang, L., Zhang, R., Sang, X., Chan, C.-y., Lin, Z., and Cao, J.: Characteristics and applications of size-segregated biomass burning tracers in China's Pearl River Delta region, *Atmos. Environ.*, 102, 290-301, <http://dx.doi.org/10.1016/j.atmosenv.2014.12.009>, 2015.

Responses to the comments of referee#3

Interactive comment on “Fine particulate matter associated with monsoonal effect and the responses of biomass fire hotspots in the tropical environment” by M. F. Khan et al.

Anonymous Referee #3

Received and published: 13 September 2015

1. The title is not appropriate. Title should include the terms “source apportionment” and also “health effect”. As, authors one of the main conclusion is on the health impact. Overall, the associated cancer risk posed by the exposure of toxic metals in PM_{2.5} is three to four in 1 000 000 people in this location.

Authors’ response

We do agree with the review’s suggestion and have revised the title as shown below:

“Fine particulate matter in the tropical environment: monsoonal effects, source apportionment and health risk assessment”

2. Authors used quartz microfiber filters for trace metals analyzer. Quartz microfiber filters are not the great filters for trace metals. They may have some influences on the trace metals content. Though they did blank correction for the filters.

Authors’ response

We agree with the reviewer's observation. Despite possible influence of trace metal content, several researchers still use the microfibre quartz filter but with strict QA/QC and blank corrected to account for possible interferences. Taking into account the reviewer's comment, we have included in the text several references to corroborate our choice of filter. The QA/QC aspect has been mentioned in the text (page 22221, line 8)

3. Authors measured only water-soluble inorganic ions and trace metals in $PM_{2.5}$. But the main component of $PM_{2.5}$ is organic (OC) and black carbon (BC or EC) and also water-soluble organic ions are missing. Trace metals may be up to 1% and water-soluble inorganic ions may be up to 20% but carbonaceous species may be up to 50% of the fine particles are missing in this chemical composition.

Authors' response

We acknowledge reviewer's concern and we will definitely take this into consideration in future studies. At this stage, we added an explanation highlighted in turquoise color as follows (page 22232, line 3):

“The reported result of the mass closure model is based on the analyzed chemical components of filter samples (~ 55%). As described in the mass closure, a large portion of $PM_{2.5}$ mass (~ 45%) was left unidentified; this unidentified component is believed to be the organics or carbonaceous species. Elemental carbon (EC), organic carbon (OC) and water-soluble organics were not measured due to the lack of instrumentation. Other possible reasons for the un-identified portion are: i) unaccounted for mineral oxides as they are abundant in $PM_{2.5}$, and ii) water associated with salts.”

4. Author cut off point for EF is 1. Whereas Sun et al. (2006) suggested the threshold of $EF > 5$ to differentiate between sources from the Earth's crust and from anthropogenic sources. In contrast, Mohd Tahir et al. (2013) proposed the EF cut-off of ten to identify crustal and natural origin of heavy metals. What is the basis of their cut off point???

How did thy calculate EF? I mean which for formula please refer that not my reference like Taylor (1964) but with formula or equation.

Authors' response

We understand and accept the reviewer's concern. We also made similar answer to the comments of previous reviewer#2. We acknowledge that the use of EF is subjected to variation depending on the reference crustal values. A larger margin is more often used for atmospheric samples (EF >10) to account for this uncertainty and also possible regional variation and the conservation of crustal element ratios through various environmental compartments. Therefore, we have revised the cut-off point to 10 with explanation pertaining to that highlighted in turquoise color as follows (Page 22223; lines 11-24):

“Chester et al. (2000); Cheung et al. (2012); Khan et al. (2010); Mohd Tahir et al. (2013); Torfs and Van Grieken (1997) proposed a EF cut-off of ten to differentiate between crustal and natural and anthropogenic origins of heavy metals. Thus, we consider EF = 10 as the cut-off point. Therefore, a good number of the metals (Zn, Cr, Rb, Be, V, Fe, Ca, Co, Sr, Pb, As and Bi) in PM_{2.5} in this study can be assumed to originate from anthropogenic sources; Fig. 5a. These heavy metals were not natural or from the Earth's crust. No seasonal differences were observed in the EF of the heavy metals.”

The manuscript has been revised with following equation of EF (page 22223, line 9):

“The EF of each element can be defined using the following equation:

$$EF = \frac{\left(\frac{E}{Al}\right)_{PM\ 2.5}}{\left(\frac{E}{Al}\right)_{Crust}} \quad (1)$$

where, E/Al is the concentration ratio of element, E, to the reference metal, Al.”

5. Figure caption is inadequate. e.g., Figure 3 should have the source (e.g., HYSPLIT,), backward or forward, starting and ending time etc. Though authors mentioned them in the text. They should also write

in the figure caption.

Authors' response

Revised caption of Figure 3 as shown below:

“Figure 3 The location of biomass fire hotspots and the monthly mean cluster of backward trajectories by HYSPLIT 4.9 model for 120 h and 500 m releasing height starting from 16:00 UTC during the south-westerly and the north-easterly monsoon”

6. Authors are using $PM_{2.5}$ MC with equation 4. It would be great if they can discuss $PM_{2.5}$ MC and $PM_{2.5}$ mass.

Authors' response

We agree with the reviewer's suggestion. We have added an explanation and a correlation plot of $PM_{2.5}$ (HVS) and $PM_{2.5}$ (MC) highlighted in turquoise color as bellows (page 22232, line 3):

“The average value of reconstructed $PM_{2.5}$ by mass closure (MC) is $14.12 \pm 4.32 \mu\text{g m}^{-3}$ with a minimum of $6.70 \mu\text{g m}^{-3}$ and a maximum of $24.19 \mu\text{g m}^{-3}$. On the other hand, the measured $PM_{2.5}$ determined gravimetrically by HVS was $25.13 \pm 9.21 \mu\text{g m}^{-3}$ with a range of 7.01 to $42.78 \mu\text{g m}^{-3}$. A correlation plot of $PM_{2.5}$ (MC) and measured $PM_{2.5}$ (HVS) is shown in Fig. 5e. The correlation shows a good fit ($r^2 = 0.98$) with a slope of 0.46 and an intercept of 1.93. The results of the fit parameters suggest that the $PM_{2.5}$ mass (MC) concentration was underestimated compared to $PM_{2.5}$ (HVS). The reported result of the mass closure model is based on the analyzed chemical components of filter samples (~ 55%). As described in the mass closure, a large portion of $PM_{2.5}$ mass (~ 45%) was left unidentified; this unidentified component is believed to be the organics or carbonaceous species. Elemental carbon (EC), organic carbon (OC) and water-soluble organics were not measured due to the lack of instrumentation. Other possible reasons for the un-identified portion are: i) unaccounted for mineral oxides as they are abundant in $PM_{2.5}$, and ii) water associated with salts.”




Figure 5 (a) Enrichment factor (EF) of heavy metals in PM_{2.5}, (b) correlation plot of nss-K⁺ and total-K⁺, (c) reconstructed mass concentration of PM_{2.5} by mass closure model, (d) correlation plot of K⁺ and Na⁺, and (e) correlation plot of estimated PM_{2.5} (MC) and measured PM_{2.5} (HVS)

7. It is kind of weird to discuss the carcinogenicity on the basis on of sources. E.g., if a metal Ni came from coal may be more carcinogen than if it is came from traffic. Please try to rewording then section Table 3 and discussion 3.4.

Authors' response

We agreed with the reviewer's comment; we did not mean to say that metals in one source are more carcinogenic than the same metals in other sources. We apologies for the confusion and we have reworded our discussion to reflect that carcinogenic effects are due to metals present in PM_{2.5} that are associated with that particular source (pages 22237-22239):

“Table 3. Hazard quotient (HQ) and lifetime carcinogenic risk (LCR) for selected heavy metals in PM_{2.5} associated with sources”

Revised Table 3

Table 3. Hazard quotient (HQ) and lifetime carcinogenic risk (LCR) for selected heavy metals in PM_{2.5} associated with sources

Inhalation	Mineral/road dust		Motor vehicle/biomass		Coal burning		PM _{2.5}	
	HQ	LCR	HQ	LCR	HQ	LCR	HQ	LCR
Pb	-	4.0×10 ⁻⁸	-	1.0×10 ⁻⁷	-	4.1×10 ⁻⁸	-	2.5×10 ⁻⁷
As	1.8×10 ⁻³	1.1×10 ⁻⁷	4.9×10 ⁻²	1.1×10 ⁻⁶	6.6×10 ⁻²	1.5×10 ⁻⁶	15.9×10 ⁻²	3.5×10 ⁻⁶
Cd	4.6×10 ⁻⁴	8.2×10 ⁻⁹	9.1×10 ⁻³	5.6×10 ⁻⁸	5.5×10 ⁻³	3.4×10 ⁻⁸	2.2×10 ⁻²	1.4×10 ⁻⁷
Cu	-	-	-	-	-	-	-	-
Mn	-	-	7.0×10 ⁻³	-	5.1×10 ⁻³	-	3.3×10 ⁻²	-
Zn	-	-	-	-	-	-	-	-
Ni	2.0×10 ⁻²	2.5×10 ⁻⁷	2.3×10 ⁻²	9.5×10 ⁻⁸	2.6×10 ⁻²	1.1×10 ⁻⁷	14.3×10 ⁻²	5.9×10 ⁻⁷
THR (HI and LCR)	2.3×10 ⁻²	1.6×10 ⁻⁷	8.8×10 ⁻²	1.2×10 ⁻⁶	10.2×10 ⁻²	1.5×10 ⁻⁶	35.7×10 ⁻²	3.9×10 ⁻⁶

Pb*: pb (acetate), As*: As (Inorganic), Cd*: Cd (Diet), Mn*: Mn (Diet), Zn*: Zn (Metallic), Ni*: Ni (Refinery Dust), "-": no data, **THR: Total Health Risk, HI: Hazard Index, HQ: Hazard quotient (non-cancer risk); LCR: Lifetime Cancer Risk**

1 “Table 3 shows the non-carcinogenic (represented as HQ) and carcinogenic risks posed by several
2 selected metals (Pb, As, Cd, Cu, Mn, Zn and Ni) in PM_{2.5} mass concentration through inhalation
3 exposure associating with sources. The HQ values for As and Ni in PM_{2.5} mass concentration are
4 15.9×10^{-2} and 14.3×10^{-2} , respectively, suggesting the non-carcinogenic health risks posed by these
5 metals might be higher compared to other metals. The HQ for four selected metals (Pb, As, Cd and
6 Ni) in PM_{2.5} mass was highest in the PM_{2.5} mass originating from the coal burning source and least in
7 PM_{2.5} originating from the mineral/road dust source. The cut-off point for significant health risks or
8 the safe level to the exposed population is HQ > 1. Our results showed that the sum of HQ for each
9 metal are lower than the safe level (= 1) in PM_{2.5} mass concentration originating from each source.
10 The sum of HQ for PM_{2.5} is 35.7×10^{-2} , which is lower than the HQs of PM_{2.5} reported by Hu et al.
11 (2012) in Nanjing, China (2.96); Cao et al. (2014) in Shanxi Province, China ($1.06 \times 10^{+1}$); and Taner
12 et al. (2013) in a non-smoking restaurant in Turkey (4.09). A study by Hu et al. (2012), reported HQ
13 values for As and Ni in PM_{2.5} as 4.14×10^{-1} and 1.73×10^{-1} , respectively, in Nanjing, China. However,
14 the HQs of PM_{2.5} estimated after inhalation at two sites in Nanjing City, China (0.88 (Xianlin) and
15 0.79 (Gulou)) were close to the safe level (= 1) according to a study by Li et al. (2015). At two urban
16 locations in Yangtze River Delta, China, the HQ for Cr in PM_{2.5} was within the acceptable limit but
17 higher for Mn (Niu et al., 2015). Although the HQ calculated for As was the highest, it was below 1,
18 thus the non-carcinogenic health risk was estimated to be at a safe level. In addition, the hazard index
19 (total - hazard quotient) of PM_{2.5} calculated for the four heavy metals (As, Cd, Mn, Ni) from the
20 different sources (Table 3) showed an insignificant health risk.

21 The carcinogenic risks from the carcinogenic heavy metals Pb, As, Cd and Ni in PM_{2.5} are shown
22 in Table 3. Similar to the non-carcinogenic risks, the lifetime carcinogenic risk level is estimated for
23 PM_{2.5} mass concentration and may be contributed to by several heavy metals from different sources:
24 mineral/road dust, motor vehicle emissions/biomass burning and coal combustion. The total life-time
25 cancer risk (LCR) from heavy metals in the PM_{2.5} mass concentration was calculated at 3.9×10^{-6}
26 which is a significant cancer risk. The main carcinogenic heavy metal of concern to the health of
27 people at the current location is As; the other heavy metals (Ni, Pb, and Cd) did not pose a significant
28 cancer risk. Thus, the LCR from the PM_{2.5} mass concentration originating from motor
29 vehicle/biomass and coal burning sources showed a value of 1×10^{-6} , slightly above the acceptable
30 cancer risk level as recommended by USEPA, while the total LCR from PM_{2.5} mass concentration
31 from all sources was estimated to be 4×10^{-6} which is also slightly above the acceptable cancer risk.
32 The carcinogenic risk posed by As (3.66×10^{-3}) in PM_{2.5} in Shanxi Province, China (Cao et al., 2014)
33 was higher than the guideline value set by USEPA. A study by Niu et al. (2015) of PM_{2.5}-bound
34 metals showed a high cancer risk in Yangtze River Delta, China (2.47×10^{-4}). A study by Pandey et al.
35 (2013) conducted in the vicinity of human activities observed that the concentrations of Cd, Cr, Ni
36 and Pb in PM_{2.5} showed higher excess cancer risk (ECR) due to those particle-bound metals compared

37 to guideline level set by USEPA. Satsangi et al. (2014) also reported a higher cancer risk from Cr, Ni
38 and Cd in PM_{2.5} compared to the USEPA guideline. The integrated carcinogenic risk of six metals
39 (Cr, As, Co, Pb, Ni and Cd) in PM_{2.5} in Tianjin, China were in the range $3.4 \times 10^{-3} - 4.1 \times 10^{-3}$ which is
40 reportedly beyond the tolerance level (Zhang et al., 2014). The total ECRs based on the average
41 values of As, Cd, Cr, Ni and Pb in PM_{2.5} is 4.34×10^{-5} in Delhi, India, implying that four or five people
42 might get cancer out of 100,000 people after exposure to toxic metals in PM_{2.5} (Khanna et al., 2015).
43 Our findings showed that the life-time cancer risk posed by the exposure of heavy metals in PM_{2.5}
44 mass concentration is three to four per 1,000,000 people at this location. This significant cancer risk
45 warrants further investigation. Our findings showed that an insignificant non-carcinogenic risk and
46 significant cancer risk is posed to the population from exposure to PM_{2.5} at this location. Detailed
47 exposure assessment of the PM_{2.5} at the specific sources and the health risks posed by individual
48 hazardous elements of concern may help to improve understanding about the exposure pathways as
49 well as the detailed risk factors involved in both carcinogenic and non-carcinogenic risk.”

50

51 8. Overall: Abstract is not well written. Conclusion is okay. Authors spent more times on
52 methodology not on the results and discussions.

53 **Author's responses**

54 We appreciate reviewer's observation. We have already modified our abstract. We have revised the
55 Health risk part in the “Results and Discussion”. Based on the comments from reviewer#2, we also
56 have enriched our “Results and Discussion” in the revised manuscript file highlighted in turquoise
57 color. The revised version of the abstract is shown below (page 22217, lines 2-20):

58

59 “The health implications of PM_{2.5} in the tropical region of Southeast Asia (SEA) are significant as
60 PM_{2.5} can pose serious health concerns. PM_{2.5} concentration and sources here are strongly influenced
61 by changes in the monsoon regime from the southwest quadrant to the northeast quadrant in the
62 region. In this work, PM_{2.5} samples were collected at a semi-urban area using a high volume air
63 sampler at different seasons on 24 h basis. Analysis of trace elements and water-soluble ions was
64 performed using inductively coupled plasma mass spectroscopy (ICP-MS) and ion chromatography
65 (IC), respectively. Apportionment analysis of PM_{2.5} was carried out using the United States
66 Environmental Protection Agency (US EPA) positive matrix factorization (PMF) 5.0 and a mass
67 closure model. We quantitatively characterized the health risks posed to human populations through
68 the inhalation of selected heavy metals in PM_{2.5}. 48% of the samples collected exceeded the World
69 Health Organization (WHO) 24 h PM_{2.5} guideline but only 19% of the samples exceeded 24 h US

70 EPA National Ambient Air Quality Standard (NAAQS). The PM_{2.5} concentration was slightly higher
71 during the north-east monsoon compared to south-west monsoon. The main trace metals identified
72 were As, Pb, Cd, Ni, Mn, V and Cr while the main ions were SO₄²⁻, NO₃⁻, NH₄⁺ and Na. The mass
73 closure model identified four major sources of PM_{2.5} that accounts for 55% of total mass balance. The
74 four sources are mineral matter (MIN) (35%), secondary inorganic aerosol (SIA) (11%), sea salt (SS)
75 (7%), and trace elements (TE) (2%). PMF 5.0 elucidated five potential sources: motor vehicle
76 emissions coupled with biomass burning (31%) were the most dominant, followed by marine/sulfate
77 aerosol (20%), coal burning (19%), nitrate aerosol (17%), and mineral/road dust (13%). The hazard
78 quotient (HQ) for four selected metals (Pb, As, Cd and Ni) in PM_{2.5} mass was highest in PM_{2.5} mass
79 from the coal burning source and least in PM_{2.5} mass originating from the mineral/road dust source.
80 The main carcinogenic heavy metal of concern to health at the current location was As; the other
81 heavy metals (Ni, Pb, and Cd) did not pose a significant cancer risk in PM_{2.5} mass concentration.
82 Overall, the associated lifetime cancer risk posed by the exposure of hazardous metals in PM_{2.5} is
83 three to four in 1,000,000 people at this location.”

84

85

86 **Reference:**

87

88 Cao, S., Duan, X., Zhao, X., Ma, J., Dong, T., Huang, N., Sun, C., He, B., and Wei, F.: Health risks
89 from the exposure of children to As, Se, Pb and other heavy metals near the largest coking
90 plant in China, *Sci. Total. Environ.*, 472, 1001-1009,
91 <http://dx.doi.org/10.1016/j.scitotenv.2013.11.124>, 2014.

92 Chester, R., Nimmo, M., Fones, G. R., Keyse, S., and Zhang, Z.: Trace metal chemistry of particulate
93 aerosols from the UK mainland coastal rim of the NE Irish sea, *Atmos. Environ.*, 34, 949-
94 958, [http://dx.doi.org/10.1016/S1352-2310\(99\)00234-4](http://dx.doi.org/10.1016/S1352-2310(99)00234-4), 2000.

95 Cheung, K., Shafer, M. M., Schauer, J. J., and Sioutas, C.: Historical trends in the mass and chemical
96 species concentrations of coarse particulate matter in the Los Angeles Basin and relation to
97 sources and air quality regulations, *J. Air Waste Manage. Assoc.*, 62, 541-556,
98 [10.1080/10962247.2012.661382](http://dx.doi.org/10.1080/10962247.2012.661382), 2012.

99 Hu, X., Zhang, Y., Ding, Z., Wang, T., Lian, H., Sun, Y., and Wu, J.: Bioaccessibility and health risk
100 of arsenic and heavy metals (Cd, Co, Cr, Cu, Ni, Pb, Zn and Mn) in TSP and PM_{2.5} in

101 Nanjing, China, Atmos. Environ., 57, 146-152,
102 <http://dx.doi.org/10.1016/j.atmosenv.2012.04.056>, 2012.

103 Khan, M. F., Shirasuna, Y., Hirano, K., and Masunaga, S.: Urban and suburban aerosol in Yokohama,
104 Japan: a comprehensive chemical characterization, Environ. Monit. Assess., 171, 441-456,
105 10.1007/s10661-009-1290-1, 2010.

106 Khanna, I., Khare, M., and Gargava, P.: Health Risks Associated with Heavy Metals in Fine
107 Particulate Matter: A Case Study in Delhi City, India, J. Geosci. Environ. Protec., 3, 72,
108 10.4236/gep.2015.32012, 2015.

109 Li, Y., Zhang, Z., Liu, H., Zhou, H., Fan, Z., Lin, M., Wu, D., and Xia, B.: Characteristics, sources
110 and health risk assessment of toxic heavy metals in PM_{2.5} at a megacity of southwest China,
111 Environ. Geochem. Health, 1-10, 10.1007/s10653-015-9722-z, 2015.

112 Mohd Tahir, N., Suratman, S., Fong, F. T., Hamzah, M. S., and Latif, M. T.: Temporal distribution
113 and chemical characterization of atmospheric particulate matter in the eastern coast of
114 Peninsular Malaysia, Aerosol Air Qual. Res., 13, 584-595, 10.4209/aaqr.2012.08.0216, 2013.

115 Niu, L., Ye, H., Xu, C., Yao, Y., and Liu, W.: Highly time- and size-resolved fingerprint analysis and
116 risk assessment of airborne elements in a megacity in the Yangtze River Delta, China,
117 Chemosphere, 119, 112-121, <http://dx.doi.org/10.1016/j.chemosphere.2014.05.062>, 2015.

118 Pandey, P., Patel, D. K., Khan, A. H., Barman, S. C., Murthy, R. C., and Kisku, G. C.: Temporal
119 distribution of fine particulates (PM_{2.5}, PM₁₀), potentially toxic metals, PAHs and Metal-
120 bound carcinogenic risk in the population of Lucknow City, India, J. Environ. Sci. Health.,
121 Part A, 48, 730-745, 10.1080/10934529.2013.744613, 2013.

122 Satsangi, P. G., Yadav, S., Pipal, A. S., and Kumbhar, N.: Characteristics of trace metals in fine
123 (PM_{2.5}) and inhalable (PM₁₀) particles and its health risk assessment along with in-silico
124 approach in indoor environment of India, Atmos. Environ., 92, 384-393,
125 <http://dx.doi.org/10.1016/j.atmosenv.2014.04.047>, 2014.

126 Taner, S., Pekey, B., and Pekey, H.: Fine particulate matter in the indoor air of barbeque restaurants:
127 Elemental compositions, sources and health risks, Sci. Total. Environ., 454-455, 79-87,
128 <http://dx.doi.org/10.1016/j.scitotenv.2013.03.018>, 2013.

129 Torfs, K., and Van Grieken, R.: Chemical relations between atmospheric aerosols, deposition and
130 stone decay layers on historic buildings at the mediterranean coast, Atmos. Environ., 31,
131 2179-2192, [http://dx.doi.org/10.1016/S1352-2310\(97\)00038-1](http://dx.doi.org/10.1016/S1352-2310(97)00038-1), 1997.

132 Zhang, N., Han, B., He, F., Xu, J., Niu, C., Zhou, J., Kong, S., Bai, Z., and Xu, H.: Characterization,
133 health risk of heavy metals, and source apportionment of atmospheric PM2.5 to children in
134 summer and winter: an exposure panel study in Tianjin, China, *Air Qual. Atmos. Health*, 1-
135 11, 10.1007/s11869-014-0289-0, 2014.

136

137

138 **Fine particulate matter in the tropical environment: monsoonal**
139 **effects, source apportionment and health risk assessment**

140

141 **Md Firoz Khan^{1,2,*}, Mohd Talib Latif^{1,3}, Wuan Hwa Saw¹, Norhaniza Amil^{1,4},**
142 **Mohd Shahrul Mohd Nadzir^{1,2}, Mazrura Sahani⁵, Norhayati Mohd Tahir^{6,7}, Jing**
143 **Xiang Chung¹**

144

145 [1]{School of Environmental and Natural Resource Sciences, Faculty of Science and
146 Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia}

147

148 [2]{Centre for Tropical Climate Change System (IKLIM), Institute for Climate Change,
149 Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia}

150

151 [3]{Institute for Environment and Development (LESTARI), Universiti Kebangsaan
152 Malaysia, 43600 Bangi, Selangor, Malaysia}

153

154 [4]{School of Industrial Technology (Environmental Division), Universiti Sains Malaysia,
155 11800 Penang, Malaysia}

156 [5]{Environmental Health and Industrial Safety Program, School of Diagnostic Science and
157 Applied Health, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Jalan Raja
158 Muda Abdul Aziz, 50300 Kuala Lumpur, Malaysia}

159

160 [6]{Environmental Research Group, School of Marine Science and Environment, Universiti
161 Malaysia Terengganu, 21030 Kuala Terengganu, Terengganu, Malaysia}

162

163 [7]{Institute of Oceanography, Universiti Malaysia Terengganu, 21030 Kuala Terengganu,
164 Terengganu, Malaysia}

165

166 *Corresponding author

167 Tel: +603-89213822, fax: +603-89253357

168 Email: mdfiroz.khan@ukm.edu.my; mdfiroz.khan@gmail.com (M. F. Khan)

169

Abstract

170 The health implications of PM_{2.5} in the tropical region of Southeast Asia (SEA) are
171 significant as PM_{2.5} can pose serious health concerns. PM_{2.5} concentration and sources here
172 are strongly influenced by changes in the monsoon regime from the southwest quadrant to the
173 northeast quadrant in the region. In this work, PM_{2.5} samples were collected at a semi-urban
174 area using a high volume air sampler at different seasons on 24 h basis. Analysis of trace
175 elements and water-soluble ions was performed using inductively coupled plasma mass
176 spectroscopy (ICP-MS) and ion chromatography (IC), respectively. Apportionment analysis
177 of PM_{2.5} was carried out using the United States Environmental Protection Agency (US EPA)
178 positive matrix factorization (PMF) 5.0 and a mass closure model. We quantitatively
179 characterized the health risks posed to human populations through the inhalation of selected
180 heavy metals in PM_{2.5}. 48% of the samples collected exceeded the World Health
181 Organization (WHO) 24 h PM_{2.5} guideline but only 19% of the samples exceeded 24 h US
182 EPA National Ambient Air Quality Standard (NAAQS). The PM_{2.5} concentration was
183 slightly higher during the north-east monsoon compared to south-west monsoon. The main
184 trace metals identified were As, Pb, Cd, Ni, Mn, V and Cr while the main ions were SO₄²⁻,
185 NO₃⁻, NH₄⁺ and Na. The mass closure model identified four major sources of PM_{2.5} that
186 accounts for 55% of total mass balance. The four sources are mineral matter (MIN) (35%),
187 secondary inorganic aerosol (SIA) (11%), sea salt (SS) (7%), and trace elements (TE) (2%).
188 PMF 5.0 elucidated five potential sources: motor vehicle emissions coupled with biomass
189 burning (31%) were the most dominant, followed by marine/sulfate aerosol (20%), coal
190 burning (19%), nitrate aerosol (17%), and mineral/road dust (13%). The hazard quotient
191 (HQ) for four selected metals (Pb, As, Cd and Ni) in PM_{2.5} mass was highest in PM_{2.5} mass
192 from the coal burning source and least in PM_{2.5} mass originating from the mineral/road dust
193 source. The main carcinogenic heavy metal of concern to health at the current location was
194 As; the other heavy metals (Ni, Pb, and Cd) did not pose a significant cancer risk in PM_{2.5}
195 mass concentration. Overall, the associated lifetime cancer risk posed by the exposure of
196 hazardous metals in PM_{2.5} is three to four per 1,000,000 people at this location.

197

198 **1 Introduction**

199 Atmospheric fine particles ($PM_{2.5}$, $dP \leq 2.5 \mu m$), a mixture of many inorganic and organic
200 components, reside for a long time in the atmosphere and can penetrate deep into the lung.
201 Prolonged exposure to $PM_{2.5}$ can cause adverse health impacts and premature mortality in
202 humans (Betha et al., 2014). Potential health benefits and an improvement in general
203 mortality could be expected if the control policies were implemented (Boldo et al., 2011).
204 The adverse effects of $PM_{2.5}$ can reach intercontinental scales (Anenberg et al., 2014) due to
205 the potential transport of $PM_{2.5}$ over hundreds to thousands of kilometers (Seinfeld and
206 Pandis, 2012). The sources of $PM_{2.5}$, particularly motor vehicle emissions, are associated
207 with an increase in hospital admissions (Kioumourtzoglou et al., 2014). A study by Bell et al.
208 (2014) suggested that controlling some of the sources of $PM_{2.5}$ could protect public health
209 more efficiently than the regulation of particle concentration. Thus, the possible reduction in
210 health risks from the predominant sources of $PM_{2.5}$ is desired as part of the mitigation
211 strategy. Diesel emissions and biomass burning, as the primary risk sources of $PM_{2.5}$, should
212 be closely monitored and regulated (Wu et al., 2009).

213 The identification of $PM_{2.5}$ sources is becoming a widely-recognized way to protect human
214 health as well as the environment. Multivariate receptor models are very useful in the source
215 apportionment of $PM_{2.5}$. Widely used multivariate methods are: a) a chemical mass balance
216 model (CMB) (Watson et al., 1990), b) positive matrix factorization (PMF) (Paatero, 1997;
217 Paatero and Tapper, 1994), c) Unmix (Henry, 1987), d) principal component analysis coupled
218 with absolute principal component score (PCA/APCS) (Thurston and Spengler, 1985), e)
219 pragmatic mass closure (PMC) (Harrison et al., 2003) and f) a new source-type identification
220 method for $PM_{2.5}$ known as Reduction and Species Clustering Using Episodes (ReSCUE)
221 (Vedantham et al., 2014). PMF is the most reliable method for source-type identification for
222 the following reasons: i) it uses a weighted least-squares fit and estimates error of the
223 measured data and can impose non-negativity constraints weighing each data point
224 individually (Paatero, 1997; Paatero and Tapper, 1994), ii) *a priori* knowledge of pollutants
225 is not necessary and iii) it is able to deal with missing values, noisy data, outliers, and values
226 below detection limit (Baumann et al., 2008; Khan et al., 2012; Khan et al., 2015b; Polissar et
227 al., 1998a; Polissar et al., 1998b). A recent study by Gibson et al. (2014) suggested that PMF
228 can resolve $PM_{2.5}$ concentrations even below $2 \mu g m^{-3}$ more accurately compared to PMC and
229 CMB.

230 Source apportionment studies of $PM_{2.5}$ based on monsoonal seasonal changes in Malaysia are
231 of widespread interest due to the influence of local sources as well as trans-boundary haze
232 pollution. This haze pollution reaches its worst level during the south-west (SW) and north-
233 east (NE) monsoons each year. Therefore, the main objectives of this work are to investigate
234 a) the monsoonal effect on the variability of $PM_{2.5}$ and its chemical composition, b) factors
235 influencing the sources of $PM_{2.5}$ and c) to quantitatively characterize the non-carcinogenic
236 and carcinogenic risks to the potentially-exposed human populations by selected heavy
237 metals in $PM_{2.5}$ released from the particular sources. The $PM_{2.5}$ mass concentration
238 contributed by each source will be calculated using PMF 5.0.

239

240 **2 Methodologies**

241 **2.1 Description of the study area**

242 Figure 1 shows the sampling location which is on the roof top of the Biology Building of the
243 Faculty of Science and Technology (FST), University Kebangsaan Malaysia (UKM),
244 Malaysia (2° 55' 31.91" N, 101° 46' 55.59" E, about 65 m above sea level). This site is less
245 than 1 km from the main Bangi road.

246 **2.2 Sampling and analysis of $PM_{2.5}$ samples**

247 Sampling was carried out on a 24 h basis for a period from July to September 2013 and
248 January to February 2014 for a total of 27 samples. The $PM_{2.5}$ samples were collected on
249 quartz microfiber filters (203 mm×254 mm, WhatmanTM, UK) through a $PM_{2.5}$ high volume
250 sampler (HVS, Tisch, USA) at a flow rate of 1.13 m³ min⁻¹. Several others researchers also
251 conducted sampling using the quartz microfiber filters for the analysis of trace metals (Li et
252 al., 2015a; Martins et al., 2016; Kholdebarin et al., 2015; Cusack et al., 2015; Sánchez-
253 Soberón et al., 2015). Prior to use, the filters were preheated at 500 °C for 3 h to remove any
254 deposited organic compounds. All filter papers either blank or exposed were conditioned in a
255 desiccator for 24 h before and after sampling prior to weighing using a 5-digit high-resolution
256 electronic balance (A&D, GR-202, Japan) with a 0.01 mg detection limit. The filter samples
257 were then stored at -18 °C until the extraction procedure. A microwave-assisted digestion
258 system (Start D, Milestone, Germany) was employed for the preparation of the trace element
259 samples. The microwave was operated at two temperature stages, 180 °C for 20 min and
260 ramping to 220 °C for 15 min. The power was set at 500 watts during the procedure when the

261 number of samples \leq three. However, the power was set at 1000 watts if the number of
262 samples exceeded three. A 4:1 ratio of 12 mL nitric acid (65%, Merck KGaA, Germany) and 3
263 mL hydrogen peroxide (40%, Merck KGaA, Germany) was used as the reagent in this
264 digestion process. A portion of the filter was soaked in the tetrafluoromethaxil (TFM) vessels
265 (SK-10, Milestone, Germany) of the microwave where total mass of the sample and reagent
266 was maintained below 0.25 g for quality assurance purposes. Upon completion, the samples
267 were filtered using a syringe filter (Acrodisc[®], 0.2 μ m, Pall Gelman Laboratory, MI, USA)
268 with a 50 cc/mL Terumo syringe (Terumo[®], Tokyo, Japan) before dilution to 25 mL using
269 ultrapure water (UPW, 18.2 M Ω cm, Easypure[®] II, Thermo Scientific, Canada). For the
270 preparation of samples for water-soluble ion analysis, a portion of the filter samples was cut
271 into small pieces and placed directly into 50 mL centrifuge tubes with UPW. For this
272 extraction, a combination of ultrasonic vibration, centrifuge and mechanical shaking were
273 applied. The samples were first sonicated in an ultrasonic bath (Elmasonic S70H, Elma,
274 Germany) for 20 min. Then, the extraction solutions were centrifuged at 2500 rpm (Kubota
275 5100, Japan) for 10 min before shaken using a vortex mixer for 10 min. The sonication and
276 centrifuged steps were repeated for two more times before the extract was filtered through
277 glass microfiber filters (Whatman[™], UK). Both the trace elements and water-soluble ion
278 extracts were refrigerated at 4 °C until further analysis. The trace elements (Al, Ba, Ca, Fe,
279 Mg, Pb, Zn, Ag, As, Cd, Cr, Li, Be, Bi, Co, Cu, Mn, Ni, Rb, Se, Sr and V) were determined
280 by Inductively Coupled Plasma Mass Spectroscopy (ICP-MS, PerkinElmer ELAN 9000,
281 USA) while the water-soluble ionic composition (Na^+ , NH_4^+ , K^+ , Ca^{2+} , Mg^{2+} , Cl^- , NO_3^- and
282 SO_4^{2-}) was determined using Ion Chromatography (Metrohm 850 model 881 Compact IC
283 Pro, Switzerland). Metrosep A-Supp 5-150/4.0 and C4-100/4.0 columns were used in the
284 determination of cations and anions, respectively. 1.7 mmol L⁻¹ nitric and 0.7 mmol L⁻¹
285 dipicolinic acid (Merck KGaA, Germany) were prepared to be used as eluents for cations.
286 Eluents of 6.4 mmol L⁻¹ sodium carbonate (Na_2CO_3) (Merck KGaA, Germany) and 2.0 mmol
287 L⁻¹ sodium bicarbonate (NaHCO_3) (Merck KGaA, Germany) were prepared and used to
288 measure anions (Cl^- , NO_3^- and SO_4^{2-}) with a flow rate of 0.7 mL min⁻¹. 100 mmol L⁻¹
289 Suprapur[®] sulfuric acid (H_2SO_4) (Merck KGaA, Germany) was also prepared to use as a
290 suppressor regenerant and ions were detected by a conductivity detector.

291 **2.3 Quality assurance and quality control (QA/QC)**

292 As part of QA/QC, the concentrations of the composition of PM_{2.5} were corrected from the
293 reagent and filter blanks samples, which were treated with a similar procedure to the exposed
294 filters. To determine the recovery (%) of the heavy metals, a standard reference material
295 (SRM), Urban Particulate Matter SRM 1648a obtained from the National Institute of
296 Standards and Technology (NIST), USA, was treated using the procedures outlined above.
297 The method detection limit (MDL) for trace elements is calculated as three times the standard
298 deviation of ten replicates of the reagent blank. Three samples of filter blanks were used to
299 calculate the MDL of water-soluble ions. Overall MDL were as reported in Table 1. During
300 the trace element analysis by ICP-MS, two modes of analysis were applied with updated
301 calibration curves each time. Based on trial runs and SRM1648a, the elements were initially
302 screened for concentration levels which resulted in two modes analysis: (a) a set of metals
303 (Al, Ca, Fe, Mg, Zn and Mn) with high concentrations (with several dilution factors); and (b)
304 a set of metals (Ba, Pb, Ag, As, Cd, Cr, Li, Be, Bi, Co, Cu, Ni, Rb, Se, Sr and V) with low
305 concentrations.

306 **2.4 Local circulation of wind and biomass fire hotspots**

307 Each year, Peninsular Malaysia experiences two monsoon regimes, the south-west (SW)
308 monsoon (June-September) and the north-east (NE) monsoon (December-March). During the
309 SW monsoon, south-west winds dominate the wind pattern in Peninsular Malaysia, inducing
310 drier weather. During the NE monsoon, strong north-east winds dominate over the Peninsular
311 Malaysia, bringing more rainfall to the east coast. To investigate this, the regional synoptic
312 wind field 10 m above the surface and resolution of $0.25 \times 0.25^\circ$ ranging from latitude: -10° ,
313 25° N, longitude: 85° , 125° E was plotted using Open Grid Analysis and Display System
314 (GrADS version 2.0.2). The wind field used to demonstrate the monsoon regimes in this
315 study is a gridded product produced by the global atmospheric reanalysis known as ERA-
316 Interim, by the European Centre for Medium-Range Weather Forecasts (ECMWF) (Dee et
317 al., 2011).

318 The ERA-Interim 10 m surface wind vectors (January 2004 to June 2014) show the two
319 opposite monsoon regimes experienced by Peninsular Malaysia (Fig. 2). It can be seen that
320 the south-west wind, from June to August, blew from Sumatra Island, Indonesia to Peninsular
321 Malaysia was generally weaker with wind speed around $1-2 \text{ m s}^{-1}$. Whereas the north-east
322 wind, from November to January, was much stronger, with wind speeds of around $5-7 \text{ m s}^{-1}$
323 (Fig. 2).

324 Biomass fire hotspots and the travel path of the monthly back trajectories of each season were
325 also plotted (Fig. 3). The mean clusters of back trajectories were produced using the Hybrid
326 Single-Particle Lagrangian Integrated Trajectory Model (HYSPLIT 4.9), and were re-plotted
327 using the graphical software, IGOR Pro 6.0.1 (WaveMetrics, OR, USA). A release height of
328 about 500 m for 120 h back trajectories with 6 h intervals was chosen. Trajectory start time
329 was chosen at 16:00 (UTC) to represent 24:00 (local time). The cluster mean of trajectories
330 was regarded by numeral number and colour (1-red, 2-green, 3-turquoise, and 4-purple). The
331 fire hotspot data of the Moderate Resolution Imaging Spectroradiometer (MODIS) was used
332 to investigate the biomass burning hotspots in the specific area of interest. The data was
333 downloaded from the National Aeronautics and Space Administration (NASA)-Land
334 Atmosphere near Real-Time Capability for Earth Observing System (EOS)-Fire Information
335 for Resource Management System (NASA LANCE FIRMS) fire archive covering an area
336 from 15°S to 30°N and 80°W to 130°E. In addition, to investigate the variability of the
337 boundary layer height around the region of Peninsular Malaysia, ERA-Interim boundary
338 layer height (BLH) gridded data from January 2000 to December 2014 was downloaded from
339 the European Centre for Medium-Range Weather Forecasts (ECMWF). The resolution of this
340 data was $0.5^{\circ} \times 0.5^{\circ}$, covering the domain of the Peninsular Malaysia (lat: 99 – 105°; lon: 0 –
341 9°). Yearly daily means of the ERA-Interim BLH data were calculated using the Climate Data
342 Operators (CDO) version 1.6.9 software (<https://code.zmaw.de/projects/cdo>) developed by
343 the Max-Plank-Institute, by first calculating the area mean.

344 2.5 Enrichment Factor (EF)

345 The EF of the heavy metals was calculated based on the abundance of elements in the Earth's
346 crust published by Taylor (1964). The EF of each element can be defined using the following
347 equation:

$$348 \quad EF = \frac{\left(\frac{E}{Al}\right)_{PM\ 2.5}}{\left(\frac{E}{Al}\right)_{Crust}} \quad (1)$$

349 where E/Al is the concentration ratio of element, E, to the reference metal, Al. Al was
350 selected as the reference element to calculate the annual and seasonal EF. Several other
351 researchers also used Al as the reference element (Birmili et al., 2006; Khan et al., 2010a;
352 Sun et al., 2006). Chester et al. (2000); Cheung et al. (2012); Khan et al. (2010a); Mohd Tahir
353 et al. (2013); Torfs and Van Grieken (1997) proposed a EF cut-off of ten to differentiate
354 between crustal and natural and anthropogenic origins of heavy metals. Thus, we consider

355 EF = 10 as the cut-off point. Therefore, a good number of the metals (Zn, Cr, Rb, Be, V, Fe,
356 Ca, Co, Sr, Pb, As and Bi) in PM_{2.5} in this study can be assumed to originate from
357 anthropogenic sources; Fig. 5a. These heavy metals were not natural or from the Earth's
358 crust. No seasonal differences were observed in the EF of the heavy metals.

359 2.6 Mass closure model

360 A study by Harrison et al. (2003) introduced a PMC model for the source apportionment of
361 particulate matter, which is the basis for this study. The variables were grouped into the
362 following four sub-classes: i) mineral matter (MIN), ii) sea salts (SS), iii) secondary
363 inorganic aerosol (SIA), iv) trace elements (TE) and v) undefined (UD). MIN is derived from
364 the sum of Al, Mg, K, Ca, and Fe multiplied by the appropriate factors to convert them into
365 their corresponding oxides as described by the following Eq. (2). Ca was multiplied by a
366 factor of 1.95 to account for CaO and CaCO₃ as this metal is assumed to be present in these
367 two forms (Remoundaki et al., 2013; Sillanpää et al., 2006; Terzi et al., 2010)

368

$$369 \text{MIN} = 1.89\text{Al} + 1.66\text{Mg} + 1.21\text{K} + 1.95\text{Ca} + 1.43\text{Fe} \quad (2)$$

370

371 The contribution of SS was estimated by assuming that soluble Na⁺ in PM_{2.5} samples
372 originated solely from the marine source and is based on the composition of seawater,
373 ignoring potential atmospheric transformation (Seinfeld and Pandis, 2012). Following Terzi
374 et al. (2010), the composition of sea salt comprised of the following Eq. (3)

375

$$376 \text{SS} = [\text{Na}^+] + [\text{ss-Cl}^-] + [\text{ss-Mg}^{2+}] + [\text{ss-K}^+] + [\text{ss-Ca}^{2+}] + [\text{ss-SO}_4^{2-}] \quad (3)$$

377

378 where, ss-Cl⁻ = 1.8*Na⁺, ss-Mg²⁺ = 0.12*Na⁺, ss-K⁺ = 0.036*Na⁺, ss-Ca²⁺ = 0.038*Na⁺ and
379 ss-SO₄²⁻ = 0.252*Na⁺. Meanwhile, SIA can be estimated by the sum of non-sea salt-sulfate
380 (nss-SO₄²⁻), NO₃⁻ and NH₄⁺ as explained by Remoundaki et al. (2013); Terzi et al. (2010)
381 with the following Eq. (4)

382

$$383 \text{SIA} = [\text{nss-SO}_4^{2-}] + [\text{NO}_3^-] + [\text{NH}_4^+] \quad (4)$$

384

385 Finally, TE is calculated by the sum of rest of the metals analysed in this study and UD
386 represents unidentified gravimetric mass of PM_{2.5}. Therefore, the overall mass closure
387 equation applied in this work can be expressed as the following Eq (5)

388

$$\begin{aligned}
& PM_{2.5} MC = MIN(1.89Al + 1.66Mg + 1.21K + 1.95Ca + 1.43Fe) + \\
389 & SS([Na^+] + [ss-Cl^-] + [ss-Mg^{2+}] + [ss-K^+] + [ss-Ca^{2+}] + [ss-SO_4^{2-}]) + \\
390 & SIA([nss-SO_4^{2-}] + [NO_3^-] + [NH_4^+]) + TE + UD \quad (5)
\end{aligned}$$

391 2.7 Source Apportionment of PM_{2.5} using PMF

392 Source apportionment of PM_{2.5} was conducted using the US EPA PMF 5.0 model of the
393 United States Environmental Protection Agency (US EPA) as suggested by Norris et al.
394 (2014). The PMF model is a mathematical factor-based receptor model that interprets source
395 types with a robust uncertainty estimate. Two sets of data were run through the PMF model:
396 i) concentration and ii) uncertainty. The concentration of each element was pretreated and
397 validated based on the outliers, missing values and/or values below MDL. In particular,
398 variables with outliers were excluded. Species with concentrations below MDL were
399 replaced with the half of the MDL (Baumann et al., 2008; Polissar et al., 1998a; Polissar et
400 al., 1998b). The uncertainty value of each variable of each sample was calculated following
401 the empirical formula Eq. (6):

$$402 \sigma_{ij} = 0.01(X_{ij} + \bar{X}_j) \quad (6)$$

403 Where σ_{ij} is the estimated measurement error for jth species in the ith sample, X_{ij} is the
404 observed elements concentration and \bar{X}_j is the mean value. The factor 0.01 was determined
405 through trial and error procedures following by Ogulei et al. (2006a). Thus, the measurement
406 of uncertainty (S_{ij}) can be computed with Eq. (7) as applied by Chueinta et al. (2000):

$$407 S_{ij} = \sigma_{ij} + CX_{ij} \quad (7)$$

408 Where σ_{ij} the estimation of measurement error (Eq. 6) and C is a constant. In this study, we
409 used a value of 0.4 for C which, according to Ogulei et al. (2006b), produced the best Q value
410 as it is the closest to theoretical value and physically interpretable results. Other main
411 researchers have also applied this procedure for the calculation of uncertainty (Harrison et al.,
412 2011; Hedberg et al., 2005; Khan et al., 2015b). An additional 5% uncertainty was added to
413 cover any methodological errors during the preparation of filter papers, gravimetric mass
414 measurements and preparing the calibration curves.

415 Initially, PMF factors were resolved using the numbers of 20 runs with a seed value of 9. The
416 number of factors was changed to optimize the goodness-of-fit parameter of Q over the

417 theoretical Q. Five factors were decided upon based on the lowest Q (Robust) and Q (True)
418 value of 180.26 with the Q (true)/Q_{exp} value of 0.50 after 604 computational steps and the
419 convergence of the PMF results. The Q/Q_{exp} ratio for most of the variables was < 5 to 0.92
420 which indicates that the Q values were very similar to the expected value. Some of the
421 variables, however, showed a ratio of 0.5 because the computed Q value were smaller than
422 the expected Q value. A study by Brown et al. (2012) described this discrepancy as
423 contributing to the increase of global uncertainty. However, the sharp drop for PM_{2.5} mass
424 ratio (0.03) was due to the down-weighting of the signal to noise (S/N) values. To show the
425 stability of the results, we estimated the error of the concentration for each variable using
426 bootstrap, displacement (DISP) and a combination of BS-DISP. A comparison of the error
427 estimates with base model runs are demonstrated in the supplementary Fig. S1. The five-
428 factor results were relatively stable with meaningful physical interpretation and satisfactorily
429 comparable with the bootstrap analysis. Fe and Cr were reported as outliers and therefore
430 excluded in the calculation. Referring to Table 2, the overall PM_{2.5} concentration is well
431 explained within ±10% by the PMF 5.0 considering the F_{peak} = 0.

432 **2.8 Health risk assessment (HRA) of PM_{2.5} and associated various sources**

433 The human health risk posed by heavy metals may occur through inhalation of PM_{2.5}. We
434 applied the US EPA supplemented guidance to estimate the risk posed by heavy metals in
435 PM_{2.5} mass concentration and their various sources. As part of the HRA, we considered
436 lifetime non-carcinogenic and carcinogenic risk. USEPA (2011) describes the exposure
437 concentration (EC) by the following equation:

$$438 \quad EC_{inh} = C \times \frac{ET \times EF \times ED}{ATn} \quad (8)$$

439 Where *C* is the concentration of metals in PM_{2.5} estimated for each source with µg m⁻³ unit
440 for the estimation of *EC_{inh}*; EF is the exposure frequency (151 days year⁻¹) representing July,
441 August, September, January and February; ED is exposure duration (24 years for adult); BW
442 is the average body weight (70 kg for adult); ET is the exposure time (h/day); ATn is the
443 average time (ATn = ED × 365 days × 24 h/day for non-carcinogenic and ATn = 70 year ×
444 365 days/year × 24 h/day for carcinogenic risk). ED, BW and AT values are based on the
445 study by Hu et al. (2012).

446 Further, we examined the non-carcinogenic risk (presented by the hazard quotient (HQ)) and
447 lifetime carcinogenic risk (LCR) of selected heavy metals as classified by the International

448 Agency for Research on Cancer (IARC). The following equations were involved for the
449 calculation of HQ and LCR:

$$450 \quad HQ = \frac{EC}{(RfCi \times 1000 \mu g m^{-3})} \quad (9)$$

$$451 \quad LCR = IUR \times EC \quad (10)$$

452 Where, RfC_i is the inhalation reference concentration ($mg m^{-3}$); IUR is the inhalation unit risk
453 ($(\mu g m^{-3})^{-1}$). The non-carcinogenic risk or HQ represents the observable health effects from
454 exposure to the $PM_{2.5}$ based on the dose-response relationship principles. The cut-off point
455 for significant health risks to the exposed population is $HQ > 1$. The carcinogenic risk refers
456 to a person's chance of developing cancer from exposure to any carcinogenic agent. LCR
457 represents the excess lifetime cancer risk is described in terms of the probability that an
458 exposed individual will develop cancer because of that exposure by age 70 as defined by US
459 EPA Risk Communication
([http://www.epa.gov/superfund/community/pdfs/toolkit/risk_communicati-
460 onattachment6.pdf](http://www.epa.gov/superfund/community/pdfs/toolkit/risk_communicationattachment6.pdf)). The carcinogenic risk from the lifetime exposure of those hazardous
461 metals is regulated by the acceptable or tolerance level (1×10^{-6}) set by the US EPA which
462 corresponds to lifetime exposure to an unpolluted environment (Satsangi et al., 2014).

464

465

466 **3 Results and Discussions**

467 **3.1 Concentration of $PM_{2.5}$ and its chemical composition**

468 Table 1 summarizes the statistics from the SW monsoon, the NE monsoon and overall
469 concentrations of $PM_{2.5}$, heavy metals and major ions. Overall, the 24 h average values of
470 $PM_{2.5}$ (avg = $25.13 \mu g m^{-3}$) in the study area are slightly higher than that of the WHO 24 h
471 guideline ($25 \mu g m^{-3}$) but lower than that of 24 h US EPA National Ambient Air Quality
472 Standard (NAAQS) ($35 \mu g m^{-3}$). Of the samples taken during the day, 48% exceeded the
473 WHO 24 h guideline while 19% of them exceeded the US EPA 24 h NAAQS for $PM_{2.5}$
474 (Currently Malaysia has no set guidelines for $PM_{2.5}$). If we compare the $PM_{2.5}$ overall value
475 of $25.13 \mu g m^{-3}$ with yearly mean of US EPA NAAQS ($15 \mu g m^{-3}$), WHO ($10 \mu g m^{-3}$),
476 European Union (EU) ($25 \mu g m^{-3}$), DoE (Australia) ($8 \mu g m^{-3}$), the concentration of $PM_{2.5}$ is
477 much higher with respect the guideline set by all regulatory bodies. The average value of
478 $PM_{2.5}$ during the NE monsoon was slightly higher than the SW monsoon. During the

479 southwest monsoon season, PM_{2.5} was mainly carried by the prevailing southwest wind from
480 the Sumatra Island of Indonesia which is located at the Southwest quadrant of the SEA
481 region. On the other hand, during the northeast monsoon season, the PM_{2.5} sources can be
482 traced back to the Chinese mainland, Indochina region and the Philippines. This is due to the
483 prevailing northeast wind transporting PM_{2.5} from these locations to the tropical region of
484 SEA. The *Student T-test* for paired independent samples showed that the mean during these
485 two monsoons varies insignificantly ($t = 1.19, p > 0.05$). However, the monsoonal changes in
486 this region as displayed in Fig. 3, showed that air masses of different origins transport
487 different pollutants to the area. The back trajectory plots showed that there were high
488 numbers of biomass fire hotspots during both seasons but from different regions (Fig. 3). The
489 period of June to September is the dry season each year in Malaysia and Sumatra of
490 Indonesia. During this dry season, biomass fire hotspots are densely located in this area due
491 to the burning of agricultural waste and forest fires. Several other researchers also reported
492 the high number of biomass fire - related hotspots to these regions (Khan et al., 2015c;
493 Sahani et al., 2014). On the other hand, December to March is usually the wet season in
494 Malaysia. However, the backward trajectories showed that air masses were transported from
495 Mainland China and neighbouring regions. In Mainland China and neighbouring regions, this
496 is dry season. During the dry season in this region there are a lot of fires, as reported by
497 Zhang et al. (2015) and Ho et al. (2014), and this influences the pollution of air masses
498 transported to the present location. This scenario of biomass fire hotspots is commonly
499 noticed in these two seasons. In past years Malaysia and Singapore have experienced
500 intensified haze episodes in this particular season, e.g. 1997, 2005, 2013 and 2015. A study of
501 this area by Kanniah et al. (2014) observed that during the dry season (June to September)
502 aerosols mainly originated from the west and south-west (i.e. Sumatra, Indonesia), while
503 during the wet season (November to March), aerosols were mostly associated with the NE
504 monsoon winds coming from the South China Sea. Also, the variability of BLH and WS
505 were able to influence the concentration of the pollutants at a particular location. Fig. 4 shows
506 the day to day variation of BLH and WS with respect to the 24 h average of PM_{2.5}
507 concentration. From the plot, it is revealed that the daily average PM_{2.5} concentration is
508 inversely proportional to the BLH. Therefore, while factors such as traffic volume, industrial
509 emissions, power plants, land use and population size can alter the concentration of PM_{2.5},
510 meteorological factors which govern the day to day variation of BLH in Peninsular Malaysia
511 might play a crucial role too. These meteorological factors can include strong local
512 convection, which is a very common meteorological feature in this region, and also the

513 movement of air via a land-sea breeze due to the sea surrounding Peninsular Malaysia. A
514 study by Lelieveld et al. (2001) reported that strong convection can ventilate the daily BLH.
515 The small expansion of BLH that was observed during NE monsoon was most likely due to
516 the higher magnitude of WS to Peninsular Malaysia during this season, as demonstrated in
517 Fig. 2.

518 In comparison, our results of $PM_{2.5}$ here on the west coast of Peninsular Malaysia (avg =
519 $25.13 \mu\text{g m}^{-3}$) are higher compared to the east coast of Peninsular Malaysia at $14.3 \mu\text{g m}^{-3}$
520 (Mohd Tahir et al., 2013). This $PM_{2.5}$ concentration in this study area was similar to the
521 annual concentration of $PM_{2.5}$ measured in Petaling Jaya, Kuala Lumpur ($26.85 \mu\text{g m}^{-3}$) by
522 Rahman et al. (2011), Petaling Jaya ($33 \mu\text{g m}^{-3}$) and Gombak ($28 \mu\text{g m}^{-3}$) by Keywood et al.
523 (2003) and Singapore ($27.2 \mu\text{g m}^{-3}$) as reported by Balasubramanian et al. (2003). The yearly
524 mean value of $PM_{2.5}$ in the Bandung urban area and suburban location in Lembang of
525 Indonesia are 14.03 and $11.88 \mu\text{g m}^{-3}$, respectively (Santoso et al., 2008), which are much
526 lower concentration compared to the this study. However, Lestari and Mauliadi (2009)
527 reported that the $PM_{2.5}$ concentration of $43.5 \mu\text{g m}^{-3}$ in the Bandung city, Indonesia, was
528 about 1.7 times larger than that of the current location and by Budhavant et al. (2015) showed
529 $19 \mu\text{g m}^{-3}$ in Male, Maldives (urban) which is lower as well compared to this study. A
530 comparative study conducted in Bangkok ($34 \mu\text{g m}^{-3}$), Beijing ($136 \mu\text{g m}^{-3}$), Chennai ($44 \mu\text{g}$
531 m^{-3}), Bandung ($45.5 \mu\text{g m}^{-3}$), Manila ($43.5 \mu\text{g m}^{-3}$) and Hanoi ($78.5 \mu\text{g m}^{-3}$) showed
532 consistently higher $PM_{2.5}$ pollution in the Southeast and South Asian cities as compared to
533 this study (Kim Oanh et al., 2006). From Table 1, it can be seen that the highest concentration
534 of anions species was found for SO_4^{2-} followed by NO_3^- . A study by Zhang et al. (2012)
535 suggested that the photochemical conversion of SO_2 to H_2SO_4 is the main reason for the
536 changes of sulfate concentration in $PM_{2.5}$ and that higher temperatures reduce the nitrate
537 concentration by the partitioning of nitrate into the gas phase. However, lower temperatures
538 and a stable atmosphere favours the formation of NO_3^- aerosol reacting with NH_4^+ , i.e.
539 shifting the gas phase nitrate into the particle phase (Mariani and de Mello, 2007). The
540 formation of NH_4NO_3 normally occurs at high humidity with lower temperatures (Morales
541 and Leiva, 2006). The average molar ratios of SO_4^{2-} to NO_3^- were 6.0 with a range of 0.16 –
542 38.24 which suggests that the sulfate aerosol is more dominant over the nitrate aerosol and
543 may have been transported from trans-boundary sources. A similar observation was found in
544 a study in the UK by Abdalmogith and Harrison (2006). This ion balance ratio indicates the
545 possible sources of aerosol, and stationary sources dominate over the mobile sources as

546 explained by Arimoto et al. (1996). The average ratio of SO_4^{2-} to NH_4^+ was 1.28 which is
547 much higher than the ratio of NO_3^- to NH_4^+ (0.63), confirming that the sulfate aerosol is more
548 stable in tropical conditions as compared to the nitrate aerosol. A study by Maenhaut et al.
549 (2008) described a similar observation. The formation of ammonium sulfate is favoured in
550 the fine fraction (Khan et al., 2010b). For the cations, the highest concentration was NH_4^+
551 followed by Na^+ . However, the average molar ratio of Cl^- to Na^+ did not reflect the seawater
552 ratio. “Cl loss” may be the cause of the drop in Cl^- to Na^+ ratio. Boreddy et al. (2014) also
553 noticed a chlorine depletion due to atmospheric processing in the western north Pacific.
554 According to Finlayson-Pitts and Pitts Jr. (2000), sulfuric and nitric acids have a tendency to
555 react rapidly with NaCl, the major component of sea salt particles, to produce gaseous HCl
556 under 50–100% relative humidity conditions. For heavy metals, the predominant metal
557 reported was Fe with concentrations in the range 2171 - 4567 ng m^{-3} . Ca showed the second-
558 highest concentrations with the concentration range of below MDL - 3149 ng m^{-3} . A study
559 by Yin and Harrison (2008) suggested that Fe originates from non-traffic sources and that
560 iron and calcium are released into ambient air through the resuspension of surface dust.
561 Among other heavy metals of particular health concern, the average concentrations of As, Pb,
562 Cd, Ni, Mn, V and Cr were 5.76, 21.84, 0.54, 4.03, 17.24, 5.13 and 107.68 ng m^{-3} ,
563 respectively. The As concentration was nearly equal to the WHO and US EPA guideline
564 values of 6.6 and 6 ng m^{-3} , respectively. Therefore, As may be of significant health concern.
565 The concentrations of other hazardous metals were well below the WHO and European
566 commission guidelines. The EF reveals that all metals of $\text{PM}_{2.5}$ can be assumed to originate
567 from anthropogenic sources with no seasonal differences observed (Fig. 5a).

568 **3.2 Mass closure model**

569 The $\text{PM}_{2.5}$ was reconstructed by the use of a PMC model (Harrison et al., 2003). Employing
570 the mass closure model outlined in the previous section, the four major classes of chemical
571 components contributing to $\text{PM}_{2.5}$ were: i) MIN, ii) SIA, iii) SS, iv) TE and UD. As shown in
572 Fig. 5c, the overall reconstructed masses of MIN, SIA, SS, TE and UD were 8970, 2841,
573 1727, 626.2 and 11511 ng m^{-3} , respectively. MIN is released from soil or crustal sources and
574 represents the oxide form of the metals (Remoundaki et al., 2013; Sillanpää et al., 2006; Terzi
575 et al., 2010). The MIN component comprises 35% of the $\text{PM}_{2.5}$ concentration. SIA, which
576 accounts for 11%, is comprised of the most abundant secondary ions (nss-SO_4^{2-} , NO_3^- and
577 NH_4^+). These are formed in the atmosphere from the precursor gases (SO_2 , NH_3 and NO_x)

578 through a gas-to-particle conversion (Sillanpää et al., 2006) and therefore are assumed to be
579 in the form of $(\text{NH}_4)_2\text{SO}_4$ and NH_4NO_3 in the aerosol phase (Joseph et al., 2012). It is
580 important to mention that the behaviour of the formation pattern of the SIA in this South East
581 Asia region may differ from other regions due to the nearly constant temperature throughout
582 the year. SS and TE accounted for 7% and 2% of the $\text{PM}_{2.5}$. The UD, undefined fraction,
583 accounted for 45% of $\text{PM}_{2.5}$.

584 The average value of reconstructed $\text{PM}_{2.5}$ by mass closure (MC) is $14.12 \pm 4.32 \mu\text{g m}^{-3}$ with a
585 minimum of $6.70 \mu\text{g m}^{-3}$ and a maximum of $24.19 \mu\text{g m}^{-3}$. On the other hand, the measured
586 $\text{PM}_{2.5}$ determined gravimetrically by HVS was $25.13 \pm 9.21 \mu\text{g m}^{-3}$ with a range of 7.01 to
587 $42.78 \mu\text{g m}^{-3}$. A correlation plot of $\text{PM}_{2.5}$ (MC) and measured $\text{PM}_{2.5}$ (HVS) is shown in Fig.
588 5e. The correlation shows a good fit ($r^2 = 0.98$) with a slope of 0.46 and an intercept of 1.93.
589 The results of the fit parameters suggest that the $\text{PM}_{2.5}$ mass (MC) concentration was
590 underestimated compared to $\text{PM}_{2.5}$ (HVS). The reported result of the mass closure model is
591 based on the analyzed chemical components of filter samples (~ 55%). As described in the
592 mass closure, a large portion of $\text{PM}_{2.5}$ mass (~ 45%) was left unidentified; this unidentified
593 component is believed to be the organics or carbonaceous species. Elemental carbon (EC),
594 organic carbon (OC) and water-soluble organics were not measured due to the lack of
595 instrumentation. Other possible reasons for the un-identified portion are: i) unaccounted for
596 mineral oxides as they are abundant in $\text{PM}_{2.5}$, and ii) water associated with salts.

597 During the SW monsoon, the UD showed the higher concentrations; this can be explained by
598 the annual biomass haze episodes experienced in this area. Thus, a large proportion of the UD
599 of $\text{PM}_{2.5}$ is probably formed from the organic fraction. Such findings are consistent with a
600 study conducted by Abas and Simoneit (1996) which also found that the concentrations of
601 organic compounds observed were greater during the haze episodes than any other periods in
602 a year, and that some of them are suspected to be transported from trans-boundary sources.

603 The seasonal variability of the results obtained from the mass closure model is shown in Fig.
604 5c. The reconstructed masses of MIN, SIA and SS were higher in the NE than the SW
605 monsoon. These haze events were very likely caused by the slash-and-burn activities
606 practiced by the agriculture industries, and the occurrence of forest fires during this dry
607 season. The regional trans-boundary pollution during the NE and SW monsoon is the
608 underlying reason for the change in the chemical component concentrations as well as the
609 overall $\text{PM}_{2.5}$.

610

611 3.3 Identification and apportionment of PM_{2.5} sources

612 Using US EPA PMF 5.0, the five identified sources of PM_{2.5} were i) mineral and road dust,
613 ii) motor vehicle emissions and biomass burning, iii) nitrate aerosol, iv) coal burning and v)
614 marine and sulfate aerosol. Each of the source profiles is shown in Fig. 6a which
615 demonstrates the concentration and percentage of the variables to each factor. The reported
616 PMF analysis is based on the chemical components of filter samples. As described in the
617 mass closure, a large portion of the PM_{2.5} mass fraction (about 45%) was not apportioned. In
618 the PMF 5.0 procedure, the contributions of five factors were estimated and then the
619 integrated contribution of the five factors was regressed over the measured PM_{2.5} (HVS). The
620 regression fit line was forced through the origin. Thus, our regression of the PM_{2.5} (PMF) and
621 PM_{2.5} (HVS) showed that the PM_{2.5} had been reproduced by PMF 5.0 with an error of less
622 than 10% and the correlation of PM_{2.5} (PMF) and PM_{2.5} (HVS) showed a strong and
623 significant correlation (slope = 0.91, $r^2 = 0.88$, $p < 0.01$) (Fig. 6b). To evaluate the results of
624 the PMF model, the regression between predicted and observed data for each variable is
625 shown during the operation. A linear correlation between the predicted and measured mass
626 represents the goodness-of-fit of linear regression. Our values strongly suggested that the five
627 identified sources could be readily interpreted.

628

629 Factor component one: the predominant tracers are Mg, Zn, Cu, Ni and Ca²⁺. The mineral or
630 natural fugitive dust component is identified based on the presence of Mg (52% of the Mg
631 mass), Ca²⁺ (42% of Ca²⁺ mass), Ca (28% of Ca mass) and Al (19% of Al mass), as shown in
632 Table 2. Many other researchers cite these metals as markers for a mineral dust source
633 (Dall'Osto et al., 2013; Moreno et al., 2013; Mustaffa et al., 2014; Viana et al., 2008; Waked
634 et al., 2014). The possible cause of the mineral dust is the rapid development activities such
635 as construction, renovation of road surface etc, around this suburban region. Airborne soil
636 and construction material are the key sources of mineral dust (Dai et al., 2013; Gugamsetty et
637 al., 2012; Huang et al., 2014). Cu, Zn and Ba are associated with road dust due to the release
638 of these metal markers from cars from non-exhaust sources (Amato et al., 2011). Several
639 studies identified that Cu is released from brake wear or the brake pads/tailpipes of cars
640 (Wåhlin et al., 2006) while Zn originates from tire wear (Dall'Osto et al., 2013) and additives
641 in cars as lubricant (Ålander et al., 2005). A study by Wang and Hopke (2013) suggested that
642 Ni was emitted from gasoline engine and road dust sources. Ni (18% of Ni mass) and V (20%
643 of V mass) are moderately presented in this factor component which shows the existence of

644 heavy lubricating oil combustion (Amato et al., 2011). The average contribution of mineral or
645 natural fugitive and road dust sources to the PM_{2.5} was 3.17 µg m⁻³ or 13%.

646 Factor component two: contains substantial Pb, NH₄⁺ and K⁺. Motor vehicle emissions and
647 biomass burning sources accounted for 7.47 µg m⁻³ or 31% of the total PM_{2.5} concentration,
648 which makes these the largest sources contributing to the PM_{2.5} concentration. Pb along with
649 the moderately enriched metals As, Cd, Zn, Ni and V (Fig. 5a, refer to previous section for
650 detail), represents a motor vehicle emission source (Wu et al., 2007). The brake wear dust of
651 motor vehicles contains Pb (Garg et al., 2000). A study by Begum et al. (2010) conducted in
652 Dhaka and by Santoso et al. (2013) at roadsides in Jakarta defined Pb in PM_{2.5} releasing from
653 the pre-existing road dust by PMF. Choi et al. (2013) also introduced Pb in PM_{2.5} as a tracer
654 for the motor vehicle source. Zn is released from the wear and tear of tyres
655 (Srimuruganandam and Shiva Nagendra, 2012). Further, Zn in PM_{2.5} appeared to have a
656 motor vehicle source as resolved by PMF, due to its use as fuel detergent and anti-wear
657 additive (Brown et al., 2007). Ni and V were widely reported in the literature as markers for
658 the combustion of engine oil or residual oil combustion (Gugamsetty et al., 2012; Han et al.,
659 2006; Huang et al., 2014; Yu et al., 2013). Pb is no longer used as an additive in gasoline
660 fuel. Thus, the Pb does not reflect the emissions from engine combustion but does reflect
661 those from a non-exhaust traffic source. A study conducted by Rahman et al. (2011) in Kuala
662 Lumpur investigating Pb in PM_{2.5} found that it originated from the soil dust source, indicating
663 the influence of road dust. Also, coal combustion is a predominant source of Pb (Tao et al.,
664 2014). The K⁺ ion has been widely cited in the literature as an excellent tracer representing a
665 wood or biomass burning source (Dall'Osto et al., 2013; Kim and Hopke, 2007; Mustaffa et
666 al., 2014; Wahid et al., 2013). The biomass burning source is generally comprised of either
667 wood burning as residential fuel, agriculture residue/waste, and/or wild forest fires. In Kuala
668 Lumpur, the biomass burning source was described due to the presence of K from PM_{2.5}
669 measured by Particle Induced X-ray Emission (Rahman et al., 2011). During the episode of
670 biomass burning in Chengu, China, K⁺ and other related tracers in PM_{2.5} were increased by a
671 factor of 2 -7. In this suburban region, the smoke emissions released due to the burning of
672 wheat straw, rape straw and other biomass fuel for domestic cooking or heating purposes
673 (Tao et al., 2013). K⁺ is also mainly emitted from biomass burning in the suburb of
674 Shenzhen, China (Dai et al., 2013), Beijing, China (Yu et al., 2013; Zhang et al., 2013) and
675 Colombo, Sri Lanka (Seneviratne et al., 2011). In Seoul, Korea, biomass burning is
676 characterised by the presence of K and other related markers in PM_{2.5}. The character of
677 burning in this East Asian city is typically post-harvest field burning, biofuel burning for

678 heating and cooking as well as forest fire from the outside of the city (Heo et al., 2009). Thus,
679 the local and regional transport of smoke from the burning sources contribute to this factor.
680 Hong Kong experiences the influence of biomass burning in PM_{2.5} due to its transboundary
681 origin (Huang et al., 2014). During the sampling period in the SW monsoon, the MODIS
682 detected a very high number of fire counts over the Sumatra Island. In this monsoon season,
683 the wind will consistently travel from the southwest direction, bringing air masses from these
684 burning areas to Peninsular Malaysia. During the NE monsoon, on the other hand, the wind
685 will travel from the northeast direction, bringing air masses from the China mainland,
686 Indochina and the Philippines to Peninsular Malaysia. In this period of time, a high density of
687 fire locations were found on the Indo-China and China mainland. Zhang et al. (2015)
688 demonstrate that during the dry season there is important biomass burning activity in the
689 Pearl River Delta (China), which can result in trans-border transport and a regional scale
690 character of biomass burning. Therefore under the north-east monsoonal regime it is possible
691 that outflow from that area can maybe influence the specific area. A study by Streets et al.
692 (2003) estimated that China contributes 25% to the total biomass burning in Asia and showed
693 a good agreement between national estimate of biomass burning and adjusted fire count.
694 Yang et al. (2013) applied spatial-temporal features of fire counts and observed that the study
695 area of Heilongjiang Province, China was seriously affected by forest fires during 2000-2011.
696 Reid et al. (2013) reported a high intensity of fire counts in Vietnam/China region in April
697 and in Indonesia during September. Khan et al. (2015a) also reported a high density of fire
698 locations in Thailand, Vietnam and Laos during February and Sahani et al. (2014) reported
699 many in the same regions during June-September. The biomass burning is the dominant
700 source of trace gas and particulate matter and the fire emissions are mainly concentrated in
701 Indonesia, Thailand, Myanmar and Cambodia (Chang and Song, 2010). Further, a
702 comparison of nss-K⁺ with the respective total K⁺ is shown in Fig. 5b. The correlation of nss-
703 K⁺ as a function of total K⁺ showed a strong correlation coefficient ($r^2 = 0.95$) which suggests
704 that K⁺ can be used as a biomass tracer. K⁺ may also be emitted from local fire sources.
705 Additionally, the molar equivalent of K⁺ and Na⁺, as shown in Fig 5c, demonstrated
706 significant correlation ($r^2 = 0.70$) with a slope value of 0.34 which is much higher compared
707 to 0.0225-0.230 and 0.0218, reported by Wilson (1975) and Hara et al. (2012), respectively.
708 The higher molar ratio of K⁺ and Na⁺ indicates that at the current location, Na⁺ depletion was
709 high and the K⁺ might also release from other dominant sources. Additional significant
710 sources of K⁺ which may attribute to the mass are soil dust, sea salt, vegetation and meat
711 cooking (Zhang et al., 2010).

712 Factor component three: this factor is mainly dominated by the concentration of the
713 nitrate ion (75% of NO_3^- mass) suggesting that this source is strongly related to the formation
714 of nitrate aerosol. NO_3^- is mainly formed from the conversion of NO_x , which is emitted from
715 the exhaust of motor vehicles (Dai et al., 2013). Huang et al. (2014) also identified a nitrate
716 source in $\text{PM}_{2.5}$ by the use of PMF in suburban areas of Hong Kong. In Beijing, a nitrate
717 source appeared in $\text{PM}_{2.5}$ when source apportionment performed by PMF (Song et al., 2006).
718 This source is also contributed to by the small amount of Al, Mn and Ca^{2+} . Overall, it
719 accounted for $4.11 \mu\text{g m}^{-3}$ or about 17% of the $\text{PM}_{2.5}$ concentration.

720 Factor component four: this fourth source has an abundance of As, Ba and Sr (Se moderately
721 contributed) and thus is classified as coal burning source. As constitutes the most to this
722 fourth component at 53% (of As mass), which gives an indication that this source is related to
723 the coal combustion. In Malaysia, several power plants are operating on the west coast of
724 Peninsular Malaysia, e.g. Port Dickson, Kapar and Manjung. The power plants located at Port
725 Dickson and Kapar are about 50 km far from the sampling station. These plants use coal as
726 the raw material to generate electric power. Other researchers have also used As a tracer for
727 the coal burning source (Moreno et al., 2013) and As and Se by Meij and te Winkel (2007);
728 Querol et al. (1995). As and Se are categorized as of great concern and Ba and Sr are of
729 moderate environmental concern in the utilization of coal, as reported by Vejahati et al.
730 (2010). However, Ba is an indicator of brake wear and tear from motor vehicles (Gietl et al.,
731 2010). Overall, the coal burning source accounts for $4.60 \mu\text{g m}^{-3}$ or about 19% of $\text{PM}_{2.5}$.

732 Factor component five: this component features Na^+ (50% of Na^+ mass), Cl^- (46% of Cl^-
733 mass) and sulfate (64% of SO_4^{2-} mass) suggesting the presence of marine as well as sulfate
734 aerosol. Begum et al. (2010) identified sea salt in $\text{PM}_{2.5}$ by PMF in Dhaka, based on the
735 appearance of Na and Cl. Choi et al. (2013) defined a sea salt source in Seoul, Korea due to
736 the high contribution of Na^+ and Cl^- in $\text{PM}_{2.5}$. Several other studies in East, Southeast and
737 South Asia assigned a sea salt source in $\text{PM}_{2.5}$ considering Na^+ and Cl^- from the model output
738 of PMF (Lee et al., 1999; Santoso et al., 2008; Santoso et al., 2013; Seneviratne et al., 2011).
739 For sulfate, it shows that nss- SO_4^{2-} contributed 93% to the total sulfate concentration while
740 ss- SO_4^{2-} accounted for only 6%. Therefore, the sulfate aerosol in $\text{PM}_{2.5}$ is released as a
741 product from the photochemical conversion of SO_2 , which mainly originates from
742 anthropogenic large point sources as observed by Heo et al. (2009) in Seoul, South Korea. A
743 secondary sulfate source in $\text{PM}_{2.5}$ was also identified by Huang et al. (2014) in a suburban
744 area of Hong Kong and by Song et al. (2006) in Beijing. The marine and sulfate aerosol, as
745 the final identified source, accounts for $4.99 \mu\text{g m}^{-3}$ or about 20% of the total $\text{PM}_{2.5}$

746 concentration. A study by Kim and Hopke (2007) defined a sea salt source by the high
747 concentration of Na^+ and Cl^- , while sulfate sources are based on the high concentration of
748 sulfate. The secondary aerosol fraction is an important source worldwide, which is also the
749 case here. It generally constitutes a predominant portion of $\text{PM}_{2.5}$, which splits into two
750 modes i.e. the nitrate-rich and sulfate-rich factors. Studies by Chen et al. (2007) and
751 McGinnis et al. (2014) also identified the major contribution of the secondary aerosol
752 fraction to $\text{PM}_{2.5}$.

753 3.4 Health risk implications

754 Table 3 shows the non-carcinogenic (represented as HQ) and carcinogenic risks posed by
755 several selected metals (Pb, As, Cd, Cu, Mn, Zn and Ni) in $\text{PM}_{2.5}$ mass concentration through
756 inhalation exposure associating with sources. The HQ values for As and Ni in $\text{PM}_{2.5}$ mass
757 concentration are 15.9×10^{-2} and 14.3×10^{-2} , respectively, suggesting the non-carcinogenic
758 health risks posed by these metals might be higher compared to other metals. The HQ for
759 four selected metals (Pb, As, Cd and Ni) in $\text{PM}_{2.5}$ mass was highest in the $\text{PM}_{2.5}$ mass
760 originating from the coal burning source and least in $\text{PM}_{2.5}$ originating from the mineral/road
761 dust source. The cut-off point for significant health risks or the safe level to the exposed
762 population is $\text{HQ} > 1$. Our results showed that the sum of HQ for each metal are lower than
763 the safe level (= 1) in $\text{PM}_{2.5}$ mass concentration originating from each source. The sum of HQ
764 for $\text{PM}_{2.5}$ is 35.7×10^{-2} , which is lower than the HQs of $\text{PM}_{2.5}$ reported by Hu et al. (2012) in
765 Nanjing, China (2.96); Cao et al. (2014) in Shanxi Province, China ($1.06 \times 10^{+1}$); and Taner et
766 al. (2013) in a non-smoking restaurant in Turkey (4.09). A study by Hu et al. (2012), reported
767 HQ values for As and Ni in $\text{PM}_{2.5}$ as 4.14×10^{-1} and 1.73×10^{-1} , respectively, in Nanjing,
768 China. However, the HQs of $\text{PM}_{2.5}$ estimated after inhalation at two sites in Nanjing City,
769 China (0.88 (Xianlin) and 0.79 (Gulou)) were close to the safe level (= 1) according to a
770 study by Li et al. (2015b). At two urban locations in Yangtze River Delta, China, the HQ for
771 Cr in $\text{PM}_{2.5}$ was within the acceptable limit but higher for Mn (Niu et al., 2015). Although the
772 HQ calculated for As was the highest, it was below 1, thus the non-carcinogenic health risk
773 was estimated to be at a safe level. In addition, the hazard index (total - hazard quotient) of
774 $\text{PM}_{2.5}$ calculated for the four heavy metals (As, Cd, Mn, Ni) from the different sources (Table
775 3) showed an insignificant health risk.

776 The carcinogenic risks from the carcinogenic heavy metals Pb, As, Cd and Ni in $\text{PM}_{2.5}$
777 are shown in Table 3. Similar to the non-carcinogenic risks, the lifetime carcinogenic risk

778 level is estimated for PM_{2.5} mass concentration and may be contributed to by several heavy
779 metals from different sources: mineral/road dust, motor vehicle emissions/biomass burning
780 and coal combustion. The total life-time cancer risk (LCR) from heavy metals in the PM_{2.5}
781 mass concentration was calculated at 3.9×10^{-6} which is a significant cancer risk. The main
782 carcinogenic heavy metal of concern to the health of people at the current location is As; the
783 other heavy metals (Ni, Pb, and Cd) did not pose a significant cancer risk. Thus, the LCR
784 from the PM_{2.5} mass concentration originating from motor vehicle/biomass and coal burning
785 sources showed a value of 1×10^{-6} , slightly above the acceptable cancer risk level as
786 recommended by USEPA, while the total LCR from PM_{2.5} mass concentration from all
787 sources was estimated to be 4×10^{-6} which is also slightly above the acceptable cancer risk.
788 The carcinogenic risk posed by As (3.66×10^{-3}) in PM_{2.5} in Shanxi Province, China (Cao et
789 al., 2014) was higher than the guideline value set by USEPA. A study by Niu et al. (2015) of
790 PM_{2.5}-bound metals showed a high cancer risk in Yangtze River Delta, China (2.47×10^{-4}). A
791 study by Pandey et al. (2013) conducted in the vicinity of human activities observed that the
792 concentrations of Cd, Cr, Ni and Pb in PM_{2.5} showed higher excess cancer risk (ECR) due to
793 those particle-bound metals compared to guideline level set by USEPA. Satsangi et al. (2014)
794 also reported a higher cancer risk from Cr, Ni and Cd in PM_{2.5} compared to the USEPA
795 guideline. The integrated carcinogenic risk of six metals (Cr, As, Co, Pb, Ni and Cd) in PM_{2.5}
796 in Tianjin, China were in the range $3.4 \times 10^{-3} - 4.1 \times 10^{-3}$ which is reportedly beyond the
797 tolerance level (Zhang et al., 2014). The total ECRs based on the average values of As, Cd,
798 Cr, Ni and Pb in PM_{2.5} is 4.34×10^{-5} in Delhi, India, implying that four or five people might
799 get cancer out of 100,000 people after exposure to toxic metals in PM_{2.5} (Khanna et al.,
800 2015). Our findings showed that the life-time cancer risk posed by the exposure of heavy
801 metals in PM_{2.5} mass concentration is three to four per 1,000,000 people at this location. This
802 significant cancer risk warrants further investigation. Our findings showed that an
803 insignificant non-carcinogenic risk and significant cancer risk is posed to the population from
804 exposure to PM_{2.5} at this location. Detailed exposure assessment of the PM_{2.5} at the specific
805 sources and the health risks posed by individual hazardous elements of concern may help to
806 improve understanding about the exposure pathways as well as the detailed risk factors
807 involved in both carcinogenic and non-carcinogenic risk.

808

809 4 Conclusions

810 PM_{2.5} samples were collected using a high volume sampler in a semi-urban site on Peninsular
811 Malaysia. The results obtained for PM_{2.5} showed that 48% of the samples exceeded the WHO
812 24 h guideline and 19% exceeded the US EPA 24 h NAAQS for PM_{2.5}. The average value of
813 PM_{2.5} is higher than a previous study carried out on the east coast of Peninsular Malaysia.
814 PM_{2.5} concentration during the NE monsoon was slightly higher than the SW monsoon with
815 air masses from different origins. For the SW monsoon the air masses originated from
816 Sumatera, Indonesia and for the NE monsoon from Southern China. Major carcinogenic
817 metals, namely As, Pb, Cr, Ni and Cd, have contributed to PM_{2.5} by a significant portion. **Of**
818 **all carcinogenic metals**, As was close to the values set by the WHO and EU guidelines.
819 Results of the EF analysis suggested that a large number of the heavy metals in PM_{2.5} were
820 emitted from anthropogenic sources. No seasonal differences were found in the EF of the
821 heavy metals. The mass closure model results showed higher MIN, SIA and SS in the NE
822 than the SW monsoon. Further analysis revealed that sulfate is relatively more stable in
823 tropical climates compared to nitrate aerosol, indicating the dominance of static sources over
824 mobile sources. However, the average molar ratio of Cl⁻ to Na⁺ does not reflect the seawater
825 ratio. "Cl loss" may be the cause of the drop in Cl⁻ to Na⁺ ratio. The five sources of PM_{2.5}
826 obtained by the PMF 5.0 model were dominated by motor vehicle emissions/biomass burning
827 (7.47 µg m⁻³, 31%). The other four sources were mineral/road dust; nitrate aerosol; coal
828 burning; and marine-sulfate aerosol with an overall contribution of 3.17 µg m⁻³ (13%), 4.11
829 µg m⁻³ (17%), 4.60 µg m⁻³ (19%), and 4.99 µg m⁻³ (20%), respectively.

830 Using the PMF-identified sources as the basis, **the hazard quotient (HQ) for four selected**
831 **metals (Pb, As, Cd and Ni) in PM_{2.5} mass was highest in PM_{2.5} originating from a coal**
832 **burning source and least in that originating from a mineral/road dust** source. The non-
833 carcinogenic cancer risk posed by the exposure of PM_{2.5} was at a considerably safer level
834 compared to the South and East Asian region. The **lifetime cancer risk (LCR)** followed the
835 order of As > Ni > Pb > Cd for mineral/road dust, coal burning sources and PM_{2.5} **mass**
836 concentration, and As > Pb > Ni > Cd for motor vehicle/biomass burning. Among the trace
837 metals studied, As predominantly showed the largest **LCR in PM_{2.5} mass concentration** as
838 well as **its** associated sources, implying the largest risk after exposure of PM_{2.5} to people at
839 the current location. The associated **LCR** posed by the exposure of **the hazardous** metals in
840 PM_{2.5} **mass concentration** was three to four per 1,000,000 people at this location. This
841 significant LCR warrants further investigation. **Detailed exposure assessment of the PM_{2.5} at**
842 **the specific sources and the health risks posed by individual hazardous elements** may help to
843 improve understanding about the exposure pathways as well as the detailed **risk** factors

844 involved in both carcinogenic and non-carcinogenic risk. **Therefore**, the motor vehicle
845 emissions and regional trans-boundary pollution were the major underlying reasons for the
846 change in the chemical component of PM_{2.5} in tropical Peninsular Malaysia, which
847 potentially leads to different health threats.

848

849 **Acknowledgements**

850 The authors would like to thank the Malaysian Ministry of Higher Education for funding via
851 Fundamental Research Grants (FRGS/1/2013/SPWN01/UKM/02/) and Universiti
852 Kebangsaan Malaysia for University Research Grant (DIP-2014-005). Special thanks to Dr
853 Rose Norman for assistance with the proofreading of this manuscript

854

855 **References**

856

857 Abas, M. R., and Simoneit, B. R. T.: Composition of extractable organic matter of air
858 particles from malaysia: Initial study, *Atmos. Environ.*, 30, 2779-2793,
859 [http://dx.doi.org/10.1016/1352-2310\(95\)00336-3](http://dx.doi.org/10.1016/1352-2310(95)00336-3), 1996.

860 Abdalmogith, S. S., and Harrison, R. M.: An analysis of spatial and temporal properties of
861 daily sulfate, nitrate and chloride concentrations at UK urban and rural sites, *J.*
862 *Environ. Monit.*, 8, 691-699, 10.1039/B601562J, 2006.

863 Ålander, T., Antikainen, E., Raunemaa, T., Elonen, E., Rautiola, A., and Torkkell, K.:
864 Particle Emissions from a Small Two-Stroke Engine: Effects of Fuel, Lubricating Oil,
865 and Exhaust Aftertreatment on Particle Characteristics, *Aerosol Sci. Technol.*, 39,
866 151-161, 10.1080/027868290910224, 2005.

867 Amato, F., Viana, M., Richard, A., Furger, M., Prévôt, A. S. H., Nava, S., Lucarelli, F.,
868 Bukowiecki, N., Alastuey, A., Reche, C., Moreno, T., Pandolfi, M., Pey, J., and
869 Querol, X.: Size and time-resolved roadside enrichment of atmospheric particulate
870 pollutants, *Atmos. Chem. Phys.*, 11, 2917-2931, 10.5194/acp-11-2917-2011, 2011.

871 Anenberg, S., West, J. J., Yu, H., Chin, M., Schulz, M., Bergmann, D., Bey, I., Bian, H.,
872 Diehl, T., Fiore, A., Hess, P., Marmer, E., Montanaro, V., Park, R., Shindell, D.,
873 Takemura, T., and Dentener, F.: Impacts of intercontinental transport of

874 anthropogenic fine particulate matter on human mortality, *Air Qual. Atmos. Health*,
875 1-11, 10.1007/s11869-014-0248-9, 2014.

876 Arimoto, R., Duce, R. A., Savoie, D. L., Prospero, J. M., Talbot, R., Cullen, J. D., Tomza, U.,
877 Lewis, N. F., and Ray, B. J.: Relationships among aerosol constituents from Asia and
878 the North Pacific during PEM-West A, *J. Geophys. Res.*, 101, 2011-2023,
879 10.1029/95JD01071, 1996.

880 Balasubramanian, R., Qian, W. B., Decesari, S., Facchini, M. C., and Fuzzi, S.:
881 Comprehensive characterization of PM_{2.5} aerosols in Singapore, *J. Geophys. Res.*,
882 108, 4523, 10.1029/2002JD002517, 2003.

883 Baumann, K., Jayanty, R. K. M., and Flanagan, J. B.: Fine Particulate Matter Source
884 Apportionment for the Chemical Speciation Trends Network Site at Birmingham,
885 Alabama, Using Positive Matrix Factorization, *J. Air Waste Manage. Assoc.*, 58, 27-
886 44, 10.3155/1047-3289.58.1.27, 2008.

887 Begum, B. A., Biswas, S. K., Markwitz, A., and Hopke, P. K.: Identification of sources of
888 fine and coarse particulate matter in Dhaka, Bangladesh, *Aerosol Air Qual. Res.*, 10,
889 345-353, 10.4209/aaqr.2009.12.0082 2010.

890 Bell, M. L., Ebisu, K., Leaderer, B. P., Gent, J. F., Lee, H. J., Koutrakis, P., Wang, Y.,
891 Dominici, F., and Peng, R. D.: Associations of PM_{2.5} Constituents and Sources with
892 Hospital Admissions: Analysis of Four Counties in Connecticut and Massachusetts
893 (USA) for Persons \geq 65 Years of Age, *Environ. Health Persp.*, 122, 138,
894 10.1289/ehp.1306656, 2014.

895 Betha, R., Behera, S. N., and Balasubramanian, R.: 2013 Southeast Asian Smoke Haze:
896 Fractionation of Particulate-Bound Elements and Associated Health Risk, *Environ.*
897 *Sci. Technol.*, 48, 4327-4335, 10.1021/es405533d, 2014.

898 Birmili, W., Allen, A. G., Bary, F., and Harrison, R. M.: Trace Metal Concentrations and
899 Water Solubility in Size-Fractionated Atmospheric Particles and Influence of Road
900 Traffic, *Environ. Sci. Technol.*, 40, 1144-1153, 10.1021/es0486925, 2006.

901 Boldo, E., Linares, C., Lumbreras, J., Borge, R., Narros, A., García-Pérez, J., Fernández-
902 Navarro, P., Pérez-Gómez, B., Aragonés, N., Ramis, R., Pollán, M., Moreno, T.,
903 Karanasiou, A., and López-Abente, G.: Health impact assessment of a reduction in
904 ambient PM_{2.5} levels in Spain, *Environ. Int.*, 37, 342-348,
905 <http://dx.doi.org/10.1016/j.envint.2010.10.004>, 2011.

906 Boreddy, S. K. R., Kawamura, K., and Jung, J.: Hygroscopic properties of particles nebulized
907 from water extracts of aerosols collected at Chichijima Island in the western North

908 Pacific: An outflow region of Asian dust, *Journal of Geophysical Research:*
909 *Atmospheres*, 119, 2013JD020626, 10.1002/2013JD020626, 2014.

910 Brown, S. G., Frankel, A., Raffuse, S. M., Roberts, P. T., Hafner, H. R., and Anderson, D. J.:
911 Source Apportionment of Fine Particulate Matter in Phoenix, AZ, Using Positive
912 Matrix Factorization, *J. Air Waste Manage. Assoc.*, 57, 741-752, 10.3155/1047-
913 3289.57.6.741, 2007.

914 Brown, S. G., Lee, T., Norris, G. A., Roberts, P. T., Collett Jr, J. L., Paatero, P., and
915 Worsnop, D. R.: Receptor modeling of near-roadway aerosol mass spectrometer data
916 in Las Vegas, Nevada, with EPA PMF, *Atmos. Chem. Phys.*, 12, 309-325,
917 10.5194/acp-12-309-2012, 2012.

918 Budhavant, K., Andersson, A., Bosch, C., Kruså, M., Murthaza, A., Zahid, and Gustafsson,
919 Ö.: Apportioned contributions of PM_{2.5} fine aerosol particles over the Maldives
920 (northern Indian Ocean) from local sources vs long-range transport, *Sci. Total.*
921 *Environ.*, 536, 72-78, <http://dx.doi.org/10.1016/j.scitotenv.2015.07.059>, 2015.

922 Cao, S., Duan, X., Zhao, X., Ma, J., Dong, T., Huang, N., Sun, C., He, B., and Wei, F.:
923 Health risks from the exposure of children to As, Se, Pb and other heavy metals near
924 the largest coking plant in China, *Sci. Total. Environ.*, 472, 1001-1009,
925 <http://dx.doi.org/10.1016/j.scitotenv.2013.11.124>, 2014.

926 Chang, D., and Song, Y.: Estimates of biomass burning emissions in tropical Asia based on
927 satellite-derived data, *Atmos. Chem. Phys.*, 10, 2335-2351, 10.5194/acp-10-2335-
928 2010, 2010.

929 Chen, L. W. A., Watson, J. G., Chow, J. C., and Magliano, K. L.: Quantifying PM_{2.5} Source
930 Contributions for the San Joaquin Valley with Multivariate Receptor Models,
931 *Environ. Sci. Technol.*, 41, 2818-2826, 10.1021/es0525105, 2007.

932 Chester, R., Nimmo, M., Fones, G. R., Keyse, S., and Zhang, Z.: Trace metal chemistry of
933 particulate aerosols from the UK mainland coastal rim of the NE Irish sea, *Atmos.*
934 *Environ.*, 34, 949-958, [http://dx.doi.org/10.1016/S1352-2310\(99\)00234-4](http://dx.doi.org/10.1016/S1352-2310(99)00234-4), 2000.

935 Cheung, K., Shafer, M. M., Schauer, J. J., and Sioutas, C.: Historical trends in the mass and
936 chemical species concentrations of coarse particulate matter in the Los Angeles Basin
937 and relation to sources and air quality regulations, *J. Air Waste Manage. Assoc.*, 62,
938 541-556, 10.1080/10962247.2012.661382, 2012.

939 Choi, J.-k., Heo, J.-B., Ban, S.-J., Yi, S.-M., and Zoh, K.-D.: Source apportionment of PM_{2.5}
940 at the coastal area in Korea, *Sci. Total. Environ.*, 447, 370-380,
941 <http://dx.doi.org/10.1016/j.scitotenv.2012.12.047>, 2013.

942 Chueinta, W., Hopke, P. K., and Paatero, P.: Investigation of sources of atmospheric aerosol
943 at urban and suburban residential areas in Thailand by positive matrix factorization,
944 *Atmos. Environ.*, 34, 3319-3329, [http://dx.doi.org/10.1016/S1352-2310\(99\)00433-1](http://dx.doi.org/10.1016/S1352-2310(99)00433-1),
945 2000.

946 Cusack, M., Talbot, N., Ondráček, J., Minguillón, M. C., Martins, V., Klouda, K., Schwarz,
947 J., and Ždímal, V.: Variability of aerosols and chemical composition of PM10, PM2.5
948 and PM1 on a platform of the Prague underground metro, *Atmos. Environ.*, 118, 176-
949 183, <http://dx.doi.org/10.1016/j.atmosenv.2015.08.013>, 2015.

950 Dai, W., Gao, J., Cao, G., and Ouyang, F.: Chemical composition and source identification of
951 PM2.5 in the suburb of Shenzhen, China, *Atmos. Res.*, 122, 391-400,
952 <http://dx.doi.org/10.1016/j.atmosres.2012.12.004>, 2013.

953 Dall'Osto, M., Querol, X., Amato, F., Karanasiou, A., Lucarelli, F., Nava, S., Calzolari, G.,
954 and Chiari, M.: Hourly elemental concentrations in PM2.5 aerosols sampled
955 simultaneously at urban background and road site during SAPUSS – diurnal
956 variations and PMF receptor modelling, *Atmos. Chem. Phys.*, 13, 4375-4392,
957 [10.5194/acp-13-4375-2013](https://doi.org/10.5194/acp-13-4375-2013), 2013.

958 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U.,
959 Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de
960 Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J.,
961 Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P.,
962 Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J. J.,
963 Park, B. K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J. N., and Vitart, F.:
964 The ERA-Interim reanalysis: configuration and performance of the data assimilation
965 system, *Quart. J. Roy. Meteor. Soc.*, 137, 553-597, [10.1002/qj.828](https://doi.org/10.1002/qj.828), 2011.

966 Finlayson-Pitts, B. J., and Pitts Jr., J. N.: *Chemistry of the Upper and Lower Atmosphere:*
967 *Theory, Experiments, and Applications*, Wiley, New York, USA, 2000.

968 Garg, B. D., Cadle, S. H., Mulawa, P. A., Groblicki, P. J., Laroo, C., and Parr, G. A.: Brake
969 Wear Particulate Matter Emissions, *Environ. Sci. Technol.*, 34, 4463-4469,
970 [10.1021/es001108h](https://doi.org/10.1021/es001108h), 2000.

971 Gibson, M. D., Haelssig, J., Pierce, J. R., Parrington, M., Franklin, J. E., Hopper, J. T., Li, Z.,
972 and Ward, T. J.: A comparison of four receptor models used to quantify the boreal
973 wildfire smoke contribution to surface PM2.5 in Halifax, Nova Scotia during the
974 BORTAS-B experiment, *Atmos. Chem. Phys. Discuss.*, 14, 24043-24086,
975 [10.5194/acpd-14-24043-2014](https://doi.org/10.5194/acpd-14-24043-2014), 2014.

976 Gietl, J. K., Lawrence, R., Thorpe, A. J., and Harrison, R. M.: Identification of brake wear
977 particles and derivation of a quantitative tracer for brake dust at a major road, *Atmos.*
978 *Environ.*, 44, 141-146, <http://dx.doi.org/10.1016/j.atmosenv.2009.10.016>, 2010.

979 Gugamsetty, B., Wei, H., Liu, C.-N., Awasthi, A., Hsu, S.-C., Tsai, C.-J., Roam, G.-D., Wu,
980 Y.-C., and Chen, C.-F.: Source characterization and apportionment of PM₁₀, PM_{2.5}
981 and PM_{0.1} by using positive matrix factorization, *Aerosol Air Qual. Res.*, 12, 476-
982 491, [10.4209/aaqr.2012.04.0084](https://doi.org/10.4209/aaqr.2012.04.0084), 2012.

983 Han, J., Moon, K., Lee, S., Kim, Y., Ryu, S., Cliff, S., and Yi, S.: Size-resolved source
984 apportionment of ambient particles by positive matrix factorization at Gosan
985 background site in East Asia, *Atmos. Chem. Phys.*, 6, 211-223, [1680-7324/acp/2006-](https://doi.org/10.19754/atmoschemphys-2006-06-211)
986 [6-211](https://doi.org/10.19754/atmoschemphys-2006-06-211), 2006.

987 Hara, K., Osada, K., Yabuki, M., and Yamanouchi, T.: Seasonal variation of fractionated sea-
988 salt particles on the Antarctic coast, *Geophysical Research Letters*, 39, L18801,
989 [10.1029/2012GL052761](https://doi.org/10.1029/2012GL052761), 2012.

990 Harrison, R. M., Jones, A. M., and Lawrence, R. G.: A pragmatic mass closure model for
991 airborne particulate matter at urban background and roadside sites, *Atmos. Environ.*,
992 37, 4927-4933, <http://dx.doi.org/10.1016/j.atmosenv.2003.08.025>, 2003.

993 Harrison, R. M., Beddows, D. C. S., and Dall'Osto, M.: PMF Analysis of Wide-Range
994 Particle Size Spectra Collected on a Major Highway, *Environ. Sci. Technol.*, 45,
995 5522-5528, [10.1021/es2006622](https://doi.org/10.1021/es2006622), 2011.

996 Hedberg, E., Gidhagen, L., and Johansson, C.: Source contributions to PM₁₀ and arsenic
997 concentrations in Central Chile using positive matrix factorization, *Atmos. Environ.*,
998 39, 549-561, <http://dx.doi.org/10.1016/j.atmosenv.2004.11.001>, 2005.

999 Henry, R. C.: Current factor analysis receptor models are ill-posed, *Atmos. Environ.*, 21,
1000 1815-1820, [http://dx.doi.org/10.1016/0004-6981\(87\)90122-3](http://dx.doi.org/10.1016/0004-6981(87)90122-3), 1987.

1001 Heo, J. B., Hopke, P. K., and Yi, S. M.: Source apportionment of PM_{2.5} in Seoul, Korea,
1002 *Atmos. Chem. Phys.*, 9, 4957-4971, [10.5194/acp-9-4957-2009](https://doi.org/10.5194/acp-9-4957-2009), 2009.

1003 Ho, K. F., Engling, G., Ho, S. S. H., Huang, R., Lai, S., Cao, J., and Lee, S. C.: Seasonal
1004 variations of anhydrosugars in PM_{2.5} in the Pearl River Delta Region, China, 2014,
1005 [10.3402/tellusb.v66.22577](https://doi.org/10.3402/tellusb.v66.22577), 2014.

1006 Hu, X., Zhang, Y., Ding, Z., Wang, T., Lian, H., Sun, Y., and Wu, J.: Bioaccessibility and
1007 health risk of arsenic and heavy metals (Cd, Co, Cr, Cu, Ni, Pb, Zn and Mn) in TSP
1008 and PM_{2.5} in Nanjing, China, *Atmos. Environ.*, 57, 146-152,
1009 <http://dx.doi.org/10.1016/j.atmosenv.2012.04.056>, 2012.

1010 Huang, X. H., Bian, Q., Ng, W. M., Louie, P. K., and Yu, J. Z.: Characterization of PM_{2.5}
1011 major components and source investigation in suburban Hong Kong: A one year
1012 monitoring study, *Aerosol Air Qual. Res.*, 14, 237-250, 10.4209/aaqr.2013.01.0020,
1013 2014.

1014 Joseph, A. E., Unnikrishnan, S., and Kumar, R.: Chemical characterization and mass closure
1015 of fine aerosol for different land use patterns in Mumbai city, *Aerosol Air Qual. Res.*,
1016 12, 61-72, 10.4209/aaqr.2011.04.0049, 2012.

1017 Kanniah, K. D., Lim, H. Q., Kaskaoutis, D. G., and Cracknell, A. P.: Investigating aerosol
1018 properties in Peninsular Malaysia via the synergy of satellite remote sensing and
1019 ground-based measurements, *Atmos. Res.*, 138, 223-239,
1020 <http://dx.doi.org/10.1016/j.atmosres.2013.11.018>, 2014.

1021 Keywood, M. D., Ayers, G. P., Gras, J. L., Boers, C. P., and Leong: Haze in the Klang Valley
1022 of Malaysia, *Atmos. Chem. Phys.*, 3, 591-605, 10.5194/acp-3-591-2003, 2003.

1023 Khan, M. F., Shirasuna, Y., Hirano, K., and Masunaga, S.: Urban and suburban aerosol in
1024 Yokohama, Japan: a comprehensive chemical characterization, *Environ. Monit.*
1025 *Assess.*, 171, 441-456, 10.1007/s10661-009-1290-1, 2010a.

1026 Khan, M. F., Shirasuna, Y., Hirano, K., and Masunaga, S.: Characterization of PM_{2.5},
1027 PM_{2.5-10} and PM_{>10} in ambient air, Yokohama, Japan, *Atmos. Res.*, 96, 159-172,
1028 <http://dx.doi.org/10.1016/j.atmosres.2009.12.009>, 2010b.

1029 Khan, M. F., Hirano, K., and Masunaga, S.: Assessment of the sources of suspended
1030 particulate matter aerosol using US EPA PMF 3.0, *Environ. Monit. Assess.*, 184,
1031 1063-1083, 10.1007/s10661-011-2021-y, 2012.

1032 Khan, M. F., Latif, M. T., Amil, N., Juneng, L., Mohamad, N., Nadzir, M. S. M., and Hoque,
1033 H. M. S.: Characterization and source apportionment of particle number concentration
1034 at a semi-urban tropical environment, *Environ. Sci. Pollut. Res.*, 1-16,
1035 10.1007/s11356-015-4541-4, 2015a.

1036 Khan, M. F., Latif, M. T., Lim, C. H., Amil, N., Jaafar, S. A., Dominick, D., Mohd Nadzir,
1037 M. S., Sahani, M., and Tahir, N. M.: Seasonal effect and source apportionment of
1038 polycyclic aromatic hydrocarbons in PM_{2.5}, *Atmos. Environ.*, 106, 178-190,
1039 <http://dx.doi.org/10.1016/j.atmosenv.2015.01.077>, 2015b.

1040 Khan, M. F., Latif, M. T., Lim, C. H., Amil, N., Jaafar, S. A., Dominick, D., Nadzir, M. S.
1041 M., Sahani, M., and Tahir, N. M.: Seasonal effect and source apportionment of
1042 polycyclic aromatic hydrocarbons in PM_{2.5}, *Atmos. Environ.*, 106, 178-190,
1043 10.1016/j.atmosenv.2015.01.077, 2015c.

1044 Khanna, I., Khare, M., and Gargava, P.: Health Risks Associated with Heavy Metals in Fine
1045 Particulate Matter: A Case Study in Delhi City, India, *J. Geosci. Environ. Protec.*, 3,
1046 72, 10.4236/gep.2015.32012, 2015.

1047 Kholdebarin, A., Biati, A., Moattar, F., and Shariat, S.: Outdoor PM10 source apportionment
1048 in metropolitan cities—a case study, *Environ. Monit. Assess.*, 187, 1-12,
1049 10.1007/s10661-015-4294-z, 2015.

1050 Kim, E., and Hopke, P. K.: Source characterization of ambient fine particles in the Los
1051 Angeles basin, *J. Environ. Eng. Sci.*, 6, 343-353, 10.1139/s06-054, 2007.

1052 Kim Oanh, N. T., Upadhyay, N., Zhuang, Y. H., Hao, Z. P., Murthy, D. V. S., Lestari, P.,
1053 Villarin, J. T., Chengchua, K., Co, H. X., Dung, N. T., and Lindgren, E. S.: Particulate
1054 air pollution in six Asian cities: Spatial and temporal distributions, and associated
1055 sources, *Atmos. Environ.*, 40, 3367-3380,
1056 <http://dx.doi.org/10.1016/j.atmosenv.2006.01.050>, 2006.

1057 Kioumourtzoglou, M.-A., Coull, B. A., Dominici, F., Koutrakis, P., Schwartz, J., and Suh,
1058 H.: The impact of source contribution uncertainty on the effects of source-specific
1059 PM2.5 on hospital admissions: A case study in Boston, MA, *J. Expos. Sci. Environ.
1060 Epidemiol.*, 24, 365-371, 10.1038/jes.2014.7, 2014.

1061 Lee, E., Chan, C. K., and Paatero, P.: Application of positive matrix factorization in source
1062 apportionment of particulate pollutants in Hong Kong, *Atmos. Environ.*, 33, 3201-
1063 3212, [http://dx.doi.org/10.1016/S1352-2310\(99\)00113-2](http://dx.doi.org/10.1016/S1352-2310(99)00113-2), 1999.

1064 Lelieveld, J., Crutzen, P. J., Ramanathan, V., Andreae, M. O., Brenninkmeijer, C. A. M.,
1065 Campos, T., Cass, G. R., Dickerson, R. R., Fischer, H., de Gouw, J. A., Hansel, A.,
1066 Jefferson, A., Kley, D., de Laat, A. T. J., Lal, S., Lawrence, M. G., Lobert, J. M.,
1067 Mayol-Bracero, O. L., Mitra, A. P., Novakov, T., Oltmans, S. J., Prather, K. A.,
1068 Reiner, T., Rodhe, H., Scheeren, H. A., Sikka, D., and Williams, J.: The Indian Ocean
1069 Experiment: Widespread Air Pollution from South and Southeast Asia, *Science*, 291,
1070 1031-1036, 10.1126/science.1057103, 2001.

1071 Lestari, P., and Mauliadi, Y. D.: Source apportionment of particulate matter at urban mixed
1072 site in Indonesia using PMF, *Atmos. Environ.*, 43, 1760-1770,
1073 <http://dx.doi.org/10.1016/j.atmosenv.2008.12.044>, 2009.

1074 Li, H., Wang, J., Wang, Q. g., Qian, X., Qian, Y., Yang, M., Li, F., Lu, H., and Wang, C.:
1075 Chemical fractionation of arsenic and heavy metals in fine particle matter and its
1076 implications for risk assessment: A case study in Nanjing, China, *Atmos. Environ.*,
1077 103, 339-346, <http://dx.doi.org/10.1016/j.atmosenv.2014.12.065>, 2015a.

- 1078 Li, Y., Zhang, Z., Liu, H., Zhou, H., Fan, Z., Lin, M., Wu, D., and Xia, B.: Characteristics,
1079 sources and health risk assessment of toxic heavy metals in PM_{2.5} at a megacity of
1080 southwest China, *Environ. Geochem. Health*, 1-10, 10.1007/s10653-015-9722-z,
1081 2015b.
- 1082 Maenhaut, W., Raes, N., Chi, X., Cafmeyer, J., and Wang, W.: Chemical composition and
1083 mass closure for PM_{2.5} and PM₁₀ aerosols at K-puszta, Hungary, in summer 2006,
1084 *X-Ray Spectrom.*, 37, 193-197, 10.1002/xrs.1062, 2008.
- 1085 Mariani, R. L., and de Mello, W. Z.: PM_{2.5-10}, PM_{2.5} and associated water-soluble
1086 inorganic species at a coastal urban site in the metropolitan region of Rio de Janeiro,
1087 *Atmos. Environ.*, 41, 2887-2892, <http://dx.doi.org/10.1016/j.atmosenv.2006.12.009>,
1088 2007.
- 1089 Martins, V., Moreno, T., Minguillón, M. C., van Drooge, B. L., Reche, C., Amato, F., de
1090 Miguel, E., Capdevila, M., Centelles, S., and Querol, X.: Origin of inorganic and
1091 organic components of PM_{2.5} in subway stations of Barcelona, Spain, *Environmental*
1092 *Pollution*, 208, Part A, 125-136, <http://dx.doi.org/10.1016/j.envpol.2015.07.004>,
1093 2016.
- 1094 McGinnis, J. E., Heo, J., Olson, M. R., Rutter, A. P., and Schauer, J. J.: Understanding the
1095 sources and composition of the incremental excess of fine particles across multiple
1096 sampling locations in one air shed, *J. Environ. Sci.*, 26, 818-826, 10.1016/S1001-
1097 0742(13)60508-2, 2014.
- 1098 Meij, R., and te Winkel, H.: The emissions of heavy metals and persistent organic pollutants
1099 from modern coal-fired power stations, *Atmos. Environ.*, 41, 9262-9272,
1100 <http://dx.doi.org/10.1016/j.atmosenv.2007.04.042>, 2007.
- 1101 Mohd Tahir, N., Suratman, S., Fong, F. T., Hamzah, M. S., and Latif, M. T.: Temporal
1102 distribution and chemical characterization of atmospheric particulate matter in the
1103 eastern coast of Peninsular Malaysia, *Aerosol Air Qual. Res.*, 13, 584-595,
1104 10.4209/aaqr.2012.08.0216, 2013.
- 1105 Morales, R., and Leiva, G.: Distribution and Critical Concentration of PM in the city of
1106 Santiago, Chile (in Spanish), in: Morales RGE (ed) *Atmospheric Urban Pollution:*
1107 *Critical episodes of the Environmental pollution in the City of Santiago, Chile* (in
1108 Spanish), 1st edn. Editorial Universitaria SA, Santiago, 2006.
- 1109 Moreno, T., Karanasiou, A., Amato, F., Lucarelli, F., Nava, S., Calzolari, G., Chiari, M., Coz,
1110 E., Artíñano, B., Lumbreras, J., Borge, R., Boldo, E., Linares, C., Alastuey, A.,
1111 Querol, X., and Gibbons, W.: Daily and hourly sourcing of metallic and mineral dust

1112 in urban air contaminated by traffic and coal-burning emissions, *Atmos. Environ.*, 68,
1113 33-44, <http://dx.doi.org/10.1016/j.atmosenv.2012.11.037>, 2013.

1114 Mustaffa, N., Latif, M., Ali, M., and Khan, M.: Source apportionment of surfactants in
1115 marine aerosols at different locations along the Malacca Straits, *Environ. Sci. Pollut.*
1116 *Res.*, 21, 6590-6602, [10.1007/s11356-014-2562-z](https://doi.org/10.1007/s11356-014-2562-z), 2014.

1117 Niu, L., Ye, H., Xu, C., Yao, Y., and Liu, W.: Highly time- and size-resolved fingerprint
1118 analysis and risk assessment of airborne elements in a megacity in the Yangtze River
1119 Delta, China, *Chemosphere*, 119, 112-121,
1120 <http://dx.doi.org/10.1016/j.chemosphere.2014.05.062>, 2015.

1121 Norris, G., Duvall, R., Brown, S., and Bai, S.: EPA positive matrix factorization (PMF) 5.0
1122 fundamentals & user guide, Prepared for the US Environmental Protection Agency,
1123 Washington, DC, by the National Exposure Research Laboratory, Research Triangle
1124 Park, 2014.

1125 Ogulei, D., Hopke, P. K., and Wallace, L. A.: Analysis of indoor particle size distributions in
1126 an occupied townhouse using positive matrix factorization, *Indoor Air*, 16, 204-215,
1127 [10.1111/j.1600-0668.2006.00418.x](https://doi.org/10.1111/j.1600-0668.2006.00418.x), 2006a.

1128 Ogulei, D., Hopke, P. K., Zhou, L., Patrick Pancras, J., Nair, N., and Ondov, J. M.: Source
1129 apportionment of Baltimore aerosol from combined size distribution and chemical
1130 composition data, *Atmos. Environ.*, 40, Supplement 2, 396-410,
1131 <http://dx.doi.org/10.1016/j.atmosenv.2005.11.075>, 2006b.

1132 Paatero, P., and Tapper, U.: Positive matrix factorization: A non-negative factor model with
1133 optimal utilization of error estimates of data values, *Environmetrics*, 5, 111-126,
1134 [10.1002/env.3170050203](https://doi.org/10.1002/env.3170050203), 1994.

1135 Paatero, P.: Least squares formulation of robust non-negative factor analysis, *Chemometr.*
1136 *Intell. Lab.*, 37, 23-35, [http://dx.doi.org/10.1016/S0169-7439\(96\)00044-5](http://dx.doi.org/10.1016/S0169-7439(96)00044-5), 1997.

1137 Pandey, P., Patel, D. K., Khan, A. H., Barman, S. C., Murthy, R. C., and Kisku, G. C.:
1138 Temporal distribution of fine particulates (PM_{2.5}, PM₁₀), potentially toxic metals,
1139 PAHs and Metal-bound carcinogenic risk in the population of Lucknow City, India, *J.*
1140 *Environ. Sci. Health., Part A*, 48, 730-745, [10.1080/10934529.2013.744613](https://doi.org/10.1080/10934529.2013.744613), 2013.

1141 Polissar, A. V., Hopke, P. K., Malm, W. C., and Sisler, J. F.: Atmospheric aerosol over
1142 Alaska: 1. Spatial and seasonal variability, *J. Geophys. Res.*, 103, 19035-19044,
1143 [10.1029/98JD01365](https://doi.org/10.1029/98JD01365), 1998a.

1144 Polissar, A. V., Hopke, P. K., Paatero, P., Malm, W. C., and Sisler, J. F.: Atmospheric
1145 aerosol over Alaska: 2. Elemental composition and sources, *J. Geophys. Res.*, 103,
1146 19045-19057, [10.1029/98JD01212](https://doi.org/10.1029/98JD01212), 1998b.

1147 Querol, X., Fernández-Turiel, J., and López-Soler, A.: Trace elements in coal and their
1148 behaviour during combustion in a large power station, *Fuel*, 74, 331-343,
1149 [http://dx.doi.org/10.1016/0016-2361\(95\)93464-O](http://dx.doi.org/10.1016/0016-2361(95)93464-O), 1995.

1150 Rahman, S. A., Hamzah, M. S., Wood, A. K., Elias, M. S., Salim, A., Ashifa, N., and Sanuri,
1151 E.: Sources apportionment of fine and coarse aerosol in Klang Valley, Kuala Lumpur
1152 using positive matrix factorization, *Atmos. Pollut. Res.*, 2, 197-206,
1153 [10.5094/APR.2011.025](https://doi.org/10.5094/APR.2011.025), 2011.

1154 Reid, J. S., Hyer, E. J., Johnson, R. S., Holben, B. N., Yokelson, R. J., Zhang, J., Campbell, J.
1155 R., Christopher, S. A., Di Girolamo, L., Giglio, L., Holz, R. E., Kearney, C.,
1156 Miettinen, J., Reid, E. A., Turk, F. J., Wang, J., Xian, P., Zhao, G., Balasubramanian,
1157 R., Chew, B. N., Janjai, S., Lagrosas, N., Lestari, P., Lin, N.-H., Mahmud, M.,
1158 Nguyen, A. X., Norris, B., Oanh, N. T. K., Oo, M., Salinas, S. V., Welton, E. J., and
1159 Liew, S. C.: Observing and understanding the Southeast Asian aerosol system by
1160 remote sensing: An initial review and analysis for the Seven Southeast Asian Studies
1161 (7SEAS) program, *Atmos. Res.*, 122, 403-468,
1162 <http://dx.doi.org/10.1016/j.atmosres.2012.06.005>, 2013.

1163 Remoundaki, E., Kassomenos, P., Mantas, E., Mihalopoulos, N., and Tsezos, M.:
1164 Composition and mass closure of PM_{2.5} in urban environment (Athens, Greece),
1165 *Aerosol Air Qual. Res.*, 13, 72-82, [10.4209/aaqr.2012.03.0054](https://doi.org/10.4209/aaqr.2012.03.0054), 2013.

1166 Sahani, M., Zainon, N. A., Mahiyuddin, W. R. W., Latif, M. T., Hod, R., Khan, M. F., Tahir,
1167 N. M., and Chan, C.-C.: A case-crossover analysis of forest fire haze events and
1168 mortality in Malaysia, *Atmos. Environ.*, 96, 257-265,
1169 [10.1016/j.atmosenv.2014.07.043](https://doi.org/10.1016/j.atmosenv.2014.07.043), 2014.

1170 Sánchez-Soberón, F., Rovira, J., Mari, M., Sierra, J., Nadal, M., Domingo, J. L., and
1171 Schuhmacher, M.: Main components and human health risks assessment of PM₁₀,
1172 PM_{2.5}, and PM₁ in two areas influenced by cement plants, *Atmos. Environ.*, 120,
1173 109-116, <http://dx.doi.org/10.1016/j.atmosenv.2015.08.020>, 2015.

1174 Santoso, M., Hopke, P. K., Hidayat, A., and Diah Dwiana, L.: Sources identification of the
1175 atmospheric aerosol at urban and suburban sites in Indonesia by positive matrix
1176 factorization, *Sci. Total. Environ.*, 397, 229-237,
1177 <http://dx.doi.org/10.1016/j.scitotenv.2008.01.057>, 2008.

1178 Santoso, M., Lestiani, D. D., and Markwitz, A.: Characterization of airborne particulate
1179 matter collected at Jakarta roadside of an arterial road, *J. Radioanal. Nucl. Chem.*,
1180 297, 165-169, 10.1007/s10967-012-2350-5, 2013.

1181 Satsangi, P. G., Yadav, S., Pipal, A. S., and Kumbhar, N.: Characteristics of trace metals in
1182 fine (PM_{2.5}) and inhalable (PM₁₀) particles and its health risk assessment along with
1183 in-silico approach in indoor environment of India, *Atmos. Environ.*, 92, 384-393,
1184 <http://dx.doi.org/10.1016/j.atmosenv.2014.04.047>, 2014.

1185 Seinfeld, J. H., and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to*
1186 *climate change*, John Wiley & Sons, 2012.

1187 Seneviratne, M., Waduge, V. A., Hadagiripathira, L., Sanjeevani, S., Attanayake, T.,
1188 Jayaratne, N., and Hopke, P. K.: Characterization and source apportionment of
1189 particulate pollution in Colombo, Sri Lanka, *Atmos. Pollut. Res.*, 2,
1190 10.5094/APR.2011.026, 2011.

1191 Sillanpää, M., Hillamo, R., Saarikoski, S., Frey, A., Pennanen, A., Makkonen, U., Spolnik,
1192 Z., Van Grieken, R., Braniš, M., Brunekreef, B., Chalbot, M.-C., Kuhlbusch, T.,
1193 Sunyer, J., Kerminen, V.-M., Kulmala, M., and Salonen, R. O.: Chemical
1194 composition and mass closure of particulate matter at six urban sites in Europe,
1195 *Atmos. Environ.*, 40, Supplement 2, 212-223,
1196 <http://dx.doi.org/10.1016/j.atmosenv.2006.01.063>, 2006.

1197 Song, Y., Xie, S., Zhang, Y., Zeng, L., Salmon, L. G., and Zheng, M.: Source apportionment
1198 of PM_{2.5} in Beijing using principal component analysis/absolute principal component
1199 scores and UNMIX, *Sci. Total. Environ.*, 372, 278-286,
1200 <http://dx.doi.org/10.1016/j.scitotenv.2006.08.041>, 2006.

1201 Srimuruganandam, B., and Shiva Nagendra, S. M.: Application of positive matrix
1202 factorization in characterization of PM₁₀ and PM_{2.5} emission sources at urban
1203 roadside, *Chemosphere*, 88, 120-130,
1204 <http://dx.doi.org/10.1016/j.chemosphere.2012.02.083>, 2012.

1205 Streets, D., Yarber, K., Woo, J. H., and Carmichael, G.: Biomass burning in Asia: Annual
1206 and seasonal estimates and atmospheric emissions, *Global Biogeochem. Cy.*, 17,
1207 1099, 10.1029/2003GB002040, 2003, 2003.

1208 Sun, Y., Zhuang, G., Tang, A., Wang, Y., and An, Z.: Chemical Characteristics of PM_{2.5} and
1209 PM₁₀ in Haze–Fog Episodes in Beijing, *Environ. Sci. Technol.*, 40, 3148-3155,
1210 10.1021/es051533g, 2006.

1211 Taner, S., Pekey, B., and Pekey, H.: Fine particulate matter in the indoor air of barbeque
1212 restaurants: Elemental compositions, sources and health risks, *Sci. Total. Environ.*,
1213 454–455, 79-87, <http://dx.doi.org/10.1016/j.scitotenv.2013.03.018>, 2013.

1214 Tao, J., Zhang, L., Engling, G., Zhang, R., Yang, Y., Cao, J., Zhu, C., Wang, Q., and Luo, L.:
1215 Chemical composition of PM_{2.5} in an urban environment in Chengdu, China:
1216 Importance of springtime dust storms and biomass burning, *Atmos. Res.*, 122, 270-
1217 283, <http://dx.doi.org/10.1016/j.atmosres.2012.11.004>, 2013.

1218 Tao, J., Gao, J., Zhang, L., Zhang, R., Che, H., Zhang, Z., Lin, Z., Jing, J., Cao, J., and Hsu,
1219 S. C.: PM_{2.5} pollution in a megacity of southwest China: source apportionment and
1220 implication, *Atmos. Chem. Phys.*, 14, 8679-8699, 10.5194/acp-14-8679-2014, 2014.

1221 Taylor, S. R.: Abundance of chemical elements in the continental crust: a new table,
1222 *Geochim. Cosmochim. Ac.*, 28, 1273-1285, 1964.

1223 Terzi, E., Argyropoulos, G., Bougatioti, A., Mihalopoulos, N., Nikolaou, K., and Samara, C.:
1224 Chemical composition and mass closure of ambient PM₁₀ at urban sites, *Atmos.*
1225 *Environ.*, 44, 2231-2239, <http://dx.doi.org/10.1016/j.atmosenv.2010.02.019>, 2010.

1226 Thurston, G. D., and Spengler, J. D.: A quantitative assessment of source contributions to
1227 inhalable particulate matter pollution in metropolitan Boston, *Atmos. Environ.*, 19, 9-
1228 25, [http://dx.doi.org/10.1016/0004-6981\(85\)90132-5](http://dx.doi.org/10.1016/0004-6981(85)90132-5), 1985.

1229 Torfs, K., and Van Grieken, R.: Chemical relations between atmospheric aerosols, deposition
1230 and stone decay layers on historic buildings at the mediterranean coast, *Atmos.*
1231 *Environ.*, 31, 2179-2192, [http://dx.doi.org/10.1016/S1352-2310\(97\)00038-1](http://dx.doi.org/10.1016/S1352-2310(97)00038-1), 1997.

1232 Vedantham, R., Landis, M. S., Olson, D., and Pancras, J. P.: Source Identification of PM_{2.5}
1233 in Steubenville, Ohio Using a Hybrid Method for Highly Time-Resolved Data,
1234 *Environ. Sci. Technol.*, 48, 1718-1726, 10.1021/es402704n, 2014.

1235 Vejehati, F., Xu, Z., and Gupta, R.: Trace elements in coal: Associations with coal and
1236 minerals and their behavior during coal utilization – A review, *Fuel*, 89, 904-911,
1237 <http://dx.doi.org/10.1016/j.fuel.2009.06.013>, 2010.

1238 Viana, M., Kuhlbusch, T. A. J., Querol, X., Alastuey, A., Harrison, R. M., Hopke, P. K.,
1239 Winiwarter, W., Vallius, M., Szidat, S., Prévôt, A. S. H., Hueglin, C., Bloemen, H.,
1240 Wählin, P., Vecchi, R., Miranda, A. I., Kasper-Giebl, A., Maenhaut, W., and
1241 Hitzenberger, R.: Source apportionment of particulate matter in Europe: A review of
1242 methods and results, *J. Aerosol Sci.*, 39, 827-849,
1243 <http://dx.doi.org/10.1016/j.jaerosci.2008.05.007>, 2008.

- 1244 Wahid, N. B. A., Latif, M. T., and Suratman, S.: Composition and source apportionment of
1245 surfactants in atmospheric aerosols of urban and semi-urban areas in Malaysia,
1246 *Chemosphere*, 91, 1508-1516, <http://dx.doi.org/10.1016/j.chemosphere.2012.12.029>,
1247 2013.
- 1248 Wåhlin, P., Berkowicz, R., and Palmgren, F.: Characterisation of traffic-generated particulate
1249 matter in Copenhagen, *Atmos. Environ.*, 40, 2151-2159,
1250 <http://dx.doi.org/10.1016/j.atmosenv.2005.11.049>, 2006.
- 1251 Waked, A., Favez, O., Alleman, L. Y., Piot, C., Petit, J. E., Delaunay, T., Verlinden, E.,
1252 Golly, B., Besombes, J. L., Jaffrezo, J. L., and Leoz-Garziandia, E.: Source
1253 apportionment of PM10 in a north-western Europe regional urban background site
1254 (Lens, France) using positive matrix factorization and including primary biogenic
1255 emissions, *Atmos. Chem. Phys.*, 14, 3325-3346, 10.5194/acp-14-3325-2014, 2014.
- 1256 Wang, Y., and Hopke, P. K.: A ten-year source apportionment study of ambient fine
1257 particulate matter in San Jose, California, *Atmos. Pollut. Res.*, 4, 398-404,
1258 10.5094/APR.2013.045, 2013.
- 1259 Watson, J. G., Robinson, N. F., Chow, J. C., Henry, R. C., Kim, B. M., Pace, T. G., Meyer, E.
1260 L., and Nguyen, Q.: The USEPA/DRI chemical mass balance receptor model, CMB
1261 7.0, *Environ. Softw.*, 5, 38-49, [http://dx.doi.org/10.1016/0266-9838\(90\)90015-X](http://dx.doi.org/10.1016/0266-9838(90)90015-X),
1262 1990.
- 1263 Wilson, T. R.: Salinity and the major elements of sea-water, in *Chemical Oceanography*,
1264 edited by J. P. Riley and G. Skirrow, pp. 365–413, Academic, San Diego, Calif.,
1265 1975.
- 1266 Wu, C.-f., Wu, S.-y., Wu, Y.-H., Cullen, A. C., Larson, T. V., Williamson, J., and Liu, L. J.
1267 S.: Cancer risk assessment of selected hazardous air pollutants in Seattle, *Environ.*
1268 *Int.*, 35, 516-522, <http://dx.doi.org/10.1016/j.envint.2008.09.009>, 2009.
- 1269 Wu, Y.-S., Fang, G.-C., Lee, W.-J., Lee, J.-F., Chang, C.-C., and Lee, C.-Z.: A review of
1270 atmospheric fine particulate matter and its associated trace metal pollutants in Asian
1271 countries during the period 1995–2005, *J. Hazard. Mater.*, 143, 511-515,
1272 <http://dx.doi.org/10.1016/j.jhazmat.2006.09.066>, 2007.
- 1273 Yang, W., Zhang, S., Tang, J., Bu, K., Yang, J., and Chang, L.: A MODIS time series data
1274 based algorithm for mapping forest fire burned area, *Chin. Geogr. Sci.*, 23, 344-352,
1275 10.1007/s11769-013-0597-6, 2013.

1276 Yin, J., and Harrison, R. M.: Pragmatic mass closure study for PM1.0, PM2.5 and PM10 at
1277 roadside, urban background and rural sites, *Atmos. Environ.*, 42, 980-988,
1278 <http://dx.doi.org/10.1016/j.atmosenv.2007.10.005>, 2008.

1279 Yu, L., Wang, G., Zhang, R., Zhang, L., Song, Y., Wu, B., Li, X., An, K., and Chu, J.:
1280 Characterization and source apportionment of PM2.5 in an urban environment in
1281 Beijing, *Aerosol Air Qual. Res.*, 13, 574-583, 2013.

1282 Zhang, H., Li, J., Ying, Q., Yu, J. Z., Wu, D., Cheng, Y., He, K., and Jiang, J.: Source
1283 apportionment of PM2.5 nitrate and sulfate in China using a source-oriented chemical
1284 transport model, *Atmos. Environ.*, 62, 228-242,
1285 <http://dx.doi.org/10.1016/j.atmosenv.2012.08.014>, 2012.

1286 Zhang, N., Han, B., He, F., Xu, J., Niu, C., Zhou, J., Kong, S., Bai, Z., and Xu, H.:
1287 Characterization, health risk of heavy metals, and source apportionment of
1288 atmospheric PM2.5 to children in summer and winter: an exposure panel study in
1289 Tianjin, China, *Air Qual. Atmos. Health*, 1-11, 10.1007/s11869-014-0289-0, 2014.

1290 Zhang, R., Jing, J., Tao, J., Hsu, S. C., Wang, G., Cao, J., Lee, C. S. L., Zhu, L., Chen, Z.,
1291 Zhao, Y., and Shen, Z.: Chemical characterization and source apportionment of
1292 PM2.5 in Beijing: seasonal perspective, *Atmos. Chem. Phys.*, 13, 7053-7074,
1293 10.5194/acp-13-7053-2013, 2013.

1294 Zhang, X., Hecobian, A., Zheng, M., Frank, N. H., and Weber, R. J.: Biomass burning impact
1295 on PM 2.5 over the southeastern US during 2007: integrating chemically speciated
1296 FRM filter measurements, MODIS fire counts and PMF analysis, *Atmos. Chem.*
1297 *Phys.*, 10, 6839-6853, 10.5194/acp-10-6839-2010, 2010.

1298 Zhang, Z., Gao, J., Engling, G., Tao, J., Chai, F., Zhang, L., Zhang, R., Sang, X., Chan, C.-y.,
1299 Lin, Z., and Cao, J.: Characteristics and applications of size-segregated biomass
1300 burning tracers in China's Pearl River Delta region, *Atmos. Environ.*, 102, 290-301,
1301 <http://dx.doi.org/10.1016/j.atmosenv.2014.12.009>, 2015.

Table 1 The statistical parameters of the PM_{2.5} and its compositions

Species	Overall (n=27)		SW ^b (n=9)		NE ^c (n=18)		MDL ^d	Recovery (%)
	Mean±SD ^a	Range	Mean±SD	Range	Mean±SD	Range		
Al	267.6±145.0	98.48-826.6	203.1±118.42	98.48-416.09	299.8±149.2	101.9-826.6	0.70	70 (54-97)
Ba	1660±1501	319.2-6092	1372±1480	319.2-5187	1804±1532	447.6-6092	0.02	-
Ca	1770±725.9	n.d.-3150	1584±325.5	1234-2154	1975±683.4	882.1-3150	2.88	33 (23-47)
Fe	3052±654.6	2171-4567	2513±239.6	2171-2893	3322±630.4	2530-4567	0.40	80 (69-95)
Mg	207.6±83.85	34.43-371.7	207.1±72.85	119.0-356.0	207.9±90.86	34.43-371.7	0.17	95 (81-111)
Pb	21.84±16.30	3.57-76.17	28.06±20.27	13.1-76.17	18.72±13.49	3.57-51.70	0.01	119 (89-134)
Zn	389.2±179.8	178.8-817.9	526.8±236.3	178.8-817.9	320.4±90.25	184.7-448.2	0.22	102 (84-129)
Ag	0.09±0.05	n.d.-0.21	0.05±0.04	0.01-0.11	0.10±0.05	0.02-0.21	0.01	120 (97-170)
As	5.76±4.74	1.10-18.33	5.22±2.93	1.55-9.79	6.04±5.49	1.10-18.33	0.45	88 (81-94)
Cd	0.54±0.29	0.13-1.15	0.44±0.22	0.13-0.81	0.58±0.32	0.17-1.15	<0.01	85 (81-89)
Cr	107.68±18.57	82.32-152.62	91.06±7.52	82.32-104.4	115.9±16.78	91.17-152.6	0.02	56 (31-87)
Li	0.22±0.12	0.04-0.43	0.11±0.06	0.04-0.21	0.28±0.10	0.07-0.43	0.09	-
Be	0.01±0.01	n.d.-0.03	0.003±0.01	n.d.-0.01	0.01±0.01	n.d.-0.03	<0.01	-
Bi	0.76±0.60	0.08-2.08	0.67±0.35	0.13-1.17	0.80±0.70	0.08-2.08	0.03	-
Co	0.85±0.47	0.39-2.36	1.16±0.61	0.39-2.36	0.70±0.30	0.39-1.38	0.08	96 (87-109)
Cu	28.33±11.02	16.83-62.55	32.39±10.08	19.78-49.27	26.30±11.17	16.83-62.55	0.30	101 (96-105)
Mn	4.03±1.91	0.23-7.18	3.13±2.07	0.23-6.08	4.49±1.71	1.46-7.18	0.95	126 (114-147)
Ni	17.24±8.55	7.86-46.70	23.59±11.11	7.86-46.70	14.06±4.66	8.84-27.03	0.67	91 (82-99)
Rb	3.59±1.08	1.74-6.16	4.14±1.29	2.23-6.16	3.32±0.87	1.74-4.69	0.13	78 (52-113)
Se	0.65±0.33	0.20-1.24	0.36±0.10	0.20-0.53	0.79±0.31	0.39-1.24	0.09	94 (78-110)
Sr	40.25±31.05	13.75-120.93	35.88±32.10	13.75-118.47	42.43±31.22	15.72-120.9	0.38	91 (75-125)
V	5.13±3.05	0.63-13.16	3.70±2.47	0.63-7.82	5.85±3.12	2.21-13.16	<0.01	85 (77-93)
Na ⁺	532.1±262.0	n.d.-1029.07	363.9±185.6	159.9-778.8	606.90	23.66-1029.1	62.68	-
NH ₄ ⁺	598.9±399.2	82.60-1622.17	542.5±320.8	82.60-1141.4	627.2±439.0	105.5-1622.2	-	-
K ⁺	343.3±183.2	70.18-696.04	307.8±103.5	175.6-484.6	361.1±212.7	70.18-696.0	2.35	-
Ca ²⁺	255.9±84.22	87.55-455.55	295.1±95.8	186.4-455.6	236.3±72.84	87.55-360.4	23.21	-
Mg ²⁺	42.26±17.57	12.70-77.60	32.61±18.32	12.70-71.94	47.09±15.49	15.65-77.60	23.71	-
Cl ⁻	56.71±44.94	4.67-151.18	67.63±24.21	40.07-107.18	51.25±52.13	4.67-151.2	0.98	-
NO ₃ ⁻	926.9±1031.8	98.66-3523.7	194.8±73.63	98.66-311.3	1293±1095	136.5-3524	16.51	-
SO ₄ ²⁺	2127±2068	n.d.-6211	n.d.	n.d.	2127±2068	350.5-6211	1.82	-
^e PM _{2.5}	25.13±9.21	7.01-42.28	22.16±9.14	7.01-35.73	26.61±9.14	12.76-42.28	-	-

^aSD: standard deviation, ^bSW: south-westerly monsoon, ^cNE: north-westerly monsoon, ^dMDL: method detection limit, ^ePM_{2.5} (μg m⁻³), n.d.: not detected, "-": no

Table 2 The contribution of sources to PM_{2.5} and the compositions estimated by PMF 5.0

Variables	Mineral/road dust		Motor vehicle emissions/		Nitrate aerosol		Coal burning		Marine/sulfate aerosol	
	ng m ⁻³	%	ng m ⁻³	%	ng m ⁻³	%	ng m ⁻³	%	ng m ⁻³	%
PM _{2.5}	3.17±0.15**	13±1	7.47±1.26**	31±5	4.11±0.47**	17±2	4.60±0.37**	19±2	4.99±0.67**	20±3
Al	42.65±3.17	19±1	45.37±3.85	20±2	69.06±2.45	31±1	29.84±1.73	13±1	36.71±2.51	16±1
Ba	269.3±205.9	22±17	32.85±146.9	3±14	166.9±71.90	13±6	661.7±246.9	52±19	117.8±116.8	10±11
Ca	445.1±32.07	28±2	235.43±37.76	15±2	350.6±35.82	22±2	303.4±30.14	19±2	267.1±26.00	17±2
Mg	92.36±5.02	52±3	47.59±21.66	27±12	25.43±12.33	14±7	10.32±6.11	6±3	1.23±5.50	1±3
Pb	3.56±0.79	20±4	9.11±2.32	50±13	0.58±0.40	3±2	3.61±0.42	20±2	1.25±1.36	7±8
Zn	157.7±17.09	48±5	45.66±30.11	14±9	60.74±21.74	18±7	50.56±19.46	15±6	14.33±8.53	4±3
As	0.18±.35	4±7	1.76±0.55	41±14	0.05±0.10	1±2	2.37±0.65	53±13	0.05±0.22	1±6
Cd	0.03±0.01	6±2	0.22±0.06	44±12	0.07±0.02	13±3	0.13±0.02	27±3	0.05±0.02	10±5
Cu	12.38±0.59	50±2	3.55±2.37	14±10	4.20±1.45	17±6	3.27±1.16	13±5	1.45±0.42	6±2
Mn	-	-	0.84±0.27	25±8	1.16±0.19	35±6	0.62±0.26	18±7	0.71±0.09	21±3
Ni	7.21±0.50	48±4	2.79±1.18	18±8	1.70±0.77	11±5	3.11±0.80	20±5	0.36±0.23	2±2
Rb	1.33±0.07	38±2	0.76±0.20	22±6	0.45±0.14	13±4	0.67±0.10	19±3	0.26±0.04	7±1
Se	0.05±0.01	8±2	0.14±0.03	24±6	0.14±0.02	23±3	0.11±0.02	19±3	0.16±0.01	27±1
Sr	8.26±4.51	25±14	1.19±3.18	4±11	4.60±1.80	14±6	15.05±5.05	45±15	3.59±2.40	11±9
V	0.19±0.08	5±2	0.81±0.24	20±6	1.25±0.20	30±6	0.59±0.32	14±7	1.28±0.17	31±4
Na ⁺	88.10±28.60	19±6	17.28±56.76	4±12	120.8±10.99	26±3	7.93±4.69	2±1	234.2±20.31	50±5
Ammonium	59.48±30.60	11±6	241.1±61.51	44±11	82.56±18.67	15±4	8.55±16.10	2±3	156.2±48.24	28±8
K ⁺	65.10±18.20	20±6	91.08±16.94	28±5	50.69±6.14	16±2	9.53±3.42	3±1	108.4±16.41	33±5
Ca ²⁺	99.79±3.69	42±1	50.52±18.74	21±8	47.25±9.79	20±4	12.39±6.67	5±3	26.39±4.03	11±2
Mg ²⁺	8.18±1.46	23±4	6.96±1.06	19±3	9.27±0.31	26±1	1.92±0.38	5±1	9.72±0.32	27±1
Cl ⁻	15.88±4.06	36±10	1.83±2.95	4±8	-	-	5.90±0.73	13±2	20.58±6.45	46±13
Nitrate	90.86±36.16	11±4	6.66±21.39	1±2	611.0±27.43	75±3	5.75±16.39	1±2	103.4±53.25	13±7
Sulfate	307.2±142.1	21±10	58.02±152.1	4±11	74.23±33.91	5±2	89.77±10.15	6±1	935.1±112.5	64±7

**unit: µg m⁻³, SD^a: standard deviation, “-“: no data

Table 3. Hazard quotient (HQ) or non-carcinogenic risk and lifetime carcinogenic risk (LCR) for selected heavy metals in PM_{2.5} based on their various sources

Inhalation	Mineral/road dust		Motor vehicle/biomass		Coal burning		PM _{2.5}	
	HQ	LCR	HQ	LCR	HQ	LCR	HQ	LCR
Pb	-	4.0×10 ⁻⁸	-	1.0×10 ⁻⁷	-	4.1×10 ⁻⁸	-	2.5×10 ⁻⁷
As	1.8×10 ⁻³	1.1×10 ⁻⁷	4.9×10 ⁻²	1.1×10 ⁻⁶	6.6×10 ⁻²	1.5×10 ⁻⁶	15.9×10 ⁻²	3.5×10 ⁻⁶
Cd	4.6×10 ⁻⁴	8.2×10 ⁻⁹	9.1×10 ⁻³	5.6×10 ⁻⁸	5.5×10 ⁻³	3.4×10 ⁻⁸	2.2×10 ⁻²	1.4×10 ⁻⁷
Cu	-	-	-	-	-	-	-	-
Mn	-	-	7.0×10 ⁻³	-	5.1×10 ⁻³	-	3.3×10 ⁻²	-
Zn	-	-	-	-	-	-	-	-
Ni	2.0×10 ⁻²	2.5×10 ⁻⁷	2.3×10 ⁻²	9.5×10 ⁻⁸	2.6×10 ⁻²	1.1×10 ⁻⁷	14.3×10 ⁻²	5.9×10 ⁻⁷
THR (HI and LCR)	2.3×10 ⁻²	1.6×10 ⁻⁷	8.8×10 ⁻²	1.2×10 ⁻⁶	10.2×10 ⁻²	1.5×10 ⁻⁶	35.7×10 ⁻²	3.9×10 ⁻⁶

Pb*: pb (acetate), As*: As (Inorganic), Cd*: Cd (Diet), Mn*: Mn (Diet), Zn*: Zn (Metallic), Ni*: Ni (Refinery Dust), "-": no data, **THR: Total Health Risk, HI: Hazard Index, LCR: Lifetime Cancer Risk**




Figure 1 Map of the study area showing the sampling site and nearby line sources




Figure 2 Monthly climatology wind vector from January 2004 to June 2014




Figure 3 The location of biomass fire hotspots and the monthly mean cluster of backward trajectories by HYSPLIT 4.9 model for 120 h and 500 m releasing height starting from 16:00 UTC during the south-westerly and the north-easterly monsoon




Figure 4 Time series of 24 h averages of PM_{2.5}, wind speed (m s⁻¹) and yearly daily mean of the boundary layer height (BLH) over the region of Malaysian Peninsula





Figure 5 (a) Enrichment factor (EF) of heavy metals in PM_{2.5}, (b) correlation plot of nss-K⁺ and total-K⁺, (c) reconstructed mass concentration of PM_{2.5} by mass closure model, (d) correlation plot of K⁺ and Na⁺, and (e) correlation plot of estimated PM_{2.5} (MC) and measured PM_{2.5} (HVS)

(a)



(b)




Figure 6 (a) The source profiles of PM_{2.5} prediction by positive matrix factorization model and (b) comparison of modeled PM_{2.5} (PMF) and measured PM_{2.5} (HVS).