



Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf¹, L. D. Ziemba¹, G. Chen¹, C. A. Corr^{1,2}, J. H. Crawford¹, G. S. Diskin¹, R. H. Moore¹, K. L. Thornhill^{1,3}, E. L. Winstead^{1,3}, and B. E. Anderson¹

¹Science Directorate, NASA Langley Research Center, Hampton, VA, USA

²NASA Postdoctoral Program, Oak Ridge Associated Universities, Oak Ridge, TN, USA

³Science Systems and Applications Inc., Hampton, VA, USA

Received: 5 August 2015 – Accepted: 12 August 2015 – Published: 28 August 2015

Correspondence to: A. J. Beyersdorf (andreas.j.beyersdorf@nasa.gov)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

In order to utilize satellite-based aerosol measurements for the determination of air quality, the relationship between aerosol optical properties (wavelength-dependent, column-integrated extinction measured by satellites) and mass measurements of aerosol loading ($PM_{2.5}$ used for air quality monitoring) must be understood. This connection varies with many factors including those specific to the aerosol type, such as composition, size and hygroscopicity, and to the surrounding atmosphere, such as temperature, relative humidity (RH) and altitude, all of which can vary spatially and temporally. During the DISCOVER-AQ (Deriving Information on Surface conditions from Column and Vertically Resolved Observations Relevant to Air Quality) project, extensive in-situ atmospheric profiling in the Baltimore, MD–Washington, DC region was performed during fourteen flights in July 2011. Identical flight plans and profile locations throughout the project provide meaningful statistics for determining the variability in and correlations between aerosol loading, composition, optical properties and meteorological conditions.

Measured water-soluble aerosol mass was composed primarily of ammonium sulfate (campaign average of 32 %) and organics (57 %). A distinct difference in composition was observed with high-loading days having a proportionally larger percentage of ammonium sulfate (up to 49 %) due to transport from the Ohio River Valley. This composition shift caused a change in the aerosol water-uptake potential (hygroscopicity) such that higher relative contributions of ammonium sulfate increased the bulk aerosol hygroscopicity. These days also tended to have higher relative humidity causing an increase in the water content of the aerosol. Conversely, low aerosol loading days had lower ammonium sulfate and higher black carbon contributions causing lower single scattering albedos (SSAs). The average black carbon concentrations were 240 ng m^{-3} in the lowest 1 km decreasing to 35 ng m^{-3} in the free troposphere (above 3 km).

Routine airborne sampling over six locations was used to evaluate the relative contributions of aerosol loading, composition, and relative humidity (the amount of water

ACPD

15, 23317–23355, 2015

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

available for uptake onto aerosols) to variability in mixed layer aerosol. Aerosol loading was found to be the predominant source accounting for 88 % on average of the measured spatial variability in extinction with lesser contributions from variability in relative humidity (10 %) and aerosol composition (1.3 %). On average, changes in aerosol loading also caused 82 % of the diurnal variability in ambient aerosol extinction. However on days with relative humidity above 60 %, variability in RH was found to cause up to 62 % of the spatial variability and 95 % of the diurnal variability in ambient extinction.

This work shows that extinction is driven to first-order by aerosol mass loadings; however, humidity-driven hydration effects play an important secondary role. This motivates combined satellite/modelling assimilation products that are able to capture these components of the AOD-PM_{2.5} link. Conversely, aerosol hygroscopicity and SSA play a minor role in driving variations both spatially and throughout the day in aerosol extinction and therefore AOD. However, changes in aerosol hygroscopicity from day-to-day were large and could cause a bias of up to 27 % if not accounted for. Thus it appears that a single daily measurement of aerosol hygroscopicity can be used for AOD-to-PM_{2.5} conversions over the study region (on the order of 1400 km²). This is complimentary to the results of Chu et al. (2015) that determined the aerosol vertical distribution from “a single lidar is feasible to cover the range of 100 km” in the same region.

1 Introduction

Aerosols are detrimental to human health and are regulated as a criteria pollutant by the United States Environmental Protection Agency (EPA, 2014) and international agencies (Vahlsing and Smith, 2012) with compliance based on measurements at ground sites. However, satellites allow for the measurement of atmospheric conditions with a larger spatial coverage than possible with a ground-based network of instruments and thus have the potential to be useful tools in diagnosing ground-level air quality, particularly of aerosols (Al-Saadi et al., 2005). Additionally, satellites have the

advantage of detecting regional air quality events in areas without historical air quality problems which thus have no or limited ground-based sensor stations.

In order to relate satellite aerosol measurements to surface air quality, the connection between aerosol optical depth (AOD) measured by satellites to ground-level fine-mode aerosol mass ($PM_{2.5}$) must be known. The relationship between AOD and $PM_{2.5}$ has been widely studied (Hoff and Christopher, 2009 and the references therein; Crumeyrolle et al., 2014 for the current region) and ground-level $PM_{2.5}$ has been estimated based on AOD measurements both empirically (Liu et al., 2005) and through the use of global models. Van Donkelaar et al. (2006) found that the relative vertical extinction profile is the most important factor in the AOD-to- $PM_{2.5}$ relationship. Thus this relationship is weakest in regions where the vertical distribution cannot be reasonably modelled and is best in regions with fairly uniform aerosol type and vertical distribution (well-mixed boundary layer with minimal free tropospheric aerosol) such as the Northeast US (Engel-Cox et al., 2004). Based on lidar measurements in the Baltimore, MD–Washington, DC region, Chu et al. (2015) suggested that a single lidar could provide adequate information on the vertical distribution to allow for retrievals of $PM_{2.5}$ from AOD measurements made within 100 km of the lidar. However, the AOD- $PM_{2.5}$ relationship is not only dependent on the aerosol vertical distribution but also variability in aerosol composition and relative humidity (RH), both of which can be large in urban areas due to the densely located nature of local and regional sources. This work is an analysis of spatial and temporal variability in aerosol loading, composition and RH in the Baltimore, MD–Washington, DC region and their effect on variability in aerosol extinction.

DISCOVER-AQ (Deriving Information on Surface conditions from Column and Vertically Resolved Observations Relevant to Air Quality) was a multi-city NASA project designed to better elucidate the connection between satellite measurements and air quality by studying the variability in gas-phase and particulate pollutants in urban environments. The first campaign was performed in the Baltimore–Washington region in July 2011 and combined remote sensing instruments on the NASA Langley UC-12 fly-

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ing at 9 km, ground-based observations at multiple sites throughout the region, and in situ airborne measurements from the NASA Wallops P-3B for the detailed analysis of atmospheric composition in the Baltimore–Washington urban airshed. The P-3B flight plans (Fig. 1) were consistent among the 14 flights over 29 days to provide meaningful statistics (Table 1).

DISCOVER-AQ provides a valuable dataset to determine the variability in aerosol extinction throughout the region. However, it is important to note that changes in aerosol extinction are not necessarily solely due to an increase or decrease in aerosol loadings but can also be indicative of variability in relative humidity and aerosol composition. Thus these data will be used to examine:

1. the influence that aerosol loading, composition and relative humidity have on variability in aerosol extinction in the Baltimore–Washington region; and
2. the spatial and temporal resolution requirements of these parameters necessary to reproduce the variability in aerosol extinction.

These questions are relevant to scientists and policy makers seeking to assess the ability of satellite AOD retrievals to diagnose ground-level air quality.

2 Experimental design

The NASA P-3B was equipped with a variety of in situ aerosol and gas-phase measurements. Air was sampled with an isokinetic inlet which efficiently collects and transmits particles with a diameter smaller than 4 μm (McNaughton et al., 2007). The current analysis uses a subset of these measurements including aerosol scattering, absorption, size-distribution and composition. Scattering coefficients at 450, 550 and 700 nm were measured with an integrating nephelometer (TSI, Inc. model 3563) and corrected for truncation errors according to Anderson and Ogren (1998), while absorption coefficients at 470, 532 and 660 nm were measured with a Particle Soot Absorption Photometer (PSAP, Radiance Research) and corrected for filter scattering according to

23321

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Virkkula (2010). In order to calculate extinction, the measured Angstrom exponent was used to adjust the scattering at 550 to 532 nm (Ziemba et al., 2013).

During sampling, the RH of the air is modified due to the temperature gradient between the outside and inside of the plane. This causes a change in the scattering coefficient due to the generally hygroscopic nature of aerosol. To provide a stable scattering signal, the sample is initially dried to approximately 20 % RH utilizing a nafion drier and then sampled with tandem nephelometers (with and without humidification) to find the dry ($\sigma_{\text{scat, dry}}$ at a RH_{dry} of approximately 20 %) and humidified scattering coefficients ($\sigma_{\text{scat, wet}}$ at a RH_{wet} of approximately 80 %). These scattering measurements are related via a single-parameter monotonic growth curve (Gasso et al., 2000)

$$\sigma_{\text{scat, wet}} = \sigma_{\text{scat, dry}} \cdot \left[\frac{100 - \text{RH}_{\text{wet}}}{100 - \text{RH}_{\text{dry}}} \right]^{-\gamma} \quad (1)$$

where γ is an experimentally determined variable of the hygroscopicity with water-uptake increasing with increasing γ . $\sigma_{\text{scat, dry}}$ was corrected to 20 % RH based on Eq. (1) to account for any variability in RH_{dry} . Once γ is determined, the scattering at ambient RH ($\sigma_{\text{scat, amb}}$, RH_{amb}) is found from

$$\sigma_{\text{scat, amb}} = \sigma_{\text{scat, dry}} \cdot \left[\frac{100 - \text{RH}_{\text{amb}}}{80} \right]^{-\gamma} \quad (2)$$

Ambient RH was calculated based on measurements of water vapor concentration by an open-path diode laser hygrometer (Diskin et al., 2002), static temperature and pressure. Aerosol extinction at ambient RH ($\sigma_{\text{ext, amb}}$) can then be found by summing $\sigma_{\text{scat, amb}}$ and absorption (σ_{abs})

$$\sigma_{\text{ext, amb}} = \sigma_{\text{scat, dry}} \cdot \left[\frac{100 - \text{RH}_{\text{amb}}}{80} \right]^{-\gamma} + \sigma_{\text{abs}} \quad (3)$$

The dependence of aerosol absorption on RH is highly uncertain (Redemann et al., 2001; Mikhailov et al., 2006; Brem et al., 2012) and is therefore not incorporated but

likely manifests as only a small uncertainty in total extinction due to the fact that absorption was only a minor component of extinction (4 % on average).

Ziemba et al. (2013) showed a good correlation (R^2 of 0.88) between extinction measurements from the P-3B and coincident measurements performed by a high spectral resolution lidar (HSRL) on the UC-12. Recent work (Brock et al., 2015; Wagner et al., 2015) have suggested an additional model for aerosol hygroscopicity known as the kappa (κ) parameterization. However, these two models (based on γ and κ) are fairly consistent (scattering within 5 %) at RHs below 85 %, a range which comprised 96 % of the data measured by the P-3B. In addition, the good agreement between HSRL and in situ data (utilizing the γ correction scheme) suggest this is a valid model for the aerosol measured in Baltimore during DISCOVER-AQ (Ziemba et al., 2013).

Single scattering albedo (SSA) describes the relationship between aerosol scattering and extinction:

$$SSA = \left(\frac{\sigma_{\text{scat, dry}}}{\sigma_{\text{ext, dry}}} \right) = \left(\frac{\sigma_{\text{scat, dry}}}{\sigma_{\text{scat, dry}} + \sigma_{\text{abs}}} \right). \quad (4)$$

SSA can vary with RH (as scattering increases) but is here defined as the SSA under dry conditions (20 % RH). Thus Eq. (3) can be rewritten as

$$\sigma_{\text{ext, amb}} = \sigma_{\text{ext, dry}} \cdot \left[1 + SSA \cdot \left(\left[\frac{100 - RH_{\text{amb}}}{80} \right]^{-\gamma} - 1 \right) \right]. \quad (5)$$

Black carbon (BC) mass was measured with a Single Particle Soot Photometer (SP2, Droplet Measurement Technologies) while a pair of Particle-Into-Liquid Samplers (PILS; Brechtel Manufacturing, Inc.; Weber et al., 2001) were used to measure water-soluble organic and inorganic species. The PILS captures particles in the sampled air flow into a liquid flow of deionized water. The first PILS was coupled to a total organic carbon (TOC) analyzer (Sievers Model 800) to give the mass of water-soluble organic carbon (WSOC) at a 10 s time resolution. The TOC analyzer reports the organic carbon mass in $\mu\text{g C m}^{-3}$ and not the total organic mass (which includes mass

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



due to bonded hydrogen and oxygen atoms). Thus, to determine total water-soluble organic matter (WSOM), a multiplier ranging from 1.6 for urban to 2.1 for non-urban aerosols must be applied (Turpin and Lim, 2001). For the present work, a value of 1.8 is used based on Hand and Malm (2007). However, it should be noted that this does not include mass from any water-insoluble organic compounds. The liquid flow from the second PILS was collected in vials at a resolution of 3.25 or 5 min for off-line ion chromatographic analysis of chloride, nitrate, nitrite, sulfate, sodium, ammonium, potassium, magnesium, and calcium mass concentrations. Complete inorganic composition data are not available from the first three flights due to contamination from the sample vials; alternate vials were used for the remainder of the campaign. Aerosol size distributions were measured with an Ultra-High Sensitivity Aerosol Spectrometer (UHSAS, Droplet Measurement Technologies) calibrated with ammonium sulfate. All data are publicly available from the NASA Langley Research Center Atmospheric Science Data Center (ASDC, 2015).

As the PILS is unable to measure insoluble aerosol, the measured aerosol mass is a lower limit for the actual mass. The PILS mass can be compared to the volume measured by the UHSAS utilizing a density determined based on the measured mass of organics (1.2 g cm^{-3} , Turpin and Lim, 2001) and ammonium sulfate (1.77 g cm^{-3}). Based on this analysis, the PILS measured approximately 82 % of the aerosol mass with the other 18 % assumed to be insoluble organic compounds. Higher insoluble organic masses are estimated for higher loadings days with insoluble loadings near zero for low loading days. While this analysis gives an approximation of possible insoluble mass, the uncertainty is high and therefore this insoluble mass estimation is not included in future analysis.

3 Results – mission overview

Each DISCOVER-AQ-Maryland flight can be broken into two to three repetitive circuits which encompassed spirals from 0.3 to 4.5 km centered over six primary ground

**Aerosol composition
and variability in the
Baltimore–
Washington, DC
region**

A. J. Beyersdorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

most completely neutralized as ammonium sulfate, $(\text{NH}_4)_2\text{SO}_4$, with minimal bisulfate, $(\text{NH}_4)\text{HSO}_4$. Composition varied between flights with polluted days (as noted in Fig. 4) exhibiting a higher fraction of ammonium sulfate. Back-trajectory analysis with the Hybrid Single Particle Lagrangian Integrated Trajectory Model (HYSPLIT; Draxler and Hess, 1998; Draxler and Rolph, 2015) suggested these high aerosol loading days were related to long-range transport from the Ohio River Valley (Fig. 6) which has enhanced sulfur dioxide emissions due to a high density of coal-fired power plants in the region (Hand et al., 2012). These days were generally associated with low pressure systems to the northwest of the study region. Conversely, low loading days tended to have northerly flow due to high pressure systems to the west.

The flights with transport from the west and higher aerosol loadings (starred in Fig. 4) were found to have relatively more ammonium sulfate (38 % of mass compared to 20 % for clean days) and less organics (polluted, 52 %; clean, 65 %). Less polluted days had higher percentages of ammonium nitrate (polluted, 1.4 %; clean, 5 %) and BC (polluted, 2.0 %; clean, 2.7 %). The higher BC mass percentage also leads to higher absorption relative to scattering and therefore lower SSA on these less polluted days (polluted, 0.98; clean, 0.93; Fig. 7). However, on an absolute basis the polluted days had higher BC and absorption than on the clean days. Average BC concentrations for the entire month were 240 ng m^{-3} in the lowest 1 km decreasing to 35 ng m^{-3} in the free troposphere (above 3 km).

The polluted flight days also had higher γ values (Fig. 7). This water-uptake is largely dependent on aerosol composition with soluble organics having lower hygroscopicity than ammonium sulfate and ammonium nitrate. This can be seen as an inverse relationship with $\gamma = 0.60 - 0.0042 \times \text{organic mass fraction}$ (Fig. 8). These values are intermediate between measurements made in other urban areas (Asia and US, Quinn et al., 2005; Texas, Massoli et al., 2009) and in the remote atmosphere (the Indian Ocean, Quinn et al., 2005). Differences are likely due to differences in the measurement of organics; the current study uses PILS to measure only water-soluble organics while the other studies use aerosol mass spectrometry or thermo-optical methods which are

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sensitive to all organic species. In addition to an elevated γ , high loading days were typically more humid ($64 \pm 7\%$ compared to $49 \pm 7\%$). These higher humidities and γ -values resulted in a higher water content of the aerosols as evident from ambient extinctions that were 25 % higher than dry values on high loading days compared to the 12 % observed on low loading days. The highest daily-averaged water content of aerosol extinction was 40 % measured during Flight 8.

Aerosol mass is the primary measurement of aerosol loading and the basis by which ground air quality is regulated. Boundary layer dry extinction, ambient extinction and AOD are additional measures of aerosol loading in combination but incorporate an increasing amount of confounding factors. For instance, dry extinction is dependent on the aerosol mass loading in addition to aerosol size and composition. Ambient extinction is dependent on these same factors plus the aerosol hygroscopicity and RH. Finally, AOD is also dependent on the vertical distribution of aerosols and RH. To a first approximation (Fig. 4), aerosol mass loading, dry extinction (not shown), ambient extinction and AOD follow similar trends showing that aerosol mass loadings are the primary factor controlling day-to-day variability in aerosol optical properties. However, aerosol mass measurements via PILS do not account for insoluble aerosol. Dry mass extinction efficiencies calculated from extinction and mass measurements were variable ranging between 3.2 and $8.3 \text{ m}^2 \text{ g}^{-1}$. The highest mass extinction efficiencies (measured on high loading days) likely are indicative of the presence of insoluble organic material. Therefore, because of the variable quantity of insoluble mass and the low time resolution of the PILS measurements, future analysis will use the dry extinction as a proxy for aerosol loadings.

4 Results – regional variability

Aerosol extinction varied not only on a temporal basis (Fig. 4) but also spatially. Because there is such a large difference in aerosol loadings, optical properties (related to composition) and RH between flights, using campaign averages would distort the spa-

tial trends. Therefore, each circuit consisting of spirals over six ground sites is treated as a separate “snapshot” of the region and the properties measured over each site are normalized to the circuit average to study the spatial variability. Data below 1 km pressure altitude were used for 34 circuits for which spirals were performed over all six sites (absorption measurements were not available for one additional circuit and therefore it was not included in this analysis).

In order to get a general overview of aerosol variability in the regional, the average normalized dry and ambient extinctions along with RH for all of the circuits are shown in Fig. 9. The data is first normalized to the average for the circuit and then the normalized values are averaged. The highest dry aerosol extinction was nearest downtown Baltimore with Site 5 extinction 5.6 % larger than the average. However, the average ambient extinction measured was highest at the north end of the region where Site 3 is 5.5 % larger. This is consistent with the observed latitudinal gradient in RH. This shows that meteorological conditions (RH) can alter spatial trends in ambient extinction. Theoretically, it is possible for the entire region to have the same aerosol loading but differing extinction due to variability in composition and RH. Conversely, it is possible that the entire region could have a gradient in aerosol loading yet the composition and RH vary in such a way that extinction is constant throughout the region.

However, in order to study aerosol variability it is important to analyze each circuit individually (and not as a campaign average as done in Fig. 9). Equation (5) shows the dependence of aerosol ambient extinction on aerosol loading ($\sigma_{\text{ext, dry}}$), composition (SSA and γ) and RH, and can be used as a simple model to determine the factors controlling aerosol ambient extinction. From this, an assessment of the accuracy needed for each of these parameters to relate aerosol extinction (which can be derived from satellite measurements) to aerosol loading can be performed. In order to determine the relative importance of aerosol loading, composition and RH on extinction, the partial

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



derivatives of Eq. (5) can be determined:

$$\frac{\partial \sigma_{\text{ext, amb}}}{\partial \sigma_{\text{ext, dry}}} = 1 + \text{SSA} \cdot \left(\left[\frac{100 - \text{RH}_{\text{amb}}}{80} \right]^{-\gamma} - 1 \right), \quad (6)$$

$$\frac{\partial \sigma_{\text{ext, amb}}}{\partial \text{SSA}} = \sigma_{\text{ext, dry}} \cdot \left(\left[\frac{100 - \text{RH}_{\text{amb}}}{80} \right]^{-\gamma} - 1 \right), \quad (7)$$

$$\frac{\partial \sigma_{\text{ext, amb}}}{\partial \text{RH}} = \frac{\sigma_{\text{ext, dry}} \cdot \text{SSA} \cdot \gamma}{80} \left[\frac{100 - \text{RH}_{\text{amb}}}{80} \right]^{-\gamma-1}, \quad (8)$$

$$\frac{\partial \sigma_{\text{ext, amb}}}{\partial \gamma} = -\sigma_{\text{ext, dry}} \cdot \text{SSA} \cdot \left[\frac{100 - \text{RH}_{\text{amb}}}{80} \right]^{-\gamma} \cdot \ln \left(\frac{100 - \text{RH}_{\text{amb}}}{80} \right). \quad (9)$$

As expected, ambient extinction is linear with dry extinction (the partial derivative does not contain $\sigma_{\text{ext, dry}}$). The positive linear dependence on SSA shows that if all other variables are held constant, as SSA increases scattering becomes a larger fraction of extinction and at any RH above 20% will cause an increase in extinction due to water uptake. The dependence on RH and γ are both non-linear and thus their effects are most important when the RH is high or the aerosol is very hygroscopic.

Equations (6) through (9) can be combined to give the total differential for $\sigma_{\text{ext, amb}}$

$$d\sigma_{\text{ext, amb}} = \frac{\partial \sigma_{\text{ext, amb}}}{\partial \sigma_{\text{ext, dry}}} \cdot d\sigma_{\text{ext, dry}} + \frac{\partial \sigma_{\text{ext, amb}}}{\partial \text{SSA}} \cdot d\text{SSA} + \frac{\partial \sigma_{\text{ext, amb}}}{\partial \text{RH}} \cdot d\text{RH} + \frac{\partial \sigma_{\text{ext, amb}}}{\partial \gamma} \cdot d\gamma. \quad (10)$$

Assuming that the four variables are independent

$$s(\sigma_{\text{ext, amb}}) = \left[\left(\frac{\partial \sigma_{\text{ext, amb}}}{\partial \sigma_{\text{ext, dry}}} \cdot s(\sigma_{\text{ext, dry}}) \right)^2 + \left(\frac{\partial \sigma_{\text{ext, amb}}}{\partial \text{SSA}} \cdot s(\text{SSA}) \right)^2 + \left(\frac{\partial \sigma_{\text{ext, amb}}}{\partial \text{RH}} \cdot s(\text{RH}) \right)^2 + \left(\frac{\partial \sigma_{\text{ext, amb}}}{\partial \gamma} \cdot s(\gamma) \right)^2 \right]^{1/2} \quad (11)$$

where $s(x)$ is the standard deviation in x which is used as a measure of the variability in measurements made at the six sites during one circuit. Each term signifies the explained variance due to each of the four properties. Thus the relative contribution (RC)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of dry aerosol scattering to the variability in ambient extinction in the region can then be found by:

$$RC(\sigma_{\text{ext, dry}}) = \frac{\left(\frac{\partial\sigma_{\text{ext, amb}}}{\partial\sigma_{\text{ext, dry}}} \cdot s(\sigma_{\text{ext, dry}})\right)^2}{s(\sigma_{\text{ext, amb}})^2}. \quad (12)$$

Using this method, the RC for each of the four variables can be determined for each circuit.

In order to determine the relative contribution of each factor on the variability in ambient aerosol extinction, each circuit was analyzed separately. Shown in Fig. 10 are two extreme cases. During Flight 1, ambient relative humidity was low ($37 \pm 4\%$) resulting in little water uptake (the shaded portion on the upper panel). Thus variability in dry extinction (aerosol loading) is the major contributor ($RC(\sigma_{\text{ext, dry}}) = 99\%$) to variability in ambient extinction. The second case during Flight 14 shows a period of high RH ($64 \pm 8\%$). Water uptake was substantial and greatest at Site 3 where the RH is the highest. In this case, the variability in aerosol extinction is not only dependent on variability in dry extinction (41%) but also relative humidity (57%).

On average, aerosol loading (dry extinction) accounted for 88% of the spatial variability in extinction, with 27 of the 34 complete circuits having $RC(\sigma_{\text{ext, dry}})$ above 80% (Fig. 11). Variability in RH only accounted for 10% of the ambient extinction variability on average with only 5 circuits having $RC(\text{RH})$ greater than 20%. Four of these cases where RH had a large effect on ambient extinction variability corresponded to days with high RH (above 60%). This is due to the non-linearity of extinction with respect to RH (Eq. 8). Thus at low relative humidities, changes in RH minimally impact ambient extinction. Conversely, when RHs are high, small changes can produce large variations in ambient extinction. Changes in γ and SSA were smaller contributors to ambient extinction variability (1.3% and less than 0.1% on average, respectively).

5 Results – diurnal variability

A similar analysis can be performed to examine the diurnal variability of aerosol extinction. For this analysis, each variable was averaged for each of the six sites during each flight. This produced data at each spiral site approximately every 2 h during each flight period (3 to 4 values per site per day); the comparison between these values were then used to determine the diurnal variability in each parameter over the course of each flight. Figure 12 shows data at Site 4 from the same flights used for the regional variability analysis. For Flight 1, little water uptake occurred during the flight period so more than 99 % of the diurnal change in ambient extinction is due to changes in aerosol loading. In contrast, during Flight 14, extinction variability is dependent on both changes in aerosol loading and RH (51 and 49 %, respectively). From the first to second circuit, ambient extinction dropped as a result of an RH change from 70 to 59 %. After 16:00 local time, the RH continued to drop but ambient extinction increased due to an increase in dry aerosol extinction. Thus in this case, knowledge of the aerosol loading and RH trends are needed to interpret the aerosol extinction diurnal trends. On average, diurnal extinction variability was dominated by changing aerosol loading (82 %) with smaller contributions from changes in RH, γ and SSA (16, 1.6 and less than 0.1 %, respectively). However, the RC (RH) was as high as 95 % during Flight 9, a day with high RH and highly variable RH (Fig. 11).

6 Discussion

The conversion of extinction at ambient RH to extinction at a reduced (“dry”) RH is important in relating remote sensing measurements of ambient extinction to dry aerosol mass. Though the analysis above shows that variability in γ and SSA are only minor contributors to ambient extinction variability, converting between ambient and dry extinction requires knowledge of both parameters, as evident by Eq. (3). However, both γ and SSA are not routinely measured at air quality monitoring sites. So the question

ACPD

15, 23317–23355, 2015

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

could be asked “At what frequency (both spatially and temporally) do γ and SSA need to be known to determine the proper RH conversion?” This can be examined by analyzing the DISCOVER-AQ-Maryland data recorded below 1 km and determining how using more averaged data yields differing ambient aerosol extinctions.

As a result of changes in composition seen in Fig. 4, γ varied between 0.14 (Flight 1) and 0.47 (Flight 8) with an average of 0.32 (Fig. 13). Comparing the ambient extinction calculated during each spiral with the extinction calculated using the daily average γ resulted in a bias of $\pm 1.6\%$ in ambient extinction with no clear trend with respect to aerosol extinction. Using the monthly average for the entire region causes a bias of $\pm 6.8\%$ (Table 2) with deviations of up to 27% at high aerosol extinction because γ tended to be higher on high aerosol loading days (Fig. 8). We conclude that spatial γ differences in the Baltimore region are not large enough to cause significant biases in deriving dry extinction from ambient values. However, day-to-day variability in γ can cause large discrepancies. Thus it appears that a single daily measurement of γ (or one based on compositional measurements, Fig. 8) is able to be used for AOD-to-PM_{2.5} correlations over the study region (on the order of 1400 km²) within an uncertainty of 2%.

A similar analysis can be performed to evaluate the importance of SSA in retrieving dry extinction from ambient extinction (Fig. 14 and Table 2). SSA varied from 0.91 to 0.99 during the mission with higher SSA measured on high aerosol loading days due to the increased loading of ammonium sulfate and other secondary aerosols which are typically more scattering than primary aerosols. Comparing the ambient extinction calculated during each spiral with the extinction calculated using the daily average SSA resulted in a bias of $\pm 0.2\%$ in ambient extinction showing that regional variability in SSA was not high enough to make a significant difference. Using the monthly average for the entire region produces biases of $\pm 0.5\%$ with deviations of up to 1.0% at high aerosol extinction.

Doing the same analysis for dry aerosol extinction or RH show markedly different results (Figs. 15 and 16, Table 2). The use of a daily average dry extinction causes a bias

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of $\pm 22\%$ showing that regional variation in aerosol loading must be accounted for. Utilizing a monthly average extinction causes discrepancies of $\pm 111\%$ due to the large day-to-day variability in aerosol loading. Biases based on limited knowledge of RH were smaller with $\pm 6.2\%$ for daily and 11% for monthly RH. Thus, Table 2 gives a hierarchy of factors for variability in extinction measurements: loading $>$ RH $>$ γ $>$ SSA.

An analysis of the effects of aerosol and meteorological parameters on AOD in the Southeastern US based on 37 airborne profiles (Brock et al., 2015) show similar trends in the significance of factors with aerosol mass the most important. Relative humidity had a non-linear significance on AOD with the greatest significance for extremely humid conditions (the 90th percentile RH profiles). Varying aerosol size parameters and the vertical distribution of the aerosols resulted in moderate AOD changes; while AODs were largely insensitive to refractive index in a fashion similar to the present findings of SSA as a minor contributor to extinction variability.

7 Conclusions

Measurements made in the Baltimore–Washington DC region during DISCOVER-AQ in July 2011 can be generalized as such: on days influenced by transport from the Ohio River Valley, aerosol loadings were higher (aerosol mass concentrations of $18.7 \pm 4.4 \mu\text{g m}^{-3}$ and AODs of 0.43 ± 0.12) and the aerosol were more hygroscopic (γ of 0.36 ± 0.07) because of a larger percentage of ammonium sulfate (38 % of water-soluble mass) in comparison to days impacted by northerly transport (aerosol masses of 5.4 ± 1.3 , AODs of 0.08 ± 0.03 , γ of 0.26 ± 0.09 , 20 % ammonium sulfate). In both cases, the regional and diurnal variability in aerosol extinction are controlled primarily by changes in aerosol loadings. However, on days associated with westerly transport (which also were more humid) variability in RH also contributed significantly to the regional (14 %) and diurnal (22 %) variability in extinction. Thus changes in AOD cannot directly be seen as changes in $\text{PM}_{2.5}$ but must take into account spatial and temporal variability in RH.

**Aerosol composition
and variability in the
Baltimore–
Washington, DC
region**

A. J. Beyersdorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Brem, B. T., Mena Gonzalez, F. C., Meyers, S. R., Bond, T. C., and Rood, M. J.: Laboratory-measured optical properties of inorganic and organic aerosols at relative humidities up to 95 %, *Aerosol Sci. Tech.*, 46, 178–190, 2012.
- Chu, D. A., Ferrare, R., Szykman, J., Lewis, J., Scarino, A., Hains, J., Burton, S., Chen, G., Tsaif, T., Hostetler, C., Hair, J., Holben, B., and Crawford, J.: Regional characteristics of the relationship between columnar AOD and surface PM_{2.5}: application of lidar aerosol extinction profiles over Baltimore–Washington Corridor during DISCOVER-AQ, *Atmos. Environ.*, 101, 338–349, 2015.
- Crumeyrolle, S., Chen, G., Ziemba, L., Beyersdorf, A., Thornhill, L., Winstead, E., Moore, R. H., Shook, M. A., Hudgins, C., and Anderson, B. E.: Factors that influence surface PM_{2.5} values inferred from satellite observations: perspective gained for the US Baltimore–Washington metropolitan area during DISCOVER-AQ, *Atmos. Chem. Phys.*, 14, 2139–2153, doi:10.5194/acp-14-2139-2014, 2014.
- Diskin, G. S., Podolske, J. R., Sachse, G. W., and Slate, T. A.: Open-path airborne tunable diode laser hygrometer, *Proc. SPIE*, 4817, 196–204, 2002.
- Draxler, R. R. and Hess, G. D.: An overview of the HYSPLIT-4 modeling system of trajectories, dispersion, and deposition, *Aust. Meteorol. Mag.*, 47, 295–308, 1998.
- Draxler, R. R. and Rolph, G. D.: NOAA Air Resources Laboratory HYSPLIT (HYbrid Single-Particle Lagrangian Integrated Trajectory), available at: <http://ready.arl.noaa.gov/HYSPLIT.php> (last access: 5 August 2015), 2015.
- Engel-Cox, J. A., Holloman, C. H., Coutant, B. W., and Hoff, R. M.: Qualitative and quantitative evaluation of MODIS satellite sensor data for regional and urban scale air quality, *Atmos. Environ.*, 38, 2495–2509, 2004.
- EPA, 2014: National Ambient Air Quality Standards, available at: <http://www.epa.gov/air/criteria.html> (last access: 5 August 2015), 2015.
- Gasso, S., Hegg, D. A., Covert, D. S., Collins, D., Noone, K. J., Ostrom, E., Schmid, B., Russell, P. B., Livingston, J. M., Durkee, P. A., and Jonsson, H.: Influence of humidity on the aerosol scattering coefficient and its effect on the upwelling radiance during ACE-2, *Tellus B*, 52, 546–567, 2000.
- Hand, J. L. and Malm, W. C.: Review of aerosol mass scattering efficiencies from ground-based measurements since 1990, *J. Geophys. Res.*, 112, D16203, doi:10.1029/2007JD008484, 2007.

**Aerosol composition
and variability in the
Baltimore–
Washington, DC
region**

A. J. Beyersdorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Hand, J. L., Schichtel, B. A., Malm, W. C., and Pitchford, M. L.: Particulate sulfate ion concentration and SO₂ emission trends in the United States from the early 1990s through 2010, *Atmos. Chem. Phys.*, 12, 10353–10365, doi:10.5194/acp-12-10353-2012, 2012.
- Hoff, R. M. and Christopher, S. A.: Remote sensing of particulate pollution from space: have we reached the promised land?, *JAPCA J. Air Waste Ma.*, 59, 645–675, 2009.
- Liu, Y., Sarnat, J. A., Kilaru, V., Jacob, D. J., and Koutrakis, P.: Estimating ground-level PM_{2.5} in the Eastern United States using satellite remote sensing, *Environ. Sci. Technol.*, 39, 3269–3278, 2005.
- Massoli, P., Bates, T. S., Quinn, P. K., Lack, D. A., Baynard, T., Lerner, B. M., Tucker, S. C., Brioude, J., Stohl, A., and Williams, E. J.: Aerosol optical and hygroscopic properties during TexAQS-GoMACCS 2006 and their impact on aerosol direct radiative forcing, *J. Geophys. Res.*, 114, D00F07, doi:10.1029/2008JD011604, 2009.
- McNaughton, C. S., Clarke, A. D., Howell, S. G., Pinkerton, M., Anderson, B., Thornhill, L., Hudgins, C., Winstead, E., Dibb, J. E., Scheuer, E., and Maring, H.: Results from the DC-8 Inlet Characterization Experiment (DICE): airborne versus surface sampling of mineral dust and sea salt aerosols, *Aerosol Sci. Tech.*, 41, 136–159, 2007.
- Mikhailov, E. F., Vlasenko, S. S., Podgorny, I. A., Ramanathan, V., and Corrigan, C. E.: Optical properties of soot–water drop agglomerates: an experimental study, *J. Geophys. Res.*, 111, D07209, doi:10.1029/2005JD006389, 2006.
- Quinn, P. K., Bates, T. S., Baynard, T., Clarke, A. D., Onasch, T. B., Wang, W., Rood, M. J., Andrews, E., Allan, J., Carrico, C. M., Coffman, D., and Worsnop, D.: Impact of particulate organic matter on the relative humidity dependence of light scattering: a simplified parameterization, *Geophys. Res. Lett.*, 32, L22809, doi:10.1029/2005GL024322, 2005.
- Redemann, J., Russell, P. B., and Hamill, P.: Dependence of aerosol light absorption and single-scattering albedo on ambient relative humidity for sulfate aerosols with black carbon cores, *J. Geophys. Res.*, 106, 27485–27495, 2001.
- Turpin, B. J. and Lim, H.-J.: Species contributions to PM_{2.5} mass concentrations: revisiting common assumptions for estimating organic mass, *Aerosol Sci. Tech.*, 35, 602–610, 2001.
- Vahlsing, C. and Smith, K. R.: Global review of national ambient air quality standards for PM₁₀ and SO₂ (24 h), *Air Qual. Atmos. Health*, 5, 393–399, 2012.
- van Donkelaar, A., Martin, R. V., and Rokjin, J. P.: Estimating ground-level PM_{2.5} using aerosol optical depth determined from satellite remote sensing, *J. Geophys. Res.*, 111, D21201, doi:10.1029/2005JD006996, 2006.

**Aerosol composition
and variability in the
Baltimore–
Washington, DC
region**

A. J. Beyersdorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Virkkula, A.: Correction of the calibration of the 3-wavelength particle soot absorption photometer (3 λ PSAP), *Aerosol Sci. Tech.*, 44, 706–712, 2010.

Wagner, N. L., Brock, C. A., Angevine, W. M., Beyersdorf, A., Campuzano-Jost, P., Day, D., de Gouw, J. A., Diskin, G. S., Gordon, T. D., Graus, M. G., Holloway, J. S., Huey, G., Jimenez, J. L., Lack, D. A., Liao, J., Liu, X., Markovic, M. Z., Middlebrook, A. M., Mikoviny, T., Peischl, J., Perring, A. E., Richardson, M. S., Ryerson, T. B., Schwarz, J. P., Warneke, C., Welti, A., Wisthaler, A., Ziemba, L. D., and Murphy, D. M.: In situ vertical profiles of aerosol extinction, mass, and composition over the southeast United States during SENEX and SEAC⁴RS: observations of a modest aerosol enhancement aloft, *Atmos. Chem. Phys.*, 15, 7085–7102, doi:10.5194/acp-15-7085-2015, 2015.

Weber, R. J., Orsini, D., Daun, Y., Lee, Y.-N., Klotz, P. J., and Brechtel, F.: A particle-into-liquid collector for rapid measurement of aerosol bulk chemical composition, *Aerosol Sci. Tech.*, 35, 718–727, 2001.

Ziemba, L. D., Thornhill, K. L., Ferrare, R., Barrick, J., Beyersdorf, A. J., Chen, G., Crumeyrolle, S. N., Hair, J., Hostetler, C., Hudgins, C., Obland, M., Rogers, R., Scarino, A. J., Winstead, E. L., and Anderson, B. E.: Airborne observations of aerosol extinction by in situ and remote-sensing techniques: evaluation of particle hygroscopicity, *Geophys. Res. Lett.*, 40, 417–422, 2013.

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. Percent bias in ambient extinction based on daily and monthly averaging of contribution variables.

Variable	Percent bias based on averaging	
	Daily	Monthly
Dry Extinction	22	111
RH	6.2	10.7
γ	1.6	6.8
SSA	0.21	0.49

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.




Figure 1. Flight path for Flight 1. Portions below 1 km are shown in red and those above in black. Flights originated at NASA Wallops Flight Facility (southeast of the area shown) to ground Sites 1 through 6 in order with a spiral performed at each site. The circuit was typically flown 3 times per flight before returning to Wallops. Water is denoted as blue with the Chesapeake Bay at the center and the Delaware Bay on the right edge.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.




Figure 2. Time series of extinction (at ambient RH and 532 nm) and altitude (gray dashed line) for Flight 9 (upper panel). Extinction measurements during each circuit are highlighted by differing background color. Each circuit is then plotted in the bottom panel to show the changes in aerosol between the circuits. Profile locations correspond to those shown in Fig. 1.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.




Figure 3. Vertical profiles of aerosol extinction (at ambient RH and 532 nm) for Flight 9 segregated by circuit and profile site. Horizontal lines represent the boundary layer (solid line) and buffer layer (dashed line) heights during each circuit at Site 2 based on airborne measurements of the potential temperature profile.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.




Figure 4. Average AOD (at ambient RH) along with boundary layer (below 1 km) extinction, aerosol mass, effective radius and composition for each of the fourteen flights. Aerosol mass and composition data are not available for the first three flights. Flights with predominantly westerly transport from the Ohio River Valley are indicated by stars at the top of the plots.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.




Figure 5. Average vertical profiles of aerosol extinction (at ambient RH and 532 nm) for all flights with Flights 9 and 14 highlighted (left panel). These profiles can then be normalized to the total aerosol loading (AOD) to get the normalized vertical profile (left panel, arbitrary units).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Aerosol composition
and variability in the
Baltimore–
Washington, DC
region**

A. J. Beyersdorf et al.




Figure 6. 72 h back-trajectories based on HYSPLIT for the first circuit of each flight at Site 5 at an altitude of 1 km colored by the average AOD measured during that flight.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.




Figure 7. Average profiles for extinction (at ambient RH and 532 nm), γ , SSA and composition for all flights (black line), days with predominantly westerly transport from the Ohio River Valley (red line), and days with northerly transport (blue line).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.



	Slope	Y-Intercept
Baltimore	-0.42	0.60
Texas	-0.5	0.84
Western Pacific	-0.7	0.9
Northeast U.S.	-0.5	0.8
Indian Ocean	-0.3	0.5

Figure 8. Relationship between γ (at 532 nm) and organic mass fraction for the present study (data below 1 km), Texas (Massoli et al., 2009), the western Pacific, the northeast US, and the Indian Ocean (Quinn et al., 2005). The organic mass fraction is found by dividing the WSOM by the total mass measured by the PILS and SP2. Other studies used organic mass measured by aerosol mass spectrometer or thermo-optical methods. The ratio of scattering at 80 % RH to 20 % [$f(\text{RH})$] is shown on the right-axis (note the irregular spacing).

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.




Figure 9. Average normalized 532 nm dry extinction (left panel), RH (center) and 532 nm ambient extinction (right) for all of the circuits (data is normalized to the average value for that circuit). The site with the maximum value is labelled.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.




Figure 10. Average 532 nm ambient extinction, dry extinction and RH below 1 km during spirals over the six sites during Flights 1 and 14.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.




Figure 11. Relative contribution of dry extinction and RH on the spatial variability in ambient extinction as a function of RH (left) and on the diurnal variability (right). Diamonds represent the average relative contributions for 10 % RH increments.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.




Figure 12. Trends in 532 nm ambient extinction, dry extinction and RH below 1 km during spirals at Site 4 during Flights 1 and 14.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.




Figure 13. Average γ for each flight (top) along with estimated ambient extinction and percent bias if the flight-average (left) and campaign-average (right) γ are used.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.




Figure 14. Average SSA for each flight (top) along with estimated ambient extinction and percent bias if the flight-average (left) and campaign-average (right) SSA are used.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.




Figure 15. Average dry extinction for each flight (top) along with estimated ambient extinction and percent bias if the flight-average (left) and campaign-average (right) dry extinction are used.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Aerosol composition and variability in the Baltimore–Washington, DC region

A. J. Beyersdorf et al.




Figure 16. Average RH for each flight (top) along with estimated ambient extinction and percent bias if the flight-average (left) and campaign-average (right) RH are used.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

