

Reply to comment RC C7582

The manuscript by Tesdal et al. uses a global atmospheric GCM, which includes a representation of aerosol chemistry and processes, to evaluate the effect of various DMS emission estimates on sulfate burden and the subsequent radiative impact. The study builds incrementally on pre-existing work. The manuscript is well written and structured, and the content is appropriate for ACP. However, before publication I recommend that the authors consider the points below, particularly about evaluating the model's ability to reproduce observed quantities.

We thank Referee #1 for his/her insightful comments. In particular, we appreciate the point that the article did not sufficiently address the issue of model assessment, to which we have responded by adding a new subsection comparing simulated sulfate concentrations with observations. We have revised the manuscript to address all of the referee's points and to clarify our terminology in line with the suggestions made. Specific responses to all of the referee's points follow:

Major comments

The suitability of the model for the study is not demonstrated. If we are to have faith in the results of the model's response to different perturbations (as part of the sensitivity study), we need to first know that the model is capable of representing the quantities in question. Has the model ever been evaluated against observations of DMS, SO₂, sulphate, CCN, CDN, radiation...? Particularly in regions which are dominated by marine aerosol (Southern Ocean). Does the model simulate the observed seasonal cycles of these quantities adequately?

The model uses a basic (bulk) representation of aerosol which does not fully simulate aerosol microphysics (e.g. competition between condensation and new particle formation, coagulation / interaction between different aerosol species etc). Bellouin et al. (2013, ACP 13: 2027) find stark differences between a bulk and a microphysical aerosol scheme in how they simulate aerosol direct and indirect effects, including the response to a DMS perturbation. Some discussion of the limitations of the aerosol scheme is therefore necessary, and would be particularly useful when connected to an evaluation of model skill vs observations. Calculation of the 'CCN sensitivity' for comparison with the equivalent values calculated in Woodhouse et al. (2010) and others would also be valuable.

We would like to respond to issues raised in the first and second paragraph together since they are related.

We have added a comparison (new Section 3.1) of simulated and observed sulphate concentrations over the ocean in order to address the concern about a lack of model validation, with a focus on regions that are expected to be strongly affected by DMS emissions. These results provide evidence for realistic simulations of atmospheric DMS sources and aerosol removal processes in the marine atmosphere. In addition, the relationship between sulphate aerosol concentrations and cloud microphysical properties in simulations with CanAM and observations were previously investigated by Ma et al. (2010). Simulated aerosol concentrations,

cloud microphysical properties, and sulphate aerosol forcings (unpublished but assessed as part of the model development process) are all well within the range of uncertainty so that we have confidence that simulated impacts of DMS emissions on sulphate aerosol radiative forcings are meaningful. This is discussed in the revised manuscript (Section 3.1).

We agree that bulk and microphysical schemes typically tend to be very different in terms of the processes that are considered and results for radiative forcings. According to our own experience with the CCCma bulk and aerosol microphysics schemes, large differences exist for some of the results. However, we consider the bulk approach to be appropriate and perhaps even desirable in the context of the current study. The bulk aerosol scheme is a well-documented approach which produces results that are relatively straightforward to analyze. Similar bulk schemes have been widely used in different models in the past and are used in some CMIP5 models. Based on the few available studies that are based on results from different models with bulk and microphysics schemes we do not see evidence for considerable improvements in radiative forcing estimates based on simulations with microphysics schemes relative to bulk schemes (Schulz et al., 2006; Koch et al., 2009; Quaas et al., 2009). The manuscript has been revised to make this point (see end of Section 2.1 in revised manuscript). The Discussion section of the revised text also explicitly raises the point that we consider a single AGCM in this study, and neglect uncertainties in the radiative effect of DMS based on different model formulations.

Unfortunately, we are unable to calculate a "CCN sensitivity" for comparisons with results from Woodhouse et al. (2010) owing to the fact that the model does not simulate CCN concentrations. The simple parameterization of aerosol indirect effects in the model supplies cloud droplet number concentration as a function of sulphate concentration.

Relevant changes

page 5, lines 19-22: Added "The bulk scheme considered in this study is simpler than approaches that consider aerosol microphysics in detail (e.g., Bellouin et al., 2013). However, based on the few available studies comparing results from different models with bulk and microphysics schemes, we do not see evidence of considerable improvement in radiative forcing estimates based on simulations with microphysics schemes relative to bulk schemes (Schulz et al., 2006; Koch et al., 2009; Quaas et al., 2009)."

pages 6-8: Added new Section 3.1.

pages 24-25: Added new Figures 2 & 3.

I was surprised that greater prominence was not given to the air resistance result and the impact that can potentially have on DMS flux / aerosol. That seems like an important conclusion, and the biggest contribution.

We agree that this is a significant result from our modelling exercise. We have emphasized it further in our conclusion.

Relevant changes

page 15, lines 4-7: Added "Neglect of air-side resistance in the flux parameterization was shown to have a larger effect on net TOA radiation than suppressing spatial or temporal structure in the DMS concentration field. From the perspective of global climate, accurate formulation of surface fluxes is as or more important than accurate representation of sea surface DMS concentrations."

The word 'forcing' is used incorrectly in the manuscript (including in the title). Radiative forcing is the difference in radiation budget between two time periods (e.g. preindustrial and present day). I think the term 'radiative effect' is what is meant, as the manuscript in question doesn't consider different time periods (except in one paragraph on page 23948: line 25).

In the paper, radiative forcings were defined as net changes in the TOA energy balance due to changes in sulfate aerosol concentrations which are produced by DMS emissions and subsequent oxidation of DMS in the atmosphere. A distinction between present-day and preindustrial time periods is different than our analysis of the impacts of different DMS emission data sets on climate. We believe that our approach is consistent with the definition of radiative forcings given in the IPCC AR5 report:

"RF is the net change in the energy balance of the Earth system due to some imposed perturbation. It is usually expressed in watts per square meter averaged over a particular period of time and quantifies the energy imbalance that occurs when the imposed change takes place. Though usually difficult to observe, calculated RF provides a simple quantitative basis for comparing some aspects of the potential climate response to different imposed agents, especially global mean temperature, and hence is widely used in the scientific community. Forcing is often presented as the value due to changes between two particular times, such as pre-industrial to present-day, while its time evolution provides a more complete picture." (Myhre et al., 2013)

We generally agree that 'forcing' has been used incorrectly in the original text and removed the term "radiative forcing" as per the referee's request. In order to account for the ambiguous usage of the term "radiative forcing" in previous literature and to clarify what we mean by "radiative effect", we added the following text (Introduction):

In the following, net changes in the energy balance at the top of the atmosphere due to changes in concentrations of DMS-derived sulfate will be referred to as the radiative effect, which is sometimes also referred to as radiative forcing.

In many places, however, we did not feel it was necessary to substitute the one phrase for the other. We rather modified the text wherever the term appeared, so as to include enough detail about the underlying processes that the meaning should be clearer than in the original text, with the term "radiative forcing" removed. In one place we retained the term radiative forcing, because it refers to specific outputs of particular models, which are documented in the literature cited.

Relevant changes

- page 1: Changed title to "Sensitivity of modelled sulfate aerosol and its radiative effect on climate to ocean DMS concentration and air-sea flux".
- page 1, line 10: Changed "radiative forcing" to "radiative effect of sulfate"
- page 1, line 13: Changed "net radiative forcing" to "the net radiative effect"
- page 3, lines 5-6: Added "In the following, net changes in the energy budget at the top of the atmosphere (TOA) due to changes in concentrations of DMS-derived sulfate will be referred to as the radiative effect, which is sometimes also referred to as radiative forcing"
- page 3, line 9: Changed "forcing" to "effect"
- page 3, line 10: Changed "radiative forcing" to "TOA radiative budget"
- page 5, lines 17-18: Changed "difference between the total radiative forcing and clear-sky forcing" to "difference between the net radiative fluxes for all-sky and clear-sky only"
- page 5, line 25: Changed "radiative forcing" to "net TOA radiation"
- page 9, line 13: Changed "radiative forcing" to "radiative effects"
- page 9, line 14: Changed "top of the atmosphere (TOA) radiative forcing" to "TOA net radiation"
- page 9, line 18: Changed "forcing" to "effect"
- page 9, line 19: Changed "forcing" to "effect"
- page 9, line 21: Changed "radiative forcing" to "net radiation"
- page 9, line 24: Changed "forcing" to "effect"
- page 9, line 27: Changed "radiative forcing" to "the radiation budget"
- page 9, line 33: Changed "radiative forcing" to "effects on the atmospheric radiation budget"
- pages 9-10, lines 34-1: Changed "in radiative forcing" to "of radiative perturbations"
- page 10, line 2: Changed "forcing" to "effect"
- page 10, line 3: Changed "radiative forcing" to "perturbations to net TOA radiation"
- page 10, line 8: Removed "radiative forcing"
- page 10, line 10: Changed "radiative forcing" to "net TOA radiation"
- page 10, lines 11-12: Changed "radiative forcing" to "radiation"
- page 10, line 19: Changed "forcing" to "effect"
- page 10, line 20: Changed "forcing" to "effect"
- page 10, line 21: Changed "radiative forcing" to "the radiative effects"
- page 10, line 27: Changed "forcing" to "effects"
- page 10, lines 28-29: Changed "resulting radiative forcing" to "concentration of sulfate aerosol and its effect on the TOA radiation budget"
- page 11, line 23: Changed "forcing" to "effect"
- page 12, line 20: Changed "mean radiative forcing" to "net TOA radiation"

page 12, line 23: Changed "forcing response to" to "effect of"
page 12, line 32: Changed "radiative forcing" to "the radiative effect of DMS emissions"
page 13, line 3: Removed "on radiative forcing"
page 13, line 12: Changed "forcing" to "effect"
page 13, line 13: Changed "radiative forcing" to "net TOA radiation"
page 13, line 16: Changed "radiative forcing" to "net TOA radiation"
page 13, lines 22-23: Changed "radiative forcing" to "the radiative effect of DMS"
page 14, line 31: Changed "radiative forcing" to "their effect on the planetary radiation budget"
page 14, line 33: Changed "radiative forcing associated with DMS" to "effect of DMS on net TOA radiation"
page 15, lines 9-10: Changed "DMS flux and uncertainty in radiative forcing" to "DMS flux, aerosol burdens and their radiative effects"
page 27: Changed "Radiative forcing difference (change" to "Change"

It would be very informative to see the spatial responses (change in DMS, sulphate, CDN, radiation... as a map) to the DMS perturbations. Presenting this information would make it much easier to compare and interpret the results against previous work.

Since our work is primarily concerned with global changes, we do not feel it is within the scope of the current contribution to include these maps in the main text. But we agree that these can be very informative for the interested reader, so we have added them as supplementary material.

Minor comments

These questions have helpfully pointed toward needed clarification, and we have added further detail to the corresponding sections.

Page 23939, line 19: How is the ensemble created? Varying initial conditions?

Following sentence has been added:

Each ensemble member uses the exact same model configuration, but in each a different seed was used in the random number generator used in the radiation code.

Page 23939, line 22: 'realizations' = 'ensemble members'?

Yes. We changed 'realizations' to 'ensemble members'.

Page 23939, line 22: It would be useful to state what the spread was between the realizations, for direct comparison to the response from altered DMS emissions.

Please note that the spreads of the ensemble members for each model run are also shown in

Figure 2 and 3. We added the following in the result section:

The largest ensemble spread in DMS emissions among the simulations is less than $0.06 \mu\text{mol m}^{-2} \text{d}^{-1}$, which is negligible compared to the overall range of DMS emissions of the different model runs ($3.15 \mu\text{mol m}^{-2} \text{d}^{-1}$).

Page 23944, line 28: Has the statistical significance of the changes been calculated?

Statistical significance is determined from Figure 9 by comparing the mean differences among the model runs with the corresponding error bars. We have added a reference to Figure 9 in the revised text and added the following to the caption of Figure 9 (Please note that Figure 9 in the revised manuscript is Figure 7 in the previously submitted version.):

To derive the error estimates, all treatments (control, temporally invariant, spatially uniform, and no air-side resistance) were pooled after their separate means were removed; error bars are \pm two standard deviations of the pooled data ($n = 12$). Statistical significance is determined by comparing the mean differences among the model runs with the corresponding error bars.

Page 23945, line 7: Discussion or presentation of the spatial patterns would be useful in understanding the increase in SO₂ but decrease in sulphate. Are any further model diagnostics available to probe this further? It's an interesting outcome and one which it would be useful to understand.

As discussed above, we consider a detailed analysis of spatial patterns to be beyond the scope of the current contribution, but the relevant maps are presented as supplementary material.

Page 23948, line 20: Vallina et al. (2007) don't go as far as calculating the atmospheric/ climate response.

We agree. Vallina et al. (2007) mainly looked at the response of DMS given a change in climate. With "weak effect" bias we are referring to a link between DMS and climate, both as the influence of DMS on climate and vice versa. We acknowledge that this was not clear in the original manuscript and the text has been changed accordingly:

Previous studies have found a relatively weak link between DMS fluxes and climate (e.g., Kloster et al., 2007; Vallina et al., 2007; Woodhouse et al., 2010).

Page 23948, line 20: Wouldn't a low DMS flux result in a lower background aerosol concentration, thus making the system more sensitive to DMS perturbations...?

The referee's question points to a lack of clarity in the wording, which we have revised to prevent misunderstanding. The effect described by the reviewer could conceivably occur, but it would require a large change in the relative flux, and there is no particular reason for that not to occur in cases with a higher 'baseline' flux as well. Because the radiative effect on climate is more or less linear with the total global DMS source, perturbing a DMS concentration field that has a higher global total flux will generally have a greater effect than perturbing a field with a lower total flux.

Reply to comment RC C9192:

This paper presents sensitivity runs using a GCM with an explicit aerosol scheme in order to quantify the impacts of using different seawater DMS climatologies and sea-air flux parameterizations. This study adds to the numerous studies in literature which have had similar aims at understanding the effect of DMS and impacts of changes in the DMS flux.

Unfortunately, in its current form, the manuscript does not add significantly to the literature and hence I cannot recommend its publication. Most of the work presented here has been done before and simple sensitivity studies do not add much to our current knowledge on the effect of DMS, especially when there is no comparison done with observations.

We thank Referee #2 for his/her constructive comments. We agree that we did not sufficiently clarify the advances made in our paper. In particular, our study adds to existing literature by assessing the relative importance of spatial and temporal variation of DMS fluxes to atmospheric properties as compared with the magnitude of the global total flux. No other systematic analysis of the relationship between spatio-temporal distribution of DMS emissions and aerosol radiative effects on climate has been published to date, despite large uncertainties in specified DMS data sets in models. We recognize, however, that this motivation was not clear in the previous version of the study. In order to clarify that this is our objective, we have added some discussion of this point to the Introduction, emphasizing how the study adds to the existing literature. We also note that the Discussion paper has already been cited by Mungall et al. (2016), "Dimethyl sulfide in the summertime Arctic atmosphere: measurements and source sensitivity simulations", doi:10.5194/acp-16-6665-2016. (none of the authors of the present paper were involved in that study.)

The manuscript has been revised to address the referee's points and to clarify how the study adds to existing literature. Discussion of all of the referee's points follow:

1) The suitability of the model to study the impact of DMS on aerosols, cloud properties and radiation changes has not been demonstrated. Have the authors compared the model results to any observations? Several climatologies of sulphate are available, and locations with atmospheric DMS and SO₂/H₂SO₄ observations can be also compared to show that the model is indeed close to the observations or highlight the major shortcomings of the model.

A similar concern was raised by the other referee and we would like to repeat our response: We have added a comparison of simulated and observed sulfate concentrations over the ocean in order to address the concern about a lack of model validation, with a focus on regions that are expected to be strongly affected by DMS emissions. These results provide evidence for realistic simulations of atmospheric DMS sources and aerosol removal processes in the marine atmosphere. In addition, the relationship between sulfate aerosol concentrations and cloud microphysical properties in simulations with CanAM and observations were previously investigated by Ma et al. (2010). Simulated aerosol concentrations, cloud microphysical properties, and unpublished results for sulfate aerosol forcings are all well within the range of uncertainty. So we have confidence that simulated impacts of DMS emissions on sulfate aerosol radiative effects on climate are meaningful.

Relevant changes

pages 6-8: Added new Section 3.1.

pages 24-25: Added new Figures 2 & 3.

2) What is the reason behind including sensitivity experiments based on seawater climatologies K99, K00 and L10? L10 is an updated version of the K00 and K00 climatology. It seems unnecessary to include all three considering the L10 climatology is an upgrade and covers all the data that went into the K99 and K00 climatologies. This would save a lot of discussion, which could be focused on other important features of the results. It is important to note that they are not different climatologies, but rather upgrades and hence only the latest should be used.

We agree that the L10 climatology is an update to the K99 and K00 climatologies, and thus L10 comprises the K99 and K00 data and is more reliable. However, by comparing these climatologies, we are able to demonstrate the need to resolve uncertainties in the DMS concentration fields. By demonstrating the sensitivity of the climate system to global DMS flux, we demonstrate the importance of updates like L10 (as did Mahajan et al. 2015), and the need for continued improvement to estimates of DMS concentrations. Furthermore, K99 and K00 were used in previous studies (e.g., Thomas et al., 2010, 2011; Woodhouse et al., 2010, 2013), and only recently published studies use L10. So, by including K99 and K00, we allow for the comparison of our results with other modeling studies. In order to clarify this to the reader, we have added the above argument to Section 2.2 in the revised manuscript.

Relevant changes

page 5 (line 32) - page 6 (line 4): Added "The L10 climatology is an update to the K99 and K00 climatologies, incorporating a larger set of DMS observations. By comparing these three climatologies, we can assess the consequences of using an updated DMS concentration climatology for air-sea fluxes and climate response (Mahajan et al., 2015), helping quantify the importance of continued improvements to estimates of the seawater DMS field. Furthermore, K99 and K00 have been used in a number of previous studies (e.g., Thomas et al., 2010; Thomas et al 2011; Woodhouse et al., 2010; Woodhouse et al., 2013). By including K99 and K00, we allow for the comparison of our results with those of previous studies."

3) Also, is there a specific reason the authors include the AN01 seawater DMS climatology? This climatology, in particular, does not compare well with observations of seawater DMS, which shows that estimating DMS using global fields of chlorophyll, nutrients and light is not an accurate method considering the complexity in the oceanic DMS cycle. Additionally, a similar exercise has been by the authors in another paper (Tesdal et al., Environ. Chem., 2015).

We agree that the AN01 seawater DMS climatology does not compare well with observations, as is the case for all empirical models analyzed by Tesdal et al. (2016). However, AN01

produces global mean emissions similar to L10, and while it is a bit of an outlier among the empirical models in terms of its spatial pattern, it is the most skillful relative to observations by some measures. AN01 is a useful addition in our sensitivity study, given the study's focus on the relative importance of spatial and temporal variability compared with global annual mean flux. AN01 produces global mean emissions very similar to L10 but its spatial and temporal pattern is very different. This point has been made in the revised manuscript (Section 2.2)

4) Lana et al. 2011 did a comparison of the total flux of DMS using different climatologies, similar to that presented in table 3. A recent publication, Mahajan et al. 2015 (which is not cited) has done similar comparisons to those presented in this paper, but in further detail including spatial features, quantifying the impact of a change in the DMS climatology. Woodhouse et al. (2013) have studied the effect of changing the DMS flux on regional scales. This raises the question as to what is new in this study. The differences by changing the sea-air parameterisations are not surprising, while changes due to different climatologies have been studied before using different models.

In addition to investigating the sensitivity of radiative effects on climate to different DMS climatologies, our study also focuses on a number of questions not answered by other studies. As mentioned above, the study analyzes the relationship between spatio-temporal distributions of DMS and aerosol radiative effects and addresses the question how the spatial or temporal distribution of DMS flux influences the climate system compared to the global annual mean flux. It also presents evidence that the inclusion of air-side resistance in sea-air parameterizations is a significant factor in calculating emissions and aerosols, which was remarked upon by the other referee. The reference to Mahajan et al. (2015) is greatly appreciated, and we have mentioned the results from that study in the revised manuscript. However, while Mahajan et al. (2015) consider the effect of spatial features of DMS climatologies on atmospheric properties, they do so only in the context of the differences between the K00 and L10 climatologies, and not as a comparison of the relative importance of spatial variation to atmospheric properties in general (nor do they consider the effect of temporal variation). Further, Lana et al. (2011) provide comparisons of total flux but do not include information on any other atmospheric burdens, included in Table 3 of this study. Woodhouse et al. (2013) did not look at radiative effects on climate (as this study does) but rather focused on changes in cloud condensation nuclei (CCN).

Relevant changes

page 3, lines 15-21: Added "A recent study demonstrated substantial impacts on cloud properties and the radiation budget by using the updated L10 climatology versus the previous K00 (Mahajan et al., 2015). Even though L10 represents a substantial extension of K00, the spatial and temporal patterns are still fairly similar compared to most other available DMS reconstructions (Tesdal et al., 2016). However, the substitution of L10 for K00 results in increases of the aerosol effect on net TOA radiation of about 20%. The potentially large sensitivity of climate response to DMS climatology demonstrated by Mahajan et al. (2015) motivates the question of how important spatial and temporal variations are to atmospheric properties and radiative effects relative to changes in the global total DMS flux."

5) Why do the authors focus only on global and annual means? Considering the large spatial and seasonal differences between the climatologies, one would expect large regional difference (e.g., Mahajan et al., 2015) and these features would be smoothed out by taking global and annual means. This could be a reason for the linear dependence of the results, when in reality regions with different background aerosol loading, or relatively lower anthropogenic SO₂ emission would show highly nonlinear dependence considering the multistep chemistry-physics interactions to go from seawater DMS to cloud properties.

The focus of our analysis is on impacts of DMS emissions on the global and annual mean radiation budget. Global and annual mean radiative effects of aerosols can be used to estimate changes in global energy budgets and temperature changes in the climate system. They thereby provide a simple basis for quantifying aspects of the climate response to imposed forcing agents, especially global mean temperature, and hence are widely used in the scientific community (Myhre et al., 2013).

We did not attempt to analyze regional and sub-annual variations. These are much more difficult to analyze in a statistically robust way than global and annual mean effects owing to natural variations in regional climate processes. In general, relationships between variations in radiative effects and temperature responses are highly non-linear and non-trivial in the climate system. They are addressed in only relatively few studies, often under highly idealized conditions. We therefore consider an analysis of these relationships to be well beyond the scope of this study.

6) Does the model include ternary nucleation of H₂SO₄ and MSA with organics, which have recently been estimated to play a major role in the conversion of DMS to aerosols? (Dawson, PNAS et al., 2012; Riccobono et al., Science, 2014).

The model does not employ any parameterization of aerosol nucleation. Only bulk aerosol parameterizations are used, as described in the paper, and in more detail by von Salzen et al. (2013). The bulk aerosol scheme is a well-documented approach which produces results that are relatively straightforward to analyze. Similar bulk schemes have been widely used in different

models in the past and are used in some CMIP5 models. On the basis of the few available studies that discuss results from different models with bulk and microphysics schemes, we do not see evidence for considerable improvements in radiative forcing estimates based on simulations with microphysics schemes relative to bulk schemes (Schulz et al., 2006; Koch et al., 2009; Quaas et al., 2009). The manuscript has been revised to make this point (see end of Section 2.1 in revised manuscript).

Relevant changes

page 5, lines 19-22: Added "The bulk scheme considered in this study is simpler than approaches that consider aerosol microphysics in detail (e.g., Bellouin et al., 2013). However, based on the few available studies comparing results from different models with bulk and microphysics schemes, we do not see evidence of considerable improvement in radiative forcing estimates based on simulations with microphysics schemes relative to bulk schemes (Schulz et al., 2006; Koch et al., 2009; Quaas et al., 2009)."

References

- Kloster, S., Six, K. D., Feichter, J., Maier-Reimer, E., Roeckner, E., Wetzzel, P., Stier, P., and Esch, M.: Response of dimethylsulfide (DMS) in the ocean and atmosphere to global warming, *J. Geophys. Res.-Biogeosci.*, 112, G03005, doi:10.1029/2006JG000224, 2007.
- Koch, D., Schulz, M., Kinne, S., McNaughton, C., Spackman, J. R., Balkanski, Y., Bauer, S., Bernsten, T., Bond, T. C., Boucher, O., Chin, M., Clarke, A., De Luca, N., Dentener, F., Diehl, T., Dubovik, O., Easter, R., Fahey, D. W., Feichter, J., Fillmore, D., Freitag, S., Ghan, S. J., Ginoux, P., Gong, S., Horowitz, L., Iversen, T., Kirkevåg, A., Klimont, Z., Kondo, Y., Krol, M., Liu, X., Miller, R., Montanaro, V., Moteki, N., Myhre, G., Penner, J. E., Perlwitz, J., Pitari, G., Reddy, S., Sahu, L., Sakamoto, H., Schuster, G., Schwarz, J. P., Seland, Ø., Stier, P., Takegawa, N., Takemura, T., Textor, C., van Aardenne, J. A., and Zhao, Y.: Evaluation of black carbon estimations in global aerosol models, *Atmos. Chem. Phys.*, 9, 9001–9026, doi:10.5194/acp-9-9001-2009, 2009.
- Lana, A., Bell, T. G., Simó, R., Vallina, S. M., Ballabrera-Poy, J., Kettle, A. J., Dachs, J., Bopp, L., Saltzman, E. S., Stefels, J., Johnson, J. E., and Liss, P. S.: An updated climatology of surface dimethylsulfide concentrations and emission fluxes in the global ocean, *Global Biogeochem. Cy.*, 25, GB1004, doi:10.1029/2010GB003850, 2011.
- Ma, X., von Salzen, K., and Cole, J.: Constraints on interactions between aerosols and clouds on a global scale from a combination of MODIS-CERES satellite data and climate simulations, *Atmos. Chem. Phys.*, 10, 9851–9861, doi:10.5194/acp-10-9851-2010, 2010.
- Mahajan, A. S., Fadnavis, S., Thomas, M. A., Pozzoli, L., Gupta, S., Royer, S.-J., Saiz-Lopez, A., and Simó, R.: Quantifying the impacts of an updated global dimethyl sulfide climatology on cloud microphysics and aerosol radiative forcing, *J. Geophys. Res.-Atmos.*, 120, 2524–2536, doi:10.1002/2014JD022687, 2015.
- Mungall, E. L., Croft, B., Lizotte, M., Thomas, J. L., Murphy, J. G., Levasseur, M., Martin, R. V., Wentzell, J. J. B., Liggio, J., and Abbatt, J. P. D.: Dimethyl sulfide in the summertime Arctic atmosphere: measurements and source sensitivity simulations. *Atmos. Chem. Phys.*, 16(11), 6665–6680, doi:10.5194/acp-16-6665-2016, 2016.
- Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestad, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T., and Zhang, H.: Anthropogenic and natural radiative forcing, in: *Climate Change 2013: The physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- Quaas, J., Ming, Y., Menon, S., Takemura, T., Wang, M., Penner, J. E., Gettelman, A., Lohmann, U., Bellouin, N., Boucher, O., Sayer, A. M., Thomas, G. E., McComiskey, A., Feingold, G., Hoose, C., Kristjánsson, J. E., Liu, X., Balkanski, Y., Donner, L. J., Ginoux,

- P. A., Stier, P., Grandey, B., Feichter, J., Sednev, I., Bauer, S. E., Koch, D., Grainger, R. G., Kirkevåg, A., Iversen, T., Seland, Ø., Easter, R., Ghan, S., Rasch, P. J., Morrison, H., Lamarque, J.-F., Iacono, M. J., Kinne, S., and Schulz, M.: Aerosol indirect effects - general circulation model intercomparison and evaluation with satellite data, *Atmos. Chem. Phys.*, 9, 8697–8717, doi:10.5194/acp-9-8697-2009, 2009.
- Schulz, M., Textor, C., Kinne, S., Balkanski, Y., Bauer, S., Berntsen, T., Berglen, T., Boucher, O., Dentener, F., Guibert, S., Isaksen, I. S. A., Iversen, T., Koch, D., Kirkevåg, A., Liu, X., Montanaro, V., Myhre, G., Penner, J. E., Pitari, G., Reddy, S., Seland, Ø., Stier, P., and Takemura, T.: Radiative forcing by aerosols as derived from the AeroCom present-day and pre-industrial simulations, *Atmos. Chem. Phys.*, 6, 5225–5246, doi:10.5194/acp-6-5225-2006, 2006.
- Tesdal, J.-E., Christian, J. R., Monahan, A. H., and von Salzen, K.: Evaluation of diverse approaches for estimating sea-surface DMS concentration and air–sea exchange at global scale, *Environ. Chem.*, 13, 390–412, doi:10.1071/EN14255, 2016.
- Thomas, M. A., Suntharalingam, P., Pozzoli, L., Rast, S., Devasthale, A., Kloster, S., Feichter, J., and Lenton, T. M.: Quantification of DMS aerosol-cloud-climate interactions using the ECHAM5-HAMMOZ model in a current climate scenario, *Atmos. Chem. Phys.*, 10, 7425–7438, doi:10.5194/acp-10-7425-2010, 2010.
- Thomas, M. A., Suntharalingam, P., Pozzoli, L., Devasthale, A., Kloster, S., Rast, S., Feichter, J., and Lenton, T. M.: Rate of non-linearity in DMS aerosol-cloud-climate interactions, *Atmos. Chem. Phys.*, 11, 11175–11183, doi:10.5194/acp-11-11175-2011, 2011.
- Vallina, S. M., Simó, R., and Manizza, M.: Weak response of oceanic dimethylsulfide to upper mixing shoaling induced by global warming, *Proc. Natl. Aca. Sci. USA*, 104, 16004–16009, doi:10.1073/pnas.0700843104, 2007.
- von Salzen, K., Scinocca, J. F., McFarlane, N. A., Li, J., Cole, J. N. S., Plummer, D., Verseghy, D., Reader, M. C., Ma, X., Lazare, M., and Solheim, L.: The Canadian Fourth Generation Atmospheric Global Climate Model (CanAM4). Part I: Representation of physical processes, *Atmosphere-Ocean*, 51, 104–125, doi:10.1080/07055900.2012.755610, 2013.
- Woodhouse, M. T., Carslaw, K. S., Mann, G. W., Vallina, S. M., Vogt, M., Halloran, P. R., and Boucher, O.: Low sensitivity of cloud condensation nuclei to changes in the sea-air flux of dimethyl-sulphide, *Atmos. Chem. Phys.*, 10, 7545–7559, doi:10.5194/acp-10-7545-2010, 2010.
- Woodhouse, M. T., Mann, G. W., Carslaw, K. S., and Boucher, O.: Sensitivity of cloud condensation nuclei to regional changes in dimethyl-sulphide emissions, *Atmos. Chem. Phys.*, 13, 2723–2733, doi:10.5194/acp-13-2723-2013, 2013.

~~Sensitivity of modelled sulfate radiative forcing to DMS concentration and air-sea flux formulation~~ Sensitivity of modelled sulfate aerosol and its radiative effect on climate to ocean DMS concentration and air-sea flux

J.-E. Tesdal^{1,a}, J. R. Christian^{2,3}, A. H. Monahan¹, and K. von Salzen³

¹School of Earth and Ocean Sciences, University of Victoria, Victoria, British Columbia, Canada

²Fisheries and Oceans Canada, Sidney, British Columbia, Canada

³Canadian Centre for Climate Modelling and Analysis, Environment Canada, Victoria, British Columbia, Canada

^anow at: Lamont-Doherty Earth Observatory, Columbia University, Palisades, New York, USA

Correspondence to: J.-E. Tesdal (tesdal@ldeo.columbia.edu)

Abstract.

Dimethylsulfide (DMS) is a well-known marine trace gas that is emitted from the ocean and subsequently oxidizes to sulfate in the atmosphere. Sulfate aerosols in the atmosphere have direct and indirect effects on the amount of solar radiation reaching the Earth's surface. Thus, as a potential source of sulfate, ocean efflux of DMS needs to be accounted for in climate studies. Seawater concentration of DMS is highly variable in space and time, which in turn leads to high spatial and temporal variability of ocean DMS emissions. Because of sparse sampling (in both space and time), large uncertainties remain regarding ocean DMS concentration. In this study, we use an atmospheric general circulation model with explicit aerosol chemistry (CanAM4.1) and several climatologies of surface ocean DMS concentration to assess uncertainties about the climate impact of ocean DMS efflux. Despite substantial variation in the spatial pattern and seasonal evolution of simulated DMS fluxes, the global mean radiative ~~forcing~~ effect of sulfate is approximately linearly proportional to the global mean surface flux of DMS; the spatial and temporal distribution of ocean DMS efflux has only a minor effect on the global radiation ~~balance~~ budget. The effect of the spatial structure, however, generates statistically significant changes in the global mean concentrations of some aerosol species. The effect of seasonality on ~~net-radiative forcing~~ the net radiative effect is larger than that of spatial distribution, and is significant at global scale.

15 **1 Introduction**

The global shortwave radiation budget is influenced by sulfate aerosols in two ways: directly via scattering, and indirectly through changes to the radiative properties of clouds (as sulfate droplets act as cloud condensation nuclei, CCN) (Charlson et al., 1987; Andreae and Crutzen, 1997). An important natural source of atmospheric sulfate is the oxidation of biogenic dimethylsulfide (DMS) which has outgassed from the ocean surface (Andreae and Raemdonck, 1983; Bates et al., 1992).
20 Particular interest in the role of DMS in the atmospheric sulfur cycle arose following the hypothesis by Charlson et al.

(1987) of a negative feedback on ocean surface temperature changes mediated by cloud albedo and phytoplankton productivity: the so-called “CLAW hypothesis”. However, subsequent studies have suggested that the influence of DMS on CCN formation may be weak (Quinn and Bates, 2011; Woodhouse et al., 2010, 2013) and that the associated albedo changes are uncertain (Stevens and Feingold, 2009). Furthermore, a comprehensive understanding of the physical and biogeochemical processes that control the production of DMS and its removal from the ocean has not yet been established. The production and consumption of DMS in the water column involve a range of biotic and abiotic processes (Stefels et al., 2007). While outgassing of DMS from the ocean surface is of interest because of its climatic influence, it is a relatively minor term in the ocean DMS budget. Potentially as little as 1–10 % of ocean DMS production reaches the atmosphere (Mallin et al., 1992; Bates et al., 1994). While some model experiments have found evidence of enhanced DMS fluxes under global warming (Cameron-Smith et al., 2011; Gabric et al., 2004, 2005), others have suggested that the changes are weak (~~Bopp et al., 2003; Vallina and Simó, 2007~~) ([Bopp et al., 2003; Vallina et al., 2007](#)) or might actually be negative (Kloster et al., 2007; Six et al., 2013). While the strength and character of the influence of DMS on global climate are uncertain, little work has been done to quantify the contribution of individual components of this uncertainty. The present study uses a comprehensive global atmospheric circulation model to quantify the uncertainty associated with surface concentration fields of DMS and [surface-air-sea](#) flux parameterizations.

Kettle et al. (1999) (K99) compiled a global DMS database for the development of DMS climatologies and of parameterizations for use in modelling studies (Halloran et al., 2010). However, spatial and temporal variations in DMS concentration are not well constrained by this database, because the number of available observations is still relatively small. There are large temporal and spatial variations in the sea surface concentration of DMS (Asher et al., 2011; Tortell, 2005; Tortell et al., 2011), and the current observational dataset provides only sparse information from wide expanses of the ocean. In the absence of measurements uniformly distributed in space and time to fully characterize its spatial and temporal variability, interpolation and extrapolation schemes are required to construct continuous, observationally-based global fields of DMS ~~concentrations~~ [concentration](#) (Kettle et al., 1999; Lana et al., 2011). While the estimates generally indicate continuously elevated concentrations in tropical latitudes in contrast to low winter and high summer concentrations in middle and high latitudes, these fields remain highly uncertain due to inadequate sampling. For example, observationally-based climatologies such as those of K99, and Lana et al. (2011) (L10, released in 2010) show “bulls-eye” maxima that likely do not reflect the real distribution of DMS. The range of possible surface DMS fields increases when climatologies based on diagnostic or prognostic models are considered (Tesdal et al., 2016).

The parameterization of air-sea fluxes is also uncertain. Several different parameterizations of the piston velocity in terms of wind speed have been used in modelling studies (e.g. Liss and Merlivat, 1986; Wanninkhof, 1992; Nightingale et al., 2000), leading to substantially different flux fields for a given concentration field (Tesdal et al., 2016). Furthermore, it has been found that neglect of air-side resistance in the flux formulation (as is often done) can change estimates of fluxes by about 10 % (McGillis et al., 2000; Tesdal et al., 2016). The large differences in DMS sea surface concentration fields between different climatologies and in flux parameterizations can cause substantial variation in estimated fluxes (Tesdal et al., 2016).

An important question is how the uncertainty in fluxes translates into an uncertainty in the climate response. Although DMS fields show large differences in spatial pattern and seasonality, the differences in global- and annual-mean fluxes are considerably smaller. As well, the climatic significance of relatively small-scale concentration features remains uncertain, given the large-scale structure of the winds which drive the fluxes and the subsequent transport and oxidation to sulfate aerosol.

5 In the following, net changes in the energy budget at the top of the atmosphere (TOA) due to changes in concentrations of DMS-derived sulfate will be referred to as the radiative effect, which is sometimes also referred to as radiative forcing.

Comprehensive atmospheric general circulation models are the natural tool for assessing the uncertainty in the climatic influence of oceanic DMS fluxes. Using different DMS concentration fields as boundary conditions, the resulting changes in the atmospheric burden of sulfur species and radiative ~~forcing~~ effect can be assessed. Previous modelling studies have
10 focused on the effect of DMS on aerosol, CCN and ~~radiative forcing~~ TOA radiation budget by scaling a single DMS field (~~e.g., Gunson et al., 2006; Thomas et al., 2011~~) (e.g., Gunson et al. 2006; Thomas et al. 2011). Using a coupled atmosphere-ocean model, Grandey and Wang (2015) estimated a radiative perturbation of -1.3 to -1.5 W m^{-2} when increasing global total
15 DMS flux from 18 to 46 TgS y^{-1} . There has not been much discussion of the climatic effect of differences in the spatial and temporal structure of DMS flux (Woodhouse et al., 2013).

15 A recent study demonstrated substantial impacts on cloud properties and the radiation budget by using the updated L10 climatology versus the previous K00 (Mahajan et al., 2015). Even though L10 represents a substantial extension of K00, the spatial and temporal patterns are still fairly similar compared to most other available DMS reconstructions (Tesdal et al., 2016). However, the substitution of L10 for K00 results in increases of the aerosol effect on net TOA radiation of about 20%. The potentially large sensitivity of climate response to DMS climatology demonstrated by Mahajan et al. (2015) motivates the ques-
20 tion of how important spatial and temporal variations are to atmospheric properties and radiative effects relative to changes in the global total DMS flux. This question is addressed in the present study using the fourth generation ~~of the~~ Canadian Atmospheric Global Climate Model (~~AGCM~~), CanAM4.1).

Previous simulations with the Canadian ~~AGCM atmosphere model~~ used ocean emissions of DMS calculated from one specific climatology (K99) and one gas flux parameterization (~~that of Liss and Merlivat, 1986~~), that of Liss and Merlivat (1986)
25 (LM86). In this study, we assess the uncertainty in the climatic influence of DMS with simulations ~~of~~ with CanAM4.1 using different surface concentration climatologies and flux parameterizations. As our baseline reference, we use the recently-developed observationally-based climatology of L10. These simulation results are compared to those obtained using three different climatologies: K99, an updated version of K99 from Kettle and Andreae (2000) (K00), and the empirical model of Anderson et al. (2001) (AN01) ~~which was shown in~~. AN01 was shown by Tesdal et al. (2016) to produce global-mean DMS fluxes sim-
30 ilar to those associated with the observationally-based climatologies. To further assess the importance of spatial and temporal structure in the DMS concentration fields, simulations were carried out with the L10 climatology replaced with its spatial mean (retaining month-to-month changes) and with its annual mean (retaining spatial variability). Two flux parameterizations are considered: LM86 and that of Nightingale et al. (2000) (N00). Section 2 describes the AGCM and the details of the numerical experiments. The results of the simulations are presented in ~~Seet.~~ Section 3, followed by a discussion of the results in
35 ~~Seet.~~ Section 4. Conclusions are presented in ~~Seet.~~ Section 5.

2 Methods

2.1 Model description

All model simulations presented in this study were made with the ~~fourth-generation Canadian Atmosphere~~ [fourth generation Canadian Atmospheric Global Climate](#) Model (CanAM4.1), the atmosphere component of the Canadian Earth System Model (~~CanESM2~~). CanAM4.1 is a slightly newer version of CanAM4 (von Salzen et al., 2013) with improved diagnostic capabilities. Model dynamics are computed spectrally with a horizontal resolution of T63, equivalent to a 128×64 linear grid. The model has 49 layers in the vertical extending from the surface to 1 hPa, with a spacing of about 100 m at the surface and increasing monotonically at higher altitudes.

Figure 1 presents a schematic of the sulfur cycle and the radiative effects of sulfate aerosols as represented in CanAM4.1. The ocean efflux of DMS is a source of aerosols via oxidation to sulfur dioxide (SO_2), which in turn is oxidized to form sulfate (SO_4^{2-}). The air-sea gas transfer of DMS is calculated with wind speed from the model, while ice cover and sea surface temperature (SST) are specified using a climatological dataset from the Atmospheric Model Intercomparison Project (AMIP) (Hurrell et al., 2008). In addition to the ocean source, the model also accounts for DMS fluxes from the terrestrial biosphere using specified monthly-mean fields (Spiro et al., 1992). Besides DMS, the model also includes additional terrestrial sources of sulfur to the atmosphere: monthly mean emissions of gas phase SO_2 from fires (i.e., biomass burning) and anthropogenic sources, as well as volcanic emissions (Dentener et al., 2006). Anthropogenic aerosol and aerosol precursor emissions are used based on ~~the~~ Representative Concentration Pathway ([RCP4.5](#)) ~~scenario~~ [RCP 4.5](#) from the fifth Coupled Model Intercomparison Project (CMIP5; Lamarque et al., 2010; Moss et al., 2010).

Transport, dry and wet deposition, and chemical transformations of sulfur species are all accounted for in CanAM4.1 (von Salzen et al., 2013). DMS is oxidized to SO_2 by hydroxyl radicals (OH) during daylight hours and by nitrate radicals (NO_3) at night. Sulfate aerosol (SO_4^{2-}) production is modelled by in-cloud and gas-phase (clear-sky) oxidation of SO_2 . In-cloud production is treated differently in layer and convective clouds. The presence of ozone (O_3) and hydrogen peroxide (H_2O_2) as oxidants is a requirement for both types of clouds, and oxidation rates are modelled as pH-dependent (von Salzen et al., 2000). The in-cloud oxidation rate in deep convective clouds is calculated in proportion to the cloud fraction, which is determined based on Slingo (1987). As CanAM4.1 does not have a fully interactive chemical transport module, it uses specified oxidant concentrations (OH, NO_3 , O_3 , H_2O_2) from the Model for Ozone and Related Chemical Tracers (MOZART Brasseur et al., 1998). Ammonia (NH_3) and ammonium (NH_4^+) concentration fields are also specified (Dentener and Crutzen, 1994).

The removal of sulfate aerosol takes place through wet and dry deposition. The dry deposition flux of sulfate simply depends on the concentration within the model layer adjacent to the surface along with a defined dry deposition velocity (Lohmann et al., 1999). Wet deposition, as with the in-cloud oxidation outlined above, is treated separately for layer and convective clouds. Within convective clouds, scavenging is modelled as a function of precipitation (von Salzen et al., 2000). Wet deposition fluxes from in-cloud scavenging of aerosols in layer clouds depend on local rates of conversion of cloud water to rainwater (Croft et al., 2005). Scavenging by falling rain droplets beneath convective clouds is parameterized using a mean collection efficiency (Berge, 1993).

CanAM4.1 accounts for sulfate aerosol, organic carbon aerosol, black carbon, sea salt, and dust as separate species using a bulk aerosol scheme (Lohmann et al., 1999; Croft et al., 2005). In the CanAM4.1 version used in this study, the cloud droplet number concentration (CDNC) depends only on the local concentration of sulfate. The empirical parameterization of Dufresne et al. (2005) is used. This parameterization relates CDNC to the concentration of sulfate as:

$$5 \quad \text{CDNC} = 60[\text{SO}_4^{2-}]^{0.2}, \quad (1)$$

where CDNC is in number per cm^{-3} and $[\text{SO}_4^{2-}]$ is the sulfate concentration in $\mu\text{g m}^{-3}$. For this relationship, a lower bound on CDNC of 1 cm^{-3} is used.

CanAM4.1 calculates the direct radiative effect of scattering by aerosols and the first indirect radiative effect in which cloud optical properties are influenced by aerosol concentrations. Effects of aerosols on the conversion of cloud water to precipitation (second indirect effect) are not considered in the current version of CanAM4.1. Direct effect calculations account for scattering and absorption using Mie theory. These processes depend on aerosol mass and relative humidity: sulfate aerosols scatter radiation more efficiently at higher relative humidity as they swell in size to establish thermodynamic equilibrium according to Raoult's law. The overall efficiency of the scattering effect also varies with wavelength and aerosol concentration. The first indirect effect is computed by determining the effective radius of cloud droplets based on the relationship between sulfate aerosol and CDNC described above. Smaller droplets are more efficient at scattering solar radiation than larger droplets. Given the much greater cloud fraction of layer (stratiform) clouds compared to convective clouds, the indirect effect is only applied in layer clouds. Within each model grid cell, the cloud forcing is determined as the difference between the ~~total radiative forcing~~ net radiative fluxes for all-sky and clear-sky ~~forcing~~ only.

The bulk scheme considered in this study is simpler than approaches that consider aerosol microphysics in detail (e.g., Bellouin et al., 2013). However, based on the few available studies comparing results from different models with bulk and microphysics schemes, we do not see evidence of considerable improvement in radiative forcing estimates based on simulations with microphysics schemes relative to bulk schemes (Schulz et al., 2006; Koch et al., 2009; Quaas et al., 2009).

2.2 Description of the model experiments

A series of model experiments was conducted to investigate the effects of different sea surface DMS climatologies and gas transfer formulations on ~~radiative forcing net TOA radiation~~ radiative forcing net TOA radiation and the atmospheric burdens of DMS, SO_2 , and sulfate aerosol. These experiments are listed in Table 1. The surface concentration fields considered are the observationally-derived K99, K00, and L10 and the empirical algorithm AN01, which computes DMS concentration from chlorophyll ~~and~~ and nutrient concentrations and solar irradiance (Anderson et al., 2001). Of the various diagnostic and prognostic models of DMS used in global models, AN01 was found to produce global-mean DMS fluxes closest to L10 (although the spatial structures of the fluxes differ considerably; Tesdal et al., 2016). As well, we consider simulations with the L10 climatology replaced by its spatial mean (but retaining the seasonal cycle) and with L10 replaced by its annual mean (retaining the spatial structure).

The L10 climatology is an update to the K99 and K00 climatologies, incorporating a larger set of DMS observations. By comparing these three climatologies, we can assess the consequences of using an updated DMS concentration climatol-

[ogy for air-sea fluxes and climate response \(Mahajan et al., 2015\), helping quantify the importance of continued improvements to estimates of the seawater DMS field. Furthermore, K99 and K00 have been used in a number of previous studies \(e.g., Thomas et al., 2010, 2011; Woodhouse et al., 2010, 2013\). By including K99 and K00, we allow for the comparison of our results with those of previous studies.](#)

5 Because the wind speed and DMS concentration are correlated, the fluxes associated with the temporally invariant or spatially uniform concentration fields do not equal the global mean flux associated with the spatially and temporally varying concentration. Because we wish to distinguish the direct climatic influence of spatial and temporal structure in DMS fluxes from the global-mean flux, the temporally or spatially uniform DMS concentration fields were rescaled to produce the same total flux as ~~model simulations with temporally or spatially varying concentrations~~[the reference simulation](#). The scaling factors were determined with offline calculations using ERA-Interim reanalysis wind, sea ice, and SST (Dee et al., 2011). For 10 the temporally-invariant run, a single scaling factor was determined, while for the spatially uniform case scaling factors were determined for each monthly field. Two additional simulations were conducted with spatial and temporal patterns given by climatologies other than L10 (K99 and AN01), but scaled to have the same global mean flux as L10 (Table 1).

15 The DMS flux formulations considered are L86 and N00 (Liss and Merlivat, 1986; Nightingale et al., 2000). For N00, we conducted simulations with and without air-side resistance (γ_a) accounted for in the flux formulation. A detailed discussion of different DMS concentration climatologies and flux formulations is presented in Tesdal et al. (2016).

20 The control simulation (L10 & N00 & γ_a) was carried out using the L10 DMS concentration field with the N00 wind parameterization scheme and accounting for air resistance (Nightingale et al., 2000; Tesdal et al., 2016). The L10 climatology was used for the control simulation as it is in closest agreement with the observational database, including observations made since it was developed (Tesdal et al., 2016).

All DMS concentration fields were prepared offline before model simulations were carried out. The AN01 climatology was constructed using observed chlorophyll, light, and nutrient fields (as outlined in Tesdal et al., 2016). Differences between the model runs result from differences in DMS concentration fields, flux parameterizations, and internal variability in the model. Other aspects of the model, such as oxidation pathways and cloud microphysics, are the same for all model experiments.

25 In order to assess internal variability, an ensemble of three 5-year long runs were produced for each model configuration. [Each ensemble member uses the exact same model configuration, but in each a different seed was used in the random number generator used in the radiation code.](#) Ensemble averages are statistically more robust estimates of the climate influence of DMS than any individual member of the ensemble. The spread among ~~realizations~~[ensemble members](#) indicates the magnitude of the response to changes in DMS fluxes relative to internal variability. All simulations are carried out ~~are~~ for the period from 30 January 2003 to December 2008, with the first year discarded as a spin-up period.

3 Results

3.1 [Assessment of simulated sulfate aerosol](#)

The assessment of simulations of natural sulfate aerosol in the marine troposphere is challenging given a lack of chemical observations in remote regions of the ocean where the contribution of DMS oxidation to sulfate aerosol concentration is most significant. Even where chemical measurements exist, the relative contribution of ocean DMS emissions to net sulfate production cannot be directly observed. In an attempt to assess model results for sulfate, observed sulfate concentrations from ship-based measurements were compiled from available data sets obtained from the NOAA PMEL Atmospheric Chemistry Data Server (<http://saga.pmel.noaa.gov/data>). Figure 2 shows a map of the cruise transects from which observational datasets were drawn. The datasets contain multiple types of sulfate, but only total non-seasalt sulfate was considered (sum of all size fractions present).

For comparisons with these data sets, simulated sulfate concentrations from the control run were compared to the measurements, matching simulated and observed near-surface concentrations according to nearest location and month of the year. This yields a correlation of 0.57 between simulated and observed concentrations (Fig. 3). The mean simulated concentration in Fig. 3 of $0.96 \pm 0.86 \mu\text{g m}^{-3}$ for all available data is lower than the corresponding observed value of $1.84 \pm 2.31 \mu\text{g m}^{-3}$ (i.e. model underestimate of 48%).

In an attempt to characterize the impact of anthropogenic pollution on the results, the data were grouped by latitude (Fig. 3a). Simulated and observed concentrations between the equator and 50°N are relatively high, which can be partly explained by contributions of anthropogenic emissions to sulfate concentrations. Without these data, the agreement between mean simulated and measured concentrations improves noticeably ($0.49 \pm 0.13 \mu\text{g m}^{-3}$ in the model versus $0.72 \pm 0.58 \mu\text{g m}^{-3}$ in observations, an underestimate of 32%, Fig. 3a).

To further characterize the impact of DMS emissions on atmospheric sulfate concentration, the fraction of sulfate from ocean DMS emissions was diagnosed based on model simulations with and without ocean DMS emissions. The diagnosed fraction from these simulations is in good overall agreement with results from Gondwe et al. (2003), which show a large influence of DMS emissions on near-surface sulfate concentration in the southern hemisphere and at high latitudes in the northern hemisphere. As shown in Fig. 3b, low sulfate concentrations tend to be associated with a high fraction of sulfur originating from DMS emissions. There is good agreement between mean simulated and observed concentrations where this fraction exceeds 40% ($0.48 \pm 0.26 \mu\text{g m}^{-3}$ in the model versus $0.72 \pm 0.55 \mu\text{g m}^{-3}$ in observations, Fig. 3a), with a Pearson correlation coefficient of 0.62.

Comparisons in Fig. 3 are influenced by the use of climatological instead of actual emissions in the model. In addition, differences between climatological and actual meteorological situations, and the relatively low spatial resolution of the model need to be considered when interpreting simulated sulfate concentrations. These factors may largely explain the lower variability in simulated concentrations compared to observations. However, even if local differences in spatial or temporal variability in sulfate concentrations exist they are unlikely to greatly influence the global climate, based on an analysis of results with variable DMS emissions in Section 3.4.

In summary, the analysis of sulfate concentrations over the ocean confirms that a substantial fraction of the sulfate concentrations in observations and the model are related to emissions of sulfur from the ocean, with good overall agreement in regions that are most strongly affected by DMS emissions. This provides evidence of realistic simulations of atmospheric DMS sources

and aerosol removal processes in the marine atmosphere. In addition, simulated relationships between sulfate aerosol concentrations and simulated cloud microphysical properties in the model agree well with relationships that are based on observed cloud properties over the ocean (Ma et al., 2010; von Salzen et al., 2013), which provides evidence for realistic responses of simulated radiative effects of sulfate aerosol to DMS emissions in the model.

5 3. 1 2 Comparison between model and reanalysis flux estimates

Before analyzing the climatic influence of differences in DMS fluxes, we will compare the ~~global and annual mean DMS flux~~ global- and annual-mean DMS fluxes in the different simulations to the fluxes calculated with the ERA-Interim reanalysis SST, sea ice, and wind speed fields over the same time period as the model simulations (Table 2). The global- and annual-mean flux is generally higher in the CanAM4.1 simulation than ~~the value resulting from the reanalysis fields~~ when reanalysis fields
10 are used: it is 22–24 % larger with N00 (with or without air resistance) and 14 % larger with LM86. These differences must result primarily from differences in the wind fields, because SST and sea ice cover are specified in all simulations with AMIP boundary conditions ~~that should be~~ and are very similar to the ERA-Interim fields. The winds are overall somewhat stronger in the model than in the reanalysis product: the annual mean surface wind speed is 17 % higher on average in CanAM4.1. The ~~probability-frequency~~ probability-frequency distribution and seasonality of the winds also differ slightly between the model and observations (not
15 shown). Fluxes are particularly sensitive to high wind speeds, and slight changes in the wind distribution can be magnified in the DMS flux. Consistent with the results of Tesdal et al. (2016), the DMS flux calculated with the L10 DMS concentration field is higher than that calculated with K99 or K00, independent of which gas transfer formulation is used.

Because the DMS fluxes computed with the model wind fields differ substantially from those computed by reanalysis winds, we expect the simulated climatic influence of DMS to be biased. ~~Because~~ However, as our focus is on the sensitivity of the
20 climatic influence of DMS to changes in DMS fluxes, ~~rather than the absolute strength of the effect, this model bias is not~~ expected to affect our results.

3. 2 3 Fluxes and atmospheric sulfur burdens

Changes in the DMS concentration climatology and the flux formulation result in substantial changes in the global mean flux (~~the sum of including~~ both ocean and terrestrial sources; Table 3). The change relative to the control simulation ranges
25 from a 37 % reduction using K99 and LM86 to an 8 % increase when neglecting air-side resistance. The largest ensemble spread in DMS emissions among the simulations is less than $0.06 \mu\text{mol m}^{-2} \text{d}^{-1}$, which is negligible compared to the overall range of DMS emissions of the different model runs ($3.15 \mu\text{mol m}^{-2} \text{d}^{-1}$). By construction, the difference from the reference simulation is negligible in the temporally-invariant and spatially-uniform simulations and in the simulations with the rescaled concentration fields K99* and AN01* (Table 1).

30 The magnitudes of the simulated sulfur sources, sinks, and atmospheric burdens are also presented in Table 3. The budgets of sulfur species are very close to equilibrium in all simulations (sources approximately equal sinks). The reduction in DMS emission for simulations using K99 relative to those using L10 results in a reduction in daytime oxidation by OH, while nighttime oxidation by NO₃ does not change much. In contrast, both daytime and nighttime oxidation rates are affected equally

when L10 is replaced with K00. The responses of oxidation rates to changes in DMS concentration patterns likely result from the distribution of the oxidants OH and NO₃, which are specified in CanAM4.1.

The relationship between changes in the simulated atmospheric burdens of sulfur species and changes in DMS flux is approximately linear (Table 3). The largest changes occur in the DMS burden: the difference of ~0.1 TgS (61 %) between L10 & N00 and K99 & LM86 is close to the difference in DMS flux (68 %) between these two simulations. The relative changes of SO₂ and sulfate burdens are smaller than those of DMS because of the large background value for SO₂ and sulfate from other sources (anthropogenic and volcanic).

The relationships of DMS, SO₂, and SO₄²⁻ burdens with DMS flux are illustrated in Fig. 2-4. There are two distinct groups of simulations, depending on which DMS field is used. Regression lines computed for simulations with L10 (blue) and with K99 (purple) are almost parallel, indicating an approximately constant offset in burden between the K99 and L10 simulations. The sensitivity of atmospheric burdens of sulfur species to the spatial and temporal structure of DMS concentration is much smaller than to the global mean flux.

3. 3.4 Relationship between radiative ~~foreing~~effects, sulfate and DMS

To a first approximation, the relationship between ~~top-of-the-atmosphere (TOA) radiative forcing~~ TOA net radiation and the global mean flux of DMS is linear (Fig. 3.5). Deviations from that linear relationship can be attributed to differences in spatial and temporal distribution among the DMS fields (~~or internal variability~~). As with atmospheric SO₂ and SO₄²⁻ burden, the relationship between the radiation fields and DMS flux can be divided into two classes of simulations using K99 or L10. The response of the radiative ~~foreing~~ effect to differences in flux is smaller for K99-based simulations than for those based on L10. K99 generally has larger radiative ~~foreing~~ effect relative to the better-constrained L10, and this difference increases with increasing flux (i.e., with increasing wind speed and/or gas exchange coefficient).

Figure 3-5 shows that there is considerable variation in TOA ~~radiative forcing~~ net radiation depending on the strength of the ocean DMS source. Across the experiments, the range ~~in-ensemble-mean-radiative-response~~ is 0.67 W m⁻² (among ensemble means). The sensitivity to air-sea flux parameterization is particularly strong: the difference between LM86 and N00 in average flux (and thus in radiative ~~foreing~~ effect) is greater than the difference among DMS concentration fields considered. The difference in net radiation between K00 and L10 is 0.33 W m⁻², very similar to the value of 0.3 W m⁻² estimated by Mahajan et al. (2015).

The spread of the individual ensemble members in Fig. 3-5 indicates the uncertainty in ~~radiative forcing~~ the radiation budget resulting from model internal variability over the 5-year period of the simulations, independent of the boundary conditions. This spread is on average 0.12 W m⁻² (ranging from 0.04 to 0.19 W m⁻²); (compared to a range of ~~ensemble means of~~ 0.67 W m⁻² across experiments).

The DMS concentration fields considered in this analysis are a relatively narrow subset of the observationally-based or modelled climatologies considered in Tesdal et al. (2016). Use of some of these very different concentration fields would be expected to result in substantially different ~~radiative forcing~~ effects on the atmospheric radiation budget. A linear regression model constructed from the subset of simulations using N00 & γ_a was used to obtain an estimate of the possible range ~~in-ra-~~

diative forcing of radiative perturbations corresponding to the entire range of DMS climatologies (Fig. 46). Offline reanalysis-based DMS fluxes were used in the estimation of DMS radiative forcing effects for those climatologies for which DMS fluxes from CanAM4.1 were not available. The range of radiative forcing perturbations to net TOA radiation across the different DMS climatologies with the same flux formulation is 0.75 W m^{-2} , with L10 at the lower end since it produces the largest flux.

5 A similar estimate can be made for variation among the available piston velocity schemes, constructing the linear regression with model runs that have the same DMS field but different air-sea flux parameterizations (not shown). Using L10 as the DMS field and considering flux estimates obtained using N00, LM86, and a third parameterization of Wanninkhof (1992) (W92) produces a radiative forcing range of 1.04 W m^{-2} .

10 Irrespective of differences in the spatial and temporal patterns of the DMS concentration field, the relationship between radiative forcing net TOA radiation and atmospheric sulfate burden is close to linear (Fig. 57). There is no evidence of distinct relationships depending on use of the L10 or K99 climatologies as seen in the relationship between DMS flux and TOA radiative forcing radiation. Evidently, these differences are associated with spatial and temporal differences in the oxidation of DMS to sulfate (Fig. 24).

15 Global means of individual radiation fields (shortwave cloud forcing, TOA clear-sky reflected flux, and TOA total reflected flux) are plotted against global mean DMS flux and global mean sulfate burden in Fig. 6-8. TOA clear-sky reflected flux represents the direct aerosol radiative effects, while shortwave cloud forcing represents the first indirect effect. In these simulations, the direct and first indirect effects are approximately equally sensitive to changes in DMS flux (or sulfate burden). The response of all-sky TOA total reflected flux to changes in global mean DMS flux and atmospheric sulfate burden (Fig. 68) is similar to the total radiative forcing effect (Figs. 3 and 5 and 7), and the range in total reflected flux is as large as that of total forcing among the different simulations. As the total radiative forcing effect includes variation in longwave radiation while the reflected solar flux accounts only for shortwave radiation, our results confirm that radiative forcing the radiative effects associated with DMS is are primarily in the shortwave.

25 The internal variability in either cloud forcing or clear-sky reflected flux is generally larger than in the total reflected flux (which is approximately the sum of the first two) (Fig. 68). While the overall radiative impact effect of DMS fluxes in the model is estimated with reasonable precision with these experiments, larger ensembles or longer integrations may be required to achieve the same level of precision for the different components of the radiative response.

3. 45 The effect of DMS spatial and temporal structure on aerosols and radiative forcing effects

30 Suppressing either the spatial or temporal variability of ocean DMS concentration changes the resulting radiative forcing concentration of sulfate aerosol and its effect on the TOA radiation budget (Figs. 3 and 65 and 8). While these changes are small, the ensemble spreads indicate that in some cases they are robust. The changes in global mean DMS flux, oxidation rates, sulfur species burdens, and radiative responses components of the TOA radiation budget between the control run and model runs with temporally-invariant and spatially-uniform DMS fields are shown in Fig. 7-9. For comparison, the changes from a simulation using L10 and N00 but neglecting the air resistance term are also shown.

The global mean burden of a species in the atmosphere over a given time period is determined by the efficiency of internal sources and sinks, and indirectly by the transport. The time-mean state is effectively in equilibrium (Table 3), so global sulfur budgets are a simple sum over all internal sources, sinks, and fluxes between sulfur species. However, a balanced budget can be achieved with different values of the individual source and sink terms. The global rates of individual flux or sink processes are determined by the spatial and temporal relationships among the chemical species involved.

By construction, the spatially-uniform and temporally-invariant DMS concentration fields yield global-mean DMS fluxes that differ only slightly from the control simulation. However, there are substantial and statistically robust changes in the sink strengths (Fig. 9). The absence of spatial or temporal structure in the DMS concentration fields has different effects during day and night: daytime oxidation of DMS by OH is decreased in these simulations, balanced by an increase in nighttime oxidation by NO₃. The simulation without air resistance shows an increase in global-mean DMS flux compared to the control of about 0.40 μmol m⁻² d⁻¹, which is balanced by an increase in both oxidation rates.

The atmospheric burdens of all sulfur species increase significantly in the simulation without air resistance. As for the simulations with spatially- or temporally-averaged DMS concentrations, only the spatially-uniform DMS simulation results in a change in oxidation patterns resulting in statistically robust increases in the burdens of both DMS and SO₂. Interestingly, the increase in SO₂ in these simulations is associated with a decrease in SO₄²⁻. A similar decrease in SO₄²⁻ burdens is also seen in the simulations with temporally-invariant DMS concentration fields, although neither the DMS nor SO₂ burdens show statistically robust changes.

For all three of these sets of simulations, there is a much stronger response in the clear-sky reflected flux than in the shortwave cloud forcing. The changes in total reflected solar flux are statistically robust for both the simulations with temporally-invariant surface concentration of DMS and those without air-side resistance. In all of these simulations, the effect on TOA cloud forcing is not significantly different from zero.

Taken together, these results indicate that the spatial and temporal distribution of DMS flux affects the aerosol direct radiative forcing effect primarily by influencing the efficiency of oxidation of DMS to SO₂ and SO₄²⁻. The effect on reflected solar fluxes of changes of in SO₄²⁻ is larger for simulations with temporally-invariant DMS concentration than for spatially-uniform concentration, despite the change in SO₄²⁻ being larger in the latter case. This fact will be discussed in more detail in the next section Discussion.

Note that the magnitudes (but not the signs) of changes in SO₄ resulting from suppressing spatial or temporal structure in the DMS concentration fields are the same as neglecting the air-side resistance term in the DMS flux formulation. Air-side resistance is often ignored in calculations of air-sea DMS fluxes. Our results indicate that the effect of neglecting this term is comparable in magnitude to the seemingly more dramatic change of entirely eliminating temporal or spatial structure in the DMS concentration fields.

4 Discussion

The results presented in [Section 3](#) demonstrate that while the magnitude of the spatial and temporal mean DMS flux is linearly related to the mean DMS burden to a good first approximation, there are deviations from this linear relationship. A simple expression for the global spatial- and temporal-mean DMS budget is

$$5 \left\langle \frac{d}{dt} \text{DMS} \right\rangle = \langle E \rangle - \langle O \times \text{DMS} \rangle, \quad (2)$$

where the angle brackets denote global space- and time-averages, E is the emission field, and O is the oxidation rate field (per unit of DMS concentration). At equilibrium, the rate of change vanishes, and

$$\langle E \rangle = \langle O \times \text{DMS} \rangle. \quad (3)$$

The upper three panels of [Fig. 7-9](#) present simulated values of $\langle E \rangle$ and $\langle O \times \text{DMS} \rangle$ for three sets of simulations (spatially-uniform, temporally-invariant, and neglecting air-side resistance).

If O and DMS did not depend on space or time, then we could decompose $\langle O \times \text{DMS} \rangle$ as $\langle O \rangle \times \langle \text{DMS} \rangle$, and an exactly linear relation between global-mean flux and global-mean atmospheric burden would exist. The deviations from this relationship evident in [Fig. 2-4](#) result from spatial and temporal correlations between the distribution of DMS and its sinks. Similarly, deviations from a purely linear relationship between spatial- and temporal-mean atmospheric burdens of SO_2 and SO_4^{2-} result from correlations between SO_2 and its oxidation rate. Atmospheric transport contributes to spatial and temporal correlations between atmospheric distributions of sulfur species and their sources and sinks. For example, some DMS emitted in the tropics will be transported by convective processes to the upper troposphere, where sinks are weaker. Similarly, the lifetime of sulfate transported to the upper troposphere is extended, as its primary sink is in low- to mid-tropospheric clouds. A detailed analysis of the spatial relationships among these processes is outside the scope of the present study.

As with the atmospheric burdens of sulfur species, the response of [mean-radiative-foreing-net TOA radiation](#) to changes in mean DMS flux is linear to a first approximation ([Fig. 35](#)), with some scatter around this relationship resulting from model internal variability and differences in the spatial and temporal structure of the DMS fluxes. ~~There are therefore~~ [Our model simulations allow us to assess the relative sizes of](#) three sources of uncertainty in the radiative ~~foreing-response-to-effect of~~ [DMS emissions: \(1\) uncertainty in total emissions, \(2\) uncertainty in spatial/temporal pattern of fluxes, and \(3\) internal variability.](#) ~~Inspection of~~ [Fig. 3-5](#) indicates that for the range of DMS climatologies and flux formulations considered, the size of the first of these uncertainties is about 0.7 W m^{-2} , while ~~that of~~ the second and third are smaller (about 0.2 W m^{-2}). While internal variability and uncertainty in spatial and temporal structure in DMS flux contribute to the overall uncertainty in the ~~radiative-re-~~ [response-net radiation budget](#), our study shows that uncertainty in the global-mean flux is the dominant contributor. [Uncertainties associated with model representations of atmospheric chemistry, cloud microphysics, and radiative transfer cannot be assessed using a single AGCM. Comparison of the magnitudes of these uncertainties to those we have considered is an interesting direction of future study.](#)

The reduction in ~~radiative-foreing-~~ [the radiative effect of DMS emissions](#) resulting from suppressing the seasonal cycle in L10 is larger than that resulting from suppressing spatial variability ([Fig. 35](#)). This is consistent with the fact that DMS

concentrations in L10 tend to be higher in ~~the~~ summer (when changes to shortwave fluxes are particularly important) at mid- to high latitudes. As atmospheric residence times of sulfur species are on the order of a few to several days and their transport is primarily zonal, DMS emitted in the mid- or high latitudes will have its strongest effect ~~on radiative forcing~~ in these latitude bands, and there will be a spatial correlation between DMS-derived sulfate aerosol concentration and aerosol radiative effects. These results suggest that for global-mean responses, resolving the correct seasonal distribution of DMS fluxes is more important than resolving the spatial distribution, although neither is as important as the global-mean flux. However, we also note that the ensembles of the spatially-uniform and temporally-invariant simulations slightly overlap and it is possible that the difference between the two is a result of internal variability.

The fact that the deviations of TOA net radiation and reflected solar flux are similar in absolute value (Figs. ~~5 and 6~~ 7 and 8) demonstrates that the climate response to DMS is dominated by shortwave fluxes. A weak response in the longwave may exist, but comparison of Figs. ~~5 and 6~~ 7 and 8 suggests that it is smaller than internal variability. Furthermore, the strongly linear relationship between the atmospheric burden of SO_4^{2-} and total radiative ~~forcing effect~~ (Fig. 57) demonstrates that simulated reductions in ~~radiative forcing net TOA radiation~~ are a direct response to ~~reduction increases~~ in the atmospheric sulfate burden. Further statements about the causal relationship between changes in DMS flux and the global radiative ~~response effect~~ are difficult because of the broad range of processes and feedbacks in the model ~~than can affect radiative fluxes~~.

Rough estimates of the range in ~~radiative forcing net TOA radiation~~ given the possible range in DMS flux are 0.75 W m^{-2} (among the range of available DMS fields) and 1.04 W m^{-2} (among all different flux parameterizations considered). Contrasting these uncertainties with the well-constrained radiative forcing of $+1.82 \pm 0.19 \text{ W m}^{-2}$ due to the increase in atmospheric CO_2 from 1750 to 2011 (Myhre et al., 2013) emphasizes the degree of uncertainty in DMS-derived aerosol forcing and the need to better constrain this quantity. Previous studies have found a relatively weak ~~influence of DMS fluxes on link between DMS fluxes and~~ climate (e.g., Woodhouse et al., 2010; Kloster et al., 2007; Vallina et al., 2007). However, these studies may have a “weak effect” bias because of a low bias in DMS flux (Fig. 46), which would translate into a low bias in ~~radiative forcing the radiative effect of DMS~~. The results of the current study show that there is a systematic deviation from the control run of up to 0.75 W m^{-2} for some DMS models and algorithms.

The uncertainty in DMS concentration estimates contributes substantially to uncertainties in present-day aerosol radiative forcing (Dentener et al., 2006; Carslaw et al., 2013), defined as the difference ~~in radiative fluxes~~ between present-day and preindustrial due to anthropogenic changes in the atmospheric aerosol burden. ~~To estimate the present-day forcing, it is necessary to have a reliable estimate of aerosol forcing in preindustrial times.~~ (Myhre et al., 2013). While observationally-based estimates ~~of present-day radiative fluxes~~ can be made for the present day, these are not available for preindustrial conditions. Current understanding of the natural sulfur cycle indicates that most preindustrial ~~forcing from sulfate aerosol was determined by DMS flux sulfate aerosol originated from DMS~~ and volcanic emissions (Carslaw et al., 2013). Uncertainty in estimates of these fluxes, which must be based on models in the absence of direct observations, will impact forcing estimates. The large uncertainty in DMS flux translates into uncertainty in preindustrial aerosol ~~forcing concentration~~, regardless of whether one assumes that DMS ~~flux remains sources remain~~ the same as or similar to preindustrial conditions. As DMS ~~fluxes emissions~~

may have changed from the preindustrial state, ~~the use of~~ using fluxes estimated from present-day conditions increases this uncertainty.

Our estimates of climatic effects of DMS obtained using CanAM4.1 could be biased due to idealized assumptions about aerosol processes and the absence of a process-based representation of the indirect aerosol effect. These biases would be expected to be especially pronounced in the parts of the atmosphere least affected by anthropogenic emissions, such as the Southern Hemisphere. ~~Future model simulations could be done with an atmosphere model that has a more physical treatment of aerosol processes and cloud microphysical properties.~~ It is possible that sensitivity to the spatial and temporal distribution of DMS would change with an improved representation of cloud microphysics. Furthermore, instead of using specified atmospheric concentrations of the oxidants, a comprehensive tropospheric chemistry scheme could be used to achieve a more realistic modelling of atmospheric DMS oxidation.

This study did not investigate climate sensitivity to DMS flux in a coupled model; all model simulations were atmosphere-only. These experiments could be repeated in a coupled model setting which would allow for the feedbacks central to the CLAW hypothesis. Furthermore, a coupled model setup would allow for the evaluation of prognostic DMS modules, as opposed to using specified (climatological) fields. Such an analysis would allow exploration of the ~~climate sensitivity~~ sensitivity of radiation and climate to specific parameters or ~~different~~ mechanisms within the prognostic DMS formulations and to distinguish this from sensitivity to other aspects of the model. Two caveats regarding such an analysis are that DMS concentration fields resulting from existing prognostic models differ substantially from observations (Tesdal et al., 2016) and that internal variability would increase due to the longer timescales of oceanic variability.

The focus of our analysis has been the influence of DMS emissions on sulfate aerosol and its radiative effects, which can be used to estimate changes in global energy budgets. These measures provide a simple basis for quantifying aspects of the climate response to imposed forcing agents, especially global mean temperature, and hence are widely used in the scientific community (Myhre et al., 2013). We did not attempt to analyze regional and sub-annual variations in radiative effects of aerosols which are more difficult to analyze in a statistically robust way because internal variability is much larger relative to the atmosphere. forced response on regional scales. In general, regional relationships between aerosols, radiation and temperature response can be complex and nonlinear. While these relationships are beyond the scope of the present study, we consider our estimates of global scale effects to be robust and relatively insensitive to regional-scale processes.

5 Conclusions

Despite more than 30 years of concerted research on the issue, fundamental uncertainties remain regarding the spatial and temporal structure of surface ocean DMS concentrations and how best to model DMS fluxes (Tesdal et al., 2016). In this study, we have used the atmospheric component of a state-of-the-art Global Climate Model (CanAM4.1) to assess the uncertainty in atmospheric sulfur burdens and ~~radiative forcing~~ their effect on the planetary radiation budget associated with uncertainties in DMS concentration fields and air-sea flux formulations. Our results indicate that to a first approximation, the global spatial and temporal mean ~~radiative forcing associated with DMS~~ effect of DMS on net TOA radiation scales linearly with the spatial and

temporal mean flux. Spatial and temporal correlations between model sulfur species (DMS, SO₂, and SO₄²⁻) and their sinks result in deviations from this linear relationship that exceed internal variability, but these deviations are relatively small. This result suggests that on a global scale, it is most important to have an accurate estimate of the global DMS flux, while resolving the exact spatial and temporal distribution is of less importance. [Neglect of air-side resistance in the flux parameterization was](#)

5 [shown to have a larger effect on net TOA radiation than suppressing spatial or temporal structure in the DMS concentration field. From the perspective of global climate, accurate formulation of surface fluxes is as or more important than accurate representation of sea surface DMS concentrations.](#)

A comprehensive view of the global scale uncertainties is important for understanding the role of DMS in the climate system. Uncertainty about the global DMS concentration translates to uncertainty about global estimates of DMS flux ~~and uncertainty~~
10 ~~in radiative forcing~~, [aerosol burdens and their radiative effects](#). These uncertainties limit the confidence with which we can make statements about the importance of ~~the role of~~ DMS in the climate system, and leave open the possibility that changes in DMS fluxes could alter future climate in as-yet-unexpected ways.

Acknowledgements. J.-E. Tesdal would like to acknowledge support from the Natural Sciences and Engineering Research Council of Canada (NSERC) CREATE Training Program in Interdisciplinary Climate Science. A. H. Monahan acknowledges support from the NSERC Discovery Grant Program. The authors also thank Kelly McCusker ~~for her~~ [and two anonymous reviewers for their](#) helpful comments on this manuscript.

The L10 data were obtained from the SOLAS website (http://www.bodc.ac.uk/solas_integration/implementation_products/group1/dms/). The K00 dataset was obtained from the NCAR Data Archive (<http://rda.ucar.edu/datasets/ds289.2/>). The data used to construct the AN01 climatology are described in detail in Tesdal et al. (2016). ERA-Interim reanalysis products were obtained from the European Centre for
20 Medium-Range Weather Forecasts (http://apps.ecmwf.int/datasets/data/interim_full_daily). [We also acknowledge the assistance from Timothy Bates and Jim Johnson in retrieving nss-SO₄²⁻ measurements from the NOAA PMEL Atmospheric Chemistry Data Server \(<http://saga.pmel.noaa.gov/data/>\).](#)

References

- Anderson, T., Spall, S., Yool, A., Cipollini, P., Challenor, P., and Fasham, M.: Global fields of sea surface dimethylsulfide predicted from chlorophyll, nutrients and light, *J. Marine Syst.*, 30, 1–20, doi:10.1016/S0924-7963(01)00028-8, 2001.
- Andreae, M. O. and Crutzen, P.: Atmospheric aerosols: Biogeochemical sources and role in atmospheric chemistry, *Science*, 276, 1052–1058, doi:10.1126/science.276.5315.1052, 1997.
- Andreae, M. O. and Raemdonck, H.: Dimethyl sulfide in the surface ocean and the marine atmosphere: A global view, *Science*, 221, 744–747, doi:10.1126/science.221.4612.744, 1983.
- Aranami, K. and Tsunogai, S.: Seasonal and regional comparison of oceanic and atmospheric dimethylsulfide in the northern North Pacific: Dilution effects on its concentration during winter, *J. Geophys. Res.-Atmos.*, 109, D12303, doi:10.1029/2003JD004288, 2004.
- 10 Asher, E. C., Merzouk, A., and Tortell, P. D.: Fine-scale spatial and temporal variability of surface water dimethylsulfide (DMS) concentrations and sea-air fluxes in the NE Subarctic Pacific, *Mar. Chem.*, 126, 63–75, doi:10.1016/j.marchem.2011.03.009, 2011.
- Aumont, O., Belviso, S., and Monfray, P.: Dimethylsulfoniopropionate (DMSP) and dimethylsulfide (DMS) sea surface distributions simulated from a global three-dimensional ocean carbon cycle model, *J. Geophys. Res.-Oceans*, 107, 3029, doi:10.1029/1999JC000111, 2002.
- Bates, T., Lamb, B., Guenther, A., Dignon, J., and Stoiber, R.: Sulfur emissions to the atmosphere from natural sources, *J. Atmos. Chem.*, 15, 315–337, doi:10.1007/BF00115242, 1992.
- Bates, T. S., Kiene, R. P., Wolfe, G. V., Matrai, P. A., Chavez, F. P., Buck, K. R., Blomquist, B. W., and Cuhel, R. L.: The cycling of sulfur in surface seawater of the Northeast Pacific, *J. Geophys. Res.-Oceans*, 99, 7835–7843, doi:10.1029/93JC02782, 1994.
- [Bellouin, N., Mann, G. W., Woodhouse, M. T., Johnson, C., Carslaw, K. S., and Dalvi, M.: Impact of the modal aerosol scheme GLOMAP-mode on aerosol forcing in the Hadley Centre Global Environmental Model, *Atmos. Chem. Phys.*, 13, 3027–3044, doi:10.5194/acp-13-3027-2013, 2013.](#)
- 20 Belviso, S., Bopp, L., Moulin, C., Orr, J., Anderson, T., Aumont, O., Chu, S., Elliott, S., Maltrud, M., and Simó, R.: Comparison of global climatological maps of sea surface dimethyl sulfide, *Global Biogeochem. Cy.*, 18, GB3013, doi:10.1029/2003GB002193, 2004.
- Belviso, S., Masotti, I., Tagliabue, A., Bopp, L., Brockmann, P., Fichot, C., Caniaux, G., Prieur, L., Ras, J., Uitz, J., Loisel, H., Dessailly, D., Alvain, S., Kasamatsu, N., and Fukuchi, M.: DMS dynamics in the most oligotrophic subtropical zones of the global ocean, *Biogeochemistry*, 110, 215–241, doi:10.1007/s10533-011-9648-1, 2012.
- Berge, E.: Coupling of wet scavenging of sulphur to clouds in a numerical weather prediction model, *Tellus Ser. B*, 45, 1–22, doi:10.1034/j.1600-0889.1993.00001.x, 1993.
- Bopp, L., Aumont, O., Belviso, S., and Monfray, P.: Potential impact of climate change on marine dimethyl sulfide emissions, *Tellus Ser. B*, 55, 11–22, doi:10.1034/j.1600-0889.2003.042.x, 2003.
- 30 Brasseur, G. P., Hauglustaine, D. A., Walters, S., Rasch, P. J., Müller, J.-F., Granier, C., and Tie, X. X.: MOZART, a global chemical transport model for ozone and related chemical tracers: 1. Model description, *J. Geophys. Res.-Atmos.*, 103, 28265–28289, doi:10.1029/98JD02397, 1998.
- Cameron-Smith, P., Elliott, S., Maltrud, M., Erickson, D., and Wingenter, O.: Changes in dimethyl sulfide oceanic distribution due to climate change, *Geophys. Res. Lett.*, 38, L07704, doi:10.1029/2011GL047069, 2011.
- 35 Carslaw, K. S., Lee, L. A., Reddington, C. L., Pringle, K. J., Rap, A., Forster, P. M., Mann, G. W., Spracklen, D. V., Woodhouse, M. T., Regayre, L. A., and Pierce, J. R.: Large contribution of natural aerosols to uncertainty in indirect forcing, *Nature*, 503, 67–71, doi:10.1038/nature12674, 2013.

- Charlson, R. J., Lovelock, J. E., Andreae, M. O., and Warren, S. G.: Oceanic phytoplankton, atmospheric sulphur, cloud albedo and climate, *Nature*, 326, 655–661, doi:10.1038/326655a0, 1987.
- Collins, W. J., Bellouin, N., Doutriaux-Boucher, M., Gedney, N., Halloran, P., Hinton, T., Hughes, J., Jones, C. D., Joshi, M., Liddicoat, S., Martin, G., O'Connor, F., Rae, J., Senior, C., Sitch, S., Totterdell, I., Wiltshire, A., and Woodward, S.: Development and evaluation of an Earth-System model – HadGEM2, *Geosci. Model Dev.*, 4, 1051–1075, doi:10.5194/gmd-4-1051-2011, 2011.
- 5 Croft, B., Lohmann, U., and von Salzen, K.: Black carbon ageing in the Canadian Centre for Climate modelling and analysis atmospheric general circulation model, *Atmos. Chem. Phys.*, 5, 1931–1949, doi:10.5194/acp-5-1931-2005, 2005.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J. J., Park, B. K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J. N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. Roy. Meteorol. Soc.*, 137, 553–597, doi:10.1002/qj.828, 2011.
- 10 Dentener, F., Kinne, S., Bond, T., Boucher, O., Cofala, J., Generoso, S., Ginoux, P., Gong, S., Hoelzemann, J. J., Ito, A., Marelli, L., Penner, J. E., Putaud, J.-P., Textor, C., Schulz, M., van der Werf, G. R., and Wilson, J.: Emissions of primary aerosol and precursor gases in the years 2000 and 1750 prescribed data-sets for AeroCom, *Atmos. Chem. Phys.*, 6, 4321–4344, doi:10.5194/acp-6-4321-2006, 2006.
- 15 Dentener, F. J. and Crutzen, P. J.: A three-dimensional model of the global ammonia cycle, *J. Atmos. Chem.*, 19, 331–369, doi:10.1007/BF00694492, 1994.
- Dufresne, J.-L., Quaas, J., Boucher, O., Denvil, S., and Fairhead, L.: Contrasts in the effects on climate of anthropogenic sulfate aerosols between the 20th and the 21st century, *Geophys. Res. Lett.*, 32, L21703, doi:10.1029/2005GL023619, 2005.
- 20 Elliott, S.: Dependence of DMS global sea-air flux distribution on transfer velocity and concentration field type, *J. Geophys. Res.-Biogeosci.*, 114, G02001, doi:10.1029/2008JG000710, 2009.
- Gabric, A., Qu, B., Matrai, P., and Hirst, A.: The simulated response of dimethylsulfide production in the Arctic Ocean to global warming, *Tellus Ser. B*, 57, 391–403, doi:10.1111/j.1600-0889.2005.00163.x, 2005.
- Gabric, A. J., Simó, R., Cropp, R. A., Hirst, A. C., and Dachs, J.: Modeling estimates of the global emission of dimethylsulfide under enhanced greenhouse conditions, *Global Biogeochem. Cy.*, 18, GB2014, doi:10.1029/2004GB002337, 2004.
- 25 [Gondwe, M., Krol, M., Gieskes, W., Klaassen, W., and de Baar, H.: The contribution of ocean-leaving DMS to the global atmospheric burdens of DMS, MSA, SO₂, and NSS SO₄⁻, *Global Biogeochem. Cy.*, 17, 1056, doi:10.1029/2002GB001937, 2003.](#)
- [Grandey, B. S. and Wang, C.: Enhanced marine sulphur emissions offset global warming and impact rainfall, *Sci. Rep.*, 5, 13055, doi:10.1038/srep13055, 2015.](#)
- 30 Gunson, J. R., Spall, S. A., Anderson, T. R., Jones, A., Totterdell, I. J., and Woodage, M. J.: Climate sensitivity to ocean dimethylsulphide emissions, *Geophys. Res. Lett.*, 33, L07701, doi:10.1029/2005GL024982, 2006.
- Halloran, P. R., Bell, T. G., and Totterdell, I. J.: Can we trust empirical marine DMS parameterisations within projections of future climate?, *Biogeosciences*, 7, 1645–1656, doi:10.5194/bg-7-1645-2010, 2010.
- Hurrell, J. W., Hack, J. J., Shea, D., Caron, J. M., and Rosinski, J.: A new sea surface temperature and sea ice boundary dataset for the Community Atmosphere Model, *J. Climate*, 21, 5145–5153, doi:10.1175/2008JCLI2292.1, 2008.
- 35 Kettle, A. J. and Andreae, M. O.: Flux of dimethylsulfide from the oceans: A comparison of updated data sets and flux models, *J. Geophys. Res.-Atmos.*, 105, 26793–26808, doi:10.1029/2000JD900252, 2000.

- Kettle, A. J., Andreae, M. O., Amouroux, D., Andreae, T., Bates, T., Berresheim, H., Bingemer, H., et al.: A global database of sea surface dimethylsulfide (DMS) measurements and a procedure to predict sea surface DMS as a function of latitude, longitude, and month, *Global Biogeochem. Cy.*, 13, 399–444, doi:10.1029/1999GB900004, 1999.
- 5 Kloster, S., Feichter, J., Maier-Reimer, E., Six, K. D., Stier, P., and Wetzel, P.: DMS cycle in the marine ocean-atmosphere system – a global model study, *Biogeosciences*, 3, 29–51, doi:10.5194/bg-3-29-2006, 2006.
- Kloster, S., Six, K. D., Feichter, J., Maier-Reimer, E., Roeckner, E., Wetzel, P., Stier, P., and Esch, M.: Response of dimethylsulfide (DMS) in the ocean and atmosphere to global warming, *J. Geophys. Res.-Biogeosci.*, 112, G03005, doi:10.1029/2006JG000224, 2007.
- [Koch, D., Schulz, M., Kinne, S., McNaughton, C., Spackman, J. R., Balkanski, Y., Bauer, S., Bernsten, T., Bond, T. C., Boucher, O., Chin, M., Clarke, A., De Luca, N., Dentener, F., Diehl, T., Dubovik, O., Easter, R., Fahey, D. W., Feichter, J., Fillmore, D., Freitag, S., Ghan, S. J., Ginoux, P., Gong, S., Horowitz, L., Iversen, T., Kirkevåg, A., Klimont, Z., Kondo, Y., Krol, M., Liu, X., Miller, R., Montanaro, V., Moteki, N., Myhre, G., Penner, J. E., Perlwitz, J., Pitari, G., Reddy, S., Sahu, L., Sakamoto, H., Schuster, G., Schwarz, J. P., Seland, Ø., Stier, P., Takegawa, N., Takemura, T., Textor, C., van Aardenne, J. A., and Zhao, Y.: Evaluation of black carbon estimations in global aerosol models, *Atmos. Chem. Phys.*, 9, 9001–9026, doi:10.5194/acp-9-9001-2009, 2009.](#)
- 10 [Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical \(1850–2000\) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, *Atmos. Chem. Phys.*, 10, 7017–7039, doi:10.5194/acp-10-7017-2010, 2010.](#)
- 15 [Lana, A., Bell, T. G., Simó, R., Vallina, S. M., Ballabrera-Poy, J., Kettle, A. J., Dachs, J., Bopp, L., Saltzman, E. S., Stefels, J., Johnson, J. E., and Liss, P. S.: An updated climatology of surface dimethylsulfide concentrations and emission fluxes in the global ocean, *Global Biogeochem. Cy.*, 25, GB1004, doi:10.1029/2010GB003850, 2011.](#)
- Liss, P. S. and Merlivat, L.: Air-Sea Gas Exchange Rates: Introduction and Synthesis, in: *The role of air-sea exchange in geochemical cycling*, edited by: Buat-Ménard, P., vol. 185 of NATO ASI Series, 113–127, Springer Netherlands, doi:10.1007/978-94-009-4738-2_5, 1986.
- Lohmann, U., von Salzen, K., McFarlane, N., Leighton, H. G., and Feichter, J.: Tropospheric sulfur cycle in the Canadian general circulation model, *J. Geophys. Res.-Atmos.*, 104, 26833–26858, doi:10.1029/1999JD900343, 1999.
- 25 [Ma, X., von Salzen, K., and Cole, J.: Constraints on interactions between aerosols and clouds on a global scale from a combination of MODIS-CERES satellite data and climate simulations, *Atmos. Chem. Phys.*, 10, 9851–9861, doi:10.5194/acp-10-9851-2010, 2010.](#)
- [Mahajan, A. S., Fadnavis, S., Thomas, M. A., Pozzoli, L., Gupta, S., Royer, S.-J., Saiz-Lopez, A., and Simó, R.: Quantifying the impacts of an updated global dimethyl sulfide climatology on cloud microphysics and aerosol radiative forcing, *J. Geophys. Res.-Atmos.*, 120, 2524–2536, doi:10.1002/2014JD022687, 2015.](#)
- 30 [Malin, G., Turner, S. M., and Liss, P. S.: Sulfur: The plankton/climate connection, *J. Phycology*, 28, 590–597, doi:10.1111/j.0022-3646.1992.00590.x, 1992.](#)
- [McGillis, W. R., Dacey, J. W. H., Frew, N. M., Bock, E., and Nelson, R. K.: Water-air flux of dimethylsulfide, *J. Geophys. Res.-Oceans*, 105, 1187–1193, doi:10.1029/1999JC900243, 2000.](#)
- Miles, C. J., Bell, T. G., and Lenton, T. M.: Testing the relationship between the solar radiation dose and surface DMS concentrations using in situ data, *Biogeosciences*, 6, 1927–1934, doi:10.5194/bg-6-1927-2009, 2009.
- 35 [Moss, R., Edmonds, J., Hibbard, K., Manning, M., Ross, S., van Vuuren, D., Carter, T., Emori, S., Kainuma, M., Kram, T., Meehl, G., Mitchell, J., Nakicenovic, N., Riahi, K., Smith, S., Stouffer, R., Thomson, A., Weyant, J., and Wilbanks, T.: The next generation of scenarios for climate change research and assessment, *Nature*, 463, 747–756, 2010.](#)

- Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestedt, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T., and Zhang, H.: Anthropogenic and natural radiative forcing, in: *Climate Change 2013: The physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- 5
Nightingale, P. D., Malin, G., Law, C. S., Watson, A. J., Liss, P. S., Liddicoat, M. I., Boutin, J., and Upstill-Goddard, R. C.: In situ evaluation of air-sea gas exchange parameterizations using novel conservative and volatile tracers, *Global Biogeochem. Cy.*, 14, 373–387, doi:10.1029/1999GB900091, 2000.
- [Quaas, J., Ming, Y., Menon, S., Takemura, T., Wang, M., Penner, J. E., Gettelman, A., Lohmann, U., Bellouin, N., Boucher, O., Sayer, A. M., Thomas, G. E., McComiskey, A., Feingold, G., Hoose, C., Kristjánsson, J. E., Liu, X., Balkanski, Y., Donner, L. J., Ginoux, P. A., Stier, P., Grandey, B., Feichter, J., Sednev, I., Bauer, S. E., Koch, D., Grainger, R. G., Kirkevåg, A., Iversen, T., Seland, Ø., Easter, R., Ghan, S., Rasch, P. J., Morrison, H., Lamarque, J.-F., Iacono, M. J., Kinne, S., and Schulz, M.: Aerosol indirect effects - general circulation model intercomparison and evaluation with satellite data, *Atmos. Chem. Phys.*, 9, 8697–8717, doi:10.5194/acp-9-8697-2009, 2009.](#)
- 10
Quinn, P. K. and Bates, T. S.: The case against climate regulation via oceanic phytoplankton sulphur emissions, *Nature*, 480, 51–56, doi:10.1038/nature10580, 2011.
- 15
[Schulz, M., Textor, C., Kinne, S., Balkanski, Y., Bauer, S., Berntsen, T., Berglen, T., Boucher, O., Dentener, F., Guibert, S., Isaksen, I. S. A., Iversen, T., Koch, D., Kirkevåg, A., Liu, X., Montanaro, V., Myhre, G., Penner, J. E., Pitari, G., Reddy, S., Seland, Ø., Stier, P., and Takemura, T.: Radiative forcing by aerosols as derived from the AeroCom present-day and pre-industrial simulations, *Atmos. Chem. Phys.*, 6, 5225–5246, doi:10.5194/acp-6-5225-2006, 2006.](#)
- 20
Simó, R. and Dachs, J.: Global ocean emission of dimethylsulfide predicted from biogeophysical data, *Global Biogeochem. Cy.*, 16, 1078, doi:10.1029/2001GB001829, 2002.
- Six, K. D. and Maier-Reimer, E.: What controls the oceanic dimethylsulfide (DMS) cycle? A modeling approach, *Global Biogeochem. Cy.*, 20, GB4011, doi:10.1029/2005GB002674, 2006.
- Six, K. D., Kloster, S., Ilyina, T., Archer, S. D., Zhang, K., and Maier-Reimer, E.: Global warming amplified by reduced sulphur fluxes as a result of ocean acidification, *Nat. Climate Change*, 3, 975–978, doi:10.1038/nclimate1981, 2013.
- 25
Slingo, J. M.: The development and verification of a cloud prediction scheme for the ECMWF model, *Q. J. Roy. Meteorol. Soc.*, 113, 899–927, doi:10.1002/qj.49711347710, 1987.
- Spiro, P. A., Jacob, D. J., and Logan, J. A.: Global inventory of sulfur emissions with $1^\circ \times 1^\circ$ resolution, *J. Geophys. Res.-Atmos.*, 97, 6023–6036, doi:10.1029/91JD03139, 1992.
- 30
Stefels, J., Steinke, M., Turner, S., Malin, G., and Belviso, S.: Environmental constraints on the production and removal of the climatically active gas dimethylsulphide (DMS) and implications for ecosystem modelling, *Biogeochemistry*, 83, 245–275, doi:10.1007/s10533-007-9091-5, 2007.
- Stevens, B. and Feingold, G.: Untangling aerosol effects on clouds and precipitation in a buffered system, *Nature*, 461, 607–613, doi:10.1038/nature08281, 2009.
- 35
Tesdal, J.-E., Christian, J. R., Monahan, A. H., and von Salzen, K.: Evaluation of diverse approaches for estimating [sea-surface](#) DMS concentration and [flux-air-sea exchange](#) at global scale, *Environ. Chem.*, ~~in press, 2015-~~ [13, 390–412](#), doi:10.1071/EN14255, [2016](#).

[Thomas, M. A., Suntharalingam, P., Pozzoli, L., Rast, S., Devasthale, A., Kloster, S., Feichter, J., and Lenton, T. M.: Quantification of DMS aerosol-cloud-climate interactions using the ECHAM5-HAMMOZ model in a current climate scenario, *Atmos. Chem. Phys.*, 10, 7425–7438, doi:10.5194/acp-10-7425-2010, 2010.](#)

- Thomas, M. A., Suntharalingam, P., Pozzoli, L., Devasthale, A., Kloster, S., Rast, S., Feichter, J., and Lenton, T. M.: Rate of non-linearity in
5 DMS aerosol-cloud-climate interactions, *Atmos. Chem. Phys.*, 11, 11175–11183, doi:10.5194/acp-11-11175-2011, 2011.
- Tortell, P., Guéguen, C., Long, M., Payne, C., Lee, P., and DiTullio, G.: Spatial variability and temporal dynamics of surface water pCO₂,
ΔO₂/Ar and dimethylsulfide in the Ross Sea, Antarctica, *Deep-Sea Res. Pt I*, 58, 241–259, doi:10.1016/j.dsr.2010.12.006, 2011.
- Tortell, P. D.: Small-scale heterogeneity of dissolved gas concentrations in marine continental shelf waters, *Geochem. Geophys. Geosyst.*, 6,
Q11M04, doi:10.1029/2005GC000953, 2005.
- 10 Vallina, S. M. and Simó, R.: Strong relationship between DMS and the solar radiation dose over the global surface ocean, *Science*, 315,
506–508, doi:10.1126/science.1133680, 2007.
- Vallina, S. M., Simó, R., and Manizza, M.: Weak response of oceanic dimethylsulfide to upper mixing shoaling induced by global warming,
Proc. Natl. Aca. Sci. USA, 104, 16004–16009, doi:10.1073/pnas.0700843104, 2007.
- Vogt, M., Vallina, S. M., Buitenhuis, E. T., Bopp, L., and Le Quééré, C.: Simulating dimethylsulphide seasonality with the dynamic green
15 ocean model PlankTOM5, *J. Geophys. Res.-Oceans*, 115, C06021, doi:10.1029/2009JC005529, 2010.
- von Salzen, K., Leighton, H. G., Ariya, P. A., Barrie, L. A., Gong, S. L., Blanchet, J.-P., Spacek, L., Lohmann, U., and Kleinman, L. I.:
Sensitivity of sulphate aerosol size distributions and CCN concentrations over North America to SO_x emissions and H₂O₂ concentrations,
J. Geophys. Res.-Atmos., 105, 9741–9766, doi:10.1029/2000JD900027, 2000.
- von Salzen, K., Scinocca, J. F., McFarlane, N. A., Li, J., Cole, J. N. S., Plummer, D., Verseghy, D., Reader, M. C., Ma, X., Lazare, M.,
20 and Solheim, L.: The Canadian Fourth Generation Atmospheric Global Climate Model (CanAM4). Part I: Representation of physical
processes, *Atmosphere-Ocean*, 51, 104–125, doi:10.1080/07055900.2012.755610, 2013.
- Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean, *J. Geophys. Res.-Oceans*, 97, 7373–7382,
doi:10.1029/92JC00188, 1992.
- Woodhouse, M. T., Carslaw, K. S., Mann, G. W., Vallina, S. M., Vogt, M., Halloran, P. R., and Boucher, O.: Low sensitivity of cloud
25 condensation nuclei to changes in the sea-air flux of dimethyl-sulphide, *Atmos. Chem. Phys.*, 10, 7545–7559, doi:10.5194/acp-10-7545-
2010, 2010.
- Woodhouse, M. T., Mann, G. W., Carslaw, K. S., and Boucher, O.: Sensitivity of cloud condensation nuclei to regional changes in dimethyl-
sulphide emissions, *Atmos. Chem. Phys.*, 13, 2723–2733, doi:10.5194/acp-13-2723-2013, 2013.

Table 1. List of model sensitivity experiments.

Name	Description
L10 & N00 & γ_a	Control experiment
L10 & N00	No air resistance
L10 & LM86	LM86 flux scheme, no air resistance
K99 & LM86	Older K99 climatology instead of L10 climatology, LM86 flux scheme, no air resistance
K99 & N00 & γ_a	As control, but with K99 climatology
K00 & N00 & γ_a	As control, but with K00 climatology
K99* & N00 & γ_a	As control, but with K99 scaled to L10 global flux
AN01* & N00 & γ_a	As control, but with AN01 scaled to L10 global flux
Temporally invariant	L10 annual mean field for all months scaled to the original L10 global flux
Spatially uniform	Spatially uniform fields with monthly global mean L10 concentration scaled to the original L10 global flux

Table 2. Ocean emissions of DMS from CanAM4.1 and offline calculations with reanalysis fields. DMS flux is derived for the time period of the model simulations (January 2004 to December 2008). Quantities in parentheses are percentage changes relative to the reference run (L10 & N00 & γ_a).

Model runs	CanAM4.1		ERA-Interim	
	Mean flux $\mu\text{mol m}^{-2} \text{d}^{-1}$	Total flux TgS y^{-1}	Mean flux $\mu\text{mol m}^{-2} \text{d}^{-1}$	Total flux TgS y^{-1}
L10 & N00 & γ_a	7.02	28.9	5.72	23.6
L10 & N00	7.60	31.3 (+8 %)	6.13	25.3 (+7 %)
L10 & LM86	4.94	20.4 (-29 %)	4.34	17.9 (-24 %)
K99 & LM86	4.44	18.3 (-37 %)	3.89	16.0 (-32 %)
K99 & N00 & γ_a	6.31	25.9 (-10 %)	5.11	21.0 (-11 %)
K00 & N00 & γ_a	6.02	24.7 (-14 %)	4.90	20.3 (-15 %)

Table 3. Annual DMS emissions, oxidation rates and atmospheric burdens of DMS, SO₂ and SO₄²⁻.

Model runs	Fluxes and oxidation rates (TgS y ⁻¹)			Atmospheric burdens (TgS)		
	Emissions ^a	Oxidation by OH	Oxidation by NO ₃	DMS	SO ₂	SO ₄ ²⁻
L10 & N00 & γ_a	29.8	18.1	11.7	0.24	0.40	0.67
L10 & N00	32.2	19.6	12.6	0.26	0.42	0.69
L10 & LM86	21.2	12.9	8.31	0.17	0.35	0.59
K99 & LM86	19.1	11.2	7.98	0.16	0.34	0.55
K99 & N00 & γ_a	26.8	15.6	11.2	0.22	0.38	0.62
K00 & N00 & γ_a	25.6	15.8	9.86	0.22	0.38	0.61
K99* & N00 & γ_a	29.9	17.4	12.5	0.25	0.40	0.65
AN01* & N00 & γ_a	29.9	16.6	13.3	0.25	0.40	0.64
Temporally invariant	29.7	16.2	13.6	0.24	0.41	0.65
Spatially uniform	29.9	16.3	13.6	0.25	0.41	0.64

^a Includes terrestrial emissions**Table 4.** Diagnostic and prognostic models used to estimate DMS concentrations in Fig. 4-6. These models are discussed in detail in Tesdal et al. (2016).

Name	Reference
AN01	Anderson et al. (2001)
AT04	Aranami and Tsunogai (2004)
AU02	Aumont et al. (2002)
BE04	Belviso et al. (2004)
HadOCC	Collins et al. (2011)
HAMOCC	Kloster et al. (2006), Six and Maier-Reimer (2006)
MI09	Miles et al. (2009)
PlankTOM	Vogt et al. (2010)
PISCES	Belviso et al. (2012)
POP-TGM	Elliott (2009)
SD02	Simó and Dachs (2002)
VS07	Vallina and Simó (2007)




Figure 1. Schematic representation of the sulfur cycle and radiative effects of sulfate aerosols in CanAM4.1. In each grid cell, the model accounts for sources and sinks of sulfate aerosol (SO_4^{2-}), SO_2 , and DMS. SO_2 is emitted from volcanos, fires, and anthropogenic sources. DMS is mainly emitted from the oceans, but there are also some terrestrial sources. DMS is oxidized to SO_2 by OH during the day and by NO_3 during the night. SO_2 is oxidized to sulfate both within clouds and under clear-sky conditions. In-cloud oxidation of sulfur and wet deposition is treated separately for layer (stratiform) and convective clouds. For both types of clouds, oxidation occurs via ozone (O_3) and hydrogen peroxide (H_2O_2). Oxidation rates depend on the pH of the cloud water, which depends on the concentrations of nitric acid (HNO_3), ammonia (NH_3), and carbon dioxide (CO_2).




Figure 2 . Map showing the locations of the nss-SO_4^{2-} measurements used in the model assessment. All cruise data were obtained from the NOAA PMEL Atmospheric Chemistry Data Server (<http://saga.pmel.noaa.gov/data>) collected by ACE-1 (October-December 1995), ACE-2 (June-July 1997), ACE-ASIA (March-April 2001), Aerosols99-INDOEX (January-March 1999), ICEALOT (March-April 2008), NEAQ5 (July-August 2002), NEAQ5 (July- August 2004), TexAQ5-GoMACCS (July-September 2006), VOCALS (October-December 2008) and WACS (August 2012).




Figure 3. Correlation between observed and modelled nss-SO_4^{2-} using the control run (L10 & N00 & γ_a). Modelled nss-SO_4^{2-} are derived from mean results of the simulation during the time period from 2004 to 2008. Observed concentrations are from 1995-2012 and matched to the nearest grid point and month of the model results. Comparison is done for (a) different latitude bands and (b) different fraction of sulfate that is produced by DMS emissions.




Figure 2.4. Scatterplots of atmospheric burdens of sulfur species vs. other species and ocean DMS emissions. Each dot represents the global- and annual-mean of individual ensemble members from the model experiments summarized in Table 1. Crosses indicate ensemble means for simulations with the original, unscaled flux fields. The regression lines were computed from the individual ensemble members corresponding to these unmodified DMS flux fields. Open circles denote ensemble mean of simulations with seasonality (red) or spatial pattern (yellow) removed. Open diamonds denote ensemble averages of simulations with DMS fields different from L10 but scaled to give the same global mean flux. The first column shows atmospheric burdens of sulfur species (SO_4^{2-} , SO_2 , DMS) against ocean emission of DMS, the second column shows atmospheric burdens of SO_4^{2-} and SO_2 against DMS burden, and the third column shows atmospheric burden of SO_4^{2-} against the SO_2 burden.




Figure 35. Radiative forcing difference (change in global annual mean net radiation at the top of the atmosphere, τ -TOA) between model experiments and control experiment relative to the global annual mean flux of ocean DMS. Crosses represent the ensemble means of simulations with unmodified DMS fields. Open circles denote ensemble mean of simulations with seasonality (red) or spatial pattern (yellow) removed. Open diamonds denote simulations with DMS fields different from L10 but scaled to give the same global mean flux. Individual ensemble members for each experiment are shown as dots of the same colour. Only data from individual runs with unmodified K99 (purple) or L10 (blue) DMS emissions are used for the corresponding regression lines.




Figure 46. Estimated difference in global annual mean net radiation at the top of the atmosphere (TOA) for the different climatologies considered in Tesdal et al. (2016) and the control simulation plotted against the global ocean efflux of DMS. DMS fluxes were computed offline using fields from the ERA-Interim reanalysis with N00 & γ_a as the air-sea transfer scheme (large filled circles). A linear regression for these runs only (grey dashed line) is used to derive estimates for other experiments (small red dots on regression line). The different climatologies considered are listed in Table 4.




Figure 57. Deviation in global annual mean net radiation at TOA from control plotted against the global annual mean atmospheric burden of SO_4^{2-} . Symbols are as in Fig. 3-5. All data points are used for the linear regression (grey dashed line).




Figure 6.8. Deviation in global means of cloud forcing (upper panels), clear-sky reflected (middle panels), and total reflected irradiance (lower panels) at TOA from control plotted against global annual mean ocean DMS flux (left) and global annual mean atmospheric burden of SO_4^{2-} (right). Symbols are as in Fig. 3-5.




Figure 7.9. Absolute differences in global mean flux, oxidation rates, sulfur burdens, and radiation between the control run and model runs with seasonally invariant (red) or spatially uniform (yellow) DMS concentration, and the L10 & N00 model experiment (blue). Fluxes and oxidation rate of DMS are shown in the upper panels. The global mean DMS flux includes terrestrial sources to ensure mass balance. The only sink for DMS is oxidation to SO₂, which is shown for both oxidation pathways (oxidation by OH and NO₃ radicals). Absolute changes in the atmospheric sulfur burdens of DMS, SO₂, and SO₄²⁻ are shown in the middle panels. Bottom panels show absolute changes in cloud forcing, clear-sky reflected and total reflected shortwave flux. Total reflected flux is the sum of cloud and clear-sky reflected flux. To derive the error estimates, all treatments (control, temporally invariant, spatially uniform, and no air-side resistance) were pooled after their separate means were removed; error bars are ± two standard deviations of the pooled data ($n = 12$). [Statistical significance is determined by comparing the mean differences among the model runs with the corresponding error bars.](#)