# 1 Evaluation of observed and modelled aerosol lifetimes using radioactive

2 tracers of opportunity and an ensemble of 19 global models

- 3
- N. I. Kristiansen<sup>1</sup>, A. Stohl<sup>1</sup>, D. J. L. Olivié<sup>2</sup>, B. Croft<sup>3</sup>, O. A. Søvde<sup>4</sup>, H. Klein<sup>2</sup>, T.
- 5 Christoudias<sup>5</sup>, D. Kunkel<sup>6</sup>, S. J. Leadbetter<sup>7</sup>, Y.H. Lee<sup>8</sup>, K. Zhang<sup>9</sup>, K. Tsigaridis<sup>10</sup>,
- 6 T. Bergman<sup>11</sup>, N. Evangeliou<sup>1,12</sup>, H. Wang<sup>9</sup>, P.-L. Ma<sup>9</sup>, R. C. Easter<sup>9</sup>, P. J. Rasch<sup>9</sup>,
- 7 X. Liu<sup>13</sup>, G. Pitari<sup>14</sup>, G. Di Genova<sup>14</sup>, S. Y. Zhao<sup>15</sup>, Y. Balkanski<sup>12</sup>, S. E. Bauer<sup>10</sup>, G.
- 8 S. Faluvegi<sup>10</sup>, H. Kokkola<sup>11</sup>, R. V. Martin<sup>3</sup>, J. R. Pierce<sup>16,3</sup>, M. Schulz<sup>2</sup>, D. Shindell<sup>8</sup>,
- 9 H. Tost<sup>6</sup> and H. Zhang<sup>15</sup>
- 10
- 11 [1] {NILU-Norwegian Institute for Air Research, Kjeller, Norway}
- 12 [2] {Norwegian Meteorological Institute, Oslo, Norway}
- 13 [3] {Department of Physics and Atmospheric Science, Dalhousie University, Halifax, Canada}
- 14 [4] {Center for International Climate and Environmental Research Oslo (CICERO), Oslo, Norway}
- 15 [5] {The Cyprus Institute, Nicosia, Cyprus}
- 16 [6] {Institute for Atmospheric Physics, Johannes Gutenberg-University Mainz, Germany}
- 17 [7] {Met Office, Exeter, UK}
- 18 [8] {Earth and Ocean Sciences, Nicholas School of the Environment, Duke University, Durham, NC, USA}
- 19 [9] {Pacific Northwest National Laboratory (PNNL), Richland, WA, USA}
- 20 [10] {Center for Climate Systems Research, Columbia University, and NASA Goddard Institute for Space
- 21 Studies, New York, NY, USA}
- 22 [11] {Finnish Meteorological Institute, Kuopio, Finland}
- 23 [12] {Laboratoire des Sciences du Climat et de l'Environnement, CEA-CNRS-UVSQ, Gif-sur-Yvette,
- 24 France }
- 25 [13] {Department of Atmospheric Science, University of Wyoming, Laramie, WY, USA}
- 26 [14] {University of L'Aquila, L'Aquila, Italy}
- 27 [15] {Laboratory for climate studies, National Climate Center, Chinese Meteorological Administration,
- 28 Beijing, China}
- 29 [16] {Department Atmospheric Science, Colorado State University, Fort Collins, CO, USA}
- 30
- 31 Correspondence to: N. I. Kristiansen (nik@nilu.no)

## 1 Abstract

Aerosols have important impacts on air quality and climate, but the processes affecting their removal from 2 the atmosphere are not fully understood and are poorly constrained by observations. This makes modelled 3 aerosol lifetimes uncertain. In this study, we make use of an observational constraint on aerosol lifetimes 4 provided by radionuclide measurements and investigate the causes of differences within a set of global 5 6 models. During the Fukushima Dai-Ichi nuclear power plant accident of March 2011, the radioactive isotopes cesium-137 (<sup>137</sup>Cs) and xenon-133 (<sup>133</sup>Xe) were released in large quantities. Cesium attached to 7 particles in the ambient air, approximately according to their available aerosol surface area. <sup>137</sup>Cs size 8 distribution measurements taken close to the power plant suggested that accumulation-mode (AM) sulphate 9 aerosols were the main carriers for the cesium. Hence, <sup>137</sup>Cs can be used as a proxy tracer for the AM 10 sulphate aerosol's fate in the atmosphere. In contrast, the noble gas <sup>133</sup>Xe behaves almost like a passive 11 transport tracer. Global surface measurements of the two radioactive isotopes taken over several months 12 after the release allow the derivation of a lifetime of the carrier aerosol. We compare this to the lifetimes 13 simulated by 19 different atmospheric transport models initialized with identical emissions of <sup>137</sup>Cs that 14 were assigned to an aerosol tracer with each model's default properties of AM sulphate, and <sup>133</sup>Xe emissions 15 that were assigned to a passive tracer. We investigate to what extent the modelled sulphate tracer can 16 reproduce the measurements, especially with respect to the observed loss of aerosol mass with time. 17 Modelled <sup>37</sup>Cs and <sup>133</sup>Xe concentrations sampled at the same location and times as station measurements 18 allow a direct comparison between measured and modelled aerosol lifetime. The e-folding lifetime  $\tau_{e}$ , 19 calculated from station measurement data taken between two and nine weeks after the start of the emissions, 20 is 14.3 days (95% confidence interval 13.1-15.7 days). The equivalent modelled  $\tau_e$  lifetimes have a large 21 spread, varying between 4.8 and 26.7 days with a model median of 9.4±2.3 days, indicating too fast removal 22 in most models. Because sufficient measurement data were only available from about two weeks after the 23 release, the estimated lifetimes apply to aerosols that have undergone long-range transport, i.e. not for 24 freshly emitted aerosol. However, modelled instantaneous lifetimes show that the initial removal in the first 25 two weeks was quicker (lifetimes between 1-5 days) due to the emissions occurring at low altitudes and co-26 location of the fresh plume with strong precipitation. Deviations between measured and modelled aerosol 27 lifetimes are largest for the northernmost stations and at later time periods, suggesting that models do not 28 transport enough of the aerosol towards the Arctic. The models underestimate passive tracer (<sup>133</sup>Xe) 29 concentrations in the Arctic as well but to a smaller extent than for the aerosol (<sup>137</sup>Cs) tracer. This indicates 30 that in addition to too fast aerosol removal in the models, errors in simulated atmospheric transport towards 31 the Arctic in most models also contribute to the underestimation of the Arctic aerosol concentrations. 32

#### 1 **1 Introduction**

Aerosols play an important role for air quality and influence the global climate (Friedlander, 1977; Seinfeld 2 and Pandis, 1998; Ramanathan et al., 2001) but the processes affecting their removal from the atmosphere 3 are not fully understood and are poorly constrained by observations. Generally, aerosol concentrations are 4 affected by emissions, transport, removal, and physico-chemical transformation (e.g., Pöschl, 2005), and the 5 atmospheric lifetime of aerosols is a function of the various removal processes, such as dry deposition by 6 impaction and sedimentation, as well as wet deposition. For accumulation-mode (AM) aerosols (0.1-2 µm 7 diameter), the dominant removal process is wet deposition. The uncertainties and lack of observational 8 constraints on these removal processes affect the ability to model aerosol concentrations correctly and make 9 modelled aerosol lifetimes uncertain. The uncertainty in the effects of aerosols on climate further affects the 10 ability to diagnose how sensitive the climate is to greenhouse gas emissions (e.g. Andreae, 2007). 11

Observation-based estimates of aerosol lifetimes are sparse due to the difficulty of obtaining measurements 12 that cover a sufficient geographical area and time period for robust analysis. Reported observation-based 13 aerosol lifetimes range from a few days to about a month in the troposphere (Williams et al., 2002; Paris et 14 al., 2009; Schmale et al., 2011). Other aerosol lifetime estimates are derived from radionuclides produced by 15 cosmic rays, radon decay or nuclear bomb tests, and vary from 4 days to more than a month (Giorgi and 16 Chameides, 1986), reflecting the different origin (e.g., surface or stratospheric) of radionuclide tracers. 17 Aerosol residence times of about 4 days in the lower troposphere and about 12 days in the middle to upper 18 troposphere may be seen as typical (Moore et al., 1973) but higher values of 8 days for the lower 19 troposphere have been reported as well (Papastefanou, 2006). Following the Chernobyl nuclear accident, the 20 exponential decline of the radionuclide concentrations indicated a residence time of 7 days (Cambray et al., 21 1987). Models report global average residence times of AM aerosol in the atmosphere on the order of 3–7 22 days for species emitted near the surface (Chin et al., 1996; Feichter et al., 1996; Stier et al., 2005; Berglen 23 et al., 2004; Liu et al., 2005; Bourgeois and Bey, 2011; Chung and Seinfeld, 2002; Koch and Hansen, 2005; 24 Textor et al., 2006). The differences in reported lifetimes from observations and models can partly be 25 attributed to the applied definition of lifetime (i.e. characteristic time of exponential decay vs ratio of burden 26 to deposition or emissions). These lifetime definitions are only equivalent if the decay has a constant e-27 folding time over the considered time period (Croft et al., 2014). Several definitions of lifetime and 28 residence time (the ratio of burden to deposition/emission/production) exist but the terminologies are often 29 used inconsistently. We encourage future studies to give clear information about which lifetime definitions 30 that are used. 31

Many modelling studies have analysed the global distribution, transport and lifetime of aerosols, particularly within the Aerosol Comparisons between Observations and Models (AeroCom) initiative (e.g., Textor et al., 2006; Koch et al., 2009; Samset et al., 2014). It has been demonstrated that large differences exist for aerosol dispersal and removal between models. Samset et al. (2014) found that, compared to aircraft

measurements models seem to overestimate black carbon (BC) aerosol concentrations in the middle and 1 upper troposphere, and thus, a short aerosol lifetime appears necessary to reproduce such observations. On 2 the other hand, the models generally underestimate the aerosol concentrations closer to the surface, and this 3 would get worse with a shorter model lifetime. In particular, models struggle to capture the high aerosol 4 concentrations in the Artic related to the Arctic Haze season (e.g., Shindell et al., 2008; Koch et al., 2009). A 5 6 general underestimation of surface aerosol concentrations in the Arctic is found during the Arctic haze season, while an overestimation is often found in the summer (e.g., Eckhardt et al., 2015). Models are 7 underestimating poleward transport, removing aerosols too efficiently, or not confining pollution sufficiently 8 to the lowest model levels due to excessive vertical diffusion (Koch et al., 2009), but it is not clear which is 9 the main cause. It has also not been fully quantified how these model issues evolve during transport to the 10 Arctic. Browse et al. (2012) and Wang et al. (2013b) found that scavenging parameterizations play a 11 significant role in modelling Arctic aerosol concentrations. In particular, slow scavenging by ice clouds in 12 winter and enhanced scavenging by drizzle in summer are important for modelling the annual cycle of 13 Arctic aerosol concentrations. Using a source-tagging technique, Wang et al. (2014b) found that the annual 14 mean lifetime of Arctic AM BC aerosol has very strong source-region dependence, varying by a factor of 15 four. Zhang et al. (2015) further showed that the lifetime also depends on season and emission type. 16 Substantially lower BC lifetime is found in summer, likely due to relatively strong wet removal, than in 17 other seasons, and open-fire emissions that have higher initial injection heights which leads to generally 18 longer lifetime than emissions from the surface. 19

In this study, we evaluate modelled aerosol lifetimes slightly differently than in previous studies. We use a 20 unique single event with relatively well-known emissions to determine the lifetime of aerosols in the 21 atmosphere. Previous studies report the mean lifetime of aerosols from simulations with higher uncertainty 22 in the emission terms. Specifically, emissions and lifetimes are sometimes tuned to obtain what is thought to 23 be "reasonable" concentrations. In this exercise, we use emissions of radionuclides from the Fukushima Dai-24 Ichi nuclear power plant (FD-NPP) accident in March 2011 as "tracers of opportunity". The cesium (<sup>137</sup>Cs) 25 released during the accident attached to the particles in the ambient air, approximately in proportion to their 26 surface area (Papastefanou, 2008). The peak of the aerosol surface area distribution is generally in the AM, 27 which in the area of FD-NPP is typically dominated by sulphate. Kaneyasu et al. (2012) performed 28 measurements close to FD-NPP and confirmed that <sup>137</sup>Cs was attached to or included in aerosols (internally 29 mixed with other aerosol components), and their <sup>137</sup>Cs size distribution measurements showed that AM 30 sulphate aerosols were the main transport carriers for the cesium. They further explained that elemental 31 carbon (EC) or BC particles were unlikely to be the transport carriers because flaming fires did not continue 32 during the FD-NPP accident, large-scale forest fires were not reported around FD-NPP, and because the 33 local residents had refrained from the burning of firewood in fear of the re-emission of radionuclides since 34 the accident. However, they could not exclude the possibility that water-insoluble organic carbon (OC) 35 could have acted as a transport medium of cesium. Even though it is possible that some of the <sup>137</sup>Cs attached 36

- to other aerosol than sulphate, these aerosol components were likely mixed internally with the dominant AM
  sulphate aerosol and therefore should have similar removal properties. Miyamoto et al. (2014) reported <sup>137</sup>Cs
  size distribution measurements taken in an earlier phase (6 days after the accident) and closer to FD-NPP,
  which showed activity median aerodynamic diameter (AMAD) of <sup>137</sup>Cs around 1.5-1.6 µm, in agreement
  with the results of Kaneyasu et al. (2012). Masson et al. (2013) found that after long-range transport to
  Europe, AMAD of <sup>137</sup>Cs ranged between 0.25 and 0.71 µm, thus again in the AM of the ambient aerosols.
- 7 Hence, <sup>137</sup>Cs can be used as a proxy tracer for the AM sulphate aerosol's fate in the atmosphere.
- Cesium from FD-NPP was measured in the Northern Hemisphere for more than three months after its 8 release and traced the fate of its carrier-aerosol in the atmosphere. These measurements provided a unique 9 opportunity to estimate the lifetime of AM aerosols in the atmosphere, as presented by Kristiansen et al. 10 (2012). In that study, measurements of the two radionuclides xenon ( $^{133}$ Xe) and cesium ( $^{137}$ Cs) were used, 11 both released in large quantities from FD-NPP. The noble gas xenon  $(^{133}Xe)$  was used as a passive transport 12 tracer. Notice that both radionuclides have very low background concentrations caused by emissions from 13 nuclear facilities (Wotawa et al., 2010) or, in the case of <sup>137</sup>Cs, resuspension of deposited radiocesium. 14 Background values were subtracted from all measured values, as described by Kristiansen et al. (2012). 15 These authors used the measured ratios of the aerosol  $(^{137}Cs)$  to the passive tracer  $(^{133}Xe)$  enhancements in 16 the surface concentrations, to compensate for variability in transport, and estimated an AM aerosol lifetime 17 of 10-14 days. This is longer than the mean lifetimes of AM aerosols obtained from most aerosol models 18 (typically in the range of 3–7 days). The disagreement could be partly due to the fact that the emissions were 19 from a single location and during a specific season, the measurements were all ground-based, and thus the 20 data were not fully representative of the global and annual mean aerosol lifetime, as well as using definitions 21 of lifetime that were not equivalent under the considered conditions (Croft et al., 2014). In the current study, 22 we try to resolve this issue and investigate to what extent aerosol models can reproduce the observations, 23 especially with respect to the observed loss of aerosol mass with time. 24
- We use the term "aerosol lifetime" throughout the paper to indicate the lifetime of AM aerosol and primarily sulphate. We assume that the <sup>137</sup>Cs attached mostly to the dominant AM sulphate aerosol, confirmed by measurements. The lifetimes apply to aerosols that have undergone long-range transport (after about 2-3 weeks), i.e. the results presented cannot directly constrain the lifetime of freshly emitted aerosols.
- The paper is organized as follows. In Sect. 2 we describe the measurements used in the study, followed by an overview of all participating models in Sect. 3. The methods are described in Sect. 4, and the main results presented in Sect. 5. In Sect. 6 we further discuss some important aspects of the results and compare our results to other recent studies. Main conclusions are summarized in Sect. 7. The paper also includes supplementary information in three appendices A to C.
- 34

## 1 2 Observations

We have used atmospheric surface measurements of activity concentrations of the noble gas 133Xe and the 2 aerosol-bound radionuclide <sup>137</sup>Cs available from several stations (Fig. 1) operated by the Comprehensive 3 Nuclear-Test-Ban Treaty Organization (CTBTO). For collecting particulate radionuclides, about 20.000 m<sup>3</sup> 4 of air is blown through a filter over a period of 24 h. The different radionuclides are measured with high-5 6 resolution germanium detectors (Schulze et al., 2000; Medici, 2001). The minimum detectable activity concentration of <sup>137</sup>Cs is 1 µBq m<sup>-3</sup>. During the International Noble Gas Experiment (INGE), noble gas 7 measurement systems have been set up worldwide (Wernsberger and Schlosser, 2004; Saey and de Geer, 8 2005) at CTBTO stations. The collection period of the xenon samples is typically 12 h. The most prevalent 9 and important isotope is <sup>133</sup>Xe, which is measured with an accuracy of about 0.1 mBg m<sup>-3</sup> All measured 10 radionuclide concentrations were corrected for radioactive decay relative to the time of the earthquake on 11 11 March 05:46 UTC that triggered the nuclear accident. The measurements were further converted from 12 activity per norm cubic meter at standard temperature and pressure (273.15 K and 101325 Pa) to activity per 13 cubic meter (using meteorological analysis data) for comparison with the model results. The measurements 14 and their uncertainties are described in more detail by Stohl et al. (2012) and Kristiansen et al. (2012). 15

16

#### 17 3 Model simulations

A total of 19 atmospheric transport models have simulated the transport and removal of the radioactive
isotopes released during the FD-NPP accident. The models are classified as either Lagrangian Particle
Dispersion Models (LPDM), Aerosol Transport Models (ATM; models which rely entirely on
meteorological input data) and Aerosol Circulation Models (ACM; models which calculate their own
meteorology or are nudged towards (re)analysis data). Table 1 shows an overview of the models included in
the experiment including their type, meteorology and model resolutions. More details on each model's
treatment of aerosols are given in Appendix A, Table A1.

All model simulations were initiated with identical emissions of cesium  $(^{137}Cs)$  and xenon  $(^{133}Xe)$  as 25 determined from inverse modelling by Stohl et al. (2012). The simulations extended from 11 March until at 26 least 5 June 2011 for when the last measurement of the radionuclides were taken. A total of 36.6 PBg of 27 <sup>137</sup>Cs and 15.3 EBg of <sup>133</sup>Xe were released by all models. The emission rates vary significantly over the 28 emission period considered (11 March – 20 April 2011) but the major emissions of the radionuclides 29 occurred over about 5 days (11-15 March 2011). The emissions of cesium continued until 20 March after 30 which they dropped significantly. The releases were divided into three vertical layers between 0–1000 m 31 above ground level (Stohl et al., 2012). Croft et al. (2014) have shown that the e-folding lifetimes derived 32 from their GEOS-Chem model simulations of the Fukushima emissions do not depend very much on the 33 exact specification of the emissions (e.g., their altitude, location and time). This is because radionuclides 34 could be measured in the atmosphere for several months, long after the emissions practically had ceased 35

(Kristiansen et al., 2012). Therefore, after the end of the emissions the decrease in measured concentrations
can be attributed solely to aerosol removal. Biases in the emissions affect the absolute model-simulated
values, but not the lifetime estimate. In the analysis, it might be expected that the FLEXPART model will
perform relatively well since the source terms used by all models were estimated using inverse modelling
with the help of FLEXPART. However, the source term was constrained using measurements from several
other stations than those utilized in the current study, and both airborne and deposition data.

The cesium was treated as sulphate aerosols in the model simulations, i.e. undergoing the same wet and dry 7 deposition as AM sulphate aerosols. Xenon was treated as a passive tracer without wet and dry removal 8 processes. In order to evaluate the removal of aerosols due to wet and dry deposition processes only, no 9 radioactive decay of <sup>137</sup>Cs (half-life 30 years) and <sup>133</sup>Xe (half-life 5.25 days) was simulated by the models or 10 the model simulation was decay-corrected to the start of the nuclear accident (11 March 05:46 UTC). Both 11 the emissions used for the model simulations and the atmospheric concentration measurements were also 12 decay-corrected. Modelled cesium and xenon concentrations were sampled at the location of the 11 CTBTO 13 sites (Fig. 1), at times when measurements were available. This allows a direct comparison to the 14 measurements and observation-based lifetime evaluations (Kristiansen et al., 2012). Modelled total 15 atmospheric burdens as a function of time were also calculated and evaluated. 16

The transport of the radioactive cloud across the Northern Hemisphere is illustrated in Fig. 2, as simulated by the FLEXPART model using meteorological analysis data from the Global Forecast System (GFS) model of the National Centers for Environmental Prediction (NCEP), in the weeks following the initial release at FD-NPP. While transport patterns depend on the meteorological data set used (e.g., some models generate their own meteorology), it can be seen that 3-4 weeks after the start of the emissions, the radionuclides were already distributed fairly homogeneously over the entire Northern Hemisphere.

23

## 24 4 Methods

We use the same basic approaches as in Kristiansen et al. (2012) to evaluate measured and modelled loss of 25 aerosol mass with time, i.e. aerosol lifetimes. Measured aerosol lifetimes derived directly from the decay of 26 station measurements of cesium (<sup>137</sup>Cs) and xenon (<sup>133</sup>Xe) (Kristiansen et al., 2012), are compared to 27 modelled aerosol lifetimes determined in exactly the same way from modelled station concentrations for the 28 same time periods as the observations. To reduce the variability caused by atmospheric transport, we 29 normalize the  ${}^{137}$ Cs values by the  ${}^{133}$ Xe values, i.e. the ratio of the aerosol ( ${}^{137}$ Cs) to the passive tracer 30 (<sup>133</sup>Xe) is used throughout all lifetime evaluations. This largely compensates for variability in transport but 31 not completely because the source terms for <sup>137</sup>Cs and <sup>133</sup>Xe are not perfectly correlated. Additionally, we 32 use global burdens estimated from measurement data as in Kristiansen et al. (2012) and compare these to 33 modelled global burdens. 34

Several definitions of atmospheric lifetime exist. In Kristiansen et al. (2012), the e-folding lifetime was 1 used as a measure of aerosol lifetime. Croft et al. (2014) document, compare and explain differences 2 between global mean aerosol lifetime, the definition typically reported for aerosol- and climate model 3 simulations, and e-folding times from their GEOS-Chem transport simulations of the FD-NPP accident 4 emissions. They show that the two definitions are not directly comparable for the FD-NPP case. The lifetime 5 6 results for the former definition were heavily influenced by initial quick removal, giving a much shorter lifetime than the e-folding lifetime for the hemispherically relatively well-mixed phase starting about 3 7 weeks after the main emissions. They also show that the global mean aerosol lifetime strongly depends on 8 the altitude where emissions are assumed to occur, while the e-folding lifetime is relatively insensitive to 9 emissions parameters such as altitude, location and time, suggesting that e-folding times allow a robust 10 comparison between modelled and measurement-based lifetimes. For the purpose of this study we consider 11 lifetime as equivalent to residence time. 12

In this exercise, we will use the e-folding lifetime (as in Kristiansen et al., 2012) and the instantaneous lifetime (as in Croft et al., 2014) as means of evaluation. The two lifetimes are typically used to evaluate exponentially decreasing aerosol concentrations after an emissions pulse.

16 The e-folding lifetime is defined as

17 
$$\tau_e = \frac{-t_i}{ln \frac{\mathcal{C}(t_i)}{\mathcal{C}(t_0)}}$$
(1)

where  $C(t_i)$  is the concentration at time  $t_i$ ,  $C(t_0)$  the initial concentration at time  $t_0$ , and  $t_i$  is the time since  $t_0$ . Instantaneous lifetime is defined as

20 
$$\tau_{inst} = \frac{t_{(i-1)} - t_i}{ln \frac{\mathcal{C}(t_i)}{\mathcal{C}(t_{i-1})}}$$
(2)

21 where  $t_i$  and  $t_{i-1}$  are adjacent time steps, here in this study separated by 1 day.

Further, we calculate the mean and median of all the modelled lifetimes. These are calculated from the lifetimes obtained for each model. Since the distribution of the modelled lifetime data are not symmetric but have outliers, the mean is not the best representative of the centre of the data, and therefore the median is preferred. The variability is given as the standard deviation (STD) from the mean, and the median absolute deviation (MAD) from the median. The MAD is defined as the median of the absolute deviations from the data's median:

28  $MAD = \text{median} (|\tau_i - \text{median}(\tau_i)|, i = 1, ..., N)$ (3)

where *N* is the total number of models and  $\tau_i$  is the lifetime (either e-folding or instantaneous) obtained from model *i*.

#### 1 5 Results

## 2 5.1 Direct comparison of measured and modelled aerosol lifetimes

Here we present a direct comparison of measured and modelled aerosol lifetimes based on an observational 3 constraint provided by radionuclide measurements of the FD-NPP emissions. We investigate the causes of 4 differences between the measurements and models. The measured and modelled aerosol lifetimes are 5 estimated from the ratios of the aerosol ( $^{137}$ Cs) to the passive tracer ( $^{133}$ Xe) surface concentrations (Sect. 4) 6 at 11 CTBTO measurement sites (Sect. 3) as shown in Fig. 3. Both the measurements and all model 7 simulations show a decrease in these ratios with time due to removal of aerosols. The aerosol lifetimes are 8 estimated by fitting an exponential decay model (grey lines in Fig. 3) to the daily median ratios over all 9 stations (black triangles;  $\omega$  in Fig. 3), between 15 and 65 days after the start of the emissions on 11 March. 10 During this time period, measurement data exist from at least five stations each day, i.e. the daily median 11 value  $\omega$  is calculated from at least five values. The data density before day 15 and after day 65 is quite 12 sparse and there are sampling biases. Before day 15, the measurements are mainly from the closest stations 13 as the plume has not yet reached all sites and mixed through the Northern Hemisphere. After day 65, valid 14 measurement data become sparse because concentrations start to drop below the detection limit and all data 15 below detection limit were discarded. The remaining valid data after day 65 are mainly from Yellowknife 16 and Spitsbergen with values near the detection limit. If these periods were considered, the lifetimes would be 17 biased high, possibly due to a latitude effect discussed later. 18

The measured aerosol (<sup>137</sup>Cs) to the passive tracer (<sup>133</sup>Xe) ratios decay with an e-folding lifetime  $\tau_e$  (Eq. 1) 19 of 14.3 days (Fig. 3 and Table 2) with a 95% confidence interval of 13.1-15.7 days (Appendix B and Table 20 B1). The equivalent modelled  $\tau_e$  lifetimes vary between 4.8 and 26.7 days with a model mean (± standard 21 22 deviation) of 10.7 $\pm$ 5.5, and a model median ( $\pm$  MAD, Eq. 3) of 9.4 $\pm$ 2.3 days (Table 2). Thus, the model mean and median lifetimes are shorter than the lifetime based on the measurements, indicating too quick a 23 24 removal of the aerosols in the models. However, some individual models have longer lifetimes than measured, and the model mean  $\pm$  standard deviation encompasses the measured lifetime. Large variations in 25 26 the modelled lifetimes are expected due to differences in the simulated transport and especially tropospheric removal. 27

28 Table 2 shows the lifetime estimates using data only from stations below and above 50°N separately. Both 29 measured and mean modelled lifetimes are shorter below 50°N than at high latitudes, suggesting less efficient aerosol removal at high latitudes during the time period investigated (March-June). However, a few 30 models (NorESM, ULAQ-CCM, EMAC-1, EMAC-2 and NAME) also have somewhat shorter lifetimes at 31 higher latitudes than at lower latitudes, and two models (CAM5 and FLEXPART) show no change in 32 lifetime with latitude. Wang et al. (2013b) found that CAM5, with the shortest lifetimes of all models, 33 overestimates aerosol wet removal by super-cooled liquid in mixed-phase clouds, which was improved in 34 35 CAM5 PNNL. There is a larger spread in model lifetimes and larger deviations from the measurements at

high latitudes compared to at lower latitudes, indicating larger uncertainties in the simulation of aerosols at 1 higher latitudes. Similarly, lifetime estimates from the earlier phases (25-45 days after the start of the 2 release, a period when the major emissions were over) are compared to those for later time periods (days 45-3 65) (Table 2). Both the measurements and the median model give slightly longer lifetimes (by about 0.3-1.0 4 days) for the earlier phases than for later time periods. This is perhaps unexpected as the radionuclides are 5 6 initially located in the lower troposphere where they are strongly affected by precipitation scavenging and dry deposition, and later when the radionuclides are mostly located in the relatively dry upper troposphere 7 (or even stratosphere) the removal processes are much less effective which would yield an increase in 8 lifetime. However, since the data used to derive the lifetimes are ground-based and thus biased towards that 9 part of the aerosol population residing in the lower troposphere, neither the measurements nor the models 10 may fully reflect the increase in lifetime due to transport in the upper troposphere. Differences in lifetime 11 between tracers emitted near the surface and those coming from the stratosphere may be substantial, as 12 discussed in Kristiansen et al. (2012). Furthermore, most of the radionuclides were uplifted immediately 13 after the release (Stohl et al., 2012) and thus quite well mixed in the troposphere already for the earlier time 14 period (days 25-45), as also shown in Fig. 2. In addition, station data from the later time periods (day 45-65) 15 include fewer data points in the calculations than in the earlier phases. Lastly, there are larger variations in 16 the modelled lifetimes for the later phase, and a larger model to measurement deviation, indicating larger 17 uncertainty in the modelled lifetimes for the later phase than for the earlier phase. 18

In summary, both the mean and median modelled aerosol lifetimes are somewhat shorter than the observed aerosol lifetime, although mostly agreeing within the standard deviations of the modelled lifetimes. The underestimation of lifetimes by the models is smaller than what was indicated by Kristiansen et al. (2012) based on reported mean modelled lifetimes from the literature, consistent with the findings of Croft et al. (2014). The deviations between observed and modelled aerosol lifetimes are largest for the northernmost stations and at later time periods indicating higher uncertainties in the simulation of aerosols and aerosol lifetimes at higher latitudes and later times.

26

#### 27 5.2 Model evaluation of latitudinal variations in transport and scavenging

The previous section indicated that disagreements between the measurements and the models are largest at 28 29 higher latitudes and at later time periods. This agrees with several previous studies (e.g., Shindell et al., 2008; Koch et al., 2009), which suggest that models do not capture the Arctic aerosol variations very well. 30 In this study, we have the possibility to evaluate the models' abilities to simulate the aerosol concentrations 31 and transport at different latitudes, by comparing the measured and modelled aerosol (<sup>137</sup>Cs) and the passive 32 transport tracer (<sup>133</sup>Xe) data. In this way, we can directly explore how well the models are able to reproduce 33 the observations at the northernmost stations (i.e. above 60°N; Yellowknife 62.5°N and Spitsbergen 78°N) 34 35 compared to at lower latitudes. We also examine whether there are any temporal variations in the modelobservation deviations. The analyses are limited by the low number of Arctic/Subarctic stations and the 36

1 particular season and time-period (March-June 2011) of the measurements. Some indications may still be

given which might help explain shortcomings with simulated scavenging and/or transport in the models, and
if one is the more likely cause for deviations to observations.

- We use the ratio of the modelled to measured surface concentrations as means of model score, where a 4 ratio > 1 indicates that the model over-predicts, and a ratio < 1 represents under-predictions (Fig. 4 and 5). 5 Figure 6 shows the ratio of modelled to observed aerosol (green circles) and passive transport tracer (blue 6 circles) concentrations as a function of latitude of the measurement stations. Generally, the models 7 reproduce the observed passive tracer ( $^{133}$ Xe) values better than the observed aerosol ( $^{137}$ Cs) values, 8 indicating that transport is reasonably well represented in the models. Scavenging affects only the aerosols, 9 causing more variability and a larger extent of over- or under-prediction. On average, over all stations and 10 models, the median ratio of modelled to measured passive tracer values is 0.79 (Fig. 4) and for the aerosols 11 0.49 (Fig. 5), suggesting a general under-prediction of both aerosol and passive tracer concentrations. The 12 aerosol underestimation is consistent with the lifetime results from the previous section (5.1) which showed 13 that modelled aerosol lifetimes are too short compared to the measurements, and this would yield too much 14 removal and thus under-predictions of the aerosol concentrations. 15
- Furthermore, at the southernmost stations (Wake Island and Oahu) many models over-predict both the 16 aerosol and passive tracer concentrations, whilst under-predicting both at the stations further north (Fig. 4 17 and 5). The exception is Schauinsland (48°N) where the models generally over-predict the passive tracer but 18 under-predict the aerosol values. We find the largest under-prediction at the northernmost stations; 19 Yellowknife for the passive tracer and Spitsbergen for the aerosol values. The modelled to measured aerosol 20 ratios have a pronounced latitudinal trend with ratios decreasing (i.e. larger under-predictions) with 21 increasing latitude (Fig. 5 and 6).. This indicates that there is too much aerosol removal occurring en route to 22 the Arctic. There is also a slight decrease in modelled to measured passive tracer ratios with latitude (Fig. 4 23 and 6), suggesting that transport to the high latitudes is also not strong enough in the models and this 24 contributes to the aerosol underestimates at high latitudes. However, the main reason for the aerosol 25 underestimates at high latitudes must be a too efficient simulated removal. 26
- Finally, Fig. 6 illustrates the temporal variations of modelled to measured ratios. The values are medians
  over 14 day time periods and the size of the circles indicates the time; the larger the circle the later the time
  period. Particularly for the aerosol values, some models tend to show a stronger underestimation with time,
  also evident in the median model.
- Our results suggest that modelling of both scavenging and transport are causes for disagreements with observations. Transport to the high latitudes is not strong enough in the models and this contributes to the model underestimates also of the aerosol tracer at high latitudes. However, the modelled to observed ratios decrease much more strongly with latitude for the aerosol than for the passive tracer and this indicates that problems with the aerosol scavenging is the major reason for disagreement between the simulations and the

measurements. Scavenging in the models depends strongly on the representation of clouds (temporal and
spatial occurrence and dimensions) as well as on the amount of precipitation, which are likely to be the
underlying issues. This helps explain the larger deviations between measured and modelled aerosol lifetimes
for the northernmost stations and at later time periods as seen in the previous section (Table 2).

5

## 6 5.3 Instantaneous lifetimes

The time variations in aerosol lifetimes can be examined further by considering the instantaneous lifetimes. 7  $\tau_{inst}$  (defined in Sect. 4). In this section we use global modelled data as opposed to the previous sections that 8 focused on modelled concentrations at the measurements stations. The estimate of instantaneous lifetimes 9 uses the global aerosol ( $^{137}$ Cs) and passive tracer ( $^{133}$ Xe) burdens as simulated by the models. Kristiansen et 10 al. (2012) also provide an estimate for the burdens using the station measurements and a 1D box model. This 11 estimate is however limited by the assumption of a well-mixed state in the atmosphere and only valid 12 approximately after about 3 weeks after the release until about 9 weeks (Appendix C). Therefore, global 13 burdens estimated from the measurement data are only available for a limited time period. The global 14 burdens are shown in Appendix C and Fig. C1. The modelled global aerosol (<sup>137</sup>Cs) burdens increase 15 initially due to continuous cesium emissions but decrease as the aerosols are removed from the atmosphere, 16 while the passive tracer (<sup>133</sup>Xe) global burdens stay approximately constant in the models after the xenon 17 emissions cease. In the following, we analyse the instantaneous lifetimes obtained from the global burdens. 18

We use the ratios of the aerosol ( $^{137}$ Cs) to the passive tracer ( $^{133}$ Xe) global burdens, and calculate the 19 instantaneous lifetimes (Eq. 2) from these ratios, as shown in Table 3 and Fig. 7. In the first week after the 20 start of the release, the modelled instantaneous lifetimes are about 2 days (Table 3), much shorter than the e-21 folding lifetimes presented in Sect. 5.1 for later time periods. This illustrates that the initial removal was 22 23 quicker, as previously suggested by Kristiansen et al. (2012) and illustrated in Croft et al. (2014). This is due to the emissions occurring at low altitudes and co-location of the plume with strong precipitation. After 24 about three weeks, estimates for the instantaneous lifetimes from the measurement data are also possible and 25 give a lifetime of 9.3-10.9 days, which is mostly longer than the median model lifetime of about 6.0-10.4 26 days (Table 3). After about seven weeks (day 45-50), the instantaneous lifetimes reach some quasi-steady 27 state with less variability (Fig. 7). The occurrence of this less variable steady state is related to the 28 emissions. Between day 10-40 there are low-level emissions of <sup>137</sup>Cs, about one order of magnitude less than 29 during the highest emissions around day 5. Because the emission was a point source (and not global) and its 30 geographical distribution is rather local, one continues to see the impact of single deposition events, making 31 the instantaneous lifetime fluctuate considerably. On day 40 all emissions of <sup>137</sup>Cs end in the model 32 simulations. The PBL is soon emptied from most <sup>137</sup>Cs, and the remaining <sup>137</sup>Cs (mainly in the free 33 troposphere) is more widely spread and gives rise to a more stable instantaneous lifetime, and thus a quasi-34 steady state with less variability. The modelled instantaneous lifetimes over days 49 to 63 (weeks 8-9; 35

shaded area in Fig. 7) vary between 2.8 and 19.0 days with a model median of 10.4±2.5 days. The median instantaneous lifetime for the same time period based on the measurements is 9.6±0.7 days. In this period, the median modelled lifetimes are closer to, but slightly longer than those based on measurement data, in contrast to earlier phases. This could be due to more prominent modelled than real mixing into the stratosphere where lifetimes are longer. This stratospheric mixing is not evident in the surface measurement data.

The further steady increase in the instantaneous lifetimes in the later phases after about day 65 (Fig. 7) is 7 attributed to transport and mixing into regions where the lifetimes are longer, such as the upper troposphere 8 and the stratosphere. After some time, the troposphere is mainly cleaned by wet scavenging while aerosols 9 that have been transported into the stratosphere remain, as the removal there is inefficient, especially for AM 10 aerosols for which also gravitational settling is very slow. The total aerosol mass and the lifetimes are then 11 dominated by the stratospheric loading, giving a continuous increase in lifetime (Cassiani et al., 2013). The 12 large variations in the steady increase of instantaneous lifetime between models might be related to how the 13 models simulate transport into the upper troposphere and stratosphere. The longest lifetimes at later stages 14 are probably given by models that inject a fraction of the aerosols quickly into the stratosphere (probably 15 even by a single event) and have an efficient removal in the troposphere, which would give a large (and 16 increasing) fraction of the aerosols residing in the upper troposphere and stratosphere. Furthermore, models 17 with diffusive advection schemes may excessively "leak" aerosols into the stratosphere. Thus, the 18 instantaneous lifetimes at later stages are probably more indicative of the upper tropospheric/stratospheric 19 aerosol fraction and are less useful to constrain tropospheric removal. In addition, the well-mixed 20 assumption required for the measurement-derived lifetimes breaks down once a substantial fraction of the 21 material is in the stratosphere. This means that measurement-derived and model-derived instantaneous 22 lifetimes for the later phases are not entirely comparable, as they are not derived in a consistent way. It is 23 expected that the measurement-derived instantaneous lifetimes are systematically lower, since they are 24 based on surface measurements with the assumption of a constant scale height. Calculating the global 25 burdens based on the modelled station data and the box model reveals that for some models the box-model 26 estimates indeed underestimate the full global burdens (see Appendix C and Fig. C1). 27

In summary, we find that the modelled instantaneous lifetimes were initially short (about 1-2 days) and the initial removal quicker than for later time periods. At later times (weeks 4-7) the instantaneous lifetimes obtained by measurements were generally longer than obtained from the models, as also seen in Sect. 5.1 from station locations. After that, the modelled instantaneous lifetimes are influenced by mixing into the stratosphere and not representative anymore of scavenging processes in the troposphere.

## 1 6 Discussion

## 2 6.1 Interpretation of the aerosol lifetime estimates

To interpret our estimates of aerosol lifetime, we first discuss an exemplified evolution of the aerosol burden 3 after a unit pulse emission (Fig. 8). In our example, the decay of the burden is not perfectly exponential but 4 consists of different characteristic timescales. There is a fast initial removal with a short time decay  $\tau_1$  which 5 represents the time period when the aerosols are mostly residing in the planetary boundary layer (PBL) and 6 are very susceptible to dry and wet deposition. The second period has slower removal with a time scale  $\tau_2$ , 7 and represents the period when most of the aerosols are residing in the free troposphere. The last time 8 period, with the slowest decay time scale  $\tau_3$ , is the period when most of the aerosols are removed from the 9 troposphere and only the fraction that was transported into the stratosphere remains. 10

The exemplified curve would depend on where on the globe the emissions take place, during which season, 11 and might differ from year to year. Thus, the time scales will vary greatly for different cases, and also for 12 different models handling the meteorology for the given case differently.  $\tau_1$  would be very different for all 13 cases, probably depending on a few precipitation events, while different cases and models would have a 14 smaller spread for  $\tau_2$  and  $\tau_3$  which are affected by processes on a more global scale. Global mean lifetimes 15 (e.g., Croft et al., 2014) would be mainly determined by the initial behaviour ( $\tau_1$ ) and it could be argued that 16 knowing the initial lifetime is important for estimating the aerosol burden, the radiative forcing and climate 17 sensitivity. 18

In this study we focus mainly on the intermediate timescale  $\tau_2$  because sufficient measurement data were 19 only available from about two weeks after the start of the release, until about day 65 (Sect. 5.1). We do not 20 have sufficient measurements to constrain  $\tau_1$ , but it is likely that some of the same processes contribute to 21 determine both  $\tau_1$  and  $\tau_2$ . While  $\tau_1$  is probably mainly impacted by processes within the PBL,  $\tau_2$  is 22 23 additionally influenced by how the aerosols are mixed out of the PBL, transported in the free troposphere and finally brought back into the PBL. That way it is possible that models that well capture  $\tau_2$  also perform 24 well for  $\tau_1$ . In our case, another difference between  $\tau_1$  and  $\tau_2$  is that the former was mostly determined by a 25 few individual precipitation events near Fukushima, while the latter is determined by removal occurring 26 worldwide. 27

28

29 The evolution of the burden in Fig. 8 can be approximated by the sum of three exponential functions, i.e.

30 
$$f(t) = f_1 \exp \frac{-t}{\tau_1} + f_2 \exp \frac{-t}{\tau_2} + f_3 \exp \frac{-t}{\tau_3}$$
(4)

31 where  $f_1 + f_2 + f_3 = 1$ .

32 A mean lifetime can be calculated as

$$\tau_{\rm m} = f_1 \tau_1 + f_2 \tau_2 + f_3 \tau_3 \tag{5}$$

2 which corresponds to the area under the curve f(t).

Using a optimizing convolution approach we can find the coefficients in Eq. 4 which give the best agreement 3 between the aerosol burdens obtained from the model simulations (Appendix C) and the expression in Eq. 4. 4 Table 4 shows the characteristic timescales  $\tau_1$ ,  $\tau_2$  and  $\tau_3$  obtained from the convolution approach, calculated 5 for the different models from their aerosol ( $^{137}$ Cs) burdens, as well as the mean lifetime  $\tau_m$  (Eq. 5). The initial 6 time scale  $\tau_1$  is fastest and around 2 days, which is related to the low altitude emissions of hydrophilic aerosols. 7 The intermediate time scale  $\tau_2$  is around 12 days, and the longest time scale  $\tau_3$  about 150-200 days. The 8 estimated  $\tau_3$  is very uncertain for most models as it is very close to the a priori estimate of 200 days. The 9 exceptions are the NorESM and GISS-ModelE models which had longer time series available. The model 10 mean and median for  $\tau_3$  are therefore calculated based on these two models only. The mean lifetime  $\tau_m$  is rather 11 short, on the order of 2-3 days, and is strongly affected by  $\tau_1$ , which was also demonstrated by Croft et al. 12 (2014). The estimate for  $\tau_2$  shows good agreement with the instantaneous lifetime  $\tau_{inst}$  over weeks 8-9 from 13 Fig. 7 and Table 3, and for most models the e-folding lifetime  $\tau_e$  from Fig. 3 and Table 2 also fit well with  $\tau_2$ . 14 This clearly demonstrate that in this study the focus is on the intermediate timescale  $\tau_2$ . 15

16

#### 17 6.2 Causes of model-observation deviations and model differences

Our results show that there are significant deviations between aerosol lifetimes obtained from models and those obtained from observations. The modelled lifetimes have a large spread but are generally shorter than the observed lifetime. In addition, large biases in model to observed surface concentrations were found, with an overall under-prediction of both the aerosol (<sup>137</sup>Cs) and passive tracer (<sup>133</sup>Xe) concentrations. The modelobservation deviations can be due to inaccuracies in the source emissions, errors in scavenging and convective transport, and incorrect diffusivity of the models. These issues are discussed more in the following.

Inaccuracies in the source emissions will affect the absolute model-simulated values and therefore the biases 25 in the model to observed concentrations (Figs. 4-6). It cannot be ruled out that our <sup>137</sup>Cs emissions are too 26 low, or there were errors in the injection heights. However, the <sup>137</sup>Cs source term used is rather on the high 27 side compared to others (e.g., Chino et al., 2011). Also, the <sup>133</sup>Xe source term has been confirmed 28 independently (Stohl et al., 2012b) and has rather low uncertainty. Overall, uncertainties in the assumed 29 source term and its implementation are likely not the main reasons for the general underestimation of the 30 aerosol (<sup>137</sup>Cs) and passive tracer (<sup>133</sup>Xe) concentrations. The lifetime estimates (Fig. 3 and Table 2) are not 31 affected by the absolute values of the source term used. However, these lifetime estimates would be affected 32 by possible late emissions of radionuclides long after the start of the release. Such late additional releases 33 could be either direct late emissions from FD-NPP or indirect releases by resuspension of deposited 34

radionuclides. Kristiansen et al. (2012) discussed this uncertainty and found no evidence for such late
 emissions, neither in the measurement data nor in the existing literature on the FD-NPP accident.

Most models have treated cesium solely as sulphate aerosols, while it is possible that some of the cesium 3 attached to other aerosol components with different deposition properties. Initially, many non-soluble 4 aerosols could have been present around the power plant, which might be removed less efficiently from the 5 6 atmosphere in the first days after emission before mixing internally with soluble aerosol components. The assumption that cesium attached solely to sulphate could result in too strong scavenging during the first few 7 days after the release, and can therefore contribute to the overall underestimations of the aerosol (<sup>137</sup>Cs) 8 concentrations (Fig. 5). The e-folding lifetimes (Table 2) should not be affected by errors in the modelled 9 scavenging efficiency in the initial phase ( $\tau_1$  in Fig. 8), as the e-folding lifetimes have been derived only for 10 later periods (after 15 days, i.e.,  $\tau_2$  in Fig. 8) when the aerosols are likely internally mixed and aerosol 11 components undergo similar removal. 12

More importantly, the models' treatment of aerosol scavenging is essential for both the modelled lifetime 13 and absolute concentrations. The modelled representation and distribution of clouds, as well as rain patterns 14 and rain intensity, are important factors since they determine where and how much aerosol is removed. 15 Further, the horizontal resolution of the model and meteorological input data might have a large impact in 16 terms of clouds and precipitation (intensity and spatial extent). Higher horizontal resolution can in general 17 lead to more resolved large-scale and less sub-grid convective clouds (Hagemann et al., 2006). The 18 scavenging parameterizations, including cloud and rain definitions and occurrences, differ greatly between 19 the models (Table A1), and are likely the major cause of model differences and deviations between models 20 and observations. This issue is further discussed in Sect. 6.6. 21

22 Additionally, the coupling between aerosol scavenging and the convection scheme in the models may represent a large source of uncertainty and thus contributes to both model-observation deviations and 23 differences between models. Different aerosol amounts may reach the dry upper troposphere depending on 24 how aerosols are scavenged in convective updrafts and whether updrafts are allowed in the convection 25 scheme to overshoot the level of neutral buoyancy. This has an immediate impact on lifetime via the 26 27 convective scavenging and also longer term effects, depending on how much aerosol can reach the upper troposphere. Finally, the diffusivity of the models, i.e. either pure numerical diffusivity or apparent 28 diffusivity caused by less representative winds, may be an important cause of the deviations between the 29 models and observations, and the differences between models. If the models are too diffusive, they may 30 unphysically transport aerosols to higher altitudes where they are preserved. Once too much aerosol mass 31 reaches high altitudes, e-folding lifetimes would be very long. At the same time too excessive vertical 32 diffusion would mean that the aerosols are not confined sufficiently to the lowest model levels (Koch et al., 33 2009) and thus the models will probably underestimate mass near the surface. These topics of coupling 34 convection with scavenging, and diffusivity should be investigated further. 35

## 2 6.3 Modelled aerosol lifetime versus aerosol concentration bias

Comparing the lifetime estimates in Table 2 and the aerosol concentration bias values in Fig. 5, it seems that 3 some models have a large bias to the measured aerosol concentrations, yet still have a realistic lifetime. E.g., 4 OsloCTM3 has a relatively small bias (ratio 0.76, i.e. -24% relative bias) and a lifetime of 8.8 days, while 5 OsloCTM2 highly underestimates the aerosol concentrations (ratio 0.02, i.e. -98% relative bias) and has a 6 lifetime estimate of 11.2 days, closer to the lifetime estimate from the observations (14.3 days). This is 7 probably because the concentrations are strongly influenced by the treatment of the quick removal in the 8 first few days when the plume was at low altitude and co-located with areas of strong precipitation, whereas 9 the lifetimes derived for later time periods are insensitive to this initial removal. Depending on how the 10 models simulate this initial episode (e.g., how the aerosol plume and precipitation areas align), different 11 amounts of aerosols can survive the first few days. This is clearly seen in Fig. 9, which shows how the total 12 atmospheric aerosol (<sup>137</sup>Cs) burdens for the different models vary over the first three weeks after the initial 13 release. As an example, OsloCTM2 has stronger aerosol removal than OsloCTM3 during the first three 14 weeks. Further, even relatively small differences in lifetimes can lead to large concentration differences after 15 several multiples of the lifetime. For instance, assume tracers with 6 and 10 days e-folding lifetimes. After 16 60 days, total atmospheric burdens of these two tracers will differ by a factor 50 (reduced to 1/403 and 17 1/22026, respectively, of the initial burden). This might also explain why aerosol models differ so widely 18 from each other at later times (e.g., in remote regions like the Arctic), whereas they agree much more closely 19 in earlier time periods and closer to the source regions (e.g. Fig. 5). 20

21

Additionally, by comparing the instantaneous lifetimes in Table 3 with the aerosol concentration bias values 22 in Fig. 5, the models with positive biases (greater than one, i.e. overestimation of the aerosol concentrations) 23 tend to show longer instantaneous lifetime in the first few weeks (Fig. 10a). The models that are strongly 24 25 underestimating the aerosol concentrations have a shorter instantaneous lifetime in the first weeks. This confirms that the initial quick removal is modelled differently by the various models, which affects the 26 absolute concentrations and bias calculations. Furthermore, there seems to be a positive correlation of 27 lifetimes for different time periods (Fig. 10b). Most models that have short instantaneous lifetimes in the 28 first three weeks also have short instantaneous lifetimes for later time periods. 29

30

## 31 6.4 Differences in modelled transport and scavenging

Our results show that there is a large spread in the model results. This is due to the different ways the models handle transport and scavenging. We note that the range of models in our analysis is quite wide. Many are developed for climate applications and may have been adapted for longer lifetime aspects, while others are more focused on dispersion and transport on a shorter time scale. In addition, some models are related to 1 each other, which will affect the model ensemble. However, as some models are related by driving

2 meteorology, others are related by aerosol module, which makes it difficult to categorize this appropriately.

Sensitivity simulations with the NorESM model (not shown) indicate that between five ensemble members, 3 all simulating their own meteorological conditions, there were no large differences in lifetime. This suggests 4 that the lifetime estimates are independent of the meteorological situation, although the difference between 5 the different models might be greater than a perturbed ensemble of one model, and resolution effects might 6 also play a role. Croft et al. (2014) showed that the e-folding lifetimes do not depend very much on the exact 7 model setup (emission altitude, location and time). Therefore, the model's scavenging parameterizations are 8 likely to be the main cause of inter-model differences in e-folding times in this study, especially at times 9 periods around 15-65 days after the release start. 10

11

The vertical distribution of the modelled aerosols will likely differ substantially between the various model 12 simulations. At later time periods, the troposphere would be mainly cleaned by wet scavenging while 13 aerosols that have been transported into the stratosphere (or upper troposphere) remain, as the removal there 14 is inefficient. The total aerosol mass and the lifetimes are then dominated by the upper tropospheric and 15 stratospheric loading, giving an increase in lifetime (Cassiani et al., 2013). The different stratospheric 16 loadings might therefore be a cause of large model differences. For example, in the OsloCTM2 and 17 OsloCTM3 simulations, about 70-80% of the burden is stratospheric at the end of the simulation. 18 19 OsloCTM2 has a higher stratospheric loading than OsloCTM3, which might be related to its coarser resolution and thus stronger diffusion, which partly explains why OsloCTM2 generally has a longer lifetime 20 (Table 2). By calculating tropospheric burdens and the associated lifetimes (burden relative to wet 21 deposition; not shown), this yields the lifetime with respect to scavenging without the effect of stratospheric 22 mixing. Then the lifetime for OsloCTM3 is longer than for OsloCTM2, highlighting the impact of the 23 different stratospheric aerosol loadings at later times. 24

25

The vertical distribution of the aerosols is explored further for four different models which span the range in 26 modelled lifetimes; FLEXPART with a short lifetime (~6 days); EEMEP with the longest (~27 days); and 27 OsloCTM3 and NorESM with lifetimes closer to, but still lower than the measurements (8.8 and 10.5 days, 28 respectively). Figure 11 shows the total aerosol (<sup>137</sup>Cs) mass as a function of latitude for these four models, 29 in four different height layers (surface, 0-5 km, 5-10 km and above 10 km) at three different time steps (14, 30 28 and 41 days after the first emissions). The aerosol concentrations in the surface layer (lowest model level) 31 peak around the latitude of the emission pulse for all models. FLEXPART and NorESM generally have 32 more aerosols at the surface than OsloCTM3 and EEMEP and this is particularly evident at the northernmost 33 latitudes. The same trend can be seen in the lower troposphere (0-5 km) except OsloCTM3 is closer to or 34 above FLEXPART while NorESM has consistently more mass at these heights. At higher altitudes (5-10 km 35 36 and above 10 km) OsloCTM3 and NorESM have the most aerosol mass at highest altitudes (above 10 km).

1 This indicates that models with lifetimes closer to the measurements have more mass at higher altitudes. The

2 FLEXPART and EEMEP models with lifetimes that deviate more from the observations show lower

- 3 fractions of the aerosol mass at higher altitudes.
- 4

The passive tracer also shows large differences between models (e.g. Fig. 4). Even models from the same 5 "family" (e.g. GISS, CAM and ECHAM) give quite different results for the passive tracer. In addition, the 6 four models based on ECHAM5 seem to simulate the passive tracer transport quite differently. This could be 7 due to differences in the driving meteorological fields or advection schemes used in the models. The tracer 8 transport is affected by four processes: 1) large-scale transport, 2) convective transport, 3) turbulent 9 transport, and 4) artificial numerical diffusion. All four processes are sensitive to horizontal and vertical 10 resolution, even if the models are all nudged towards the same reanalysis. Another possibility is that the 11 implementation of the point source was treated differently in the models, e.g. the flux was added at slightly 12 different times (before or after certain physical processes) in the model (see Fig. 9). To identify the exact 13 reasons why there are such large differences warrants a further study. 14

15

## 16 6.5 Sensitivity of e-folding times to measurement sampling

In this study we have used modelled surface concentrations at the times when observation data exist, 17 amounting to 433 data points over all 11 stations and measurement times (March to June 2011) considered. 18 Croft et al. (2014) did a similar lifetime estimate but derived from modelled surface concentrations of <sup>137</sup>Cs 19 from continuous time series at the 11 stations during March to June 2011 using about 19.000 hourly values. 20 They found a lifetime for GEOS-Chem of about 16.7 days, while in this study the lifetime estimate for 21 GEOS-Chem is longer (20.2 days). For five other models (GISS-modelE, FLEXPART, BCC SGCM CAM, 22 EMAC-1 and EMAC-2) we calculated the difference in the estimated lifetime using continuous (1-hourly or 23 3-hourly) time series at the stations, compared to using only the 433 data points at the times of the 24 measurements. For all five models, the lifetime is slightly shorter (by 0.2 to 0.9 days) when using the 25 26 continuous data than when using only data at the measurement times, but the difference is much smaller than for GEOS-Chem (3.5 days). GEOS-Chem is sensitive to the initial scavenging and transport as apparent in 27 Fig. 3 by the anomalous values at Wake Island in the early time period. Insufficient scavenging at mid-high 28 latitudes may also play a role, as seen in the overestimation of aerosol concentrations in Fig. 5, and this 29 30 seems to contribute to the longer lifetime for the selection of points used in this study. Further discussion on the scavenging parameterization in GEOS-Chem is given in Sect. 6.6. 31

Another issue with the measurement sampling is how to deal with measurement values below detection limit (BDL). In our analysis we have discarded all data BDL, however, this might bias the measurements high. To address this, we made a sensitivity test where all measurements values BLD were set to 1% of the detection limit, and included in the analysis. This increased the number of station values from 433 to 495. The stations with the most data BDL are Ulan Bator, Ashland and St. Johns, and the BDL values occur mostly between

days 40 to 70. During the days 15 to 65, the time period of the lifetime estimate in Fig. 3, there are 38 BDL 1 values mostly at Ulan Bator and Ashland. Using also BDL values in the analysis, the e-folding lifetime over 2 days 15-65 decrease by 0.7 days (from 14.3 to 13.6 days) compared to when discarding BDL values. This 3 illustrates that a small bias might be present when removing values BDL, but the difference is not significant 4 indicating that the measurement numbers are robust to BDL values. Besides, since the exact same stations 5 6 data points are used for the derivation of the e-folding lifetimes from the measurement data and the modelled data, it is likely that the same bias is present in both the modelled and measurement data sets and 7 therefore the comparison of modelled and measured lifetimes remains valid. 8

9

#### 10 6.6 Sensitivity of e-folding times and aerosol bias to wet removal parameterizations

A sensitivity test with the NAMEIII model was done (not shown) where the model's scavenging coefficients were decreased by a factor 10. It is not suggested that the initial scavenging coefficients were a factor 10 too high, but it was done to demonstrate the results of running the model with different scavenging coefficients. Decreasing the scavenging coefficients by a factor of 10 increased the e-folding lifetime from 5.5 days to 13.1 days (not shown), illustrating the dependency of estimated lifetimes on scavenging treatment.

Further, recent revisions to GEOS-Chem that introduce an explicit dependence of wet removal efficiency on 16 (1) cloud liquid and ice water content, (2) cloud temperature and (3) cloud fraction from the assimilated 17 meteorological fields have vielded about 40% lower mean lifetimes in the Arctic spring and summer (Croft 18 et al. (in prep.); Wang et al., 2014). Two revised simulations based on these changes, GEOS-Chem\_T258 19 and GEOS-Chem\_allT, are included here (Fig. 12 and Table 5) for comparison to the default scavenging in 20 GEOS- Chem (GEOS-Chem\_Orig, same simulation as labelled GEOS-Chem in all previous tables and 21 figures). Simulation GEOS-Chem T258 updates the scavenging scheme relative to GEOS-Chem Orig 22 through processes (1) - (3), but preserves the restriction for large-scale in-cloud scavenging of aerosols to 23 temperatures above 258K. Simulation GEOS-Chem T258 produces a significantly lower aerosol 24 concentration bias (1.69 vs 5.53) relative to observations (Fig. 12). As well, the e-folding lifetimes for the 25 station data are shorter by up to 6 days (Table 5). These improvements imply that using an explicit 26 dependence of wet removal rates on cloud properties may be necessary to properly capture aerosol lifetimes. 27 Simulation GEOS-Chem T258 yields the greatest change in the lifetimes for earlier periods (about 30% for 28 days 25-45, Table 5), indicating that these scavenging revisions influence removal efficiency in or near the 29 boundary layer more than in the free troposphere. The simulated station e-folding lifetimes for days 45-65 30 are insensitive to these scavenging revisions at temperatures warmer than 258K, suggesting that the day 46-31 65 e-folding lifetimes characterize scavenging efficiency at colder temperatures in the free troposphere. 32 Implementing these large-scale scavenging revisions at all temperatures (simulation GEOS-Chem\_allT) 33 yields the closest agreement with the observed day 45-65 e-folding lifetimes (13.2 days vs. 14.7 days), 34 although the aerosol concentration bias relative to observations become negative (ratio 48, i.e. -0.52 relative 35

1 bias). This illustrates the important control of scavenging efficiency in the free troposphere and at

2 temperatures colder than 258K on these e-folding lifetimes.

3

## 4 6.7 Comparison to other recent studies

The present study suggests that models generally have a too short sulphate lifetime, while in a recent study 5 Samset et al. (2014) find that models seem to have a too long black carbon (BC) lifetime, in line with other 6 recent studies (Bauer et al., 2013; Wang et al., 2014). These and other studies (e.g., Wang et al., 2013b) 7 8 using vertical profiles of BC from aircraft measurements have shown that the models tend to underestimate the concentrations close to ground, while overestimating at higher altitudes, particularly in the upper 9 troposphere. The overall tendency for the atmospheric column is an overestimation and hence a conclusion 10 that models require a shorter lifetime and more wet removal to re-produce the measurements. However, a 11 uniform reduction of the modelled BC lifetime would lead to an even larger underestimation at the surface 12 in the Arctic (Eckhardt et al., 2015). This points to considerable regional and vertical differences in 13 lifetimes, which was clearly illustrated by Croft et al. (2014) using the GEOS-Chem model. They showed 14 that modelled lifetimes were lowest and less than 5 days in the boundary layer below 2 km, and increased by 15 several orders of magnitude with altitude as wet removal mechanisms became increasingly less efficient 16 with altitude. Lastly, transport issues may cause the apparent misfit between surface and higher altitudes. In 17 18 our study, we have shown that some models underestimate transport particularly to the northernmost stations. 19

Eckhardt et al. (2015) compared both surface- and aircraft measurements of sulphate and black carbon (BC) 20 in the Arctic to model output from eleven different models. They found that the models generally 21 underestimate the surface concentrations of BC and sulphate in winter/spring, whereas concentrations in 22 summer were overestimated. For sulphate, they found very large differences in the model ensemble, with an 23 apparent anti-correlation between modelled surface concentrations and total atmospheric columns. They also 24 found a strong correlation between surface measured sulphate and BC concentrations, which indicated that 25 the sources contributing to sulphate and BC are similar throughout the Arctic, and that the aerosols are 26 internally mixed and undergo similar removal. Neither Eckhardt et al. (2015) nor Samset et al. (2014) found 27 an obvious factor (model type or aerosol treatment) that could explain why some models performed better 28 29 than others.

Our findings suggest that the models tend to underestimate aerosol lifetimes compared to radionuclide observations, in contrast to what is found for BC from some of the other studies mentioned above. Of course, the differences may arise from the different properties of sulphate and BC. The removal time of BC depends strongly on its mixing state and emitted size distribution (Reddington et al., 2013). Specifically, in models the BC removal time scale is also dependent on the aging parameterizations, which are often very simple and might be a confounding factor. This could mean that our results are not representative for BC

and are certainly not representative for freshly emitted BC. However, the high correlation and similar 1 behaviour of sulphate and BC found by Eckhardt et al. (2015) suggests that their fate in the remote 2 atmosphere is similar. The disagreement between models and observations is particularly evident for the 3 Arctic regions, and there seems to be a mismatch between the surface and at higher altitudes. Therefore, we 4 recommend that radionuclides, in particular <sup>133</sup>Xe and <sup>137</sup>Cs, be measured in the free troposphere in the days 5 6 and weeks following any detection coming from abnormal atmospheric releases. Such measurement will allow us to understand if the models can reproduce the atmospheric burden and vertical profiles of the 7 aerosols, and better constrain modelled aerosol lifetimes. 8

9

#### 10 7 Conclusions

In this study, we have compared measured and modelled accumulation-mode (AM) aerosol lifetimes, using 11 radioactive isotopes released during the Fukushima Dai-Ichi nuclear power plant accident of March 2011. 12 The radioactive isotope <sup>137</sup>Cs released in large quantities during the accident, attached to particles in the 13 ambient air, approximately according to the particle surface area, which is generally dominated by the 14 accumulation mode (AM). Measurements suggested that sulphate aerosols were the main carrier for the 15 cesium, and cesium was therefore used as a tracer for the AM sulphate aerosol's fate in the atmosphere. In 16 contrast, the noble gas xenon (<sup>133</sup>Xe), also released during the accident, behaved almost like a passive tracer 17 and served as a reference species for atmospheric transport. Global measurements of the two radioactive 18 isotopes taken over several months allowed the quantification of the lifetime of the carrier aerosols. The 19 lifetimes apply to aerosols that have undergone long-range transport (after about 2-3 weeks), i.e. the results 20 presented cannot directly constrain the lifetime of freshly emitted aerosols or secondary aerosols produced in 21 22 the boundary layer.

19 global models simulated the transport of the radioactive isotopes using identical emissions. We
investigated to what extent the models could reproduce the observations, especially with respect to the
observed loss of aerosol mass with time. Model results sampled at exactly the same location and times as
station measurements allowed a direct comparison between measured and modelled aerosol decay and
provided a strong constraint on modelled aerosol lifetime. Concentrations at measurement sites and global
atmospheric burdens were used to evaluate the modelled lifetime of the aerosols.

#### 29 Our main findings are

The e-folding lifetime τ<sub>e</sub> estimated from station measurements taken about two to nine weeks after
 the start of the emissions, is 14.3 days (95% confidence interval 13.1-15.7 days), and serve as an
 estimate for the lifetime of AM sulphate aerosol. The equivalent modelled τ<sub>e</sub> lifetimes have a large
 spread, varying between 4.8 and 26.7 days with a model median of 9.4±2.3 days. Modelled
 instantaneous lifetimes show that the initial removal was quicker due to the emissions occurring at

low altitudes and co-location with strong precipitation. Both e-folding and instantaneous lifetime
 estimates show that the models generally give a slightly shorter aerosol lifetime than observed. This
 is in contrast to recent findings for black carbon, which showed too long modelled lifetimes
 compared to aircraft measurements. More measurements of both radionuclides and different aerosol
 components, particularly in the free troposphere, are needed to better constrain modelled lifetimes,
 and understand these inconsistencies.

- Deviations between measured and modelled aerosol lifetimes are largest for the northernmost
   stations and at later time periods. Comparisons of measured and modelled aerosol (<sup>137</sup>Cs) and passive
   tracer (<sup>133</sup>Xe) concentrations at each measurement station show a general underestimation at high
   latitudes, suggesting that both scavenging and transport are causes for disagreements with
   observations. The underestimation is largest for the aerosols (<sup>137</sup>Cs), suggesting that scavenging and
   related cloud processes are the major reasons for the deviation from observations.
- Some models have a large bias to the measured aerosol concentrations, while at the same time 13 showing a realistic lifetime. This is likely due to the treatment of the initial quick removal that affects 14 the aerosol concentrations but not the lifetime estimate, which is derived only for later periods (after 15 two weeks). Different approaches might be needed for tackling over- or under-scavenging close to 16 the source emission in the early phase, and global-scale scavenging at later time periods. Revisions 17 to one model that introduced an explicit dependence of wet removal efficiency on cloud liquid and 18 ice water content, cloud temperature and cloud fraction from the assimilated meteorological fields, 19 yielded significantly improved results. 20
- How representative or diffusive the transport schemes of the models are, and to what extent they
   produce too fast vertical transport, should be investigated further. Excessive diffusion in the models
   would mean that too much aerosol mass would be present at high altitudes, thereby increasing e folding lifetimes independently of scavenging, while at the same time the models will likely
   underestimate aerosol concentrations near the surface.
- 26

## 1 Appendix A: Aerosol modules

2 Table A1 gives an overview of each model's treatment of the aerosols, i.e. the aerosol module.

3

## 4 Appendix B: Confidence intervals and R<sup>2</sup> statistics for lifetime estimates

Table B1 gives the 95% confidence interval (C.I.) for the e-folding lifetime  $\tau_e$  (Eq.1) from Table 2. Note that r<sub>e</sub> is not the centre of the C.I. as explained by Kristiansen et al. (2012). The table also gives the coefficients of determination R<sup>2</sup> which measure how close the data are to the fitted exponential decay models, with 1.0 indicating a perfect fit, and values that are not statistically significant at the 99.9% level are marked with an asterisk.

10

## 11 Appendix C: Global aerosol burden

The aerosol lifetime can also be assessed through the decay of the global atmospheric aerosol ( $^{137}$ Cs) 12 burden. Figure C1 shows different estimates of measured and modelled atmospheric burdens of both the 13 aerosol ( $^{137}$ Cs) and the passive tracer ( $^{133}$ Xe). First, global burdens are estimated with a 1D-box model using 14 the station measurements (Kristiansen et al., 2012). The box model assumes that the station concentrations at 15 the ground are representative for the depth of the tropospheric column and for the latitude band a certain 16 station is located in. This requires an assumption of a well-mixed state, which occurs after about 20 days 17 (Kristiansen et al., 2012). The same box-model calculations were performed on the modelled station data 18 19 (dark green and blue lines in Fig. C1). Second, modelled global burdens were also calculated from the full 3D model output (light green and turquoise lines in Fig. C1). 20

21 Comparing the burdens calculated by applying the box model to the model simulated station data (dark green and blue lines in Fig. C1) with the modelled global burdens based on the full 3D model output (bright 22 green and turquoise lines) serves as an evaluation of the box model assumption. For some models the global 23 burdens from the surface sites extrapolated with the box model compare well with the burdens from the full 24 model output directly, while for almost half of the models (GISS-ModelE, ULAO-CCM, CAM5, EMAC-1, 25 ECHAM-HAM2, EEMEP, OsloCTM2, NAME) the box-model estimates do not reconstruct the full global 26 burdens very well and a general underestimation from the box-model estimates are found, which is expected 27 since they are based on surface measurements. When estimating the lifetime from the modelled global 28 29 burdens, we choose to use the global burdens based on the full model output rather than those based on the modelled station data and the box model, while for the lifetime estimate from the measurements we must 30 rely on the box model and measured station data. This makes the comparison of lifetimes derived from the 31 global burden data not entirely consistent since the measured and modelled burdens are not calculated in the 32 exact same way. 33

The modelled passive tracer (<sup>133</sup>Xe) burdens (turquoise lines in Fig. C1) increase in the first few days due to 1 xenon emissions, which continue until day 5. Over the next months, the passive tracer burdens stay 2 approximately constant since it is not lost by any physical removal such as wet scavenging (and radioactive 3 decay is corrected for). The passive tracer global burdens based on the station data and box model (dark blue 4 lines) are only shown from day 20 onwards due to the required well-mixed assumption. Some slight 5 6 decrease seen in the measurements is related to leakage into the stratosphere or outside the domain (i.e., the Southern Hemisphere) considered for the box model estimates. The modelled global aerosol (<sup>137</sup>Cs) burdens 7 (bright green lines) also increase initially due to continuous cesium emissions which peak on day 3. Until 8 about day 40, the aerosol burdens vary as they are affected by the emissions. After about 10 days, the cesium 9 emissions drop significantly, but emissions continue between day 10 and 40 at about two orders of 10 magnitude lower than the largest emissions. Hence, occasionally the aerosol burdens are increasing after a 11 decrease up until day 40. After this day, the cesium burdens decrease as the aerosols are removed from the 12 atmosphere or are lost by mixing into the stratosphere. The ratio of the aerosol (<sup>137</sup>Cs) to the passive tracer 13 (<sup>133</sup>Xe) burdens (black lines) decreases due to loss of aerosols, mostly dominated by wet deposition. 14

The e-folding lifetime (Eq. 1) is calculated from the ratio of the aerosol ( $^{137}Cs$ ) to the passive tracer ( $^{133}Xe$ ) 15 burdens, by fitting an exponential decay model to the ratios between day 45 and day 65 (grey lines). Before 16 day 45, the global aerosol burdens vary due to continuous emissions and reach a quasi-steady state after 17 about 45-50 days when the burden values are becoming less variable. After day 65, the measurement 18 uncertainty becomes more substantial as the measured concentrations approach the instruments' detection 19 limit. The  $\tau_e$  lifetime based on the global burdens from the station measurements and the box model is 12.9 20 days, while the lifetime based on the modelled global burdens (from the full 3D model output) vary between 21 2.6 and 16.3 days with a model mean ( $\pm$  standard deviation) of 10.0 $\pm$ 4.1 days, and a model median ( $\pm$  MAD, 22 Eq. 3) of 9.4±2.4 days. 23

The lifetime derived from the box-model based burdens is more uncertain than the lifetime estimate from the 24 station data in Sect. 5.1 due to the required assumption of a well-mixed state of the atmosphere. The 25 assumption is violated due to slow mixing into the stratosphere. Once a large fraction of the material is in 26 the stratosphere the well-mixed assumption breaks down. This effect becomes more prominent at later times 27 as the fraction of aerosols (<sup>137</sup>Cs) in the stratosphere increases while most of the tropospheric aerosols 28 (<sup>137</sup>Cs) have been removed by scavenging. Loss to the stratosphere cannot be distinguished from 29 tropospheric removal. Errors due to variability caused by the relatively poor representivity of the 30 measurement network may also exist in the box-model estimates. The modelled global burdens calculated 31 from the full 3D model outputs should not have this problem. That means that measurement-derived and 32 model-derived lifetimes for the later phases are not entirely comparable, as they are not derived in a 33 34 consistent way.

- 1 In summary, the lifetime estimates based on global atmospheric burdens show that the modelled lifetimes
- 2 are somewhat shorter than those based on the measurements. This is the same trend as seen for the station
- 3 data in Sect. 5.1. However, the lifetime estimates based on the box-model calculations are more uncertain
- 4 due to the required well-mixed assumption.

## **1** Author contribution

2 N. I. K. and A. S. designed the experiment, performed the FLEXPART simulations and prepared the

- 3 manuscript. The NorESM simulations and the optimizing convolution approach calculations were done by
- 4 D. J. L. O. The GEOS-Chem simulations were conducted by B. C. with input from J. R. P. and R. V. M.
- 5 The OsloCTM2 and OsloCTM3 simulations were done by O. A. S., while the EEMEP simulations were
- 6 carried out by H. K. The EMAC-1 simulations were done by T. C. and the EMAC-2 simulations by D. K.
- 7 with help from H. T. The NAMEIII simulations were performed by S. J. L., and the GISS-ModelE2-
- 8 TOMAS simulations by Y.H. L. with input from D. S. The ECHAM5-HAM2 calculations were done by K.
- 9 Z., and K. T. completed the GISS-ModelE simulations with input from S. E. B. The ECHAM5-SALSA
- 10 simulations were conducted by T. B. with guidance from H. K, and the LMDZORINCA simulations were
- done by N. E. with input from Y. B. The CAM5 and CAM5\_PNNL simulations were carried out by H. W.
- 12 together with P.-L. M., R. C. E. and P. J. R., and the CAM5\_NDG simulations were done by K. Z., H. W.,
- and X. L. The ULAQ-CCM simulations were completed by G. P. and G. D G., and the
- 14 BCC\_AGCM\_2.0.1\_CAM simulations were conducted by S. Y. Z. with guidance from H. Z. Additional
- 15 input to the interpretation of the analysis was provided by M. S. and Y. B. The analysis of model results and
- 16 preparation of all figures and tables were done by N. I. K. All authors provided input to the manuscript.
- 17

#### 18 Acknowledgements

We would like to thank all the scientists who produced the CTBTO measurement data and made them 19 available to us. The research leading to these results has received partial funding from the Norwegian 20 Research Council under the NORKLIMA and KLIMAFORSK program (project "AeroCom-P3"). H. Zhang 21 and S. Y. Zhao are supported by the National Basic Research Program of China (Grant No.: 22 2011CB403405). H. Wang, R.C. Easter, P.-L. Ma, P.J. Rasch acknowledge support from the U.S. 23 Department of Energy (DOE), Office of Science, Biological and Environmental Research as part of the 24 Earth System Modeling Program. T. Bergman and H. Kokkola were supported by the Academy of Finland 25 Centre of Excellence (project no. 272041). The ECHAM-HAMMOZ model is developed by a consortium 26 composed of ETH Zurich, Max Planck Institut für Meteorologie, Forschungszentrum Jülich, University of 27 Oxford, the Finnish Meteorological Institute and the Leibniz Institute for Tropospheric Research, and 28 managed by the Center for Climate Systems Modeling (C2SM) at ETH Zurich. The GISS model 29 group acknowledges resources supporting this work, provided by the NASA High-End Computing (HEC) 30 Program through the NASA Center for Climate Simulation (NCCS) at Goddard Space Flight Center. 31

### 1 References

Adams, P. J., and Seinfeld, J. H.: Predicting global aerosol size distributions in general circulation models, J.
Geophys. Res., 107, doi:10.1029/2001JD001010, 2002.

Andreae, M. O.: Atmospheric aerosols versus greenhouse gases in the twenty-first century, Phil. Trans. R.
Soc. A (2007) 365, 1915–1923 doi:10.1098/rsta.2007.2051, 2007.

- Bauer, S. E., Bausch, A., Nazarenko, L., Tsigaridis, K., Xu, B. Q., Edwards, R., Bisiaux, M., and
  McConnell, J.: Historical and future black carbon deposition on the three ice caps: ice core measurements
  and model simulations from 1850 to 2100, J. Geophys. Res., 118, 7948–7961, doi:10.1002/Jgrd.50612,
  2013.
- 10 Berglen, T. F.; T. K. Berntsen, I. S. A. Isaksen, and J. K. Sundet: A global model of the coupled
- 11 sulfur/oxidant chemistry in the troposphere: The sulfur cycle. J. Geophys. Res., 109(D19310),
- 12 doi:10.1029/2003JD003948, 2004.
- 13 Bergman, T., Kerminen, V.-M., Korhonen, H., Lehtinen, K. J., Makkonen, R., Arola, A., Mielonen, T.,
- Romakkaniemi, S., Kulmala, M., and Kokkola, H.: Evaluation of the sectional aerosol microphysics module
  SALSA implementation in ECHAM5-HAM aerosol-climate model, Geosci. Model Dev., 5, 845-868,
  doi:10.5194/gmd-5-845-2012, 2012.
- 17 Bentsen, M., Bethke, I., Debernard, J. B., Iversen, T., Kirkevåg, A., Seland, Ø., Drange, H., Roelandt, C.,
- 18 Seierstad, I. A., Hoose, C., and Kristjánsson, J. E.: The Norwegian Earth System Model, NorESM1-M Part
- 19 1: Description and basic evaluation of the physical climate, Geosci. Model Dev., 6, 687–720,
- 20 doi:10.5194/gmd-6-687-2013, 2013.
- Bey, I., Jacob, D. J., Logan, J. A., and Yantosca, R. M.: Asian chemical outflow to the Pacific in spring:
  Origins, pathways and bud- gets, J. Geophys. Res., 106, 23097–23113, doi: 10.1029/2001JD000806, 2001.
- Bourgeois, Q. and Bey, I.: Pollution transport efficiency toward the Arctic: Sensitivity to aerosol scavenging
  and source regions, J. Geophys. Res., 116, D08213, doi:10.1029/2010JD015096, 2011.
- 25 Browse, J., K. S. Carslaw, S. Arnold, K. J. Pringle and O. Boucher, The scavenging processes controlling
- the seasonal cycle in Arctic sulfate and black carbon aerosol, Atmos. Chem. Phys., 12, 6775 6798, doi:
  10.5194/acp-12-6775-2012, 2012.
- 28 Cambray, R. S., Cawse, P. A., Garland, J. A., Gibson, J.A. B., John- son, P., Lewis, G. N. J., Newton, D.,
- Salmon, L., and Wade, B. O.: Observations on radioactivity from the Chernobyl accident, Nuclear Energy
  26, 77–101, 1987.

- Cassiani, M., Stohl, A., and Eckhardt, S.: The dispersion characteristics of air pollution from the world's
   megacities, Atmos. Chem. Phys., 13, doi:10.5194/acp-13-9975-2013, 9975–9996, 2013.
- Chin, M., Jacob, D. J., Gardner, G. M., Foreman-Fowler, M. S., Spiro, P. A., and Savoie, D. L.: A global
  three-dimensional model of tropospheric sulfate, J. Geophys. Res., 101, 18667–18690, doi:
  10.1029/96JD01221, 1996.
- 5 10.1029/9001201201, 1990.
  - 6 Chino, M., Nakayama, H., Nagai, H., Terada, H., Katata, G., and Yamazawa, H.: Preliminary estimation of
    7 release amounts of 1311 and 137Cs accidentally discharged from the Fukushima Daiichi nuclear power plant
  - 8 into the atmosphere, J. Nuc. Sci. Tech., 48, 1129–1134, , doi: 10.1080/18811248.2011.9711799, 2011.
- 9 Christoudias, T. and Lelieveld, J.: Modelling the global atmospheric transport and deposition of
- 10 radionuclides from the Fukushima Dai-ichi nuclear accident, Atmos. Chem. Phys., 13, 1425-1438,
- 11 doi:10.5194/acp-13-1425-2013, 2013.
- 12 Chung, S. H. and Seinfeld, J. H.: Global distribution and climate forcing of carbonaceous aerosols, J.
- 13 Geophys. Res., 107, 4407, doi:10.1029/2001JD001397, 2002.
- Croft, B., Martin, R. V., Leaitch, W. R., Breider, T., D'Andrea, S. D., and Pierce, J. R.: Processes controlling
  the seasonal cycle of arctic aerosol number and size distributions, in prep., 2015.
- Croft, B., Pierce, J. R., and Martin, R. V.: Interpreting aerosol lifetimes using the GEOS-Chem model and
  constraints from radionuclide measurements, Atmos. Chem. Phys., 14, 4313–4325, doi:10.5194/acp-144313-2014, 2014.
- 19 Eckhardt, S., Quennehen, B., Olivié, D. J. L., Berntsen, T. K., Cherian, R., Christensen, J. H., Collins, W.,
- 20 Crepinsek, S., Daskalakis, N., Flanner, M., Herber, A., Heyes, C., Hodnebrog, Ø., Huang, L.,
- 21 Kanakidou, M., Klimont, Z., Langner, J., Law, K. S., Massling, A., Myriokefalitakis, S., Nielsen, I. E.,
- 22 Nøjgaard, J. K., Quaas, J., Quinn, P. K., Raut, J.-C., Rumbold, S. T., Schulz, M., Skeie, R. B., Skov, H.,
- 23 Lund, M. T., Uttal, T., von Salzen, K., Mahmood, R., and Stohl, A.: Current model capabilities for
- simulating black carbon and sulfate concentrations in the Arctic atmosphere: a multi-model evaluation using
- a comprehensive measurement data set, Atmos. Chem. Phys. Discuss., 15, 10425-10477, doi:10.5194/acpd15-10425-2015, 2015.
- 27 Evangeliou, N., Balkanski, Y., Cozic, A., and Møller A. P.: Global Transport and Deposition of 137Cs
- Following the Fukushima Nuclear Power Plant Accident in Japan: Emphasis on Europe and Asia Using
- 29 High–Resolution Model Versions and Radiological Impact Assessment of the Human Population and the
- 30 Environment Using Interactive Tools, Environ. Sci. Technol., 47, 5803-5812, doi: 10.1021/es400372u,
- 31 2013.

- 1 Feichter, J., Kjellstrom, E., Rodhe, H., Dentener, F., Lelieveld, J., and Roelofs, G.-J.: Simulation of the
- 2 tropospheric sulphur cycle in a global climate model, Atmos. Environ., 30, 1693–1707, 1996.
- Friedlander, S. K.: Smoke, dust and haze: Fundamentals of aerosol behavior, New York, Wiley-Interscience,
  333 pp., doi: 10.1016/1352-2310(95)00394-0, 1977.
- 5 Giorgi, F. and Chameides, W. I.: Rainout lifetimes of highly soluble aerosols and gases as inferred from
- 6 simulations with a General Circulation Model, J. Geophys. Res , 91, 14367–14376, doi:
- 7 10.1029/JD091iD13p14367, 1986.
- Gong, S. L., Barrie, L. A., Blanchet, J.-P., von Salzen, K., Lohmann, U., Lensins, G., Spacek, L., Zhang, L.
  M., Girard, E., Lin, H., Leaitch, R., Leighton, H., Chylek, P., and Huang, P.: Canadian Aerosol Module: A
  size-segregated simulation of atmospheric aerosol processes for climate and air quality models 1. Module
  development, J. Geophys. Res., 108, D1, 4007, 10.1029/2001JD002002, 2003.
- Hagemann, S., Arpe, K. and Roeckner E.: Evaluation of the Hydrological Cycle in the ECHAM5 Model. J.
  Climate, 19, 3810–3827.doi: http://dx.doi.org/10.1175/JCLI3831.1, 2006.
- 14 Iversen, T., Bentsen, M., Bethke, I., Debernard, J. B., Kirkevåg, A., Seland, Ø., Drange, H., Kristjánsson, J.
- 15 E., Medhaug, I., Sand, M., and Seierstad, I. A.: The Norwegian Earth System Model, NorESM1-M Part 2:
- Climate response and scenario projections, Geosci. Model Dev., 6, 389–415, doi:10.5194/gmd-6-389-2013,
  2013.
- Kaneyasu, N., Ohashi, H., Suzuki, F., Okuda, T., and Ikemori, F.: Sulfate aerosol as a potential transport
  medium of radiocesium from the Fukushima nuclear accident, Environ. Sci.Technol., 46, 5720–5726
  doi:10.1021/es204667h, 2012.
- 21 Kerkweg, A., Buchholz, J., Ganzeveld, L., Pozzer, A., Tost, H., and Jöckel, P.: Technical Note: An
- 22 implementation of the dry removal processes DRY DEPosition and SEDImentation in the Modular Earth
- 23 Submodel System (MESSy), Atmos. Chem. Phys., 6, 4617-4632, doi:10.5194/acp-6-4617-2006, 2006.
- 24 Kirkevåg, A., Iversen, T., Seland, Ø., Hoose, C., Kristjánsson, J. E., Struthers, H., Ekman, A. M. L., Ghan,
- 25 S., Griesfeller, J., Nilsson, E. D., and Schulz, M.: Aerosol climate interactions in the Norwegian Earth
- 26 System Model NorESM1-M, Geosci. Model Dev., 6, 207–244, doi:10.5194/gmd-6-207-2013, URL
- 27 http://www.geosci-model-dev.net/6/207/2013/, 2013.
- 28 Koch, D. and Hansen, J.: Distant origins of Arctic black carbon, A Goddard Institute for Space Studies
- 29 ModelE experiment, J. Geophys. Res., 110, D04204, doi:10.1029/2004jd005296, 2005.
- 30 Koch, D., Schmidt, G. A., and Field, C. V.: Sulfur, sea salt, and radionuclide aerosols in GISS ModelE, J.
- 31 Geophys. Res., 111, D06206, doi:10.1029/2004jd005550, 2006.

- 1 Koch, D., Schulz, M., Kinne, S., McNaughton, C., Spackman, J. R., Balkanski, Y., Bauer, S., Berntsen, T.,
- 2 Bond, T. C., Boucher, O., Chin, M., Clarke, A., De Luca, N., Dentener, F., Diehl, T., Dubovik, O.,
- 3 Easter, R., Fahey, D. W., Feichter, J., Fillmore, D., Freitag, S., Ghan, S., Ginoux, P., Gong, S., Horowitz, L.,
- 4 Iversen, T., Kirkevåg, A., Klimont, Z., Kondo, Y., Krol, M., Liu, X., Miller, R., Montanaro, V., Moteki, N.,
- 5 Myhre, G., Penner, J. E., Perlwitz, J., Pitari, G., Reddy, S., Sahu, L., Sakamoto, H., Schuster, G.,
- 6 Schwarz, J. P., Seland, Ø., Stier, P., Takegawa, N., Takemura, T., Textor, C., van Aardenne, J. A., and
- 7 Zhao, Y.: Evaluation of black carbon estimations in global aerosol models, Atmos. Chem. Phys., 9, 9001-
- 8 9026, doi:10.5194/acp-9-9001-2009, 2009.
- 9 Kristiansen, N. I., A. Stohl, and G. Wotawa, Atmospheric removal times of the aerosol-bound
- 10 radionuclides 137Cs and 131I measured after the Fukushima Dai-ichi nuclear accident a constraint for air
- 11 quality and climate models, Atmos. Chem. Phys., 12, 10759-10769, doi:10.5194/acp-12-10759-2012,
- **12** 2012.
- 13 Kunkel, D. Global modeling of pollutant transport and deposition from anthropogenic emission hot spots on
- 14 global scale, PhD Dissertation, Johannes Gutenberg-Universität in Mainz, 2012.
- 15 Kunkel, D., Tost, H., and Lawrence, M. G.: Aerosol pollution potential from major population centers,
- 16 Atmos. Chem. Phys., 13, 4203-4222, doi:10.5194/acp-13-4203-2013, 2013.
- 17 Laakso, A., Kokkola, H., Partanen, A.-I., Niemeier, U., Timmreck, C., Lehtinen, K., Hakkarainen, H.
- and Korhonen, H.: Radiative and climate impacts of a large volcanic eruption during stratospheric sulfur
  geoengineering, submitted to ACPD, 2015.
- Leadbetter, S., Hort, M., Jones, A., Webster, H., and Draxler R.: Sensitivity of the deposition of Caesium137 from Fukushima Dai-ichi nuclear power plant on the wet deposition parameterisation in NAME. J. Env.
  Rad., 139, 200-211, doi: 10.1016/j.jenvrad.2014.03.018., 2015.
- 23 Lee, Y. H., Adams, P. J., and Shindell, D. T.: Evaluation of the global aerosol microphysical ModelE2-
- TOMAS model against satellite and ground-based observations, Geosci. Model Dev., 8, 631-667,
- 25 doi:10.5194/gmd-8-631-2015, 2015.
- Liu, H., Jacob, D. J., Bey, I. and Yantosca, R. M.: Constraints from 210Pb and 7Be on wet deposition and
  transport in a global three-dimensional chemical tracer model driven by assimilated meteorological fields, J.
  Geophys. Res., 106, 12109–12128, 2001.
- 29 Liu, X., Penner, J. E., and Herzog, M.: Global modeling of aerosol dynamics: Model description, evaluation,
- 30 and interactions between sulfate and nonsulfate aerosols, J. Geophys. Res., 110, D18206,
- doi:10.1029/2004JD005674, 2005.

- 1 Liu, X., Easter, R. C., Ghan, S. J., Zaveri, R., Rasch, P., Shi, X., Lamarque, J.-F., Gettelman, A.,
- 2 Morrison, H., Vitt, F., Conley, A., Park, S., Neale, R., Hannay, C., Ekman, A. M. L., Hess, P.,
- 3 Mahowald, N., Collins, W., Iacono, M. J., Bretherton, C. S., Flanner, M. G., and Mitchell, D.: Toward a
- 4 minimal representation of aerosols in climate models: description and evaluation in the Community
- 5 Atmosphere Model CAM5, Geosci. Model Dev., 5, 709-739, doi:10.5194/gmd-5-709-2012, 2012.
- 6 Ma, P.-L., P. J. Rasch, H. Wang, K. Zhang, R. C. Easter, S. Tilmes, J. D. Fast, X. Liu, J.-H. Yoon, and J.-F.
- 7 Lamarque: The role of circulation features on black carbon transport into the Arctic in the Community
- 8 Atmosphere Model version 5 (CAM5), J. Geophys. Res. Atmos., 118, 4657–4669, doi:10.1002/jgrd.50411,
  9 2013.
- 10 Masson, O., Ringer, W., Malá, H., Rulik, P., Dlugosz-Lisiecka, M., Eleftheriadis, K., Meisenberg, O., De
- 11 Vismes-Ott, A., and Gensdarmes, F.: Size Distributions of Airborne Radionuclides from the Fukushima
- 12 Nuclear Accident at Several Places in Europe, Environ. Sci. Technol., 47 (19), pp 10995–11003,
- 13 doi: 10.1021/es401973c, 2013.
- Medici, F.: The IMS radionuclide network of the CTBT, Radiat. Phys. Chem., 61, 689–690, doi:
  10.1016/S0969-806X(01)00375-9, 2001.
- 16 Miyamoto, Y., Yasuda, K., Magara, M.: Size distribution of radioactive particles collected at Tokai, Japan 6
- 17 days after the nuclear accident, , Environ. Sci. Technol., 132, pp 1–7,
- 18 http://dx.doi.org/10.1016/j.jenvrad.2014.01.010, 2014.
- Moore, H. E., Poet, S. E., and Martell, E. A.: 222Rn, 210Pb, 210Bi, and 210Po profiles and aerosol
  residence times versus altitude, J. Geophys. Res., 78(30), 7065–7075, doi: 10.1029/JC078i030p07065, 1973.
- 21 Neu, J. L. and Prather, M. J.: Toward a more physical representation of precipitation scavenging in global
- chemistry models: cloud overlap and ice physics and their impact on tropospheric ozone, Atmos. Chem.
  Phys., 12, 3289–3310, doi:10.5194/acp-12-3289-2012, 2012.
- Papastefanou, C.: Residence time of tropospheric aerosols in association with radioactive nuclides, Applied
  Radiation and Isotopes, 64, 93–100, doi:10.1016/j.apradiso.2005.07.006, 2006.
- 26 Papastefanou, C.: Radioactive aerosols, Radioactivity in the environment, Volume 12, Series Editor: M. S.
- 27 Baxter, Elsevier, ISBN: 978-0-08-044075-0, 2008.
- 28 Paris, J.-D., Stohl, A., Nédélec, P., Arshinov, M. Yu., Panchenko, M. V., Shmargunov, V. P., Law, K. S.,
- 29 Belan, B. D., and Ciais, P.: Wildfire smoke in the Siberian Arctic in summer: source characterization and
- 30 plume evolution from airborne measurements, At- mos. Chem. Phys., 9, 9315–9327, doi:10.5194/acp-9-
- 31 9315-2009, 2009.

1 Pitari, G., V. Aquila, B. Kravitz, A. Robock, S. Watanabe, I. Cionni, N. De Luca, G. Di Genova, E. Mancini,

2 and S. Tilmes: Stratospheric Ozone Response to Sulfate Geoengineering: Results from the Geoengineering

- 3 Model Intercomparison Project (GeoMIP), J. Geophys. Res., 119, 2629-2653, doi:10.1002/2013JD020566,
- 4 2014.
- 5 Pringle, K. J., Tost, H., Message, S., Steil, B., Giannadaki, D., Nenes, A., Fountoukis, C., Stier, P.,
- 6 Vignati, E., and Lelieveld, J.: Description and evaluation of GMXe: a new aerosol submodel for global
- 7 simulations (v1), Geosci. Model Dev., 3, 391-412, doi:10.5194/gmd-3-391-2010, 2010.
- 8 Pöschl, U.: Atmospheric Aerosols: Composition, Transformation, Climate and Health Effects. Angew.
- 9 Chem. Int. Ed., 44: 7520–7540. doi: 10.1002/anie.200501122, 2005.
- Ramanathan, V., P. J. Crutzen, J. T. Kiehl, D. Rosenfeld: Aerosols, Climate, and the Hydrological Cycle,
  Science, Vol. 294 no. 5549 pp. 2119-2124, doi: 10.1126/science.1064034, 2001.
- 12 Reddington, C. L., McMeeking, G., Mann, G. W., Coe, H., Frontoso, M. G., Liu, D., Flynn, M., Spracklen,
- 13 D. V., and Carslaw, K. S.: The mass and number size distributions of black carbon aerosol over Europe,
- 14 Atmos. Chem. Phys., 13, 4917-4939, doi:10.5194/acp-13-4917-2013, 2013.
- Saey, P. R. J., and de Geer, L.-E.: Notes on radioxenon measurements for CTBT verification purposes,
  Appl. Radiat. Isot., 63, 765–773, doi: 10.1016/j.apradiso.2005.05.035, 2005.
- 17 Samset, B. H., Myhre, G., Herber, A., Kondo, Y., Li, S.-M., Moteki, N., Koike, M., Oshima, N.,
- 18 Schwarz, J. P., Balkanski, Y., Bauer, S. E., Bellouin, N., Berntsen, T. K., Bian, H., Chin, M., Diehl, T.,
- 19 Easter, R. C., Ghan, S. J., Iversen, T., Kirkevåg, A., Lamarque, J.-F., Lin, G., Liu, X., Penner, J. E.,
- 20 Schulz, M., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., and Zhang, K.: Modeled black
- 21 carbon radiative forcing and atmospheric lifetime in AeroCom Phase II constrained by aircraft observations,
- 22 Atmos. Chem. Phys., 14, 12465-12477, doi:10.5194/acp-14-12465-2014, 2014.
- 23 Schmale, J., Schneider, J., Ancellet, G., Quennehen, B., Stohl, A., Sodemann, H., Burkhart, J. F.,
- 24 Hamburger, T., Arnold, S. R., Schwarzenboeck, A., Borrmann, S., and Law, K. S.: Source identification and
- airborne chemical characterisation of aerosol pollution from long-range transport over Greenland during
- 26 POLAR- CAT summer campaign 2008, Atmos. Chem. Phys., 11, 10097–10123, doi:10.5194/acp-11-
- 27 10097-2011, 2011.
- 28 Schmidt, G. A., et al., Configuration and assessment of the Giss ModelE2 contributions to the CMIP5
- 29 archive, J. Adv. Model. Earth Syst., 6, 141–184, doi:10.1002/2013MS000265, 2014.
- 30 Schulze, J., Auer, M., and Werzi, R.: Low level radioactivity measurement in support of the CTBTO, Appl.
- 31 Radiat. Isot., 53, 23–30, 10.1016/j.apradiso.2005.05.035, 2000.

- 1 Seinfeld, J. H. and Pandis, S. N.: Atmospheric Chemistry and Physics: From Air Pollution to Climate
- 2 Change. New York, John Wiley and Sons, 1998.
- 3 Seland, Ø., Iversen, T., Kirkevåg, A., and Storelvmo, T.: Aerosol-climate interactions in the CAM-Oslo
- 4 atmospheric GCM and investigation of associated basic shortcomings, Tellus, 60A, 459–951, doi:
- 5 10.1111/j.1600-0870.2008.00318.x, 2008.
- 6 Shindell, D. T., Chin, M., Dentener, F., Doherty, R. M., Faluvegi, G., Fiore, A. M., Hess, P., Koch, D. M.,
- 7 MacKenzie, I. A., Sanderson, M. G., Schultz, M. G., Schulz, M., Stevenson, D. S., Teich, H., Textor, C.,
- 8 Wild, O., Bergmann, D. J., Bey, I., Bian, H., Cuvelier, C., Duncan, B. N., Folberth, G., Horowitz, L. W.,
- 9 Jonson, J., Kaminski, J. W., Marmer, E., Park, R., Pringle, K. J., Schroeder, S., Szopa, S., Takemura, T.,
- 10 Zeng, G., Keating, T. J., and Zuber, A.: A multi-model assessment of pollution transport to the Arctic,
- 11 Atmos. Chem. Phys., 8, 5353-5372, doi:10.5194/acp-8-5353-2008, 2008.
- 12 Stier, P., Feichter, J., Kinne, S., Kloster, S., Vignati, E., Wilson, J., Ganzeveld, L., Tegen, I., Werner, M.,
- 13 Balkanski, Y., Schulz, M., Boucher, O., Minikin, A., and Petzold, A.: The aerosol-climate model ECHAM5-
- 14 HAM, Atmos. Chem. Phys., 5, 1125–1156, doi:10.5194/acp-5-1125-2005, 2005.
- Stohl, A., Forster, C., Frank, A., Seibert, P., and Wotawa, G.: Technical note: The Lagrangian particle
  dispersion model FLEXPART version 6.2, Atmos. Chem. Phys., 5, 2461–2474, doi:10.5194/acp-5-24612005, 2005.
- 18 Stohl, A., P. Seibert, G. Wotawa, D. Arnold, J. F. Burkhart, S. Eckhardt, C. Tapia, A. Vargas, and T. J.
- 19 Yasunari: Xenon-133 and caesium-137 releases into the atmosphere from the Fukushima Dai-ichi nuclear
- 20 power plant: determination of the source term, atmospheric dispersion, and deposition, Atmos. Chem. Phys.,
- 21 12, 2313-2343, doi:10.5194/acp-12-2313-2012, 2012.
- Stohl, A., P. Seibert, and G. Wotawa, The total release of xenon-133 from the Fukushima Dai-ichi nuclear
  power plant accident. J. Environ. Radioact. 112, 155-159, doi:10.1016/j.jenvrad.2012.06.001, 2012b.
- 24 Søvde, O. A., M, Gauss, S. Smyshlyaev, and I. S. A. Isaksen: Evaluation of the chemical transport model
- 25 Oslo CTM2 with focus on Arctic winter ozone depletion. J. Geophys. Res., vol. 113, D09304,
- 26 doi:10.1029/2007JD009240, 2008.
- Søvde, O. A.; M. J. Prather, I. S. A. Isaksen, T. K. Berntsen, F. Stordal, X. Zhu, C. D. Holmes and J. Hsu:
  The chemical transport model Oslo CTM3, Geosci. Model Dev., 5, 1441-1469, doi:10.5194/gmd-5-14412012, 2012.
- 30 Textor, C., Schulz, M., Guibert, S., Kinne, S., Balkanski, Y., Bauer, S., Berntsen, T., Berglen, T., Boucher,
- O., Chin, M., Dentener, F., Diehl, T., Easter, R., Feichter, H., Fillmore, D., Ghan, S., Ginoux, P., Gong, S.,
- 32 Grini, A., Hendricks, J., Horowitz, L., Huang, P., Isaksen, I., Iversen, I., Kloster, S., Koch, D., Kirkevåg, A.,

- Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Liu, X., Montanaro, V., Myhre, G., Penner, J., 1
- Pitari, G., Reddy, S., Seland, Ø., Stier, P., Takemura, T., and Tie, X.: Analysis and quantification of the 2
- diversities of aerosol life cycles within AeroCom, Atmos. Chem. Phys., 6, 1777-1813, doi:10.5194/acp-6-3 1777-2006, 2006.
- 4
- Tsigaridis, K., Koch, D., and Menon, S.: Uncertainties and importance of sea spray composition on aerosol 5 6 direct and indirect effects, J. Geophys. Res.-Atmos., 118, 220-235, doi:10.1029/2012jd018165, 2013.
- Vignati, E., J. Wilson, and P. Stier: M7: An efficient size-resolved aerosol microphysics module for large-7 scale aerosol transport models, J. Geophys. Res., 109, D22202, doi:10.1029/2003JD004485, 2004. 8
- Wang, Q., Jacob, D. J., Spackman, J. R., Perring, A. E., Schwarz, J. P., Moteki, N., Marais, E. A., Ge, C., 9
- Wang, J., and Barrett, S. R. H.: Global budget and radiative forcing of black carbon aerosol: Constraints 10 from pole-to-pole (HIPPO) observations across the Pacific, J. Geophys. Res.-Atmos., 119, 195–206,
- 11 doi:10.1002/2013jd020824, 2014. 12
- Wang, Z., Zhang, H., Jing, X., Wei, X.: Effect of non-spherical dust aerosol on its direct radiative forcing. 13 Atmospheric Research, 120:121, 112-126, doi: 10.1016/j.atmosres.2012.08.006, 2013a. 14
- Wang, H., Easter, R. C., Rasch, P. J., Wang, M., Liu, X., Ghan, S. J., Qian, Y., Yoon, J.-H., Ma, P.-L., and 15
- Vinoj, V.: Sensitivity of remote aerosol distributions to representation of cloud-aerosol interactions in a 16
- global climate model, Geosci. Model Dev., 6, 765-782, doi:10.5194/gmd-6-765-2013, 2013b. 17
- Wang, H., Rasch, P. J., Easter, R. C., Singh, B., Zhang, R., Ma, P. L., Qian, Y., and Beagley, N.: Using an 18
- explicit emission tagging method in global modeling of source-receptor relationships for black carbon in the 19
- Arctic: Variations, Sources and Transport pathways, J. Geophys. Res.-Atmos., 119, 12888–12909, doi: 20
- 10.1002/2014JD022297, 2014b. 21
- Webster H. and D. Thomson.: The NAME wet deposition scheme. Forecasting Research Technical Report 22
- 584, Met Office, UK, http://www.metoffice.gov.uk/archive/forecasting-research-technical-report-584, 2014. 23
- Wernsberger, B., and Schlosser, C.: Noble gas monitoring within the international monitoring system of the 24
- Comprehensive Nuclear Test-Ban Treaty, Radiation Phys. Chem., 71, 775–779, doi: 25
- 10.1016/j.radphyschem.2004.04.088, 2004. 26
- Williams, J., de Reus, M., Krejci, R., Fischer, H., and Ström, J.: Application of the variability-size 27
- relationship to atmospheric aerosol studies: estimating aerosol lifetimes and ages, Atmos. Chem. Phys., 2, 28
- 133-145, doi:10.5194/acp-2-133-2002, 2002. 29
- Wotawa, G., Becker, A., Kalinowski, M., Saey, P., Tuma, M., and Zähringer, M.: Computation and analysis 30
- 31 of the global distribution of the radioxenon isotope 133133 based on emissions from nuclear power plants

1 and radioisotope production facilities and its relevance for the verification of the Nuclear-Test-Ban treaty,

2 Pure Appl. Geophys., 167, 541–557, 2010.

- Zhang, K., O'Donnell, D., Kazil, J., Stier, P., Kinne, S., Lohmann, U., Ferrachat, S., Croft, B., Quaas, J.,
- 4 Wan, H., Rast, S., and Feichter, J.: The global aerosol-climate model ECHAM-HAM, version 2: sensitivity
- 5 to improvements in process representations, Atmos. Chem. Phys., 12, 8911-8949, doi: 10.5194/acp-12-
- 6 8911-2012, 2012a.
- Zhang, H., Jing, X., Li, J., Application and evaluation of a new radiation code under McICA scheme in
  BCC\_AGCM2.0.1, Geosci. Model Dev., 7, 737–754, doi:10.5194/gmd-7-737-2014, 2014.
- 9 Zhang, H., Wang, Zh., Wang, Za., Liu, Q., Gong, S., Zhang, X., Shen, Z., Lu, P., Wei, X., Che, H., Li, L.:
- Simulation of direct radiative forcing of aerosols and their effects on East Asian climate using an interactive
   AGCM-aerosol coupled system. Clim. Dyn. 38: 1675-1693, doi: 10.1007/s00382-011-1131-0, 2012b.
- 12 Zhang, K., Wan, H., Liu, X., Ghan, S. J., Kooperman, G. J., Ma, P.-L., Rasch, P. J., Neubauer, D., and
- 13 Lohmann, U.: Technical Note: On the use of nudging for aerosol–climate model intercomparison studies,
- 14 Atmos. Chem. Phys., 14, 8631-8645, doi:10.5194/acp-14-8631-2014, 2014.
- 15 Zhang, R., Wang, H., Qian, Y., Rasch, P. J., Easter, R. C., Ma, P.-L., Singh, B., Huang, J., and Fu, Q.:
- 16 Quantifying sources, transport, deposition, and radiative forcing of black carbon over the Himalayas and
- 17 Tibetan Plateau, Atmos. Chem. Phys., 15, 6205-6223, doi:10.5194/acp-15-6205-2015, 2015.

1 Table 1. List of models. ATM: Aerosol Transport Models (models which rely entirely on meteorological

2 input data), ACM: Aerosol Circulation Models (models which calculate their own meteorology or are

3 nudged towards (re)analysis data), LPDM: Lagrangian Particle Dispersion Models, \*AEROCOM Phase II

4 models.

	Model	Туре	Meteorology	Model output resolution H: Horizontal (degrees lat x lon) V: Vertical, T: Temporal	ReferencesKirkevåg et al. (2013), Iversen et al. (2013), Bentsen et al. (2013)Adams and Seinfeld (2002), Lee et al. (2015)Koch et al. (2006), Tsigaridis et al. (2013), Schmidt et al. (2014)Pitari et al. (2014)Pitari et al. (2014)Gong et al. (2012b), Zhang et al. (2014)Evangeliou et al. (2013)Liu et al. (2012), 	
1	NorESM	ACM	Internal (generated online)	H: 1.875°×2.5° V: 26 levels up to ~2.2 hPa. T: 3 h mean (model calc. 30 min)	Kirkevåg et al. (2013), Iversen et al. (2013), Bentsen et al. (2013)	
2	GISS-ModelE2-TOMAS*	ACM	NCEP reanalysis horizontal winds every 6 h	H: $2.0^{\circ} \times 2.5^{\circ}$ V: 40 levels up to 0.1 hPa T: 3 h mean (model calc. 30 min)	Adams and Seinfeld (2002), Lee et al. (2015)	
3	GISS-ModelE*	ACM	NCEP reanalysis horizontal winds every 6 h	H: $2.0^{\circ} \times 2.5^{\circ}$ V: 40 levels up to 0.1 hPa T: 3 h mean (model calc. 30 min)	Koch et al. (2006), Tsigaridis et al. (2013), Schmidt et al. (2014)	
4	ULAQ-CCM	ACM	Internal (generated online)	H: $5.0^{\circ} \times 6.0^{\circ}$ (T21) V: 126 levels up to mesosphere T: 45 min	Pitari et al. (2014)	
5	BCC_AGCM_2.0.1_CAM*	ACM	Online coupled. NCEP/NCAR reanalysis as initial field	H: 2.8° x 2.8° V: 26 levels up to 2.9 hPa T: 3 h (model calc. 20 min)	Gong et al. (2003), Zhang et al. (2012b), Zhang et al. (2014)	
6	LMDZORINCA	ACM	Nudged to ERA-Interim reanalysis wind fields every 6 h with a relaxation time of 10 days	H: 2.5°×1.27° V: 39 levels up to ~78 km T: 3 h mean	Evangeliou et al. (2013)	
7	CAM5*	ACM	ERA-Interim reanalysis every 6 h (re- gridded to the model grid resolution)	H: 1.9°×2.5° V: 56 levels up to 1.9 hPa T: 30 min	Liu et al. (2012), Ma et al. (2013)	
8	CAM5_PNNL*	ACM	Same as CAM5	Same as CAM5	Wang et al. (2013b)	
9	CAM5_NDG	ACM	ERA-Interim reanalysis horizontal winds (re-gridded to the model grid) at every model time step, with a relaxation time scale of 6 h	H: 1.9°×2.5° V: 30 levels up to 3.6 hPa T: 30 min	Liu et al. (2012), Zhang et al. (2014)	
10	ECHAM5-MESSy- Atmospheric Chemistry Model, v1.92 (EMAC-1)	ACM	ERA-Interim every 6 h on the model grid resolution. Nudged variables are divergence and vorticity of the wind, temperature, and the logarithm of the surface pressure	H: $1.1^{\circ} \times 1.1^{\circ}$ V: 31 levels up to 10hPa T: 3 h (model calc. 6 min)	Christoudias et al. (2013)	
11	ECHAM5-MESSy- Atmospheric Chemistry Model, v2.50 (EMAC-2)	ACM	ERA-Interim every 6 h on the model grid resolution. Nudged variables are divergence and vorticity of the wind, temperature, and the logarithm of the surface pressure	H: 1.9°x1.9° V: 31 levels up to 10hPa (~30 km) T: 1 h (model calc. 15 min)	Kunkel (2012), Kunkel et al. (2013)	
12	ECHAM5-HAM2*	ACM	Nudged towards ERA-Interim reanalysis at every time step. Relaxation time scales are 6h for vorticity, 48 h for divergence, 24 h for temperature, and 24 h for surface pressure	H: 2.8°×2.8° V: 19 levels (top layer centre at 10 hPa, including stratosphere) T: 30 min	Zhang et al. (2012a)	
13	ECHAM5-SALSA*	ACM	ERA-Interim data at 6 h intervals	H: 1.9°×1.9° V: 31 levels up to 10 hPa (~30km) T: 1 h mean (model calc. 12 min)	Bergman et al. (2012), Laakso et al. (2015)	
14	GEOS-Chem v09-01-03	ATM	GMAO GEOS-5.2.0 assimilated meteorology, 0.67x0.5 degree horizontal grid, 6 h, re-gridded to model resolution	H: 2.0°x2.5° V: 47 levels up to 0.01 hPa T: 1 h	www.geos-chem.org Croft et al. (2014), Bey et al. (2001)	
15	EEMEP v2533	ATM	ECMWF IFS cycle 36, 3 h forecasts	H: 1.0°×1.0° V: 20 levels up to 100hPa (~14 km) T: 30 min	http://www.emep.int/msc w/models.html	
16	OsloCTM2*	ATM	ECMWF IFS cycle 36, 3 h forecasts	H: 2.8°×2.8° V: 60 levels up to 0.1 hPa T: 1 h	Søvde et al. (2008), Berglen et al. (2004)	
17	OsloCTM3	ATM	ECMWF IFS cycle 36, 3 h forecasts	H: 1.1°×1.1° V: 60 levels up to 0.1 hPa T: 1 h	Søvde et al. (2012)	
18	NAME III	LPDM	Met Office Unified Model analysis, $0.35^{\circ} \times 0.23^{\circ}$ and 3 h resolution.	H: 1.0°×1.0° V: 2 km (upper level at 20 km) T: 3 h mean	Leadbetter et al. (2015), Webster and Thomson (2014)	
19	FLEXPART v9.0	LPDM	National Centers for Environmental Prediction (NCEP) Global Forecast System (GFS) analyses, 0.5°×0.5° and 3 h resolution	H: 2.0°×2.0° V: 100 m surf conc.+ total column T: 3 h mean	Stohl et al. (2005), Stohl et al. (2012)	

1 Table 2. E-folding aerosol lifetimes (Eq. 1) estimated from the decay of the ratios between the aerosol

2  $(^{137}Cs)$  and the passive tracer  $(^{133}Xe)$  surface concentrations at 11 station locations. Variations in lifetime for

different latitude regions (below and above 50°N) and time periods are also given. Mean and median

4 modelled lifetimes are calculated from the lifetimes obtained for each model and the variability is given as

5 Standard Deviation (STD) and Median Absolute Deviation (MAD, Eq. 3). The data are shown in Fig. 3.

6 95% confidence intervals and  $R^2$  statistics are given in Appendix B, Table B1. Values that are not

- 7 statistically significant are marked with a star and were not considered for the mean and median
- 8 calculations.

		e-folding lifetime									
				$\tau_e$ [days]							
		Days 15-65	Days 15-65	Days 15-65	Days 25-45	Days 45-65					
	MODEL	All stations	Stations below 50°N	Stations above 50°N	All stations	All stations					
1	NorESM	10.5	11.2	9.0	10.1	10.9					
2	GISS-ModelE2-TOMAS	9.6	9.3	10.1	9.5	10.2					
3	GISS-ModelE	8.0	7.7	8.5	7.5	8.6					
4	ULAQ-CCM	6.0	6.4	5.7	5.7	4.3					
5	BCC_AGCM	16.7	17.7	18.5	17.9	12.6					
6	LMDZORINCA	11.5	10.0	16.1	6.9	17.5					
7	CAM5	4.8	5.1	5.1	14.6*	2.8					
8	CAM5_PNNL	12.5	12.7	18.7	14.7	15.1					
9	CAM5_NDG	7.7	7.3	9.8	6.1	6.2					
10	EMAC-1	11.7	13.9	8.3	8.7	8.1					
11	EMAC-2	8.1	8.1	7.8	19.9*	3.9					
12	ECHAM-HAM2	6.4	6.3	8.8	5.7	6.2					
13	ECHAM5-SALSA	6.9	7.5	7.8	8.6	34.0*					
14	GEOS-Chem	20.4	16.6	33.0	18.6	23.3					
15	EEMEP	26.7	20.2	50.0*	15.6	23.6*					
16	OSLO-CTM2	11.2	11.1	19.9	10.8	8.3					
17	OSLO-CTM3	8.8	9.1	9.3	13.0	7.8					
18	NAME	9.4	11.0	8.7	7.3	4.5					
19	FLEXPART	5.8	5.9	5.9	6.5	5.0					
	MODEL MEAN ± STD	$10.7 \pm 5.4$	$10.4 \pm 4.2$	11.7±7.0	10.1±4.3	9.1±5.5					
	MODEL MEDIAN $\pm$ MAD	9.4±2.3	9.3±2.0	8.9±1.2	8.7±2.2	8.1±3.1					
	OBSERVATIONS	14.3	13.5	15.0	15.0	14.7					

- 1 Table 3. Median instantaneous lifetimes calculated from the ratios of the aerosol (<sup>137</sup>Cs) to the passive tracer
- 2 (<sup>133</sup>Xe) global burdens during different time periods (weeks) after the start of the emissions (11 March
- 3 2011). The variability is given as the Median Absolute Deviation (MAD, Eq. 3) for the individual models, as
- 4 well as for the observations and the model median, while the Standard deviation (STD) is given of the model
- 5 mean. These variabilities are calculated from the instantaneous lifetimes over each of the time periods. The
- 6 data are shown in Fig. 7.

		Instantaneous lifetime										
				$\tau_{inst}$ [days]								
	MODEL	Week 1 (11-18 Mar; days 1-7)	Weeks 2-3 (18 Mar–1 Apr; days 7-21)	Weeks 4-5 (1-15 Apr; days 21-35)	Weeks 6-7 (15-29 Apr; days 35-49)	Weeks 8-9 (29 Apr-13 May; days 49-63)						
1	NorESM	$1.6 \pm 0.4$	$6.8 \pm 2.4$	$8.7 \pm 2.8$	$6.9\pm2.4$	9.1 ± 0.3						
2	GISS-ModelE2-TOMAS	$2.5\pm0.8$	$10.1\pm3.6$	$9.5\pm3.6$	$7.4 \pm 1.5$	$9.9\pm0.8$						
3	GISS-ModelE	$2.2 \pm 1.3$	$8.7\pm5.9$	$7.9\pm3.5$	$5.5\pm0.8$	$7.9\pm0.3$						
4	ULAQ-CCM	$2.4\pm1.5$	$5.7\pm1.0$	$10.0\pm2.5$	$8.5\pm1.4$	$11.2\pm1.3$						
5	BCC_AGCM	$2.8\pm2.0$	$9.8\pm2.3$	$9.5\pm1.7$	$9.7\pm0.8$	$11.6\pm0.4$						
6	LMDZORINCA	$2.5\pm2.1$	$11.3\pm9.0$	$8.2\pm3.6$	$8.6\pm3.6$	$19.0\pm1.3$						
7	CAM5	$3.6\pm2.6$	$3.6\pm1.7$	$2.6\pm1.3$	$2.1\pm0.8$	$2.8\pm0.3$						
8	CAM5_PNNL	$2.5\pm0.8$	$11.7\pm5.4$	$10.3\pm5.2$	$11.0\pm3.5$	$17.3\pm1.1$						
9	CAM5_NDG	$2.1\pm0.7$	$6.0\pm1.6$	$7.9\pm4.1$	$6.7\pm2.6$	$12.8\pm1.4$						
10	EMAC-1	$2.4\pm0.7$	$6.8\pm4.1$	$8.3\pm6.1$	$5.5\pm0.8$	$6.9\pm0.7$						
11	EMAC-2	$1.7 \pm 1.2$	$4.4\pm2.2$	$4.4\pm3.1$	$3.0\pm0.4$	$4.2\pm0.2$						
12	ECHAM-HAM2	$2.8 \pm 1.5$	$6.9\pm4.8$	$6.4\pm3.1$	$5.1 \pm 1.9$	$10.4\pm0.7$						
13	ECHAM5-SALSA	$2.2\pm0.9$	$4.1\pm0.4$	$5.3\pm1.6$	$10.4\pm1.0$	$14.4\pm2.4$						
14	GEOS-Chem	$1.4\pm0.3$	$10.7\pm8.4$	$11.8\pm5.0$	$11.6\pm2.1$	$16.4\pm0.9$						
15	EEMEP	$1.0\pm0.4$	$1.4\pm0.5$	$2.7\pm1.8$	$3.0\pm1.6$	$8.1\pm1.9$						
16	OSLO-CTM2	$1.3\pm0.5$	$3.7\pm2.2$	$4.6\pm3.0$	$3.9\pm1.8$	$18.1\pm1.9$						
17	OSLO-CTM3	$1.5\pm0.8$	$11.9\pm6.7$	$6.3\pm3.7$	$6.0\pm1.4$	$10.4\pm2.0$						
18	NAME	$0.8\pm0.3$	$1.8\pm0.7$	$2.2\pm1.4$	$3.2\pm1.2$	$9.2\pm3.6$						
19	FLEXPART	$1.5\pm0.2$	$6.6\pm4.1$	$3.7\pm1.7$	$3.7\pm1.4$	$6.3\pm0.4$						
	MODEL MEAN $\pm$ STD	$2.0\pm0.7$	$6.4\pm3.6$	$6.9\pm2.9$	$6.4\pm2.9$	$10.8\pm4.6$						
	MODEL MEDIAN $\pm$ MAD	$2.2\pm0.6$	$6.6\pm2.9$	$7.9\pm2.1$	$6.0\pm2.5$	$10.4\pm2.5$						
	OBSERVATIONS + box model	N/A	N/A	$10.9\pm4.9$	9.3 ± 1.2	$9.6\pm0.7$						

1 Table 4. Characteristic timescales  $\tau_1$ ,  $\tau_2$  and  $\tau_3$  estimated with a convolution approach (Eq. 4) based on the

2 different models' aerosol (<sup>137</sup>Cs) burdens, as well as the mean lifetime  $\tau_m$  (Eq. 5), and the instantaneous

3 lifetime  $\tau_{inst}$  over weeks 8-9 from Fig. 7 and Table 3, and the e-folding lifetime  $\tau e$  from Fig. 3 and Table 2.

4 Values in brackets for  $\tau_3$  indicate high uncertainty and are discarded in the mean and median calculations.

		$ au_1$	$ au_2$	$\tau_3$	$ au_{ m m}$	$ au_{ ext{inst}}$	$ au_{ m e}$
	MODEL					Week 8-9	Days 15-65,
	MODEL					(Table 3)	(Table 2)
1	NorESM	1.8	10.4	192.9	2.0	9.1	10.5
2	GISS-ModelE2-	4.1	11.4	[314.8]	4.4	9.9	9.6
3	GISS-ModelE	3.8	8.5	116.8	4.1	7.9	8.0
4	ULAQ-CCM	7.1	18.8	[200.0]	7.4	11.2	6.0
5	BCC_AGCM	1.1	11.0	[200.2]	2.3	11.6	16.7
6	LMDZORINCA	2.5	20.5	[200.5]	2.6	19.0	11.5
7	CAM5	1.1	3.0	[ 11.9]	1.1	2.8	4.8
8	CAM5_PNNL	1.7	15.9	[200.2]	2.2	17.3	12.5
9	CAM5_NDG	2.8	14.3	[206.2]	2.9	12.8	7.7
10	EMAC-1	1.8	7.0	[200.1]	2.2	6.9	11.7
11	EMAC-2	1.1	4.4	[ 38.2]	1.2	4.2	8.1
12	ECHAM-HAM2	2.2	18.6	[224.4]	2.2	10.4	6.4
13	ECHAM5-SALSA	2.4	19.2	[472.3]	2.4	14.4	6.9
14	GEOS-Chem	1.0	15.2	[200.1]	1.7	16.4	20.4
15	EEMEP	0.6	8.2	[297.1]	0.7	8.1	26.7
16	OSLO-CTM2	2.3	18.9	[203.0]	2.4	18.1	11.2
17	OSLO-CTM3	4.5	16.3	[200.1]	4.6	10.4	8.8
18	NAME	0.9	6.5	[309.7]	1.0	9.2	9.4
19	FLEXPART	2.0	6.8	[183.9]	2.1	6.3	5.8
	MODEL MEAN $\pm$ STD	2.4±1.6	12.4±5.6	154.9±53.8	2.6±1.6	$10.8 \pm 4.6$	$10.7 \pm 5.4$
	MODEL MEDIAN $\pm$	$2.0\pm0.9$	11.4±4.6	$154.8 \pm 38.1$	$2.2 \pm 0.5$	$10.4 \pm 2.5$	$9.4 \pm 2.3$

5

- 1 Table 5. Same as Table 2, but including two revised simulations with the GEOS-Chem model; GEOS-
- 2 Chem\_allT and GEOS-Chem\_T258 (see text and Fig. 12 for description of simulations).

			e	-folding lifetim	e	
				$\tau_e$ [days]		
		Days 15-65	Days 15-65	Days 15-65	Days 25-45	Days 45-65
	MODEL	All stations	Stations below 50°N	Stations above 50°N	All stations	All stations
1	GEOS-Chem_Orig	20.8	17.7	30.8	19.1	21.8
2	GEOS-Chem_T258	19.7	16.2	32.2	13.1	19.5
3	GEOS-Chem_allT	13.7	12.0	17.2	10.6	13.2
	OBSERVATIONS	14.3	13.5	15.0	15.0	14.7

## 1 Table A1. Specification of the Aerosol Modules

	Model	Aerosol module
		size distribution, density, microphysics (external/internal mixing, nucleation, condensation, coagulation), dry and wet removal (in-cloud, below-cloud)
1	NorESM	Cesium was modelled as accumulation mode sulphate, with median radius of $0.1 \times 10^{-6}$ m, and a density of 1769 kgm <sup>-3</sup> . The aerosol module (Seland et al., 2008; Kirkevåg et al., 2013) calculates mass-concentrations of aerosol species that are tagged according to production mechanisms in clear and cloudy air and four size- classes (nucleation, aitken, accumulation, and coarse modes). These processes are primary emission, gaseous and aqueous chemistry (cloud processing), nucleation, condensation, and coagulation. Loss terms are dry deposition, in-cloud and below-cloud scavenging. The chemical components included are sulphate (SO4), black carbon (BC), organic matter (OM), sea-salt (SS), and mineral dust (DU). This adds up to 20 aerosol components in addition to two gaseous precursors (SO2 and dimethyl sulphide, DMS). Dry and wet deposition for aerosols: (i) Hygroscopic growth of particles is included. (ii) The dry deposition velocity depends on particles size, and the relative humidity influences this dependence for hygroscopic particles. (iii) Gravitational settling is included for coarse particles. (iv) Wet deposition: both in-cloud and below-cloud scavenging is taken into account for both stratiform and convective clouds.
2	GISS-ModelE2-TOMAS	TOMAS (TwO-Moment Aerosol Sectional) microphysics model (Adams and Seinfeld, 2002; Lee et al. 2015) has 12 bins, covering from 10 nm to 10 $\mu$ m in a dry particle diameter: 10 bins from 10 nm to 1 $\mu$ m and 2 bins from 1 $\mu$ m to 10 $\mu$ m. We assumed model cesium to follow a size distribution of sulphate particles. Cesium and sulphate emissions are assumed to have a bimodal log-normal distribution: 5% of emissions as a nucleation mode with geometric number mean diameter (GSD) of 10 nm and a geometric number mean diameter (GSD) of 1.6 and 95% as an Aitken mode with GMD of 70 nm and GSD of 2. Coagulation and condensation are used, but aerosol nucleation is turned off to avoid re-partitioning cesium mass to smaller size. Density of sulphate is assumed to be 1.78 g cm <sup>-3</sup> . Internal mixing assumption except for wet deposition. Dry deposition velocity is computed based on a resistance-in-series approach with size-resolved resistance in the quasi-laminar sublayer and size-resolved gravitational settling velocity. Wet deposition: a modified Köhler theory is used for in-cloud scavenging, and a size-resolved first-order removal scheme for below-cloud scavenging.
3	GISS-ModelE	Using the mass based aerosol scheme OMA, cesium was modelled as accumulation mode sulphate. The model transports sulphate as a mass tracer with a constant mean dry radius of $0.3 \times 10^{-6}$ m. The wet deposition includes in and below cloud scavenging, which depends on the cloud cover, the hygroscopicity of aerosols, sulphate aerosols are assumed to be fully hygroscopic, and the precipitation and evaporation rates. Convective and large-scale clouds are treated separately, with convective clouds handling individual convective plumes.
4	ULAQ-CCM	Size distribution explicitly predicted from a microphysical code with gas-particle interactions (external mixing, homogeneous and heterogeneous nucleation, condensation/evaporation, coagulation, gravitational settling, large-scale transport), dry and wet removal (in-cloud, below-cloud).
5	BCC_AGCM_2.0.1_CAM	Canadian Aerosol Module (CAM). The size spectrum of sulphate was discretized into 12 bins from 0.005 to 20.48 µm. Density of sulphate is about 1.8 g/cm <sup>3</sup> . Internal mixing except for the freshly emitted insoluble components. Coagulation: only binary collisions of particles are considered, coagulation coefficient is calculated from contributions of Brownian, turbulence and gravitational settling movements. No Nucleation and Condensation in order to avoid false source of cesium. Dry deposition: a dry deposition velocity for seasalt on the ocean was modified to treat multi-component aerosols and complicate land properties. Wet deposition: includes in and below cloud scavenging, which depend on cloud cover, the properties of both aerosols and hydrometeors, precipitation and evaporation rates.
6	LMDZORINCA	INCA (INteraction with Chemistry and Aerosols) model. Cesium was treated as a submicronic (diameters < 1um) aerosol in accumulation mode following a lognormal particle size distribution. No microphysics. The dry deposition of cesium was computed using the analogy of surface resistance. The deposition velocity is defined as the inverse of the sum of an aerodynamic resistance and a surface resistance placed in series. Wet scavenging for large-scale stratiform precipitation is calculated adopting the falling raindrop approach. Scavenging by convective precipitation is calculated as part of the upward convective mass flux.
7	CAM5	Cesium was assumed to attach to particles in the accumulation mode that includes internally mixed sulphate, black carbon, primary and secondary organic matter, mineral dust, and sea salt. Particles in this mode are described by a lognormal size distribution with a prescribed geometric standard deviation of 1.8 and a dry diameter size range of 58-270 nm. The evolution of aerosol particles is controlled by various processes including emission, transport, aerosol microphysics (e.g., nucleation, condensation, and coagulation), cloud chemistry, and dry/wet removal processes (Liu et al., 2012). Aerosol wet removal is parameterized separately for stratiform and convective clouds by in-cloud and below-cloud processes. For stratiform clouds, the in-cloud removal process involves explicit aerosol activation to form cloud droplets and subsequent removal of cloud-borne aerosols due to the conversion of cloud droplets to precipitation. Aerosol dry deposition velocities are calculated with model provided aerodynamic resistance, friction velocity, and surface properties. Gravitational settling is also treated.
8	CAM5_PNNL	Cesium is treated in the same way as in the CAM5 model. However, a new unified treatment of vertical transport and in-cloud wet removal processes in convective clouds is applied, which has a more detailed treatment of aerosol activation in convective updrafts. Also, a mechanism is added for laterally entrained aerosols to be activated and then removed. In addition, a few other changes have been introduced to wet removal processes to reduce some known biases in the remote aerosol distributions predicted by the default CAM5 model (Wang et al. 2013b).
9	CAM5_NDG	Same as CAM5

	Model	Aerosol module size distribution density microphysics (external/internal mixing nucleation condensation coagulation) dry and wet
		removal (in-cloud, below-cloud)
10	ECHAM5-MESSy-	Cesium is modelled as a water-soluble aerosol with a standard lognormal distribution with mean radius 0.25
	Atmospheric Chemistry	$\mu$ m and a Henry's law coefficient equal to 1.0 mol L <sup>-1</sup> atm <sup>-1</sup> and a density of 1000.0 kg m <sup>-3</sup> . Scavenging by
	Model v1 92 (EMAC-1)	turbulent transfer and uptake processes onto the earth's surface. The denosition velocity is calculated using a
		resistance model. Sedimentation: Calculated separately based on the mass of the aerosol particles.
11	ECHAM5-MESSy-	Cesium is treated as quasi-passive aerosol particles with sulphate characteristics, i.e. molar mass of 96 g/mol
	Atmospheric Chemistry	and a density of 2 g/cm <sup>3</sup> . The aerosols have an initial diameter of 0.5 $\mu$ m but can grow due to coagulation,
	Model x2 50 (EMAC 2)	coating and condensation, and are internally mixed. Thus, the aerosols are affected by the background but do not interact with the background. Acrossl microphysics are calculated with CMXs (Pringle et al. 2010).
	Wodel, $V2.30$ (EWAC-2)	Aerosol removal: Scavenging by impaction (below cloud) and nucleation (in-cloud), depending on aerosol
		size and solubility and rain intensity and droplet size. Dry deposition: removal by turbulent transfer and
		uptake processes onto the earth's surface. Dry deposition for aerosols follows the big-leaf approach. The
		deposition velocity is calculated using a resistance model. Sedimentation: Calculated separately based on the
10		mass of the aerosol particles (Kerkweg et al., 2006).
12	ECHAM5-HAM2	(2012a). Cesium is assumed to attach to accumulation-mode aerosols. In-cloud and below-cloud wet
		scavenging of cesium is considered for both stratiform and convective clouds. Dry deposition is also
		considered, but gravitational sedimentation is neglected for cesium.
13	ECHAM5-SALSA	SALSA describes aerosol size distribution for diameters ranging from 3nm to 10um using 17 internally and
		externally mixed sections. The aerosol microphysical processes considered include condensation,
		as well as sedimentation (Bergman et al., 2012, Laakso et al., 2015). Wet deposition is based on the size and
		composition dependent scavenging parameter, rain intensity and type of precipitation, which are used to In-
		cloud and below-cloud scavenging separately (for the coefficients see Bergman et al. (2012)). Dry
		deposition uses a big leaf method, which resolves aerosol deposition to surfaces on-line. Aerosol deposition
		velocity is calculated using aerosol number and mass together with wet radius, density, turbulence and surface cover (more detailed explanation can be found in Stier et al. (2005) and Kerkweg et al. (2006)
		Gravitational sedimentation is calculated using the Stokes law. Cesium is assumed to have density of 1.83
		g/cm3. It attaches to the particles immediately and thus it is allocated to model size bins with diameters
		ranging from 50nm to 10um using a lognormal distribution with a mean diameter of 150nm and standard deviation of 1.59
14	GEOS-Chem v09-01-03	Cesium is treated as accumulation mode sulphate. No aerosol microphysics. Dry Deposition: Resistance in
11		series scheme with dry deposition velocity dependent on surface type, as described in Wang et al. (2011).
		Wet Deposition: Separate treatment for large-scale and convective clouds (Liu et al., 2001). Below -cloud
		scavenging: Depends on precipitation rate, separate scavenging coefficients for accumulation and coarse
		mode aerosols, separate treatment for rain and snow. In-cloud scavenging: Depends on aerosol
15	EEMED 2522	Casium treated as aerosal particles ( <b>PPM2 5</b> ) with no sedimentation. No microphysics. Wet deposition
15	EEMEP V2533	depends on precipitation intensity and species-specific scavenging coefficients. Different coefficients are
		specified for in-cloud (rain out) and below-cloud scavenging. Dry deposition: impaction of particles on
		surface roughness elements and turbulent diffusion to surface treated using the resistance method to estimate
		a deposition velocity.
16	OsloCTM2	precipitation Removed in both convective and large-scale precipitation. Dry deposition is not included
17	OsloCTM3	Cesium is treated as sulphate, following Berglen et al. (2004), but large-scale wet scavenging follows Neu et
17	Obloc This	al. (2012), while convective wet scavenging uses the CTM2 approach. Both are described by Søvde et al.
		(2012). Dry deposition is not included.
18	NAME III	Cesium is treated as aerosol particles with no sedimentation. No microphysics. Wet deposition depends on
		in-cloud (rain out) and below-cloud (wash out) and rain and snow (Webster and Thomson 2014)) Dry
		deposition: impaction of particles on surface roughness elements and turbulent diffusion to surface treated
		using the resistance method to estimate a deposition velocity.
19	FLEXPART v9.0	Cesium is treated as aerosol particles with mean particle diameter 0.4 µm and density 1.9 g/m <sup>3</sup> . No
		microphysics. Wet deposition depends on precipitation intensity and determined by species-specific scavenging coefficients. In cloud (rain out) and below cloud (wash out) scavenging are tracted differently
		as well as large-scale and convective scale precipitation. Dry deposition: impaction of particles on surface
		roughness elements and turbulent diffusion to the surface treated using the resistance method to estimate a
		deposition velocity, which also includes a sedimentation velocity dependent on the particle size.
1		

- 1 Table B1. Confidence intervals [days] for the e-folding aerosol lifetimes in Table 2, and the coefficients of
- 2 determination  $R^2$ . Values that are not statistically significant at the 99.9% level are marked with an asterisk.

		Statistics for e-folding lifetime										
						$ au_{ m e}$ [c	lays]					
		Days 15-	-65	Days 15	-65	Days 15	5-65	Days 25	5-45	Days 45	5-65	
	MODEL	All stations		Stations below 50°N		Stations above 50°N		All stations		All stations		
		C.I.	$\mathbb{R}^2$	C.I.	$\mathbb{R}^2$	C.I.	$\mathbb{R}^2$	C.I.	$\mathbb{R}^2$	C.I.	$\mathbb{R}^2$	
1	NorESM	[9.8,11.3]	0.94	[10.3,12.4]	0.91	[8.5,9.6]	0.96	[8.3,13.0]	0.83	[8.5,15.0]	0.77	
2	GISS-ModelE2-TOMAS	[9.2,10.0]	0.98	[8.9,9.8]	0.97	[9.4,10.9]	0.94	[8.2,11.3]	0.91	[8.5,12.7]	0.86	
3	GISS-ModelE	[7.6,8.4]	0.97	[7.3,8.2]	0.96	[7.7,9.5]	0.89	[6.5,8.8]	0.92	[6.9,11.5]	0.79	
4	ULAQ-CCM	[5.7,6.3]	0.97	[5.8,7.1]	0.90	[5.4,6.0]	0.97	[5.2,6.4]	0.96	[3.9,4.7]	0.96	
5	BCC_AGCM	[15.6,18.0]	0.94	[16.1,19.7]	0.90	[16.8,20.4]	0.90	[14.0,24.7]	0.76	[10.8,15.0]	0.90	
6	LMDZORINCA	[10.5,12.8]	0.89	[9.2,11.1]	0.91	[12.9,21.5]	0.57	[6.0,8.2]	0.91	[10.2,61.8]	0.32	
7	CAM5	[4.4,5.4]	0.88	[4.5,5.9]	0.84	[4.4,6.0]	0.77	[8.3,60.2]	0.30*	[2.4,3.6]	0.86	
8	CAM5_PNNL	[11.5,13.5]	0.93	[11.7,13.8]	0.93	[15.3,24.0]	0.63	[11.2,21.3]	0.71	[11.0,24.4]	0.63	
9	CAM5_NDG	[7.2,8.2]	0.95	[6.8,7.9]	0.94	[8.1,12.5]	0.64	[5.2,7.3]	0.89	[5.2,7.6]	0.88	
10	EMAC-1	[10.7,12.8]	0.91	[12.1,16.3]	0.80	[7.4,9.6]	0.84	[7.6,10.3]	0.91	[6.6,10.4]	0.83	
11	EMAC-2	[7.0,9.6]	0.78	[6.9,9.8]	0.74	[6.6,9.6]	0.70	[11.1,98.8]	0.28*	[3.2,5.1]	0.82	
12	ECHAM-HAM2	[5.9,7.0]	0.92	[5.7,7.0]	0.90	[7.4,10.7]	0.72	[4.7,7.3]	0.84	[4.5,10.0]	0.63	
13	ECHAM5-SALSA	[6.1,8.0]	0.81	[6.4,8.9]	0.77	[6.4,9.9]	0.64	[6.4,12.9]	0.68	[11.4,-34.2]	0.06*	
14	GEOS-Chem	[18.1,23.3]	0.84	[14.8,18.9]	0.86	[25.6,46.4]	0.50	[13.8,28.4]	0.67	[14.6,57.5]	0.41	
15	EEMEP	[20.1,39.7]	0.44	[15.1,30.5]	0.44	[25.5,1233]	0.08*	[9.3,50.1]	0.34	[10.2,-75.2]	0.12*	
16	OSLO-CTM2	[9.9,12.9]	0.84	[9.6,13.1]	0.79	[13.9,35.4]	0.31	[6.9,24.7]	0.44	[6.0,13.8]	0.61	
17	OSLO-CTM3	[8.1,9.5]	0.94	[8.2,10.4]	0.86	[8.1,10.8]	0.80	[10.5,17.0]	0.81	[6.1,10.8]	0.76	
18	NAME	[7.7,11.9]	0.65	[8.9,14.4]	0.61	[6.6,12.6]	0.46	[4.5,19.6]	0.39	[3.2,7.9]	0.57	
19	FLEXPART	[5.5,6.1]	0.97	[5.5,6.3]	0.94	[5.5,6.5]	0.92	[5.6,7.7]	0.91	[4.2,6.4]	0.85	
	OBSERVATIONS	[13.1,15.7]	0.91	[12.2,15.2]	0.88	[12.9,17.7]	0.77	[11.7,20.9]	0.76	[10.1,26.8]	0.54	



Figure 1. Measurement station network. CTBTO stations (red markers) measuring particulates (<sup>137</sup>Cs) and
 noble gases (<sup>133</sup>Xe). The position of the Fukushima Dai-Ichi nuclear power plant (FD-NPP) is shown by a
 green marker.







1

Figure 3. Time-series of measured and modelled ratios of the aerosol ( $^{137}Cs$ ) to the passive tracer ( $^{133}Xe$ ) surface concentrations at 11 CTBTO station locations.  $\omega$  values (median, black triangles) represent the daily median ratios (median concentration for each day over all stations). Fits of exponential decay models to the  $\omega$  data are shown as grey lines with e-folding time scales  $\tau_e$  (Eq. 1) as indicated. The fits are made over days 15 to 65 for which data exist from at least five stations each day.



2 Figure 3. Continued.

	1
	L
-	L

	Station (latitude)/	Wak	Oah	۵ch	Cha	llss	St I	Sha	Illa	Sto	Vel	Sni	ΔΠ
	Model	(19°N)	(22°N)	(37°N)	(38°N)	(44°N)	(48°N)	(48°N)	(48°N)	(59°N)	(62°N)	(78°N)	STATIONS
1	NorESM	1.58	1.22	0.76	0.80	0.54	0.47	1.19	0.61	0.47	0.48	0.58	0.61
2	GISS-ModelE2-TOMAS	0.49	0.92	1.00	1.29	1.24	1.14	2.12	1.01	1.30	0.94	1.27	1.16
3	GISS-ModelE	0.51	1.02	1.20	1.33	1.24	1.13	2.11	1.03	1.29	0.98	1.27	1.17
4	ULAQ-CCM	13.09	4.14	0.73	0.46	0.80	0.43	0.60	0.56	0.37	0.66	0.58	0.61
5	BCC_AGCM2.0.1_CAM	1.49	1.72	0.93	0.91	0.70	0.65	1.54	0.64	0.65	0.57	0.68	0.72
6	LMDZORINCA	1.60	2.04	1.69	1.61	1.08	0.97	2.08	1.11	0.90	0.90	1.09	1.15
7	CAM5	0.70	1.10	0.91	0.96	0.85	0.83	1.37	0.80	0.88	0.71	0.90	0.86
8	CAM5_PNNL	0.68	1.07	0.91	0.97	0.85	0.84	1.35	0.81	0.88	0.71	0.90	0.86
9	CAM5_NDG	1.72	0.72	0.48	0.73	0.60	0.55	0.91	0.50	0.59	0.40	0.57	0.58
10	EMAC-1	0.23	0.30	0.35	0.33	0.94	0.66	1.18	0.76	0.85	0.60	0.93	0.70
11	EMAC-2	1.20	1.22	0.39	0.35	0.87	0.69	0.37	0.62	0.46	0.59	0.84	0.70
12	ECHAM-HAM2	0.65	0.90	0.68	0.71	0.68	0.65	1.18	0.65	0.69	0.55	0.72	0.68
13	ECHAM_salsa	0.71	0.78	0.04	0.21	0.29	0.20	0.12	0.15	0.07	0.07	0.19	0.15
14	GEOS-Chem	1.00	1.08	0.88	0.84	0.83	0.81	1.51	0.74	0.88	0.72	0.93	0.87
15	EEMEP	2.13	2.15	1.45	1.70	1.45	1.30	2.36	1.33	1.51	1.21	1.52	1.48
16	OsloCTM2	1.20	1.07	0.70	0.68	0.68	0.64	1.41	0.70	0.66	0.55	0.90	0.76
17	OsloCTM3	1.05	1.24	0.79	0.82	0.72	0.63	1.49	0.77	0.67	0.60	0.76	0.77
18	NAME	1.13	0.88	0.60	0.73	0.66	0.46	0.76	0.50	0.38	0.31	0.17	0.51
19	FLEXPART	1.17	1.35	0.81	0.93	0.95	1.03	1.91	1.03	0.91	0.72	1.62	1.03
	MODEL MEDIAN	1.05	1.08	0.77	0.81	0.77	0.67	1.46	0.75	0.71	0.63	0.85	0.79

Figure 4. Median ratio of the modelled to measured passive tracer  $(^{133}Xe)$  surface concentrations, for each

4 model and each station as well as the model median and over all stations. Values greater than 1.2

5 (substantial over-predictions) are shown in red, and values less than 0.8 (substantial under-predictions) in

6 green.

	Station (latitude)/	Wak.	Oah.	Ash.	Cha.	Uss.	St.J	Sha.	Ula.	Sto.	Yel.	Spi.	ALL
	Model	(19°N)	(22°N)	(37°N)	(38°N)	(44°N)	(48°N)	(48°N)	(48°N)	(59°N)	(62°N)	(78°N)	STATIONS
1	NorESM	0.34	1.04	5.42	2.00	0.97	0.51	1.13	1.70	0.61	0.60	1.28	1.13
2	GISS-ModelE2-TOMAS	6.18	9.04	20.45	13.35	7.45	11.44	13.65	11.30	12.65	5.50	10.14	10.17
3	GISS-ModelE	1.10	3.61	4.09	3.32	2.10	3.64	2.48	1.82	2.14	0.90	1.53	2.06
4	ULAQ-CCM	60.73	3.60	2.20	0.41	367.45	0.08	0.19	46.43	0.39	0.19	0.80	0.80
5	BCC_AGCM2.0.1_CAM	9.42	18.99	18.74	8.03	6.03	4.03	10.46	5.29	3.41	1.31	3.14	5.50
6	LMDZORINCA	0.00	2.17	2.52	0.99	0.24	0.55	1.27	0.80	0.30	0.34	0.09	0.56
7	CAM5	0.95	0.54	0.08	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
8	CAM5_PNNL	15.71	12.89	6.04	3.67	2.32	1.81	5.89	5.26	2.78	0.93	1.13	3.16
9	CAM5_NDG	10.72	3.37	2.69	1.76	0.19	0.29	0.91	0.88	0.60	0.16	0.07	0.69
10	EMAC-1	0.01	0.11	0.29	0.06	0.32	0.34	0.67	0.58	0.71	0.14	0.51	0.33
11	EMAC-2	0.02	0.59	0.51	0.11	0.18	0.04	0.04	0.03	0.02	0.03	0.02	0.05
12	ECHAM-HAM2	5.43	1.80	0.87	0.28	0.11	0.02	0.20	0.15	0.03	0.02	0.00	0.11
13	ECHAM_salsa	2.51	2.47	0.38	0.67	0.07	0.24	0.17	0.12	0.02	0.05	0.01	0.15
14	GEOS-Chem	0.14	4.14	7.52	6.32	4.58	6.49	8.74	7.50	5.55	3.25	6.89	5.53
15	EEMEP	0.00	0.05	0.03	0.02	0.01	0.01	0.02	0.02	0.01	0.01	0.01	0.01
16	OsloCTM2	0.06	0.21	0.30	0.06	0.01	0.01	0.05	0.01	0.01	0.00	0.00	0.02
17	OsloCTM3	0.69	2.73	4.96	2.62	0.63	0.51	1.28	1.19	0.39	0.34	0.08	0.76
18	NAME	0.03	0.13	0.02	0.03	0.01	0.01	0.00	0.01	0.01	0.00	0.00	0.01
19	FLEXPART	2.54	2.81	1.00	0.64	0.21	0.19	0.57	0.42	0.46	0.03	0.17	0.35
	MODEL MEDIAN	0.91	1.67	1.99	0.64	0.41	0.19	0.74	0.78	0.32	0.15	0.13	0.49

3 Figure 5. Same as Fig. 4 but for aerosol  $(^{137}Cs)$  surface concentrations.



1

Figure 6. Ratio of modelled to observed aerosol (green) and passive tracer (blue) surface concentrations as a function of the latitude of the measurement station and time; values at each measurement station are the median values over 14 day time periods (11-24 March, 25 March-7 April, 8-21 April, 22 April-5 May, 6-19 May, 20 May-2 June, 3-17 June 2011). The size of the circles indicate the time; the larger the circle the later the time period. Values above the dotted 1-line are over-predicted by the model and values below one are under-predictions. Notice different scales on the y-axes.







Figure 7. Instantaneous lifetimes calculated from the ratios of the aerosol ( $^{137}$ Cs) to the passive tracer ( $^{133}$ Xe) global burdens. The median instantaneous lifetimes  $\tau_{inst}$  over days 49-63 (weeks 8-9; 29 April-13 May; shaded area) are indicated, and represent the time period when the ratios have reached a quasi-steady state and influence from transport into the stratosphere is less significant than for later times. The measurement + box model refers to the estimate from Kristiansen et al. (2012) for the burdens using the station measurements and a 1D-box model (see Appendix C and Fig. C1).







- 1
- 2 Figure 8. Examplified evolution of the aerosol burden after a unit emission, with characteristic time scales
- 3 indicated by  $\tau_1$ ,  $\tau_2$  and  $\tau_3$ .



Figure 9. Source emissions of <sup>137</sup>Cs and modelled total atmospheric aerosol (<sup>137</sup>Cs) burdens, for the initial
three weeks after the first release.



2 Figure 10. a) Modelled instantaneous lifetimes averaged over weeks 1-3 (see Fig. 7) versus aerosol

- 3 concentration bias (modelled to measured aerosol  $(^{137}Cs)$  station concentrations, see Fig. 5). The vertically
- 4 dotted line is the 1-line (perfect match between modelled and observed aerosol concentrations), b)
- 5 Correlation of instantaneous lifetimes at different time periods; modelled instantaneous lifetime averaged
- 6 over weeks 1-3 versus weeks 4-6 (see Fig. 7).



2 Figure 11. Modelled total aerosol (<sup>137</sup>Cs) mass as a function of latitude for four models (FLEXPART,

3 OsloCTM3, NorESM and EEMEP), in four different height layers (surface, 0-5 km, 5-10 km and > 10 km)

- 4 and at three different time steps (14, 28 and 41 days after the first emissions). Note the different vertical
  - 5 axes.





2 Figure 12. Same as Fig. 6, but including two revised simulations with the GEOS-Chem model. GEOS-

- 3 Chem\_Orig is the same simulation as in Fig. 6, GEOS-Chem\_allT has an updated scavenging scheme
- 4 relative to GEOS-Chem\_Orig through processes related to cloud liquid and ice water content, cloud
- 5 temperature and using cloud fraction from the assimilated meteorological field. GEOS-Chem\_T258 includes
- 6 the same revisions, but restricts the large-scale in-cloud scavenging of aerosols to temperatures above 258K,
- 7 same restriction for simulation GEOS-Chem\_Orig.



7

time scales  $\tau_e$  (Eq. 1) as indicated.

Figure C1. Time series of global atmospheric burdens of aerosol (<sup>137</sup>Cs; green lines), passive tracer (<sup>133</sup>Xe; blue and turquoise lines) and their ratio (grey lines). Dark green and dark blue lines represent global burdens estimated by applying a 1D-box model to the measured or model simulated station data. Bright green and turquoise lines show the modelled global atmospheric burdens calculated from the full 3D model output. Fits of exponential decay models to the ratios between day 45 and 65 are shown as black lines, with e-folding



2 Figure C1. Continued.