

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Stratospheric ozone change and related climate impacts over 1850–2100 as modelled by the ACCMIP ensemble

F. Iglesias-Suarez, P. J. Young, and O. Wild

Lancaster Environment Centre, Lancaster University, Lancaster, UK

Received: 3 July 2015 – Accepted: 20 August 2015 – Published: 15 September 2015

Correspondence to: F. Iglesias-Suarez (n.iglesiassuarez@lancaster.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Discussion Paper | Discussion Paper

Stratospheric ozone change and related climate impacts over 1850–2100

F. Iglesias-Suarez et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Stratospheric ozone and associated climate impacts in the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP) simulations are evaluated in the recent past (1980–2000), and examined in the long-term (1850–2100) using the Representative Concentration Pathways low and high emission scenarios (RCP2.6 and RCP8.5, respectively) for the period 2000–2100. ACCMIP multi-model mean total column ozone (TCO) trends compare favourably, within uncertainty estimates, against observations. Particularly good agreement is seen in the Antarctic austral spring ($-11.9\% \text{dec}^{-1}$ compared to observed $\sim -13.8 \pm 11\% \text{dec}^{-1}$), although larger deviations are found in the Arctic's boreal spring ($-2.1\% \text{dec}^{-1}$ compared to observed $\sim -5.3 \pm 3\% \text{dec}^{-1}$). The simulated ozone hole has cooled the lower stratosphere during austral spring in the last few decades (-2.2Kdec^{-1}). This cooling results in Southern Hemisphere summertime tropospheric circulation changes captured by an increase in the Southern Annular Mode (SAM) index (1.27hPa dec^{-1}). In the future, the interplay between the ozone hole recovery and greenhouse gases (GHGs) concentrations may result in the SAM index returning to pre-ozone hole levels or even with a more positive phase from around the second half of the century (-0.4 and 0.3hPa dec^{-1} for the RCP2.6 and RCP8.5, respectively). By 2100, stratospheric ozone sensitivity to GHG concentrations is greatest in the Arctic and Northern Hemisphere midlatitudes (37.7 and 16.1 DU difference between the RCP2.6 and RCP8.5, respectively), and smallest over the tropics and Antarctica continent (2.5 and 8.1 DU respectively). Future TCO changes in the tropics are mainly determined by the upper stratospheric ozone sensitivity to GHG concentrations, due to a large compensation between tropospheric and lower stratospheric column ozone changes in the two RCP scenarios. These results demonstrate how changes in stratospheric ozone are tightly linked to climate and show the benefit of including the processes interactively in climate models.

15, 25175–25229, 2015

ACPD

Stratospheric ozone
change and related
climate impacts over
1850–2100

F. Iglesias-Suarez et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
	
	
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



1 Introduction

The Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP) (Lamarque et al., 2013b) was designed to evaluate the long-term (1850–2100) atmospheric composition changes (e.g. ozone) to inform the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (IPCC, 2013), supplementing phase 5 of the Coupled Model Intercomparison Project (CMIP5) (Taylor et al., 2012), where the focus was more on physical climate change. In addition, ACCMIP is the first model intercomparison project in which the majority of the models included chemical schemes appropriate for stratospheric and tropospheric chemistry. Due to the absorption of shortwave radiation, stratospheric ozone is important for determining the stratospheric climate (e.g. Randel and Wu, 1999) and has a strong influence on tropospheric ozone through stratosphere-to-troposphere transport (e.g. Collins et al., 2003; Sudo et al., 2003; Zeng and Pyle, 2003; Hegglin and Shepherd, 2009). In addition, changes in stratospheric ozone can affect atmospheric circulation and climate, reaching to the lower troposphere in the case of the Antarctic ozone hole (e.g. Thompson and Solomon, 2002; Gillett and Thompson, 2003). This study evaluates stratospheric ozone changes and associated climate impacts in the ACCMIP simulations, quantifying the evolution since the pre-industrial period through to the end of the 21st century.

Stratospheric ozone represents approximately 90 % of ozone in the atmosphere and absorbs much of the ultraviolet solar radiation harmful for the biosphere (e.g. WMO, 2014; UNEP, 2015). Anthropogenic emissions of ozone depleting substances (ODS) such as chlorofluorocarbons and other halogenated compounds containing chlorine and bromine have played a key role in depleting stratospheric ozone during the latter half of the 20th century (e.g. WMO, 2014). Although present globally averaged TCO levels are only ~ 3.5 % lower than pre-1980 values, about half the TCO is depleted over Antarctica between September and November (austral spring) each year (Forster et al., 2011). Halogen loading peaked around 1998 and started to decrease afterwards due to the implementation of the Montreal Protocol and its Amendments and Adjustments

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
	
	
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



(e.g. WMO, 2007, 2014). As a result, stratospheric ozone is expected to recover and return to pre-industrial values during the 21st century (Austin and Wilson, 2006; Eyring et al., 2007, 2010a, 2013; Austin et al., 2010). Although anthropogenic ODS are the main cause of ozone depletion over the last decades, other species such as methane, nitrous dioxide (N_2O) and carbon dioxide (CO_2) affect stratospheric ozone chemistry as well (Haigh and Pyle, 1982; Portmann et al., 2012; Revell et al., 2012; Reader et al., 2013; Dietmüller et al., 2014). Randeniya et al. (2002) argued that increasing concentrations of methane can amplify ozone production in the lower stratosphere via photochemical production, though increases of water vapour from methane oxidation may have the opposite effect (Dvortsov and Solomon, 2001). Nitrogen oxides (NO_x) chemistry is important in the middle-upper stratosphere for ozone; thus, variations and trends in the source gas (N_2O) may have a substantial influence on ozone levels (e.g. Ravishankara et al., 2009; Portmann et al., 2012; Revell et al., 2012).

As ODS levels slowly decrease, projected climate change will likely play a key role in stratospheric ozone evolution through its impacts on temperature and atmospheric circulation (e.g. IPCC, 2013). The impact of climate change on ozone in the stratosphere further complicates the attribution of the recovery (e.g. Waugh et al., 2009a; Eyring et al., 2010b) since increases in CO_2 levels cool the stratosphere, slowing gas-phase ozone loss processes (e.g. reduced NO_x abundances; reduced HO_x -catalysed ozone loss) resulting in ozone increases, particularly in the middle-upper stratosphere and high latitudes (e.g. Haigh and Pyle, 1982; Randeniya et al., 2002; Rosenfield et al., 2002; Jonsson et al., 2004). Further, an acceleration of the equator-to-pole Brewer–Dobson Circulation (BDC) has been predicted in many model studies under high GHG concentrations (e.g. Butchart et al., 2006, 2010; Garcia and Randel, 2008; Li et al., 2008; Shepherd, 2008), although its strength can only be inferred indirectly from observations, meaning that there are large uncertainties in recent trends (e.g. Engel et al., 2009; Bönisch et al., 2011; Young et al., 2011; Stiller et al., 2012). This BDC acceleration enhances transport in the atmosphere and stratospheric–tropospheric exchange (STE), and is likely to have a substantial role throughout the 21st century (e.g. Butchart,

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Stratospheric ozone
change and related
climate impacts over
1850–2100**

F. Iglesias-Suarez et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

sphere (e.g. Prather et al., 2001; Gregg et al., 2003; Jerrett et al., 2009). Its abundance in the troposphere is determined from the balance of STE and photochemistry production involving the oxidation of hydrocarbons and carbon monoxide (CO) in the presence of NO_x , vs. chemical destruction and deposition to the surface (e.g. Lelieveld and Dentener, 2000; Wild, 2007). These terms depend in turn on climate system dynamics (e.g. STE) and on the magnitude and spatial distribution of ozone precursors emissions such as, volatile organic compounds (VOCs), NO_x and CO (e.g. chemical production and destruction) (e.g. Wild, 2007). Several studies found tropospheric ozone increases due to climate change via enhanced STE (e.g. Collins et al., 2003; Sudo et al., 2003; Hegglin and Shepherd, 2009). Other studies have shown positive relationship between anthropogenic emissions and tropospheric ozone abundance (Stevenson et al., 2006; Young et al., 2013a). However, the ultimately net impact of climate and emissions changes remains unclear (Stevenson et al., 2006; Isaksen et al., 2009; Jacob and Winner, 2009), and it may differ substantially by region, altitude or season (e.g. Myhre et al., 2013).

Further, the ozone hole influences surface climate via temperature and circulation changes (e.g. Thompson and Solomon, 2002; Gillett and Thompson, 2003) owing to direct radiative effects (e.g. Randel and Wu, 1999; Forster et al., 2011). The ozone layer heats the stratosphere by absorbing incoming ultraviolet solar radiation, hence, trends and variations on ozone would impact stratospheric dynamics (e.g. Ramaswamy et al., 2006; Randel et al., 2009; Gillett et al., 2011). In the Southern Hemisphere (SH), stratospheric circulation changes associated to ozone depletion have been linked to tropospheric circulation changes primarily during austral summer (lagging the former 1–2 months), based on observations (Thompson and Solomon, 2002) and model simulations (Gillett and Thompson, 2003). These SH extratropical circulation changes could be described by the leading mode of variability or the SAM (e.g. Thompson and Wallace, 2000). Previous studies based on CCMs simulations reported positive trends in the SAM over the ozone depletion period (Sexton, 2001; Gillett and Thompson, 2003; Shindell and Schmidt, 2004; Arblaster and Meehl, 2006; Son et al., 2008; Polvani et al., 2010; Sigmond et al., 2010; McLandress et al., 2011). Projected ozone recovery should

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

have the opposite effect than ozone depletion (i.e. a negative trend in the SAM), and this is important as it opposes the effect of increasing GHG concentrations. Some studies suggest that these effects will largely cancel out each other during the next several decades in austral summer owing to these competing forces (Shindell and Schmidt, 5 2004; Perlitz et al., 2008; Son et al., 2009, 2010; Arblaster et al., 2011; Polvani et al., 2011; Barnes et al., 2013; Gillett and Fyfe, 2013).

Multi-model experiments are useful for evaluating model differences in not fully understood processes and associated feedbacks, and for identifying agreements and disagreements between various parameterisations (e.g. Shindell et al., 2006; Stevenson et al., 2006). While CMIP5 provides a framework towards a more Earth System approach to intercompare model simulations and enables their improvement, it lacks comprehensive information on atmospheric composition and models with full interactive chemistry (Lamarque et al., 2013b). ACCMIP aims to fill this gap by evaluating how atmospheric composition drives climate change, and provides a gauge of the uncertainty by different physical and chemical parameterisations in models (Myhre et al., 15 2013). In this study we quantify the evolution of stratospheric ozone and related climate impacts in the ACCMIP simulations from pre-industrial times (1850), recent past (1980) and present day (2000) to the near-future (2030) and the end of the 21st century (2100). First, we evaluate recent past and present-day ACCMIP stratospheric ozone simulations with observations and other model based products. Then, we assess ozone 20 projections and ozone sensitivity to GHG concentrations. Finally, a description of the associated impacts of stratospheric ozone depletion and projected recovery in the climate system is presented, with a focus in the SH. In addition, this study compares ACCMIP simulations with those from CMIP5 and CCMVal2 and identifies agreements 25 and disagreements among different parameterisations. This paper complements previous analysis of the ACCMIP simulations on tropospheric ozone evolution (Young et al., 2013a; Parrish et al., 2014), radiative forcing (Bowman et al., 2013; Shindell et al., 2013a; Stevenson et al., 2013), hydroxyl radical and methane lifetime (Naik et al., 2013b; Voulgarakis et al., 2013), historical black carbon evaluation (Lee et al., 2013),

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



nitrogen and sulfur deposition (Lamarque et al., 2013a), and climate evaluation (Lamarque et al., 2013b).

The remainder of this paper is organised as follows. Section 2 describes the models and simulations used here, with a focus on the various ozone chemistry schemes.

- 5 In Sect. 3, ozone is examined in the recent past against observations, and analysed from 1850 to 2100 under the low and high RCPs emission scenarios for those models with interactive chemistry-climate feedback. Section 4 explores past and future stratospheric ozone evolution and climate interactions. A discussion of the results is presented in Sect. 5, followed by a brief summary and main conclusions in Sect. 6.

10 **2 Models, simulations and analysis**

In this section we describe main details of the ACCMIP models, simulations, and analysis conducted in this paper. A comprehensive description of the models and simulations along with further references are provided by Lamarque et al. (2013b).

2.1 ACCMIP models

- 15 Table 1 summarises the ACCMIP models analysed in this study and their important features. We considered 8 models that had time-varying stratospheric ozone, either prescribed (offline) or interactively calculated (online). From the full ACCMIP ensemble (Lamarque et al., 2013b), we have excluded: EMAC, GEOSCCM and GISS-E2-TOMAS, as these did not produce output for all the scenarios and time periods analysed here (see Sect. 2.2); CICERO-OsloCTM and LMDzORINCA, as these used a constant climatological value of stratospheric ozone; MOCAGE and STOC-HadAM3, which showed poor stratospheric ozone chemistry performance compared to observations; and NCAR-CAM5.1, as this model was focused on aerosol output and did not save ozone fields.
- 20

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
	
	
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

The ACCMIP models included in this study are CCMs (7) or chemistry general circulation models (1) with atmospheric chemistry modules. The CCMs implemented a coupled composition-radiation scheme, whereas the chemistry and radiation was not coupled in UM-CAM (see Table 1). Both sea surface temperatures and sea-ice concentrations were prescribed, except in GISS-E2-R which interactively calculated them. Similarly to Eyring et al. (2013), we group the models into two categories: 6 models with full atmospheric chemistry (CHEM), and 2 models with online tropospheric chemistry but with prescribed ozone in the stratosphere (NOCHEM) (Fig. 4 of Lamarque et al., 2013b). All CHEM models included ODS (with Cl and Br) and the impact of polar stratospheric clouds (PSCs) on heterogeneous chemistry, although a linearised ozone chemistry parameterisation was implemented in CESM-CAM-Superfast (McLinden et al., 2000; Hsu and Prather, 2009). The other two models, HadGEM2 and UM-CAM, prescribed stratospheric ozone concentrations from the IGAC/SPARC database (Cionni et al., 2011).

A final important distinction among the models is how stratospheric changes are able to influence photolysis rates. The simplest scheme is for UM-CAM, where the photolysis rates are derived from a look-up table as a function of time, latitude and altitude only, and using a climatological cloud and ozone fields (i.e. the rates are the same for all simulations) (e.g. Zeng et al., 2008, 2010). HadGEM2 also employs a look-up table, but the rates are additionally a function of the model overhead ozone column, and are therefore sensitive to stratospheric ozone change (Collins et al., 2011; Martin et al., 2011). The look-up table is more complex still with CESM-CAM-Superfast (Gent et al., 2010), CMAM (Scinocca et al., 2008), GFDL-AM3 (Donner et al., 2011; Griffies et al., 2011) and NCAR-CAM3.5 (Gent et al., 2010; Lamarque et al., 2012), where an adjustment is applied to take surface albedo and cloudiness into account, which couples with the simulated aerosols. Fully online photolysis calculations were only made for MIROC-CHEM (Watanabe et al., 2011) and GISS-E2-R (Schmidt et al., 2006; Shindell et al., 2013b).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

As per Young et al. (2013a), all models were interpolated to a common grid (5° by 5° latitude/longitude and 24 pressure levels).

2.2 ACCMIP scenarios and simulations

The ACCMIP simulations were designed to span the pre-industrial period to the end of the 21st century. In this study, time slices from the years 1850, 1980 and 2000 comprise historical projections (hereafter Hist), whereas time slices from the years 2030 and 2100 future simulations. The latter follow the climate and composition/emission projections prescribed by the Representative Concentration Pathways (RCPs) (van Vuuren et al., 2011; Lamarque et al., 2012), named after their nominal radiative forcing at the end of the 21st century relative to 1750. Here we consider RCP2.6 (referring to 2.6 W m^{-2}) and RCP8.5 (8.5 W m^{-2}), since they bracket the range of warming in the ACCMIP simulations, and are the scenarios that have been completed by the greatest number of models.

Future ODS (the total organic chlorine and bromine compounds) in CHEM models follow the RCPs values from Meinshausen et al. (2011), which does not include the early phase-out of hydrochlorofluorocarbons agreed in 2007 by the Parties to the Montreal Protocol. Note that ODS may be specified as concentrations (CMAM, GFDL-AM3 and NCAR-CAM3.5) or emissions (CESM-CAM-superfast, GISS-E2-R, MIROC-CHEM) in different models, though these were the same within each time slice simulation (except for GISS-E2-R; see below). This is slightly different from the modified halogen scenario of WMO (2007) used in the IGAC/SPARC ozone database employed by the NOCHEM models. Nevertheless, halogen concentrations in both future scenarios peak around the year 2000 and decline afterwards, although slightly different timing of ozone returning to historical levels may be found. Tropospheric ozone precursors emissions follow Lamarque et al. (2010) for the historical period, and Lamarque et al. (2013b) for the RCPs.

Most models completed time slice simulations for each period and scenario, usually 10 years average about each time slice (e.g. 1975–1984 for the Hist 1980 time slice,

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
	
	
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



although other models simulated time slices ranging from 5 to 11 years). Notice that interannual variability for a given time slice is generally small (Young et al., 2013a). The exception is GISS-E2-R, which ran transient simulations with a coupled ocean. Equivalent time slice means were calculated by averaging 10 years centred on the desired time slice, (1975–1984 for 1980 and so forth), except for the 1850 time slice (1850–1859 mean).

2.3 CMIP5 and CCMVal2 simulations

We also include CMIP5 and CCMVal2 simulations as a benchmark for the former models. We use a subset of five “high” top CMIP5 models, defined here as those models that represented and saved ozone output above 10 hPa for the historical (1850–2005, most of the models), and future (RCP2.6 and RCP8.5, 2005–2100) emission scenarios: CESM1-WACCM, GFDL-CM3, MPI-ESM-LR, MIROC-ESM, and MIROC-ESM-CHEM. Only high top models are considered here due to the implications the upper stratosphere has on, among other factors, stratospheric dynamical variability (Charlton-Perez et al., 2013), and tropospheric circulation (Wilcox et al., 2012). Moreover, we will show how, in the tropics, upper stratospheric ozone plays a key role on TCO projections during the 21st century (see Sect. 3.2). Again, we group the models into two categories: 3 models with full atmospheric chemistry (CHEM: CESM1-WACCM, GFDL-CM3 and MIROC-ESM-CHEM), and 2 models with prescribed ozone (NOCHEM: MPI-ESM-LR and MIROC-ESM). A detailed description of the models, simulations and ozone concentrations are presented by Taylor et al. (2012) and Eyring et al. (2013).

In addition, we include 14 CCMVal2 models that represented ozone under the REF-B1 scenario (1960–2006, most of the models): CAM3.5, CCSRNIES, CMAM, E39CA, EMAC, GEOSCCM, LMDZrepro, Niwa-SOCOL, SOCOL, ULAQ, UMETRAC, UMUKCA-METO, UMUKCA-UCAM and WACCM. All these models had interactive stratospheric chemistry and coupled composition-climate feedback, although simplified or absent chemistry in the troposphere. Morgenstern et al. (2010) describe in detail CCMVal2 models and REF-B1 simulations.

In contrast to ACCMIP time slice simulations, these data sets were based on transient experiments, which may result in slightly different ozone levels, as simulations depart from initial conditions. Nevertheless, equivalent time slice means were calculated in the same manner as above for consistency purposes throughout all analysis involving trends or ozone changes. Note, however, that calculating trends using least-squares linear fits from their transient runs would not have a significant impact on the results. A caveat is that TCO was calculated from the ozone mixing ratio field, which may slightly differ ($\sim 1.5\%$) from that of the model's native TCO (Eyring et al., 2013).

2.4 Tropopause definition

- For the purpose of comparing the outputs among models, a tracer tropopause definition has been argued to be suitable (Wild, 2007). This study follows Young et al. (2013a) method, in which the tropopause is based on the 150 ppbv ozone contour, after Prather et al. (2001). The definition is fitted for all time slices using ozone from the Hist 1850 time slice for each model and month; meaning that the “troposphere” is defined as a fixed volume region of the atmosphere. On the one hand, Young et al. (2013a) argued that using a monthly mean tropopause from the 1850 time slice prevents issues with different degrees of ozone depletion among the models, especially for SH high latitudes. On the other hand, this neglects the fact that the tropopause height may vary with time due to climate change (e.g. Santer et al., 2003a, b). Nevertheless, Young et al. (2013a) have shown that using ozone from the Hist 2000 time slice to define the tropopause across all time slices, generally results in tropospheric ozone columns of $\pm 5\%$ compared to the Hist 1850 time slice.

3 Long-term total column ozone evolution in the ACCMIP models

This section presents an evaluation of the present-day (Hist 2000) TCO distribution and recent (1980–2000) ozone trends against observations and observationally-derived

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
	
	
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
	
	
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



data. The evolution of TCO from the pre-industrial period (1850) to the end of the 21st century (2100) is also discussed, with a particular focus on the different contribution of trends in the tropical tropospheric, lower stratospheric, and upper stratospheric columns to the total column trend. Previously, Young et al. (2013a) have shown that 5 TCO distribution changes in the ACCMIP multi-model mean agree well with the Total Ozone Mapping Spectrometer (TOMS) for the last few decades (their Fig. S7). However, ACCMIP models simulate weaker (not significant) ozone depletion in early boreal spring over the Arctic between Hist 1980 and 2000 compared to TOMS (see also Sects. 3.1 and 5).

10 3.1 Evaluation of ozone trends, 1980–2000

Figure 1 shows TCO decadal trends between 1980 and 2000 for the global mean, and a number of latitude bands. The figure compares the ACCMIP, CMIP5 and CCMVal2 models against the Bodeker Scientific TCO data set (Bodeker Scientific TCO – version 2.8), combining a number of different satellite-based instruments (Bodeker et al., 2005; Struthers et al., 2009), and observations from the Solar Backscatter Ultraviolet (SBUV – version 15 8.6) merged ozone data sets (McPeters et al., 2013). In addition, Fig. 1 includes trends from the IGAC/SPARC ozone data set (Cionni et al., 2011) which was used by the majority of the models with prescribed ozone concentrations (both ACCMIP and CMIP5). The different data sets trends are broadly comparable but differ slightly in their calculation and uncertainty determination. For ACCMIP, CMIP5 and CCMVal2 models, the 20 trends are for the differences between the Hist 1980 and 2000 time slices with the range shown as box/whisker plots (central 50 % of trends as the box; 95 % confidence intervals as the whiskers). Trends for the observations and IGAC/SPARC database are linear trends with error bars indicating the 95 % confidence level based on the standard error for the fit, and corrected for lag-1 autocorrelation for the former (Santer et al., 25 2000). The annual mean is used in evaluations for the global, tropical and midlatitudes regions. Additional evaluations are made for the boreal spring in the Arctic (March, April

and May) and the austral spring in the Antarctic (September, October and November) when strongest ozone depletion occurs.

Within uncertainty, the overall response for ACCMIP is in good agreement with observational data sets in terms of decadal trends and absolute values, with the Northern Hemisphere (NH) being the region where models differ most. In line with CMIP5 and CCMVal2 models, strongest changes are found over Antarctica in austral spring associated to the ozone hole, and smallest over the tropics where ODS are least effective. ACCMIP NOCHEM models typically simulate smaller decadal trends than CHEM models, consistent with the possible underestimation of SH ozone depletion trends in the IGAC/SPARC ozone data set (Hassler et al., 2013; Young et al., 2014). However, outside extratropical SH regions, IGAC/SPARC tends to show better agreement with observations than CHEM models. ACCMIP CHEM and CMIP5 CHEM models show very similar TCO decadal trends in all regions ($\pm 0.10\text{--}0.24\,\%\text{dec}^{-1}$), although differing somewhat more at high latitudes in the SH, where ozone depletion is greatest ($\pm 2.9\,\%\text{dec}^{-1}$). ACCMIP NOCHEM and CMIP5 NOCHEM models show more disparate trends ($\pm 0.46\text{--}2.11\,\%\text{dec}^{-1}$), which may be related to different ozone data sets and the implementation method on each model (i.e. online tropospheric chemistry in ACCMIP models).

Figure 2 compares vertically resolved ozone decadal trends for the same period, regions and seasons, for the ACCMIP multi-model mean and individual models against the Binary Database of Profiles (BDBP version 1.1.0.6) data set, using the so-called Tier 0 and Tier 1.4 data (Bodeker et al., 2013). Tier 0 includes ozone measurements from a wide range of satellite and ground-based platforms, whereas Tier 1.4 is a regression model fitted to the same observations. Uncertainty estimates for the BDBP Tier 1.4 trends are from the linear least square fits, as for the observations in Fig. 1. ACCMIP shows most disagreement with the BDBP data in the lower and middle stratosphere region and best agreement with Tier 1.4 in the upper stratosphere.

In the Tropics (Fig. 1b), within uncertainty all data sets agree fairly well with observations. Although ACCMIP, CMIP5 and CCMVal2 simulate small decadal trends

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- (-0.42 , -0.66 and -0.85 \% dec^{-1} respectively), the spread of the models at the 95 % confidence interval stays within the negative range. However, uncertainty estimates in TCO in the SBUV and BodSci TCO data sets embrace trends of different sign (-0.65 ± 1.52 , and $-0.43 \pm 2.27\text{ \% dec}^{-1}$ respectively). IGAC/SPARC presents slightly stronger decadal trends than observations in this region. CMIP5 and CCMVal2 multi-model means show slightly stronger decadal trends and agree better with observations than ACCMIP in this region. In terms of absolute values, the spread of the ACCMIP models overlaps the observed TCO for the Hist 2000 time slice, though most models differ by more than the observational standard deviation (7 out of 8). Biases in TCO may be attributed to different altitude regions (Fig. 2b). ACCMIP models fail to represent observed ozone depletion occurring in the lower and middle stratosphere region, which may be linked to a poor representation of the HO_x and NO_x catalytic loss cycles (e.g. Lary, 1997; Nedoluha et al., 2015).

In the NH midlatitudes (Fig. 1c), ACCMIP simulates smaller decadal trends than CMIP5 and CCMVal2 (-0.84 , -1.36 and -1.38 \% dec^{-1} respectively), though all data sets are at the low end of the observational uncertainties ($-1.64\text{--}2.45 \pm 1.2\text{ \% dec}^{-1}$ respectively). TCO decadal trends for IGAC/SPARC and NOCHEM models, show better agreement with observations than CHEM models in this region. Due to the BDC, the abundance of ozone at midlatitudes is affected by the relatively ozone-rich air coming from the upper stratosphere over the tropics. The ACCMIP Hist 2000 simulation agrees fairly well with observations in terms of absolute values, however, once again most models diverge by more than the observational standard deviation (7 out of 8). The ACCMIP multi-model mean falls within the BDBP Tier 1.4 uncertainty estimates for most of the lower and middle stratosphere, though simulates weaker ozone depletion in the lower stratosphere and fails to capture a small positive trend between 10–5 hPa, likely associated to tropospheric upwelling and ozone catalytic loss cycle via NO_x , respectively (Fig. 2c).

Over the Arctic in boreal spring (Fig. 1e), again the ACCMIP, CMIP5 and CCMVal2 data sets show weaker decadal trends than observations (-2.01 , -3.08 and

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

3.2 Past modelled and future projected total column ozone

In this section, the evolution of past modelled TCO (from 1850 to 2000) and the sensitivity of ozone to future GHG emissions (from 2030 to 2100) under the lower and higher RCPs scenarios are discussed for the regions and seasons presented in the evaluation section. In the tropical region, TCO evolution is further analysed by looking at the stratospheric (split into upper and lower regions) and tropospheric columns ozone. Historical and future global annual mean of TCO and associated uncertainty (± 1 standard deviation) for the ACCMIP and CMIP5 CHEM models and the IGAC/SPARC data set is given in Table 2.

To probe how different emissions of GHG affect stratospheric ozone, we only include in this section ACCMIP and CMIP5 models with full ozone chemistry (CHEM). In addition, we compare these results with the IGAC/SPARC database, generally used by those models with prescribed stratospheric ozone. Note that tropospheric column ozone under the RCPs at the end of the 21st century could lead to differences in TCO around 20 DU, due to differences in ozone precursors emissions (e.g. methane) (Young et al., 2013a). Again, vertical resolved ozone changes are presented to give insight on the vertical distribution of ozone changes (for the 1850–2100 and 2000–2100 periods).

Figure 3 shows, except for the extratropical regions in the SH, an increase in TCO from the pre-industrial period (Hist 1850) to the near-past (Hist 1980) owing to ozone precursors emissions. In the SH extratropical, due to special conditions (e.g. greater isolation from the main sources of ozone precursors and stratospheric cold temperatures during austral winter and early spring), there is a decrease in TCO that is particularly pronounced over Antarctica (−12.4 %). Between near-past and present-day (Hist 2000), a period characterised by ODS emissions, the TCO decreases everywhere, with the magnitude being dependent on the region. Thus, the relative change of TCO between the present-day and pre-industrial periods varies across different regions, mainly due to the competing effects of ozone precursors and ODS emissions (approximately,

[Title Page](#)[Abstract](#) [Introduction](#)[Conclusions](#) [References](#)[Tables](#) [Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Stratospheric ozone
change and related
climate impacts over
1850–2100**

F. Iglesias-Suarez et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

from 2.9 % in the NH midlatitudes and –34.9 % over Antarctica). Notice, however, that minimal stratospheric ozone depletion occurs before the 1960s.

Future TCO projected for the RCPs 2100 time slices relative to present-day are affected by the impact of the Montreal Protocol on limiting ODS emissions, climate change and ozone precursors emissions. TCO changes between 2000 and 2100 relative to the pre-industrial period for the low and high emission scenarios are in the range of approximately from –1.2 to 2.0 % in the tropics and 28.3–31.7 % over Antarctica, respectively. Ozone “super-recovery”, defined here as higher stratospheric ozone levels than those during pre-ozone depletion (1850), is found for ACCMIP CHEM models in RCP8.5 2100 in all regions and seasons, with the exception in the tropics and over Antarctica during austral spring. As expected from the above climate impacts, the biggest super-recovery is found, in the order of 12.6 % over the Arctic during boreal spring, and between 3.8–6.5 % at midlatitudes for the RCP8.5 2100 time slice. Similar levels of stratospheric ozone super-recovery are found in the CMIP5 CHEM models. In contrast, the IGAC/SPARC database only projects small super-recovery in the NH polar region and at midlatitudes in the SH. These ozone super-recovery results are consistent with recent findings on stratospheric ozone sensitivity to GHG concentrations (Waugh et al., 2009a; Eyring et al., 2010b).

We give special attention to TCO projections in the tropics, since an acceleration of the BDC, due to increases in GHG concentrations would lead to a rise of tropospheric ozone-poor air entering the tropical lower stratosphere (Butchart et al., 2006, 2010, 2011; SPARC-CCMVal, 2010; Eyring et al., 2013). In other words, ozone concentrations in the lower stratosphere would decrease with high GHG emissions.

Figure 4 presents upper (> 10 hPa) and lower (150–15 hPa) stratospheric and tropospheric columns ozone in the tropics, from the pre-industrial period to the end of the 21st century. Tropospheric column ozone increases with higher ozone precursors emissions during the historical period (1850–2000). Future emissions of ozone precursors (e.g. CO and NO_x) are fairly similar among the RCPs scenarios, decreasing to various degrees between the present-day and 2100 (van Vuuren et al., 2011). The ex-

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

ception is that the methane burden under the RCP8.5 scenario roughly doubles by the end of the 21st century (Meinshausen et al., 2011). Mainly due to the methane burden and the stratospheric ozone influence via STE, ACCMIP CHEM tropospheric column ozone change by 2100 relative to present-day is -5.5 and 5.8 DU, for the RCP2.6 and RCP8.5 scenarios respectively. For both stratospheric columns ozone, there is a small decrease from the pre-industrial period to present-day (-3.2 – 3.3 DU), which remained fairly constant by 2030 for both RCPs scenarios. Although ODS concentrations decrease during the 21st century, two different stories occur in the second half of the century. In the upper stratosphere, ozone amounts return to pre-industrial levels under the low emission scenario by 2100. However, RCP8.5 2100 ozone levels relative to present-day increase 8.3 DU, due to a slow down of the ozone catalytic loss cycles, linked to the stratospheric cooling (e.g. Haigh and Pyle, 1982; Portmann and Solomon, 2007; Revell et al., 2012; Reader et al., 2013; Dietmüller et al., 2014). In the lower stratosphere, ozone levels change little (-0.8 DU) by 2100 relative to the present-day for the RCP2.6, though decrease by -8.5 DU under the RCP8.5 scenario, likely due to the acceleration of the BDC. In summary, stratospheric column ozone by 2100 remains fairly similar to the present-day, although different stories are drawn in the upper and lower stratosphere. RCP2.6 and RCP8.5 2100 TCO relative to pre-industrial period is the result of changes in lower stratospheric and tropospheric columns ozone largely cancelling each other. Therefore, future TCO evolution in the tropics is primarily determined by the upper stratospheric ozone sensitivity to GHG concentrations under the RCPs considered here.

Figure 5 presents vertically resolved ozone change between the Hist 1850 and RCPs 2100 time slices and between the Hist 2000 and RCPs 2100 time slices (top and bottom rows, respectively). In the midlatitudes (Fig. 5b and d), lower stratospheric ozone is positively correlated to GHG concentrations, in contrast to the tropics, due to the influx of relatively “rich” ozone air from lower latitudes (e.g. WMO, 2011). However, middle and upper stratospheric ozone sensitivity to GHG concentrations behaves the same as in the tropics. Substantial ozone increases are simulated by 2100, in the altitude region

of the upper troposphere-lower stratosphere and the middle and upper stratosphere, relative to pre-industrial (1850) and present-day (2000) levels. We note that climate impact in ozone levels is weaker in the southern than in the northern midlatitudes for the ACCMIP and CMIP5 multi-model means, likely due to hemispheric differences in STE and ozone flux (Shepherd, 2008), which is in contrast to IGAC/SPARC data set. TCO for the RCP8.5 2100 time slice is 6.9–13.1 % higher than those simulated in the Hist 1850 time slice. While, the RCP2.6 2100 time slice in the northern midlatitudes is similar to present-day levels, in the southern midlatitudes is similar to pre-industrial levels. This is mainly due to regional differences in ozone precursors emissions and the tropospheric ozone contribution (Fig. 3c and d).

Over the Arctic in boreal spring (Fig. 3e), results similar to those in the northern mid-latitudes are found for all models, though higher stratospheric ozone sensitivity to GHG concentrations lead to approximately two times larger scenario differences for the 2100 time slice (37.7 DU between RCP2.6 and RCP8.5). In addition to the RCP8.5 emission scenario, ozone super recovery is also simulated under the RCP2.6 scenario by ACCMIP and CMIP5 CHEM models. The IGAC/SPARC data set projects similar results to those under the latter scenario. Note that the ACCMIP and CMIP5 multi-model means show a small increase in TCO by 1980 and no significant ozone depletion by 2000 relative to 1850. This is in sharp contrast to the polar region in the SH, which highlights both regional differences in ozone precursors sources and atmospheric conditions.

Over Antarctica during austral spring (Fig. 3f), TCO evolution is more isolated from GHG effects and ozone precursors than in other regions. In agreement with previous studies, ACCMIP and CMIP5 CHEM models project similar values under the lower and higher GHG scenarios (Austin et al., 2010; SPARC-CCMVal, 2010; Eyring et al., 2013). TCO in the RCPs 2100 time slices remained below 1850s levels (−3.3–6.7 %). This suggests decreasing ODS concentrations during the 21st century as the main driver of stratospheric ozone in this region and season (i.e. ozone super-recovery is found for RCP8.5 2100 in other seasons). Furthermore, vertical distribution changes of stratospheric ozone in 2100, compared to 1850 (Fig. 5f1), and 2000 (Fig. 5f2), show

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



small differences between the above scenarios (e.g. small sensitivity to GHG concentrations). Evolution of stratospheric ozone at high latitudes in the SH, particularly during spring season, has implications over surface climate due to modifications in temperature and circulation patterns.

5 **4 Stratospheric ozone changes and associated climate impacts in the Southern Hemisphere**

To probe stratospheric ozone evolution and climate interactions (1850–2100), we first examine simulated stratospheric temperatures in Sect. 4.1. SAM index evolution is presented in Sect. 4.2. Note that ozone loss over the Arctic in boreal spring is only around 25 % of the depletion observed in the Antarctic (see also Fig. 1e), and is not believed to have a significant role in driving NH surface climate (e.g. Grise et al., 2009; Eyring et al., 2010a; Morgenstern et al., 2010).

10 **4.1 Lower stratospheric temperatures changes**

Figure 6 shows recent stratospheric temperature decadal trends (1980–2000) in polar regions during springtime (March–April–May in the Arctic and October–November–December in the Antarctic). The figure compares temperature in the lower stratosphere (TLS) in the ACCMIP, CMIP5 and CCMVal2 models with observational estimates based on Microwave Sounding Units (MSU) of the Remote Sensing Systems (RSS – version 3.3) (Mears et al., 2011), the Satellite Applications and Research (STAR – version 3.0) (Zou et al., 2006, 2009), and the University of Alabama in Huntsville (UAH – version 5.4) (Christy et al., 2003) (Fig. 6a–c). The TLS vertical weighting function from RSS is used to derive MSU temperature from climate models output. Temperature vertical profile decadal trends in the ACCMIP models (Fig. 6b–d) are compared against radiosonde products of the Radiosonde Observation Correction Using Reanalyses (RAOBCORE – version 1.5), Radiosonde Innovation Composite Homogenization

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

(RICH-obs and RICH-tau – version 1.5) (Haimberger et al., 2008, 2012), the Hadley Centre radiosonde temperature product (HadAT2) (Thorne et al., 2005), and the Iterative Universal Kriging (IUK) Radiosonde Analysis Project (Sherwood et al., 2008) (version 2.01).

- 5 Over the NH polar cap in boreal spring, although ACCMIP, CMIP5 and CCMVal2 models are within observational estimates, all simulates weaker decadal trends (-0.5 , -0.1 , -0.4 K dec^{-1} , respectively) than observed ($-1.5\text{--}1.7 \pm 3.4 \text{ K dec}^{-1}$) (Fig. 6a). This is likely due to the abnormally cold boreal winters in the mid-1990s (i.e. more PSCs formation), which resulted in enhanced ozone loss during boreal spring (Newman et al.,
10 2001). Natural variability in models not constrained by observed meteorology is difficult to reproduce (Austin et al., 2003; Charlton-Perez et al., 2010, 2013; Butchart et al., 2011; Shepherd et al., 2014). Moreover, ACCMIP simulations, based on time slice experiments for most models, did not embrace that period, only those boundary conditions for 1980 and 2000 years. This weaker trend on stratospheric temperature is
15 also seen in the vertical profile above around the tropopause (Fig. 6b).

Over Antarctica in austral spring, the ACCMIP, CMIP5 and CCMVal2 multi-model means are in very good agreement (-2.2 , -2.5 , -1.9 K dec^{-1} , respectively) with satellite measurements ($-2.0\text{--}2.3 \pm 6.4 \text{ K dec}^{-1}$) (Fig. 6c). CHEM models tend to simulate stronger trends than NOCHEM models, likely due to the fact that the IGAC/SPARC
20 ozone data set is at the lower end of the observational estimates as has been shown in Solomon et al. (2012), Hassler et al. (2013), Young et al. (2014). They argued the importance of the ozone data set for appropriate representation of stratospheric temperature, and in turn SH surface climate. Although, large uncertainties exist in this region and period, all ACCMIP individual models fall within the observational error estimates (Fig. 6d). Note that observational estimates are significant at the 95 % confidence levels, if year 2000 is removed from the linear fit (-2.95 ± 2.90 , -3.02 ± 2.95 and
25 $-3.12 \pm 2.87 \text{ K dec}^{-1}$ for the RSS, STAR and UAH data sets, respectively), as this year was “anomalously” warm. The relatively large spread of the simulated stratospheric temperature trend for the observational period is consistent with the models spread

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Stratospheric ozone
change and related
climate impacts over
1850–2100**

F. Iglesias-Suarez et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

of ozone in this region (Figs. 1f and 2f). The correlation between stratospheric ozone and temperature trends becomes evident by comparing TCO trends between the Hist 1980 and 2000 time slices and TLS trends for the same period between CHEM and NOCHEM models (i.e. large ozone depletion results in stronger stratospheric cooling trends).

- Figure 7a depicts SH polar cap TLS long-term evolution (1850–2100) normalised to pre-industrial levels during austral spring. As commented above, stratospheric temperature can be perturbed by anthropogenic emissions of ODS and GHG, both having a net cooling effect. ACCMIP Hist 1980 and 2000 TLS time slices (−2.5 and −6.0 K) are driven by the combination of ozone depletion and climate change since the pre-industrial period. In future projections, ozone recovery and GHG concentrations are expected to have an opposite effect on stratospheric temperatures. The slightly increased of the TLS by 2030 in the RCPs time slices relative to present-day, is very similar between the lower and higher RCPs emission scenarios (0.9 and 0.4 K, respectively). By the end of the 21st century, the projected TLS under the RCP2.6 scenario returns to Hist 1980 levels, whereas it remains fairly unchanged under the RCP8.5 scenario relative to 2030. These two different stories suggest a key role of GHG concentration in the second half of the century, with significant implications for many aspects of the SH surface climate as reported previously (McLandress et al., 2011; Perlitz, 2011; Polvani et al., 2011); see Sect. 4.2 and Thompson et al. (2011) and Previdi and Polvani (2014) for a comprehensive review.

4.2 Southern Annular Mode evolution

- The SAM index is defined as per Gong and Wang (1999), by subtracting the zonal mean sea level pressure (SLP) at 65° S latitude from the zonal mean SLP at 40° S latitude from monthly mean output. The SAM index is a proxy of variability in the jets captured by SLP anomalies at middle and high latitudes (e.g. Thompson and Wallace, 2000).

Figure 7b shows SAM index long-term evolution (1850–2100) normalised to 1850 levels during austral summer. Observational estimates based on the Hadley Centre Sea Level Pressure data set (HadSLP2) are shown from 1970 to 2012. The ACCMIP multi-model mean shows a positive trend between Hist 1980 and 2000 time slices ($1.27 \text{ hPa dec}^{-1}$), coinciding with the highest ozone depletion period. Within uncertainty, this is weaker than observational estimates ($2.2 \pm 1.1 \text{ hPa dec}^{-1}$). ACCMIP CHEM and NOCHEM models show similar SAM index trends, although the latter presents weaker TLS trends (see Fig. 6c). As seen in Fig. 7a for the TLS in austral spring, by 2030 for both RCPs scenarios the ACCMIP multi-model mean shows a slight decrease in the SAM index relative to Hist 2000.

Two different stories are drawn from 2030 to 2100. The SAM index simulated under the RCP2.6 scenario tends to return to “normal” levels ($-0.4 \text{ hPa dec}^{-1}$), as ODS concentrations and GHG emissions decrease during the second half of the century. In contrast, under the RCP8.5 scenario GHG concentrations increase, resulting in a positive trend of the SAM index (0.3 hPa dec^{-1}). By using a paired sample Student’s *t* test, we find that SAM index changes between Hist 2000 and 2100 relative to Hist 1850, are significant for the RCP2.6 at the 5 % level, although is not significant for the RCP8.5. CMIP5 multi-model mean shows better agreement with observations during the record period (2.1 hPa dec^{-1}) than ACCMIP. During the second half of the 21st century (2030–2100), however, the CMIP5 multi-model mean shows consistent projections with the latter (-0.4 and 0.4 hPa dec^{-1} for RCP2.6 and RCP8.5, respectively).

5 Discussion

TCO trends in ACCMIP models compare favourably with observations, however, ozone trends in the tropical lower stratosphere are low biased (e.g. weaker ozone depletion). It has been argued that tropical upwelling (or the BDC) is the main driver in this region determining ozone levels (Lamarque and Solomon, 2010; Polvani and Solomon, 2012), with chemical processes playing a minor role (e.g. Meul et al., 2014). However,

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
	
	
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



observed BDC and its seasonal cycle (Fu et al., 2010; Young et al., 2011) are poorly constrained in modelling studies (e.g. Butchart et al., 2006, 2010; Garcia and Randel, 2008). This is important since ozone depletion determines to a large extent the temperatures in the lower stratosphere (e.g. Polvani and Solomon, 2012) (note that 5 temperature trends in this region are low biased in ACCMIP models compared to observations, not shown), and the latter triggers significant feedbacks in climate response (Stevenson, 2015). Models with less ozone depletion in the tropical lower stratosphere may have stronger climate sensitivity (Dietmüller et al., 2014; Nowack et al., 2015).

Long-term TCO changes relative to Hist 1850 in the ACCMIP models considered in 10 this study, are least consistent for Hist 2000 in the Antarctic springtime (i.e. the period with large ozone losses) and for RCP8.5 2100 in general. The latter is likely linked to sensitivity of ozone to future GHG emissions uncertainty (i.e. various direct and indirect processes affecting ozone amounts in the troposphere and the stratosphere). For example, CO₂ and methane mixing ratios increase by more than 3 and 4 times in RCP8.5 15 2100 relative to the pre-industrial period, respectively. Nevertheless, the ACCMIP and CMIP5 multi-model means, show consistent RCP8.5 2100 projections. Although TCO changes are relative to the Hist 1850, a period without direct measurements (e.g. estimates with large uncertainties), ACCMIP models show good agreement compared to other time slices. For example, the interquartile range (central 50 % of the data) varies 20 approximately 3–8 % of the corresponding mean value across the regions and seasons considered here.

Stratospheric ozone has been shown to be asymmetrical over the SH polar cap (Grytsai et al., 2007). Prescribing zonal mean ozone fields in CCMs may have implications on SH climate (e.g. Crook et al., 2008; Gillett et al., 2009), particularly in 25 early spring stratospheric temperatures (September–October) and, though less pronounced in November–December (Calvo et al., 2012; Young et al., 2013b). During strong depletion periods such as, in the recent past (1980–2000) and in the near-future (2000–2030), eliminating zonal asymmetry may lead to a poor representation of stratospheric and tropospheric climate trends in the SH (Waugh et al., 2009b). We

showed that NOCHEM models simulated both weaker springtime TLS negative trends over the Antarctic compared to observational estimates, and stronger positive trends in the near-future compared to CHEM models. In addition, Young et al. (2014) find large differences in SH surface climate responses when using different ozone data sets. ACCMIP CHEM and NOCHEM models show most disagreement on SAM index trends in the near-future, period with relatively strong ozone depletion ($> \text{Hist 1980}$). The former projects negligible trends compared to $-0.57 \text{ hPa dec}^{-1}$ and three times weaker negative trends than the latter, for the RCP2.6 and RCP8.5 respectively. This is consistent with CHEM and NOCHEM TLS springtime trends in this period and region. Nevertheless, ACCMIP models participating in this study agree with previous observational (e.g. Thompson and Solomon, 2002; Marshall, 2003, 2007) and modelling studies (e.g. Gillett and Thompson, 2003; Son et al., 2008, 2010, 2009; Polvani et al., 2010, 2011; Arblaster et al., 2011; McLandress et al., 2011; Gillett and Fyfe, 2013; Keeble et al., 2014) on the SH surface climate response, measured here using the SAM index.

6 Summary and conclusions

This study has analysed stratospheric ozone evolution from 1850 to 2100 from a group of chemistry climate models with either prescribed or interactively resolved time-varying ozone in the stratosphere and participated in the ACCMIP activity (8 out of 15 models). We have evaluated TCO and vertically resolved ozone trends between 1980 and 2000, and examined past and future ozone projections under the low and high RCPs future emission scenarios (RCP2.6 and RCP8.5, respectively). Finally, we have assessed TLS and temperature profile trends at high latitudes in the recent past, and analysed TLS and SH surface climate response (diagnosed using the SAM index), from the pre-industrial period to the end of the 21st century.

Within uncertainty estimates, the ACCMIP multi-model mean TCO compares favourably with recent observational trends (1980–2000), although individual models often show significant deviations, particularly those models that include interactive

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)
[▶](#)[▶](#)
[◀](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Overall, stratospheric ozone and associated climate impacts are fairly well represented by the ACCMIP ensemble mean in the recent past (1980–2000), and individual models also agree on the sign and distribution of past and future changes (1850–2100). In line with previous multi-model analyses (Son et al., 2008, 2010; Eyring et al., 2010a, 5 2013; Gillett and Fyfe, 2013), and observation studies (Thompson and Solomon, 2002; Marshall, 2003, 2007), the ACCMIP models show strong positive trends of the SAM index in austral summer during the ozone depletion period ($1.27 \text{ hPa dec}^{-1}$ 1980–2000), which is in agreement with observations ($2.2 \pm 1.1 \text{ hPa dec}^{-1}$). While this trend to a positive SAM index is additive to the SAM trend driven by GHG increases for the present-day, the projected ozone hole recovery during this century will oppose the GHG-driven SAM trend. Under the low emission scenario, the SAM index tends to return to pre-industrial levels from around the second half of the 21st century ($-0.4 \text{ hPa dec}^{-1}$ between 10 2030–2100); i.e. the impact of ozone recovery is stronger than GHG. In contrast, with the higher emission scenario, the GHG-driven SAM trend exceeds the opposing ozone recovery-driven trend, and the SAM index continues on its positive trend (0.3 hPa dec^{-1} between 2030–2100).

In this study we have presented stratospheric ozone evolution (1850–2100) using a number of models that participated in the ACCMIP activity. We have demonstrated both its key role in the present and future SH climate and the importance of how it is 15 represented in climate models (i.e. prescribed or interactively resolved). These results and work over the last decade have shown that changes in stratospheric ozone are tightly coupled to the climate, demonstrating the benefits of including these processes interactively in models. It is clear that our ability to understand future climate will depend on models that can reliably simulate these chemistry-climate feedbacks.

25 References

Arblaster, J. M. and Meehl, G. A.: Contributions of external forcings to Southern Annular Mode trends, *J. Climate*, 19, 2896–2905, doi:10.1175/JCLI3774.1, 2006.

- Arblaster, J. M., Meehl, G. A., and Karoly, D. J.: Future climate change in the Southern Hemisphere: competing effects of ozone and greenhouse gases, *Geophys. Res. Lett.*, 38, L02701, doi:10.1029/2010GL045384, 2011.
- Austin, J. and Wilson, R. J.: Ensemble simulations of the decline and recovery of stratospheric ozone, *J. Geophys. Res.*, 111, D16314, doi:10.1029/2005JD006907, 2006.
- Austin, J., Shindell, D., Beagley, S. R., Brühl, C., Dameris, M., Manzini, E., Nagashima, T., Newman, P., Pawson, S., Pitari, G., Rozanov, E., Schnadt, C., and Shepherd, T. G.: Uncertainties and assessments of chemistry-climate models of the stratosphere, *Atmos. Chem. Phys.*, 3, 1–27, doi:10.5194/acp-3-1-2003, 2003.
- Austin, J., Scinocca, J., Plummer, D., Oman, L., Waugh, D., Akiyoshi, H., Bekki, S., Braesicke, P., Butchart, N., Chipperfield, M., Cugnet, D., Dameris, M., Dhomse, S., Eyring, V., Frith, S., Garcia, R. R., Garny, H., Gettelman, A., Hardiman, S. C., Kinnison, D., Lamarque, J. F., Mancini, E., Marchand, M., Michou, M., Morgenstern, O., Nakamura, T., Pawson, S., Pitari, G., Pyle, J., Rozanov, E., Shepherd, T. G., Shibata, K., Teyssèdre, H., Wilson, R. J., and Yamashita, Y.: Decline and recovery of total column ozone using a multimodel time series analysis, *J. Geophys. Res.*, 115, D00M10, doi:10.1029/2010JD013857, 2010.
- Barnes, E. A., Barnes, N. W., and Polvani, L. M.: Delayed Southern Hemisphere climate change induced by stratospheric ozone recovery, as projected by the CMIP5 models, *J. Climate*, 27, 852–867, doi:10.1175/JCLI-D-13-00246.1, 2013.
- Bodeker, G. E., Shiona, H., and Eskes, H.: Indicators of Antarctic ozone depletion, *Atmos. Chem. Phys.*, 5, 2603–2615, doi:10.5194/acp-5-2603-2005, 2005.
- Bodeker, G. E., Hassler, B., Young, P. J., and Portmann, R. W.: A vertically resolved, global, gap-free ozone database for assessing or constraining global climate model simulations, *Earth Syst. Sci. Data*, 5, 31–43, doi:10.5194/essd-5-31-2013, 2013.
- Bönisch, H., Engel, A., Birner, Th., Hoor, P., Tarasick, D. W., and Ray, E. A.: On the structural changes in the Brewer–Dobson Circulation after 2000, *Atmos. Chem. Phys.*, 11, 3937–3948, doi:10.5194/acp-11-3937-2011, 2011.
- Bowman, K. W., Shindell, D. T., Worden, H. M., Lamarque, J. F., Young, P. J., Stevenson, D. S., Qu, Z., de la Torre, M., Bergmann, D., Cameron-Smith, P. J., Collins, W. J., Doherty, R., Dalsøren, S. B., Faluvegi, G., Folberth, G., Horowitz, L. W., Josse, B. M., Lee, Y. H., MacKenzie, I. A., Myhre, G., Nagashima, T., Naik, V., Plummer, D. A., Rumbold, S. T., Skeie, R. B., Strode, S. A., Sudo, K., Szopa, S., Voulgarakis, A., Zeng, G., Kulawik, S. S., Aghedo, A. M., and Worden, J. R.: Evaluation of ACCMIP outgoing longwave radiation from

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- tropospheric ozone using TES satellite observations, *Atmos. Chem. Phys.*, 13, 4057–4072, doi:10.5194/acp-13-4057-2013, 2013.
- Butchart, N.: The Brewer–Dobson Circulation, *Rev. Geophys.*, 52, 157–184, doi:10.1002/2013RG000448, 2014.
- Butchart, N., Scaife, A. A., Bourqui, M., de Grandpré, J., Hare, S. H. E., Kettleborough, J., Langematz, U., Manzini, E., Sassi, F., Shibata, K., Shindell, D., and Sigmond, M.: Simulations of anthropogenic change in the strength of the Brewer–Dobson Circulation, *Clim. Dynam.*, 27, 727–741, doi:10.1007/s00382-006-0162-4, 2006.
- Butchart, N., Cionni, I., Eyring, V., Shepherd, T. G., Waugh, D. W., Akiyoshi, H., Austin, J., Brühl, C., Chipperfield, M. P., Cordero, E., Dameris, M., Deckert, R., Dhomse, S., Frith, S. M., Garcia, R. R., Gettelman, A., Giorgetta, M. A., Kinnison, D. E., Li, F., Mancini, E., McLandress, C., Pawson, S., Pitari, G., Plummer, D. A., Rozanov, E., Sassi, F., Scinocca, J. F., Shibata, K., Steil, B., and Tian, W.: Chemistry–Climate Model simulations of twenty-first century stratospheric climate and circulation changes, *J. Climate*, 23, 5349–5374, doi:10.1175/2010JCLI3404.1, 2010.
- Butchart, N., Charlton-Perez, A. J., Cionni, I., Hardiman, S. C., Haynes, P. H., Krüger, K., Kushner, P. J., Newman, P. A., Osprey, S. M., Perlitz, J., Sigmond, M., Wang, L., Akiyoshi, H., Austin, J., Bekki, S., Baumgaertner, A., Braesicke, P., Brühl, C., Chipperfield, M., Dameris, M., Dhomse, S., Eyring, V., Garcia, R., Garny, H., Jöckel, P., Lamarque, J. F., Marchand, M., Michou, M., Morgenstern, O., Nakamura, T., Pawson, S., Plummer, D., Pyle, J., Rozanov, E., Scinocca, J., Shepherd, T. G., Shibata, K., Smale, D., Teyssèdre, H., Tian, W., Waugh, D., and Yamashita, Y.: Multimodel climate and variability of the stratosphere, *J. Geophys. Res.*, 116, D05102, doi:10.1029/2010JD014995, 2011.
- Calvo, N., Garcia, R. R., Marsh, D. R., Mills, M. J., Kinnison, D. E., and Young, P. J.: Reconciling modeled and observed temperature trends over Antarctica, *Geophys. Res. Lett.*, 39, L16803, doi:10.1029/2012gl052526, 2012.
- Charlton-Perez, A. J., Hawkins, E., Eyring, V., Cionni, I., Bodeker, G. E., Kinnison, D. E., Akiyoshi, H., Frith, S. M., Garcia, R., Gettelman, A., Lamarque, J. F., Nakamura, T., Pawson, S., Yamashita, Y., Bekki, S., Braesicke, P., Chipperfield, M. P., Dhomse, S., Marchand, M., Mancini, E., Morgenstern, O., Pitari, G., Plummer, D., Pyle, J. A., Rozanov, E., Scinocca, J., Shibata, K., Shepherd, T. G., Tian, W., and Waugh, D. W.: The potential to narrow uncertainty in projections of stratospheric ozone over the 21st century, *Atmos. Chem. Phys.*, 10, 9473–9486, doi:10.5194/acp-10-9473-2010, 2010.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Charlton-Perez, A. J., Baldwin, M. P., Birner, T., Black, R. X., Butler, A. H., Calvo, N., Davis, N. A., Gerber, E. P., Gillett, N., Hardiman, S., Kim, J., Krüger, K., Lee, Y.-Y., Manzini, E., McDaniel, B. A., Polvani, L., Reichler, T., Shaw, T. A., Sigmond, M., Son, S.-W., Toohey, M., Wilcox, L., Yoden, S., Christiansen, B., Lott, F., Shindell, D., Yukimoto, S., and Watanabe, S.: On the lack of stratospheric dynamical variability in low-top versions of the CMIP5 models, *J. Geophys. Res.*, 118, 2494–2505, doi:10.1002/jgrd.50125, 2013.
- Christy, J. R., Spencer, R. W., Norris, W. B., Braswell, W. D., and Parker, D. E.: Error estimates of version 5.0 of MSU–AMSU bulk atmospheric temperatures, *J. Atmos. Ocean. Tech.*, 20, 613–629, doi:10.1175/1520-0426(2003)20<613:EEOVOM>2.0.CO;2, 2003.
- Cionni, I., Eyring, V., Lamarque, J. F., Randel, W. J., Stevenson, D. S., Wu, F., Bodeker, G. E., Shepherd, T. G., Shindell, D. T., and Waugh, D. W.: Ozone database in support of CMIP5 simulations: results and corresponding radiative forcing, *Atmos. Chem. Phys.*, 11, 11267–11292, doi:10.5194/acp-11-11267-2011, 2011.
- Collins, W. J., Derwent, R. G., Garnier, B., Johnson, C. E., Sanderson, M. G., and Stevenson, D. S.: Effect of stratosphere-troposphere exchange on the future tropospheric ozone trend, *J. Geophys. Res.*, 108, 8528, doi:10.1029/2002JD002617, 2003.
- Collins, W. J., Bellouin, N., Doutriaux-Boucher, M., Gedney, N., Halloran, P., Hinton, T., Hughes, J., Jones, C. D., Joshi, M., Liddicoat, S., Martin, G., O'Connor, F., Rae, J., Senior, C., Sitch, S., Totterdell, I., Wiltshire, A., and Woodward, S.: Development and evaluation of an Earth-System model – HadGEM2, *Geosci. Model Dev.*, 4, 1051–1075, doi:10.5194/gmd-4-1051-2011, 2011.
- Crook, J. A., Gillett, N. P., and Keeley, S. P. E.: Sensitivity of Southern Hemisphere climate to zonal asymmetry in ozone, *Geophys. Res. Lett.*, 35, L07806, doi:10.1029/2007GL032698, 2008.
- Dietmüller, S., Ponater, M., and Sausen, R.: Interactive ozone induces a negative feedback in CO₂-driven climate change simulations, *J. Geophys. Res.*, 119, 1796–1805, doi:10.1002/2013JD020575, 2014.
- Donner, L. J., Wyman, B. L., Hemler, R. S., Horowitz, L. W., Ming, Y., Zhao, M., Golaz, J.-C., Ginoux, P., Lin, S. J., Schwarzkopf, M. D., Austin, J., Alaka, G., Cooke, W. F., Delworth, T. L., Freidenreich, S. M., Gordon, C. T., Griffies, S. M., Held, I. M., Hurlin, W. J., Klein, S. A., Knutson, T. R., Langenhorst, A. R., Lee, H.-C., Lin, Y., Magi, B. I., Malyshev, S. L., Milly, P. C. D., Naik, V., Nath, M. J., Pincus, R., Poshay, J. J., Ramaswamy, V., Seman, C. J., Shevliakova, E., Sirutis, J. J., Stern, W. F., Stouffer, R. J., Wilson, R. J., Winton, M., Witten-

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀
Back

▶
Close

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Stratospheric ozone change and related climate impacts over 1850–2100

F. Iglesias-Suarez et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



berg, A. T., and Zeng, F.: The dynamical core, physical parameterizations, and basic simulation characteristics of the atmospheric component AM3 of the GFDL Global Coupled Model CM3, *J. Climate*, 24, 3484–3519, doi:10.1175/2011JCLI3955.1, 2011.

Dvortsov, V. L. and Solomon, S.: Response of the stratospheric temperatures and ozone to past and future increases in stratospheric humidity, *J. Geophys. Res.*, 106, 7505–7514, doi:10.1029/2000JD900637, 2001.

Engel, A., Mobius, T., Bonisch, H., Schmidt, U., Heinz, R., Levin, I., Atlas, E., Aoki, S., Nakazawa, T., Sugawara, S., Moore, F., Hurst, D., Elkins, J., Schauffler, S., Andrews, A., and Boering, K.: Age of stratospheric air unchanged within uncertainties over the past 30 years, *Nature Geosci.*, 2, 28–31, doi:10.1038/ngeo388, 2009.

Eyring, V., Butchart, N., Waugh, D. W., Akiyoshi, H., Austin, J., Bekki, S., Bodeker, G. E., Boville, B. A., Brühl, C., Chipperfield, M. P., Cordero, E., Dameris, M., Deushi, M., Fioletov, V. E., Frith, S. M., Garcia, R. R., Gettelman, A., Giorgetta, M. A., Grewe, V., Jourdain, L., Kinnison, D. E., Mancini, E., Manzini, E., Marchand, M., Marsh, D. R., Nagashima, T., Newman, P. A., Nielsen, J. E., Pawson, S., Pitari, G., Plummer, D. A., Rozanov, E., Schraner, M., Shepherd, T. G., Shibata, K., Stolarski, R. S., Struthers, H., Tian, W., and Yoshiki, M.: Assessment of temperature, trace species, and ozone in chemistry-climate model simulations of the recent past, *J. Geophys. Res.*, 111, D22308, doi:10.1029/2006JD007327, 2006.

Eyring, V., Waugh, D. W., Bodeker, G. E., Cordero, E., Akiyoshi, H., Austin, J., Beagley, S. R., Boville, B. A., Braesicke, P., Brühl, C., Butchart, N., Chipperfield, M. P., Dameris, M., Deckert, R., Deushi, M., Frith, S. M., Garcia, R. R., Gettelman, A., Giorgetta, M. A., Kinnison, D. E., Mancini, E., Manzini, E., Marsh, D. R., Matthes, S., Nagashima, T., Newman, P. A., Nielsen, J. E., Pawson, S., Pitari, G., Plummer, D. A., Rozanov, E., Schraner, M., Scinocca, J. F., Semeniuk, K., Shepherd, T. G., Shibata, K., Steil, B., Stolarski, R. S., Tian, W., and Yoshiki, M.: Multimodel projections of stratospheric ozone in the 21st century, *J. Geophys. Res.*, 112, D16303, doi:10.1029/2006JD008332, 2007.

Eyring, V., Cionni, I., Bodeker, G. E., Charlton-Perez, A. J., Kinnison, D. E., Scinocca, J. F., Waugh, D. W., Akiyoshi, H., Bekki, S., Chipperfield, M. P., Dameris, M., Dhomse, S., Frith, S. M., Garny, H., Gettelman, A., Kubin, A., Langematz, U., Mancini, E., Marchand, M., Nakamura, T., Oman, L. D., Pawson, S., Pitari, G., Plummer, D. A., Rozanov, E., Shepherd, T. G., Shibata, K., Tian, W., Braesicke, P., Hardiman, S. C., Lamarque, J. F., Morgenstern, O., Pyle, J. A., Smale, D., and Yamashita, Y.: Multi-model assessment of strato-

spheric ozone return dates and ozone recovery in CCMVal-2 models, *Atmos. Chem. Phys.*, 10, 9451–9472, doi:10.5194/acp-10-9451-2010, 2010a.

Eyring, V., Cionni, I., Lamarque, J. F., Akiyoshi, H., Bodeker, G. E., Charlton-Perez, A. J.,
Frith, S. M., Gettelman, A., Kinnison, D. E., Nakamura, T., Oman, L. D., Pawson, S., and
Yamashita, Y.: Sensitivity of 21st century stratospheric ozone to greenhouse gas scenarios,
Geophys. Res. Lett., 37, L16807, doi:10.1029/2010GL044443, 2010b.

Eyring, V., Arblaster, J. M., Cionni, I., Sedláček, J., Perlitz, J., Young, P. J., Bekki, S.,
Bergmann, D., Cameron-Smith, P., Collins, W. J., Faluvegi, G., Gottschaldt, K. D.,
Horowitz, L. W., Kinnison, D. E., Lamarque, J. F., Marsh, D. R., Saint-Martin, D., Shin-
dell, D. T., Sudo, K., Szopa, S., and Watanabe, S.: Long-term ozone changes and
associated climate impacts in CMIP5 simulations, *J. Geophys. Res.*, 118, 5029–5060,
doi:10.1002/jgrd.50316, 2013.

Forster, P. M., Thompson, D. W. J., Baldwin, M. P., Chipperfield, M. P., Dameris, M., Haigh, J.
D., Karoly, D. J., Kushner, P. J., Randel, W. J., Rosenlof, K. H., Seidel, D. J., Solomon, S.,
Beig, G., Braesicke, P., Butchart, N., Gillett, N. P., Grise, K. M., Marsh, D. R., McLandress, C.,
Rao, T. N., Son, S.-W., Stenchikov, G. L., and Yoden, S.: Stratospheric changes and climate,
Chapter 4 in Scientific Assessment of Ozone Depletion: 2010, Global Ozone Research and
Monitoring Project-Report No. 52, 516 p., Geneva, Switzerland, World Meteorological Orga-
nization, 2011.

Fu, Q., Solomon, S., and Lin, P.: On the seasonal dependence of tropical lower-stratospheric
temperature trends, *Atmos. Chem. Phys.*, 10, 2643–2653, doi:10.5194/acp-10-2643-2010,
2010.

Garcia, R. R. and Randel, W. J.: Acceleration of the Brewer–Dobson Circulation due to in-
creases in greenhouse gases, *J. Atmos. Sci.*, 65, 2731–2739, doi:10.1175/2008JAS2712.1,
2008.

Gent, P., Yeager, S., Neale, R., Levis, S., and Bailey, D.: Improvements in a half degree at-
mosphere/land version of the CCSM, *Clim. Dynam.*, 34, 819–833, doi:10.1007/s00382-009-
0614-8, 2010.

Gillett, N. P. and Fyfe, J. C.: Annular mode changes in the CMIP5 simulations, *Geophys. Res.
Lett.*, 40, 1189–1193, doi:10.1002/grl.50249, 2013.

Gillett, N. P. and Thompson, D. W. J.: Simulation of recent Southern Hemisphere climate
change, *Science*, 302, 273–275, doi:10.1126/science.1087440, 2003.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)





- Gillett, N. P., Scinocca, J. F., Plummer, D. A., and Reader, M. C.: Sensitivity of climate to dynamically-consistent zonal asymmetries in ozone, *Geophys. Res. Lett.*, 36, L10809, doi:10.1029/2009GL037246, 2009.

Gillett, N. P., Akiyoshi, H., Bekki, S., Braesicke, P., Eyring, V., Garcia, R., Karpechko, A. Yu., McLinden, C. A., Morgenstern, O., Plummer, D. A., Pyle, J. A., Rozanov, E., Scinocca, J., and Shibata, K.: Attribution of observed changes in stratospheric ozone and temperature, *Atmos. Chem. Phys.*, 11, 599–609, doi:10.5194/acp-11-599-2011, 2011.

Gong, D. and Wang, S.: Definition of Antarctic Oscillation index, *Geophys. Res. Lett.*, 26, 459–462, doi:10.1029/1999GL900003, 1999.

Gregg, J. W., Jones, C. G., and Dawson, T. E.: Urbanization effects on tree growth in the vicinity of New York City, *Nature*, 424, 183–187, doi:10.1038/nature01728, 2003.

Griffies, S. M., Winton, M., Donner, L. J., Horowitz, L. W., Downes, S. M., Farneti, R., Gnanadesikan, A., Hurlin, W. J., Lee, H.-C., Liang, Z., Palter, J. B., Samuels, B. L., Wittenberg, A. T., Wyman, B. L., Yin, J., and Zadeh, N.: The GFDL CM3 Coupled Climate Model: characteristics of the ocean and sea ice simulations, *J. Climate*, 24, 3520–3544, doi:10.1175/2011JCLI3964.1, 2011.

Grise, K. M., Thompson, D. W. J., and Forster, P. M.: On the role of radiative processes in stratosphere–troposphere coupling, *J. Climate*, 22, 4154–4161, doi:10.1175/2009JCLI2756.1, 2009.

Grytsai, A. V., Evtushevsky, O. M., Agapitov, O. V., Klekociuk, A. R., and Milinevsky, G. P.: Structure and long-term change in the zonal asymmetry in Antarctic total ozone during spring, *Ann. Geophys.*, 25, 361–374, doi:10.5194/angeo-25-361-2007, 2007.

Haigh, J. D. and Pyle, J. A.: Ozone perturbation experiments in a two-dimensional circulation model, *Q. J. Roy. Meteor. Soc.*, 108, 551–574, doi:10.1002/qj.49710845705, 1982.

Haimberger, L., Tavolato, C., and Sperka, S.: Toward elimination of the warm bias in historic radiosonde temperature records – some new results from a comprehensive intercomparison of upper-air data, *J. Climate*, 21, 4587–4606, doi:10.1175/2008JCLI1929.1, 2008.

Haimberger, L., Tavolato, C., and Sperka, S.: Homogenization of the Global Radiosonde Temperature Dataset through combined comparison with reanalysis background series and neighboring stations, *J. Climate*, 25, 8108–8131, doi:10.1175/JCLI-D-11-00668.1, 2012.

Hassler, B., Young, P. J., Portmann, R. W., Bodeker, G. E., Daniel, J. S., Rosenlof, K. H., and Solomon, S.: Comparison of three vertically resolved ozone data sets: climatology, trends

**Stratospheric ozone
change and related
climate impacts over
1850–2100**

F. Iglesias-Suarez et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Lamarque, J.-F. and Solomon, S.: Impact of changes in climate and halocarbons on recent lower stratosphere ozone and temperature trends, *J. Climate*, 23, 2599–2611, doi:10.1175/2010JCLI3179.1, 2010.

5 Lamarque, J.-F., Kyle, G. P., Meinshausen, M., Riahi, K., Smith, S., van Vuuren, D., Conley, A., and Vitt, F.: Global and regional evolution of short-lived radiatively-active gases and aerosols in the representative concentration pathways, *Climatic Change*, 109, 191–212, doi:10.1007/s10584-011-0155-0, 2011.

10 Lamarque, J.-F., Emmons, L. K., Hess, P. G., Kinnison, D. E., Tilmes, S., Vitt, F., Heald, C. L., Holland, E. A., Lauritzen, P. H., Neu, J., Orlando, J. J., Rasch, P. J., and Tyndall, G. K.: CAM-chem: description and evaluation of interactive atmospheric chemistry in the Community Earth System Model, *Geosci. Model Dev.*, 5, 369–411, doi:10.5194/gmd-5-369-2012, 2012.

15 Lamarque, J.-F., Dentener, F., McConnell, J., Ro, C.-U., Shaw, M., Vet, R., Bergmann, D., Cameron-Smith, P., Dalsoren, S., Doherty, R., Faluvegi, G., Ghan, S. J., Josse, B., Lee, Y. H., MacKenzie, I. A., Plummer, D., Shindell, D. T., Skeie, R. B., Stevenson, D. S., Strode, S., Zeng, G., Curran, M., Dahl-Jensen, D., Das, S., Fritzsche, D., and Nolan, M.: Multi-model mean nitrogen and sulfur deposition from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP): evaluation of historical and projected future changes, *Atmos. Chem. Phys.*, 13, 7997–8018, doi:10.5194/acp-13-7997-2013, 2013a.

20 Lamarque, J.-F., Shindell, D. T., Josse, B., Young, P. J., Cionni, I., Eyring, V., Bergmann, D., Cameron-Smith, P., Collins, W. J., Doherty, R., Dalsoren, S., Faluvegi, G., Folberth, G., Ghan, S. J., Horowitz, L. W., Lee, Y. H., MacKenzie, I. A., Nagashima, T., Naik, V., Plummer, D., Righi, M., Rumbold, S. T., Schulz, M., Skeie, R. B., Stevenson, D. S., Strode, S., Sudo, K., Szopa, S., Voulgarakis, A., and Zeng, G.: The Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP): overview and description of models, simulations and climate diagnostics, *Geosci. Model Dev.*, 6, 179–206, doi:10.5194/gmd-6-179-2013, 2013b.

25 Lary, D. J.: Catalytic destruction of stratospheric ozone, *J. Geophys. Res.*, 102, 21515–21526, 1997.

30 Lee, Y. H., Lamarque, J.-F., Flanner, M. G., Jiao, C., Shindell, D. T., Berntsen, T., Bisiaux, M. M., Cao, J., Collins, W. J., Curran, M., Edwards, R., Faluvegi, G., Ghan, S., Horowitz, L. W., McConnell, J. R., Ming, J., Myhre, G., Nagashima, T., Naik, V., Rumbold, S. T., Skeie, R. B., Sudo, K., Takemura, T., Thevenon, F., Xu, B., and Yoon, J.-H.: Evaluation of preindustrial to present-day black carbon and its albedo forcing from Atmospheric Chemistry and

Stratospheric ozone change and related climate impacts over 1850–2100

F. Iglesias-Suarez et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- perfield, M. P., Cugnet, D., Dameris, M., Dhomse, S., Frith, S. M., Garny, H., Gettelman, A., Hardiman, S. C., Hegglin, M. I., Jöckel, P., Kinnison, D. E., Lamarque, J. F., Mancini, E., Manzini, E., Marchand, M., Michou, M., Nakamura, T., Nielsen, J. E., Olivié, D., Pitari, G., Plummer, D. A., Rozanov, E., Scinocca, J. F., Smale, D., Teyssèdre, H., Toohey, M., Tian, W., and Yamashita, Y.: Review of the formulation of present-generation stratospheric chemistry-climate models and associated external forcings, *J. Geophys. Res.*, 115, D00M02, doi:10.1029/2009JD013728, 2010.
- Myhre, G., Shindell, D., Breton, F.-M., Collins, W., Fuglestvedt, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T., and Zhang, H.: Anthropogenic and natural radiative forcing, in: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V. and Midgley, P. M., Cambridge University Press, Cambridge, UK and New York, NY, USA, 659–740, 2013.
- Naik, V., Horowitz, L. W., Fiore, A. M., Ginoux, P., Mao, J., Aghedo, A. M., and Levy, H.: Impact of preindustrial to present-day changes in short-lived pollutant emissions on atmospheric composition and climate forcing, *J. Geophys. Res.*, 118, 8086–8110, doi:10.1002/jgrd.50608, 2013a.
- Naik, V., Voulgarakis, A., Fiore, A. M., Horowitz, L. W., Lamarque, J.-F., Lin, M., Prather, M. J., Young, P. J., Bergmann, D., Cameron-Smith, P. J., Cionni, I., Collins, W. J., Dalsøren, S. B., Doherty, R., Eyring, V., Faluvegi, G., Folberth, G. A., Josse, B., Lee, Y. H., MacKenzie, I. A., Nagashima, T., van Noije, T. P. C., Plummer, D. A., Righi, M., Rumbold, S. T., Skeie, R., Shindell, D. T., Stevenson, D. S., Strode, S., Sudo, K., Szopa, S., and Zeng, G.: Preindustrial to present-day changes in tropospheric hydroxyl radical and methane lifetime from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmos. Chem. Phys.*, 13, 5277–5298, doi:10.5194/acp-13-5277-2013, 2013b.
- Nedoluha, G. E., Siskind, D. E., Lambert, A., and Boone, C.: The decrease in mid-stratospheric tropical ozone since 1991, *Atmos. Chem. Phys.*, 15, 4215–4224, doi:10.5194/acp-15-4215-2015, 2015.
- Newman, P. A., Nash, E. R., and Rosenfield, J. E.: What controls the temperature of the Arctic stratosphere during the spring?, *J. Geophys. Res.*, 106, 19999–20010, doi:10.1029/2000JD000061, 2001.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Nowack, P. J., Luke Abraham, N., Maycock, A. C., Braesicke, P., Gregory, J. M., Joshi, M. M., Osprey, A., and Pyle, J. A.: A large ozone-circulation feedback and its implications for global warming assessments, *Nature Clim. Change*, 5, 41–45, doi:10.1038/nclimate2451, 2015.
- Parrish, D. D., Lamarque, J.-F., Naik, V., Horowitz, L., Shindell, D. T., Staehelin, J., Derwent, R., Cooper, O. R., Tanimoto, H., Volz-Thomas, A., Gilge, S., Scheel, H.-E., Steinbacher, M., and Fröhlich, M.: Long-term changes in lower tropospheric baseline ozone concentrations: Comparing chemistry-climate models and observations at northern midlatitudes, *J. Geophys. Res. Atmos.*, 119, 5719–5736, doi:10.1002/2013JD021435, 2014.
- Perlwitz, J.: Atmospheric science: Tug of war on the jet stream, *Nature Clim. Change*, 1, 29–31, doi:10.1038/nclimate1065, 2011.
- Perlwitz, J., Pawson, S., Fogt, R. L., Nielsen, J. E., and Neff, W. D.: Impact of stratospheric ozone hole recovery on Antarctic climate, *Geophys. Res. Lett.*, 35, L08714, doi:10.1029/2008GL033317, 2008.
- Polvani, L. M. and Solomon, S.: The signature of ozone depletion on tropical temperature trends, as revealed by their seasonal cycle in model integrations with single forcings, *J. Geophys. Res.*, 117, D17102, doi:10.1029/2012JD017719, 2012.
- Polvani, L. M., Waugh, D. W., Correa, G. J. P., and Son, S.-W.: Stratospheric ozone depletion: the main driver of twentieth-century atmospheric circulation changes in the Southern Hemisphere, *J. Climate*, 24, 795–812, doi:10.1175/2010JCLI3772.1, 2010.
- Polvani, L. M., Previdi, M., and Deser, C.: Large cancellation, due to ozone recovery, of future Southern Hemisphere atmospheric circulation trends, *Geophys. Res. Lett.*, 38, L04707, doi:10.1029/2011GL046712, 2011.
- Portmann, R. W. and Solomon, S.: Indirect radiative forcing of the ozone layer during the 21st century, *Geophys. Res. Lett.*, 34, L02813, doi:10.1029/2006GL028252, 2007.
- Portmann, R. W., Daniel, J. S., and Ravishankara, A. R.: Stratospheric ozone depletion due to nitrous oxide: influences of other gases, *Philos. T. R. Soc. B*, 367, 1256–1264, doi:10.1098/rstb.2011.0377, 2012.
- Prather, M. J. and Ehhalt, D.: Atmospheric chemistry and greenhouse gases, in: *Climate Change 2001: The Scientific Basis*, Cambridge University Press, Cambridge, UK, 239–287, 2001.
- Previdi, M. and Polvani, L. M.: Climate system response to stratospheric ozone depletion and recovery, *Q. J. Roy. Meteorol. Soc.*, 140, 2401–2419, doi:10.1002/qj.2330, 2014.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Ramaswamy, V., Schwarzkopf, M. D., Randel, W. J., Santer, B. D., Soden, B. J., and Stenchikov, G. L.: Anthropogenic and natural influences in the evolution of lower stratospheric cooling, *Science*, 311, 1138–1141, doi:10.1126/science.1122587, 2006.
- Randel, W. J. and Wu, F.: Cooling of the Arctic and Antarctic polar stratospheres due to ozone depletion, *J. Climate*, 12, 1467–1479, doi:10.1175/1520-0442(1999)012<1467:COTAAA>2.0.CO;2, 1999.
- Randel, W. J., Shine, K. P., Austin, J., Barnett, J., Claud, C., Gillett, N. P., Keckhut, P., Langematz, U., Lin, R., Long, C., Mears, C., Miller, A., Nash, J., Seidel, D. J., Thompson, D. W. J., Wu, F., and Yoden, S.: An update of observed stratospheric temperature trends, *J. Geophys. Res.*, 114, D02107, doi:10.1029/2008JD010421, 2009.
- Randeniya, L. K., Vohralik, P. F., and Plumb, I. C.: Stratospheric ozone depletion at northern mid latitudes in the 21st century: the importance of future concentrations of greenhouse gases nitrous oxide and methane, *Geophys. Res. Lett.*, 29, 10-11–10-14, doi:10.1029/2001GL014295, 2002.
- Ravishankara, A. R., Daniel, J. S., and Portmann, R. W.: Nitrous oxide (N_2O): the dominant ozone-depleting substance emitted in the 21st century, *Science*, 326, 123–125, doi:10.1126/science.1176985, 2009.
- Reader, M. C., Plummer, D. A., Scinocca, J. F., and Shepherd, T. G.: Contributions to twentieth century total column ozone change from halocarbons, tropospheric ozone precursors, and climate change, *Geophys. Res. Lett.*, 40, 6276–6281, doi:10.1002/2013GL057776, 2013.
- Revell, L. E., Bodeker, G. E., Smale, D., Lehmann, R., Huck, P. E., Williamson, B. E., Rozanov, E., and Struthers, H.: The effectiveness of N_2O in depleting stratospheric ozone, *Geophys. Res. Lett.*, 39, L15806, doi:10.1029/2012GL052143, 2012.
- Rosenfield, J. E., Douglass, A. R., and Considine, D. B.: The impact of increasing carbon dioxide on ozone recovery, *J. Geophys. Res.*, 107, ACH 7-1–ACH 7-9, doi:10.1029/2001JD000824, 2002.
- Santer, B. D., Wigley, T. M. L., Boyle, J. S., Gaffen, D. J., Hnilo, J. J., Nychka, D., Parker, D. E., and Taylor, K. E.: Statistical significance of trends and trend differences in layer-average atmospheric temperature time series, *J. Geophys. Res.*, 105, 7337–7356, doi:10.1029/1999JD901105, 2000.
- Santer, B. D., Sausen, R., Wigley, T. M. L., Boyle, J. S., AchutaRao, K., Doutriaux, C., Hansen, J. E., Meehl, G. A., Roeckner, E., Ruedy, R., Schmidt, G., and Taylor, K. E.: Behavior

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

of tropopause height and atmospheric temperature in models, reanalyses, and observations: decadal changes, *J. Geophys. Res.*, 108, 4002, doi:10.1029/2002JD002258, 2003a.

5 Santer, B. D., Wehner, M. F., Wigley, T. M. L., Sausen, R., Meehl, G. A., Taylor, K. E., Ammann, C., Arblaster, J., Washington, W. M., Boyle, J. S., and Brüggemann, W.: Contributions of anthropogenic and natural forcing to recent tropopause height changes, *Science*, 301, 479–483, doi:10.1126/science.1084123, 2003b.

10 Schmidt, G. A., Ruedy, R., Hansen, J. E., Aleinov, I., Bell, N., Bauer, M., Bauer, S., Cairns, B., Canuto, V., Cheng, Y., Del Genio, A., Faluvegi, G., Friend, A. D., Hall, T. M., Hu, Y., Kelley, M., Kiang, N. Y., Koch, D., Lacis, A. A., Lerner, J., Lo, K. K., Miller, R. L., Nazarenko, L., Oinas, V., Perlitz, J., Perlitz, J., Rind, D., Romanou, A., Russell, G. L., Sato, M., Shin-dell, D. T., Stone, P. H., Sun, S., Tausnev, N., Thresher, D., and Yao, M.-S.: Present-day atmospheric simulations using GISS ModelE: comparison to in situ, satellite, and reanalysis data, *J. Climate*, 19, 153–192, doi:10.1175/JCLI3612.1, 2006.

15 Scinocca, J. F., McFarlane, N. A., Lazare, M., Li, J., and Plummer, D.: Technical Note: The CCCma third generation AGCM and its extension into the middle atmosphere, *Atmos. Chem. Phys.*, 8, 7055–7074, doi:10.5194/acp-8-7055-2008, 2008.

Sexton, D. M. H.: The effect of stratospheric ozone depletion on the phase of the Antarctic Oscillation, *Geophys. Res. Lett.*, 28, 3697–3700, doi:10.1029/2001GL013376, 2001.

20 Shepherd, T. G.: Dynamics, stratospheric ozone, and climate change, *Atmos.-Ocean*, 46, 117–138, doi:10.3137/ao.460106, 2008.

Shepherd, T. G., Plummer, D. A., Scinocca, J. F., Hegglin, M. I., Fioletov, V. E., Reader, M. C., Remsberg, E., von Clarmann, T., and Wang, H. J.: Reconciliation of halogen-induced ozone loss with the total-column ozone record, *Nat. Geosci.*, 7, 443–449, doi:10.1038/ngeo2155, 2014.

25 Sherwood, S. C., Meyer, C. L., Allen, R. J., and Titchner, H. A.: Robust tropospheric warming revealed by iteratively homogenized radiosonde data, *J. Climate*, 21, 5336–5352, doi:10.1175/2008JCLI2320.1, 2008.

30 Shindell, D. T. and Schmidt, G. A.: Southern Hemisphere climate response to ozone changes and greenhouse gas increases, *Geophys. Res. Lett.*, 31, L18209, doi:10.1029/2004GL020724, 2004.

Shindell, D. T., Faluvegi, G., Stevenson, D. S., Krol, M. C., Emmons, L. K., Lamarque, J. F., Pétron, G., Dentener, F. J., Ellingsen, K., Schultz, M. G., Wild, O., Amann, M., Atherton, C. S., Bergmann, D. J., Bey, I., Butler, T., Cofala, J., Collins, W. J., Derwent, R. G., Doherty, R. M.,

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Drevet, J., Eskes, H. J., Fiore, A. M., Gauss, M., Hauglustaine, D. A., Horowitz, L. W., Isaksen, I. S. A., Lawrence, M. G., Montanaro, V., Müller, J. F., Pitari, G., Prather, M. J., Pyle, J. A., Rast, S., Rodriguez, J. M., Sanderson, M. G., Savage, N. H., Strahan, S. E., Sudo, K., Szopa, S., Unger, N., van Noije, T. P. C., and Zeng, G.: Multimodel simulations of carbon monoxide: comparison with observations and projected near-future changes, *J. Geophys. Res.*, 111, D19306, doi:10.1029/2006JD007100, 2006.

Shindell, D. T., Lamarque, J.-F., Schulz, M., Flanner, M., Jiao, C., Chin, M., Young, P. J., Lee, Y. H., Rotstayn, L., Mahowald, N., Milly, G., Faluvegi, G., Balkanski, Y., Collins, W. J., Conley, A. J., Dalsoren, S., Easter, R., Ghan, S., Horowitz, L., Liu, X., Myhre, G., Naganashima, T., Naik, V., Rumbold, S. T., Skeie, R., Sudo, K., Szopa, S., Takemura, T., Voulgarakis, A., Yoon, J.-H., and Lo, F.: Radiative forcing in the ACCMIP historical and future climate simulations, *Atmos. Chem. Phys.*, 13, 2939–2974, doi:10.5194/acp-13-2939-2013, 2013a.

Shindell, D. T., Pechony, O., Voulgarakis, A., Faluvegi, G., Nazarenko, L., Lamarque, J.-F., Bowman, K., Milly, G., Kovari, B., Ruedy, R., and Schmidt, G. A.: Interactive ozone and methane chemistry in GISS-E2 historical and future climate simulations, *Atmos. Chem. Phys.*, 13, 2653–2689, doi:10.5194/acp-13-2653-2013, 2013b.

Sigmond, M., Fyfe, J. C., and Scinocca, J. F.: Does the ocean impact the atmospheric response to stratospheric ozone depletion?, *Geophys. Res. Lett.*, 37, L12706, doi:10.1029/2010GL043773, 2010.

Solomon, S., Young, P. J., and Hassler, B.: Uncertainties in the evolution of stratospheric ozone and implications for recent temperature changes in the tropical lower stratosphere, *Geophys. Res. Lett.*, 39, L17706, doi:10.1029/2012gl052723, 2012.

Son, S. W., Polvani, L. M., Waugh, D. W., Akiyoshi, H., Garcia, R., Kinnison, D., Pawson, S., Rozanov, E., Shepherd, T. G., and Shibata, K.: The impact of stratospheric ozone recovery on the Southern Hemisphere westerly jet, *Science*, 320, 1486–1489, doi:10.1126/science.1155939, 2008.

Son, S.-W., Tandon, N. F., Polvani, L. M., and Waugh, D. W.: Ozone hole and Southern Hemisphere climate change, *Geophys. Res. Lett.*, 36, L15705, doi:10.1029/2009GL038671, 2009.

Son, S. W., Gerber, E. P., Perlitz, J., Polvani, L. M., Gillett, N. P., Seo, K. H., Eyring, V., Shepherd, T. G., Waugh, D., Akiyoshi, H., Austin, J., Baumgaertner, A., Bekki, S., Braesicke, P., Brühl, C., Butchart, N., Chipperfield, M. P., Cugnet, D., Dameris, M., Dhomse, S., Frith, S.,

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)
[▶](#)[▶](#)
[◀](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Garny, H., Garcia, R., Hardiman, S. C., Jöckel, P., Lamarque, J. F., Mancini, E., Marchand, M., Michou, M., Nakamura, T., Morgenstern, O., Pitari, G., Plummer, D. A., Pyle, J., Rozanov, E., Scinocca, J. F., Shibata, K., Smale, D., Teyssèdre, H., Tian, W., and Yamashita, Y.: Impact of stratospheric ozone on Southern Hemisphere circulation change: A multimodel assessment, *J. Geophys. Res.*, 115, D00M07, doi:10.1029/2010JD014271, 2010.

SPARC CCMVal: SPARC CCMVal Report on the Evaluation of Chemistry-Climate Models, edited by: Eyring, V., Shepherd, T. G., and Waugh, D. W., SPARC Report No. 5, WCRP-132, WMO/TD-No. 1526, 2010.

Stevenson, D. S.: Atmospheric chemistry: climate's chemical sensitivity, *Nature Clim. Change*, 5, 21–22, doi:10.1038/nclimate2477, 2015.

Stevenson, D. S., Dentener, F. J., Schultz, M. G., Ellingsen, K., van Noije, T. P. C., Wild, O., Zeng, G., Amann, M., Atherton, C. S., Bell, N., Bergmann, D. J., Bey, I., Butler, T., Cofala, J., Collins, W. J., Derwent, R. G., Doherty, R. M., Drevet, J., Eskes, H. J., Fiore, A. M., Gauss, M., Hauglustaine, D. A., Horowitz, L. W., Isaksen, I. S. A., Krol, M. C., Lamarque, J. F., Lawrence, M. G., Montanaro, V., Müller, J. F., Pitari, G., Prather, M. J., Pyle, J. A., Rast, S., Rodriguez, J. M., Sanderson, M. G., Savage, N. H., Shindell, D. T., Strahan, S. E., Sudo, K., and Szopa, S.: Multimodel ensemble simulations of present-day and near-future tropospheric ozone, *J. Geophys. Res.*, 111, D08301, doi:10.1029/2005JD006338, 2006.

Stevenson, D. S., Young, P. J., Naik, V., Lamarque, J.-F., Shindell, D. T., Voulgarakis, A., Skeie, R. B., Dalsoren, S. B., Myhre, G., Berntsen, T. K., Folberth, G. A., Rumbold, S. T., Collins, W. J., MacKenzie, I. A., Doherty, R. M., Zeng, G., van Noije, T. P. C., Strunk, A., Bergmann, D., Cameron-Smith, P., Plummer, D. A., Strode, S. A., Horowitz, L., Lee, Y. H., Szopa, S., Sudo, K., Nagashima, T., Josse, B., Cionni, I., Righi, M., Eyring, V., Conley, A., Bowman, K. W., Wild, O., and Archibald, A.: Tropospheric ozone changes, radiative forcing and attribution to emissions in the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmos. Chem. Phys.*, 13, 3063–3085, doi:10.5194/acp-13-3063-2013, 2013.

Stiller, G. P., von Clarmann, T., Haenel, F., Funke, B., Glatthor, N., Grabowski, U., Kellmann, S., Kiefer, M., Linden, A., Lossow, S., and López-Puertas, M.: Observed temporal evolution of global mean age of stratospheric air for the 2002 to 2010 period, *Atmos. Chem. Phys.*, 12, 3311–3331, doi:10.5194/acp-12-3311-2012, 2012.

Struthers, H., Bodeker, G. E., Austin, J., Bekki, S., Cionni, I., Dameris, M., Giorgi, M. A., Grewe, V., Lefèvre, F., Lott, F., Manzini, E., Peter, T., Rozanov, E., and Schraner, M.: The

simulation of the Antarctic ozone hole by chemistry-climate models, *Atmos. Chem. Phys.*, 9, 6363–6376, doi:10.5194/acp-9-6363-2009, 2009.

Sudo, K., Takahashi, M., and Akimoto, H.: Future changes in stratosphere-troposphere exchange and their impacts on future tropospheric ozone simulations, *Geophys. Res. Lett.*, 30, 2256, doi:10.1029/2003GL018526, 2003.

Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An Overview of CMIP5 and the Experiment Design, *B. Am. Meteorol. Soc.*, 93, 485–498, doi:10.1175/BAMS-D-11-00094.1, 2012.

The HadGEM2 Development Team: G. M. Martin, Bellouin, N., Collins, W. J., Culverwell, I. D., Halloran, P. R., Hardiman, S. C., Hinton, T. J., Jones, C. D., McDonald, R. E., McLaren, A. J., O'Connor, F. M., Roberts, M. J., Rodriguez, J. M., Woodward, S., Best, M. J., Brooks, M. E., Brown, A. R., Butchart, N., Dearden, C., Derbyshire, S. H., Dharsi, I., Doutriaux-Boucher, M., Edwards, J. M., Falloon, P. D., Gedney, N., Gray, L. J., Hewitt, H. T., Hobson, M., Huddleston, M. R., Hughes, J., Ineson, S., Ingram, W. J., James, P. M., Johns, T. C., Johnson, C. E., Jones, A., Jones, C. P., Joshi, M. M., Keen, A. B., Liddicoat, S., Lock, A. P., Maidens, A. V., Manners, J. C., Milton, S. F., Rae, J. G. L., Ridley, J. K., Sellar, A., Senior, C. A., Totterdell, I. J., Verhoef, A., Vidale, P. L., and Wiltshire, A.: The HadGEM2 family of Met Office Unified Model climate configurations, *Geosci. Model Dev.*, 4, 723–757, doi:10.5194/gmd-4-723-2011, 2011.

Thompson, D. W. J. and Solomon, S.: Interpretation of recent Southern Hemisphere climate change, *Science*, 296, 895–899, doi:10.1126/science.1069270, 2002.

Thompson, D. W. J. and Wallace, J. M.: Annular modes in the Extratropical Circulation. Part I: month-to-month variability*, *J. Climate*, 13, 1000–1016, doi:10.1175/1520-0442(2000)013<1000:AMITEC>2.0.CO;2, 2000.

Thompson, D. W. J., Solomon, S., Kushner, P. J., England, M. H., Grise, K. M., and Karoly, D. J.: Signatures of the Antarctic ozone hole in Southern Hemisphere surface climate change, *Nature Geosci.*, 4, 741–749, doi:10.1038/ngeo1296, 2011.

Thorne, P. W., Parker, D. E., Tett, S. F. B., Jones, P. D., McCarthy, M., Coleman, H., and Brohan, P.: Revisiting radiosonde upper air temperatures from 1958 to 2002, *J. Geophys. Res.*, 110, D18105, doi:10.1029/2004JD005753, 2005.

UNEP: Environmental Effects of Ozone Depletion and its Interaction with Climate Change: 2015 Assessment, United Nations Environment Programme (UNEP), Nairobi, 2015.

van Vuuren, Detlef, P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G., Kram, T., Krey, V., Lamarque, J.-F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S.,

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- and Rose, S.: The representative concentration pathways: an overview, *Climatic Change*, 109, 5–31, doi:10.1007/s10584-011-0148-z, 2011.
- Voulgarakis, A., Naik, V., Lamarque, J.-F., Shindell, D. T., Young, P. J., Prather, M. J., Wild, O., Field, R. D., Bergmann, D., Cameron-Smith, P., Cionni, I., Collins, W. J., Dalsøren, S. B., Doherty, R. M., Eyring, V., Faluvegi, G., Folberth, G. A., Horowitz, L. W., Josse, B., MacKenzie, I. A., Nagashima, T., Plummer, D. A., Righi, M., Rumbold, S. T., Stevenson, D. S., Strode, S. A., Sudo, K., Szopa, S., and Zeng, G.: Analysis of present day and future OH and methane lifetime in the ACCMIP simulations, *Atmos. Chem. Phys.*, 13, 2563–2587, doi:10.5194/acp-13-2563-2013, 2013.
- Watanabe, S., Hajima, T., Sudo, K., Nagashima, T., Takemura, T., Okajima, H., Nozawa, T., Kawase, H., Abe, M., Yokohata, T., Ise, T., Sato, H., Kato, E., Takata, K., Emori, S., and Kawamiya, M.: MIROC-ESM 2010: model description and basic results of CMIP5-20c3m experiments, *Geosci. Model Dev.*, 4, 845–872, doi:10.5194/gmd-4-845-2011, 2011.
- Waugh, D. W., Oman, L., Kawa, S. R., Stolarski, R. S., Pawson, S., Douglass, A. R., Newman, P. A., and Nielsen, J. E.: Impacts of climate change on stratospheric ozone recovery, *Geophys. Res. Lett.*, 36, L03805, doi:10.1029/2008GL036223, 2009a.
- Waugh, D. W., Oman, L., Newman, P. A., Stolarski, R. S., Pawson, S., Nielsen, J. E., and Perlitz, J.: Effect of zonal asymmetries in stratospheric ozone on simulated Southern Hemisphere climate trends, *Geophys. Res. Lett.*, 36, L18701, doi:10.1029/2009GL040419, 2009b.
- Wilcox, L. J., Charlton-Perez, A. J., and Gray, L. J.: Trends in austral jet position in ensembles of high- and low-top CMIP5 models, *J. Geophys. Res.*, 117, D13115, doi:10.1029/2012JD017597, 2012.
- Wild, O.: Modelling the global tropospheric ozone budget: exploring the variability in current models, *Atmos. Chem. Phys.*, 7, 2643–2660, doi:10.5194/acp-7-2643-2007, 2007.
- WMO: Scientific Assessment of Ozone Depletion: 2006, World Meteorological Organization, Geneva, Switzerland, 572 pp., 2007.
- WMO: Scientific Assessment of Ozone Depletion: 2010, World Meteorological Organization, Geneva, Switzerland, 516 pp., 2011.
- WMO: Scientific Assessment of Ozone Depletion: 2014, World Meteorological Organization, Global Ozone Research and Monitoring Project, Geneva, Switzerland, 2014.
- Young, P. J., Rosenlof, K. H., Solomon, S., Sherwood, S. C., Fu, Q., and Lamarque, J.-F.: Changes in stratospheric temperatures and their implications for changes in the Brewer-

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Young, P. J., Archibald, A. T., Bowman, K. W., Lamarque, J.-F., Naik, V., Stevenson, D. S., Tilmes, S., Voulgarakis, A., Wild, O., Bergmann, D., Cameron-Smith, P., Cionni, I., Collins, W. J., Dalsøren, S. B., Doherty, R. M., Eyring, V., Faluvegi, G., Horowitz, L. W., Josse, B., Lee, Y. H., MacKenzie, I. A., Nagashima, T., Plummer, D. A., Righi, M., Rumbold, S. T., Skeie, R. B., Shindell, D. T., Strode, S. A., Sudo, K., Szopa, S., and Zeng, G.: Pre-industrial to end 21st century projections of tropospheric ozone from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmos. Chem. Phys.*, 13, 2063–2090, doi:10.5194/acp-13-2063-2013, 2013a.
- Young, P. J., Butler, A. H., Calvo, N., Haimberger, L., Kushner, P. J., Marsh, D. R., Randel, W. J., and Rosenlof, K. H.: Agreement in late twentieth century Southern Hemisphere stratospheric temperature trends in observations and CCMVal-2, CMIP3, and CMIP5 models, *J. Geophys. Res.*, 118, 605–613, doi:10.1002/jgrd.50126, 2013b.
- Young, P. J., Davis, S. M., Hassler, B., Solomon, S., and Rosenlof, K. H.: Modeling the climate impact of Southern Hemisphere ozone depletion: The importance of the ozone data set, *Geophys. Res. Lett.*, 41, 9033–9039, doi:10.1002/2014GL061738, 2014.
- Zeng, G. and Pyle, J. A.: Changes in tropospheric ozone between 2000 and 2100 modeled in a chemistry-climate model, *Geophys. Res. Lett.*, 30, 1392, doi:10.1029/2002GL016708, 2003.
- Zeng, G., Pyle, J. A., and Young, P. J.: Impact of climate change on tropospheric ozone and its global budgets, *Atmos. Chem. Phys.*, 8, 369–387, doi:10.5194/acp-8-369-2008, 2008.
- Zeng, G., Morgenstern, O., Braesicke, P., and Pyle, J. A.: Impact of stratospheric ozone recovery on tropospheric ozone and its budget, *Geophys. Res. Lett.*, 37, L09805, doi:10.1029/2010GL042812, 2010.
- Zou, C.-Z., Goldberg, M. D., Cheng, Z., Grody, N. C., Sullivan, J. T., Cao, C., and Tarpley, D.: Recalibration of microwave sounding unit for climate studies using simultaneous nadir over-passes, *J. Geophys. Res.*, 111, D19114, doi:10.1029/2005JD006798, 2006.
- Zou, C.-Z., Gao, M., and Goldberg, M. D.: Error structure and atmospheric temperature trends in observations from the Microwave Sounding Unit, *J. Climate*, 22, 1661–1681, doi:10.1175/2008JCLI2233.1, 2009.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)
[Back](#)

[▶](#)
[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Table 1. Summary of the ACCMIP models used here.

Model	Stratospheric ozone	Composition-radiation coupling	Photolysis scheme	Reference
CESM-CAM-superfast CMAM	CHEM	Yes	Adjusted look-up table	Lamarque et al. (2012)
	CHEM	Yes	Adjusted look-up table	Scinocca et al. (2008)
GFDL-AM3	CHEM	Yes	Adjusted look-up table	Donner et al. (2011); Naik et al. (2013a)
GISS-E2-R	CHEM	Yes	Online	Koch et al. (2006); Shindell et al. (2013b)
HadGEM2	NOCHEM	Yes	Look-up table + TCO overhead	Collins et al. (2011)
MIROC-CHEM	CHEM	Yes	Online	Watanabe et al. (2011)
NCAR-CAM3.5	CHEM	Yes	Adjusted look-up table	Lamarque et al. (2011, 2012)
UM-CAM	NOCHEM	No	Look-up table	Zeng et al. (2008, 2010)

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)**Table 2.** Global annual mean of TCO (DU).

Scenario	Year	ACCMIP*	CMIP5*	IGAC/SPARC
Hist	1850	294 ± 16	300 ± 19	293 ± 1
	1980	300 ± 19	306 ± 20	292 ± 2
	2000	291 ± 16	297 ± 20	281 ± 1
RCP2.6	2030	295 ± 16	301 ± 20	288 ± 1
	2100	297 ± 18	302 ± 20	294 ± 0
RCP8.5	2030	300 ± 17	306 ± 20	290 ± 1
	2100	316 ± 23	323 ± 11	304 ± 0

* For the historical period and the RCPs emission scenarios considered here as calculated from the CHEM models and the IGAC/SPARC data set (see Sect. 2). The multi-model mean is given along with uncertainties (± 1 standard deviation).

Stratospheric ozone change and related climate impacts over 1850–2100

F. Iglesias-Suarez et al.




Figure 1. Total column ozone trends from 1980 to 2000 (‰ dec^{-1}) for the annual mean (ANN) **(a)** global, **(b)** in the tropics, **(c)** in the northern midlatitudes, **(d)** in the southern midlatitudes, **(e)** for the boreal spring in the Arctic (MAM), and **(f)** for austral spring in the Antarctic (SON). The box, whiskers and line indicate the interquartile range, 95 % range and median respectively, for the ACCMIP (light grey), CMIP5 (dark grey) and CCMVal2 (magenta) models. Multi-model means are indicated by dots. CHEM (models with interactive chemistry) and NOCHEM (models that prescribe ozone) means are indicated by “plus” and “cross” symbols, respectively. Observations and IGAC/SPARC data sets are represented by error bars indicating the 95 % confidence intervals (one tail).

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**Stratospheric ozone
change and related
climate impacts over
1850–2100**

F. Iglesias-Suarez et al.




Figure 2. Vertically resolved ozone trends (‰ dec^{-1}), for ACCMIP multi-model mean, CHEM and NOCHEM models compared to BDBP Tier 1.4 (regression model fit with uncertainty estimates indicating 95 % confidence intervals, one tail) and Tier 0 (observations).

**Stratospheric ozone
change and related
climate impacts over
1850–2100**

F. Iglesias-Suarez et al.




Figure 3. Total column ozone (DU) time series from 1850 to 2100. The box, whiskers and line indicate the interquartile range, 95 % range and median respectively, for the ACCMIP CHEM models. In addition, the multi-model mean of the CMIP5 CHEM models and the IGAC/SPARC mean are shown. All data sets are normalised to Hist 1850 time slice levels.

**Stratospheric ozone
change and related
climate impacts over
1850–2100**

F. Iglesias-Suarez et al.




Figure 4. As Fig. 3, but for the upper stratosphere (10–1 hPa), lower stratosphere (150–15 hPa) and tropospheric columns ozone (DU) in the tropics.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Stratospheric ozone change and related climate impacts over 1850–2100

F. Iglesias-Suarez et al.




Figure 5. Vertically resolved ozone change between 2100 and 1850 (a to f), and 2100 and 2000 (g to l) time slices. (a–g) are for Arctic boreal spring mean, (b–h) and (d–j) for NH and SH midlatitudes annual mean respectively, (c–i) for tropical annual mean, (e–k) for Antarctic austral spring mean, and (f–l) for global annual mean.



Stratospheric ozone change and related climate impacts over 1850–2100

F. Iglesias-Suarez et al.




Figure 6. Temperature trends from 1980 to 2000 (K dec⁻¹). (a and c) represent MSU temperature lower stratosphere (TLS) for boreal spring in the Arctic (MAM) and for austral spring in the Antarctic (OND). The box, whiskers, line, dot, “plus” and “cross” symbols show the interquartile range, 95 % range, median, multi-model mean, CHEM and NOCHEM means respectively, for the ACCMIP (light grey), CMIP5 (dark grey) and CCMVal2 (magenta) models. (b and d) represent vertically resolved temperature (T) trends for the ACCMIP simulations (light grey). Observational data sets are represented by error bars indicating the 95 % confidence intervals (one tail).




Figure 7. (a) MSU temperature lower stratosphere (TLS) and (b) SAM index time series from 1850 to 2100. The box, whiskers, line, dot, “plus” and “cross” symbols show the interquartile range, 95 % range, median, multi-model mean, CHEM and NOCHEM means respectively, for the ACCMIP models. The five years average of the CMIP5 multi-model mean is shown. In addition, HadSLP2 observational data set for (b) is represented by a solid black line. The ACCMIP models are normalised to Hist 1850 time slice levels, and the HadSLP2 data set and CMIP5 models are relative to 1860–1899 climatology.