

Answer to referee #1

We thank referee #1 for his comments.

Answer to general comment

We do not consider that the issue on transport across the TTL has been settled by our work and we agree with the reviewer that the results are sensitive to the heating rates and there are significant differences among reanalysis in this respect. In order to put the emphasis on this point we have added new calculations based on MERRA which are actually showing large differences with respect to ERA-Interim and JRA-55 in the vertical distribution of sources. This is due as noticed by the referee to different heating and we show that clouds have a particularly strong effect on MERRA in convective region. See Section 4.4-Sensitivity to the reanalysis and new Figure 9.

The referee also advocate the use of Yang et al. (2010) heating rates calculated from the observed distribution of clouds. Such heating rates are obviously an interesting attempt to understand the effect of clouds onto heating rates. However, the observation are limited and the calculation of Yang et al. (2010) is based on a seasonally averaged distribution of clouds. Radiative transfer is unfortunately a very nonlinear process and does not commute with this type of averaging. There is no reason that the averaged effect of clouds can be accurately represented by the effect of the averaged clouds. This does mean that the calculations based on reanalysis are superior but they cannot be taken *a priori* as inferior.

Figure 1 shows the vertical profile of heating rates in February and August 2007, between 20°S and 20°N for the three reanalysis ERA-Interim, JRA-55 and MERRA. This figure compares to Fig. 6 of Bergman et al. (2012). The dispersion of heating rates and heating rate variances in the TTL is clear in this figure. However, none of the analysis displays a change of heating rate distribution over the tropical band between winter and summer that compares with that of Bergman et al. (2012).

The referee mentions that in Bergman et al. (2012) there is a big drop between 65% in winter and 15% in summer of the proportion of backward trajectories reaching a cloud. It seems that this is accompanied by a very strong weakening of the oceanic contribution during summer. We do not find anything similar in our calculations even with the MERRA reanalysis. Such effect is apparently a consequence of using Yang et al. (2010) heating rates. Since the upward flux of the Brewer-Dobson circulation does not vary in such proportion, it would mean a much larger recirculation of extra-tropical stratospheric air during summer to face the lack of fresh air supply from the boundary year with consequences on tracer distribution in the tropical stratosphere which are not supported by any observation to our knowledge.

It is actually very difficult to compare the calculated heating rates with observations. However, the integrated cloud radiative effect (CRE) of the three reanalysis considered here has been evaluated against CERES observations by Li and Mao (2015) who found that "spatial correlation of CREs and TOA upward radiation fluxes in ERA-Interim is the best among the three reanalyses" in spite of some discrepancies in the global mean CRE. Li and Mao (2015) also notice that all three reanalyses have difficulties to reproduce summer CREs over East Asia, a conclusion also supported by Wang et al. (2014) based on a study of radiation budgets in AMIP-5 models.

Therefore, we are led to conclude that the heating rates and the radiative effect of clouds in the Asian monsoon region are still a puzzle that requires further investigation.

We have added some paragraphs about these matters in our discussion section.

Answer to specific comments

The specific comments are reproduced to ease the reading.

1. ***Page 26240, line 8: It would be useful to explain, however briefly, how any convection (say, in the summer) with a cloud top theta of 348 can make a contribution to transport at 380K. Presumably, some of these 348K "emitted" parcels move horizontally into a region where the LZRH is less than 348K, and rise***

upward. It is not clear why the Tibetan high efficiency is a paradox. Based on the distribution of colored arrows, most heating rates in the tropics in the ERA-Interim analysis are positive at the Tibetan mode max of 365K (even if they are negative at 365K over Tibet). So, a little horizontal wandering will get the parcel into positive heating rates. Since the parcels are already so high, getting to 380K should be easier than for parcels that start at a lower theta (convective emission theta). In general, a little more mechanistic explanation of the findings in the figures would be helpful.

The LZRH is a mean value but it fluctuates a lot when taking into account the temporal and spatial variation of the cloudy component of the heating rate. Therefore parcels launched at 348 K can sample occasionally positive heating rates and a small proportion of them can make it to cross the LZRH above which it is much easier to reach the 380 K surface. See also point 3 below. We have two other articles in preparation where the mechanistic explanation will be more developed.

- 2. Figure 5: I understand the comparison between the backward trajectories for the two different satellite cloud altitude treatments (1 km offset), but am confused by the drastic difference in the forward trajectories. With lower clouds, it takes longer for emitted parcels to get to 380K, so is it a matter of integration time in the forward direction? 3 months (backward trajectories) is still enough time for parcels from either the low or high cloud case to mostly (80 plus %) get to 380K. More explanation is needed here***

It is not only a matter of integration time. While the proportion of backward trajectories reaching a cloud does not change, they do it to a restricted set of clouds and over a slightly longer time. Regarding the forward trajectories, the situation is different: many trajectories launched from the lower LZR clouds are never able to move up and they descend in the troposphere.

- 3. I assume the LZRH in Figure 3 for the different sources represents a mean LZRH for the locations of convection in that season and region (AML, SAP...). Might***

be good to make this clear. The LZRH must jump around a lot, because a lot of convective systems in the AML in DJF (for example) with cloud tops much less than 359K contribute to air at 380K.

It is true that the LZRH calculated over 3-hour interval jumps a lot but the mean is nevertheless relevant. We have added a sensitivity test where the heating rate is replaced by its moving monthly average. The transit properties are only very weakly affected by this transformation, therefore indicating that the horizontal motion is the main mechanism through which Lagrangian parcels sample the variations of heating rates. The caption of Figure 3 is modified for more clarity : "The arrows on the upper axes indicate the corresponding LZRH levels" → "the corresponding mean LZRH levels over the season and 2005–2008."

4. *There is no reference to the results of section 5 in the abstract that I can see.*

The results from this section are now mentioned in the abstract.

References

- Bergman, J. W., Jensen, E. J., Pfister, L., and Yang, Q.: Seasonal differences of vertical-transport efficiency in the tropical tropopause layer: On the interplay between tropical deep convection, large-scale vertical ascent, and horizontal circulations, *J. Geophys. Res.*, 117, D05302, doi:10.1029/2011JD016992, 2012.
- Li, Jian-Dong and Mao, Jiang-Yu: A preliminary evaluation of global and East Asian cloud radiative effects in reanalyses, *Atmospheric and Oceanic Science Letters*, 8, 100–106, doi:10.3878/AOSL20140093, 2015.
- Wang, F., Yang, S., and Wu, T.: Radiation budget biases in AMIP5 models over the East Asian monsoon region, *Journal of Geophysical Research: Atmospheres*, 119, 13,400–13,426, doi:10.1002/2014JD022243, 2014.
- Yang, Q., Fu, Q., and Hu, Y.: Radiative impacts of clouds in the tropical tropopause layer, *Journal of Geophysical Research*, 115, doi:10.1029/2009JD012393, 2010.




Figure 1. Mean heating rate profiles $d\theta/dt$ and \pm one standard deviation for February 2007 (yellow) and August 2007 (red) averaged over 20° S- 20° N for the three reanalysis ERA-Interim, MERRA and JRA-55.

Answer to referee #2

We thank referee #2 for his thorough reading of our manuscript, his comments and his numerous corrections of our poor English. We answer below only to the comments about the scientific content.

Answers to general comment

1. ***The current title is rather vague and, while accurate in a broad sense, does not properly convey the content of the manuscript. The problem is mostly in the order of the words; for example, something like “Convective sources of trajectories traversing the tropical tropopause layer” or “Convective sources of trajectories entering the tropical stratosphere” would be better. The abstract is concise and complete, but should be edited to improve the clarity.***

The title has been changed : "Transport across the tropical tropopause layer and convection" → "Convective sources of trajectories traversing the tropical tropopause layer" following the suggestion of the referee. We have edited the abstract.

2. ***As the CLAUS record has been extended to provide more overlap with the CALIPSO cloud lidar, I wonder if anyone has tried to collocate these observations to assess the actual biases in cloud top pressure derived from CLAUS. This could be done by, e.g., treating the CALIPSO track like a trajectory (which would also incorporate and be used to test interpolation-related uncertainties). Minnis et al. (2008) did a similar analysis for MODIS-derived cloud tops. One could imagine that this could quantitatively (and even qualitatively) impact estimates of the sensitivity of the source distribution (and particularly transit times) to the corrected cloud tops, as the implied correction could depend on the derived cloud top pressure and/or have regional variations. Given that the CLAUS brightness temperatures are often used in this way and such a specialized analysis would not stand on its own, I would really like to see someone provide this***

in one of these studies. Although I feel that it would be appropriate here, I propose this as a suggestion rather than a requirement.

The CLAUS dataset contains brightness temperature at $10.8\mu\text{m}$ from geostationary satellites and our derivation of the cloud top height is basically identical, for high opaque clouds, to the infrared radiance method used in algorithms deriving cloud top height from broadband infrared imagers. Kwon et al. (2010) and Hamann et al. (2014), both using SEVIRI data and comparing the derived cloud top height with CALIPSO and CLOUDSAT reach conclusions which are consistent with the earlier study of Sherwood et al. (2004) and our approximation. Such studies extend naturally to CLAUS data.

Answers to specific comments

- 1. In Fig. 2, lines in the same plot are shown on the same axis, but with that axis representing different aspects of trajectory time (tropospheric source vs stratospheric receptor). Although I don't disagree with the rationale, this choice of presentation is potentially confusing. Given the lack of major month-to-month variability in the proportion of trajectories reaching a cold cloud top, I think that the first result (maximum of 88.7% in April; minimum of 84.7% in July) could simply be reported in the text without graphical representation.***

We have modified the caption to avoid the confusion but we think it is useful to keep this curve as the stability of this proportion throughout the year is an important result which differs from the previous study of Bergman et al. (2012).

- 2. It is not paradoxical that transport from convective sources over Tibet is particularly efficient even though the LZRH is high (p.26240, l.11–12) when one considers that the prevailing winds within the anticyclone will tend to isentropically lift these trajectories above the LZRH over the subcontinent.***

Actually some cross-isentropic motion is necessary to reach the 380 K surface even if the high level of sources above the Tibetan plateau requires less heating than for other regions to perform that task. The main effect of the anticyclone is to confine the air from the Tibetan plateau within its core with limited meridional displacement. The lifting effect of the anticyclone is much stronger for the air originating from regions south of the Tibetan plateau and ventilated by the easterly branch of the anticyclone. We will discuss those aspects in a second article in progress that focuses on the horizontal displacements. We agree that the results about Tibet are not paradoxical and we have modified the sentence.

- 3. Note that one of the key differences among the studies cited on p.26250, l.16–19 is the choice of reanalysis/analysis data (as well as the representation of convection). In particular, all of the studies identified as agreeing with these results are based on ERA-Interim. Although this is only one region, this reinforces the point that the results may be more sensitive to the choice of reanalysis than indicated by the comparison of ERA-Interim and JRA-55. It's not necessary to provide more sensitivity studies here, but section 4.3 should also include relevant contextual information from previously published results.***

We have added a comparison to the MERRA reanalysis data and a discussion on the differences between our result and those of Bergman et al. (2012).

Answers to technical suggestions

We have taken into account the minor technical suggestions not mentioned below.

p.26248, 19-10: I'm not sure I quite follow this statement — is the source curve in Fig. 2a supposed to be the sum of the individual regions (which are arranged relative to intersection with convection) or the total number of trajectories coming from convection (which are arranged relative to release at 380K)? Is the shift a lead or a lag? This sentence should be clarified.

The discussion has been clarified. We now discuss the regional curves and the shift is a lag (in normal time running forward). This is expected since a parcel leaving a cloud reaches the 380 K surface after a transit time.

p.26267, figure : recommend reordering the panels so that season is top/bottom and category is left/right – although this makes it more difficult to evaluate the relative locations of the modal peaks, it makes it easier to compare the distributions (i.e., redistribution among regions) and would be consistent with the previous figures (thereby limiting confusion)

We understand this point but we have added two more panels to this figure ("smoothed heating rates") and this layout makes easier to compare the relative locations of the modal peaks.

p.26268, figure : it may be worth repeating the upper panels of Fig. 7 and showing this as a four-panel figure to facilitate comparison

We followed this advice and have also added two panels showing the vertical source distribution obtained with MERRA reanalysis.

References

- Bergman, J. W., Jensen, E. J., Pfister, L., and Yang, Q.: Seasonal differences of vertical-transport efficiency in the tropical tropopause layer: On the interplay between tropical deep convection, large-scale vertical ascent, and horizontal circulations, *J. Geophys. Res.*, 117, D05302, doi:10.1029/2011JD016992, 2012.
- Hamann, U., Walther, A., Baum, B., Bennartz, R., Bugliaro, L., Derrien, M., Francis, P. N., Heidinger, A., Joro, S., Kniffka, A., Le Gléau, H., Lockhoff, M., Lutz, H.-J., Meirink, J. F., Minnis, P., Palikonda, R., Roebeling, R., Thoss, A., Platnick, S., Watts, P., and Wind, G.: Remote sensing of cloud top pressure/height from SEVIRI: analysis of ten current retrieval algorithms, *Atmospheric Measurement Techniques*, 7, 2839–2867, doi:10.5194/amt-7-2839-2014, 2014.
- Kwon, E.-H., Sohn, B.-J., Schmetz, J., and Watts, P.: Intercomparison of height assignment methods for opaque clouds over the tropics, *Asia-Pacific Journal of Atmospheric Sciences*, 46, 11–19, doi:10.1007/s13143-010-0002-7, 2010.

Sherwood, S. C., Chae, J.-H., Minnis, P., and McGill, M.: Underestimation of deep convective cloud tops by thermal imagery, *Geophys. Res. Lett.*, 31, L11102, doi:10.1029/2004GL019699, 2004.

[..¹] Convective sources of trajectories traversing the tropical tropopause layer [..²]

A.-S. Tissier and B. Legras

Laboratoire de Météorologie Dynamique, UPMC Paris 06/CNRS/ENS/Ecole Polytechnique,
UMR 8539, Paris, France

Correspondence to: A.-S. Tissier (as.tissier@lmd.ens.fr)

¹removed: Transport across

²removed: and convection

Abstract

Transit properties across the tropical tropopause layer are studied using extensive forward and backward Lagrangian **adiabatic** trajectories between cloud tops and the reference surface 380 K. ^[.3]After dividing the tropical domain into 11 sub-regions according to the distribution of land and convection, we estimate the contribution of each region to the upward mass flux across the 380 K surface, the vertical distribution of convective sources and of transit times over the period 2005–2008. The good agreement between forward and backward statistics is the basis of the results presented here. It is found that about 85 % of the tropical parcels at 380 K originate from convective sources ^[.4]throughout the year. From November to April, the sources are dominated by the warm pool which accounts for up to 70 % of the upward flux. During summer, Asian monsoon region is the largest contributor with similar contributions from ^[.5]the maritime and continental parts of the region; however, the vertical distributions and transit times associated with these two subregions is very different^[.6]. Convective sources are generally higher over the continental part of the Asian monsoon region, with shorter transit times. ^[.7]We estimate the monthly averaged upward mass flux on the 380 K surface and show that the contribution from convective outflow accounts for 80% on average and explain most of its seasonal variations. The largest contributor to the convective flux is the South Asian Pacific region (Warm Pool) for 39% throughout the year followed by oceanic regions surrounding continental Asia for 18 % and Africa for 10.8%. Continental Asia low lands accounts for 8%. The Tibetan plateau is a mi-

³removed: The tropical domain being subdivided

⁴removed: all along

⁵removed: oceanic regions and Asian mainland, although the signature in vertical distribution and transit time

⁶removed: , Asian mainland displaying higher sources and smaller

⁷removed: The Tibetan plateau, although

nor overall contributor [..⁸](0.8%), but transport from convective sources in this region is very efficient due to its central location beneath the Asian upper level anticyclone. [..⁹]

The core results are robust to uncertainties in data and methods[..¹⁰], but the vertical source distributions and transit times [..¹¹] exhibit some sensitivity to the representations of cloud tops and heating rates. The main sensitivity is to the radiative heating rates which vary among reanalysis.

1 Introduction

The tropical tropopause layer (TTL) is a key region in the atmosphere that controls the transport of tropospheric air into the stratosphere (Highwood and Hoskins, 1998; Fueglistaler et al., 2009). Situated above the level of main convective outflow (Corti et al., 2006; Fueglistaler et al., 2009), the TTL is penetrated by deep convection which becomes increasingly rare with altitude (Liu and Zipser, 2005; Fu et al., 2007). [..¹²]Outside of convective towers, the vertical motion is weak[..¹³]. The TTL region encompasses the level of zero radiative heating (LZRH) which marks the transition from negative to positive radiative heating values, thus creating a barrier for the large-scale transport of air parcels into the stratosphere (Folkins et al., 1999). Quantifying the transport paths across the TTL [..¹⁴]and their spatial and temporal variability is important for a better understanding of the chemical composition of the air entering the stratosphere and the relation with source regions [..¹⁵]

⁸removed: , is found to be the region with the highest impact of convection at 380 K

⁹removed: We show the robustness of our main results

¹⁰removed: but we also find some sensitivity of the vertical distribution of sources

¹¹removed: to the representation

¹²removed: At the same time

¹³removed: and the

¹⁴removed: , its

¹⁵removed: (Holton et al., 1995; Fueglistaler et al., 2004; Fu et al., 2006; James et al., 2008; Park et al., 2009; Aschmann et al., 2009; Tzella and Legras, 2011; Bergman et al., 2012; Chen et al., 2012; Bergman et al., 2013; Heath and Fuelberg, 2014; Orbe et al., 2015)

](Holton et al., 1995; Fueglistaler et al., 2004; Fu et al., 2006; James et al., 2008; Park et al., 2009; Aschmann et al., 2009; Tzella and Legras, 2011; Bergman et al., 2012; Chen et al., 2012; Bergman et al., 2013; Heath and Fuelberg, 2014; Orbe et al., 2015; Bergman et al., 2015; Vogel et al., 2015).

In this work, we focus on how parcels detrained from the convective clouds are transported across the TTL and reach the tropopause, defined as the 380 K surface, upon the combined effect of heating rate fluctuations and vertical distribution of convective sources.

Although the distribution of tropical convection is well known, the location and intensity of convective sources in the TTL are still debated. The questions addressed in this work are whether the intensity is linked only to the altitude of cloud tops (Gettelman et al., 2002, 2009; Devasthale and Fueglistaler, 2010), the role of cloud heating in favoring the crossing of the LZRH (Corti et al., 2006; Tzella and Legras, 2011), the role of horizontal transport and in-mixing from extra-tropical latitudes (Ploeger et al., 2012), and the special role of continental convection during the Asian monsoon in particular above the Tibetan Plateau [..¹⁶](Fu et al., 2006; Devasthale and Fueglistaler, 2010; Bergman et al., 2013; Heath and Fuelberg, 2014; Vogel et al., 2015).

We address the whole range of these questions by performing for the first time a comprehensive set of forward and backward trajectories in the TTL between convective sources and the tropopause. We do not account for the transport from the boundary layer to convective tops which is assumed to be fast and local.

In Sect. 2, we describe the data and the methods used in this work to retrieve trajectories connecting the clouds and the 380 K potential temperature surface. Section 3 discusses the distribution of convective sources and transit times in the TTL. Section 4 discusses the sensitivity of our results [..¹⁷]to uncertainties in the data and methods. Section 5 makes a further step by calculating the mass flux across the 380 K surface and the contributions of convective sources. Section 6 offers a summary and outlook.

¹⁶removed: (Fu et al., 2006; Devasthale and Fueglistaler, 2010; Bergman et al., 2013; Heath and Fuelberg, 2014)

¹⁷removed: t

2 Lagrangian trajectories and convective sources

Lagrangian trajectories of air parcels are calculated within the TTL between the time of detrainment from convective sources and the crossing of the 380 K potential temperature surface, taken as the lower boundary of the stratospheric over-world [..¹⁸](Holton et al., 1995).

2.1 Determination of the altitude of deep convective clouds

A prerequisite of this study is a characterization of cloud tops that is both global in space and time. We use the CLAUS dataset (Hodges et al., 2000) which provides global 3-hourly maps of brightness temperature at 30 km resolution, combined with ERA-Interim data (Dee et al., 2011) to determine the pressure of the top of the convective clouds. Since we are only interested by air parcels which are able to reach the LZRH and above, we consider only cold pixels with brightness temperatures less than 230 K. In the deep tropics between 15° S and 15° N, this temperature corresponds to a pressure of about 240 hPa which is below the main detrainment level near 200 hPa and well below the all-sky LZRH which is usually located above 155 hPa. As the mean detrainment levels and the LZRH are even higher over continents and in particular over Asia during monsoon season, this is a [..¹⁹]conservative choice.

This method is, however, limited by the inability to distinguish overlying cirrus from convective tops and is known to underestimate the altitude of the deep convective clouds by about 1 km (Sherwood et al., 2004; Minnis et al., 2008). We correct the altitude provided by CLAUS by an upward shift of 1 km. [This approximation is consistent with more recent comparisons of cloud top height determined from active sounders \(Kwon et al., 2010; Hamann et al., 2014\).](#) The sensitivity to this correction is described in Sect. 4.1. [..²⁰]Another limitation is that the brightness temperature can be colder than the environment and sometimes

¹⁸removed: Holton et al. (1995)

¹⁹removed: very

²⁰removed: An other

is colder than the cold point tropopause. This is possible under fast adiabatic cooling within active convective towers (Adler and Mack, 1986; Luo et al., 2008) and is found to occur for less than 2% of the cold pixels for each month in the 20° S–40° N band. In such cases, we follow Sherwood et al. (2004) and consider that the parcels rise adiabatically from an altitude of about 40 hPa below the cold point tropopause.

2.2 Three dimensional Lagrangian trajectories

We compute forward and backward diabatic three-dimensional trajectories in the TTL using TRACZILLA, which is a modified version of FLEXPART (Stohl et al., 2005; Pisso and Legras, 2008). The horizontal part of the motion is calculated using the 3-hourly wind fields of ERA-Interim (Dee et al., 2011), combining analysis at 00:00, 06:00, 12:00 and 18:00 UT, 3 h forecast at 03:00 and 15:00 UT and 9 h forecast at 09:00 and 21:00 UT. The vertical (cross-isentropic) displacement is calculated using the 3-hourly average all-sky radiative heating rates of ERA-Interim (including the radiative effect of the clouds but excluding the latent heating) which are archived at 01:30, 04:30, 07:30, 10:30, 13:30, 16:30, 19:30 and 22:30 UT.

The forward [..²¹]calculations are meant to estimate the impact and efficiency of convective sources while the backward calculations estimate the contribution of sources to the air composition at the 380 K surface. The two calculations provide, however, very consistent results as shown below. The calculations are performed over the whole period 2005–2008 and are discussed below in terms of monthly statistics.

2.2.1 Backward simulation

In the backward calculation, parcels are launched from the 380 K potential temperature surface, every two days, between 40° S and 40° N, on a regular grid of 0.5 degree in latitude and longitude. We retain the first encounter of a parcel with the top of a cold cloud within the previous three months as in Tzella and Legras (2011) except that our criterion is based on

²¹removed: calculation

pressure rather than temperature. A parcel encounters a cloud when its pressure is larger than the pressure of the corrected top as described above. The comparison is performed 3-hourly along the trajectory of the parcel with the CLAUS pixel containing the parcel at that time. It is clear that we may miss some cloud encounters in this way because the parcels can sometimes travel by 300 km or more over 3 h, that is 10 CLAUS pixel sizes, under strong wind conditions. The sensitivity to such effect is tested in Sect. 4.2

2.2.2 Forward simulation

In the forward calculation, parcels are launched from 3-hourly CLAUS maps, with one parcel at the center of each cold pixel (brightness temperature ≤ 230 K) at the corrected altitude of the cloud top, for locations between 20° S and 40° N. Trajectories are integrated for three months and we retain the first crossing of the 380 K surface between 40° S and 40° N when it occurs. Parcels encountering other clouds along their path, that is which are found within cloudy pixels with cloud tops higher than their altitude, are discarded. On the average, between 16% (January) and 24% (August) of the launched parcels are [..²²]eliminated for this reason. Such parcels are mostly on a descending path and keeping them instead of discarding them [..²³]only marginally affects our results for parcels crossing the 380 K surface.

3 Source distribution

We focus our study on different geographical regions as defined by the color boxes on Fig. 1. The boundaries are chosen to highlight the regions where convection is intense in the tropics and subtropics and to separate land and oceanic contributions. The South Asian Pacific (SAP) region corresponds to the warm-pool. The Asian monsoon region is divided into the continental Asia Mainland (AML), the North Asian-Pacific Ocean (NAPO)

²²removed: ending their flight in this way

²³removed: affects only marginally

which includes the Bay of Bengal and the Sea of China and the ^[..²⁴] **Tibetan Plateau (Tibet)** defined as the region in Asia above 3500 m. America is divided into South America (SAm) and Central America (CAm). There is a single region for Africa (Af).

3.1 Annual cycle

The black curve in Fig. 2a shows that on ^[..²⁵] average 86 % of the backward parcels reach a cloud top within three months. The time axis for this curve is the launch time. The parcels which do not reach a cloud are to a vast majority initialized in the subtropics and ascend backward in the deep Brewer-Dobson circulation of the extra-tropics. The proportion of parcels reaching a cloud varies very little over the mean annual cycle with a maximum in April (88.7 %) and a minimum in July (84.7 %).

The other curves in the same panel show the contributions of each region among the trajectories reaching a cloud within the ensemble of all regions in Fig. 1. In order to facilitate the comparison with forward calculations, the time axis for these curves is that of the intersection of each parcel with a cloud, grouped by months.

The main feature between November and April is the ^[..²⁶] **dominance** of the SAP region which accounts alone for a maximum of 68.4 % of all sources in January. The next winter contributors are Af and SAm with a maximum contribution of 19.4 % for Africa in April.

From June to September, NAPO ^[..²⁷] **is the leading source** with a maximum share of 35.3 % in July followed by AML which ^[..²⁸] **peaks** at 19.8 % in July. Together these two regions represent from 45 to 55 % of all sources from June to September. The following contributors are SAP, CAm, North Central Pacific (NCP) and Af. Tibet is a tiny overall contributor (with ^[..²⁹] **a** maximum share of 2.5 % in July).

²⁴removed: Tibet

²⁵removed: the

²⁶removed: domination

²⁷removed: dominates

²⁸removed: caps

²⁹removed: is

is The forward estimate of the source distribution, see Fig. 2b, is calculated from the trajectories launched at cloud top level which have reached the 380 K surface. The quantity shown is the monthly ratio of the number of parcels from a given region to the total number originating from the ensemble of regions in Fig. 1. The time associated with each trajectory is that of its launch, grouped into monthly bins. There is a striking agreement between the forward and backward distribution of sources in Fig. 2a and 2b. In spite of some slight quantitative changes in the proportions, the general pattern and the ordering of sources is ^[.30]almost identical throughout the whole year. The largest change is for Tibet which displays a forward contribution of 7.6% in July and August ^[.31](three times its backward contribution), but still remains a minor overall contributor.

The similarity is not totally unexpected as forward and backward calculations are solving dual equations for the Green function of the advection-diffusion equation (Holzer and Hall, 2000; Legras et al., 2005). Here diffusion has no role because we consider averages over regions and durations much larger than the diffusion scale and ^[.32]diffusion time defined in Legras et al. (2005). It is however surprising that a somewhat coarse discretization (only one parcel per cold $30 \text{ km} \times 30 \text{ km}$ pixel in the forward case and one parcel on a half-degree grid every two days ^{in the backward case}) which under-samples quite drastically the flow is able to reach good agreement between forward and backward calculations. The sampling is made at a higher resolution than the ERA-Interim winds but the transit time between the cloud top and the 380 K surface is long enough (as shown below) to make the trajectories numerically irreversible due to the ^[.33]chaotic aspects of transport. Our results show that the sampling is good enough to reestablish the reversibility in the statistical sense as predicted by the Green function formalism for a continuous sampling (Holzer and Hall, 2000).

Figure 2c shows the proportion of forward trajectories released in a given region from high convective tops that reach the 380 K surface within 3 months. As most of the other tra-

³⁰removed: quasi-identical over

³¹removed: , that is

³²removed: time as

³³removed: transport chaos

jectories have returned back in the lower troposphere, this proportion conveys the efficiency of each region at converting convective air into stratospheric air. Tibet displays a singular behaviour by reaching an efficiency of 84.5 % in July which indicates that this proportion of air detrained at the top of clouds reaches the 380 K surface. During summer, after Tibet, AML conveys up to 53.5 % of parcels to the stratosphere while NAPO is less efficient at about 32.5 % but still the largest contributor due to its size and the frequency of high convective clouds within its domain.

The other regions exhibit [..³⁴] efficiencies lower than 30 % with SAP lying just above SAm, CAm and Af. However, when the Asian land convection [..³⁵] ramps down, the efficiency of SAP combines with the intensity of convection in this region to let it dominate the transfers over half of the year.

The high efficiency above Tibet is consistent with previous studies who found a confinement of Tibetan air within the monsoon anticyclone (Bergman et al., 2013; Heath and Fuelberg, 2014) which persists above Asia during summer, trapping tracer compounds that recirculate inside (Park et al., 2007, 2009). The contribution to the inside of the summer Asian monsoon anticyclone (AMA) has been further estimated by calculating sources over the restricted portion of the 380 K surface confined between 20 and 40° N, 25 and 125° E, and with potential vorticity smaller than $4 \times 10^6 \text{ m}^2 \text{ s}^{-1} \text{ kg}^{-1} \text{ K}$. It is found that 87 % of the AMA parcels originate from Asia in the forward calculation, see Fig. 2b, among which [..³⁶] 19.5 % from the Tibetan plateau. In the backward calculations, see Fig. 2a, the proportions are redistributed among Asian continental convection between AML and Tibet (respectively 47.5 and 19.5 % in forward and 54.3 and 8.8 % in backward, the sum varying only from 67 to 63.1 %). The NAPO contribution is unchanged.

³⁴removed: efficiency

³⁵removed: fades in, the larger

³⁶removed: 19,5

3.2 Vertical distribution of sources

We investigate now the vertical distribution of sources within each region and its relation with the LZRH. During winter, see Fig. 3a and 3b, forward and backward calculations predict that the distribution of cloud top sources in the dominating SAP region peaks at 355 K^[..³⁷]. This is located above the all-sky LZRH ^[..³⁸] at 352.6 K during that season. A proportion of 89.5 % of the sources are then located above the LZRH in the forward calculation and 76.3 % in the backward calculation. The other contributing regions (Af, SAm, NAPO and NCP) also ^[..³⁹] all exhibit a modal peak at 355 K in the forward distribution. The backward distribution exhibits a shift towards small potential temperature which affects also the mean and the median of the distribution (see Table 1).

As in Tzella and Legras (2011), a more complex pattern emerges during summer with several competing regions, see Fig. 3c and d. Both forward and backward calculations produce similar ^[..⁴⁰] distributions of sources with the main differences being a stronger contribution of Asian land regions (AML and Tibet) in the forward distribution. SAP, CAM and NCP regions are grouped in the forward calculation with a modal peak at 353 K which is below the winter peak at 355 K in SAP. A similar pattern is found in the backward calculation. The AML and Tibet modal peaks are above 360 K, reaching 367 K for Tibet. The Africa modal peak is also near 360 K with a fairly flat distribution, and NAPO modal peak is intermediate near the common winter peak at 355 K. The backward shift in the sources towards small potential temperature is smaller than during winter in SAP. Most of the sources are again located above the LZRH, up to 90 % for the forward trajectories from NAPO and CAM (see Table 1). The only exception is Tibet, for which the LZRH, much higher than in other regions at 367.3 K, is located above the modal peak, and just at the median in forward calculations. Nevertheless, Tibet was found ^[..⁴¹] as the region with highest efficiency during summer

³⁷ removed: , that is

³⁸ removed: located

³⁹ removed: exhibit all

⁴⁰ removed: distribution

⁴¹ removed: quite paradoxically

and this might be explained by the very high level of the LZRH and sources in this region. The separation of NAPO and Asian land sources explain the double peak pattern shown in Fig. 8 of Tzella and Legras (2011).

3.3 Transit time

The differences between forward and backward calculations are mostly seen in the transit time distribution shown in Fig. 4. The peak of the distribution is always shifted to smaller values in the forward calculations and the tail is also decaying much faster. As a result, the ratio backward/forward for the median and the mean is of the order of 1.5 in SAP during winter and in AML and NAPO during summer. It is lower but always larger than 1.1 in the other regions with the exception of Tibet with a near 2 factor, see Table 2. In the forward calculation, the mean transit time over contributing oceanic regions (SAP, NCP and CAm) is of the order of 30 days during summer, which is larger than the winter value of 23 days in SAP. On the contrary, Asian land regions during summer exhibit shorter transit times (21 days for AML and 15 days for Tibet) than their winter counterparts (Af and SAm) near 28 days. [..⁴²] **Transit times from Af show** very little change between winter and summer.

The backward distribution is biased by the fact that sampled backward trajectories can easily miss a cloud by passing a pixel away and then wander away in another region. The effect is the largest for small regions like Tibet or regions ventilated by [..⁴³] **intense large scale** circulation such as Asia during summer. The backward trajectories can also, at least in principle, get trapped into unstable trajectories oscillating about the LZRH from which trajectories diverge in forward time and to which they therefore converge in backward time, although this has seldom been observed. As it appears, transit times are more sensitive to sampling effects than the source distribution.

⁴²removed: Af shows

⁴³removed: large scale intense

4 Sensitivity studies

In this section, we study the sensitivity of the results presented in Sect. 3 to changes in data and the design of our calculations.

4.1 Sensitivity to the cloud top offset

As the estimate of cloud top and the +1 km correction are subject to uncertainty, we have redone the analysis without the +1 km correction. Since the LZRH is the same, the direct effect is to reduce the proportion of forward trajectories reaching the 380 K surface, see Table 3. The ratio is about 45 % for both summer and winter but for continental Asia during summer where it is smaller. Tibet is the most sensitive region with a ratio of 22 %.

Figures 5a and 5c shows the change in the vertical distribution of sources for forward calculations. Besides the overall reduction, it is visible that the modal peaks are unremoved but for NAPO and Tibet during summer where they move, respectively, from 358 to 355 K and from 367 to 362 K. The vertical distribution of sources is made narrower by reducing the [\[..⁴⁴ \]tail](#) of the distribution towards the upper end of the interval as the highest clouds are shifted down.

The proportion of backward trajectories reaching a cloud within 3 months is now 85 % during DJF and 82.7 % during JJA, that is less but close to the value when the offset is applied (87 and 85.2 % respectively), hence with much less variations than the forward efficiency. The modal peaks are slightly shifted to lower values (with larger shift for NAPO and Tibet) and the narrowing is well pronounced with almost no sources within the [range \[365, 370 K\]](#) except over continental Asia during summer.

The lower end cutoff value of sources (at about 345 K for maritime convection and 350 K for AML) is preserved in both forward and backward calculations but for Tibet. This suggest that this cutoff is determined by the transport properties across the LZRH which are unchanged by offsetting the cloud top while the cutoff above the LZRH depends strongly

⁴⁴removed: aisle

on the distribution of cloud tops. Tibet differs by being the only region where the LZRH is above the modal peak.

[..⁴⁵] In accordance with these limited changes in the sources, the transit time distribution for all regions but Tibet is weakly affected for both forward and backward calculation, see Fig. 6. The change is localized in the small time contribution in agreement with the narrowing of the source distribution, which reduces the proportion of short transit paths. Tibet is an exception with a shift by about a factor two of the forward and backward transit times towards larger values. Tibet differs from the other regions by having a very high LZRH and a distribution of sources laying mainly under this level. This effect is amplified when the cloud top correction is cancelled with almost no contribution left above the LZRH, inducing a significant shift in the transit time distribution.

4.2 Sensitivity to increase of the size of cloud pixels

In this section we address the sensitivity to the density of cloud observation and the effect of missing cloud encounters in backward calculations. We modify the encounter criterion by enlarging the pixel size by a factor three in both latitude and longitude, retaining the smallest top pressure among the 9 CLAUS pixels surrounding the parcel at a given time. This modification [..⁴⁶] enlarges high clouds and has largest effect in regions where convective systems are small and sparse. Figure [..⁴⁷] 7 (first and second rows) compares the distributions of backward sources in winter and summer 2005 with and without enlarging the pixel size. The total number of trajectories meeting a cloud is again quite insensitive increasing by 3 % during winter and 1.5 % during summer. The distribution of sources is however modified by shifting the distribution to higher potential temperatures and widening the profile. Hence, the effect is [..⁴⁸] qualitatively opposite to the lowering of the top of clouds done in Sect. 4.1.

⁴⁵removed: Accordingly to

⁴⁶removed: enlarge

⁴⁷removed: ?? compares the distribution

⁴⁸removed: quite opposite, qualitatively

4.3 Sensitivity to the daily cycle of the heating rates

Cloud radiative forcing and the resulting heating rates are *a priori* sensitive to the daily cycle of convective activity in the tropics. We test here the sensitivity to the daily cycle of heating rates by replacing the 3-hourly sampling by a time moving average. This average at time t is performed as a discretization of

$$\tilde{X}(t) = \frac{1}{2\tau} \int_{t-\tau}^{t+\tau} X(t') \left(1 + \cos \left(\pi \frac{t-t'}{\tau} \right) \right) dt', \quad (1)$$

where τ is 30 days.

Figure 7 (third row) shows that the distributions of sources are only weakly affected even at level below the LZRH. This result is in agreement with Bergman et al. (2012) who did a similar test with MERRA reanalysis. However, the maps (not shown) of the smoothed heating rates still contain a large amount of spatial variability. This suggests that the horizontal motion that samples this variability is more important than daily fluctuations of the heating rates to cross the LZRH.

4.4 Sensitivity to the reanalysis

One main source of uncertainty is the error in the reanalysis wind and heating rates. It has been shown that the heating rates differ quite significantly among reanalysis (Wright and Fueglistaler, 2013) and that the horizontal wind may [\[.49\]](#) contain large errors in tropical regions poorly covered by radio-soundings (Podglajen et al., 2014). It is therefore important to assess how our results are sensitive to a change of the reanalysis. As an extensive comparison among all available reanalysis at the time of this writing would have consumed a lot of resources, we limit the comparison to [\[.50\]](#) two reanalysis, JRA-55 (Kobayashi et al.,

⁴⁹removed: be quite erroneous

⁵⁰removed: the

2015) [⁵¹] and MERRA (Rienecker et al., 2011). JRA-55 has higher horizontal resolution than the ERA-Interim (spherical T319 truncature instead of T255) and the same number of levels. Winds and heating rates are available every 6 h at model resolution. MERRA has about the same horizontal resolution as ERA-Interim but because the heating rates are only available in this format, we use winds and heating rates on a 1.25° horizontal grid and a reduced set of vertical pressure levels every 3 hours.

Backward calculations have been performed for 2005 using the same setup as for ERA-Interim. [⁵²] Fig. [⁵³] 8 shows that in the three cases, SAP dominates during winter and NAPO is the largest contributor during summer. There are, however, significant differences. The relative contributions of NAPO is largest in JRA-55 and the distributions are narrower in MERRA. The main difference is in [⁵⁴] the vertical location. JRA-55 sources are slightly shifted upward by 2 to 3 K with respect to ERA-Interim. MERRA sources are even more shifted by up to 12 K for oceanic sources (SAP, NAPO, CAm) during summer. The [⁵⁵] shift is smaller for SAP in winter (+5 K) and for AML and Tibet in summer (+4 K). The total proportion of backward trajectories meeting a cloud remains, however, [⁵⁶] close to that of the ERA-Interim ([⁵⁷] 80.7% [⁵⁸] for JRA-55 and 74.4% against 85.4% for ERA-Interim in 2005).

In order to interpret these results, Fig. 9 compares the mean profiles of all sky heating rates among the reanalysis for each region in January and July. In addition to the three

⁵¹ removed: which

⁵² removed: The vertical source distribution shown in

⁵³ removed: ?? is to be compared with the two upper panels of Fig. ?. There is very good qualitative agreement but there are also differences. SAP is even more prominent during winter and but the largest

⁵⁴ removed: NAPO taking a larger share

⁵⁵ removed: distributions are also shifted towards higher altitudes, a result which is compatible with a generally higher LZRH in JRA-55 (not shown)

⁵⁶ removed: very

⁵⁷ removed: 81

⁵⁸ removed: against 85

reanalysis, we show also the curve for MERRA2 (Molod et al., 2015). It is clear that all the curves are always close within non convective regions (AML, CAm, SEP, Tibet during winter; SAm, SEP during summer) where heating rates are calculated from clear sky radiative transfer. In this region, ERA-Interim, however, displays larger heating rates than the two others above 370 K without affecting the LZRH.

Over convective regions, where additional cooling or heating is provided by clouds, there is a clear separation between the couple JRA-55/ERA-Interim and MERRA/MERRA2. ERA-Interim still displays larger heating rates than JRA-55 and this shifts down its LZRH by a few K. In the Asian region during summer and on SAP during winter, ERA-Interim cools less than JRA-55 below 340 K but this does not affect the LZRH. This is consistent with the shift observed in the source distribution.

MERRA/MERRA2 exhibit a very special pattern over convective regions with reduced heating near 355 K with respect to the two other reanalysis and a strong heating near 345 K, resulting in a characteristic "S" pattern. As a result, the LZRH is pushed upward and multiple LZRH occur over NAPO and CAm during summer. In all cases, MERRA2 is very close to MERRA except over Tibet during summer, where MERRA does not differ very much from ERA-Interim and JRA-55 while MERRA2 does quite unexpectedly.

5 Mass flux across the 380 K surface and regional distribution

In this section, we make a further step by determining the mass flux across the 380 K surface and the contribution of each convective region.

5.1 Method and validation

The instantaneous diabatic mass flux M across the 380 K surface, over a specific domain Γ of the sphere, can be estimated from all sky radiative heating rates as:

$$M_{\text{diab}} = \int_{\Gamma} \int \sigma_{380\text{K}} \left. \frac{d\theta}{dt} \right|_{380\text{K}} ds, \quad (2)$$

with $\sigma_{380\text{K}} = -\frac{1}{g} \frac{\partial p}{\partial \theta}$ at $\theta = 380\text{K}$, obtained from the temperature and pressure profile of ERA-Interim, with $g = 9.81 \text{ m s}^{-2}$. For practical purposes, the integration is replaced by a weighted sum over the gridded surface. From these instantaneous fluxes, one can define monthly averages by replacing $\left. \frac{d\theta}{dt} \right|_{380\text{K}}$ in Eq. (2) by its monthly average $\langle \left. \frac{d\theta}{dt} \right|_{380\text{K}} \rangle$. The monthly upward $M_{\text{diab}}^{\uparrow}$ and downward $M_{\text{diab}}^{\downarrow}(\phi)$ fluxes can then be defined by integrating separately on the sub-domains where $\langle \left. \frac{d\theta}{dt} \right|_{380\text{K}} \rangle$ is, respectively, positive and negative. It is known that upward and backward fluxes are ill-defined when the time interval goes to zero (Hall and Holzer, 2003). This is no longer the case after time smoothing which removes the noise but the flux then may depend on the applied time smoothing interval.

Another, [..⁵⁹]more traditional, method is based on the residual mean meridional circulation $(\overline{v^*}, \overline{w^*})$, with

$$\overline{v^*} = \overline{v} - \frac{1}{p} \frac{\partial}{\partial z} \left(p \frac{\overline{v'\theta'}}{\theta_z} \right) \quad (3)$$

$$\overline{w^*} = \overline{w} + \frac{1}{a \cos \phi} \frac{\partial}{\partial \phi} \left(\cos \phi \frac{\overline{v'\theta'}}{\theta_z} \right) \quad (4)$$

as defined by Andrews et al. (1987) in log pressure coordinates $z = H \log(p_0/p)$ where the [..⁶⁰]overbar indicates a zonal average. [..⁶¹]The kinematic mass flux is then given by (Appenzeller et al., 1996):

$$M_{\text{kine}} = \iint \left[\frac{p_{380\text{K}}}{gH} \left(\overline{w^*} - \frac{1}{a} \frac{\partial z}{\partial \phi} \bigg|_{380\text{K}} \overline{v^*} \right) + \frac{1}{g} \frac{\partial p}{\partial t} \bigg|_{380\text{K}} \right] ds, \quad (5)$$

with $p_{380\text{K}}$ the pressure at the 380 K surface, $a = 6371 \text{ km}$ and $H = 7 \text{ km}$. Monthly upward and downward fluxes, $M_{\text{kine}}^{\uparrow}$ and $M_{\text{kine}}^{\downarrow}$, can be separated according to the sign of the

⁵⁹removed: mode

⁶⁰removed: bars stands for the

⁶¹removed: This

monthly average term under the integral [..⁶²]Eq. (5). Gridded summation is applied in the same way as M_{diab} .

Figure 10 compares the monthly average upward fluxes calculated from [..⁶³]Eq. (2) and [..⁶⁴]Eq. (5). Here the diabatic flux is calculated from 3-hourly data and then averaged for each month over the period 2005–2008. The upward flux is calculated as indicated above for each month and then [..⁶⁵]averaged over the four years. The kinematic mass flux is calculated in the same way from monthly averages of the residual circulation and pressure at 380 K. The contribution of the pressure variation term in [..⁶⁶]Eq. (5) is then two orders of magnitude smaller than that of the residual velocity and can be neglected.

The two estimates display a similar modulation with a minimum during summer and agree with other estimates from [..⁶⁷]ERA-Interim (Abalos et al., 2012). There is, however, a shift between $M_{\text{diab}}^{\uparrow}$ and $M_{\text{kine}}^{\uparrow}$ which increases with the size of the latitude band. The mean difference is close to 25% in the three latitude bands. It is [..⁶⁸]notable that the upward flux and the total flux are identical in the 20° S–20° N band because [..⁶⁹]monthly mean diabatic heating rates on the 380 K [..⁷⁰]surface within this domain are positive throughout the year. It is also [..⁷¹]notable that there is little change in the upward flux [..⁷²]as the domain is expanded to 30° S–30° N and 40° S–40° N because most of the upward motion occurs within the [..⁷³]20° S–20° N band.

⁶²removed: (

⁶³removed: (Fig.

⁶⁴removed: (Fig.

⁶⁵removed: average

⁶⁶removed: (Fig.

⁶⁷removed: the

⁶⁸removed: noticeable

⁶⁹removed: this domain of the surface

⁷⁰removed: is heated everywhere all along the year in monthly average

⁷¹removed: noticeable

⁷²removed: between

⁷³removed: former

The discrepancy between diabatic and kinematic fluxes can be reduced by correcting the diabatic mass flux to satisfy global mass conservation. There is no physical mean [..⁷⁴] of ensuring such mass conservation when calculating heating rates (Shine, 1989) from radiative [..⁷⁵] transfer. Actually, the mean total mass flux across the 380 K provided by Eq. (2) and [..⁷⁶] ERA-Interim data is $7.9 \times 10^9 \text{ kg s}^{-1}$ over 2005–2008. Applying a uniform compensating correction to the heating rates over the sphere defines a new mass flux $M_{\text{diab,corr}}^{\uparrow}$ which reduces the discrepancy with $M_{\text{kine}}^{\uparrow}$ for all latitudes and all times. However, as this uniform correction is entirely ad hoc in the absence of any information about the spatial and temporal distribution of the errors, we refrain from applying it in the [..⁷⁷] subsequent analysis.

The definition used for the diabatic flux is consistent with what follows. However a more appropriate definition to compare with the kinematic fluxes would be to take a zonal average before applying the sign criterion defining the upward flux. The resulting upward flux (not shown) is indeed larger than $M_{\text{diab}}^{\uparrow}$ but the difference in the $40^{\circ} \text{ S} - 40^{\circ} \text{ N}$ band is of the order of 1 [..⁷⁸] to 2%. [..⁷⁹] This shift is negligible and cannot explain the shift between kinematic and diabatic fluxes.

Notice that the total mass flux across an isentropic surface does not need to vanish instantaneously unlike across an isobaric surface under the hydrostatic approximation. However the small part of the pressure variation term in the monthly flux indicates that the mass balance must be satisfied over monthly averages in the same proportion, that is within about 1%. This is in agreement with the stratospheric over-world mass variations shown by Appenzeller et al. (1996).

⁷⁴removed: to constrain

⁷⁵removed: transfers and actually

⁷⁶removed: the

⁷⁷removed: sequel

⁷⁸removed: % but in winter where it is less than

⁷⁹removed: Therefore this is taken as

Having compared the diabatic mass flux from heating rates to the kinematic flux, we [..⁸⁰] now check that the diabatic mass flux can also be retrieved from the Lagrangian trajectories. Here the mass flux is calculated from the displacement of backward Lagrangian trajectories launched at 380 K. The heating rate is estimated as $\Delta\theta/\Delta t$ where $\Delta\theta$ is the variation of the potential temperature along the trajectory during an interval Δt after the launch[..⁸¹]. Here the interval Δt is always a multiple of 24 h [..⁸²] to ensure that the averages are taken over an integer number of daily cycles. The density σ is calculated from the ERA-Interim at the location and time of the launch. Actually, due to combined horizontal and vertical motion, and the inclination of potential temperature surfaces with respect to isobars, pressure can temporarily decrease for a descending parcel with increasing potential temperature. As a general rule, pressure undergoes much larger fluctuations than potential temperature along a Lagrangian trajectory. The mass flux over a domain Γ can then be calculated as a sum over all parcels belonging to this domain

$$M_{\text{back}} = \sum_{i \in \Gamma} \sigma_i \frac{\Delta\theta_i}{\Delta t} \delta s_i \quad (6)$$

where δs_i is the surface of the $0.5 \times 0.5^\circ$ element associated with the parcel. The monthly upward flux $M_{\text{back}}^\uparrow(\phi)$ is calculated by averaging over each month the individual parcel flux at each location over the grid, selecting all grid points where the monthly flux is positive and then averaging in longitude.

Figure 11 compares the temporal evolution of M_{diab}^\uparrow and M_{back}^\uparrow over the years 2005 to 2008. It is clear that the Lagrangian estimates with the delays $\Delta t = 24$ or 48 h provides an accurate estimate of the upward flux at all latitudes.

⁸⁰removed: know

⁸¹removed: taken as

⁸²removed: in order to average

5.2 Regional distribution of the upward mass flux

Based on these premises we calculate how the monthly upward flux is distributed among regions in the following way. First ^[.83] all the grid points on the 380 K surface within the domain 40° S–40° N where the monthly average flux is positive are selected for each month. Then the contributions of parcels originating from ^[.84] each source region are quantified. Then the mass flux of each region is calculated ^[.85] as the sum of mass fluxes at the selected grid points from which backward trajectories link to this region as the source. The sum of all these regional contributions makes the convective upward flux at the 380 K surface $M_{\text{conv}}^{\uparrow}$. In this procedure, a trajectory is labeled with the time of its launch on the 380 K surface (unlike the calculations leading to Fig. 2 where the trajectories ^[.86] are labeled according to their arrival or departure from clouds).

Figure 11 shows that $M_{\text{conv}}^{\uparrow}$ basically explains the seasonal variations of $M_{\text{back}}^{\uparrow}$ ^[.87]. The difference between these two estimates has a mean of $4.8 \times 10^9 \text{ kg s}^{-1}$ and a standard deviation of $9 \times 10^8 \text{ kg s}^{-1}$ in the 40° S–40° N band ^[.88]. As a result, the ratio $M_{\text{conv}}^{\uparrow}/M_{\text{back}}^{\uparrow}$ varies between 78.7% in winter to 80.5% in summer (80% on the average). The non convective contribution is mostly accounted by parcels which are in-mixed in the TTL from the extra-tropics (Ploeger et al., 2012).

The mean annual cycle is shown in Fig. 12. ^[.89] For each region, it exhibits a maximum ^[.90] and a minimum ^[.91] which lag by about one month with respect to the ^[.92]

⁸³removed: the grid point

⁸⁴removed: a given region is sorted out and the total contribution from this given

⁸⁵removed: by summing its contributions over all the

⁸⁶removed: where labeled with

⁸⁷removed: with a difference

⁸⁸removed: This difference

⁸⁹removed: It

⁹⁰removed: in February

⁹¹removed: in September which are shifted

⁹²removed: maximum and minimum of the source curve in

corresponding source curve Fig. 2a, where the time axis is that of the intersection with convection. This is particularly clear for SAP with a maximum in February and a minimum in September that determines whose of the total convective flux. The delay is consistent with the distribution of transit time among the main sources (see Table 2).

It is only for AML that the lag is not clear, but this is also a region with short transit times. The larger winter to summer modulation of $M_{\text{conv}}^{\uparrow}$ than the total source curve in Fig. 2a suggests that this modulation is mostly due to variations of transport properties within the TTL rather than to a modulation of the properties of convective sources.

Figure 12 shows the same hierarchy among source regions as Fig. 2a with enhanced domination of the SAP contribution over the year which accounts for 39 % of the total $M_{\text{conv}}^{\uparrow}$ flux while NAPO accounts for 18 % (see Table 4). If one adds NCP, CAM which is mostly oceanic, and the small contribution from the Atlantic, Indian Ocean and South East Pacific, we see that the contribution from oceanic regions (which include some large islands) is 74.5 % a proportion in pair with the surface covered by the oceans.

We stress that the flux $M_{\text{conv}}^{\uparrow}$ is the mass flux crossing the 380 K surface that originates from the region of convective outflow within the TTL but is not necessarily purely made of air processed by convection. This air mass contains convective air detrained from the clouds in the vicinity but is also mixed with environment air which may have been transported from distant and earlier convective sources and partly originates from in mixing or extra-tropical lower stratosphere transported into the tropics (Ploeger et al., 2012). Therefore $M_{\text{conv}}^{\uparrow}$ must be taken as an upper bound of the flux of convectively processed air.

6 Summary and outlook

We have shown that a consistent vertical distribution of convective sources of stratospheric air over the tropical regions is obtained from backward and forward diabatic trajectories in the TTL.

The seasonal cycle of sources is binary with a domination of the single South Asia Pacific region (SAP) from November to April and a more complicated pattern dominated by the regions of the Asian monsoon from June to September.

The distribution of sources among regions is qualitatively robust to uncertainties in the method and the data but the quantitative distribution is somewhat sensitive to the representation of cloud tops and [..⁹³]the reanalysis used to drive the trajectories. Generally, increasing the weight of highest clouds shifts the distribution of sources towards higher altitude and wider vertical dispersion, [..⁹⁴]but it does not change the proportion of backward trajectories meeting a cloud.

There is a pronounced seasonal cycle of the monthly average upward mass flux across the 380 K surface, with a maximum in February and a minimum in September, which is shifted by about month to the seasonal cycle of sources, due to the mean time of transit of parcels across the TTL.

The forward transit times are on the average of one month with a significant standard deviation of about 15 days[..⁹⁵]. Transit times are shorter from convection over continental Asia during [..⁹⁶]boreal summer (3 weeks from Asia Mainland (AML) and two weeks for Tibet). The backward transit times are longer[..⁹⁷]. The discrepancy between forward and backward transit times is most pronounced for transport from convection over Tibet.

One of the main [..⁹⁸]motivations of this study was to study how air parcels detrained from clouds find their way across the LZRH. Our results however show that the sources are mostly [..⁹⁹](80%) located slightly above the LZRH, but not [..¹⁰⁰]by much. This implies that only the small percentage of convective events penetrating high enough in the TTL is rele-

⁹³removed: depend also on the reanalysis

⁹⁴removed: while

⁹⁵removed: but

⁹⁶removed: summer where they are shorter

⁹⁷removed: and this discrepancy culminates for

⁹⁸removed: motivation

⁹⁹removed: located

¹⁰⁰removed: far, for about 80%

vant as stratospheric source. The proximity of sources to the LZRH [¹⁰¹] provides evidence that the vertical flow separation at this level is an important factor [¹⁰²] in determining the distribution of the sources. This will be further demonstrated in a companion paper.

The high contribution of the tallest clouds [¹⁰³] raise a concern about the lack of representation of small-scale convective features in the CLAUS dataset and in the ERA-Interim heating rates. In particular, CLAUS captures very well the large anvils that form at the top of convective systems but misses the short and elusive overshooting events over these anvils. The velocities and radiative heating rates from reanalysis also miss the cross isentropic mixing provided by such event or represent them with very crude parameterisations.

The comparison between three modern reanalysis show that the heating rates can differ substantially in the TTL (see Wright and Fueglistaler (2013)) with significant consequences on its distributions of sources. While JRA-55 remains fairly close to the ERA-Interim, MERRA shifts the sources by up to 12 K. The discrepancy is even larger with the calculations of Bergman et al. (2012, 2015) who uses heating rates based on the observed distribution of clouds (Yang et al., 2010). The main differences are during summer season. Bergman et al. (2012) find a proportion of backward trajectories reaching clouds falling to 15% in summer from 60% in winter. We find instead a maximum a maximum in April (88.7%) and a minimum in July (84.7%) with ERA-Interim. The second important difference is in the fact that continental sources in Asia prevail over oceanic sources during summer in Bergman et al. (2012, 2015).

The Yang et al. (2010) heating rates are calculated from the mean distribution of clouds. As such, due to the non linearity of radiative transfer, they may differ from the mean heating rates calculated from the varying distribution of clouds. During winter, Yang et al. (2010) find that the cloud radiative effect is mostly heating in the West Pacific at 15 and 16 km. During summer, on the contrary, they find a strong cooling effect of clouds at 16 km in the Bay of Bengal and the South China Sea. Such a difference over oceanic regions between

¹⁰¹ removed: is providing

¹⁰² removed: to determine

¹⁰³ removed: rise

winter and summer is not found in any of the reanalyses, but a cooling around 350-360 K is displayed over most convective regions by MERRA and MERRA-2.

It is actually very difficult to compare the calculated heating rates with observations. However, the integrated cloud radiative effect (CRE) of the three reanalysis considered here has been evaluated against CERES observations by Li and Mao (2015) who found that "spatial correlation of CREs and TOA upward radiation fluxes in ERA-Interim is the best among the three reanalyses" in spite of some discrepancies in the global mean CRE. Li and Mao (2015) also notice that all three reanalyses have difficulties to reproduce summer CREs over East Asia, a conclusion also supported by Wang et al. (2014) based on a study of radiation budgets in AMIP-5 models. Therefore we are led to conclude that the heating rates and the radiative effect of clouds in the Asian monsoon region are still a puzzle that requires further investigations.

Our study corroborates the special role of the Tibetan plateau in providing air to the Asian Monsoon anticyclone (AMA). We find that 87% of the AMA air originates from continental Asia which agrees very well with the finding of Heath and Fuelberg (2014) (90%), but our results differ from these authors by giving a much stronger weight to the Asian continental regions outside Tibet (AML). We find, however, that this proportion is sensitive to the representation of cloud tops and that it varies a lot within the literature: Fu et al. (2006) and [..¹⁰⁴]Wright et al. (2011) find like Heath and Fuelberg (2014) a prevalent role of [..¹⁰⁵]the Tibetan plateau while Aschmann et al. (2009) and Devasthale and Fueglistaler (2010) meet [..¹⁰⁶]our conclusions. Vogel et al. (2015) finds that that during mid-summer 2012 the AMA is fed mostly from continental sources over North India and the Tibetan plateau, and stress the role of the south boundary of the AMA as a transport barrier. Nevertheless, Tibet is characterized by a very high efficiency at carrying air from the top of the clouds to the 380 K surface and short transit times. This provide a hint that Tibet is a sensitive area

¹⁰⁴removed: Wright and Fueglistaler (2013)

¹⁰⁵removed: Tibet

¹⁰⁶removed: out conclusions.

for increase of air pollution as boundary layer compounds processed by convection can be carried efficiently and rapidly to the stratosphere.

It remains that Tibet is an overall small contributor to the global transport into the stratosphere because most of the air entering the stratosphere during summer is not processed inside the AMA but is transported around, separating the location of [..¹⁰⁷]convection from the entry point into the stratosphere [..¹⁰⁸](Bannister et al., 2004; James et al., 2008; Park et al., 2009; Vogel et al., 2015).

Our results can be compared with those of Chen et al. (2012) and Orbe et al. (2015). Chen et al. (2012) uses kinematic trajectories from the boundary layer focusing on the summer season while Orbe et al. (2015) analyses impulse tracers transported by a general circulation model and provides a whole year analysis. Although both use a different set of regions within the tropics, our results regarding the regional distribution of sources and their seasonal variations are basically consistent with these two studies. There are discrepancies, however, in the transit time scales. Chen et al. (2012) shows a distribution of transit times from the boundary layer to the stratosphere which has a strong modal peak over Asia at about 3 days while, on the contrary, Orbe et al. (2015) shows a distribution which peaks at about two months. Our results lay somewhat between but such a discrepancy [..¹⁰⁹]highlights an issue that needs to be solved [..¹¹⁰]. Transit times are an important [..¹¹¹]factor in understanding and modelling the behavior of many short-lived chemical species in the TTL and the lower stratosphere.

In spite of displaying the highest cloud tops (Liu and Zipser, 2005), the continental convection above Africa (Af) and South America (SAm) is a [..¹¹²]weaker provider than the oceanic convection of SAP and North Asian Pacific Ocean (NAPO). However, our conclu-

¹⁰⁷removed: convective

¹⁰⁸removed: (Bannister et al., 2004; James et al., 2008; Park et al., 2009)

¹⁰⁹removed: creates

¹¹⁰removed: as transit

¹¹¹removed: necessary ingredient to understand and model the behaviour of many short live

¹¹²removed: much

sions do not account for the effect of small-scale overshoots which are more commonly observed over these regions than over the oceans (Corti et al., 2008; Liu et al., 2010) due to the larger available convective energy [over land](#).

Acknowledgements. These results were obtained using the CLAU archive held at the British Atmospheric Data Centre, produced using ISCCP source data distributed by the NASA Langley Data Center. We acknowledge support from the EU 7th Framework Program under grant 603557 [¹¹³StratoClim]. We acknowledge the ECMWF and the Japanese Meteorological Agency for providing ERA-Interim and JRA-55 reanalysis data. JRA-55 data was retrieved from the NCAR C1SL RDA. The MERRA and MERRA2 data used in this study/project have been provided by the Global Modeling and Assimilation Office (GMAO) at NASA Goddard Space Flight Center. We acknowledge G. Sèze for useful discussions and help.

References

- Abalos, M., Randel, W. J., and Serrano, E.: Variability in upwelling across the tropical tropopause and correlations with tracers in the lower stratosphere, *Atmos. Chem. Phys.*, 12, 11505–11517, doi:10.5194/acp-12-11505-2012, 2012.
- Adler, R. F. and Mack, R. A.: Thunderstorm Cloud Top Dynamics as Inferred from Satellite Observations and a Cloud Top Parcel Model, *J. Atmos. Sci.*, 43, 1945–1960, doi:10.1175/1520-0469(1986)043<1945:TCTDAI>2.0.CO;2, 1986.
- Andrews, D. G., Holton, J. R., and Leovy, C. B.: Middle atmosphere dynamics, no. v. 40 in *International geophysics series*, Academic Press, Orlando, 1987.
- Appenzeller, C., Holton, J. R., and Rosenlof, K. H.: Seasonal variation of mass transport across the tropopause, *J. Geophys. Res.*, 101, 15071–15071, doi:10.1029/96JD00821, 1996.
- Aschmann, J., Sinnhuber, B.-M., Atlas, E. L., and Schauffler, S. M.: Modeling the transport of very short-lived substances into the tropical upper troposphere and lower stratosphere, *Atmos. Chem. Phys.*, 9, 9237–9247, doi:10.5194/acp-9-9237-2009, 2009.
- Bannister, R., O'Neill, A., Gregory, A., and Nissen, K.: The role of the south-east Asian monsoon and other seasonal features in creating the “tape-recorder” signal in the Unified Model, *Q. J. Roy. Meteorol. Soc.*, 130, 1531–1554, doi:10.1256/qj.03.106, 2004.

¹¹³removed: “

- Bergman, J. W., Jensen, E. J., Pfister, L., and Yang, Q.: Seasonal differences of vertical-transport efficiency in the tropical tropopause layer: On the interplay between tropical deep convection, large-scale vertical ascent, and horizontal circulations, *J. Geophys. Res.*, 117, D05302, doi:10.1029/2011JD016992, 2012.
- Bergman, J. W., Fierli, F., Jensen, E. J., Honomichl, S., and Pan, L. L.: Boundary layer sources for the Asian anticyclone: Regional contributions to a vertical conduit, *J. Geophys. Res.-Atmos.*, 118, 2560–2575, doi:10.1002/jgrd.50142, 2013.
- Bergman, J. W., Pfister, L., and Yang, Q.: Identifying robust transport features of the upper tropical troposphere, *J. Geophys. Res.*, 120, 6758–6776, doi:10.1002/2015JD023523, 2015.
- Chen, B., Xu, X. D., Yang, S., and Zhao, T. L.: Climatological perspectives of air transport from atmospheric boundary layer to tropopause layer over Asian monsoon regions during boreal summer inferred from Lagrangian approach, *Atmos. Chem. Phys.*, 12, 5827–5839, doi:10.5194/acp-12-5827-2012, 2012.
- Corti, T., Luo, B. P., Fu, Q., Vömel, H., and Peter, T.: The impact of cirrus clouds on tropical troposphere-to-stratosphere transport, *Atmos. Chem. Phys.*, 6, 2539–2547, doi:10.5194/acp-6-2539-2006, 2006.
- Corti, T., Luo, B. P., de Reus, M., Brunner, D., Cairo, F., Mahoney, M. J., Martucci, G., Matthey, R., Mitev, V., dos Santos, F. H., Schiller, C., Shur, G., Sitnikov, N. M., Spelten, N., Vössing, H. J., Borrmann, S., and Peter, T.: Unprecedented evidence for deep convection hydrating the tropical stratosphere, *Geophys. Res. Lett.*, 35, L10810, doi:10.1029/2008GL033641, 2008.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., Van de Berg, L. V. D., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J. J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. Roy. Meteorol. Soc.*, 137, 553–597, doi:10.1002/qj.828, 2011.
- Devasthale, A. and Fueglistaler, S.: A climatological perspective of deep convection penetrating the TTL during the Indian summer monsoon from the AVHRR and MODIS instruments, *Atmos. Chem. Phys.*, 10, 4573–4582, doi:10.5194/acp-10-4573-2010, 2010.
- Folkens, I., Loewenstein, M., Podolske, J., Oltmans, S. J., and Proffitt, M.: A barrier to vertical mixing at 14 km in the tropics: Evidence from ozonesondes and aircraft measurements, *J. Geophys. Res.*, 104, 22095–22095, doi:10.1029/1999JD900404, 1999.

- Fu, Q., Hu, Y., and Yang, Q.: Identifying the top of the tropical tropopause layer from vertical mass flux analysis and CALIPSO lidar cloud observations, *Geophys. Res. Lett.*, 34, L14813, doi:10.1029/2007GL030099, 2007.
- Fu, R., Hu, Y., Wright, J. S., Jiang, J. H., Dickinson, R. E., Chen, M., Filipiak, M., Read, W. G., Waters, J. W., and Wu, D. L.: Short circuit of water vapor and polluted air to the global stratosphere by convective transport over the Tibetan Plateau, *P. Natl. Acad. Sci. USA*, 103, 5664–5669, doi:10.1073/pnas.0601584103, 2006.
- Fueglistaler, S., Wernli, H., and Peter, T.: Tropical troposphere-to-stratosphere transport inferred from trajectory calculations, *J. Geophys. Res.*, 109, D03108, doi:10.1029/2003JD004069, 2004.
- Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkins, I., Fu, Q., and Mote, P. W.: Tropical tropopause layer, *Rev. Geophys.*, 47, RG1004, doi:10.1029/2008RG000267, 2009.
- Gottelman, A., Salby, M. L., and Sassi, F.: Distribution and influence of convection in the tropical tropopause region, *J. Geophys. Res.*, 107, 4080–4080, doi:10.1029/2001JD001048, 2002.
- Gottelman, A., Lauritzen, P. H., Park, M., and Kay, J. E.: Processes regulating short-lived species in the tropical tropopause layer, *J. Geophys. Res.*, 114, D13303, doi:10.1029/2009JD011785, 2009.
- Hall, T. M. and Holzer, M.: Advective-diffusive mass flux and implications for stratosphere-troposphere exchange, *Geophys. Res. Lett.*, 30, 1222, doi:10.1029/2002GL016419, 2003.
- Hamann, U., Walther, A., Baum, B., Bennartz, R., Bugliaro, L., Derrien, M., Francis, P. N., Heidinger, A., Joro, S., Kniffka, A., Le Gléau, H., Lockhoff, M., Lutz, H.-J., Meirink, J. F., Minnis, P., Palikonda, R., Roebeling, R., Thoss, A., Platnick, S., Watts, P., and Wind, G.: Remote sensing of cloud top pressure/height from SEVIRI: analysis of ten current retrieval algorithms, *Atmospheric Measurement Techniques*, 7, 2839–2867, doi:10.5194/amt-7-2839-2014, 2014.
- Heath, N. K. and Fuelberg, H. E.: Using a WRF simulation to examine regions where convection impacts the Asian summer monsoon anticyclone, *Atmos. Chem. Phys.*, 14, 2055–2070, doi:10.5194/acp-14-2055-2014, 2014.
- Highwood, E. J. and Hoskins, B. J.: The tropical tropopause, *Q. J. Roy. Meteorol. Soc.*, 124, 1579–1604, doi:10.1002/qj.49712454911, 1998.
- Hodges, K. I., Chappell, D. W., Robinson, G. J., and Yang, G.: An improved algorithm for generating global window brightness temperatures from multiple satellite infrared imagery, *J. Atmos. Ocean. Tech.*, 17, 1296–1312, 2000.
- Holton, J. R., Haynes, P. H., McIntyre, M. E., Douglass, A. R., Rood, R. B., and Pfister, L.: Stratosphere-troposphere exchange, *Rev. Geophys.*, 33, 403, doi:10.1029/95RG02097, 1995.

- Holzer, M. and Hall, T. M.: Transit-Time and Tracer-Age Distributions in Geophysical Flows, *J. Atmos. Sci.*, 57, 3539–3558, doi:10.1175/1520-0469(2000)057<3539:TTATAD>2.0.CO;2, 2000.
- James, R., Bonazzola, M., Legras, B., Surlbed, K., and Fueglistaler, S.: Water vapor transport and dehydration above convective outflow during Asian monsoon, *Geophys. Res. Lett.*, 35, L20810, doi:10.1029/2008GL035441, 2008.
- Kobayashi, S., Ota, Y., Harada, Y., Ebata, A., Moriya, M., Onoda, H., Onogi, K., Kamahori, H., Kobayashi, C., Endo, H., Miyaoka, K., and Takahashi, K.: The JRA-55 Reanalysis: General Specifications and Basic Characteristics, *J. Meteorol. Soc. Jpn. Ser. II*, 93, 5–48, doi:10.2151/jmsj.2015-001, 2015.
- Kwon, E.-H., Sohn, B.-J., Schmetz, J., and Watts, P.: Intercomparison of height assignment methods for opaque clouds over the tropics, *Asia-Pacific Journal of Atmospheric Sciences*, 46, 11–19, doi:10.1007/s13143-010-0002-7, 2010.
- Legras, B., Pissot, I., Berthet, G., and Lefèvre, F.: Variability of the Lagrangian turbulent diffusion in the lower stratosphere, *Atmos. Chem. Phys.*, 5, 1605–1622, doi:10.5194/acp-5-1605-2005, 2005.
- Li, Jian-Dong and Mao, Jiang-Yu: A preliminary evaluation of global and East Asian cloud radiative effects in reanalyses, *Atmospheric and Oceanic Science Letters*, 8, 100–106, doi:10.3878/AOSL20140093, 2015.
- Liu, C. and Zipser, E. J.: Global distribution of convection penetrating the tropical tropopause, *J. Geophys. Res.*, 110, D23104, doi:10.1029/2005JD006063, 2005.
- Liu, X. M., Rivière, E. D., Maréchal, V., Durr, G., Hamdouni, A., Arteta, J., and Khaykin, S.: Stratospheric water vapour budget and convection overshooting the tropopause: modelling study from SCOUT-AMMA, *Atmos. Chem. Phys.*, 10, 8267–8286, doi:10.5194/acp-10-8267-2010, 2010.
- Luo, Z., Liu, G. Y., and Stephens, G. L.: CloudSat adding new insight into tropical penetrating convection, *Geophys. Res. Lett.*, 35, L19819, doi:10.1029/2008GL035330, 2008.
- Minnis, P., Yost, C. R., Sun-Mack, S., and Chen, Y.: Estimating the top altitude of optically thick ice clouds from thermal infrared satellite observations using CALIPSO data, *Geophys. Res. Lett.*, 35, L12801, doi:10.1029/2008GL033947, 2008.
- Molod, A., Takacs, L., Suarez, M., and Bacmeister, J.: Development of the GEOS-5 atmospheric general circulation model: evolution from MERRA to MERRA2, *Geoscientific Model Development*, 8, 1339–1356, doi:10.5194/gmd-8-1339-2015, 2015.
- Orbe, C., Waugh, D. W., and Newman, P. A.: Air-mass origin in the tropical lower stratosphere: The influence of Asian boundary layer air, *Geophys. Res. Lett.*, 42, 4240–4248, doi:10.1002/2015GL063937, 2015.

- Park, M., Randel, W. J., Gettelman, A., Massie, S. T., and Jiang, J. H.: Transport above the Asian summer monsoon anticyclone inferred from Aura Microwave Limb Sounder tracers, *J. Geophys. Res.*, 112, D16309, doi:10.1029/2006JD008294, 2007.
- Park, M., Randel, W. J., Emmons, L. K., and Livesey, N. J.: Transport pathways of carbon monoxide in the Asian summer monsoon diagnosed from Model of Ozone and Related Tracers (MOZART), *J. Geophys. Res.*, 114, D08303, doi:10.1029/2008JD010621, 2009.
- Pisso, I. and Legras, B.: Turbulent vertical diffusivity in the sub-tropical stratosphere, *Atmos. Chem. Phys.*, 8, 697–707, doi:10.5194/acp-8-697-2008, 2008.
- Ploeger, F., Konopka, P., Müller, R., Fueglistaler, S., Schmidt, T., Manners, J. C., Grooß, J.-U., Günther, G., Forster, P. M., and Riese, M.: Horizontal transport affecting trace gas seasonality in the Tropical Tropopause Layer (TTL), *J. Geophys. Res.*, 117, D09303, doi:10.1029/2011JD017267, 2012.
- Podglajen, A., Hertzog, A., Plougonven, R., and Žagar, N.: Assessment of the accuracy of (re)analyses in the equatorial lower stratosphere, *J. Geophys. Res.-Atmos.*, 119, 11166–11188, doi:10.1002/2014JD021849, 2014.
- Rienecker, M. M., Suarez, M. J., Gelaro, R., Todling, R., Bacmeister, J., Liu, E., Bosilovich, M. G., Schubert, S. D., Takacs, L., Kim, G.-K., Bloom, S., Chen, J., Collins, D., Conaty, A., da Silva, A., Gu, W., Joiner, J., Koster, R. D., Lucchesi, R., Molod, A., Owens, T., Pawson, S., Pegion, P., Redder, C. R., Reichle, R., Robertson, F. R., Ruddick, A. G., Sienkiewicz, M., and Woollen, J.: MERRA NASA's Modern-Era Retrospective Analysis for Research and Applications, *Journal of Climate*, 24, 3624–3648, doi:10.1175/JCLI-D-11-00015.1, 2011.
- Sherwood, S. C., Chae, J.-H., Minnis, P., and McGill, M.: Underestimation of deep convective cloud tops by thermal imagery, *Geophys. Res. Lett.*, 31, L11102, doi:10.1029/2004GL019699, 2004.
- Shine, K.: Sources and sinks of zonal momentum in the middle atmosphere diagnosed using the diabatic circulation, *Q. J. Roy. Meteorol. Soc.*, 115, 265–292, doi:10.1002/qj.49711548604, 1989.
- Stohl, A., Forster, C., Frank, A., Seibert, P., and Wotawa, G.: Technical note: The Lagrangian particle dispersion model FLEXPART version 6.2, *Atmos. Chem. Phys.*, 5, 2461–2474, doi:10.5194/acp-5-2461-2005, 2005.
- Tzella, A. and Legras, B.: A Lagrangian view of convective sources for transport of air across the Tropical Tropopause Layer: distribution, times and the radiative influence of clouds, *Atmos. Chem. Phys.*, 11, 12517–12534, doi:10.5194/acp-11-12517-2011, 2011.

- Vogel, B., Günther, G., Müller, R., Groß, J.-U., and Riese, M.: Impact of different Asian source regions on the composition of the Asian monsoon anticyclone and on the extratropical lowermost stratosphere, *Atmos. Chem. Phys.*, 15, 13699–13716, doi:10.5194/acp-15-13699-2015, 2015.
- Wang, F., Yang, S., and Wu, T.: Radiation budget biases in AMIP5 models over the East Asian monsoon region, *Journal of Geophysical Research: Atmospheres*, 119, 13,400–13,426, doi:10.1002/2014JD022243, 2014.
- Wright, J. S., Fu, R., Fueglistaler, S., Liu, Y. S., and Zhang, Y.: The influence of summertime convection over Southeast Asia on water vapor in the tropical stratosphere, *Journal of Geophysical Research*, 116, doi:10.1029/2010JD015416, 2011.
- Wright, J. S. and Fueglistaler, S.: Large differences in reanalyses of diabatic heating in the tropical upper troposphere and lower stratosphere, *Atmos. Chem. Phys.*, 13, 9565–9576, doi:10.5194/acp-13-9565-2013, 2013.
- Yang, Q., Fu, Q., and Hu, Y.: Radiative impacts of clouds in the tropical tropopause layer, *Journal of Geophysical Research*, 115, doi:10.1029/2009JD012393, 2010.

Table 1. Characteristic numbers of the vertical distribution of sources for the contributing regions during winter (DJF) and summer (JJA). All the quantities but the last column are **potential temperatures in units of K**. The modal peak is based on the discretized mean histogram shown in Fig. 3. The mean, median and standard deviation are calculated from a cubic spline interpolation over the 340–380 K interval. For each region and each season the upper line refers to backward calculations and the lower line to forward calculations as indicated in the modal peak column. [¹¹⁴]Regions with low contribution have been masked.

Season	Region	LZRH	Modal peak	Mean	Median	Std dev	% above LZRH
DJF	Af	353.4	355 (B)	358.1	356.8	6.3	75.8
			355 (F)	359.5	357.9	6.8	82.5
	SAP	352.6	355 (B)	356.1	355.1	5.2	76.3
			355 (F)	358.7	357.3	5.9	89.5
	SAm	353.6	353 (B)	357.8	356.2	6.8	69.2
			355 (F)	358.7	357.4	6.4	77.1
JJA	Af	355.7	361 (B)	362.7	362	7.2	83
			359 (F)	363.1	362.2	7.2	83.7
	SAP	352.1	351 (B)	354.5	353.3	5.3	62
			353 (F)	356	354.9	5.5	75.3
	AML	359	361 (B)	363.1	362.5	5.7	75.4
			363 (F)	364.9	364.2	5.7	85.9
	NAPO	352.6	357 (B)	358.5	357.7	6.2	83
			359 (F)	360.5	359.6	6.5	90.7
Tibet	367.3	363 (B)	365.4	364.7	5.2	31.4	
		367 (F)	367.9	367.4	4.5	50.8	
NCP	351.3	351 (B)	353.8	352.4	5.5	62	
		353 (F)	355.7	354.5	5.6	80	
CAm	350.7	353 (B)	357.4	355.5	7.4	82.5	
		353 (F)	358	356.3	6.8	90.3	

¹¹⁴removed: The regions

Table 2. Characteristic numbers of the distribution of transit times for the contributing regions during winter (DJF) and summer (JJA). All the quantities are in **units of days**. The modal peak is based on the discretized mean histogram shown in Fig. 4. The mean, median and standard deviation are calculated from a cubic spline interpolation over the 0–90 day interval. For each region and each season the upper line refers to backward calculations and the lower line to forward calculations as indicated in the modal peak column. [..¹¹⁵]Regions with low contribution have been masked.

Season	Region	Modal peak	Mean	Median	Std dev
DJF	Af	26 (B)	36.9	33.6	19.4
		26 (F)	27.5	25.7	14.6
	SAP	22 (B)	34.2	30.8	19.0
		18 (F)	23.4	20.6	13.3
	SAm	30 (B)	36.7	33.7	19.5
		26 (F)	28.2	26.5	14.4
JJA	Af	30 (B)	35.4	32.7	20.0
		26 (F)	29.5	27.8	16.9
	SAP	22 (B)	38.0	33.8	19.7
		22 (F)	30.9	27.6	15.3
	AML	26 (B)	32.5	29.4	19.7
		18 (F)	21.3	19.2	13.7
	NAPO	30 (B)	38.7	35.7	19.6
		26 (F)	28.4	26.6	15.8
	Tibet	14 (B)	29.5	24.3	19.6
		10 (F)	15.1	13	9.8
NCP	22 (B)	37.2	33.3	19.2	
	22 (F)	31.4	28.5	15.3	
CAm	30 (B)	37.9	35.2	19.5	
	30 (F)	32.2	30.2	15.9	

¹¹⁵removed: The regions

Table 3. Ratio of transport efficiency between forward calculations without cloud top offset and with a +1 km cloud top offset. The ratio is averaged over DJF and JJA for the 2005–2008 period and all regions. The values for weakly contributing regions, excluded from Tables 1 and 2, are based on a small number of events and are [¹¹⁶]italicized.

	Af	ITA	SAP	AML	NAPO	CAm	SAm	Tibet	IO	NCP	SEP
DJF	0.45	<i>0.38</i>	0.45	<i>0.33</i>	<i>0.37</i>	<i>0.42</i>	0.46	<i>0.62</i>	<i>0.41</i>	<i>0.43</i>	<i>0.34</i>
JJA	0.49	<i>0.43</i>	0.46	0.36	0.42	0.46	<i>0.45</i>	0.22	<i>0.34</i>	0.46	<i>0.34</i>

¹¹⁶removed: in italic

Table 4. Distribution of annual mass flux, averaged over 2005–2008, for all the regions (in %).

Af	ITA	SAP	AML	NAPO	CAm	SAm	Tibet	IO	NCP	SEP
10.8	2.4	39.2	8	18	7.5	6	0.8	1.2	5.9	0.2

Definitions of the regional boxes




Figure 1. Distribution of the geographical regions within the tropics and the subtropics. In color [\[..¹¹⁷\]](#): [\[..¹¹⁸\]](#) Africa ([\[..¹¹⁹\]](#) Af), [\[..¹²⁰\]](#) Inter Tropical Atlantic (ITA), [\[..¹²¹\]](#) South Asian Pacific ([\[..¹²¹\]](#) SAP), [\[..¹²²\]](#) Asia Mainland (AML), [\[..¹²³\]](#) North Asian-Pacific Ocean (NAPO), [\[..¹²⁴\]](#) Central America (CAm), [\[..¹²⁵\]](#) South America (SAm), [\[..¹²²\]](#) Tibetan plateau ([\[..¹²³\]](#) Tibet), [\[..¹²⁴\]](#) Indian Ocean (IO), [\[..¹²⁵\]](#) North [\[..¹²⁶\]](#) Central Pacific ([\[..¹²⁷\]](#) NCP), and [\[..¹²⁸\]](#) South [\[..¹²⁹\]](#) East Pacific ([\[..¹²⁹\]](#) SEP). Tibet is defined as the region in Asia above 3500 m.

¹¹⁷ removed: , from west to east

¹¹⁸ removed: North Central Pacific

¹¹⁹ removed: NCP

¹²⁰ removed: East

¹²¹ removed: SEP

¹²² removed: Inter Tropical Atlantic

¹²³ removed: ITA

¹²⁴ removed: Africa (Af),

¹²⁵ removed: Asia Mainland (AML), Tibetan plateau (Tibet),

¹²⁶ removed: Asian-Pacific Ocean

¹²⁷ removed: NAPO

¹²⁸ removed: Asian

¹²⁹ removed: SAP




Figure 2. (a) ¹³⁰Color lines: source distribution from backward calculations, calculated as the proportion of backward trajectories reaching a convective top within a given region among all trajectories reaching a convective top. The sum of all the contributions is 100%. Black line: proportion of backward trajectories reaching a convective top. **(b)** Source distribution from forward calculations, calculated as the proportion of forward trajectories reaching the ¹³¹380 K isentropic surface from a given region among all the trajectories reaching ¹³²this surface. **(c)** Efficiency of transport from convective tops, calculated as the proportion of parcels released in a given region that reach the ¹³³380 K isentropic surface in the forward calculations. On the right of **(a)** and **(b)** Source distribution for the inside of the Asian Monsoon Anticyclone averaged over JJA. The statistics are obtained for the period 2005–2008. The time axis refers to the instant grouped by months where a parcel leaves (forward) or meets (backward) a cloud but for the black curve in panel **(a)** for which the time axis is that of the launch of the parcels on the 380 K surface. Curves are plotted for each region, with the color code of Fig. 1.

¹³⁰removed: Source

¹³¹removed: tropopause

¹³²removed: the tropopause

¹³³removed: tropopause




Figure 3. Vertical source distribution of the parcels in winter (DJF) and summer (JJA) for forward and backward calculations as a function of potential temperature of the source. The vertical axis counts the number of parcels calculated as $[..^{134}]$ daily averages summing events at the 8 CLAUS sampling times (then averaged over DJF and JJA and over 2005–2008) and are given in K⁻¹ day⁻¹. Curves are plotted for each region, with the color code of Fig. 1. The arrows on the upper axes indicate the corresponding mean LZRH levels over the season and 2005–2008.

¹³⁴removed: monthly




Figure 4. Distribution of the transit times in winter (DJF) and summer (JJA) for forward and backward calculations. The vertical axis is a [\dots]¹³⁵ probability density function in day^{-1} . Curves are plotted according to the color code of Fig. 1 but only for the active convective regions during each season. In the upper part of each panel, the triangles and the crosses indicate, respectively, the median and the mean for each curve.

¹³⁵removed: normalized




Figure 5. Same as Fig. 3 when the +1 km altitude correction of the cloud tops is not applied.




Figure 6. Same as Fig. 4 when the +1 km altitude correction of the cloud tops is not applied.




Figure 7. Comparison of the vertical source distribution, for ERA-Interim, for winter (DJF) and summer (JJA) between the standard backward calculation for 2005 (upper two panels)[¹³⁶], when the CLAUS pixel size is enlarged by a factor 3 as described in the text (middle two panels), and when the 3-hourly sampling of daily cycle of heating rates is replaced by a time moving average as described in the text (lower two panels).

¹³⁶removed: and




Figure 8. Vertical source distribution for winter (DJF) and summer (JJA) calculated for 2005 with JRA-55 and MERRA reanalysis using the same setup as for ERA-Interim. [¹³⁷]Upper two panels, for ERA-Interim, are the same as those shown on the two upper panels of Fig. [¹³⁸]7.

¹³⁷removed: This is to be compared with

¹³⁸removed: ??




Figure 9. Mean heating rate profiles, as a function of potential temperature, for the four reanalysis : ERA-Interim (blue), MERRA (red), MERRA2 (dashed red) and JRA-55 (black), in $\text{K}^{-1} \text{day}^{-1}$. The upper panel is for January and the lower panel is for July, averaged over 2005–2008. Curves are plotted for each region defined on Fig. 1. The region "I20" represents the 20°S – 20°N band.




Figure 10. Annual variations of the monthly upward mass flux at the 380 K surface, calculated from the ERA-Interim data. M_{diab}^{\uparrow} and $M_{diab,corr}^{\uparrow}$: diabatic mass flux calculated from the radiative heating rates of ERA-Interim respectively with and without the mass conservation correction of the radiative heating rates. M_{kine}^{\uparrow} : kinematic mass flux calculated from the residual mean meridional circulation. All quantities are calculated and averaged over the years 2005 to 2008.




Figure 11. Uncorrected upward diabatic mass flux on the 380 K surface. $M_{\text{diab}}^{\uparrow}$ (red) versus the Lagrangian diabatic mass flux $M_{\text{back}}^{\uparrow}$ on the same surface calculated with delays $\Delta t = 24$ h (green) and $\Delta t = 48$ h (blue). The black curve shows $M_{\text{conv}}^{\uparrow}$, [..¹³⁹] which is the part of $M_{\text{back}}^{\uparrow}$ with $\Delta t = 48$ h [..¹⁴⁰] that originates from backward trajectories encountering a cloud within the previous 3 months.

¹³⁹removed: that it

¹⁴⁰removed: which




Figure 12. Mean annual cycles of monthly upward mass fluxes through the 380 K isentropic surface within the 40° S–40° N band attributed to each source region. Fluxes are based on backward trajectories during 2005–2008. The black curve shows $M_{\text{conv}}^{\uparrow}$. Time is defined relative to back trajectory launch at 380 K, rather than the convective source.

¹⁴¹removed: Distribution of the monthly upward mass flux at the 380

¹⁴²removed: within the band 40

¹⁴³removed: S–40

¹⁴⁴removed: N among the regions calculated from backward trajectories over 2005–2008. The black curve shows $M_{\text{conv}}^{\uparrow}$. The time axis is that of the launch on the 380

¹⁴⁵removed: surface.