

**Fossil and non-fossil source contributions to atmospheric carbonaceous aerosols during extreme spring grassland fires in Eastern Europe**

**V. Ulevicius<sup>1</sup>, S. Byčenkienė<sup>1</sup>, C. Bozzetti<sup>2</sup>, A. Vlachou<sup>2</sup>, K. Plauškaitė<sup>1</sup>, G. Mordas<sup>1</sup>, V. Dudoitis<sup>1</sup>, G. Abbaszade<sup>3</sup>, V. Remeikis<sup>1</sup>, A. Garbaras<sup>1</sup>, A. Masalaite<sup>1</sup>, J. Blees<sup>2</sup>, R. Fröhlich<sup>2</sup>, K. R. Dällenbach<sup>2</sup>, F. Canonaco<sup>2</sup>, J. G. Slowik<sup>2</sup>, J. Dommen<sup>2</sup>, R. Zimmermann<sup>3,4</sup>, J. Schnelle-Kreis<sup>3</sup>, G. A. Salazar<sup>5</sup>, K. Agrios<sup>5,6</sup>, S. Szidat<sup>5</sup>, I. El Haddad<sup>2</sup> and A. S. H. Prévôt<sup>2</sup>**

[1] Department of Environmental Research, SRI Center for Physical Sciences and Technology, Vilnius, Lithuania

[2] Laboratory of Atmospheric Chemistry, Paul Scherrer Institute (PSI), 5232 Villigen, Switzerland

[3] Helmholtz Zentrum München, German Research Center for Environmental Health (GmbH), Joint Mass Spectrometry Centre, Cooperation Group Comprehensive Molecular Analytics and Helmholtz Virtual Institute of Complex Molecular Systems in Environmental Health – Aerosol and Health (HICE), 85764 Neuherberg, Germany

[4] Analytical Chemistry & Joint Mass Spectrometry Centre, Institute of Chemistry, University of Rostock, Dr.-Lorenz-Weg 1, D-18051 Rostock /Germany

[5] Department of Chemistry and Biochemistry & Oeschger Centre for Climate Change Research, University of Bern, 3012 Bern, Switzerland

[6] Laboratory of Radiochemistry and Environmental Chemistry, PSI, 5232 Villigen, Switzerland

Correspondence to: V. Ulevicius and A. S. H. Prévôt ([uleviev@ktl.mii.lt](mailto:uleviev@ktl.mii.lt), [andre.prevot@psi.ch](mailto:andre.prevot@psi.ch))

## Abstract

In early spring the Baltic region is frequently affected by high pollution events due to biomass burning in that area. Here we present a comprehensive study to investigate the impact of biomass/grass burning (BB) on the evolution and composition of aerosol in Preila, Lithuania, during springtime open fires. Non-refractory submicron particulate matter (NR-PM<sub>1</sub>) was measured by an Aerodyne aerosol chemical speciation monitor (ACSM) and a source apportionment with the multilinear engine (ME-2) running the positive matrix factorization (PMF) model was applied to the organic aerosol fraction to investigate the impact of biomass/grass burning. Satellite observations over regions of biomass burning activity supported the results and identification of air mass transport to the area of investigation. Sharp increases in biomass burning tracers, such as levoglucosan up to 683 ng m<sup>-3</sup> and black carbon (BC) up to 17 µg m<sup>-3</sup> were observed during this period. A further separation between fossil and non-fossil primary and secondary contributions was obtained by coupling ACSM PMF results and radiocarbon (<sup>14</sup>C) measurements of the elemental (EC) and organic (OC) carbon fractions. Non-fossil organic carbon (OC<sub>nf</sub>) was the dominant fraction of PM<sub>1</sub>, with the primary (POC<sub>nf</sub>) and secondary (SOC<sub>nf</sub>) fractions contributing 26–44% and 13–23% to the [total carbon \(TC\)](#), respectively. 5–8% of the TC had a primary fossil origin (POC<sub>f</sub>), whereas the contribution of fossil secondary organic carbon (SOC<sub>f</sub>) was 4–13%. Non-fossil EC (EC<sub>nf</sub>) and fossil EC (EC<sub>f</sub>) ranged from 13–24% and 7–12%, respectively. Isotope ratios of stable carbon and nitrogen isotopes were used to distinguish aerosol particles associated with solid and liquid fossil fuel burning.

## 1 Introduction

On a global scale wood or grass burning is a major source of organic aerosol (Crutzen et al., 1979; Levine, 1996). Approximately 90% of vegetation burning is caused by human-induced fires (Baldini et al., 2002) and only a minor fraction derives from natural processes such as lightning. The composition of biomass smoke depends on the type of wood, combustion conditions (flaming versus smoldering), and ambient weather conditions (Weimer et al., 2008, Grieshop et al., 2009, Hawkins and Russell, 2010 and Akagi et al., 2012). Fine particles emitted from biomass burning include directly emitted primary particles (POA) and secondary organic aerosols (SOA), formed in the atmosphere as the plume ages through photochemical processes driven by sunlight (Capes et al., 2008, Heringa et al., 2011).

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

1 Many studies have revealed that organic matter (OM) is the largest fraction of ambient fine  
2 particles, typically comprising 20–90 % of the submicron particulate mass (Jimenez et al.,  
3 2009). Factor analysis of aerosol mass spectra from [the](#) Aerodyne aerosol mass spectrometer  
4 enables the deconvolution of OM into different factors based on their mass spectral  
5 fingerprints (Lanz et al., 2007; Aiken et al., 2009; Ulbrich et al., 2009). Such results provided  
6 valuable insights into the source and transformation processes of organic aerosols (OA) in the  
7 atmosphere (Lanz et al., 2010; Ng et al., 2011; Hildebrandt et al., 2011; Canonaco et al. 2013;  
8 Bougiatioti et al., 2014; Huang et al., 2014).

9 The main type of biomass burning in Lithuania and surrounding countries in early spring  
10 during the last years is illegal [grass burning](#) for land clearing (Ulevicius et al., 2010b;  
11 Byčenkienė et al. 2013). The north-east European countries are considered to influence  
12 significantly the microphysical, chemical and optical properties of the aerosol in the Baltic  
13 Sea region (Kikas et al., 2008; Zawadzka et al., 2013; Mann et al., 2014, Beddows et al.,  
14 2014). Long-term measurements of carbonaceous aerosols performed in this area by Ulevicius  
15 et al. (2010a, 2010b) and Byčenkienė et al. (2011, 2013) reported a yearly occurrence of high  
16 biomass burning organic aerosol (BBOA) levels during March–April related to regional  
17 transport from the Kaliningrad region, Ukraine and [the](#) southwestern part of Russia  
18 surrounding the Black Sea, but information on the nature and chemical composition of the  
19 biomass burning aerosol in Lithuania is still limited. There has been no systematic  
20 investigation of the impact of biomass burning on ambient organic aerosol levels in this  
21 region, and a quantitative estimate is needed to understand the possible impacts of BBOA on  
22 air quality in the south-eastern Baltic Sea region.

23 In many studies levoglucosan was used to assess the contribution of biomass-burning smoke  
24 to the aerosol mass concentrations (Puxbaum et al., 2007). A number of source emission  
25 studies reported that levoglucosan is not a useful tracer after long-range transport due to its  
26 transformation (Hoffmann et al., 2010, Hennigan et al., 2010; Mochida et al., 2010). In  
27 contrast to levoglucosan, determination of radiocarbon ( $^{14}\text{C}$ ) offers a unique possibility for  
28 source apportionment of carbonaceous aerosol particles, as it unambiguously distinguishes  
29 fossil from non-fossil emissions (e.g., Currie, 2000; Ceburnis et al., 2011).

30 For this study, in the framework of the Lithuanian-Swiss Cooperation Programme joint  
31 research project (AEROLIT), an [aerosol chemical speciation monitor \(ACSM\)](#) was deployed  
32 in a background area of the South Baltic Sea to measure airborne submicron particles for one  
33 month during a period of frequent grass burning pollution. The main findings include

Deleted :

investigation of OA components (Sects. 3.1–3.2), molecular markers (Sect. 3.2), source apportionment of [elemental and organic carbon](#) (EC and OC) using  $^{14}\text{C}$  data and positive matrix factorization (PMF) of the ACSM organic mass spectra (Sect. 3.3).

## 2 Methods

### 2.1. Site description and filter sampling

Continuous air monitoring and time integrated particulate matter sampling were carried out in March 2014 in Preila, Lithuania (55° 55' N, 21° 04' E 5 m a.s.l.) (Fig. 1). Preila is a representative coastal background site, an ideal location for studying the long-range transport of air pollutants in the South-eastern Baltic region due to the absence of significant local sources (Fig. 1, Table 1). It served as a “super site” for the EUSAAR-EU-funded (Integrated Infrastructures Initiatives) project. During the measurement period, strong biomass burning activities were observed on 9–10 March 2014. A high-volume sampler (Digital model Aerosol Sampler DHA-80, 500  $\text{L min}^{-1}$ ) was used to collect  $\text{PM}_{10}$  aerosol particles onto 150 mm diameter Pallflex quartz fibre filters (pre-baked for 24 h at 550 °C) over a 24-hour sampling period. Filters were stored in a freezer (at -20 °C) immediately after sampling.

### 2.2. Instrumentation

#### 2.2.1. Aerosol Chemical Speciation Monitor and data analysis

An ACSM (Aerodyne Research, Inc., Billerica, MA, USA) was deployed to measure  $\text{PM}_{10}$  components in Preila (Fig. 1, Sect. 2.1). A  $\text{PM}_{10}$  impactor-type inlet was utilized to remove coarse particles from the sample stream. The sampling air ( $1.1 \text{ L min}^{-1}$ ) passed through a vertical 2.5 m long stainless steel tube with a 6 mm i.d. and a Nafion dryer (MD-110-12S-4, PermaPure LLC, Toms River, NJ, USA) before reaching the device. Aerosol particle diffusion losses in the sampling line were less than 4.0% for particles from 40 nm to 1  $\mu\text{m}$  according to Gormley and Kennedy (Baron and Willike, 2001) and the relative humidity lower than 50% (by SATO model SK-L200TH). Thus, the used sampling line and ambient relative humidity did not affect aerosol mass concentration measured by ACSM. The transported aerosol flow was split and directed to a scanning mobility particle sizer (model 19.3.09 IFT/TT (TROPOS, Leipzig, Germany) and to the ACSM. In the ACSM particles were directed onto a resistively heated surface at ~600 °C where NR- $\text{PM}_{10}$  components are flash vaporized and the resulting gases are subsequently ionized by 70 eV electron impact. ACSM

Deleted: 13

Deleted: 1

Deleted:

Deleted:

Deleted: s

1 was operated with a time resolution of ~28 min (for typical aerosol loadings, i.e. several  $\mu\text{g}$   
2  $\text{m}^{-3}$ ) and a scan rate of  $220 \text{ ms amu}^{-1}$  from  $m/z$  10 to 140 (approximately 31.9 s per scan and  
3 1.126 s pause), 56 scans and data interval 30 min. The data acquisition software used was  
4 DAQ 1.4.4.4. The mass concentrations and mass spectra were processed using ACSM  
5 standard data analysis software (v 1.5.3.0).  
6 The instrument was calibrated using ammonium sulphate and ammonium nitrate. The  
7 determined calibration parameters were response factor (RF)  $\text{RF}_{\text{NO}_3} = 2.75 \cdot 10^{-11}$  and relative  
8 ionization efficiency (RIE)  $\text{RIE}_{\text{NH}_4} = 6.16$ ,  $\text{RIE}_{\text{SO}_4} = 0.92$ . The  $\text{RIE}_{\text{Org}} = 1.4$ ,  $\text{RIE}_{\text{Chl}} = 1.3$  were  
9 set as default. However, the ACSM collection efficiency varies depending on the acidity of  
10 aerosol particles, aerosol composition, and particle phase water (Matthew et al., 2008). Many  
11 atmospheric aerosol studies reported reasonable agreement and linear correlations were  
12 obtained with other measurements by using a collection efficiency of 0.5 (Aiken et al. 2009;  
13 Timonen et al. 2010). Middlebrook et al. (2012) had proposed a collection efficiency  
14 calculation method. The collection efficiency for each measurement and daily mean CE  
15 values were calculated. The CE variation was small during the entire measurement campaign  
16 (March 2014), so the determined mean CE value was 0.52 with a standard deviation of 0.08,  
17 which is very close to other studies (Aiken et al. 2009; Timonen et al. 2010). This is not  
18 surprising because the sampled aerosol was dried to  $\text{RH} < 50\%$ ; moreover, the nitrate fraction  
19 was quite low (15% on average) and a high acidity of aerosols was not expected at Preila  
20 station (EMEP). Thus, we used the  $\text{CE} = 0.52$  in our investigation. The time series of organic  
21 aerosol mass spectra were processed using PMF analysis.

## 22 2.2.2. PMF analysis

23 The ACSM measured data were averaged to 1-hour time resolution. A graphical user interface  
24 SoFi (Source Finder) (Canonaco et al., 2013), developed at Paul Scherrer Institute was used to  
25 perform PMF for the source apportionment of the non-refractory OA mass spectra collected  
26 during March 2014. Only signals at  $m/z < 120$  were used for PMF analysis (Paatero and  
27 Tapper, 1994; Paatero, 1997) due to the following reasons: 1) the signals above  $m/z > 120$   
28 account for a minor fraction of total signal, 2) the  $m/z$ 's  $> 120$  have larger uncertainties  
29 because of poor ion transmission and the large interferences of naphthalene signals on some  
30  $m/z$ 's (e.g.,  $m/z$  127, 128, and 129) (Sun et al., 2012). A 2-factor solution including a Primary  
31 Organic Aerosol factor (POA), and a Secondary Organic Aerosol factor (SOA) was selected  
32 for this study. 20 different PMF runs were performed using a bootstrapping approach

Deleted: ,

Deleted: works

Deleted:

Deleted: ,

Deleted: )

(Davison and Hinkley, 1997). The bootstrap creates new input data matrices by randomly resampling measured mass spectra from the original input matrices. Moreover, each PMF bootstrap run is initiated from a different pseudorandom starting-point of the algorithm (seed). The bootstrapping approach, together with the seed approach allows a reasonable exploration of the PMF solution space (Paatero et al., 2014). Higher order solutions (3 factors) were explored yielding additional primary profiles, without a significant modification of the secondary contributions. Moreover the retrieved additional profiles showed very high time correlation ( $R^2 = 0.98$ ) with the POA factor, suggesting a splitting of the same aerosol source. As the additional primary factors could not be associated to specific primary emissions, those solutions are not shown. Medium-long range transport of polluted air masses resulted in a co-variability of the sources at the sampling site, hampering a further separation of the primary organic aerosols.

### 2.2.3. 7-wavelength aethalometer and Scanning Mobility Particle Sizer

An aethalometer, Model AE31 Spectrum (Manufactured by Aerosol d.o.o., Ljubljana, Slovenia) provided continuous measurements of the black carbon (BC) mass concentrations. The aethalometer was equipped with a PM<sub>2.5</sub> impactor. The aethalometer data were recorded with a 5-minute time resolution. The optical transmission of light absorbing carbonaceous aerosol particles was measured at seven wavelengths (370, 450, 520, 590, 660, 880, and 950 nm). Measurements at 880 nm wavelength were used to determine BC mass concentration (Lavanchy et al., 1999). The aethalometer converts light attenuation measurements to BC mass using a specific attenuation absorption cross-section ( $\sigma$ ) of  $16.6 \text{ m}^2 \text{ g}^{-1}$  (at 880 nm) (Aethalometer Operations manual). The default value for the near-infrared wavelength of 880 nm was set by the manufacturer. An empirical algorithm for loading effects compensation was used (Collaud Coen et al., 2010). Ångström exponent of the absorption coefficient computed by fitting an exponential curve was evaluated.

Aerosol size distribution measurements were performed using a Scanning Mobility Particle Sizer (SMPS) model 19.3.09 IFT/TT (TROPOS, Leipzig, Germany), with automatic sheath flow, temperature and relative humidity (RH) control (SMPS setup V2.6 TT 2006) as described in Wiedensohler et al. (2012) applying a CPC UF-02M (Mordas et al., 2013). The SMPS measured particle size (8.7 to 840.0 nm) with a time resolution of 5 min having 72 channels.

Deleted: a

Deleted: ð

#### 2.2.4. OC/EC, $^{14}\text{C}$ , $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analysis

Filter measurements were performed to determine OC, EC and [total carbon \(TC\)](#) concentrations with a thermo-optical OC/EC analyser (Sunset Laboratory Inc, USA) equipped with a non-dispersive infrared (NDIR) detector. A 1.5 cm<sup>2</sup> filter punch was analysed according to the EUSAAR2 protocol (Cavalli et al., 2010). The blank filter was subtracted only from the measured OC and TC concentrations, as for the EC the corresponding blank was below the detection limit of the instrument.

$^{14}\text{C}$  in EC and TC was measured using the accelerator mass spectrometer MICADAS, equipped with a gas-capable ion source (Szidat et al., 2014).  $^{14}\text{C}$  analysis of TC was determined after combustion of filter punches in an elemental analyser, directly coupled to the MICADAS (Salazar et al., 2015). The TC  $^{14}\text{C}$  raw data were corrected for a representative field blank. For  $^{14}\text{C}$  analysis of EC, the filters were first water extracted in order to minimize charring by removing the water-soluble OC (WSOC). Then the Swiss\_4S protocol (Zhang et al., 2012) was used to remove the water-insoluble OC (WINSOC) and measure the EC  $^{14}\text{C}$ , by coupling of the Sunset instrument to the MICADAS (Agrios et al., 2015).  $^{14}\text{C}$  in OC was determined from the TC  $^{14}\text{C}$  and the EC  $^{14}\text{C}$  results with an isotope mass balance calculation. All the data from the  $^{14}\text{C}$  analysis were corrected for the decay of the  $^{14}\text{C}$  from 1950 until present. The reported uncertainty for the non-fossil fraction of EC includes both charring of OC (overestimation of EC) and EC loss (underestimation of EC) during the WINSOC removal process (Zhang et al., 2012). Non-fossil fractions of TC, EC and OC (i.e., TC<sub>nf</sub>, EC<sub>nf</sub> and OC<sub>nf</sub>) were determined from the individual  $^{14}\text{C}$  analyses and  $^{14}\text{C}$  reference values. These reference values represent emissions from purely non-fossil sources and amount to 1.06 ± 0.03 for TC and OC and 1.10 ± 0.03 for EC based on the calculation of Mohr et al. (2009). The fossil fractions of TC, EC and OC (i.e., TC<sub>f</sub>, EC<sub>f</sub> and OC<sub>f</sub>) were determined by subtraction of the respective non-fossil fractions.

Bulk  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values were derived by measuring filter pieces (1.4 cm<sup>2</sup>) wrapped in tin capsules (8\*5 mm, Elemental Microanalysis) using an elemental analyser accompanying an isotope ratio mass spectrometer (EA-IRMS, Flash EA1112—Thermo V Advantage) via a ConFlo III interface. The autosampler of the EA was continuously flushed with He (180 mL min<sup>-1</sup>) to remove all atmospheric gases. Helium flow on the oxidation column was 80 mL min<sup>-1</sup>. Flash combustion occurred in the oxidation column with the presence of O<sub>2</sub> (the O<sub>2</sub> flow was 180 mL min<sup>-1</sup> for 4 s). Formed gases were taken to the reduction column in which molecular nitrogen was obtained from any nitrogen oxides followed by a water trap

Deleted: s

Deleted: s

Deleted: l

Deleted: l

Deleted: l

(magnesium perchlorate). The nitrogen and the carbon dioxide were separated on a packed gas chromatographic (GC) column (PoraPlot, 3m\*2cm, 35 °C) and delivered to the isotope ratio mass spectrometer (via the ConFlo interface) where the measurement of carbon and nitrogen isotope ratio was made. The amount of nitrogen and carbon in the sample was determined by a thermal conductivity detector which is a part of the elemental analyser. These measurements were used in the isotope mass balance calculations (Eq. 1).

The total carbon and total nitrogen fractions of the aerosol particles were used for the isotopic ratio measurements. Stable carbon and nitrogen isotopic ratio measurements were expressed relative to the Vienna Pee Dee Belemnite (VPDB) standard using the formula:

$$\delta^{13}\text{C} = \left( \frac{R_{\text{sample}}}{R_{\text{standard}}} - 1 \right) * 1000 (\text{‰}), \quad (1)$$

where  $R_{\text{sample}}$  and  $R_{\text{standard}}$  are the ratios of  $^{13}\text{C}$  to  $^{12}\text{C}$  (or  $^{15}\text{N}$  to  $^{14}\text{N}$ ) in the sample and the standard (referred to as VPDB), respectively.

Repeated analysis of certified reference material (caffeine IAEA-600) and oil (NBS 22) gave an average  $\delta^{13}\text{C}$  value: mean  $\pm \sigma = -27.77 \pm 0.08 \text{ ‰}$  (certified value: mean  $\pm \sigma = -27.771 \pm 0.043 \text{ ‰}_{\text{VPDB}}$ ) and  $-30.03 \pm 0.09 \text{ ‰}$  (certified value: mean  $\pm \sigma = -30.031 \pm 0.043 \text{ ‰}_{\text{VPDB}}$ ) respectively. These values were used for  $\delta^{13}\text{C}$  measurements in order to evaluate an analytical precision and calibration of a reference gas ( $\text{CO}_2$ ) to VPDB. Meanwhile, the IAEA-600 standard gave an average  $\delta^{15}\text{N}$  value: mean  $\pm \sigma = 1.0 \pm 0.2 \text{ ‰}$  which was used for calibration of a reference gas ( $\text{N}_2$ ) to air (for  $\delta^{15}\text{N}$  measurements).

Stable carbon and nitrogen isotope ratios were measured in the samples with the signal intensity reaching 1000 mV or more, due to analytical restrictions (the isotope values measurements below 1000 mV did not fulfil linearity requirements of 0.07 ‰/V for the internal standard).

The mass balance equation was used to calculate the real  $\delta$  values of carbon or nitrogen of the aerosol samples (blank correction):

$$m_{\text{measured}} \times \delta X_{\text{measured}} = m_1 \times m_{\text{blank}} \times \delta X_{\text{blank}}, \quad (2)$$

where  $m_{\text{measured}}$  was the mass of measured material (carbon or nitrogen) in the measured sample,  $\delta X_{\text{measured}}$  was the measured (aerosol + filter)  $\delta$  value (carbon or nitrogen),  $m_1$  was the mass of pure material (carbon or nitrogen), and  $m_{\text{blank}}$  and  $\delta X_{\text{blank}}$  were the mass and isotope ratio (of carbon or nitrogen) of the blank filter, respectively.

**Deleted:** Repeated analysis of certified reference material (caffeine IAEA-600) and oil (NBS 22) gave an average  $\delta^{13}\text{C}$  value: mean  $\pm \sigma = -27.77 \pm 0.08 \text{ ‰}$  (certified value: mean  $\pm \sigma = -27.771 \pm 0.043 \text{ ‰}_{\text{VPDB}}$ ) and  $\delta^{13}\text{C} = -30.031 \pm 0.043 \text{ ‰}_{\text{VPDB}}$  respectively.



### 2.2.5. Radiocarbon-based source apportionment of carbonaceous aerosols

An estimate of fossil and non-fossil primary and secondary organic carbon ( $\text{POC}_f$ ,  $\text{POC}_{nf}$ ,  $\text{SOC}_f$ ,  $\text{SOC}_{nf}$ ) was achieved by coupling ACSM-PMF results,  $^{14}\text{C}$  data, and organic marker measurements using a chemical mass balance-like approach. The sensitivity of  $\text{POC}_f$ ,  $\text{POC}_{nf}$ ,  $\text{SOC}_f$ , and  $\text{SOC}_{nf}$  contributions to the assumed parameters and measurement errors are described in details in this section. The approach is based on the  $\text{POC}_{nf}$  estimate, for a subsequent determination of  $\text{SOC}_{nf}$ ,  $\text{SOC}_f$ , and  $\text{POC}_f$  as follows:

$$\text{SOC}_{nf} = \text{OC}_{nf} - \text{POC}_{nf} \quad (3)$$

$$\text{SOC}_f = \text{SOC} - \text{SOC}_{nf} \quad (4)$$

$$\text{POC}_f = \text{OC}_f - \text{SOC}_f \quad (5)$$

$^{14}\text{C}$  measurements and ACSM-PMF results were coupled as follows. Daily  $\text{OC}_{nf}$  measurements from radiocarbon analysis as well as average daily POA from ACSM-PMF results provided two upper boundaries for the daily  $\text{POC}_{nf}$  contribution. In this manner we identified a possible daily range of  $\text{POC}_{nf}$  contributions. In order to determine more precisely the  $\text{POC}_{nf}$  daily contributions within the aforementioned possible daily ranges, we performed a sensitivity analysis. Briefly, in the sensitivity analysis we considered a uniform distribution of possible  $\text{POC}_{nf}$  contributions within the identified possible daily ranges, meaning that each  $\text{POC}_{nf}$  value in the selected ranges was considered as equally probable (however, as discussed in the next section, in order to explore the influence of this assumption we also performed the same sensitivity analysis assuming a non-uniform distribution). Assuming no  $\text{POC}_{nf}$  contribution from other sources than [biomass burning organic carbon \(BBOC\)](#), each  $\text{POC}_f$  contribution in the acceptable daily ranges could be written either as  $[\text{BBOC}] = [\text{levoglucosan}]/\alpha$  or as  $[\text{BBOC}] = [\text{EC}_{nf}]/\beta$ , where  $\alpha$  represents the levoglucosan/BBOC ratio and  $\beta$  represents the  $\text{EC}_{nf}$ /BBOC ratio. In two separated sensitivity analyses we scanned broad  $\alpha$  and  $\beta$  ranges covering the possible  $\text{POC}_{nf}$  daily ranges and we retained only  $\text{POC}_{nf}$ ,  $[\text{levoglucosan}]/\alpha$ , and  $[\text{EC}_{nf}]/\beta$  combinations associated to selected acceptance criteria described in the following. From the acceptable solutions we then derived the daily probability distribution function of  $\text{POC}_f$ ,  $\text{SOC}_{nf}$ ,  $\text{SOC}_f$ ,  $\text{POC}_f$ ,  $\alpha$ , and  $\beta$ .

The assumption that each input  $\text{POC}_{nf}$  contribution in the selected possible range is equally probable (hereafter referred to as “uniform distribution approach”) has advantages and

drawbacks: while this assumption doesn't consider any a priori information about levoglucosan/POC<sub>nf</sub> and EC<sub>nf</sub>/POC<sub>nf</sub>, it considers those ratios as equally possible. To explore the influence of this assumption on our results we performed the same sensitivity analysis assuming an input levoglucosan/POC<sub>nf</sub> distribution derived from 33 profiles for combustion of hard or softwoods in domestic fireplaces or woodstoves (Fine et al. 2001, 2002, 2004a, 2004b, Schmidl et al. 2008, the approach is hereafter referred to as "non-uniform distribution approach"). We eventually derived the probability distribution functions of the levoglucosan/POC<sub>nf</sub> and EC<sub>nf</sub>/POC<sub>nf</sub> ratios relative to the acceptable solutions. The two approaches provided similar results. From the uniform distribution approach, a median levoglucosan/POC<sub>nf</sub> ratio of 0.18 (1<sup>st</sup> quartile = 0.14; 3<sup>rd</sup> quartile = 0.23) and a median EC<sub>nf</sub>/POC<sub>nf</sub> ratio of 0.32 (1<sup>st</sup> quartile = 0.28; 3<sup>rd</sup> quartile = 0.36) were retrieved, whilst from the non-uniform distribution approach a median levoglucosan/POC<sub>nf</sub> ratio of 0.15 (1<sup>st</sup> quartile = 0.13; 3<sup>rd</sup> quartile = 0.18) and a median EC<sub>nf</sub>/POC<sub>nf</sub> ratio of 0.33 (1<sup>st</sup> quartile = 0.28; 3<sup>rd</sup> quartile = 0.36) were obtained.

In the following section a technical description of the sensitivity analysis implementation is reported. For each filter sample  $i$ , 10000 random combinations ( $r$ ) of input data,  $[TC]_{i,r}$ ,  $[EC]_{i,r}$ ,  $[EC_f]_{i,r}$ ,  $[OC_f]_{i,r}$ , and  $[Levoglucosan]_{i,r}$ , were generated. In this process, we assume a normal distribution of the errors around the average  $[X]_i$  value ( $X$  being one of the input values mentioned above), and a distribution width equal to the standard deviation  $\sigma[X]_i$ :

For each random combination of input data, the corresponding  $[OC]_{i,r}$ ,  $[EC_{nf}]_{i,r}$ , and  $[OC_{nf}]_{i,r}$  values were determined as:

$$[OC]_{i,r} = [TC]_{i,r} - [EC]_{i,r}, \quad (6)$$

$$[EC_{nf}]_{i,r} = [EC]_{i,r} - [EC_f]_{i,r}, \quad (7)$$

$$[OC_{nf}]_{i,r} = [OC]_{i,r} - [OC_f]_{i,r}. \quad (8)$$

10000 random  $[SOC]_s$  values were generated by randomly selecting a daily average  $[SOA]_s$  value from one of the 20 ACSM-PMF runs ( $s$ ). The corresponding  $[SOC]_s$  values were derived as:

$$[SOC]_s = [SOA]_s / (OM/OC)_{SOA(s)} \quad (9)$$

$(OM/OC)_{SOA(s)}$  and  $\sigma(OM/OC)_{SOA(s)}$  were calculated according to Aiken et al. (2009) as function of the fractional contribution of the  $m/z$  44 ( $f_{44}$ ) to the  $SOA_s$  mass spectra. Fröhlich et al. (2015) showed a systematic difference between  $f_{44}$  measured from ACSM and AMS.

Deleted: ,

Deleted: ,

1 therefore an empirical correction factor was accordingly applied to rescale  $f44$  from ACSM  
 2 ( $f44_{ACSM}$ ) data to the corresponding AMS  $f44$  value ( $f44_{AMS}$ ). The uncertainty relative to the  
 3  $f44$  correction factor was propagated into  $\sigma(OM/OC)_{SOA(s)}$  which includes the  $O/C_s$   
 4 uncertainty as well. Each  $[SOC]_{i,r}$  value was obtained by randomly varying  $[SOC]_s$  assuming  
 5 a normal distribution of errors around the average value  $[SOC]_s$  and a distribution width equal  
 6 to  $\sigma(OM/OC)_{SOA(s)}$ .  $[BBOC]_{i,r}$  contributions for each sample  $i$  were derived as follows:

Deleted: s

$$7 \quad [BBOC]_{i,r} = [\text{levoglucosan}]_{i,r}/\alpha, \quad (10)$$

$$8 \quad [BBOC]_{i,r} = [EC_{nf}]_{i,r}/\beta, \quad (11)$$

9 where  $\alpha$  represents the levoglucosan/BBOC ratio. This ratio was systematically varied  
 10 between 0.01 and 0.31 according to Huang et al. (2014) and references therein (scan step  
 11 equals 0.01).  $\beta$  corresponds to the EC/BBOC ratio. Values of  $\beta$  were systematically varied  
 12 between 0.1 and 0.4 according to Zhang et al. (2015) and references therein (scan step equal  
 13 to 0.01). 10000  $[BBOC]_{i,r,\alpha}$  and 10000  $[BBOC]_{i,r,\beta}$  were determined as in Eq. (8) and (9). Only  
 14 acceptable  $[BBOC]_{i,r,\alpha/\beta}$  ( $= [POC_{nf}]_{i,r,\alpha/\beta}$ ) values were considered for the sensitivity analysis.

Deleted: s

15 The criteria to consider a  $[BBOC]_{i,r,\alpha/\beta}$  value as acceptable were:

$$16 \quad \text{a) } [BBOC]_{i,r,\alpha/\beta} \leq [POC]_{i,r} \text{ and b) } [BBOC]_{i,r,\alpha/\beta} \leq [OC_{nf}]_{i,r} \quad (12)$$

17  $[POC]_{i,r}$  was determined as follows:

$$18 \quad [POC]_{i,r} = [OC]_{i,r} - [SOC]_{i,r}, \quad (13)$$

19 Only acceptable  $[POC]_{i,r}$  values were considered. The criterion to consider a  $[POC]_{i,r}$  value as  
 20 acceptable was:

$$21 \quad \text{c) } [POA]_s/[POC]_{i,r} \geq 1.3 \text{ according to Mohr et al. (2009), Aiken et al. (2009).}$$

22  $[SOC_{nf}]_{i,r}$  values were then derived as:

$$23 \quad [SOC_{nf}]_{i,r} = [OC_{nf}]_{i,r} - [POC_{nf}]_{i,r} \quad (14)$$

24 Only acceptable  $[SOC_{nf}]_{i,r}$  values were considered, where

$$25 \quad \text{d) } [SOC_{nf}]_{i,r} \leq [SOC]_{i,r} \quad (15)$$

26 Only solutions where all 4 criteria a), b), c), and d) held were considered acceptable and  
 27 retained.

28 Finally,  $[SOC_f]_{i,r}$  and  $[POC_f]_{i,r}$  were calculated as:

$$29 \quad [SOC_f]_{i,r} = [SOC]_{i,r} - [SOC_{nf}]_{i,r}, \quad (16)$$

$$[\text{POC}]_{i,t} = [\text{OC}]_{i,t} - [\text{SOC}]_{i,t}. \quad (17)$$

## 2.2.6. Organic markers and satellite products

Determination of organic marker concentrations was performed using a developed in-situ derivatization thermal desorption gas chromatography time of flight mass spectrometry (IDTD-GC-MS) method (Orasche et al., 2011).

Biomass burning episodes were explored using a variety of remote sensing datasets and their derived properties. Satellite data and ground based observations of aerosol properties from the MODIS, HYSPLIT and SILAM (Sofiev et al., 2006) were coupled to analyse the variability of carbonaceous aerosols in Lithuania (Fig. 2).

The MODIS sensors on-board NASA's Terra and Aqua satellites provide multiple thermal observations of the Earth on 9–10 March 2014 at a spatial resolution of 1 km using the latest version of the MODIS Active Fire Product (MOD14/MYD14) algorithm (MODIS, 2011). To identify the influence of air masses from different transport pathways on the large biomass burning (BB) event occurring at Preila, 72-h back trajectories at an arrival height of 100, 200 and 500 m were calculated by the Hybrid Single Particle Lagrangian Integrated Trajectory (HYSPLIT) Model Version 4.8 (Stein et al., 2015). All air mass back trajectories were generated using Gridded Meteorological Data archives of the Air Resource Laboratory (ARL), National Ocean and Atmospheric Administration (NOAA) (Fig. 2A).

The Navy Aerosol Analysis and Prediction System (NAAPS) model results were used to define the distribution of BB aerosols from wildfire areas (model description and results are available from the web pages of the Naval Research Laboratory, Monterey, CA, USA; <http://www.nrlmry.navy.mil/aerosol/>) (Fig. 2B). The NAAPS model has been adapted to combine real-time observations of biomass burning based on the joint Navy/NASA/NOAA Fire Locating and Modelling of Burning Emissions system (FLAMBE, <http://www.nrlmry.navy.mil/flambe/>) (Reid et al., 2004). The method has proven helpful in previous studies of long-range and regional transport of smoke (Honrath et al., 2004). The resolution of 2.5° longitude × 2.5° latitude National Centers for Environmental Prediction (NCEP) reanalysis data (Kanamitsu et al., 2002) during the grass burning episode were analysed to illustrate the sub synoptic-scale weather feature among the biomass burning events over Lithuania issued every 6 h for March 2014 (Fig. 2C). SILAM is an air quality and emergency open code system (<http://silam.fmi.fi/>) providing PM<sub>2.5</sub> emission maps by Eulerian dynamics and a combination of basic acid and ozone chemistry with inert particles for fire

Deleted: S

Deleted: and organic markers

Deleted: ere

Deleted: s

Deleted: s

and anthropogenic primary PM emission to account for the fire induced aerosol contribution (Fig. 2D).

## Results and discussion

### 2.1 Identification of grass burning event

Massive active fires occurred throughout the Kaliningrad region (Russia), Belorussia and Ukraine (Fig. 2A) when a high atmospheric pressure system was situated over the study area, as illustrated in the weather map of Fig. 2C. The plumes from those fires covered a large area south of the Baltic region and were transported thousands of kilometres downwind affecting the background air in Lithuania (Fig. 2). Although the number of fires was similar to previous years, the impact of the fire events on the Lithuanian air quality was enhanced in March 2014 due to air mass transport of smoke entrained in deep convection by the large scale circulation around the pressure maximum of the anticyclonic system (Fig. 2C). This is consistent with the relatively high concentrations of smoke reaching Preila as predicted by NAAPS (Fig. 2B).

The weather maps showed that the high concentration of pollutants during this BB event was caused by the anticyclonic large-scale movement, which persisted throughout the lower troposphere causing stagnant conditions and extended aerosol residence time.

### 2.2 Investigation of PM<sub>1</sub> composition and ambient concentrations of organic tracers

The climatic conditions in West Europe as well as in west part of Lithuania are moderate warm climate dominating by air mass transport from Atlantic Ocean, leading to higher humidity. Annual mean temperature increases in west-east direction. The average temperature of March was ~3–4 °C. During the BB event (9–11 March) combustion products were spread over the study region by the large-scale atmospheric circulation processes. At the beginning of the BB episode, the wind speed was up to 3 m s<sup>-1</sup> on average in the daytime of 9<sup>th</sup> March, causing weaker dilution of the pollutants while the BC concentration was higher than 12 µg m<sup>-3</sup>.

During the campaign, on average, organic aerosol (46%, 3.2 µg m<sup>-3</sup> (σ = 4.8 µg m<sup>-3</sup>)) constituted the major fraction of the NR-PM<sub>1</sub> aerosol concentration composition measured by ACSM with lower contributions of sulfate (17%, 1.2 µg m<sup>-3</sup> (σ = 1.1 µg m<sup>-3</sup>)), nitrate (20%),

Deleted: that one of

Deleted: particular, as the

Deleted: ere

Deleted:

Deleted:

Deleted:

1 1.4  $\mu\text{g m}^{-3}$  ( $\sigma = 1.8 \mu\text{g m}^{-3}$ ), ammonium (15%, 1.0  $\mu\text{g m}^{-3}$  ( $\sigma = 0.9 \mu\text{g m}^{-3}$ )), and chloride  
2 (2%, 0.1  $\mu\text{g m}^{-3}$  ( $\sigma = 0.3 \mu\text{g m}^{-3}$ )). The average composition of NR-PM<sub>1</sub> showed similar  
3 dominance of organics to previous observations in Europe (e.g., Crippa et al., 2014). OA  
4 contribution to NR-PM<sub>1</sub> was found to be much higher during the grass burning period (61%,  
5 8.6  $\mu\text{g m}^{-3}$  ( $\sigma = 5.0 \mu\text{g m}^{-3}$ ), followed by sulfate (5%, 1.4  $\mu\text{g m}^{-3}$  ( $\sigma = 0.5 \mu\text{g m}^{-3}$ ), nitrate  
6 (19%, 3.0  $\mu\text{g m}^{-3}$  ( $\sigma = 1.4 \mu\text{g m}^{-3}$ ), ammonium (13%, 1.6  $\mu\text{g m}^{-3}$  ( $\sigma = 0.7 \mu\text{g m}^{-3}$ ), and  
7 chloride (3%, 0.4  $\mu\text{g m}^{-3}$  ( $\sigma = 0.3 \mu\text{g m}^{-3}$ )) (Fig. 5A).

8 The concentrations of the monosaccharide anhydrides together with those of OC and EC are  
9 presented in Fig. 3. It is evident that during the event, when grass burning was most intense,  
10 the levoglucosan concentration increased up to 680 ng m<sup>-3</sup>. That is substantially lower than  
11 values reported during the extreme event of August 2010 in Moscow – 3100 ng m<sup>-3</sup>  
12 (Popovicheva et al., 2014) and is higher than values (220–290 ng m<sup>-3</sup>) reported during a  
13 major biomass burning episode over northern Europe in Helsinki (Saarikoski et al., 2007),  
14 while background values in Nordic rural background sites were found to be 2.1–9.8 ng m<sup>-3</sup>  
15 (Yttri et al., 2011). Concentrations of mannosan varied from 3.1 to 68.0 ng m<sup>-3</sup> and those of  
16 galactosan from 1.0 to 12.0 ng m<sup>-3</sup>. The levoglucosan to mannosan (L/M), levoglucosan to  
17 galactosan (L/G) and levoglucosan to OC (L/OC) ratios were used before to separate different  
18 BB sources (Fabbri et al., 2009; Oanh et al., 2011; Harrison et al., 2012). We measured  
19 average L/M and L/G ratios of 16.4 and 135.8, respectively. This is similar to the values  
20 found by Orasche et al. (2012) from wood combustion in residential wood appliances and in  
21 the range of L/M ratios reported (2.0–33.3) for grass fires by Oros et al. (2006). Excluding the  
22 strong event days of March 9 and 10 the sugars showed a good correlation with each other ( $R^2$   
23 > 0.86). On March 9 and 10 the mannosan/galactosan was lower at 2–6, indicating a different  
24 source than on the other days. Low mannosan/galactosan ratios were observed for grass and  
25 leaves (Sullivan et al., 2014). We observed an L to OC ratio from 0.06 to 0.16 during the  
26 biomass burning period and of ~ 0.03 during the days without biomass burning events. The  
27 values observed during biomass burning are in the range of those (0.04–0.20) reported for  
28 wildland fuels (Sullivan et al., 2008). The OC/EC ratio ranged from 1.5 to 6.2 being lower on  
29 event days (2.4–3.0) indicating an aerosol composition dominated by organic aerosol. During  
30 the intensive grass burning episode, consecutive new particle formation (NPF) events were  
31 observed. Observed NPF could be attributed to the grass burning and secondary biomass  
32 burning product transformation as was evaluated in earlier studies over the same area

Deleted:

Deleted:

Deleted:

Deleted: (

Deleted: )

Deleted:

Deleted: )

Deleted:

Deleted:

Deleted: )

Deleted:

Deleted: )

Deleted:

Deleted: Quantification of

Deleted: was significantly

Deleted: are

Deleted: Concentrations of mannosan varied from 3.1 to 68.0 ng m<sup>-3</sup>, with a mean value of 19.0 ng m<sup>-3</sup>, and concentrations of galactosan varied from 1.0 to 12.0 ng m<sup>-3</sup>.

Deleted: contribution

Deleted: to

Deleted: ,

Deleted: May

Deleted: episodes

1 (Ulevicius et al., 2010b). At 13:00, **there was significant new** particle formation on 9<sup>th</sup> and  
 2 10<sup>th</sup> March followed by subsequent growth up to three hours. The total particle number  
 3 concentration with a daily mean value of 6440 cm<sup>-3</sup> (with maximum value of 13000 cm<sup>-3</sup>)  
 4 **was observed**, which was extremely much higher than the daily mean observations in non-  
 5 event days (1660 cm<sup>-3</sup>). In this area **an annual mean** total particle concentration of 2650 cm<sup>-3</sup>  
 6 was observed (Byčėnkiėnė et al., 2013). Non-event days **were** characterized by bimodal  
 7 (Aitken (geometric mean diameter ( $D_g$ ) of 44 nm) and accumulation ( $D_g$  = 128 nm))  
 8 **distributions** with a standard deviation of 1.68 and 1.87, respectively. In comparison, during  
 9 **the** biomass burning event trimodal (nucleation ( $D_g$  = 9.0 nm), Aitken ( $D_g$  = 31.0 nm) and  
 10 accumulation ( $D_g$  = 102 nm)) **distributions** with a standard deviation of 1.77, 1.71 and 1.68,  
 11 respectively. However, **the** volume distribution was characterized by **a** bimodal size  
 12 distribution for **the** non-event days ( $D_g$  = 330 and 665 nm) and **for the** event day ( $D_g$  = 250  
 13 and 590 nm).  
 14 The measured  $\delta^{13}\text{C}$  values varied from -28.2 to -26.7 ‰. The lowest stable carbon isotope  
 15 ratio values (-28.5 ‰) were detected during the period with the highest total carbon  
 16 concentration of 12.2  $\mu\text{g m}^{-3}$  (2014.03.10) and 8.5  $\mu\text{g m}^{-3}$  (2014.03.09). The highest  
 17 concentration 14.0  $\mu\text{g m}^{-3}$  of nitrogen was detected on 10 March 2014. The nitrogen isotope  
 18 ratio values varied from +1.0 to +13.0 ‰ (Fig. 4).  
 19 Stable carbon and nitrogen isotope ratios values of aerosol particles derived from biomass  
 20 burning (C3 plants) and liquid fossil fuel are overlapping (Garbaras et al., 2015, Masalaite et  
 21 al., 2015, Turekian et al., 1998). Coal derived aerosol particles are characterised by higher  
 22  $\delta^{13}\text{C}$  and lower  $\delta^{15}\text{N}$  values (Fig. 4, solid lines).  $\delta^{13}\text{C}$  values of aerosol particles during wild  
 23 grass burning events distinguish in low  $\delta^{13}\text{C}$  values (Garbaras et al., 2008; Ulevicius et. al.,  
 24 2010b). The above mentioned distribution of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values allowed excluding coal  
 25 burning as main source for aerosol particles at Preila during the investigated event. Aerosol  
 26 particles with  $\delta^{13}\text{C}$  values equal **to** -28 ‰ and below originated mainly from grass burning  
 27 events. This interpretation of the data is consistent with the radiocarbon analysis shown  
 28 below.

Deleted: a

Formatted: Highlight

Deleted: as

Deleted: modes

Deleted: modes

Deleted: ,

Deleted: the

## 29 2.3 Source apportionment of EC and OC using $^{14}\text{C}$ data

30 Relative fossil and non-fossil contributions to OC and EC were evaluated using  $^{14}\text{C}$   
 31 analysis (Szidat et al., 2014) to enable a more detailed source attribution of the carbonaceous  
 32 aerosol mass. Widely used, two-source simple models (Currie, 2000; Lemire et al., 2002;

1 Lewis et al., 2004; Szidat et al., 2004) can only distinguish fossil from non-fossil TC  
 2 emissions. Here, carbonaceous aerosol was described to be composed of the following 4  
 3 categories: OC<sub>f</sub> and EC<sub>f</sub> attributed to primary and secondary fossil fuel combustion; and OC<sub>nf</sub>,  
 4 and EC<sub>nf</sub> typically [attributed](#) to primary and secondary biomass burning, cooking, biogenic  
 5 emissions and non-fossil OC combustion (Table 2, Fig. 5). There was day-to-day variation in  
 6 the fractional contributions to TC throughout the BB event. The fraction of elemental carbon  
 7 from biomass burning EC<sub>bb</sub> (= EC<sub>nf</sub>) to total EC was found to be on average 67±3%. For EC<sub>bb</sub>  
 8 the mean relative contribution to total carbon in background areas of Northern countries was  
 9 found to be <1.5% on non-event days (Yttri et al., 2011). It was also reported that a major  
 10 peak in EC<sub>bb</sub> values between March and April was observed at the Zeppelin atmospheric  
 11 observatory (Yttri et al., 2014). [Observed](#) high values are unusual and have only been found  
 12 in wood burning dominated places like villages in Alpine valleys (Zotter et al., 2014). This  
 13 shows, together with [the](#) high levels of levoglucosan, that biomass burning contributed to a  
 14 large extent to OC<sub>nf</sub> during this event. A mean light absorption coefficient  $\alpha_{370-950}$  (the  
 15 absorption exponent calculated using the seven [wavelengths](#) Aethalometer) of 1.38 ( $\sigma = 0.11$ )  
 16 was obtained during wildfires, which is higher than the mean  $\alpha_{370-950}$  calculated for the non-  
 17 event days (1.13,  $\sigma = 0.19$ ). The light absorption exponent values were calculated with  $\lambda =$   
 18 370 – 520 nm and  $\lambda = 590 - 950$  nm wavelengths for comparison purpose. The mean values of  
 19  $\alpha_{370-520}$  and  $\alpha_{590-950}$  were found to be 1.53 ( $\sigma = 0.19$ ) and 1.32 ( $\sigma = 0.09$ ) during event days  
 20 and 1.25 ( $\sigma = 0.27$ ) and 1.13 ( $\sigma = 0.18$ ) [for the](#) non-event days, respectively. In comparison,  
 21 during [a similar](#) event in Preila higher mean values of  $\alpha_{370-520}$  and  $\alpha_{590-950}$  nm were observed  
 22 (2.4 ( $\sigma = 0.1$ ) and 1.5 ( $\sigma = 0.1$ ), respectively) in 2008, as well as during the event in 2009 (2.3  
 23 ( $\sigma = 0.1$ ) and 1.6 ( $\sigma = 0.1$ ), respectively). (Ulevicius et al., 2010a). This is an indication of [the](#)  
 24 influence of the biomass burning on the Ångström exponent of the absorption coefficient  $\alpha$ .  
 25 The impact of organic aerosols on the spectral dependence of light absorption was [already](#)  
 26 confirmed by [the](#) OC/EC ratios. PMF analysis of OA spectra resolved two OA components,  
 27 which are attributed to POA and SOA, whose mass spectra and time series are presented in  
 28 Fig. 5 B, C. Combining these results with the <sup>14</sup>C measurements as described in section 2.2.4  
 29 shows that the high grass burning pollution event is characterized by a high non-fossil organic  
 30 compound fraction, which accounts for up to ~90% of total carbon mass.  
 31 SOA showed reasonable correlation ( $R^2 = 0.62$ ) with average NH<sub>4</sub><sup>+</sup> mass concentration during  
 32 the BB event. NH<sub>4</sub><sup>+</sup> is in this case a good tracer for secondary aerosol, as it correlates well

Deleted: emitted

Deleted: by

Deleted:

Deleted:

Deleted: y

Deleted: Such

Deleted: wavelengths

Deleted: during

Deleted: the same

Deleted: (

Deleted: )

Deleted: )

Deleted: early

Deleted:



1 with the sum of  $\text{NO}_3^-$  and  $\text{SO}_4^{2-}$  ( $R^2 = 0.96$ ) (Fig. 6). There was day-to-day variation  
 2 throughout the study period with the non-fossil contribution to organic carbon between 67–  
 3 86%.  $\text{OC}_{\text{nf}}$  was estimated to be ~65% primary, while the primary fraction of the  $\text{OC}_{\text{f}}$  in Preila  
 4 was estimated to be ~93%. Conversely, when  $\text{EC}_{\text{f}}$  showed a lower contribution (2014.03.07  
 5 and 2014.03.10; 26% and 24%, respectively),  $\text{OC}_{\text{f}}$  was also lower (15%) (Table 3). The high  
 6 fraction of biomass burning was corroborated by measurements of levoglucosan. Other  
 7 molecular markers such as hopanes for traffic emissions and picene for coal combustion  
 8 (Rutter et al., 2009) were also measured in order to monitor the possible contribution of fossil  
 9 fuel combustion during the high pollution event. Although their concentrations increased  
 10 during the episode, suggesting a contribution of co-transported fossil fuel combustion  
 11 aerosols, the radiocarbon analysis revealed the contribution of this fraction to be minor ( $\text{EC}_{\text{f}}$   
 12 ranged from 0.3 to 1.1;  $\text{OC}_{\text{f}}$  ranged from 0.5 to 1.6  $\mu\text{gm}^{-3}$  (Fig. 5)). The concentrations of the  
 13 molecular markers are provided in Table S1 of the Supplementary material. The combination  
 14 of measurements and source apportionment techniques allowed a better characterization of the  
 15 carbonaceous aerosol sources. POA determined with the ACSM is mostly non-fossil and  
 16 originates from grass burning. It is shown that  $\text{POC}_{\text{nf}}$  and  $\text{SOC}_{\text{nf}}$  concentrations increase  
 17 drastically (from 1.1 to 5.4  $\mu\text{gm}^{-3}$  for  $\text{POC}_{\text{nf}}$ ; from 0.9 to 3.1  $\mu\text{gm}^{-3}$  for  $\text{SOC}_{\text{nf}}$ ) with increasing  
 18 influence of biomass burning, whereas the concentrations of the respective fossil fractions  
 19 show a smaller increase during this episode. From the acceptable solutions obtained from the  
 20 sensitivity test described in section 2.2.5, we derived the probability distribution functions of  
 21 the different daily contributions for the  $\text{POC}_{\text{f}}$ ,  $\text{SOC}_{\text{nf}}$ ,  $\text{SOC}_{\text{f}}$ ,  $\text{POC}_{\text{f}}$  fractions (Fig. 7). The  
 22 median tests are consistent and  $\text{EC}/\text{BBOC}$  ratios obtained from the sensitivity tests are  
 23 consistent with values reported in Zhang et al. (2015) and Huang et al. (2014) (Fig. 8).

24 In Zhang et al. (2015) agricultural waste combustion is considered to be the main contributor  
 25 to the total biomass burning. Note that on 5 March a different Levoglucosan/BBOC ratio was  
 26 found (0.31) compared to the non-event days (~0.15). Also, this is consistent with different air  
 27 mass back-trajectories, associated to air masses originating in the Southern and Central  
 28 Russian Federal districts, i.e., air masses with a different geographical origin and associated to  
 29 potentially different types of biomass burning.

### 30 3 Conclusions

31 In March 2014, an intensive field campaign was conducted in the marine background of  
 32 the South Eastern Baltic region during a period of intensive grass burning. This paper

Deleted: ; linear fit:  $y = 0.816x + 0.005$

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted: Values

Deleted: both techniques

Deleted: The lines in Fig. 7 represent the absolute contribution of each source during 5–10 March 2014.

Formatted: Highlight

Deleted: 7

Deleted: Levoglucosan/BBOC

Deleted: is

Deleted: wind

1 provides the biomass burning related aerosol concentrations during grass burning estimated  
2 by data that stem from a synthesis of various techniques including surface online/offline and  
3 satellite based measurements. Lidar vertical profiles allowed confirming smoke plumes from  
4 wild fire regions. Levels of source specific tracers, i.e., levoglucosan as well as  $^{14}\text{C}$  of TC, EC  
5 and OC were used as input for source apportionment of the carbonaceous aerosol. Overall, EC  
6 and OC were dominated by non-fossil sources. The total POC fraction was separated into  
7  $\text{POC}_f$  and  $\text{POC}_{nf}$ . In terms of OC mass,  $\text{POC}_{nf}$  contributes on average 56%, while the relative  
8 contribution to TC was found to be on average 39%. In case of SOC, the contribution of  $\text{OC}_f$   
9 reached on average 10.3% (non-fossil – 25%). The  $\delta^{13}\text{C}$  value of -28.5% indicated the  
10 dominance of the aerosol derived from the vegetation burning as no significant carbon isotope  
11 fractionation occurs between the aerosol particles from biomass burning and the raw biomass  
12 material.

### 13 Acknowledgements

14 This work was supported by the Lithuanian-Swiss Cooperation Programme “Research and  
15 Development” project AEROLIT (Nr. CH-3-ŠMM-01/08).

16

Deleted:

Deleted: ,

Deleted: have been

Deleted: approach

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

## 1   **References**

- 2   Agrios, K., Salazar, G. A., Zhang, Y. L., Uglietti, C., Battaglia, M., Luginbühl, M., Ciobanu,  
3   V. G., Vonwiller, M., and Szidat, S.: Online coupling of pure O<sub>2</sub> thermo-optical methods - <sup>14</sup>C  
4   AMS for source apportionment of carbonaceous aerosols study, Nucl. Instrum. Meth. Phys.  
5   Res. B., 361, 288-293, doi:10.1016/j.nimb.2015.06.008, 2015.
- 6   Aiken, A. C., Salcedo, D., Cubison, M. J., Huffman, J. A., DeCarlo, P. F., Ulbrich, I. M.,  
7   Docherty, K. S., Sueper, D., Kimmel, J. R., Worsnop, D. R., Trimborn, A., Northway, M.,  
8   Stone, E. A., Schauer, J. J., Volkamer, R. M., Fortner, E., de Foy, B., Wang, J., Laskin, A.,  
9   Shutthanandan, V., Zheng, J., Zhang, R., Gaffney, J., Marley, N. A., Paredes-Miranda, G.,  
10   Arnott, W. P., Molina, L. T., Sosa, G., and Jimenez, J. L.: Mexico City aerosol analysis  
11   during MILAGRO using high resolution aerosol mass spectrometry at the urban supersite  
12   (T0) – Part 1: Fine particle composition and organic source apportionment, Atmos. Chem.  
13   Phys., 9, 6633– 6653, doi:10.5194/acp-9-6633-2009, 2009.
- 14   Akagi, S. K., Craven, J. S., Taylor, J. W., McMeeking, G. R., Yokelson, R. J., Burling, I. R.,  
15   Urbanski, S. P., Wold, C. E., Seinfeld, J. H., Coe, H., Alvarado, M. J., and Weise, D. R.:  
16   Evolution of trace gases and particles emitted by a chaparral fire in California, Atmos. Chem.  
17   Phys., 12, 1397–1421, doi:10.5194/acp-12-1397-2012, 2012.
- 18   Baldini, G., Campadelli, P., and Fradegrada, M.: Biomass burning monitoring by scene  
19   analysis, In Proceedings of Visualization, Imaging, and Image Processing, Calgary, 2002.
- 20   Baron, P. A. and Willeke, K. (Eds.): Aerosol Measurement: Principles, Techniques, and  
21   Applications 2nd Edition, Wiley, New York, 2001.
- 22   Beddows, D. C. S., Dall'Osto, M., Harrison, R. M., Kulmala, M., Asmi, A., Wiedensohler, A.,  
23   Laj, P., Fjaeraa, A. M., Sellegri, K., Birmili, W., Bukowiecki, N., Weingartner, E.,  
24   Baltensperger, U., Zdimal, V., Zikova, N., Putaud, J.-P., Marinoni, A., Tunved, P., Hansson,  
25   H.-C., Fiebig, M., Kivekäs, N., Swietlicki, E., Lihavainen, H., Asmi, E., Ulevicius, V., Aalto,  
26   P. P., Mihalopoulos, N., Kalivitis, N., Kalapov, I., Kiss, G., de Leeuw, G., Henzing, B.,  
27   O'Dowd, C., Jennings, S. G., Flentje, H., Meinhardt, F., Ries, L., Denier van der Gon, H. A.  
28   C., and Visschedijk, A. J. H.: Variations in tropospheric submicron particle size distributions  
29   across the European continent 2008–2009, Atmos. Chem. Phys., 14, 4327-4348,  
30   doi:10.5194/acp-14-4327-2014, 2014.

1 Bougiatioti, A., Stavroulas, I., Kostenidou, E., Zarnpas, P., Theodosi, C., Kouvarakis, G.,  
2 Canonaco, F., Prévôt, A. S. H., Nenes, A., Pandis, S. N., and Mihalopoulos, N.: Processing of  
3 biomass-burning aerosol in the eastern Mediterranean during summertime, *Atmos. Chem.*  
4 *Phys.*, 14, 4793–4807, doi:10.5194/acp-14-4793-2014, 2014.

5 Byčenkienė, S., Ulevicius, V., and Kecorius, S.: Characteristics of black carbon aerosol mass  
6 concentration over the East Baltic region from two-year measurements, *J. Environ. Monit.*,  
7 13(4), 1027–1038, doi: 10.1039/C0EM00480D, 2011.

8 Byčenkienė, S., Ulevicius V., Dudoitis V., and Pauraitė J.: Identification and characterization  
9 of black carbon aerosol sources in the East Baltic region, *Adv. Meteor.*, 2013, Article ID  
10 380614, doi.org/10.1155/2013/380614, 2013.

11 Canonaco, F., Crippa, M., Slowik, J. G., Baltensperger, U., and **Prevot** A. S. H.: SoFi, an  
12 IGOR-based interface for the efficient use of the generalized multilinear engine (ME-2) for  
13 the source apportionment: ME-2 application to aerosol mass spectrometer data, *Atmos. Meas.*  
14 *Tech.*, 6(12), 3649–3661, 2013.

15 Capes, G., Johnson, B., McFiggans, G., Williams, P. I., Haywood, J., and Coe, H.: Aging of  
16 biomass burning aerosols over West Africa: aircraft measurements of chemical composition,  
17 microphysical properties, and emission ratios, *J. Geophys. Res. Atmos.*, 113, D00C15, doi:  
18 10.1029/2008JD009845, 2008.

19 Cavalli, F., Viana, M., Yttri, K. E., Genberg, J., and Putaud, J.-P.: Toward a standardised  
20 thermal-optical protocol for measuring atmospheric organic and elemental carbon: the  
21 EUSAAR protocol, *Atmos. Meas. Tech.*, 3, 79–89, doi:10.5194/amt-3-79-2010, 2010.

22 Ceburnis, D., Garbaras, A., Szidat, S., Rinaldi, M., Fahrni, S., Perron, N., Wacker, L., Leinert,  
23 S., Remeikis, V., Facchini, M. C., Prevot, A. S. H., Jennings, S. G., and O'Dowd, C. D.:  
24 Quantification of the carbonaceous matter origin in submicron marine aerosol particles by  
25 dual carbon isotope analysis, *Atmos. Chem. Phys.*, 11, 8593–8606, doi:10.5194/acp-11-8593-  
26 2011.

27 Collaud Coen, M., Weingartner, E., Apituley, A., Ceburnis, D., Fierz-Schmidhauser, R.,  
28 Flentje, H., Henzing, J. S., Jennings, S. G., Moerman, M., Petzold, A., Schmid, O., and  
29 Baltensperger, U.: Minimizing light absorption measurement artifacts of the Aethalometer:  
30 evaluation of five correction algorithms, *Atmos. Meas. Tech.*, 3, 457–474, doi: 10.5194/amt-  
31 3-457-2010, 2010.

Formatted: Highlight

1 Crippa, M., Canonaco, F., Lanz, V. A., Äijälä, M., Allan, J. D., Carbone, S., Capes, G.,  
2 Ceburnis, D., Dall'Osto, M., Day, D. A., DeCarlo, P. F., Ehn, M., Eriksson, A., Freney, E.,  
3 Hildebrandt Ruiz, L., Hillamo, R., Jimenez, J. L., Junninen, H., Kiendler-Scharr, A.,  
4 Kortelainen, A.-M., Kulmala, M., Laaksonen, A., Mensah, A. A., Mohr, C., Nemitz, E.,  
5 O'Dowd, C., Ovadnevaite, J., Pandis, S. N., Petäjä, T., Poulain, L., Saarikoski, S., Sellegri, K.,  
6 Swietlicki, E., Tiitta, P., Worsnop, D. R., Baltensperger, U., and Prévôt, A. S. H.: Organic  
7 aerosol components derived from 25 AMS data sets across Europe using a consistent ME-2  
8 based source apportionment approach, *Atmos. Chem. Phys.*, 14, 6159-6176, doi:10.5194/acp-  
9 14-6159-2014, 2014.

10 Crutzen, P. J., Heidt, L. E., Krasnec, J. P., Pollock, W. H., and Seiler, W.: Biomass burning as  
11 a source of atmospheric gases CO, H<sub>2</sub>, N<sub>2</sub>O, NO, CH<sub>3</sub>Cl, and COS, *Nature* 282, 253-256,  
12 doi:10.1038/282253a0, 1979.

13 Currie, L. A.: Evolution and multidisciplinary frontiers of <sup>14</sup>C aerosol science, *Radiocarbon*  
14 42, 115-126, 2000.

15 Davison, A. C. and Hinkley, D. V.: *Bootstrap Methods and Their Application*, Cambridge  
16 University Press, Cambridge, UK, 582 pp., 1997.

17 [EMEP \(European Monitoring and Evaluation Programme\), website:](http://www.emep.int/index.html)  
18 <http://www.emep.int/index.html>

19 Fabbri, D., Torri, C., Simoneit, B., Marynowski, L., Rushdi, A., and Fabianska, M.:  
20 Levoglucosan and other cellulose and lignin markers in emissions from burning of Miocene  
21 lignites, *Atmos. Environ.*, 43, 2286-2295, doi:10.1016/j.atmosenv.2009.01.030, 2009.

22 Fine, P. M., Cass, G. R., and Simoneit, B. R. T.: Chemical characterization of fine particle  
23 emissions from fireplace combustion of woods grown in the northeastern United States,  
24 *Environ. Sci. Technol.*, 35, 2665-2675, doi:10.1021/es001466k, 2001.

25 Fine, P. M., Cass, G. R., and Simoneit, B. R. T.: Chemical characterization of fine particle  
26 emissions from the fireplace combustion of woods grown in the southern United States,  
27 *Environ. Sci. Technol.*, 36, 1442-1451, doi: 10.1021/es0108988, 2002.

28 Fine, P. M., Cass, G. R., and Simoneit, B. R. T.: Chemical characterization of fine particle  
29 emissions from the wood stove combustion of prevalent United States tree species, *Environ.*  
30 *Eng. Sci.*, 21, 705-721, doi: 10.1089/ees.2004.21.705, 2004a.

Deleted: .

Deleted: .

1 Fine, P. M., Cass, G. R., [and](#) Simoneit, B. R. T.: Chemical characterization of fine particle  
2 emissions from fireplace combustion of woods grown in the Midwestern and Western United  
3 States, *Environ. Eng. Sci.*, 21, 387–409, doi:10.1089/109287504323067021, 2004b.

4 Fröhlich, R., Crenn, V., Setyan, A., Belis, C. A., Canonaco, F., Favez, O., Riffault, V.,  
5 Slowik, J. G., Aas, W., Aijälä, M., Alastuey, A., Artiñano, B., Bonnaire, N., Bozzetti, C.,  
6 Bressi, M., Carbone, C., Coz, E., Croteau, P. L., Cubison, M. J., Esser-Gietl, J. K.,  
7 Green, D. C., Gros, V., Heikkinen, L., Herrmann, H., Jayne, J. T., Lunder, C. R.,  
8 Minguillón, M. C., Močnik, G., O'Dowd, C. D., Ovadnevaite, J., Petralia, E., Poulain, L.,  
9 Priestman, M., Ripoll, A., Sarda-Estève, R., Wiedensohler, A., Baltensperger, U., Sciare, J.,  
10 and Prévôt, A. S. H.: ACTRIS ACSM intercomparison – Part 2: Intercomparison of ME-2  
11 organic source apportionment results from 15 individual, co-located aerosol mass  
12 spectrometers, *Atmos. Meas. Tech.*, 8, 2555–2576, doi:10.5194/amt-8-2555-2015, 2015.

13 Garbaras, A., Andriejauskiene, J., Bariseviciute, R. and Remeikis, V.: Tracing of atmospheric  
14 aerosol sources using stable carbon isotopes, *Lith. J. Phys.*, 48, 259–264,  
15 doi:10.3952/lithjphys.48309, 2008.

16 Garbaras, A., Masalaite, A., Garbariene, I., Ceburnis, D., Krugly, E., Remeikis, V., Puida, E.,  
17 Kvietkus, K., and Martuzevicius, D.: Stable carbon fractionation in size segregated aerosol  
18 particles produced by controlled biomass burning, *J. Aerosol Science*, 79, 86–96,  
19 doi:10.1016/j.jaerosci.2014.10.005, 2015.

20 Grieshop, A. P., Logue, J. M., Donahue, N. M., and Robinson, A. L.: Laboratory investigation  
21 of photochemical oxidation of organic aerosol from wood fires 1: measurement and  
22 simulation of organic aerosol evolution, *Atmos. Chem. Phys.*, 9, 1263–1277, doi:10.5194/acp-  
23 9-1263-2009, 2009.

24 Harrison, R. M., Beddows, D. C. S., Hu, L., [and](#) Yin, J.: Comparison of methods for  
25 evaluation of wood smoke and estimation of UK ambient concentrations, *Atmos. Chem.*  
26 *Phys.*, 12, 8271 – 8283, doi:10.5194/acp-12-8271-2012, 2012.

27 Hawkins, L. N. and Russell, L. M.: Oxidation of ketone groups in transported biomass  
28 burning aerosol from the 2008 Northern California Lightning Series fires, *Atmos. Environ.*,  
29 44, 4142–4154, doi:10.1016/j.atmosenv.2010.07.036, 2010.

Deleted: .

Deleted: .

1 Hennigan, C. J., Sullivan, A. P., Collett Jr., J. L., [and](#) Robinson, A. L.: Levoglucosan stability  
 2 in biomass burning particles exposed to hydroxyl radicals, *Geophys. Res. Lett.*, 37, L09806,  
 3 doi:10.1029/2010GL043088, 2010.

4 Herich, H., Gianini, M. F. D., Piot, C., Mocnik, G., Jaffrezo, J. L., Besombes, J. L., Prévôt, A.  
 5 S. H., and Hueglin, C.: Overview of the impact of wood burning emissions on carbonaceous  
 6 aerosols and PM in large parts of the Alpine region, *Atmos. Environ.*, 89, 64–75,  
 7 doi:10.1016/j.atmosenv.2014.02.008, 2014

8 Heringa, M. F., P. F. DeCarlo, R. Chirico, T. Tritscher, J. Dommen, E. Weingartner, R.  
 9 Richter, G. Wehrle, A. S. H. [Prévôt](#), and Baltensperger U.: Investigations of primary and  
 10 secondary particulate matter of different wood combustion appliances with a high-resolution  
 11 time-of-flight aerosol mass spectrometer, *Atmos. Chem. Phys.*, 11(12), 5945-5957, 2011.

12 Hildebrandt, L., Kostenidou, E., Lanz, V. A., Prévôt, A. S. H., Baltensperger, U.,  
 13 Mihalopoulos, N., Laaksonen, A., Donahue, N. M., and Pandis, S. N.: Sources and  
 14 atmospheric processing of organic aerosol in the Mediterranean: insights from aerosol mass  
 15 spectrometer factor analysis, *Atmos. Chem. Phys.*, 11, 12499–12515, doi:10.5194/acp-11-  
 16 12499-2011, 2011.

17 Hoffmann, D., Tilgner, A., Inuma, Y., and Herrmann, H.: Atmospheric stability of  
 18 levoglucosan: a detailed laboratory and modeling study, *Environ. Sci. Technol.*, 44, 694–699,  
 19 doi:10.1021/es902476f, 2010.

20 Honrath, R., Owen, R. C., Val Martin, M., Reid, J., Lapina, K., Fialho, P., Dziobak, M.,  
 21 Kleissl, J., and Westphal, D.: Regional and hemispheric impacts of anthropogenic and  
 22 biomass burning missions on summertime CO and O<sub>3</sub> in the North Atlantic lower free  
 23 troposphere, *J. Geophys. Res.*, 109, doi:10.1029/2004JD005147, 2004.

24 Huang, R.-J., Zhang, Y., Bozzetti, C., Ho, K.-F., Cao, J.-J., Han, Y., Daellenbach, K. R.,  
 25 Slowik, J. G., Platt, S. M., Canonaco, F., Zotter, P., Wolf, R., Pieber, S. M., Bruns, E. A.,  
 26 Crippa, M., Ciarelli, G., Piazzalunga, A., Schwikowski, M., Abbaszade, G., Schnelle-Kreis,  
 27 J., Zimmermann, R., An, Z., Szidat, S., Baltensperger, U., El Haddad, I., [and Prévôt](#), A. S. H.:  
 28 High secondary aerosol contribution to particulate pollution during haze events in China,  
 29 *Nature*, 514(7521), 218-222, doi:10.1038/nature13774, 2014.

30 Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., [Prévôt](#), A. S. H., Zhang, Q., Kroll, J. H.,  
 31 DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M.,

Deleted: Prevot

Deleted: Prevot

Deleted: Prevot

1 Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A.,  
2 Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara,  
3 P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, E. J.,  
4 Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y.,  
5 Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell,  
6 L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimonono, A., Sun, J. Y., Zhang, Y.  
7 M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M.,  
8 Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and  
9 Worsnop, D. R.: Evolution of organic aerosols in the atmosphere, *Science*, 326, 1525 - 1529,  
10 doi:10.1126/science.1180353, 2009.

11 Kanamitsu, M., Ebisuzaki, W., Woollen, J., Yang, S. K., Hnilo, J. J., Fiorino, M., and Potter,  
12 G. L.: NCEP-DOE AMIP-II reanalysis (R-2), *Bull. Am. Meteorol. Soc.*, 83, 1631–1643,  
13 doi:10.1175/BAMS-83-11-1631, 2002.

14 Kikas, U., Reinart, A., Pugatshova, A., Tamm, E., and Ulevicius, V.: Microphysical, chemical  
15 and optical aerosol properties in the Baltic Sea region, *Atmos. Res.*, 90(2–4), 211–222,  
16 doi:10.1016/j.atmosres.2008.02.009, 2008.

17 Lanz, V. A., M. R. Alfarra, U. Baltensperger, B. Buchmann, C. Hueglin, and A. S. H. [Prévôt](#):  
18 Source apportionment of submicron organic aerosols at an urban site by linear unmixing of  
19 aerosol mass spectra, *Atmos. Chem. Phys.*, 7, 1503–1522, 2007.

20 Lanz, V. A., Prévôt, A. S. H., Alfarra, M. R., Weimer, S., Mohr, C., DeCarlo, P. F., Gianini,  
21 M. F. D., Hueglin, C., Schneider, J., Favez, O., D’Anna, B., George, C., and Baltensperger,  
22 U.: Characterization of aerosol chemical composition with aerosol mass spectrometry in  
23 Central Europe: an overview, *Atmos. Chem. Phys.*, 10, 10453–10471, doi:10.5194/acp-10-  
24 10453-2010, 2010.

25 Lavanchy, V. M. H., Gäggeler, H. W., Schotterer, U., Schwikowski, M., and Baltensperger,  
26 U.: Historical record of carbonaceous particle concentrations from a European high-alpine  
27 glacier (Colle Gnifetti, Switzerland), *J. Geophys. Res. Atmos.*, 104, 21227–21236,  
28 doi:10.1029/1999jd900408, 1999.

29 Lemire, K. R., Allen, D. T., Klouda, G. A., [and](#) Lewis, C. W.: Fine particulate matter source  
30 attribution for southeast Texas using  $^{14}\text{C}/^{13}\text{C}$  ratios, *J. Geophys. Res.*, 107(D22), 4613,  
31 doi:10.1029/2002JD002339, 2002.

Deleted: Prevot



1 Levine, J. S. 1996. Biomass Burning And Global Change, Cambridge, MA, MIT Press.

2 Lewis, C. W., Klouda, G. A., and Ellenson, W. D.: Radiocarbon measurement of the biogenic  
3 contribution to summertime PM-2.5 ambient aerosol in Nashville, TN, Atmos. Environ., 38,  
4 6053–6061, doi:10.1016/j.atmosenv.2004.06.011, 2004.

5 Mann, G. W., Carslaw, K. S., Reddington, C. L., Pringle, K. J., Schulz, M., Asmi, A.,  
6 Spracklen, D. V., Ridley, D. A., Woodhouse, M. T., Lee, L. A., Zhang, K., Ghan, S. J., Easter,  
7 R. C., Liu, X., Stier, P., Lee, Y. H., Adams, P. J., Tost, H., Lelieveld, J., Bauer, S. E.,  
8 Tsigaridis, K., van Noije, T. P. C., Strunk, A., Vignati, E., Bellouin, N., Dalvi, M., Johnson,  
9 C. E., Bergman, T., Kokkola, H., von Salzen, K., Yu, F., Luo, G., Petzold, A., Heintzenberg,  
10 J., Clarke, A., Ogren, J. A., Gras, J., Baltensperger, U., Kaminski, U., Jennings, S. G.,  
11 O'Dowd, C. D., Harrison, R. M., Beddows, D. C. S., Kulmala, M., Viisanen, Y., Ulevicius, V.,  
12 Mihalopoulos, N., Zdimal, V., Fiebig, M., Hansson, H.-C., Swietlicki, E., and Henzing, J. S.:  
13 Intercomparison and evaluation of global aerosol microphysical properties among AeroCom  
14 models of a range of complexity, Atmos. Chem. Phys., 14, 4679–4713, doi:10.5194/acp-14-  
15 4679-2014, 2014.

16 Masalaite, A., Remeikis, V., Garbaras, A., Dudoitis, V., Ulevicius, V., and Ceburnis, D.:  
17 Elucidating carbonaceous aerosol sources by the stable carbon  $\delta^{13}\text{C}_{\text{TC}}$  ratio in size-  
18 segregated particles, Atmos. Res., 158–159, 1–12, doi:10.1016/j.atmosres.2015.01.014, 2015.

19 Matthew, B. M., Middlebrook, A. M., and Onasch, T. B.: Collection efficiencies in an  
20 Aerodyne aerosol mass spectrometer as a function of particle phase for laboratory generated  
21 aerosols, Aerosol Sci. Tech., 42, 884–898, doi:10.1080/02786820802356797, 2008.

22 Middlebrook, A. M., Bahreini, R., Jimenez, J. L., and Canagaratna, M. R.: Evaluation of  
23 composition-dependent collection efficiencies for the Aerodyne aerosol mass spectrometer  
24 using field data, Aerosol Sci. Technol., 46, 258–271, doi:10.1080/02786826.2011.620041,  
25 2012.

26 Minguillón, M. C., Perron, N., Querol, X., Szidat, S., Fahrni, S. M., Alastuey, A., Jimenez, J.  
27 L., Mohr, C., Ortega, A. M., Day, D. A., Lanz, V. A., Wacker, L., Reche, C., Cusack, M.,  
28 Amato, F., Kiss, G., Hoffer, A., Decesari, S., Moretti, F., Hillamo, R., Teinilä, K., Seco, R.,  
29 Peñuelas, J., Metzger, A., Schallhart, S., Müller, M., Hansel, A., Burkhardt, J. F.,  
30 Baltensperger, U., and Prévôt, A. S. H.: Fossil versus contemporary sources of fine elemental

Deleted: b

Deleted: a

Deleted: g

Deleted: c

Deleted: E

Deleted: A

Deleted: M

Deleted: S

Deleted: F

Deleted: P

Deleted: P

Deleted: L

Deleted: G

Deleted: A

1 and organic carbonaceous particulate matter during the DAURE campaign in Northeast Spain,  
2 Atmos. Chem. Phys., 11, 12067-12084, doi:10.5194/acp-11-12067-2011, 2011.

3 Mochida, M., Kawamura, K., Fu, P. Q., and Takemura, T.: Seasonal variation of levoglucosan  
4 in aerosols over the western North Pacific and its assessment as a biomass-burning tracer,  
5 Atmos. Environ., 44, 3511–3518, doi:10.1016/j.atmosenv.2010.06.017, 2010.

6 MODIS NASA LANCE - FIRMS, 2011, MODIS Active Fire Detections, Data set, Available  
7 on – line: <http://earthdata.nasa.gov/data/nrt-data/firms>.

8 Mohr, C., Huffman, J. A., Cubison, M. J., Aiken, A. C., Docherty, K. S., Kimmel, J. R.,  
9 Ulbrich, I. M., Hannigan, M., and Jimenez, J. L.: Characterization of Primary Organic  
10 Aerosol 25 Emissions from Meat Cooking, Trash Burning, and Motor Vehicles with High-  
11 Resolution Aerosol Mass Spectrometry and Comparison with Ambient and Chamber  
12 Observations, Environ. Sci. Technol., 43, 2443–2449, doi:10.1021/es8011518, 2009.

13 Mordas, G., Ulevicius, V., Plauškaitė, K., and Prokopčiuk, N.: Validation of the condensation  
14 particle counter UF-02M in laboratory and ambient conditions, Lith. J. Phys., 53 (3), 175-182,  
15 2013.

16 Ng, N. L., Herndon, S. C., Trimborn, A., Canagaratna, M. R., Croteau, P. L., Onasch, T. B.,  
17 Sueper, D., Worsnop, D. R., Zhang, Q., Sun, Y. L. and Jayne, J. T.: An Aerosol Chemical  
18 Speciation Monitor (ACSM) for routine monitoring of the composition and mass  
19 concentrations of ambient aerosol, Aerosol Sci. Tech., 45, 770-784,  
20 doi:10.1080/02786826.2011.560211, 2011.

21 Oanh, N. T. K., Ly, B. T., Tipayarom, D., Manandhar, B. R., Prapat, P., Simpson, C. D., and  
22 Liu, L. J. S.: Characterization of particulate matter emission from open burning of rice straw,  
23 Atmos. Environ., 45, 493 - 502, doi:10.1016/j.atmosenv.2010.09.023 2011.

24 Orasche, J., Schnelle-Kreis, J., Abbaszade, G., and Zimmermann, R.: Technical Note: In-situ  
25 derivatization thermal desorption GC-TOFMS for direct analysis of particle-bound non-polar  
26 and polar organic species, Atmos. Chem. Phys., 11, 8977-8993, doi:10.5194/acp-11-8977-  
27 2011, 2011.

28 Orasche, J., Seidel, T., Hartmann, H., Schnelle-Kreis, J., Chow, J.C, Ruppert, H., and  
29 Zimmermann R.: Comparison of emissions from wood combustion. Part 1: Emission factors

Deleted: ,

Deleted: C

Deleted: M

Deleted: R

Deleted: M

Deleted: C

Deleted: M

Deleted: C

Deleted: A

Deleted: A

Deleted: .

Deleted: E

Deleted: W

Deleted: Combustion

Deleted: F

1 and characteristics from different small-scale residential heating appliances considering  
2 particulate matter and polycyclic aromatic hydrocarbon (PAH)-related toxicological potential  
3 of particle-bound organic species, *Energy Fuels*, 26(11), 6695-6704, doi: 10.1021/ef301295k,  
4 2012.

5 Oros, D. R., Radzi bin Abas, M., Omar, N. Y. M. J., Rahman, N. A., and Simoneit, B. R. T.:  
6 Identification and emission factors of molecular tracers in organic aerosols from biomass  
7 burning: 3. Grasses, *Appl. Geochem.*, 21, 919–940, doi:10.1016/j.apgeochem.2006.01.008  
8 2006.

9 [Paatero, P.: Least squares formulation of robust non-negative factor analysis, \*Chemometr.\*](#)  
10 [Intell. Lab.](#), 37, 23–35, doi:10.1016/S0169-7439(96)00044-5, 1997.

11 Paatero, P., and Tapper, U.: Positive matrix factorization – a nonnegative factor model with  
12 optimal utilization of error-estimates of data values, *Environmetrics*, 5, 111–126,  
13 doi:10.1002/env.3170050203, 1994.

14 Paatero, P., Eberly, S., Brown, S. G., and Norris, G. A.: Methods for estimating uncertainty in  
15 factor analytic solutions, *Atmos. Meas. Tech.*, 7, 781-797, doi:10.5194/amt-7-781-2014,  
16 2014.

17 Popovicheva, O., Kistler, M., Kireeva, E., Persiantseva, N., Timofeev, M., Kopeikin, V., and  
18 Kasper-Giebl, A.: Physicochemical characterization of smoke aerosol during large-scale  
19 wildfires: extreme event of August 2010 in Moscow, *Atmos. Environ.*, 96: 405–414,  
20 doi:10.1016/j.atmosenv.2014.03.026, 2014.

21 Puxbaum, H., Caseiro, A., Sánchez-Ochoa, A., Kasper-Giebl, A., Claeys, M., Gelencsér, A.,  
22 Legrand, M., Preunkert, S., and Pio, C.: Levoglucosan levels at background sites in Europe  
23 for assessing the impact of biomass combustion on the European aerosol background, *J.*  
24 *Geophys. Res.*, 112 (D23S05), doi:10.1029/2006JD008114, 2007.

25 Reid, J. S., Prins, E. M., Westphal, D. L., Schmidt, C. C., Richardson, K. A., Christopher, S.  
26 A., Eck, T. F., Reid, E. A., Curtis, C. A., Hoffman, J. P.: Real-time monitoring of South  
27 American smoke particle emissions and transport using a coupled remote sensing/box-model  
28 approach, *Geophys. Res. Lett.*, 31, L06107, doi:10.1029/2003GL018845, 2004.

29 Rutter, A. P., Snyder, D. C., Schauer, J. J., DeMinter, J., and Shelton, B. : Sensitivity and bias  
30 of molecular marker-based aerosol source apportionment models to small contributions of  
31 coal combustion soot. *Environ. Sci. Technol.* 43, 7770-7777, 2009.

Deleted: C

Deleted: D

Deleted: S

Deleted: S

Deleted: R

Deleted: H

Deleted: A

Deleted: C

Deleted: P

Deleted: M

Deleted: P

Deleted: A

Deleted: H

Deleted: R

Deleted: T

Deleted: P

Deleted: P

Deleted: B

Deleted: O

Deleted: S

Deleted: .

Deleted: ,

Deleted: Paatero, P.: Least squares formulation of robust non-negative factor analysis, *Chemometr. Intell. Lab.*, 37, 23–35, doi:10.1016/S0169-7439(96)00044-5, 1997.¶

Deleted: .

Deleted: &

Deleted: ¶

1 Saarikoski, S., Sillanpaa, M., Sofiev, M., Timonen, H., Saarnio, K., Teinela, K., Karppinen,  
2 A., Kukkonen, J., and Hillamo, R.: Chemical composition of aerosols during a major biomass  
3 burning episode over northern Europe in spring 2006: Experimental and modelling  
4 assessments, *Atmos. Environ.*, 41, 3577–3589, doi:10.1016/j.atmosenv.2006.12.053, 2007.

5 Salazar, G., Zhang, Y. L., Agrios, K., and Szidat, S.: Development of a method for fast and  
6 automatic radiocarbon measurement of aerosol samples by online coupling of an elemental  
7 analyzer with a MICADAS AMS, *Nucl. Instrum. Meth. Phys. Res. B.*, 361, 163–167,  
8 doi:10.1016/j.nimb.2015.03.051, 2015.

9 [Schmidl, C., Marr, I. L., Caseiro, A., Kotianová, P., Berner, A., Bauer, H., Kasper-Giebl, A.,  
10 and Puxbaum, H.: Chemical characterisation of fine particle emissions from wood stove  
11 combustion of common woods growing in mid-European Alpine regions, \*Atm. Env.\*, 42\(1\),  
12 126–141, 2008.](#)

13 Sofiev, M., Siljamo, P., Valkama, I., Ilvonen, M., Kukkonen, J.: A dispersion modelling  
14 system SILAM and its evaluation against ETEX data, *Atm. Env.*, 40, 674–685,  
15 doi:10.1016/j.atmosenv.2005.09.069, 2006.

16 [Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.:  
17 NOAA's HYSPLIT atmospheric transport and dispersion modeling system, \*Bull. Amer.  
18 Meteor. Soc.\*, 96, 2059–2077, 2015.](#)

19 Sullivan, A. P., Holden, A. S., Patterson, L. A., McMeeking, G. R., Kreidenweis, S. M.,  
20 Malm, W. C., Hao, W. M., Wold, C. E. and Collett Jr. J. L.: A method for smoke marker  
21 measurements and its potential application for determining the contribution of biomass  
22 burning from wildfires and prescribed fires to ambient PM<sub>2.5</sub> organic carbon, *J. Geophys.*  
23 *Res.*, 113, D22302, doi:10.1029/2008JD010216, 2008.

24 Sullivan, A. P., A. A. May, T. Lee, G. R. McMeeking, S. M. Kreidenweis, S. K. Akagi, R. J.  
25 Yokelson, S. P. Urbanski and J. L. Collett, Jr.: Airborne characterization of smoke marker  
26 ratios from prescribed burning, *Atm. Chem. Phys.* 14(19): 10535–10545, 2014.

27 Sun, Y. L., Wang, Z. F., Dong, H. B., Yang, T., Li, J., Pan, X. L., Chen, P., Jayne, J. T.:  
28 Characterization of summer organic and inorganic aerosols in Beijing, China with an aerosol  
29 chemical speciation monitor, *Atmos. Environ.*, 51, 250–259,  
30 doi:10.1016/j.atmosenv.2012.01.013, 2012.

Deleted: M

Deleted: S

Deleted: M

Deleted: M

Deleted: P

Deleted: A

Deleted: D

Deleted: C

Deleted: B

Deleted: B

Deleted: W

Deleted: P

Deleted: F

Deleted: A

Deleted: O

Deleted: C

Deleted: A

Deleted: C

Deleted: S

Deleted: M

1 Szidat, S., Jenk, T. M., Gäggeler, H. W., Synal, H.-A., Fisseha, R., Baltensperger, U.,  
2 Kalberer, M., Samburova, V., Reimann, S., Kasper-Giebl, A., and Hajdas, I.: Radiocarbon  
3 ( $^{14}\text{C}$ )-deduced biogenic and anthropogenic contributions to organic carbon (OC) of urban  
4 aerosols from Zurich, Switzerland, *Atmos. Environ.*, 38, 4035–4044,  
5 doi:10.1016/j.atmosenv.2004.03.066, 2004.

6 Szidat, S., Salazar, G. A., Vogel, E., Battaglia, M., Wacker, L., Synal, H.-A., and Türlér, A.:  
7  $^{14}\text{C}$  analysis and sample preparation at the new Bern Laboratory for the analysis of  
8 radiocarbon with AMS (LARA), *Radiocarbon*, 56, 561-566, doi:10.2458/56.17457, 2014.

9 Timonen, H., Aurela, M., Carbone, S., Saarnio, K., Saarikoski, S., Mäkelä, T., Kulmala, M.,  
10 Kerminen, V.-M. Worsnop, D. R. and Hillamo, R.: High time-resolution chemical  
11 characterization of the water soluble fraction of ambient aerosols with PILS-TOC-IC and  
12 AMS. *Atmos. Meas. Tech.* 3:1063–1074, doi:10.5194/amt-3-1063-2010, 2010.

13 Turekian, V. C., Macko, S., Ballentine, D., Swap, R. J., and Garstang, M.: Causes of bulk  
14 carbon and nitrogen isotopic fractionations in the products of vegetation burns: laboratory  
15 studies, *Chem. Geol.*, 152, 181-192, doi:10.1016/S0009-2541(98)00105-3 1998.

16 Ulbrich, I. M., Canagaratna, M. R., Zhang, Q., Worsnop, D. R., and Jimenez, J. L.:  
17 Interpretation of organic components from Positive Matrix Factorization of aerosol mass  
18 spectrometric data, *Atmos. Chem. Phys.*, 9, 2891-2918, doi:10.5194/acp-9-2891-2009, 2009.

19 Ulevicius, V., Byčenkienė, S., Remeikis, V., Garbaras, A., Kecorius, S., Andriejauskienė, J.,  
20 Jasinevičienė, D., and Močnik, G.: Characterization of aerosol particle episodes in Lithuania  
21 caused by long-range and regional transport, *Atmos. Res.*, 98(2-4), 190-200,  
22 doi:10.1016/j.atmosres.2010.03.02, 2010a.

23 Ulevicius, V., Byčenkienė, S., Špirkauskaitė, N., and Kecorius, S.: Biomass burning impact  
24 on black carbon aerosol mass concentration at a coastal site: case studies, *Lith. J. Phys.*, 50(3),  
25 335–344, doi:10.3952/lithjphys.50304, 2010b.

26 Widory, D.: Nitrogen isotopes: Tracers of origin and processes affecting  $\text{PM}_{10}$  in the  
27 atmosphere of Paris, *Atmos. Environ.*, 41, 2382-2390, doi:10.1016/j.atmosenv.2006.11.009,  
28 2007.

29 Weimer, S., Alfarra, M.R., Schreiber, D., Mohr, M., Prévôt, A.S.H., and Baltensperger U.:  
30 Organic aerosol mass spectral signatures from wood burning emissions: Influence of burning  
31 conditions and wood type, *J. Geophys. Res.*, 113, D10304, doi:10.1029/2007JD009309, 2008.

Deleted: A

Deleted: R

Deleted: .

Deleted: T

Deleted: R

Deleted: C

Deleted: C

Deleted: W

Deleted: S

Deleted: F

Deleted: A

Deleted: A

Deleted: Prevot

1 Wiedensohler, A., Birmili, W., Nowak, A., Sonntag, A., Weinhold, K., Merkel, M., Wehner,  
2 B., Tuch, T., Pfeifer, S., Fiebig, M., Fjåraa, A. M., Asmi, E., Sellegri, K., Depuy, R., Venzac,  
3 H., Villani, P., Laj, P., Aalto, P., Ogren, J. A., Swietlicki, E., Williams, P., Roldin, P.,  
4 Quincey, P., Hüglin, C., Fierz-Schmidhauser, R., Gysel, M., Weingartner, E., Riccobono, F.,  
5 Santos, S., Gruning, C., Faloony, K., Beddows, D., Harrison, R., Monahan, C., Jennings, S. G.,  
6 O'Dowd, C. D., Marinoni, A., Horn, H.-G., Keck, L., Jiang, J., Scheckman, J., McMurry, P.  
7 H., Deng, Z., Zhao, C. S., Moerman, M., Henzing, B., de Leeuw, G., Löschau, G., and  
8 Bastian S.: Mobility particle size spectrometers: harmonization of technical standards and  
9 data structure to facilitate high quality long-term observations of atmospheric particle number  
10 size distributions. *Atmos. Meas. Tech.*, 5, 657–685. doi:10.5194/amt-5-657-2012, 2012.

11 Yttri, K. E., Simpson, D., Nøjgaard, J. K., Kristensen, K., Genberg, J., Stenström, K.,  
12 Swietlicki, E., Hillamo, R., Aurela, M., Bauer, H., Offenberg, J. H., Jaoui, M., Dye, C.,  
13 Eckhardt, S., Burkhardt, J. F., Stohl, A., and Glasius, M.: Source apportionment of the summer  
14 time carbonaceous aerosol at Nordic rural background sites, *Atmos., Chem. Phys.*, 11, 13339-  
15 13357, doi:10.5194/acp-11-13339-2011, 2011.

16 Yttri, K. E., Lund Myhre, C., Eckhardt, S., Fiebig, M., Dye, C., Hirdman, D., Ström, J.,  
17 Klimont, Z., and Stohl, A.: Quantifying black carbon from biomass burning by means of  
18 levoglucosan – a one-year time series at the Arctic observatory Zeppelin, *Atmos. Chem.*  
19 *Phys.*, 14, 6427-6442, doi:10.5194/acp-14-6427-2014, 2014.

20 Zawadzka, O., Makuch, P., Krzysztof, M., Markowicz, Z., Zieliński, T., Petelski, T., Ulevicius,  
21 V., Strzałkowska, A., Rozwadowska, A., and Gutowska, D.: Studies of aerosol optical depth  
22 with the use of microtops II sun photometers and MODIS detectors in coastal areas of the  
23 Baltic Sea, *Acta Geophys.*, 62 (2), 400-422, doi:10.2478/s11600-013-0182-5, 2013.

24 Zhang, Y. L., Perron, N., Ciobanu, V. G., Zotter, P., Minguillon, M. C., Wacker, L., [Prévôt](#),  
25 A. S. H., Baltensperger, U., and Szidat, S.: On the isolation of OC and EC and the optimal  
26 strategy of radiocarbon based source apportionment of carbonaceous aerosols, *Atmos. Chem.*  
27 *Phys.*, 12 (22), 10841–10856, doi:10.5194/acp-12-10841-2012, 2012.

28 Zhang, Y.-L., Huang, R.-J., El Haddad, I., Ho, K.-F., Cao, J.-J., Han, Y., Zotter, P., Bozzetti,  
29 C., Daellenbach, K. R., Canonaco, F., Slowik, J. G., Salazar, G., Schwikowski, M., Schnelle-  
30 Kreis, J., Abbaszade, G., Zimmermann, R., Baltensperger, U., Prévôt, A. S. H., and Szidat, S.:  
31 Fossil vs. non-fossil sources of fine carbonaceous aerosols in four Chinese cities during the

Deleted: Prevot

1 extreme winter haze episode of 2013, *Atmos. Chem. Phys.*, 15, 1299-1312, doi:10.5194/acp-  
2 15-1299-2015, 2015.

3 Zotter, P., Ciobanu, V. G., Zhang, Y. L., El-Haddad, I., Macchia, M., Daellenbach, K. R.,  
4 Salazar, G. A., Huang, R. J., Wacker, L., Hueglin, C., Piazzalunga, A., Fermo, P.,  
5 Schwikowski, M., Baltensperger, U., Szidat, S. and Prevot A. S. H.: Radiocarbon analysis of  
6 elemental and organic carbon in Switzerland during winter-smog episodes from 2008 to 2012-  
7 Part 1: Source apportionment and spatial variability, *Atmos. Chem. Phys.*, 14(24), 13551-  
8 13570, 2014.

9

10

The Baltic Sea Area



Figure 1. A) Map of the observation site, Preila (indicated by the red mark). Nearest major cities are Klaipeda (40 km north) and Kaliningrad (90 km south), B) Environmental pollution research station Preila and site surroundings (C).

Deleted: er



1 Table 1. Preila site surroundings 10 km

Site altitude	5 (m) a.s.l.	Terrain below site <sup>1</sup>	50.0 (%)
Median altitude	0 (m) a.s.l.	Standard deviation of altitude <sup>2</sup>	7 (m)
Total population	6831	Standard deviation of population	159
Mean population density	20 (km <sup>-2</sup> )	Standard deviation of population density <sup>3</sup>	13 (km <sup>-2</sup> )
Local population density	29.5 (km <sup>-2</sup> )		
Dominating land cover types (based on GLC2000)			
Water bodies (natural & artificial) (20)			84.9 (%)
Tree cover, needle-leaved, evergreen (4)			13.2 (%)
Tree cover, mixed leaf type (6)			1.1 (%)
Herbaceous cover, closed-open (13)			0.4 (%)


2

3

<sup>1</sup>Percentage of terrain within 10 km radius from the site that lies at lower altitudes than the site itself based on GLOBE 30" (arc-second) topography data. For an elevated site this percentage will be large (close to 100 %), while for sites within valleys or basins this percentage will be small. For sites within homogeneous terrain the percentage will be 50 %. Such sites can be assumed to be more representative for a larger area, while for sites in more complex terrain small circulation systems might influence the surface concentration field and introduce large heterogeneities.

<sup>2</sup>Standard deviation of population density within a 10 km radius from the site based on GPW3 2.5' (arc-minutes) population data. Large variations within the population density pattern around a site might introduce large differences in the pollutant levels depending on wind direction. Measurements at sites with small standard deviation of population density in the surroundings are therefore thought to be more representative of a larger domain and the mean population density.

<sup>3</sup>Mean population density within a 10 km radius from the site based on GPW3 2.5' (arc-minutes) population data. Small population densities are usually connected with little emissions. Measurements at sites with small population density in the surroundings are therefore thought to be more representative of a larger domain.



1  
2  
3 Figure 2. (A) Combined MODIS images observed from the Aqua satellite on 10 March 2014,  
4 showing numerous fires due to seasonal grass burning and 72-hour air mass backward  
5 trajectories from the fire regions arriving at Preila at 100 (red), 200 (blue) and 500 (green) m  
6 above ground level (AGL). (B) NAAPS model results showing surface smoke concentrations  
7 for the strongest stage (10 March 2014) (the color scale (from blue to purple) corresponds to  
8 the 7 levels of the contours that indicate the smoke mass mixing ratio ( $\mu\text{g m}^{-3}$ ) at the surface).  
9 Smoke optical depth at a wavelength of 0.55 microns. The contouring begins at  $1 \mu\text{g m}^{-3}$  and

1 doubles in magnitude for each successive contour. (C) Pressure level in Pa at the surface for  
2 2.5 degree latitude × 2.5 degree longitude global grids (NCEP/NCAR Reanalysis 1, 10 March  
3 2014). (D) PM<sub>2.5</sub> concentration (μg m<sup>-3</sup>) forecast utilized by the SILAM chemical transport  
4 model during the event of grass fires. (E, F) ACSM organics concentration (μg m<sup>-3</sup>)  
5 (measured in Preila) weighted air mass back trajectories of 48 h (for an arrival on 8 (E) and 10  
6 (F) March 2014) with an altitude endpoint of 500 m AGL.

Deleted: at

Deleted: in Pa

Deleted: ,

Deleted:

Deleted: ACSM organic concentration

Deleted: left

Deleted: right

Formatted: Font color: Black





Figure 3. Average daily concentration during event days (from 5 to 10 March 2014) and non-event days (14, 21, 23 and 27 March 2014) for levoglucosan, galactosan, mannosan (in  $\text{ng m}^{-3}$ ) and for elemental carbon (EC) and organic carbon (OC) in  $\mu\text{g m}^{-3}$ .

1



2

3 Figure 4. Stable carbon and nitrogen isotope ratio values of PM<sub>1</sub> in Preila station. Vertical and  
 4 horizontal lines represent carbon and nitrogen, respectively, isotope ratio characteristic values  
 5 for the sources of aerosol particles (Garbaras et al., 2008, 2015; Ulevicius et al., 2010a;  
 6 Widory 2007).

7

1 Table 2. Variation of the fractions of EC<sub>f</sub>, EC<sub>nf</sub>, OC<sub>f</sub>, OC<sub>nf</sub> and TC, EC and OC values during  
2 the study periods.

	EC <sub>f</sub>	EC <sub>nf</sub>	OC <sub>f</sub>	OC <sub>nf</sub>	TC	EC	OC
Date of collection	$\mu\text{g m}^{-3}$	$\mu\text{g m}^{-3}$	$\mu\text{g m}^{-3}$	$\mu\text{g m}^{-3}$	$\mu\text{g m}^{-3}$	$\mu\text{g m}^{-3}$	$\mu\text{g m}^{-3}$
2014.03.05	0.25±0.04	0.33	0.47±0.10	2.34±0.18	3.39±0.18	0.59±0.17	2.80±0.18
2014.03.07	0.21±0.04	0.61	0.39±0.12	2.80±0.20	4.01±0.23	0.81±0.24	3.31±0.20
2014.03.08	0.15±0.05	0.26	0.56±0.07	1.46±0.12	2.43±0.13	0.41±0.18	2.24±0.15
2014.03.09	0.46±0.16	0.95	0.95±0.18	4.98±0.36	7.28±0.43	1.36±0.63	6.32±0.35
2014.03.10	0.56±0.18	1.64	1.64±0.28	7.77±0.50	11.72±0.64	2.31±0.75	9.47±0.51

3

4




Figure 5. Average chemical composition and time series of NR-PM<sub>1</sub> OA for the entire study (A), B) Time series of the POA factor and percent contribution of the corresponding tracer species (levoglucosan, picene and hopanes) to total OA, C) Time series of the SOA factor, D) Relative source apportionment of TC during the BB event. Numbers indicate the total carbon absolute concentrations in µgm<sup>-3</sup>, variations of the mass concentrations of the SOC<sub>f</sub> and SOC<sub>nf</sub> (the whiskers above and below the boxes indicate the 1<sup>st</sup> and 3<sup>rd</sup> quartiles).




Figure 6. Mass spectra of SOA and POA, error bars represent the standard deviation of 20 PMF runs (A,C) and the scatter plots illustrate the relationship between SOA and  $\text{NH}_4^+$  (B) and POA with BC (D).



1 Table 3. Average percentage contributions of different sources

Relative contributions [%] to TC	POC <sub>f</sub>	POC <sub>nf</sub>	SOC <sub>f</sub>	SOC <sub>nf</sub>	EC <sub>f</sub>	EC <sub>nf</sub>	TC to PM1
2014.03.05	5.07	43.23	6.69	22.51	9.66	12.83	28.44
2014.03.07	6.26	43.58	5.66	19.14	6.58	18.79	37.62
2014.03.08	7.73	26.28	13.36	18.65	12.60	21.38	24.80
2014.03.09	4.55	41.30	4.38	13.12	12.48	24.17	51.27
2014.03.10	6.82	43.02	5.9	14.83	7.18	22.26	43.91
Relative contributions [%] to OC	POC <sub>f</sub>	POC <sub>nf</sub>	SOC <sub>f</sub>	SOC <sub>nf</sub>			
2014.03.05	6.55	55.78	8.63	29.04			
2014.03.07	8.39	58.38	7.59	25.64			
2014.03.08	11.70	39.80	20.24	28.25			
2014.03.09	7.18	65.20	6.91	20.71			
2014.03.10	9.66	60.96	8.36	21.02			

Deleted: P

Deleted: C

Deleted: D

Deleted: S

2

3




Figure 7. Probability distribution functions of the absolute daily contribution of POC<sub>f</sub> (A), POC<sub>nf</sub> (B), SOC<sub>nf</sub> (C), SOC<sub>f</sub> (D).




Figure 8. Probability distribution functions of Levoglucosan/BBOC (A) and EC<sub>nf</sub>/BBOC (B).