

We would like to thank the reviewers for their detailed comments and time that they have put into providing their feedback. The process of considering and adapting to all of these comments has considerably tightened and improved the paper. For this reason, we want to thank you both again.

**Reviewer #1:**

1. Agreed. This has been done. Other reductions in the abstract have also been made.
2. This is an excellent and well thought out suggestion. A significant amount of text has been added in these sections. While the LAI product is more robust overall, identifying its variance is harder over deforested regions due to the smaller magnitudes. On the other hand, while it is less robust in terms of number of channels, the ratio of the variance of the NDVI to the absolute NDVI can be indicative of rapid changes over both forested and non-forested regions, and seems to help provide an extension into regions which are partially but not fully disturbed.
3. Thank you for this update. It has been done.
4. Thank you very much for helping to clarify these important points about the measurements. All of these points have been added. Additionally, extra text describing the impacts and magnitude of these errors, as well as the model errors has been added. This has really helped the contents to improve.
5. Thank you for catching this important oversight. This reference has been corrected.
6. Thank you for talking more in depth about the spatial coverage issues of MISR. You are absolutely correct that in general this is the case. However, interestingly over this region of the world, MISR actually has a much better correlation with AERONET and a lower error. It is partially due to the fact that MODIS cannot clear the clouds as well, and also due to the fact that at very high AOD levels ( $AOD > 2$ ) that MODIS assumes that there is cloud, when in fact there is aerosol. I have added a reference to support this. I have also made careful observations with MODIS, AERONET, and MISR and found this to be the case, on monthly time scales. However, in order to obtain data at weekly or daily scales, the swath width is a huge issue, and in fact MODIS is far superior. Although I am sure that at the global scale, you are right, based on my work in Southeast Asia and my reading and talking with experts on the ground there, I believe that in this case, MISR is an appropriate instrument to use. I have added in extensive cautions however, as you have mentioned. It will be interesting to see how this works in future studies to see if others can also confirm these findings.
7. This is also an interesting point. There is recovery in greenness, as reflected in the NDVI, which is related to the chlorophyll, and in the moisture content of the canopy and the soil, which is better represented in the LAI term. However, most of the fires are occurring in regions that are heavily managed or that are reasonable unmanaged but are being converted to managed regions. Furthermore, when the Monsoon Rains arrive, there will then be sufficient water to overcome the moisture issue. For these reasons, any lag between the variables would be expected to occur on time scales which are being managed, and with the complete destruction posed by the extensive fires, are not expected to be so obvious. Perhaps a different approach to looking at the regrowth, for example in areas that are managed differently would yield an interesting solution, or in regions that use different types of crops to regrow after the fires, but that is beyond the scope of the study done here.
8. There are two ways in which to look at this issue. The first is, what fraction of the total variance is covered by the mode, and the second is what physical amount of change in the underlying variable is contained by this change in the variance. Clearly, and variance captured which is smaller than the error of the measurement itself is highly circumspect and to be confident, is ignored as not being meaningful in this analysis. Furthermore, if the absolute change in the magnitude of the associated pattern represented in the high variance region is sufficiently small, such as major changes in a mostly random manner, then it is likely to be more representative of noise than a signal. In both cases, the 5% value chosen here is related to the error in the measurements themselves. This is now more clearly made in the text and the reference is updated as well. In reality, such a strong constraint may not be required, especially since the bias in the measurement is low, but the point is to be more conservative scientifically. Thank you again for helping to clarify this.
9. This has been clarified in the text.

10. Considerable work has been put into re-arranging the figures, updating the fonts, etc. Additionally, some figures have been moved to supplementary data or have been added as new supplementary data.
11. This plot has been removed and the data has been presented in a different manner through other figures and tables, both in the main paper and the supplementary material. Thank you for the suggestion!
12. Corrected.
13. Corrected.

**Reviewer #2:**

1. 2015 was an El-Nino impacted year, starting in the latter period (August-October) in the Maritime Continent. The data as analyzed in this paper however ended in 2014. Given the strength of the El-Nino was even stronger than in 2006, it would make an interesting follow-up to see if this approach will be able to replicate the results.
2. An additional 2 supplementary figures have been added, showing the spatial distribution of the fires in January 2013 and September 2013. "Maps of the fires in January 2013 and September 2013 respectively are given in Appendices C1 and C2."
3. Agreed. Additional references have been added.
4. The daily GFED datasets have been used in a previous study, while FINN on the other hand as not been attempted by this author, and will be considered for future. However, both are still based on similar satellite-based approaches, although the specific satellites used do differ. But ,the results were found to not be dramatically different, since the underlying processes that were used to construct both of these datasets still suffer from the same basic issues of pixels obscured by cloud/smoke, events that are too small to be measured as Fire Hotspots, etc. will still be overlooked. For example, the author has personally observed and photographed multiple sizable fires in 2014 that show up in the AOD measurements, but do not appear in any of these datasets nor in the hotspot measurements of any of the satellites used by FINN and GFED.
5. Fire emissions inventories relate changes based on burned area and/or fire hotspots. They are not based on changes in LAI/NDVI/EVI. Yes, over regions where hotspots and burned area are correctly representing the whole of the fires, there is a relationship. However, given the errors involved with measurement of hotspots, especially due to their transient nature, in this region of the world, there is a significant mis-match between the spatial and temporal distribution of the land-use change. This is further exacerbated based on the underlying assumptions that go into the hotspot measurements, especially when anthropogenic land use change is such a prominent factor. In this region of the world, one of the major findings here is that this relationship is not the best way to proceed if one is interested to understand the fire emissions.
6. Agreed. This has been moved and re-worded.
7. The newer Collection 6 was not a well known product and was also not fully validated throughout the entire time period of data analyzed in this paper. It would be interesting to look into this in the future, to see what impact it has. It has been recommended not to mix the various different collection data types by NASA.
8. Yes, I have looked into this, and actually have a paper currently under review that analyzes this issue in much greater depth, using a different approach and set of measurements. The conclusion is relatively robust that the fire hotspots are indeed biased due to clouds, even over otherwise "cloud free" regions of Southeast Asia in many cases. I do not know how I can reference my own under-review paper however.
9. There is definitive seasonality in LAI. The drops associated with the fires are statistically significant across the entire dataset in this region. There are other drops which are considerably smaller and which are not always statistically significant as well. It is an interesting topic for further investigation. Perhaps grouping individual biomes or geographic regions will allow this to be more closely investigated, so as to obtain a baseline of what the natural variation is, albeit small in comparison for the region under investigation.
10. The range of fire counts, on a monthly basis, is from as low as 0 to as high as 5000 at level 8 and 600 at level 9. The article has been correspondingly updated.
11. Citations added.

12. There is a lag between the variables. However, the lag is not so obvious when weekly averaged, and is not-observable when monthly averaged. This is likely a reason why the weekly average product performs well. The fact of the matter is that the data is too sparse on a day-to-day basis to obtain reasonable statistics, while on a weekly basis, one can determine changes of magnitude and position. It is interesting to see if there are oddly-timed lags, such as around El-Nino and other large-scale phenomena, although these have been observed over the regions specifically examined in this work.
13. The fire hot-spot map is similar enough to GFED and FINN, while Cohen (2014) and Cohen and Wang (2014) provide additional maps from additional perspectives. The author has access to yet another map, but given that it is still under review, it cannot yet be cited.
14. Rubbish burning is an interesting question over this region. While I have not studied it specifically, there is a highly variable component. There are communities and regions where it is much more prevalent, while there are others where it is completely taboo.
15. Yes, these checks were done, as reflected in the text now.
16. A few references have been included.
17. Agreed, it is an interesting point, worthwhile of future study. Text has been added here to clarify this as an important issue, and what small step we have done to start to address this.
18. A very good question, which we can not easily answer here, since a lot of the burning is done by people and the regrowth is intentionally in a species different from what was there before. However, with a little additional work, this could make a great topic for future study.
19. Additional references have been added here.
20. This has been added.
21. This section has been made into a new numbered section.
22. This section has been combined into the section above.
23. A figure link has been placed here.
24. An additional reference relating to the use of NDVI in this manner has been added.
25. This has been clarified in the text.
26. This is explained in detail in the text. The timing is constant from year to year, and based on the calendar date, not the time of any fires that may exist in the surrounding area. This allows for proper comparisons across different years and times of the years to be made.
27. Agreed. This has been moved and the appropriate numbers updated.
28. These figures have been combined and updated in the text where they appear.
29. This is a good point. A sentence and additional references have been added. While this model still underestimates the measurements, it actually performs much better than most models in heavily biomass burning regions, which require a scaling factor of much more than the magnitude of this error. This is now made clear.
30. The text has been considerably condensed and additional points and emphasis have both been made, as recommended.
31. This figure has been removed and the text re-written.
32. This point has been acknowledged. It can already be seen in December 2013, so that is not made clear, and it is mentioned that this would have maximized in February 2014, although it is beyond the length of the data specifically used in this analysis.
33. This point has been made very clearly. Thank you. This makes the overall flow and conclusion stronger and more precise.
34. The sentence has been re-written.
35. Corrected.
36. The term has been explained in more detail, and the marker has been changed accordingly.
37. The figure order has been updated to reflect the text. Also, additional figures have been combined when they make sense to do so.
38. Corrected.
39. Changed.
40. Clarified.
41. Completed.
42. This has been appended and the terms have been clarified, including the indices.
43. Tables have been made more self-describing. Others have been re-numbered. While others have been combined or moved into the supplementary materials.

Decadal-Scale Relationship Between Measurements of Aerosols, Land-Use Change, and Fire over  
Southeast Asia

Jason Blake Cohen<sup>1\*</sup>, Eve Lecoœur<sup>2</sup>, and Daniel Hui Long Ng<sup>2</sup>

[1] Sun Yat-Sen University, School of Atmospheric Sciences, Guangzhou, China

[2] National University of Singapore, Tropical Marine Sciences Institute, Singapore

\* Corresponding Author: Jason Blake Cohen (email: [jasonbc@alum.mit.edu](mailto:jasonbc@alum.mit.edu))

## Abstract.

A simultaneous analysis of 13 years of remotely sensed data of land cover, fires, precipitation, and aerosols from the MODIS, TRMM, and MISR satellites and the AERONET network over Southeast Asia is performed, leading to a set of robust relationships between land-use change and fire being found on inter-annual and intra-annual scales over Southeast Asia, reflecting the heavy amounts of anthropogenic influence over land use change and fires in this region of the world. First, we find that fires occur annually, but with a considerable amount of variance in their onset, duration, and intensity from year to year, and from two separate regions within Southeast Asia from each other. ~~This variability is already partially understood from previous works, including the impacts of both inter-annually and intra-annually occurring influences such as the Monsoon and El-Nino events, but yet there are other as of yet unknown influences that also are found to strongly influence the results.~~ Second, we show that a simple regression-model of the land-cover, fire, and precipitation data can be used to recreate a robust representation of the timing and magnitude of measured AOD from multiple measurements sources of this region using either 8-day (better for onset and duration) or monthly based (better for magnitude) measurements, but not daily measurements. We find that the reconstructed AOD matches the timing and intensity from AERONET measurements to within 70% to 90% and the timing and intensity of MISR measurements from to within 50% to 95%. This is a unique finding in this part of the world, since cloud-covered regions are large, yet the ~~robustness of the model is still robustly is still~~ ~~capable of holding over many of these regions, including over regions~~ where ~~otherwise~~ no fires are observed and hence no emissions ~~would be expected to source-contributeion~~ to AOD ~~would otherwise be thought to occur~~. Third, we determine that while Southeast Asia is a source region of such intense smoke emissions, that ~~portions of it are it is~~ also impacted by ~~transport of smoke transported~~ from other regions ~~as well~~. There are regions in Northern Southeast Asia which have two annual AOD peaks, one during the local fire season, and the second smaller peak corresponding to a combination

of some local smoke sources as well as transport of aerosols from fires in Southern Southeast Asia, and possibly even from anthropogenic sources in South Asia. ~~Conversely, we show that Southern Southeast Asia is affected exclusively by its own local fire sources during its own local fire season.~~

Overall, this study highlights the importance of taking into account a simultaneous use of land-use, fire, and precipitation for understanding the impacts of fires on the atmospheric loading and distribution of aerosols in Southeast Asia over both space and time. Furthermore, it highlights that there are significant advantages using 8-day and monthly average values (instead of daily data), in order to better quantify the magnitude and timing of Southeast Asia fires.

## 1 Introduction

Southeast Asia has been experiencing major haze events over the past three to five decades, due to a combination of increased urbanization (Cohen and Wang, 2014; Cohen and Prinn, 2011) and large-scale conversion of forests by fire (Cohen, 2014; van der Werf et al., 2008; Taylor, 2010; Dennis, 2005). The underlying connections and mechanisms relating the sources and strength of fire-based emissions and observed intra-annual, inter-annual, and inter-decadal variations of fire events, with meteorology, land-use change, and anthropogenic driving factors are not well understood (van der Werf et al., 2006; Giglio et al., 2006; Hansen et al., 2008; Field et al., 2009). Moreover, recent studies have shown that the impacts these events have on the atmospheric loading of aerosols and the larger climate are becoming greater in both absolute terms and frequency (Langmann et al., 2009; Nakajima et al., 1999; Podgorny et al., 2003; Rosenfeld, 1999). Some of the heaviest events, which previously in the literature were only associated only with strong El-Nino induced drying events, are now being found to occur in connection with other, less extreme impacts on precipitation and even surface moisture, occurring at various scales including but not limited to the Indian Ocean Dipole, the Madden-Julian Oscillation, the shifting of the Inter-Tropical Conversion Zone, mountain induced waves, the land-sea breeze and localized convection (Fuller and Murphy, 2006; Wooster et al., 2012; Natalia Hasler and Avissar, 2009; Reid et al., 2013). The fact that so many factors are capable of influencing these large-scale events is likely to make prediction much more challenging, as is seen by the fact that since 2000, there were extreme events, of varying intensity, length, and duration, occurring in 2002, 2004, 2006, 2009, 2013, and 2014 in Southern Southeast Asia, the region covering Indonesia, Malaysia, Singapore, and Brunei, and every year except 2003 in Northern Southeast Asia, the region covering Thailand, Myanmar, Cambodia, Vietnam, and Laos (Neale and Slingo, 2003; Chang et al., 2005; Aldrian et al., 2007; Cohen, 2014; Wooster et al., 2012). To date, other than (Cohen, 2014), there have been no other studies that have looked at Southeast Asian fires both robustly and holistically, to the extent of being

able to reproduce both the extreme and low levels of aerosols at both the monthly and the decadal scale. Furthermore, other than (Cohen, 2014; Cohen and Wang, 2014) there are no other works that have been able to satisfactorily estimate the emissions of aerosols over this region of the world, from the fundamentals and over the entire time period, without scaling or other statistical enhancement techniques, to match with atmospheric column measurements, such as aerosol optical depth (AOD) and absorbing aerosol optical depth (AAOD).

Knowledge of the spatial, temporal, and magnitude of the emissions and atmospheric loadings is essential for our improved understanding of the environmental impacts of the fires. Emissions of aerosols and gases from these fires, include significant sources of black carbon (BC), organic carbon (OC), and ozone, and therefore contribute greatly towards impacting human health (Afroz et al., 2003), atmospheric radiative forcing (Wang, 2007; Jacobson, 2001; Ming et al., 2010; Ramanathan and Carmichael, 2008; Cohen et al., 2011), and cloud and precipitation properties (Huang et al., 2006; Tao et al., 2012; Wang, 2013). Furthermore, given the general circulation of the Earth, and the lack of precipitation during the dry season in the tropics, coupled with intense localized convection, a large portion of the emitted pollutants will spread widely in space and time, entering into the global-scale circulation patterns (Wang, 2007). Therefore, emissions from these regions during these times of the year may have a significant impact on people and the environment thousands of kilometers away from their source.

AOD can be used to quantify the emissions from the fires, since it is the non-dimensional vertical integral of the atmospheric extinction (the sum of scattering and absorbance) of solar radiation due to aerosols. AOD is useful since it can be measured by a combination of land-based and space-borne instruments (Holben et al., 1998; Petrenko et al., 2012; Dubovik et al., 2000). The extinction is in turn a function of the vertical aerosol mass and size distributions as well as chemical, physical, and optical properties. These values in turn are a function of the emissions and gasses from fires and other various anthropogenic sources, in-situ processing, washout from



precipitation, and atmospheric transport. Hence, the emissions of primary BC and OC from these fires, coupled with other secondary species, has a functional relationship with the change in the AOD, which otherwise would not have occurred over these fire regions and downwind, at these specific times, if the fires were not present.

This paper uses these relationships and goes one step further, to make the link between measurements of land-use change and fires directly with the atmospheric column measurements, ~~since for with fires, the intermediary step between the two with emissions being the by-product in the middle~~. This is because rapid conversion of forests, agricultural lands, and associated waste products by burning is one of the primary sources of aerosols throughout Southeast Asia (Langmann et al., 2009; Miettinen et al., 2013). However, little is known about the exact spatial and temporal distribution of these fires (Fu et al., 2012; Chen et al., 2016; Zhou et al., 2016). Furthermore, the inter-annual and intra-annual variability of biomass burning and its associated underlying mechanisms are also not well understood or constrained by measurements, leading to the current poor understanding of fires impact on the local and global aerosol climatology (van der Werf et al., 2006). Furthermore, Southeastern Asia is often covered with clouds, which further complicates detecting both fire and the pollution that comes from it (Miettinen et al., 2013; Giglio et al., 2006; Remer et al., 2013). A few studies have looked at this and give estimates that the emissions are underestimated, up to a factor of 4 times (Giglio et al., 2003, 2006; Petrenko et al., 2012; Cohen, 2014; Cohen and Wang, 2014).

Given that large-scale fires lead to abrupt and definitive changes in the vegetative properties, we employ a set of measures of land surface properties which have a long time-record, such as LAI (Leaf Area Index), NDVI (Normalized Difference Vegetation Index), and the number of 1km by 1km pixels with a measured fire (~~FireCount~~Fire Count). While we know that some changes may be masked, obscured, or otherwise missing, any observed abrupt changes in these variables or the land's properties itself must be linked at a minimum with any observed changes in the AOD itself.

Moreover, since the onset and the offset on the Asian Monsoon controls the start and end of the fire seasons by rapidly changing from relatively dry to intensely wet and visa versa (Hansen et al., 2008), large scale changes in the monthly-scale precipitation is a proxy for the ability of the fires to occur, as well as washout of aerosols. Therefore, precipitation is also intimately linked with measured AOD over Southeastern Asia. This is even more important given that there are only very few studies that have been able to quantify emissions over this region successfully, over the decadal scale, without resorting to statistical scaling, in relation to measured AOD and AAOD. Furthermore, the few emissions datasets that have been made are not capable of working at a higher frequency than monthly. Additionally, they have not been directly linked to the changes in the land surface properties that should be driving them (Cohen and Wang, 2014; Cohen, 2014). ~~One of the most important findings we make is to carefully examine the validity of looking at the data on a daily, versus weekly, versus monthly basis. Although most of the published literature looking at these interactions leans towards using high frequency daily data (or higher frequency data still, where available), we determine and validate that using weekly or monthly average data leads to a far better ability to accurately reproduce the measured values, explain why that is the case, and then quantify some of the impacts and limitations of this result.~~

## 2 Data and Methods

Several remotely sensed and surface measurements of the surface land properties (LAI and NDVI), the number of active fires (Fire Count), aerosol (AOD), and precipitation (rainfall) are used in this study. These are used in conjunction with advanced analytical procedures to determine the regions which contribute the most to the variance of the impact of fires on the atmosphere loading of aerosols as observed by the AOD. This analysis, in addition to its own results, leads to the production of a simple statistical multi-year constrained model, which is shown to be capable of reproducing the AOD as a function of the land use, fire, and precipitation measurements, even in

additional years, and even as tested against measurements of AOD from different sources. All of the details of the measurements used, the procedures and methods employed, and the statistical and analytical techniques employed are detailed below.

## 2.1 Geography

The domain of interest for this study is Southeast Asia, which we define here as the region spreading from 90°E to 130°E in longitude, and from 14°S to 23°N in latitude (see Figure 31). The subregion defined as Northern Southeast Asia is defined by a mostly large continental land masses and a single Wet Season each year, and consists of Thailand, Myanmar, Cambodia, Laos, Vietnam, and parts of Southern Greater China. The subregion defined as Southern Southeast Asia is defined by a mixture of land and water and has two Wet Seasons each year, and consists of Malaysia, Indonesia, Brunei, and Singapore. Maps of the fires in January 2013 and September 2013 respectively are given in Appendices C1 and C2.

## 2.2 Measured Data

For the basic remotely sensed measurements used in the analysis, model construction, and results, we use remotely sensed variables from the MODIS instrument on both the TERRA and AQUA satellites. Measurements of AOD (Remer et al., 2005; Levy et al., 2013) are from Collection 6, Level 2 product, swath-by-swath at 0.55 micron, and consist of both over land and over ocean, cloud-cleared pixels, measured daily with a spatial resolution of 10km by 10km at nadir. Each swath of only quality controlled pixels of AOD data, from January 1, 2001 through December 31, 2013, has been interpolated onto a consistent and standardized 0.1° by 0.1° square grid.

It has been shown that there is an  $\pm 0.02 - 0.10 \cdot \text{AOD} + 0.04 + 0.1 \cdot \text{AOD} \pm 0.05 \cdot \text{AOD}$  biased uncertainty in the measurement of AOD of  $-0.02 - 0.10 \cdot \text{AOD} + 0.04 + 0.1 \cdot \text{AOD} \pm 0.05 \cdot \text{AOD}$  over the ocean and  $\pm 0.05 + 0.15 \cdot \text{AOD}$  over the land (Remer; Levy et al., 2013; Sayer et al., 2012). However, over this region, the magnitude of the

“noisy floor” is small compared to the linear term, given that the AOD in polluted regions goes as high as 1.5 to 2.0. And while this linear term seem to not be too small, it is actually quite small compared to the difference between the peaks and the troughs as obtained by the variance maximizing technique. Additionally, ~~However,~~ as shown by (Cohen and Wang, 2014) and others, this error is sufficiently small as to not impact the end results, especially when compared with the uncertainties in the current best-generation of models and the dynamics of the atmosphere itself. In these cases, the models tend to be lucky to obtain measurements within a 20% to 30% range of the measurements, and often perform more poorly than this (e.g. Cohen and Prinn, 2011; Cohen and Wang 2014). AOD is a measure of the vertical sum of the extinction of sunlight (scattering plus absorption) through the atmosphere due to aerosol particles, and therefore is a function of the atmospheric loading of aerosols, washout from precipitation, and the vertical, size, and optical properties of the aerosols. Hence, there is a physical relationship between measured changes in AOD and the emissions and subsequent in-situ atmospheric processing of aerosols. It has been shown that strong spatial and temporal variability in AOD measurements over this part of the world are due to biomass burning from this region of the world, while large measurements of AOD which mostly only co-vary only with precipitation (washout) are more consistent with urban emissions (Cohen, 2014; Cohen and Wang, 2014).

To estimate the land-surface and fire responses we also use the measured values of LAI, NDVI, and ~~FireCount~~ Fire Count from MODIS (Nightingale et al., 2008; Yang et al., 2006; Huete et al., 1999; Giglio et al., 2003). Measurements of LAI and ~~FireCount~~ Fire Count (Collection 5.1, Level 2 product) are made on an 8-day average basis at 1km by 1km horizontal resolution. While for NDVI the measurements are on a 16-day average basis at 1km by 1km horizontal resolution. Each product is then aggregated onto the same consistent and standardized 0.1° by 0.1° square grid used for the AOD. All measurements only use data which has been quality assured to be cloud free. However, in this region, there are some optically thin clouds that will not be picked up, and this

may significantly bias the measurements of **FireCount** **Fire Count**, which are inherently based on IR measurements, but should not be as impacting on LAI and NDVI, which both depend mostly on measurements in the visible bands.

LAI is chosen since it represents the amount of leaf material in an ecosystem and hence is useful both for identifying if there was a sudden change in the amount of vegetation available and its condition (Asner et al., 2003), such as expected after leaves are consumed in a fire. It is geometrically defined as the total one-sided area of photosynthetic tissue per unit ground surface area. LAI values range from 0 for bare ground, to the range of 1 to 4 for grassland and crops, to the range of 5 to 9 for plantations, and as high as 10 for dense conifer forests. **One of the large drawbacks of using LAI in this region of the world is that it is hard to analyze the variance in the LAI over areas that are used for non-forest agriculture. This is because the LAI is considerably lower than the tropical primary and secondary forests. Hence, after a burning event, the absolute magnitude of the LAI and hence the amount of variance is lower. Yet, this is the more robust land surface variable, given that it uses many of the wavebands from MODIS. For this reason, the variance in the LAI is most helpful in determining deforestation from fire, particularly in regions which are not found to have hotspots.**

**FireCount** **Fire Count** determines how many of the pixels within the area have an active fire. It is based on a two factors, first if there is a sufficient amount of infrared emissions to determine that there is an absolute detection of a fire of sufficient strength. The second factor is whether the detected surface temperature is sufficiently variable as compared to the surrounding pixels. Given the complexity involved with using infrared and visible streams for the fire count, as well as the possibility of thin clouds obstructing this measurement, we only use quality assured **FireCount** **Fire Count** values, those with a value corresponding to 7 or more. **In this study, it is found that the number of Fire Count can vary from 0 to more than 5000 (with a corresponding value of 8) or more than 600 (with a corresponding value of 9) on a monthly basis.**

NDVI is also chosen since it represents a measure of the health of the vegetation. NDVI is mathematically calculated from the visible (VIS) and near-infrared light reflected (NIR) by the vegetation as follows:  $NDVI = \frac{NIR - VIS}{NIR + VIS}$ . Healthy vegetation absorbs most of the visible light that

hits it, and reflects a large portion of the near-infrared light. On the other hand, unhealthy or sparsely healthy vegetation, such as after being burned, reflects more visible light and less near-infrared light. Given this formula, a value close to zero (-0.1 to 0.1) implies that the land is barren with respect to living and green vegetation, whereas values close to +1.0 correspond to the highest density of healthy green leaves. NDVI is an ideal way to search for the ratio of the magnitude of the variance to the absolute mean. This is because the variance in the change in the health is actually proportionate to the initial value. In this case, while the overall variance is not too much throughout the region, the ratio is considerably high in regions which undergo rapid change such as from burning. However, such changes are not very useful for looking at small changes over large periods of time, and more are useful at looking at changes occurring over short periods of time. This is one way to overcome the issue of regeneration, either due to natural regrowth or due to anthropogenic planting.

Furthermore, since the onset of the monsoon brings sufficiently large amounts of precipitation that it usually leads to the end of the fire season (Cohen, 2014; Natalia Hasler and Avissar, 2009), knowledge of the rainfall rate is important. For this, we use TRMM measurements of precipitation, as generated by the 3B42 algorithm. This produces daily average precipitation measurements at 0.25° by 0.25° spatial resolution over the areas of interest for this work.

To validate the results, we also use two additional measurement platforms for AOD from AERONET and MISR. From AERONET (Holben et al., 1998) we either use available AOD at 0.55 microns or interpolate the surrounding wavelength-specific measurements to 0.55 microns, at 9 different stations (see Figure 3) located in the region of interest. We use all individual Level 2.0 data

points, cloud screened and validated, and then averaged to form a daily value, where a sufficient amount of data is available. At the four stations where there is insufficient data, we use individual Level 1.5 data points. However, before forming the daily average value in the case of Level 1.5 data, we only retain the AOD measurements when the corresponding Angstrom Exponent is larger than 0.2, giving us reassurance that the product is relatively cloud-free. This has been tested by varying the sensitivity from 0.1 to 0.4 (the minimum physically acceptable value must be positive) and there is little change in the end result. Although AERONET is the most precise measurement platform for AOD, it is limited in spatial coverage. Therefore we also use measurements from MISR (Diner et al., 1998; Kahn et al., 2010; Holben et al., 1998) of AOD at 0.55 microns, with a monthly temporal resolution and a  $0.5^\circ$  by  $0.5^\circ$  spatial resolution. The reason for choosing MISR is that it has a smaller error with respect to AERONET over this region of the world than any other satellite platform, (Petrenko and Ichoku, 2013) which allows us to provide spatially distributed validation. Although MISR has a more narrow swath-width than MODIS, in this region of the world, there are actually more data points that are retrieved at the AERONET stations and that the error is lower in comparison to the AERONET Measurements. This is partially due to the fact that it is able to cloud process and clear more efficiently than MODIS due to the ~~This is in part due to a combination of the more narrow swath width, as well as its ability to approximate the spherical~~ fraction. Additionally, the fact that MISR is able to measure AOD levels greater than 2.0 allows it to actually obtain more pixels on a monthly basis over this region than MODIS. However, ~~the~~ major downside is that only at a monthly average or lower frequency is available, with the monthly dataset having from 4 to 8 data points per measurement. It is therefore effective over this region at obtaining a spatial distribution upon which to extend the more precise AERONET results. However, this helps quite a bit with the cloud clearing statistics. Combining these together allows the use of the higher quality AERONET data as an anchor, where it is available, to evaluate any errors in the magnitude between the model and the measurements, even away from the source, so long as it is

still in the same geographical region (as described below). It also provides a means for investigating how error propagation between various different measurement sources can be quantified.

All of the data used has been taken from from January 2001 through December 2013. In the case of remotely sensed data, it was first interpolated (in the case of AOD) or aggregated (in the case of ~~FireCount~~ Fire Count, NDVI, and EVI) onto a  $0.1^\circ$  by  $0.1^\circ$  square grid, using only quality assured data. These gridded, data sets, were then aggregated or interpolated respectively to the temporal resolution used, either 1-day, 8-day, or monthly average temporal resolution, to make them consistent. AERONET measurements have also been taken using whatever data was available over the same respective 1-day, 8-day, and monthly periods, and have been considered to be representative of the entire corresponding  $0.1^\circ$  by  $0.1^\circ$  box in which they are located. One of the significant advances of using this approach is the ability to analyze how the results are improved by using data with different temporal variability.

### **2.3 Variance Maximizing Technique**

Aerosol emissions and resulting changes in AOD in the Southeastern Asia region mainly comes from two types of sources: urban/anthropogenic and fires. Emissions of aerosols from urban/anthropogenic include those from cities, transportation, and industrial processes, which generally include temporally and geographically regular combustion of coal, oil, and natural gas throughout the year. On the other hand, emissions of aerosols from fires, which include clearing of forests, agriculture, peat, and rubbish, are more highly irregular over space and time, preferentially occurring under certain economic conditions as well as during periods of dryness, due to either changes in irrigation or under the influence of various meteorological/climatological conditions (Cohen, 2014). As the ultimate goal of this study is to develop an understanding and constraint on the absolute source of aerosol emissions, and since fire is the most uncertain contribution in this region, therefore the analytical technique must target the large amount of variance in the measured



fields of the AOD. ~~In specific,~~ Those regions which both contribute the most to the variance of the AOD field as well as correspond to a large annual amount ~~of of AOD on an~~ absolute basis, are the regions which which are most likely fires. A simple check of the geography ~~has been performed to~~ eliminate any false positives that are known to be urban or industrial regions, ~~of which there are at least 3 in the regions under study: in Vietnam, Indonesia, and Malaysia.~~ However, it is possible that rapidly developing industrial uses of the land, such as new large mill-towns in Indonesia (as witnessed by the author), were not fully identified. Further, observed land-use changes were considered to be reasonable if they corresponded to ~~should correspond~~ reasonable changes in the values of ~~, and the~~ NDVI and LAI. ~~are used for this purpose.~~

To achieve these goals, we first employ the Empirical Orthogonal Functions/Principal Component Analysis technique (EOF/PCA) on the 8-day average AOD product. This is one of the beautiful things about using the EOF approach: patterns in the variance of the data search for the set of the relative maxima. Therefore, since the process searches for the highest and lowest values and gradients in space and time, any unbiased error in the measurements, will not significantly impact the result. Furthermore, the 8-day average product was chosen, so that it could take full advantage of the higher frequency of the MODIS data, when compared with the MISR data. Additionally, the lifetime of the aerosol plume is roughly on order of this period of time, given the low amount of precipitation and the high amount of aerosols lofted into the atmosphere due to the heat from the fires, making source/sink and overall statistical properties robust (Cohen, 2014; Lin et al., 2009; Lin et al., 2014).

The specific EOF/PCA analysis decomposes the 8-day AOD data  $F$  into subcomponents. Each subcomponent is orthogonal to the whole, and can be ordered based on the overall contribution to the fractional amount of the overall variability (Bjornsson and Venegas, 1997). This is done by decomposing the measurements into independent (orthogonal) spatial/geographic modes  $S_i$  and their associated temporal/time modes  $T_i$ , as explained in **EQUATIONS 1-5**, where  $a_{ij}$  are

the individual measurements ( $i$  is the marker indicating the ordered number in latitude/longitude, and  $j$  is the individual marker indicating the marker in time), and  $c_i$  and  $y_i$  which are the corresponding decomposed values of the spatial and temporal maps accordingly.

$$F = \begin{pmatrix} a_{11} & \cdots & a_{1M} \\ \vdots & \ddots & \vdots \\ a_{N1} & \cdots & a_{NM} \end{pmatrix} \quad (1)$$

$$F^T F = CY^T YC^T \quad (2)$$

$$S_i = \begin{pmatrix} c_{1i} \\ \vdots \\ c_{Mi} \end{pmatrix} \quad (3)$$

$$T_i = \begin{pmatrix} y_{1i} \\ \vdots \\ y_{Ni} \end{pmatrix} \quad (4)$$

$$F = YC^T = T_1 S_1^T + \cdots + T_N S_N^T \quad (5)$$

## 2.4 Regression-Fit Model Connecting Land Use Change to AOD

Along with the analysis, we also employ a simple multi-variable linear regression model to predict AOD from measured land-use and meteorological variables. This approach is adapted because of the physical nature of the relationship between these variables. Fires lead to a direct drop in LAI in currently growing vegetation through the combustion process. In the case of agriculture which has already been harvested, the LAI would have previously dropped, while the dried products are left to burn. Similarly if there is a change in the vegetation/agricultural state after the fire, this should show up by a restored LAI, although at a different magnitude. NDVI would similarly be

impacted, as the chlorophyll is combusted along with the plant material that is associated with it.

Furthermore, the hypothesized loss of efficiency of the land surface associated with the fires would show up as a lower-frequency change in the NDVI. Due to these reasons, there may be a lag expected between the occurrence of the fire and the change in the land-use variable. However, given the rapid rate of regrowth over this region, and the high degree of cloud-cover, it is found that day-to-day passes or information are not very reliable. It is uncertain how much lag would be expected in weekly-averaged or two-weekly averaged products. Although our results have determined that in fact the relationship which is based on no lag produces the best fit. Hence, one of the objectives is to quantify the impacts of different averaging kernels applied to the measurements.

There is also evidence that in regions where dryness is an issue, which it certainly is during the extremes of the dry season throughout Southeast Asia, that NDVI recovers slower than LAI. This would certainly be the case in regions in which peat is being drained or has recently been drained, such as the Southern Southeast Asia Region, or in regions where there is little to no irrigation, such as the Northern Southeast Asian Region (Hope et al, 2007; ; Morawitz et al., 2006; Cuevas et al., 2008). The clear indication here is that the rate of greenness regrowth, as observed by the change in the NDVI may not relate to the canopy and soil moisture regrowth, which is more related to the LAI due to the additional bands in the NIR. However, in regions in the topics that are either managed or are fed by the arrival of the monsoon, this is not expected to be a significant issue, and hence a possible reason why little lag is actually observed.

During the dry season, based on how dry it is, will impact the amount, intensity, and duration of the fires as a whole. In practice, years with wetter dry season or a drier dry season should have a reduction in the intensity of the fires as well as their geographic spread, although it will not necessarily lead to them being altogether suppressed. This relationship is slightly more complex, since there are cases where anthropogenic water due to irrigation, burning occurring on very wet peat, or fast-moving thunderstorms, can make the ground quite wet, but still continue to

burn, thereby leading to an increase in emitted aerosol, and hence AOD, due to a switch of the type of fire from flaming to smoldering (Field et al., 2009; Saatchi et al., 2013). However, these cases are over and beyond the approach taken here, and are still not yet fully understood. It is thought that surface wetness is critical for this switch, although in theory this is partially a function of the LAI, NDVI, and precipitation, and hence could be approximated to first order using the approach employed here, with the physical variable itself at least being partially captured (Fisher et al., 2009; Phillips et al., 2010; Wohl et al., 2012). Another advantage is that low-temperature fires, which may otherwise go undetected, can still be represented, since they still impact changes in terms of AOD, LAI, and NDVI.

To ensure that the impact of fires is physically as expected on AOD, in which an increase in fire should lead to an increase in emissions and hence AOD, we employ two different regression equations. Both equations use LAI, NDVI and Precipitation as predictive variables R2 (Equation 7), while only one R1 (Equation 6) also uses FireCount. The regression coefficients  $\alpha_i$ ,  $\beta_i$ ,  $\gamma_i$ , and  $\delta_i$  are computed by minimizing the root-mean-square errors of the equations EQUATIONS 6,7. Using these constrained values, the AOD can be approximated during different seasons or over different areas, such as those which are cloud-covered and hence do not show measurements. These reconstructed values are generated and specifically compared against AOD values from other measurement platforms, specifically MISR and AERONET.

$$AOD_{MODIS,R1} = \alpha_1 * LAI + \beta_1 * NDVI + \gamma_1 * RAIN + \delta_1 * FireCount \quad (6)$$

$$AOD_{MODIS,R2} = \alpha_2 * LAI + \beta_2 * NDVI + \gamma_2 * RAIN \quad (7)$$

Since the nature of the land-use change, the amount of precipitation, the state of native vegetation, and the strengths and timing of the AOD signal are different over the two regions S1 and S2, we compute the fitting for AOD over each region separately. This helps us better quantify and understand the functional relationships between these variables under the different land use types,

land-use management practices, and climatologies, when the fires actually do occur. This is especially important in Southern Southeast Asia, where there is stronger year-to-year variability, the issue of cloud-cover is much more pronounced close to the equator, there are only very few ground station measurement sites, ~~and~~ vastly different sets of anthropogenic land use policies in different regions, and different magnitudes of fire emissions.

To test separately only the fire-occurring seasons, we define fire activity periods over each region as the days during which which  $T1$  and  $T2$  respectively are above a certain threshold  $\tau_{\text{North}}$  and  $\tau_{\text{South}}$ . Different thresholds for  $T1$  and  $T2$  are tested, based on the percentile  $P$  of the time series beneath the point  $\tau$  if the time series were to be regrouped and sorted. We hence use the points  $P=\{0.90,0.835,0.75\}$ . Since this method is testing for the extreme values in the AOD variance, or when the fires are occurring, this method proves to be methodologically suitable, as it is further providing a constraint on the more extreme conditions, and when the pattern is most significant. The values chosen are not arbitrary, as they are based on the statistical robustness of the magnitude of the fields  $S_i \cdot T_i$ . However, the point of the sensitivity analysis is to quantify at what point the errors in the analytical technique are no longer able to statistically retrieve the maximum contributions to the variance of signal, as compared to just picking up the variance induced by the unbiased errors in the measurements themselves.

### 3. Results

#### 3.1 Decadal-Scale Analysis of Remotely Sensed Measurements of AOD and Land Surface

##### Properties

The subsequent analysis performed using the variance maximizing analytical technique only retains those modes  $S_i, T_i$  that explain at least 5% of the total variability. This is to ensure that any signal found is larger than the uncertainty in the measurements themselves, and hence should be physically relevant. Specifically, since the MODIS AOD uncertainty is 5%, therefore we need at

least 5% of the variance for a mode to represent something useful (Bjornsson and Venegas, 1997).

Using this constraint, there are two modes  $i=\{1,2\}$  that explain the variability in the 8-day AOD

measurements (see Figure 23). 38% of the variability in the AOD field maps to region  $i = 1$  as shown in Figure 2a, and which we will hence refer to as Northern Southeast Asia. 13% of the variability in the AOD field maps to region  $i=2$  as shown in Figure 2b, which we will hence refer to as Southern Southeast Asia. The next largest mode contributes less than 5% to the total variance in the AOD field, and therefore is indistinguishable from other sources of variability and error, such as non-linear effects of El-Nino, planetary dynamical events such as the MJO, regional dynamical events, small-scale perturbations, short-term anthropogenic events, un-accounted for variations in cloud-cover, bias in the data, new urbanization around the edges of the growing megacities, and such.

The physical relevance of these mathematical modes is established by correlating the computed measured average AOD over the respective regions  $S_i$ , as a time series, with the respective Principal Component  $T_i$ . The modes are found to be highly correlated with both the AOD over Northern Southeast Asia ( $R^2=0.86$ ,  $p<0.01$ ) and the AOD over Southern Southeast Asia ( $R^2=0.86$ ,  $p<0.01$ ), as shown in Figures 2c and 2d.

Over Northern Southeast Asia there is a partially bi-annual peak, with some years having a single peak and others have two peaks. The major peak, which is the more pronounced or sole peak, occurs every year in the measured AOD averaged over  $T_1$  during the latter part of the local dry season (from mid-February to late-April). Looking at the average value of the time series of the AOD measurements over  $S_1$ , it is found that the AOD peaks at the same time as  $T_1$  peaks, and that the average AOD ranges from 0.46 to 0.86, depending on the year. The smaller peak occurs in August and September as shown in  $T_1$  in most of the years (but not in 2008, 2010, and 2011). Similarly, the average of the measured AOD over the region  $S_1$  during the same months and years has a corresponding peak ranging from 0.40 to 0.63 during the years when the second peak occurs.

The only disagreement between **T1** and the measured time series of averaged AOD over **S1** occurs during 2003, which has already been noted previously by (Cohen, 2014), although none of the variables used in this study can explain why.

Over Southern Southeast Asia, there is a one-to-one agreement between the peaks in **T2** and the peaks in the averaged measurements of AOD over **S2**, with the peaks occurring in 6 years (2001, 2002, 2004, 2006, 2009, and 2012) and not occurring in the other 7 years. The measured peak in the average AOD ranges from 0.5 to 1.2, indicating that when these events occur, their impact on the aerosol loading is larger than in Northern Southeast Asia. The timing of the peaks is also wider and less well constrained than in Northern Southeast Asia, corresponding to most of the entire dry season, from early-August to the end of October. Furthermore, there is no observed second or smaller peak.

However, the issue of cloud cover leading to missed positives is observed in Southern Southeast Asia. While this method was able to pick up the high haze and pollution years of 2002, 2004, the El-Nino in 2006, and 2009, two additional high haze and pollution years of 2010 and 2013 were not captured. As already shown in (Cohen, 2014), which was capable of capturing 2010 and 2013, the likely cause is cloud cover. We have confirmed that the MODIS cloud cover is in fact the culprit, with there being fewer than 10% of pixels containing measurements of AOD over the regions given by **S2**. In fact, the only time during these years that the results are found for these years is in  $S_i$  where  $i$  is greater than 2, and thus are under the threshold used for statistical robustness. This reconfirms the afore mentioned results that MISR is in fact better at dealing with cloudiness over this region.

Careful consideration of **T1** (see Figure 2c) shows that it is considerably more noisy than **T2** (see Figure 2c), and there are three explanations for this. First is because part because the emissions from the region are more complex. In addition to the fires, there are large urban sources from three megacities: Bangkok, Ho Chi Minh City, Hanoi, as well as many highly populated and inhabited

areas outside of these cities throughout the countryside. The emissions from these cities is consistent throughout the year, and therefore the high frequency noise in these emissions, such as day/night differences, weekday/weekend differences, etc. tends to make the signal slightly more noisy. Secondly is that the fires in this region are due to combination of a few factors, which occur on different scales and have various different size holdings in each case, meaning that small differences in timing, intensity, and duration are to be expected from when the people decide to burn and how long they decide to burn for (Taylor, 2010). There is agricultural/straw burning in Thailand, subsistence burning in Cambodia, forest clearing in Myanmar and Laos, and urban and agricultural expansion in Vietnam, with some of these agricultural regions, especially related to rice, have 2 crops a year, and hence the possibility of being burned more than once (Dennis, 2005; Tipayarom and Oanh, 2007). Thirdly, the dry season here tends to be extremely dry, without even occasional rainstorms. Therefore, any emitted particles tend to have a very long lifetime. Hence, the impact of secondary chemistry is important. This chemistry tends to be very sensitive to the emissions ratios, to clouds, and to any non-linearly emitted secondary species from urban areas as the plums proceeds downwind. On the other hand, in Southern Southeast Asia, the population is also large, but in many of the places in Indonesia and Malaysia that are source regions, the cities are large and well contained, while the countryside is still relatively empty. Secondly, in this region, the major cause of burning is the clearing of primary forests, and much of this is done by a smaller number of large-land holders, further reducing the variability. This is especially so on a year-to-year basis, during some years which there is relatively little burning at all. Finally, even during the dry season, there is still a considerable amount of small scale convective precipitation and day/night sea/land breezes and rain. Hence, the lifetime of the particles and secondary precursors tends to be slightly shorter, and the impacts of non-linear secondary processing is also reduced. Hence, the fact that Southern Southeast Asia often has an even higher average AOD, means that the emissions must



be considerably larger in terms of magnitude from year to year, although not necessarily more variable within each year, as also found in (Cohen, 2014).

These results are clearly consistent with the time-averaged values of the land-use measurements of LAI and NDVI when averaged over regions **S1** and **S2** respectively (Figure 2). Over **S1**, we can clearly see that much of the region either has an average LAI which is far too low to correspond to native or secondary forest, implying that the land is now agriculture. In other cases, there is still a high average LAI value with a corresponding reduction in NDVI, implying that primary forest is being deforested in exchange for some type of commercial agricultural tree crop, such as palm oil, rubber, or wood for paper. However, the region over which this second category is occurring is smaller in size than the first region with the simultaneous decrease in both LAI and NDVI (Huete et al., 2002; Myneni et al., 2002, 2007). On the other hand, over the region **S2** we find that the LAI is still generally quite high throughout the region of interest, while the average NDVI is falling at an even faster rate than the drop over the smaller region in **S1** in which a similar type of condition is occurring. This is completely consistent with the known large-scale deforestation occurring throughout Indonesia and Malaysia where mostly primary forest is burned and replaced with large-scale agricultural tree-based crops (Dennis, 2005; Phillips et al., 2010; Taylor, 2010; Wooster et al., 2012; Field et al., 2009).

A spatial mapping of the climatological mean and standard deviations of LAI and NDVI over Southeastern Asia are displayed in Figure 34. First, it is observed that the LAI is smaller in average over Northern Southeast Asia (LAI=2.3) than over Southern Southeast Asia (LAI=3.5). Similarly for NDVI, the average value over Northern Southeast Asia is (NDVI=0.61) while it over Southern Southeast Asia it is (NDVI=0.70). This is consistent with the knowledge that in Northern Southeast Asia, the land has been more altered from its base tropical rainforest state (Natalia Hasler and Avissar, 2009; Taylor, 2010). In fact, there is a considerable amount of rice and other agriculture which has completely replaced trees with crops. Also, the pace of forest clearing is quite rapid in

those regions which still retain a considerable amount of native forest. The only considerably widespread regions of native forests are left only in Laos and at the frontier regions near the intersection of Laos, Thailand, and Myanmar.

### 3.42 Influence of Measured Fires

To look at the impacts of measured fires, we fit the relationships between LAI, NDVI, Precipitation and AOD in two cases, both with and without the inclusion of the **FireCountFire Count** variable using REG1 and REG2. This is done separately over both the Northern and Southern regions with the corresponding different thresholds. A comparison of the time series of the region averaged AOD from each EOF region, the 4 model predicted AOD values, and the measured averaged AOD is made. The average statistical error and average statistical correlation (**coefficient of determination, R<sup>2</sup>**) between the datasets and the regression-fit model predicted AOD used to determine which threshold  $\tau$  is ultimately used for the purpose of determining the best fit coefficients for  $\alpha_i$ ,  $\beta_i$ ,  $\gamma_i$ , and  $\delta_i$ . The resulting statistics are displayed in Table 1.

As expected, including the **FireCountFire Count** variable significantly increases the performance of the algorithm in terms of correlations: on average the correlation increases from 70% to 79% in the Northern region, and from 66% to 75% in the Southern region. However, there is no improvement in the mean error between the reconstructed data and the original measured AOD.

This means that ~~the existence of~~ if there is a hotspot measurement available, it will improve the ability to predict the spatial and temporal distribution of the fires, but provides no help in terms of estimating the ~~fire offers an improvement in determining the spatial and temporal timing of the~~ fires, but does not help to estimate the intensity of the AOD or ~~hence the~~ emissions. This is physically consistent, since the actual emissions should be a more complex function of the type of burning, the material burned, and the conditions under it was burned, not just the existence of a fire.

Additionally, this is consistent because the **FireCountFire Count** product only quantifies the

likelihood of a fire occurring within the given pixel, but provides no information on the intensity of the fire. Furthermore, the results of the fitting of the regression coefficient associated to

**FireCount** (Figure 4) show that the coefficient is strongly positive over the regions where fire are the most important and AOD variability the strongest (regions within the dots). Thus, the results are found to be consistent with what is understood, that **FireCount** is a reasonable predictor of emissions of aerosols from fires, but that this factor is only useful as a predictor of the effect, not as a means of understanding the magnitude of the effect.

The best fit regression coefficients associated with NDVI make more physical sense in the case where the **FireCount** predictor is used REG1 (Figure A2a) than in the case where it is not REG2 (Figure A2b). **In all cases, the timing is based on the 8-day period in the year, from day 1 to 8 being the first data point, from day 9 to 16 being the second data point, etc. In this way, multi-year variances in the climatology can be rigorously analyzed.** In general a negative coefficient is found, which implies that regions will lose NDVI as a result of an increase in AOD, which is consistent with the health of the land decreasing during a fire. A similar gain is also found in terms of the best fit coefficients for LAI in the regions which are not rice dominant (rice has a significantly low LAI so that the signal to noise ratio from the satellite product is too low to produce a statistically significant result over these regions). The regression coefficients are thus consistent and for this reason, we only refer to REG1 from this point forward.

Making comparisons between the regression constructed AOD and the measured AODMODIS over Northern Southeast Asia leads to the determination that in average, using  $\tau_{\text{North}} = P75(\text{PC1})$  as the threshold of fire activity leads to the best results, as shown in Table 1. This leads to the reasonable conclusion that in order to represent the AOD during the fire season well, there must be greater access to data, while to represent the AOD during the non-burning or low-burning seasons, that less data is required. This is consistent with the variability being considerably larger during the burning season in both space and time over the region of interest.

On the other hand, for Southern Southeast Asia, using a very small value of  $\tau_{\text{South}} = P12.5(\text{PC}2)$  gives the best statistics. This means that using less data improves the fit during the fire season as compared to the use of more data which better constrains the fit over the whole year. This is not intuitive and is only consistent with the case that either (a) the data is more likely to be of low quality during the burning season (i.e. the data is corrupted by clouds), or that there is a considerable amount of data missing during the burning season (which is also possible due to the widespread distribution of clouds over much of both Borneo and Sumatra). This view is also consistent with the year-to-year and decadal scale of variability, wherein some years will have little to no fire, and hence data is required over a considerably longer period of time, including both high- and low-fire years in order to properly reproduce the observed patterns. For the remaining of this analysis, we only consider (1) that the reconstructed data set of AOD over the Northern region has been computed by using  $\tau_{\text{North}} = P75(\text{PC}1)$  as fire threshold, and (2) that the reconstructed data set of AOD over the Southern region has been computed by using  $\tau_{\text{South}} = P12.5(\text{PC}2)$  as fire threshold. These two data sets will be referred as AOD<sub>North,REC</sub> and AOD<sub>South,REC</sub>.

### 3.23 Comparing AERONET measurements over Northern Southeast Asia

Seven stations from AERONET are situated within the Northern region (Chiang Mai, Pimai, Bac Giang, Nghia Do, Vientiane, Mukdahan, and Ubon Ratchathani) and four stations are located inside the Southern region (Jambi, Kuching, Palangkaraya, and Singapore). The location of those stations is displayed in Figure 3 and ~~complementary information is available in~~ Table D2. Of these stations, three are urban sites located downwind from burning regions: Singapore, Bac Giang and Nghia Do, while the remaining sites are located directly in or adjacent to burning areas.—

—Figures 5 and 6 displays the temporal series of the AERONET AOD (black curve) and regression-fit modeled AOD (blue curve) at the seven stations situated within the Northern region. Table 4 displays the statistics of the goodness of fit between the measured AOD and the

reconstructed AOD respectively, AODMODIS and AODNorth, in terms of reproducing the AERONET measured AOD signal. ~~These statistics are computed over both the entire time series that the respective AERONET station is measuring (Table 4) and well as only during high pollution episodes.~~

The first general observation is that all AERONET stations in Northern Southeast Asia have an annual peak in their AOD which occurs during the fire season, ~~with this peak occurring~~ from February through April each year. Additionally, each station has a smaller second peak over many of the years, but not annually, occurring ~~sometime~~ in August or September.—

————~~There are~~ At the two remote stations, ~~which are in regions which are neither urban nor in the process of urbanizing~~: Pimai (Figure 5c) and Ubon Ratchathani (Figure 5d), ~~At both stations,~~ AOD reaches its maximum value of over 0.5 during the fire season, while generally the values are ~~considerably~~ clean throughout the rest of the year, ~~as shown in Figures 5c and 5d. At these stations, the high AOD events occur every year in February to April and a except for second local maximum occurs from September to October in roughly half of the years: 2001 to 2006, and 2009, with the a second local maximum of AOD peak being of~~ around 0.46 in September and October. At Pimai, the AERONET data shows high pollution during the fire season every year from 2003 to 2008. The model captures all these events correctly in terms of duration, with the onset and end times slightly off, leading to a correlation of 43%, ~~with an intensity mean error of. However, the intensity (mean error of -0.12) (see Table 4).~~ At Ubon Ratchathani, the AERONET data shows high pollution events during the fire season of the years 2010 to 2012. The model captures all these events in terms of duration (correlation of 80%) but ~~also~~ underestimates ~~its-the~~ intensity by a slightly larger (mean error of -0.22) ~~(see Table 4)~~. A large peak of high AOD can be seen ~~on the AERONET data at Ubon Ratchathani~~ in September 2012 corresponding to a high-pollution event ~~also observed~~ in Singapore ~~(see Figure 7d)~~. This peak, which has a maximum AOD value of 0.6, is captured by the model. During the common years of data between AERONET and AODNorth (2008 to 2013), we

can see that calculate that the model captures the fire season and the pollution that is generated by it well, both in terms of duration (correlation is 64%) and less well, but not in terms of intensity (mean error of -0.26) (see Table 4). Although the error in the intensity is not insignificant, it is still significantly better than most other errors from model studies over heavily biomass burning influenced areas of the world, the mean error is still quite good, since most admit to requiring a scaling factor from 1.7 to as much as 5 (e.g.: Wu et al., 2011; Cohen and Wang, 2014; Hodnebrog et al., 2014).

There are three stations which are situated at medium-sized urban sites which are also adjacent to or directly upwind from fire burning regions: Chiang Mai, (Figure 5e6a), Mukdahan (Figure 5f6b), and Vientiane (Figure 6e5g). All It is shown that there is a strong annual peak during the fire season from February to April at these stations. At Chiang Mai and Mukdahan, which are both nearer to the the up-wind regions where the agricultural fires occur, the maximum value for of AOD is around 0.5, while it is around 0.6 at Vientiane, which is located near the downwind edge of further downwind and hence able to undergo additional secondary processing the agricultural burning regions. Figures 5e6a, 5f6b, and 5g6e also show smaller peaks during other parts of the year: from September to October for the years 2001 to 2006 at Chiang Mai, with a maximum AOD value of 0.4; from July/August and to October/November (depending on the years) for the years 2001 to 2007, 2009, and 2010 at Mukdahan, with a maximum AOD value of 0.44; and from September to October for the years 2001 to 2007, and 2009 at Vientiane, with a maximum AOD value of 0.59. The nature of these secondary peaks are not annual in occurrence, and an explanation will be explored in more detail later on. At Mukdahan, the AERONET data demonstrates the fire season peak for every year the data exists: 2004, 2006, 2007, 2008, and 2009. The regression-fit model reproduces the high pollution every year ( $R^2=0.69$ ), while also reproducing the intensity correctly in 2007 and 2009. While there is only very sparse AERONET data at Vientiane, the

regression-fit model reproduces the signal well ( $R^2=0.64$  and  $RMS=-0.07$ ) (see Table 4). Finally, the model also captures the high pollution events measured in March, April, and September 2012.

As expected, there is a considerable amount of variability at stations which are in or near large urban areas (Megacities), due to the combination of both the fire signal as well as local emissions and in-situ secondary processing. In particular, the signals at the two stations near to the rapidly growing urban megacity of Hanoi; ~~the capital of Vietnam~~, Bac Giang (Figure 5a) and Nghia Do (Figure 5b) are very similar. These stations have a much higher annual average AOD than the other stations in the region, with the daily average value as well as long-term mean measured AOD being frequently ~~in the polluted range (AOD larger than 0.4)~~, while the ~~and the~~ annual high AOD peak ~~having~~ a yearly maximum of at least 0.9 at both of these stations (see Table 3). Figures 5a and 5b also show smaller AOD peaks (maximum value of around 0.7) during other parts of the year (from July through November depending on the year). During the fire seasons in 2004 and 2007 at Bac Giang, the timing of the high pollution events are well-captured by the regression-fit model, in terms of onset, duration, and end time, although the model intensity is underestimated.

In 2006, the Southern Southeast Asian fire season produced an extensive and massive amount of emissions T2 due to extremely dry and warm conditions brought on by the El-Nino conditions. Various models and measurements have shown that the fires from these emissions have spread from S2 throughout the Indian and Pacific Oceans (Podgorny et al., 2003). However, we have also found that the signal is clearly present at all of the stations located in S1, ~~in terms of AERONET measurements as well as regression-fit models~~. At Chiang Mai, Mukdahan, and Pimai both the intensity of the 2006 season as well as its onset, duration, and conclusion are all well reproduced in both the AERONET measurements and the regression-fit model. Even at the urban megacities Bac Giang and Nghia Do the AERONET measurements also display a high pollution peak ( $AOD=1.2$ ) around September 2006, while the regression models at both of these stations capture the measured onset, duration, and ending of this event. The only issue is that the magnitude

of the regression-fit model AOD underestimates the measured value by as much as 33% at Bac

Giang. ~~Unfortunately the other AERONET stations do not have measurements available during this event.~~

Given the intimate connection between fires and the ensuing rapid changes of the land surface which occur at the same time, ~~we now explore how the as expected,~~ LAI and NDVI have changed at the same locations as the AERONET stations. First, they show a correspondingly higher value ~~in both of these variables~~ during the second, localized peak, than at the major annual peak, with a maximum value of around 0.9 at these stations (see Table 3). Figures 5a and 5b also show smaller AOD peaks (maximum value of around 0.7) during other parts of the year (July through November depending on the year), see Tables ~~D2~~ and 3. This is indicative that the second peak, which does not occur year- to-year, may not be attributed to large-scale local burning, unless ~~either~~ the local fires are much less extensive, and thus do not lead to significant change in the land surface, but happen to just be upwind of these measurement stations in these given years, or that the local fires are much more polluting per unit of land use change, and hence still contribute to the AOD to some extent. The other possible explanations are that the pollution during these times is actually transported from other place, or are intensified due to some sort of secondary processing. However, it is also found that these changes in the year-to-year LAI and NDVI values do not vary in a one-to-one manner with T2, which has some covariance during the big fire years of 2002, 2004, 2006, and 2009, but not during other years in which the peak occurs, such as 2001, 2003, 2005, and 2007.

~~Hence, we are able to conclude that the~~ Overall, we find that the annual peak in AOD ~~as measured at these AERONET stations throughout S1 -has an annual peak which~~ is clearly due to fires, and that this is true for both urban, partially urban, and remote sites. Further, during these fire events, the dominant source contributing to the peak in AOD is from the burning itself, even in ~~the~~ urban areas ~~where it may be one of two dominant sources~~. Additionally, there is a second peak found at these stations, which is both smaller in magnitude, and only occurs in certain years. This



secondary peak is very likely not due to local burning, and instead it is shown that a significant number of these years co-vary with analyzed large-scale fires from region **S2**, **indicative of long-range transport**. However, since there are a few years during which this is also not the case, it is possible that other sources of long-range transport or secondary production of aerosols, such as from South Asia, ~~could also contribute~~.

### **3.34 Comparing AERONET measurements over Southern Southeast Asia**

In Southern Southeast Asia, **S2**, the majority of the emissions come from a small number of well-defined major urban centers, transport lines through the waterways, and wide-spread sources from fires, with much of the region still continuing primary forest or dense secondary forest. As a consequence, the major source of the variation in the AOD is a combination of the emissions from fires and precipitation (as it is the major source of the aerosols removal from the atmosphere). This is demonstrated in Figure 67, demonstrating a smoother and less variable set of measurements during the wet season than at sites over Northern Southeast Asia, 5 and 6. Consequently, the AERONET site in Singapore, the sole large urban area in **S2**, is very different from the other stations of this subregion.

Unlike in Northern Southeast Asia, in general, the AOD signal in Southern Southeast Asia tends to only peak once a year (except for in 2009 and 2014, which are special cases to be discussed later that had 2 peaks due to primary fire emissions). This primary peak, as shown in **T2** always occurs during the local fire season from August through October/November, without any additional second peak occurring during a non-burning period, as in **T1**. Effectively, this implies that emissions from **S1** are not contributing to the variance in the measured AOD over **S2** and that long-range transport from Northern Southeast Asia is not efficient in contributing to the high peaks in AOD found over **S2**.

Additionally, Southern Southeast Asia has an important source of uncertainty and bias in the

measurements over the region. Specifically, the impact of intense cloud cover is also determined to be very important, in terms of being able to capture all of the known large-scale fire based events. We observe that in a few special cases where known large-scale pollution events have occurred over S2 as measured both on the ground and by MISR measurements of AOD (Cohen, 2014), that MODIS was not able to successfully capture the events (for example: June 2013). A careful examination of the cloud cover fields and ~~FireCount~~ Fire Count measurements show that this is clearly the case, at least for June 2013; the region S2 was almost completely masked by clouds (over 80% of all pixels) in the day-to-day tracks, with more than 90% of pixels in the 8-day average fields over this period of time being masked.

The AERONET station in Singapore, is located in a highly urban environment, with sizable sources of aerosol emissions related to shipping, a high energy using population, and refineries. It is clear that there are no wild-fires occurring within Singapore. At the Singapore station, we observe an annual signal except for every year, although during the years 2008 and 2010, the signal is less intense than in the other measured years. There is a considerable amount of variation in the magnitude, the onset, and the duration of the peak, as well as a considerable amount of noise. However, the maximum measured AOD here on an 8-day average basis, ranges from a low year of 0.55 to a high year of 0.81 in 2006. Even though the fires were quite distant, it is clearly observed that then most intense event in 2006 is readily captured here, further supporting that even in an urban environment, Singapore offers a reasonable downwind signal site for observing the impacts of the fires.

On the other hand, the other AERONET stations in this region, including in Kuching, Jambi, and Palangkaraya, are situated in remote and mostly heavily jungle/forested regions of Borneo and Sumatra islands (see Table D-2). These sites are all located close-by to where the fire sources originate, in the jungles and forests of Borneo and Sumatra. The AERONET station in Jambi, situated on Sumatra Island, has an annual signal of high AOD occurring once a year, every year,

except in 2010 (where there were no ~~corresponding~~ measurements ~~during the peak season~~) ~~as given by Figure 10~~). However, the magnitude, onset, and duration of these high pollution events is highly variable from year to year. The AOD maximum value ranges from a low of 0.67 (in 2007) to a high of 1.49 (in 2006) (see Table 5). The AERONET station in Kuching, ~~is situated Northern Borneo, in Malaysia, also~~ has an annual peak signal in AOD every year that measurements are available (there were no measurements during the corresponding peak times in 2008, 2010, and 2013). The magnitude, start, and duration of this peak is again highly variable from year to year, with the maximum in measured AOD ranging from a low of 0.68 in 2007 to a maximum of 1.36 in 2006. At Palangkaraya, which is situated in Western Borneo in Indonesia, there is also a single high peak occurring every year, except for 2010 (which ~~again~~ did not have any measurements) ~~during the high fire season~~). Similar to the other stations, the intensity, onset, and duration of the high AOD signal was very variable from year to year.

The regressive-fit model based on the MODIS measurements at each of the remote sites in Southern Southeast Asia: Jambi, Kuching, and Palangkaraya, is capable of reproducing the major heavily polluted years as found in the measurements, such 2002 (max AOD of 1.24 in Jambi, 1.0 in Kuching, and 1.94 in Palangkaraya), 2004 (max AOD of 0.99 in Jambi, 0.85 in Kuching, and 1.18 in Palangkaraya), 2006 (max AOD of 1.49 in Jambi, 1.4 in Kuching, and 1.98 in Palangkaraya), and 2009 (max AOD of 0.95 in Jambi, 0.87 in Kuching, and 1.02 in Palangkaraya). At Jambi and Palangkaraya, the regressive-fit model reproduces the high AOD event of late 2012 well, with a better correlation with the AERONET measurements ( $R^2=76\%$  at Jambi and  $R^2=74\%$  at Palangkaraya) than MODIS AOD at the same grid point ( $R^2=51\%$  at Jambi and  $R^2=71\%$  at Palangkaraya), as given in Table 6, although the intensity in these years is slightly low. On the other hand, the regressive-fit model reproduces the AOD well in terms of intensity, onset, and duration at Kuching (RMS error of 0.13,  $R^2=66\%$ ) (see Table 6). However, the regressive-fit model is still basically constrained by the cloud cover issue. It is for this reason that the know high values of

aerosols in the atmosphere over Singapore in June of 2013 (as based on surface measurements and personal observation) is not captured in AERONET measurements, MODIS measurements, or the regressive-fit model. In addition to June 2013, we also find that MODIS AOD and the regressive fit model are both not capable of capturing the 2010 fire season peak either. However, the issues of cloud cover seem to be less important in other years, and we find the onset, duration, and intensity are all well matched between the regressive-fit model and AERONET measurements at Singapore during the fire seasons of the years 2007, 2008, 2009, 2011, and 2012 (see Table 6 for statistics).

### 3.45 Comparisons versus measurements from the MISR satellite

MISR satellite measurements of AOD are at lower spatial and temporal resolution than MODIS and AERONET measurements, and thus to use them as a basis for comparison, the values from MODIS and AERONET will be averaged to a monthly-basis as well as at  $0.5^{\circ} \times 0.5^{\circ}$ . Over Northern Southeast Asia, the time series of the regression-fit model AOD compares very well with the time series of the average MISR AOD over the same region ( $R^2=0.77$  over all of S1, and  $R^2=0.85$  over the region of highest variability). While there is some underestimation of the absolute AOD as compared to the MISR measurements, that underestimation is always less than 0.1, and therefore is not far from the order of magnitude of the error in the measurements themselves. One of the important reasons why the agreement is so good is that this region is generally cloud-free during the dry season when the fires occur, and hence there is a quite large and representatively similar sampling size between MODIS, MISR, and AERONET during the fire periods in this region. This establishes that indeed the MODIS based regression-fit model matches well against MISR, and is able to reproduce the variability and magnitude of the AOD over Northern Southeast Asia (Figure 7).

Not surprisingly, when fitting the results of the MODIS regression-fit model using 8-day average data, the overall fits are less good when comparing against MISR. Part of the issue is the

additional variability, but more importantly is the lack of sufficient data due to cloud coverage. Specifically, over the region S1, the correlation rises from  $R^2=0.66$  to  $R^2=0.81$  when increasing from 8-day to monthly averaging. Similarly, the comparison between the AERONET data and MISR AOD also increases from  $R^2=0.59$  to  $R^2=0.79$  when comparing 8-day averages and monthly averages respectively. Overall, the regression-fit model is able to reproduce the variation of AOD at all the stations in Northern Southeast Asia, both in terms of duration and intensity concerning high pollution events (see Figures B3 and B4).

As expected, the spatial comparison between MISR and the regression-fit model over Southern Southeast Asia is less good. The first thing to note is that the spatial extent of the region from MODIS, given with the relatively level of high certainty by S2, is considerably smaller than a similar spatial distribution of the smoke extent over this same region, when analyzed in the same way using data from MISR measurements (Cohen, 2014). This is explained in part due to the larger cloud-covered fraction in the MODIS measurements when compared with MISR, as well as the shorter averaging period with the MODIS measurements, leading to a situation where there is insufficient information at each averaging time step over much of the region. It is found that the RMS error between MISR and the regression-fit model ranges from a minor and relatively insignificant (as compared to the measurement errors) model overestimate of 0.1 in AOD, to a substantial and significant model underestimate in the AOD of up to 0.5. This regression-fit model underestimates as compared to MISR measurements is significantly larger than the AERONET and MISR disagreement over this region, which is less than 0.3 (Cohen, 2014; Shi et al., 2011) and further, this error occurs especially and exclusively during the intense fire-burning years. On the other hand, the overall temporal correlation between the regression-fit model and the time-average AOD from MISR is  $R^2=0.72$  over all and is as high as  $R^2=0.79$  over the region of highest AOD variability. This means that the inter-annual and intra-annual variation is relatively captured by the MODIS measurements and the resulting regression-fit model.

#### 4. Conclusions

An in-depth analysis of multiple measurements from MODIS, MISR, TRMM, and AERONET measurements has been performed over a 13-year period over Southeast Asia. Using MODIS AOD, the spatial and temporal patterns of the contribution of fires to the atmospheric loading of aerosols was established. Two distinct regions, with vastly different properties were observed: one in Northern Southeast Asia, which had a strong annual signal with some inter-annual variability, and another in Southern Southeast Asia, which had a strong signal with inter-annual and intra-annual variability. Northern Southeast Asia shows an annual high AOD during the fire season (varying roughly from February through April), with a smaller nearly annual peak occurring during the exact timing when Southern Southeast Asia has its fire season. Southern Southeast Asia is affected every year by their own fires (from roughly August through October), without any observed secondary peak except for during two exceptionally dry years during the second very short dry season in February 2009 and the very end of 2013 (which would up maximizing in February 2014, although it is beyond the end of the data analyzed in this paper). The representation in terms of the timing of the fires of Northern Southeast Asia was consistently good in terms of start time, length of the burning season, cessation of the burning, when compared against AERONET and MISR measurements. The representation in terms of timing over Southern Southeast Asia was not as good, but still quite acceptable when compared against AERONET and MISR measurements, with the duration of the fire season well captured in strong fire years, and the strongest part of the fire season captured in low fire years.

Bringing in different simultaneous measurements of land-surface variables, fires, precipitation, and column aerosol measurements, allows us to confirm that these patterns exist and are consistent with land-use burning. Given the difference in the timing and durations of the major monsoon seasons over these regions, the results are consistent. From this point, a simple regression-

fit model was established to predict the AOD from measurements of land-use change variables, fires, and precipitation, which should be the basis upon which fires start in the environment. These simple regression-fit models (based on MODIS and TRMM) reproduced the onset, duration, and magnitude of the measured AOD from other measured sources (MISR and AERONET) well over Northern Southeast Asia. The results of this regression-fit model demonstrate the ability to predict the AOD as observed by AERONET and MISR, using only measurements of land-use change variables and fires from MODIS, and precipitation from TRMM, measurements of some of the important and fundamental underlying factors controlling the fires.

These simple regression-fit models reproduced the onset, duration, cessation, and even the magnitude of the measured AOD from AERONET and MISR very well in Northern Southeast Asia. These simple regression-fit models also reproduced the onset, duration, and cessation, of the measured AOD from AERONET and MISR well well in Southern Southeast Asia, especially during the more intense burning years. The main issue in Southern Southeast Asia, however, was that the magnitude over this region was strongly underestimated. **These results still underestimate the column loading, but by a magnitude of 30% or less, which is far better than the typical scaling factors applied of 1.7 (70%) or more, and consistent with the results in Cohen and Wang (2014) and Cohen (2014) which show that there is an underestimate in both the overall magnitude as well as in the fire magnitude, and that correcting for the former leads to an underestimate in the latter of 20% to 30%, The result is not only larger in magnitude than the GFED emissions products, but include regions which are considered to have zero emissions in the GFED data set, a worrying conclusion, since a value of 0 cannot be scaled up by a scaling factor.** Some reasons for this include emissions sources which are more variable in space and time, such as the clearing of primary forests, peat burning, and rapid development; and other limiting reasons such as increased cloud cover reducing the number of available measurements over large portions of this region by a significant amount. Further, the inter-seasonal periods in Southern Southeast Asia tend to be both more rainy and more

cloud-covered than in Northern Southeast Asia, due to large scale convection and other regional disturbances like the MJO and the IOD.

There is a strong and consistent change in the land use variables occurring during the local fire season over both Northern and Southern Southeast Asia, although these relationships, as expected, are different over the two regions due to different types of land-use change. The relationships between burning of primary forests, grasslands or crops, and peat should all be different. Additionally, there is an important secondary use for these relationships, determining whether the observed smoke is locally produced or transported from far upwind. For example, it is clearly noted that the land-use changes are much smaller during the second non-annually occurring peak in Northern Southeast Asia, implying that while there may be some contribution from local sources, that there is also a large amount of smoke which is transported from other regions. This comes from the idea that if the land itself did not change very much, then the emissions of smoke produced must have been considerably lower. The timing of this smaller peak matches the timing of the fire occurrence over Southern Southeast Asia with a very high level of correlation. Additionally, it also cannot be ruled out that the smoke could be urban pollution from South Asia. On the other hand, there is no evidence that any of the smoke in Southern Southeast Asia originates from any region other than its own sources.

Further, we explored the added value of using higher temporal resolution data, which is usually thought to add improved value. Due to the large amount of cloudiness encountered, there was a much reduced number of measurements available over Southern Southeast Asia during the fire season using 1-day average values as compared to 8-day average values, leading to less statistical relevance. In the end, it was not possible to have a reasonable reproduction of the measured AERONET and MISR values of the onset, duration, and ending of the fires using 1-day average MODIS and TRMM data as compared to when using 8-day average MODIS and TRMM measurements to develop the regression-fit relationships. Even with the 8-day average data and the



associated regression-fit relationships, the magnitude of AOD during Southern Southeast Asia's fire season is significantly too low, although in Northern Southeast Asia, it is low but not more than the magnitude of the uncertainty of the input measurements themselves. The correlation between the regression-fit model AOD and AERONET stations over the entire decadal time period, using 8-day average MODIS data, ranges from  $R^2=0.42$  to  $R^2=0.75$ . While monthly-average data from MODIS does not provide as fine resolution for the duration, onset, and end times of the fires, it provides the best match in terms of the magnitude of the AOD measurements from AERONET and MISR. However, when using MODIS data on a monthly average basis, the regression-fit model AOD gives a better performance with the correlation coefficient between AOD and AERONET stations ranging from  $R^2=0.70$  to  $R^2=0.90$ . Furthermore, the correlation over the regions of interest S1 and S2 between the regression-fit model and MISR measurements of AOD ranges from  $R^2=0.57$  to  $R^2=0.81$ . This is due partially to less under-representation of very high short-term peaks, as well as additional data points being available in the MODIS fire and land use products at longer average time durations. This is a counter-intuitive result, with many in the community stressing the added value of higher frequency measurements, but one which is consistent with the fact that such space-born measurements are severely limited by clouds over this region of the world during the fire season. MISR has shown to represent the magnitude of the AOD well, with the measurements from monthly-average MISR measurements and monthly-average AERONET measurements being basically the same. Therefore, the ability of the regression-fit model to capture the monthly-average AOD from both MISR and AERONET, in terms of both the inter-annual and intra-annual variability in the fire seasons, is significant, and shows that indeed the changes in the land surface and the impacts of precipitation are what are driving the atmospheric loading of AOD and hence the impact of the fires over this region on the decadal scale. Further, as it is widely known, peat can burn and smolder for an extended period of time after any measured fire has gone away, and therefore, by extending the average value for the fire, it allows for a better matching with the total emissions,

which will continue to often be produced for weeks after any visible flame or surface heat is observed. Thusly, one of the important findings is to examine the most ideal temporal resolution at which to use the data, whether it be daily, weekly, or monthly. While most of the published literature leans towards using high frequency daily data (or individual swath-by-swath data, where available), we determine and validate that using weekly or monthly average data leads to a better ability to accurately reproduce the measured values, explain why that is the case, and then quantify some of the impacts and limitations of this result.

This study highlights the importance of taking into account land-use variable and precipitation for estimating AOD correctly both in time and magnitude, even if magnitude remains hard to capture on a 8-day basis. One significant bias in the magnitude of the results must be due to problems of the relationships over the region being not properly captured, such as the different anthropogenic driving forces of the land-clearing being significantly different over the two regions. A second significant bias in the magnitude is due to the fact that there is a significantly more cloud cover over the two regions during their local burning seasons(Giglio et al., 2003). These results support the efficacy of the approach introduced here: that it is appropriate to use measured changes in the land, precipitation, and active fires from MODIS and TRMM to reproduce a working model of the atmospheric aerosol loading. Furthermore, other than (Cohen, 2014; Cohen and Wang, 2014) there are no other works that have been able to satisfactorily estimate the loadings of or AOD associated with emissions aerosols over this region of the world, without using some type of scaling. This method is able to reproduce the magnitudes by introducing physical parameterizations of scaling, and doing so based on a more fundamental driver-based approach. This allows us to improve our understanding of the relationships, both in terms of how they vary over space and time, on one hand, and in terms of physical drivers, on the other.

## Acknowledgements

This work was supported by the School of Atmospheric Sciences, Sun Yat-Sen University,

~~ACCT#XXXXX,~~

the Singapore National Research Foundation (NRF) through a grant to the Center for Environmental Sensing and Monitoring (CENSAM) of the Singapore-MIT Alliance for Research and Technology (SMART), and the Tropical Marine Sciences Institute. The authors also want to thank all of the Principal Investigators of MISR, AERONET, NOAA, and MODIS, and TRMM for making the data available.

## References

- Afroz, R., Hassan, M. N., and Ibrahim, N. A.: Review of air pollution and health impacts in Malaysia, *Environmental research*, 92, 71–77, 2003.
- Aldrian, E., Gates, L. D., and Widodo, F.: Seasonal variability of Indonesian rainfall in ECHAM4 simulations and in the reanalyses: The role of ENSO, *Theoretical and Applied Climatology*, 87, 41–59, 2007.
- Asner, G. P., Scurlock, J. M., and A Hicke, J.: Global synthesis of leaf area index observations: implications for ecological and remote sensing studies, *Global Ecology and Biogeography*, 12, 191–205, 2003.
- Bjornsson, H. and Venegas, S.: A Manual for EOF and SVD Analyses of Climate Data. Department of Atmospheric and Oceanic Sciences and Centre for Climate and Global Change Research, Tech. rep., McGill University, Technical Report, 1997.
- Chang, C., Wang, Z., McBride, J., and Liu, C.-H.: Annual cycle of Southeast Asia-Maritime Continent rainfall and the asymmetric monsoon transition, *Journal of climate*, 18, 287–301, 2005.
- Chen, W., Wang, X., Cohen, J. B., Zhou, S., Zhang, Z., Chang, M., and Chan, C.-Y.: Properties of aerosols and formation mechanisms over southern China during the monsoon season, *Atmos. Chem. Phys.*, 16, 13271–13289, doi:10.5194/acp-16-13271-2016, 2016.
- Cohen, J. B.: Quantifying the occurrence and magnitude of the Southeast Asian fire climatology, *Environmental Research Letters*, 9, 114018, 2014.
- Cohen, J. B. and Prinn, R. G.: Development of a fast, urban chemistry metamodel for inclusion in global models, *Atmospheric Chemistry and Physics*, 11, 7629–7656, 2011.
- Cohen, J. B. and Wang, C.: Estimating global black carbon emissions using a top-down Kalman Filter approach, *Journal of Geophysical Research: Atmospheres*, 119, 307–323, 2014.
- Cohen, J. B., Prinn, R. G., and Wang, C.: The impact of detailed urban-scale processing on the composition, distribution, and radiative forcing of anthropogenic aerosols, *Geophysical Research Letters*, 38, 2011.
- Cuevas, G. M., Gerard, F., Balzter, H., Riano, D.: Analysing forest recovery after wildfire disturbance in boreal Siberia using remotely sensed vegetation indices, *Climate Change Biology*, Nov 3, 2008, 10.1111/j.1365-2486.2008.01784.x.
- Dennis, R. A., Grahame, J.M., Unna, A., Chokkalingam, U., Pierce, C.J., Kurniawan, I., Lachowski, H., Maus, P., Permana, R.P., Ruchiat, Y., Stolle, F., Suyanto, Tomich, T.P.: Fire, People and Pixels: Linking Social Science and Remote Sensing to Understand Underlying Causes and Impacts of Fires in Indonesia, *Human Ecology*, 33, 465–504, doi:10.1007/s10745-005-5156-z, <http://dx.doi.org/10.1007/s10745-005-5156-z>, 2005.
- Diner D J et al 1998 Multi-angle imaging spectroradiometer (MISR) instrument description and experiment overview *IEEE Trans. Geosci. Remote Sens.* 36 1072–87–
- Dubovik, O., Smirnov, A., Holben, B., King, M., Kaufman, Y., Eck, T., and Slutsker, I.: Accuracy assessments of aerosol optical properties retrieved from Aerosol Robotic Network (AERONET) Sun and sky radiance measurements, *Journal of Geophysical Research: Atmospheres* (1984–2012), 105, 9791–9806, 2000.
- Field, R. D., van der Werf, G. R., and Shen, S. S.: Human amplification of drought-induced biomass burning in Indonesia since 1960, *Nature Geoscience*, 2, 185–188, 2009.
- Fisher, J. B., Malhi, Y., Bonal, D., Da Rocha, H. R., De Araujo, A. C., Gamo, M., Goulden, M. L., Hirano, T., Huete, A. R., Kondo, H., et al.: The land–atmosphere water flux in the tropics, *Global Change Biology*, 15, 2694–2714, 2009.

- Fu J S, Hsu N C, Gao Y, Huang K, Li C, Lin N-H and Tsay S-C 2012 Evaluating the influences of biomass burning during 2006 BASE-ASIA: a regional chemical transport modeling *Atmos. Chem. Phys.* 12 3837–55
- Fuller, D. and Murphy, K.: The Enso-Fire Dynamic in Insular Southeast Asia, *Climatic Change*, 74, 435–455, doi:10.1007/s10584-006-0432-5, <http://dx.doi.org/10.1007/s10584-006-0432-5>, 2006.
- Giglio, L., Descloitres, J., Justice, C. O., and Kaufman, Y. J.: An enhanced contextual fire detection algorithm for MODIS, *Remote sensing of environment*, 87, 273–282, 2003.
- Giglio, L., Csiszar, I., and Justice, C. O.: Global distribution and seasonality of active fires as observed with the Terra and Aqua Moderate Resolution Imaging Spectroradiometer (MODIS) sensors, *Journal of Geophysical Research: Biogeosciences* (2005–2012), 111, 2006.
- Hansen, M. C., Stehman, S. V., Potapov, P. V., Loveland, T. R., Townshend, J. R., DeFries, R. S., Pittman, K. W., Arunarwati, B., Stolle, F., Steininger, M. K., et al.: Humid tropical forest clearing from 2000 to 2005 quantified by using multitemporal and multiresolution remotely sensed data, *Proceedings of the National Academy of Sciences*, 105, 9439–9444, 2008.
- Hodnebrog, O., Myhre, G., and Samset, B: How shorter black carbon lifetime alters its climate effect. *Nature Communications* 5, Article number: 5065 (2014) doi:10.1038/ncomms6065.–
- Holben, B., Eck, T., Slutsker, I., Tanre, D., Buis, J., Setzer, A., Vermote, E., Reagan, J., Kaufman, Y., Nakajima, T., et al.: AERONET: A federated instrument network and data archive for aerosol characterization, *Remote sensing of environment*, 66, 1–16, 1998.
- Huang, J., Minnis, P., Lin, B., Wang, T., Yi, Y., Hu, Y., Sun-Mack, S., and Ayers, K.: Possible influences of Asian dust aerosols on cloud properties and radiative forcing observed from MODIS and CERES, *Geophysical Research Letters*, 33, 2006.
- Huete, A., Justice, C., and Van Leeuwen, W.: MODIS vegetation index (MOD13), Algorithm theoretical basis document, 3, 213, 1999.
- Huete, A., Didan, K., Miura, T., Rodriguez, E. P., Gao, X., and Ferreira, L. G.: Overview of the radiometric and biophysical performance of the MODIS vegetation indices, *Remote sensing of environment*, 83, 195–213, 2002.
- Jacobson, M. Z.: Strong radiative heating due to the mixing state of black carbon in atmospheric aerosols, *Nature*, 409, 695–697, 2001.
- Kahn R. A., B. J. Gaitley, M. J. Garay, D. J. Diner, T. F. Eck, A. Smirnov, and B. N. Holben (2010), Multiangle Imaging SpectroRadiometer global aerosol product assessment by comparison with the Aerosol Robotic Network, *J. Geophys. Res.*, 115, D23209, doi:10.1029/2010JD014601.
- Langmann, B., Duncan, B., Textor, C., Trentmann, J., and van der Werf, G. R.: Vegetation fire emissions and their impact on air pollution and climate, *Atmospheric Environment*, 43, 107–116, 2009.
- Levy, R. C., Mattoo, S., Munchak, L. A., Remer, L. A., Sayer, A. M., Patadia, F., and Hsu, N. C.: The Collection 6 MODIS aerosol products over land and ocean, *Atmos. Meas. Tech.*, 6, 2989–3034, doi:10.5194/amt-6-2989-2013, 2013.
- Lin, C.-Y., Hsu, H.-m., Lee, Y. H., Kuo, C. H., Sheng, Y.-F., and Chu, D. A.: A new transport mechanism of biomass burning from Indochina as identified by modeling studies, *Atmos. Chem. Phys.*, 9, 7901–7911, doi:10.5194/acp-9-7901-2009, 2009.
- Lin, C.-Y., Zhao, C, Xiaohong, L., Lin, N.-H., and Chen, W.: Modelling of Long-Range Transport of Southeast Asia Biomass-Burning Aerosols to Taiwan and their Radiative Forcings over East Asia, *Tellus B*, 2014, 66, 23733, <http://dx.doi.org/10.3402/tellusb.v66.23733>.
- Miettinen, J., Hyer, E., Chia, A. S., Kwoh, L. K., and Liew, S. C.: Detection of vegetation fires and burnt areas by remote sensing in insular Southeast Asian conditions: current status of knowledge and future challenges, *International Journal of Remote Sensing*, 34, 4344–4366, 2013.

- Ming, Y., Ramaswamy, V., and Persad, G.: Two opposing effects of absorbing aerosols on global-mean precipitation, *Geophysical Research Letters*, 37, 2010.
- Morawitz, D. F., Blewett, T. M., Cohen, A., and Alberti, M.: Using NDVI to Assess Vegetative Land Cover Change In Central Puget Sound, *Environmental Monitoring and Assessment* (2006) 114: 85–106.
- Myneni, R., Hoffman, S., Knyazikhin, Y., Privette, J., Glassy, J., Tian, Y., Wang, Y., Song, X., Zhang, Y., Smith, G., et al.: Global products of vegetation leaf area and fraction absorbed PAR from year one of MODIS data, *Remote sensing of environment*, 83, 214–231, 2002.
- Myneni, R. B., Yang, W., Nemani, R. R., Huete, A. R., Dickinson, R. E., Knyazikhin, Y., Didan, K., Fu, R., Juárez, R. I. N., Saatchi, S. S., et al.: Large seasonal swings in leaf area of Amazon rainforests, *Proceedings of the National Academy of Sciences*, 104, 4820–4823, 2007.
- Nakajima, T., Higurashi, A., Takeuchi, N., and Herman, J. R.: Satellite and ground-based study of optical properties of 1997 Indonesian Forest Fire aerosols, *Geophysical Research Letters*, 26, 2421–2424, 1999.
- Natalia Hasler, D. W. and Avissar, R.: Effects of tropical deforestation on global hydroclimate: a multimodel ensemble analysis, *J. Climate*, 22, 1124–1141, doi:10.1175/2008JCLI2157.1, 2009.
- Neale, R. and Slingo, J.: The maritime continent and its role in the global climate: A GCM study, *Journal of Climate*, 16, 834–848, 2003.
- Nightingale, J., Nickeson, J., Justice, C., Baret, F., Garrigues, S., Wolfe, R., et al.: Global validation of EOS land products, lessons learned and future challenges: A MODIS case study, in: Available from. *Proceedings of 33rd International Symposium on Remote Sensing of Environment: Sustaining the Millennium Development Goals*, Stresa, Italy. [http://landval.gsfc.nasa.gov/pdf/ISRSE\\_Nightingale.pdf](http://landval.gsfc.nasa.gov/pdf/ISRSE_Nightingale.pdf), p. 4, 2008.
- Hope, A., Tague, C., and Clark, R.: Characterizing post-fire vegetation recovery of California chaparral using TM/ETM+ time-series data: *International Journal of Remote Sensing* 28, 6, 2007. <http://dx.doi.org/10.1080/01431160600908924>.
- Petrenko, M. and Ichoku, C.: Coherent uncertainty analysis of aerosol measurements from multiple satellite sensors, *Atmos. Chem. Phys.*, 13, 6777–6805, doi:10.5194/acp-13-6777-2013, 2013.
- Petrenko, M., Kahn, R., Chin, M., Soja, A., Kucsera, T., et al.: The use of satellite-measured aerosol optical depth to constrain biomass burning emissions source strength in the global model GOCART, *Journal of Geophysical Research: Atmospheres* (1984–2012), 117, 2012.
- Phillips, O. L., Van der Heijden, G., Lewis, S. L., López-González, G., Aragão, L. E., Lloyd, J., Malhi, Y., Monteagudo, A., Almeida, S., Dávila, E. A., et al.: Drought–mortality relationships for tropical forests, *New Phytologist*, 187, 631–646, 2010.
- Podgorny, I., Li, F., and Ramanathan, V.: Large aerosol radiative forcing due to the 1997 Indonesian forest fire, *Geophysical Research Letters*, 30, 2003.
- Ramanathan, V. and Carmichael, G.: Global and regional climate changes due to black carbon, *Nature geoscience*, 1, 221–227, 2008.
- Reid, J. S., Hyer, E. J., Johnson, R. S., Holben, B. N., Yokelson, R. J., Zhang, J., Campbell, J. R., Christopher, S. A., Di Girolamo, L., Giglio, L., et al.: Observing and understanding the Southeast Asian aerosol system by remote sensing: An initial review and analysis for the Seven Southeast Asian Studies (7SEAS) program, *Atmospheric Research*, 122, 403–468, 2013.
- Remer, L. A., Kaufman, Y., Tanré, D., Mattoo, S., Chu, D., Martins, J. V., Li, R. R., Ichoku, C., Levy, R., Kleidman, R., et al.: The MODIS aerosol algorithm, products, and validation, *Journal of the atmospheric sciences*, 62, 947–973, 2005.
- Remer, L. A., Mattoo, S., Levy, R. C., and Munchak, L. A.: MODIS 3 km aerosol product: algorithm and global perspective, *Atmospheric Measurement Techniques*, 6, 1829–1844, doi:10.5194/amt-6-1829-2013, <http://www.atmos-meas-tech.net/6/1829/2013/>, 2013.

- Rosenfeld, D.: TRMM observed first direct evidence of smoke from forest fires inhibiting rainfall, *Geophysical Research Letters*, 26, 3105–3108, 1999.
- Saatchi, S., Asefi-Najafabady, S., Malhi, Y., Aragao, L., Anderson, L., Myneni, R., and Nemani, R.: Persistent effects of a severe drought on Amazonian forest canopy, *PNAS*, January 8, 2013, 110, 2, p565–570.
- Sayer, A. M., Smirnov, A., Hsu, N. C., Munchak, L. A., and Holben, B. N.: Estimating marine aerosol particle volume and number from Maritime Aerosol Network data, *Atmos. Chem. Phys.*, 12, 8889–8909, doi:10.5194/acp-12-8889-2012, 2012.
- Shi, Y., Zhang, J., Reid, J. S., Hyer, E. J., Eck, T. F., Holben, B. N., and Kahn, R. A.: A critical examination of spatial biases between MODIS and MISR aerosol products; application for potential AERONET deployment, *Atmospheric Measurement Techniques*, 4, 2823–2836, doi: 10.5194/amt-4-2823-2011, <http://www.atmos-meas-tech.net/4/2823/2011/>, 2011.
- Tao, W.-K., Chen, J.-P., Li, Z., Wang, C., and Zhang, C.: Impact of aerosols on convective clouds and precipitation, *Reviews of Geophysics*, 50, 2012.
- Taylor, D.: Biomass burning, humans and climate change in Southeast Asia, *Biodiversity and Conservation*, 19, 1025–1042, doi:10.1007/s10531-009-9756-6, <http://dx.doi.org/10.1007/s10531-009-9756-6>, 2010.
- Tipayarom, D. and Oanh, N. K.: Effects from open rice straw burning emission on air quality in the Bangkok Metropolitan Region, *Science Asia*, 33, 339–345, 2007.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S., and Arellano Jr, A. F.: Interannual variability in global biomass burning emissions from 1997 to 2004, *Atmospheric Chemistry and Physics*, 6, 3423–3441, 2006.
- van der Werf, G. R., Dempewolf, J., Trigg, S. N., Randerson, J. T., Kasibhatla, P. S., Giglio, L., Murdiyarso, D., Peters, W., Morton, D. C., Collatz, G. J., Dolman, A. J., and DeFries, R. S.: Climate regulation of fire emissions and deforestation in equatorial Asia, *Proceedings of the National Academy of Sciences*, 105, 20350–20355, doi:10.1073/pnas.0803375105, <http://www.pnas.org/content/105/51/20350.abstract>, 2008.
- Wang, C.: Impact of direct radiative forcing of black carbon aerosols on tropical convective precipitation, *Geophysical research letters*, 34, 2007.
- Wang, C.: Impact of anthropogenic absorbing aerosols on clouds and precipitation: A review of recent progresses, *Atmospheric Research*, 122, 237–249, 2013.
- Wohl, E., Barros, A., Brunzell, N., Chappell, N. A., Coe, M., Giambelluca, T., Goldsmith, S., Harmon, R., Hendrickx, J. M., Juvik, J., et al.: The hydrology of the humid tropics, *Nature Climate Change*, 2, 655–662, 2012.
- Wooster, M. J., Perry, G. L. W., and Zoumas, A.: Fire, drought and El Niño relationships on Borneo (Southeast Asia) in the pre-MODIS era (1980–2000), *Biogeosciences*, 9, 317–340, doi:10.5194/bg-9-317-2012, <http://www.biogeosciences.net/9/317/2012/>, 2012.
- Wu, L., H. Su, and J. H. Jiang (2011), *Regional simulations of deep convection and biomass burning over South America: 1. Model evaluations using multiple satellite data sets*, *J. Geophys. Res.*, 116, D17208, doi:10.1029/2011JD016105.–
- Yang, W., Tan, B., Huang, D., Rautiainen, M., Shabanov, N. V., Wang, Y., Privette, J. L., Huemmrich, K. F., Fensholt, R., Sandholt, I., et al.: MODIS leafareaindex products: From validation to algorithm improvement, *Geoscience and Remote Sensing, IEEE Transactions on*, 44, 1885–1898, 2006.
- Zhou S.Z., Perry K.D., Wang, X., Cohen, J.B., Liang, J., Huang, M., Fan, Q., Chen, W., Chang, M., Ancelet, T., and Trompeter, W.J.: High time-resolved elemental components in fine and coarse particles in the Pearl River Delta region of Southern China: Dynamic variations and effects of

meteorology. Science of the Total Environment, <http://dx.doi.org/10.1016/j.scitotenv.2016.05.194> ,2016.



**Table 1.** Average error and correlation between AOD at 0.55 microns from MODIS and reconstructed AOD with different thresholds :  $\tau = P90(PC)$ ,  $\tau = P87.5(PC)$ ,  $\tau = P83.5(PC)$ , and  $\tau = P75(PC)$  on the Northern and Southern regions, obtained by using REG1 and REG2.

Stations	AOD <sub>MODIS</sub>	AOD <sub>North</sub>	AOD <sub>South</sub>
	Err/Corr (%)	Err/Corr (%)	
Chiang Mai All (218/598)	-0.1/83	-0.1/75	
Bac Giang All (154/598)	-0.03/74	-0.06/42	
Mukdahan All (238/598)	-0.01/79	-0.01/69	
Nghia Do All (79/598)	-0.12/74	-0.14/42	
Pimai All (120/598)	0.0/77	-0.01/57	
Ubon Ratchathani All (99/598)	-0.01/88	-0.02/61	
Vientiane All (36/598)	-0.08/83	-0.07/64	
Chiang Mai Fire (62/151)	-0.26/91	-0.26/64	
Bac Giang Fire (46/151)	-0.07/75	-0.24/33	
Mukdahan Fire (74/151)	-0.08/86	-0.15/49	
Nghia Do Fire (15/151)	-0.08/75	-0.5/62	
Pimai Fire (45/151)	-0.03/75	-0.12/43	
Ubon Ratchathani Fire (23/151)	-0.07/88	-0.22/80	
Vientiane Fire (8/151)	-0.14/93	-0.33/92	
Jambi All (64/598)	-0.12/51		-0.26/76
Kuching All (91/598)	0.06/75		0.13/66
Palangkaraya All (65/598)	-0.11/71		-0.11/74
Singapore All (279/598)	-0.02/29		0.01/44
Jambi Fire (6/74)	-0.51/80		-0.54/71
Kuching Fire (10/74)	-0.28/80		-0.03/-9
Palangkaraya Fire (6/74)	-0.5/85		-0.45/31
Singapore Fire (24/74)	-0.23/-21		-0.09/8

~~**Table 2.** Complementary information on AERONET stations: geographical location, data availability, average ( $\mu$ ) and standard deviation ( $\sigma$ ) values for AOD, LAI, and NDVI from MODIS, and environment description.~~

**Table 2.** Statistics over the respective Northern and Southern regions compared to the AERONET stations. Overlapped periods between the reconstructed AOD AODNorth/South and AERONET are stated in parenthesis. Fire denotes data analyzed only during the fire season, while All denotes the entire data set.

Region	t = P90(PC)	t = P87:5(PC)	t = P83:5(PC)	t = P75(PC)
	Err/Corr(%)	Err/Corr(%)	Err/Corr(%)	Err/Corr(%)
North (w/ Fire Count)	-0.02/76	-0.02/78	-0.02/80	<b>-0.01/83</b>
North (w/o Fire Count)	-0.02/69	-0.02/70	-0.02/71	-0.02/71
South (w Fire Count)	-0.01/77	<b>-0.01/78</b>	-0.01/75	0.01/69
South (w/o Fire Count)	-0.01/71	-0.01/70	-0.01/66	-0.01/57

\* not statistically significant at the  $p = 0.05$  level.

**Table 3.** Average values of maximum AOD and average LAI and NDVI during the two annual AOD peaks over the Northern region for the 2001-2013 period.

Stations	Maximum AOD	Average LAI	Average NDVI
	1 <sup>st</sup> Peak (2 <sup>nd</sup> Peak)	1 <sup>st</sup> Peak (2 <sup>nd</sup> Peak)	1 <sup>st</sup> Peak (2 <sup>nd</sup> Peak)
Bac Giang	0.89/0.74	0.44/1.1	0.37/0.58
Chiang Mai	0.5/0.4	2.3/2.97	0.56/0.7
Mukdahan	0.53/0.44	0.96/1.62	0.45/0.67
Nghia Do	0.9/0.71	0.87/1.45	0.39/0.54
Pimai	0.5/0.46	0.54/1.22	0.42/0.61
Ubon Ratchani	0.51/0.46	1.09/1.14	0.48/0.55
Vientiane	0.62/0.59	2.13/2.39	0.52/0.63
Jambi	0.98	2.92	0.68
Kuching	0.66	4.16	0.75
Palangkaraya	1.05	3.72	0.68
Singapore	0.87	1.71	0.4

**Table 4.** Statistics over the Northern region compared to the AERONET stations. Overlapped periods between the reconstructed AOD (AOD<sub>North</sub>) and AERONET are stated in parenthesis. Fire denotes data analyzed only during the fire season, while All denotes the entire data set.

**Table 5.** Average values of maximum AOD and average LAI and NDVI during the annual AOD peak over the Southern region for the 2001-2013 period.

**Table 6.** Statistics over the Southern region compared to the AERONET stations. Overlapped periods between AODSouth and AERONET are stated in parenthesis. Fire denotes data analyzed only during the fire season, while All denotes the entire data set.

\*—— not statistically significant at the  $p = 0.05$  level.

**Table 7.** Error and correlation between the reconstructed AOD versus AERONET and MISR on a monthly basis over the Northern region for the whole 2001-2013 period. Overlapped periods between AODSouth and AERONET, on one hand, and between AODNorth and MISR on the other hand, are stated in parenthesis.


**Table 8.** Error and correlation between the reconstructed AOD versus AERONET and MISR on a monthly basis over the Northern region for the whole 2001-2013 period. Overlapped periods between AODSouth and AERONET, on one hand, and between AODSouth and MISR on the other hand, are stated in parenthesis.

\_\_\_\_\_


\*\_\_\_\_\_ not statistically significant at the  $p = 0.05$  level.




**Figure 1.** Domain with the two EOF regions highlighted and the location of the AERONET stations. Climatological values of LAI (first column) and NDVI (second column) for the 2001-2013 period. Average values are displayed on the first line, while the standard deviation is displayed on the second line.




**Figure 2.** First line: EOF1 (38.2% of variance) (a) and EOF2 (13.3% of variance) (b) of the AOD (2001-2013). Regions of highest AOD variability are delineated by black dots. Second line: PC1 (cutoff 0.006, PC on the left hand axis, AOD on the right hand axis) (c) (red curve) and their associated AOD (green curve) averaged on the region. Third line: PC2 (cutoff 0.01, PC on the left hand axis, AOD on the right hand axis) (d) (red curve) and their associated AOD (green curve) averaged on the region.




**Figure 3.** Climatological values of LAI (first column) and NDVI (second column) for the 2001-2013 period. Average values are displayed on the first line, while the standard deviation is displayed on the second line. Domain with the two EOF regions highlighted and the location of the AERONET stations.




**Figure 4.** Regression Coefficients( $\delta 1$ ) associated to ~~FireCount~~ Fire Count for REG1. Regions of highest AOD variability from the EOF analysis are delineated by black dots.




**Figure 5.** Temporal series of 8-day AERONET AOD (black) and AODNorth (blue) at Bac Giang (a1), Nghia Do (b1), Pimai (c1), and Ubon Ratchathani (d1), Chiang Mai (a2), Mukdahan (b2), and Vientiane (c2) (2001-2013). All x-axes are the time coordinate from Jan 2001 through Dec 2013. All y-axes are the AOD.




(a) Bac Giang




(b) Nghia Do




(c) Pimai




(d) Ubon Ratchathani



(a) Chiang Mai




(b) Mukdahan




(c) Vientiane

**Figure 6.** Temporal series of 8-day AERONET AOD (black) and AODNorth (blue) at Chiang Mai (a), Mukdahan (b), and Vientiane (c) (2001-2013).


**Figure 67.** Temporal series of 8-day AERONET AOD (black) and AODSouth (blue) at Jambi (a), Kuching (b), Palangkaraya (c), and Singapore (d) (2001-2013).




(a) Jambi



(b) Kuching




(c) Palangkaraya



(d) Singapore




**Figure 78.** Basic statistics between MISR and AODNorth (a,b) and AODSouth (a,b) on a monthly basis (2001-2013). Regions of highest AOD variability from the EOF analysis are delineated by black dots. Within the Northern Region ~~dots~~, the mean correlation is 84.8% and the average is ~~77.3%~~ ~~in average~~ (the mean errors is ~~are~~ 0.06 ~~in both cases~~), while in the Southern Region the mean correlation is 79.2% and the average is 72.4% (mean error is 0.08).- The mean errors are given on the left hand side, while the correlations are given on the right hand side.




**Figure 9.** Basic statistics between MISR and AODSouth on a monthly basis (2001–2013). Regions of highest AOD variability from the EOF analysis are delineated by black dots. Within these dots, the mean correlation is 79.2%, while it is 72.4% in average (the mean errors are 0.08 in both cases).

**Appendix A:** Regression coefficients for REG1 and REG2 associated with LAI and NDVI

**Figure A1.** Regression coefficients associated to LAI for REG1 (a) and REG2 (b).




**Figure A2.** Regression coefficients associated to NDVI for REG1 (a) and REG2 (b).




**Appendix B:** Results at the AERONET stations on a monthly basis


**Figure B1.** Temporal series of AERONET AOD (black), AODNorth (blue), and AOD from MISR (red) at four stations of the Northern region (2001-2013).




(a) Bac Giang



(b) Chiang Mai




(c) Mukdahan




(d) Nghia Do


**Figure B2.** Temporal series of AERONET AOD (black), AODNorth (blue), and AOD from MISR (red) at three stations of the Northern region (2001-2013).



(a) Pimai




(b) Ubon Ratchathani




(c) Vientiane


**Figure B3.** Temporal series of AERONET AOD (black), AODNorth (blue), and AOD from MISR (red) at two stations of the Southern region (2001-2013).




(a) Jambi



(b) Kuching





(c) Palangkaraya



(d) Singapore



**Appendix C: Geospatial locations of fires.**

**Figure C1.** Geospatial aggregate of all fires in January 2013 over Northern Southeast Asia and Southern Southeast Asia respectively.





**Figure C2.** Geospatial aggregate of all fires in September 2013 over Northern Southeast Asia and Southern Southeast Asia respectively.



**Appendix D: Table D.** Complementary information on AERONET stations: geographical location, data availability, average ( $\mu$ ) and standard deviation ( $\sigma$ ) values for AOD, LAI, and NDVI from MODIS, and environment description.

Stations	Availability	AOD	LAI	NDVI	Other information
Bac Giang, VN (North)	2003-2009	m=0.57, r=0.28	m=0.68, r=0.43	m=0.45, r=0.15	Rural, surrounded by crops and industrial parks
Chiang Mai, TH (North)	2006-2013	m=0.29, r=0.17	m=2.19, r=0.77	m=0.61, r=0.07	Urban, surrounded by agricultural fields
Mukdahan, TH (North)	2003-2009	m=0.32, r=0.16	m=1.13, r=0.37	m=0.55, r=0.09	Rural, surrounded by agricultural fields
Nghia Do, VN (North)	2010-2013	m=0.57, r=0.29	m=0.92, r=0.59	m=0.42, r=0.15	Urban
Pimai, TH (North)	2003-2007	m=0.33, r=0.17	m=0.72, r=0.27	m=0.49, r=0.1	Rural, surrounded by agricultural fields
Ubon Ratchathani, Th (North)	2009-2012	m=0.33, r=0.17	m=1.03, r=0.33	m=0.52, r=0.07	Semi-urban, surrounded by agricultural fields
Vientiane, LA (North)	2011-2012	m=0.35, r=0.21	m=2.03, r=0.45	m=0.55, r=0.08	Semi-urban, surrounded by agricultural fields
Jambi, ID (South)	2012-2013	m=0.36, r=0.31	m=2.72, r=1.48	m=0.66, r=0.11	Rural, surrounded by jungle
Kuching, MY (South)	2011-2013	m=0.31, r=0.32	m=3.75, r=1.65	m=0.7, r=0.1	Rural, surrounded by jungle
Palangkaraya, ID (South)	2012-2013	m=0.3, r=0.37	m=3.21, r=1.43	m=0.69, r=0.11	Rural, surrounded by jungle
Singapore, SG (South)	2006-2013	m=0.34, r=0.25	m=2.38, r=1.29	m=0.42, r=0.06	Urban

**Appendix E: Table E.** Error and correlation between the reconstructed AOD versus AERONET and MISR on a monthly basis over the Northern region for the whole 2001-2013 period. Overlapped periods between AODNorth, AODSouth, and AERONET, on one hand, and between AODNorth, AODSouth, and MISR on the other hand, are stated in parenthesis.

Stations	AERONET	MISR
	Err/Corr(%)	Err/Corr(%)
Chiang Mai (65/156) – (129/156)	-0.09/82	0.02/77
Bac Giang (49/156) – (127/156)	-0.03/73	0.1/72
Mukdahan (72/156) – (135/156)	0.0/78	0.04/79
Nghia Do (28/156) – (133/156)	-0.12/83	0.07/71
Pimai (41/156) – (145/156)	0.02/70	0.03/66
Ubon Ratchathani (32/156) – (136/156)	0.0/90	0.05/74
Vientiane (15/156) – (127/156)	-0.01/76	0.02/81
Jambi (18/156) – (79/156)	-0.14/57	0.06/69
Kuching (25/156) – (114/156)	-0.06/71	0.1/64
Palangkaraya (18/156) – (102/156)	-0.11/73	0.04/79
Singapore (78/156) – (122/156)	-0.11/-2*	0.04/57

\* not statistically significant at the  $p = 0.05$  level.