



Supplement of


How skillfully can we simulate drivers of aerosol direct climate forcing at the regional scale?

P. Crippa et al.

Correspondence to: P. Crippa (paola.crippa@ncl.ac.uk)


The copyright of individual parts of the supplement might differ from the CC-BY 3.0 licence.

1 Figure S1. Empirical quantile-quantile plot of simultaneous measurements of AE at 500 nm
 2 between MODIS (Terra) and AERONET (where the AERONET station longitude (E) and
 3 latitude (N) are given in the legend). As shown, the MODIS data tend to a bimodal distribution,
 4 while in the AERONET observations AE is a continuous variable (or nearly so). Thus, while
 5 in comparison with WRF-Chem simulations AE from AERONET is treated as a continuous
 6 variable, in the majority of comparisons with MODIS a threshold of 1 is applied to identify the
 7 dominance of coarse mode ($AE < 1$) versus fine mode ($AE > 1$).



9 Figure S2. Mean fraction bias (MFB) of near-surface daily mean $PM_{2.5}$ concentrations as
10 simulated by WRF-Chem and observed at EPA sites during (a) winter, (b) spring, (c) summer
11 and (d) fall. As shown, $PM_{2.5}$ concentrations from WRF-Chem exhibit a positive bias (MFB >
12 0) for most sites and in most seasons, but the bias is largest over the southern states during
13 summer. Note also that the MFB in $PM_{2.5}$ concentrations greatly exceeds that for either AOD
14 or AE (see Fig. 2).

15




16

17

18

19 Figure S3. (a) Taylor diagram for monthly accumulated precipitation during 2008 as simulated
20 by WRF-Chem and in the gridded observations (Matsuura and Willmott, 2009), after applying
21 a linear interpolation to match the WRF-Chem grid. Panels (b) and (c) show the difference
22 [mm] between observed and simulated accumulated precipitation during the month of (b)
23 September and (c) October 2008. Values larger than zero indicate the observed precipitation is
24 higher than the simulated one.

25




26

27


28

29

30 Figure S4. Spatial coherence in the identification of extreme AOD (i.e. the occurrence of AOD
 31 above the 75th percentile value) from WRF-Chem and MODIS Terra and Aqua on a monthly
 32 basis. Green areas denote grid cells defined as experiencing extreme AOD in the WRF-Chem
 33 simulations, blue pixels indicate extreme values as diagnosed using MODIS, while red pixels
 34 indicate areas where the occurrence of extreme values is indicated by both the WRF-Chem
 35 simulations and the MODIS observations.



37 Figure S5. Fraction of near-surface monthly averaged mass concentration of nitrate versus the
38 sum of aerosol nitrate and sulfate as simulated in the accumulation mode by WRF-Chem for
39 each calendar month during 2008.



40
41