

1 **Inter-comparison of stratospheric mean-meridional**
2 **circulation and eddy mixing among six reanalysis datasets**

3

4 **Kazuyuki Miyazaki¹, Toshiki Iwasaki², Yoshio Kawatani¹, Chiaki Kobayashi³,**
5 **Satoshi Sugawara⁴, Michaela I. Hegglin⁵**

6

7 [1]{Japan Agency for Marine-Earth Science and Technology, Yokohama, Japan}

8 [2]{Department of Geophysics, Graduate School of Science, Tohoku University, Sendai,
9 Japan}

10 [3]{Meteorological Research Institute, Tsukuba, Japan}

11 [4]{Department of Science Education, Miyagi University of Education, Sendai, Japan}

12 [5]{Department of Meteorology, University of Reading, Reading, UK}

13 Correspondence to: K. Miyazaki (kmiyazaki@jamstec.go.jp)

14

15 **Abstract**

16 The stratospheric mean-meridional circulation (MMC) and eddy mixing are compared among
17 six meteorological reanalysis datasets: NCEP-NCAR, NCEP-CFSR, ERA-40, ERA-Interim,
18 JRA-25, and JRA-55 for the period 1979–2012. The reanalysis datasets produced using
19 advanced systems (i.e., NCEP-CFSR, ERA-Interim, and JRA-55) generally reveal a weaker
20 MMC in the Northern Hemisphere (NH) compared with those produced using older systems
21 (i.e., NCEP/NCAR, ERA-40, and JRA-25). The mean mixing strength differs largely among
22 the data products. In the NH lower stratosphere, the contribution of planetary-scale mixing is
23 larger in the new datasets than in the old datasets, whereas that of small-scale mixing is
24 weaker in the new datasets. Conventional data assimilation techniques introduce analysis
25 increments without maintaining physical balance, which may have caused an overly strong
26 MMC and spurious small-scale eddies in the old datasets. At the NH mid-latitudes, only
27 ERA-Interim reveals a weakening MMC trend in the deep branch of the Brewer–Dobson
28 circulation (BDC). The relative importance of the eddy mixing compared with the mean
29 meridional transport in the subtropical lower stratosphere shows increasing trends in ERA-

1 Interim and JRA-55; this together with the weakened MMC in the deep branch may imply an
2 increasing Age-of-Air (AoA) in the NH middle stratosphere in ERA-Interim. Overall,
3 discrepancies between the different variables and trends therein as derived from the different
4 reanalyses are still relatively large, suggesting that more investments into these products are
5 needed in order to obtain a consolidated picture of observed changes in the BDC and the
6 mechanisms that drive them.

7

8 **1 Introduction**

9 The Brewer–Dobson circulation (BDC), which was discovered by Brewer (1949) and Dobson
10 (1929; 1956), consists of the mean-meridional circulation (MMC) and eddy mixing in the
11 stratosphere. The stratospheric MMC is composed of ascending motions in the tropics,
12 poleward motions toward mid and high latitudes, and descending motions at high latitudes.
13 Planetary waves that propagate from the troposphere break and cause eddy mixing primarily
14 in the stratospheric surf zone surrounding the polar vortex in the winter hemisphere (McIntyre
15 and Palmer, 1983). Because the BDC motion is too slow to measure directly, the detailed
16 structure and long-term variations of the BDC are not well understood.

17 Stratospheric age-of-air (AoA) (Waugh and Hall, 2002) derived from observation data of
18 long-lived chemical compounds is frequently used as a surrogate for the combined effects of
19 the MMC and eddy mixing in order to investigate the structure and long-term variations in the
20 BDC. Observational studies have found a positive AoA trend in the Northern Hemisphere
21 (NH) mid-latitudes in the middle to upper stratosphere based on balloon measurements of SF₆
22 and CO₂ for 1975–2005 (Engel et al., 2009), MIPAS SF₆ measurements for 2002–2010
23 (Stiller et al., 2012), balloon-based measurements of SF₆ and CO₂ for 1975–2012 (Ray et al.,
24 2014), and a merged long-term satellite data record of water vapour for 1986–2010 (Hegglin
25 et al., 2014). In the Southern Hemisphere (SH) mid-latitude lower stratosphere, Stiller et al.
26 (2012) found in contrast a negative AoA trend, a result confirmed by Hegglin et al. (2014).
27 The study by Hegglin et al. (2014) also finds a negative AoA trend in the NH lower
28 stratosphere in contrast to Stiller et al. (2012), with the difference likely explainable by the
29 different time periods considered. However, these observed trends are not fully consistent
30 with simulated results using general circulation models (GCMs), which show instead an
31 acceleration in the mean BDC throughout the stratosphere (e.g., Butchart et al., 2011). The
32 acceleration of the mean BDC strength in the model simulation is associated with enhanced

1 wave driving (Garcia and Randel, 2008) and shifting critical levels for wave breaking
2 (Shepherd and McLandress, 2011). The reasons for the inconsistency between the
3 measurements and models, especially in the middle and upper stratosphere, have not yet been
4 investigated.

5 MMC and eddy mixing in the BDC are intrinsically linked (Dunkerton, 1978) and hence play
6 both important roles in determining distributions of long-lived chemical species and AoA in
7 the stratosphere. Interpretation of AoA variations therefore needs to take into account changes
8 in both MMC and eddy mixing. Several recent studies have now quantified the effects of eddy
9 mixing on AoA variations in more detail (Ray et al., 2010; Garny et al., 2014; Ploeger et al.,
10 2015a, 2015b). Garny et al. (2014) stated that eddy mixing causes recirculation of air in the
11 stratosphere, and acts to increase the mean value of AoA. Mixing inside the surf zone
12 modifies the latitudinal distribution of AoA and chemical species, whereas mixing across the
13 subtropical transport barrier is suggested to be important for the mean AoA value above the
14 mixing level (Garny et al., 2014; Ploeger et al., 2015a).

15 Meteorological reanalyses provide a realistic meteorological field by combining model
16 information with actual observations, and have the potential to provide an alternative tool to
17 study the BDC and AoA variations. Iwasaki et al. (2009) compared four reanalysis datasets
18 (NCEP-NCAR, NCEP-DOE, ERA-40, and JRA-25) and found large differences in their
19 representation of the BDC. Wright and Fueglistaler (2013) showed large differences in the
20 simulated diabatic heat budget in the tropical upper troposphere and lower stratosphere in five
21 reanalysis models (NCEP-NCAR, NCEP-CFSR, JRA-25, ERA-Interim, and MERRA), with
22 substantial implications for representation of transport and mixing. Recently, Abalos et al.
23 (2015) compared the MMC in three reanalyses (ERA-Interim, JRA-55, and MERRA) using
24 three different estimates: from the transformed Eulerian mean (TEM, Andrews and McIntyre,
25 1976) residual circulation and based on momentum and thermodynamic balances. They
26 showed a relatively large spread (around 40%) among the estimates of the magnitude of
27 tropical upwelling. Monge-Sanz et al. (2007, 2012) highlighted that the representation of the
28 BDC (including eddy mixing) has become much more realistic in ERA-Interim than in ERA-
29 40, thanks to large investments made into improving the reanalysis product. They showed that
30 AoA derived using ERA-Interim displays an increasing trend in the NH mid-latitude
31 stratosphere above 25 km in 1989–2010, consistent with the findings by Hegglin et al. (2014)
32 for 1986–2010. Ploeger et al. (2015a) quantified the effects of mixing and residual circulation

1 on mean age trends using ERA-Interim, and showed the importance of eddy mixing on the
2 increasing AoA trend in the NH middle stratosphere in 2002–2012. Diallo et al. (2012)
3 showed that the AoA trend derived using ERA-Interim in the middle stratosphere is positive
4 over the 1989–2010 period. In order to gain confidence in the ERA-Interim results, it is now
5 important to assess their consistency with long-term variations in the BDC obtained from
6 other reanalysis products.

7 Differences in the forecast model, assimilated measurements, and data assimilation technique
8 used for producing reanalysis datasets can lead to differences in their representation of the
9 BDC. Model simulations without any assimilation produce meteorological fields that follow
10 the dynamical and thermodynamic balance of the forecast model. Data assimilation analysis
11 increments, introduced by using conventional data assimilation techniques such as the three-
12 dimensional variational (3D-VAR) one, may upset this balance and degrade the expression of
13 momentum budget and wave structures. This is because they introduce an additional force,
14 without maintaining physical balance, as a result of its isotropic and instantaneous analysis
15 increment. In the 3D-VAR analysis, mean ascending motions in the tropics and mixing in the
16 subtropics in the stratosphere were found to be excessively strong (Schoeberl et al., 2003; Tan
17 et al., 2004; Scheele et al., 2005). Advanced data assimilation techniques such as the four-
18 dimensional variational method (4D-VAR) are capable of assimilating observations at the
19 exact time while maintaining the dynamical balance because of the use of flow-dependent
20 analysis, which are expected to improve the representation of both MMC and eddy mixing.

21 In this paper, we compare MMC and eddy mixing in the stratosphere for six reanalysis
22 datasets; NCEP-NCAR, NCEP-CFSR, ERA-40, ERA-Interim, JRA-25, and JRA-55. The
23 analysis is conducted for the 34 years from 1979 to 2012 based on mass-weighted isentropic
24 zonal means that allow accurate analysis of Lagrangian-mean motions and eddy mixing.
25 Based on the comparison of the mean and eddy components in the BDC, we discuss whether
26 any of the reanalysis data have the potential to reveal useful information on long-term AoA
27 variations.

28 **2 Methodology**

29 **2.1 Data**

30 The six reanalysis datasets used in our comparison can be described as follows: 1) NCEP-
31 NCAR – the National Centers for Environmental Prediction (NCEP)-National Center for

1 Atmospheric Research (NCAR) reanalysis product (Kalnay et al., 1996), with a model grid
 2 resolution of T62L28 produced using a 3D-VAR technique; 2) NCEP-CFSR – the NCEP
 3 Climate Forecast System Reanalysis (Saha et al., 2010) with a model grid resolution of
 4 T382L64 produced using a 3D-VAR technique; 3) ERA-40 – the 40-yr ECMWF Re-Analysis
 5 (Simmons and Gibson, 2000) with a model grid resolution of T159L60 produced using a 3D-
 6 VAR technique; 4) ERA-Interim – a continuously updated reanalysis since 1979 (Simmons et
 7 al., 2007), with a model grid resolution of T225L60 produced using a 4D-VAR technique; 5)
 8 JRA-25 – the Japanese 25-year reanalysis product (Onogi et al., 2007), with a model grid
 9 resolution of T106L40 provided using a 3D-VAR technique; and 6) JRA-55 – the Japanese
 10 55-year reanalysis product (Kobayashi et al., 2015), with a model grid resolution of T319L60
 11 provided using a 4D-VAR technique.

12 We here classify NCEP-NCAR, JRA-25, and ERA-40 as old datasets, and NCEP-CFSR,
 13 JRA-55, and ERA-Interim as new datasets because of improvements in these latter made by
 14 using updated forecast models, updated bias correction algorithms, and advanced data
 15 assimilation analysis. We use reanalyses results for the 34 years after 1979, when satellite
 16 measurements were assimilated into the reanalysis. For ERA-40, the mean state and linear
 17 trend are estimated for the 24 years from 1979 to 2002, since the data are not available after
 18 2002.

19 **2.2 Analysis framework**

20 The analysis of MMC and eddy mixing is based on mass-weighted isentropic zonal means
 21 (hereafter referred to as MIM analysis; Iwasaki 1989, 1992; Miyazaki and Iwasaki 2005). The
 22 MIM zonal mean is defined as:

$$23 \quad \overline{A(\phi, \theta, t)}^* = \frac{1}{2\pi} \int A(\lambda, \phi, \theta, t) \left(\frac{\partial p}{\partial \theta} / \frac{\partial \bar{p}}{\partial \theta} \right) d\lambda, \quad (1)$$

24 where ϕ is the latitude, θ is the potential temperature, t is the time, λ is the longitude, and p is
 25 the pressure. The asterisks and overbars represent mass-weighting and isentropic zonal means,
 26 respectively. Eddies are defined as departures from the mass-weighted zonal means,

$$27 \quad A' \equiv A - \overline{A}^*. \quad (2)$$

28 Their correlations are given by

1 $\overline{(A'B')}^* = \overline{(AB)}^* - \overline{A^*B^*}$, (3)

2

3 We use isentropic zonal mean pressure for the vertical coordinate,

4 $p_{\dagger} \equiv \bar{p}$. (4)

5 Unlike other isentropic coordinate analyses (Andrews, 1983; Tung, 1982, 1986), in the MIM
6 analysis, the mass-weighting is considered not only for meridional circulation but also for
7 other variables such as zonal wind following Johnson (1980). The MIM analysis thus
8 expresses the conservative nature of momentum, heat, and minor constituents, including the
9 exact lower boundary conditions and non-geostrophic effects (Iwasaki, 1989, 1992; Miyazaki
10 and Iwasaki, 2008). The MIM analysis also exactly specifies the eddy diabatic and adiabatic
11 transport terms (Miyazaki and Iwasaki, 2005).

12 The TEM (Andrews and McIntyre, 1976) provides a useful framework for understanding
13 mean and eddy transports; however, the estimation of the transport fluxes is limited and
14 complicated. Randel et al. (1994), Strahan et al. (1996), and Abalos et al. (2013) estimated
15 eddy transport terms based on eddy flux vectors for small amplitude eddies following
16 Andrews et al. (1987), while some studies estimated this term as residuals considering the
17 uncertainty and difficulty in the eddy transport term estimations (e.g., Randel et al., 1998).
18 Miyazaki et al. (2008) found a significant (>30%) difference in the mean vertical velocity
19 around the Antarctic polar vortex between the TEM residual vertical velocity and the MIM
20 MMC analysis, which can be attributed to the assumptions applied for the TEM Stokes
21 corrections. Most of the disadvantages of conventional analysis methods such as TEM (e.g.,
22 complicated and inaccurate representation of transport by both mean and eddy motions, lower
23 boundary conditions, and mass conservations) can be avoided using the MIM analysis
24 (Iwasaki, 1989; Tanaka et al., 2004; Miyazaki and Iwasaki, 2005).

25 We here focus on two altitudes; 440 K (at approximately 90–80 hPa) and 560 K (40–30 hPa)
26 as representatives of the shallow and deep branches of the BDC, respectively. Birner and
27 Bönisch (2011) reported that the shallow branch extends to about 50 hPa, and the deep branch
28 is located above that altitude. The analysis results are presented on isentropic coordinates for
29 diagnosing adiabatic and diabatic transport components. Note that if the potential temperature
30 at a constant pressure changes with time, there would not be exact agreement between the

1 estimated trends in pressure and isentropic coordinates. For example, in the last 30 years
 2 (2008–2012 mean minus 1979–1983 mean), potential temperature at 70 hPa decreased by
 3 about 2.5 K at low and mid-latitudes in both hemispheres in ERA-Interim. Nevertheless, the
 4 general structure of the linear trend was similar between the two coordinates (and hence will
 5 not be shown here). The long-term linear trend is estimated based on the least-squares fitting.
 6 The statistical significance is determined for the 95% confidence level using the Mann-
 7 Kendall test (Mann, 1945; Kendall, 1975).

8 2.2.1 Mean-meridional circulation (MMC)

9 The mass stream function χ in the MIM analysis is calculated from integrating meridional
 10 velocity with respect to p_{\dagger} :

$$11 \quad \chi = a \cos \phi \int_0^{p_{\dagger}} \overline{v^*} dp_{\dagger}. \quad (5)$$

12 where a is the Earth's radius, and v is the meridional wind. $\overline{v^*}$ is calculated from meridional
 13 wind data with consideration of the mass-weighted isentropic zonal means. The diagnostic
 14 form of the zonal mean continuity equation without an eddy term in the MIM analysis
 15 confirms that the mean meridional circulation can be expressed by the nondivergent mass
 16 stream function (Iwasaki, 1989). The mean vertical velocity $\overline{w_{\dagger}^*}$ is obtained from the mass
 17 stream function χ as follows:

18

$$19 \quad \overline{w_{\dagger}^*} = \frac{1}{2\pi a \rho_0 \cos \phi} \frac{\partial \chi}{\partial \phi}. \quad (6)$$

20 where ρ_0 is the reference atmospheric density. The mean vertical velocity $\overline{w_{\dagger}^*}$ can be related
 21 to the diabatic heating $\overline{\dot{\theta}^*}$ (Eq. (2.7) in Tanaka et al., 2004).

22

23 2.2.2 Eddy mixing

24 By assuming a flux-gradient linear relationship, the diffusion coefficient provides a measure
 25 of eddy mixing. The isentropic diffusion coefficient K_{yy} can be derived from the eddy
 26 meridional flux and meridional potential vorticity (PV) gradient on isentropic surfaces (Tung,

1 1986; Newman et al., 1988; Bartels et al., 1998; Miyazaki and Iwasaki, 2005; Miyazaki et al.,
 2 2010b) by neglecting the influence of slant diffusion:

$$3 \quad \left[\overline{(v'q')} \right]_l \approx -K_{yy}(l) \left[\left(\frac{\partial \overline{q^*}}{\partial \varphi} \right)_\theta \right]_l, \quad (7)$$

4 where q is the PV, and $[\]_l$ denotes the time average.

5 Under frictionless and adiabatic conditions, the PV acts as an atmospheric passive tracer
 6 (Hoskins et al., 1985). Miyazaki et al. (2010a) demonstrated that the diabatic source–sink
 7 effect on the PV budget is much smaller than the transport effects in the subtropical and
 8 extratropical stratosphere. In the K_{yy} estimation, steady conservative wave motions projected
 9 onto a meridional plane can cause apparent diffusion in addition to the true diffusion caused
 10 by dissipative wave motions, which may lead to significant differences between the estimated
 11 K_{yy} and the true eddy mixing. In order to reduce this effect in our estimates, a time-averaging
 12 window t is applied to the eddy PV flux, as in Miyazaki and Iwasaki (2005) and Miyazaki et
 13 al. (2010b). As a result, we expect that the estimated K_{yy} provides information on eddy mixing
 14 characteristics similar to estimates of the effective diffusivity (Nakamura, 1996; Haynes and
 15 Shuckburgh, 2000).

16 The absolute value of K_{yy} is influenced by the choice of l (set to one month in this study). We
 17 confirmed that, in all datasets, the estimated K_{yy} becomes smaller by increasing l from 6 hours
 18 to 10 days, and it becomes nearly constant when l is set to be more than 10 days. For instance,
 19 K_{yy} ($l > 10$ days) is approximately 15–20 % smaller than that of K_{yy} ($l = 6$ hour) in the NH
 20 middle stratospheric surf zone in December to February (DJF), for the case of ERA-Interim.
 21 These constant values are considered to represent the diffusion coefficient due to the true
 22 diffusions.

23 We should note that, even after eliminating the influence of apparent diffusion in K_{yy}
 24 estimates, there are limitations in elucidating eddy mixing from these estimates. As discussed
 25 by Nakamura (2008), whilst a part of the Eulerian eddy diffusivity (e.g., K_{yy}) can be attributed
 26 to instantaneous, irreversible mixing in a way similar to effective diffusivity, the Eulerian
 27 eddy diffusivity and eddy diffusivity are fundamentally different, both qualitatively and
 28 quantitatively. This is because of difficulties associated with representing eddy advective
 29 transport in the Eulerian formulation. In addition, the results of the K_{yy} analysis and related
 30 variables are presented here in the geometric latitude coordinate system (where a zonal

1 average is taken for air parcels with different PVs), whereas effective diffusivity is presented
 2 in the equivalent latitude (EL) coordinate system (based on the latitude circle that encloses the
 3 same area as the PV contour). Latitudinal variations of zonal-mean eddy mixing and
 4 associated fields (e.g., strong eddy mixing outside the polar vortex) are more clearly presented
 5 in the EL coordinate system.

6 2.2.3 The relative importance of mean and eddy transports

7 The zonal mean equation in the MIM analysis can accurately separate meridional transport
 8 into mean transport by Lagrangian-mean circulation and eddy (diffusion) transport. In the
 9 MIM analysis, the mean and eddy PV fluxes are defined as follows:

$$10 \quad Mean = \left(\overline{v^* q^*}, \overline{\dot{\theta}^* q^*} \right), \quad (8)$$

$$11 \quad Eddy = \left(\overline{(v'q')^*}, \overline{(\dot{\theta}'q')^*} \right). \quad (9)$$

12 where $\overline{v^*}$ is calculated from meridional wind data (c.f., Sec. 2.2.1). The mean transport fluxes
 13 are parallel to isopleths of the mass streamfunction, whereas the eddy transport fluxes are
 14 parallel to the isentropes under diabatic conditions. Only the meridional components are
 15 analyzed in this study.

16 In order to evaluate the relative importance of eddy transport in each reanalysis product, we
 17 estimate the ratio of eddy and total meridional transport fluxes of PV as follows:

$$18 \quad \frac{\left| \left[\overline{(v'q')^*} \right]_l \right|}{\left| \left[\overline{(v'q')^*} \right]_l \right| + \left| \left[\overline{v^* q^*} \right]_l \right|}. \quad (10)$$

19 If the transport ratio is larger (smaller) than 0.5, the eddy transport (mean transport)
 20 dominates the meridional transport on isentropic surfaces. As in the K_{yy} estimation (c.f., Eq.
 21 (7)), a time average ($l=1$ month) was applied to the mean and eddy transport fluxes.

1 **3 Results**

2 **3.1 Eddy mixing**

3 Figure 1 shows the seasonal mean isentropic diffusion coefficient K_{yy} ($l=1$ month) (hereafter
4 referred to as K_{yy}). Large K_{yy} values reveal strong isentropic mixing in the stratospheric surf
5 zone in both hemispheres, which is stronger in the NH than in the SH. The hemispheric
6 asymmetry can be attributed to the differences in planetary wave activity (e.g., Shepherd,
7 2000). The small K_{yy} equatorward of about 30° is indicative of a barrier to horizontal transport
8 between the tropics and mid-latitudes in both hemispheres (Plumb, 1996). At SH high
9 latitudes, K_{yy} becomes large after the break-up of the Antarctic polar vortex (figure not
10 shown). The eddy mixing is strongly suppressed near the core of the subtropical jet stream,
11 but shows clear maxima at its upper flank (equatorwards) and also at its lower flank
12 (poleward, during DJF). These general characteristics are commonly found in all the datasets,
13 and are generally consistent with the analysis of effective diffusivity (e.g., Haynes and
14 Shuckburgh, 2000). It is noted that the counter-gradient transport is found in the negative
15 region of K_{yy} (shaded in white) in the summer hemisphere, which is associated with almost
16 flat PV gradients.

17 The mean value and linear trend of K_{yy} are estimated in the surf zone ($40\text{--}60^\circ$) and subtropics
18 ($15\text{--}25^\circ$) around the overturning latitude of the MMC (Table 1 and Fig. 2). Isentropic mixing
19 in the surf zone influences the latitudinal gradient of tracers, whereas the mixing across the
20 subtropics is considered to be important for the stratospheric mean AoA because it
21 recirculates old air from the extratropics to the tropics (Garny et al., 2014; Ploeger et al.,
22 2015a). The anomalously low value in NCEP-NCAR in the middle stratosphere is associated
23 with the low top height of the reanalysis product.

24 K_{yy} shows an increasing trend in the NH middle stratosphere (i.e., 560 K) surf zone in all the
25 datasets over the time period 1979–2012 (+0.5 to +5.2 %/decade), with relatively weak trends
26 in the newer datasets (+0.5 to +2.2 %/decade). The trend in the NH at 440 K is positive in
27 ERA-Interim and JRA-55 in 1979–2012 and also in ERA-40 in 1979–2001. In the SH surf
28 zone, the K_{yy} shows an increasing trend from the lower to middle stratosphere in all the
29 datasets except for NCEP-NCAR and ERA-40. The positive trend is large in JRA-25 and
30 JRA-55 at 440 K (+18.0 and +16.1 %/decade, respectively) and in JRA-25 at 560 K
31 (+13.9 %/decade). The intensified surf zone mixing could be associated with changes in the

1 critical level. Climate model simulations demonstrate that long-term changes in zonal wind
2 such as the shift of zero wind line in response to climate change can enhance the upward
3 propagation of westward-propagating waves (Kawatani et al., 2011; Shepherd and
4 McLandress, 2011). Further investigations are required to comprehend the relationship
5 between changes in the critical level, wave forcing, and mixing strength in the reanalysis
6 products. The standard deviation in the middle stratosphere surf zone is smaller in the new
7 datasets than in the old datasets in both hemispheres by about 30–60% in both hemispheres.

8 In the NH winter subtropical lower stratosphere, K_{yy} shows increasing trends in NCEP-NCAR,
9 NCEP-CFSR, and ERA-40 (+0.5 to +9.0 %/decade) and decreasing trends in the other
10 datasets (-0.6 to -1.7 %/decade). The mean K_{yy} value differs largely among the data products
11 in the NH subtropical lower stratosphere (0.56–0.81). The large difference in the subtropical
12 mixing among the datasets could be associated with different representations of the Quasi-
13 Biennial Oscillation (QBO) in different reanalysis products (Pawson and Fiorino, 1998; Kim
14 and Chun, 2015; Kawatani et al., 2016). In the SH subtropical lower stratosphere, the K_{yy}
15 trend is positive only in ERA-Interim (+1.2 %/decade) and NCEP-NCAR (+8.0 %/decade),
16 and shows a large negative value in NCEP-CFSR (-10.0 %/decade) and ERA-40 (-
17 11.2 %/decade). Because of the large interannual variations, the estimated trends are not
18 statistically significant for most cases in the subtropics and the mid-latitudes.

19 The K_{yy} trend in the NH middle stratosphere surf zone is nearly zero or positive (-1.8 to
20 +8.2 %/decade) in the first 22 years in datasets, except for NCEP-NCAR and ERA-40, then
21 changes to large negative values (-5.7 to -31.5 %/decade) in the last 12 years in all the
22 datasets (Table 2). The negative trend in the latter period is larger in the old datasets (-23.5 to
23 -31.5 %/decade) than in the new datasets (-5.7 to -16.8 %/decade). These decadal scale
24 changes in the mixing trend seem to be consistent with those in the tropical upward mass flux
25 and MMC in the NH (c.f., Section 3.2). This suggests that variations in wave forcing lead to
26 decadal scale changes in both the mean and eddy transports as is expected, given that these
27 two features are intrinsically connected with each other (c.f., Section 4.1). In the SH middle
28 stratosphere surf zone, the K_{yy} trend is positive (+4.4 to +14.0 %/decade, except for ERA-40)
29 in the first 22 years and becomes negative (-24.4 to -40.4 %/decade) in the last 12 years.

30 In the NH subtropics (not shown in table), K_{yy} shows a greater positive trend in the last 12
31 years than in the first 22 years in ERA-Interim (from +1.8 to +17.1 %/decade) and JRA-55

1 (from +2.4 to +7.0 %/decade), whereas the trend is almost constant during both periods in
2 NCEP-CFSR (from +8.1 to +8.9 %/decade) and JRA-25 (+1.8 to +3.9 %/decade).

3 **3.2 Mean-meridional circulation (MMC)**

4 Figure 3 compares the meridional cross section of the mass stream function averaged over
5 DJF and June to August (JJA) in 1979–2012. Note that ERA-40 has been averaged over the
6 shorter time period 1979–2002 due to limited availability of the dataset. Comparison of ERA-
7 40 and ERA-Interim averaged over the same (shorter) time period, however, indicates that the
8 results are comparable. The general structure of the MMC, such as the poleward motion from
9 low latitudes of the summer hemisphere to mid and high latitudes of the winter hemisphere,
10 and the descending motions at high latitudes of winter hemispheres, is commonly found in all
11 the datasets. However, details regarding the structure, intensity, and trend of the MMC differ
12 among the data. In NCEP-NCAR, the MMC is noisy and unrealistically distorted in the
13 middle stratosphere (altitudes above about 550 K), and also in the SH lower stratosphere
14 during JJA. In ERA-40, it is relatively strong throughout the stratosphere in both hemispheres,
15 as already pointed out by Wohltmann and Rex (2008). Similarly, JRA-25 exhibits an MMC
16 that is somewhat stronger in the middle to upper stratosphere than that found in the other
17 datasets. In NCEP-CFSR, the tropical mean upward motion is distorted in the lower
18 stratosphere, so that mean air trajectories do not ascent in a straight line. Significant diversity
19 of the tropical mean upward motions (not shown) may be related to differences in simulated
20 diabatic heating rates in the forecast models (Fueglistaler et al., 2009).

21 Figure 4 compares the seasonal variation of the mass stream functions at 560 K. The strong
22 mean poleward motions are present from low latitudes to mid and high latitudes from autumn
23 to spring. They are relatively suppressed at the polar vortex edge. In NCEP-NCAR, the cross-
24 equatorial mean-meridional flow is relatively weak throughout the year. Also, the strong
25 poleward flow at mid-latitudes does not exhibit the seasonality apparent in the other
26 reanalyses. In ERA-40 and JRA-25, the mean poleward flow from the subtropics to the mid-
27 latitudes is relatively strong in both hemispheres throughout the year.

28 The MMC long-term trend differs greatly among the reanalysis datasets (Fig. 5). In NCEP-
29 NCAR, the spatial structure of the linear trend slope is very noisy, likely reflecting
30 inhomogeneities in the reanalysis product. In NCEP-CFSR, the MMC shows a strengthening
31 trend from the tropics to the mid-latitudes in the lower and middle stratosphere in both winter

1 hemispheres. JRA-25 reveals a strengthening trend in the lower and middle stratosphere
2 equatorward of 60N in the NH during winter and throughout the lower stratosphere in the SH
3 in both seasons, whereas it shows a weakening trend above 500 K in the SH in DJF. JRA-55
4 shows a weak positive trend in the lower stratosphere in winter. ERA-40 reveals a
5 strengthening trend throughout the lower and middle stratosphere over 1979–2002 in the SH
6 in JJA, whereas the trend pattern is noisy in the NH in both seasons. Only in ERA-Interim, the
7 MMC in the NH middle stratosphere (or above about 480 K) tends to weaken during winter.
8 A negative trend is also found from the tropics to the mid-latitudes in both hemispheres in JJA
9 in the middle stratosphere in ERA-Interim. The BDC shallow branch mostly shows a
10 strengthening trend in ERA-Interim as in other datasets. Abalos et al. (2015) revealed that the
11 acceleration in the MMC is a qualitatively robust result across different estimates (from the
12 TEM residual circulations and based on momentum and thermodynamic balances). Obvious
13 structural changes (i.e., that the shallow and deep branches changed differently) in the
14 wintertime MMC can be found only in ERA-Interim and to some extent also in JRA-25.
15 There are only a few statistically significant regions in the BDC trends. The low statistical
16 significance could be partly attributed to the fact that large variances associated with ENSO
17 and QBO, as pointed out by Abalos et al. (2015), are not removed in our trend estimates.

18 Figure 6 compares the time series of the mass stream function at mid-latitudes (40–60°) in the
19 lower stratosphere (440 K) and the middle stratosphere (560 K) during the winter seasons.
20 The mean value and linear trend are summarized in Table 3. In the NH winter mid-latitudes at
21 both altitudes, the mean MMC strength is weaker in the new data than in the old data. At 440
22 K, NCEP-NCAR is about 60% larger compared to NCEP-CFSR, ERA-40 was 30% larger
23 compared to ERA-Interim, and JRA-25 is 30% larger compared to JRA-55; at 560 K, the
24 mean MMC strength differences are 40%, 15%, and 30%, respectively. In the NH in DJF, the
25 linear trend slope is positive (indicating a strengthening of the MMC) in all datasets at 440 K
26 (+0.9 to +2.5 %/decade in 1979–2012 and +3.4 %/decade in 1979–2001 for ERA-40) and in
27 all the datasets except for ERA-Interim and ERA-40 at 560 K (+0.9 to +4.0 %/decade). The
28 rate of increase is larger at 440 K than at 560 K in NCEP-CFSR, JRA-25, and JRA-55.

29 In the SH mid-latitudes in JJA, the mean MMC strength does not reveal any systematic
30 difference between the new and old datasets, whereas the standard deviation (i.e., year-to-year
31 variability) is smaller in the new datasets than in the old datasets by 10–30%. In all the
32 datasets except for ERA-Interim, the MMC tends to strengthen at 440 K; large trends in

1 1979–2012 are present in NCEP-NCAR, JRA-25, and JRA-55 (-5.4 to -8.4 %/decade). At 560
2 K, the MMC tends to weaken in ERA-Interim, JRA-25, and NCEP-CFSR (+1.1 to
3 +4.0 %/decade); the largest trend is thereby found in ERA-Interim (+4.0 %/decade). Both the
4 shallow and deep branches of BDC in the SH show strengthening trends in NCEP-NCAR,
5 ERA-40, and JRA-55.

6 The total upward mass flux was estimated by differencing the maximum and minimum values
7 of the mass stream function along the constant isentropic surfaces (Table 4 and Fig. 7). The
8 annual mean upward mass flux is smaller in the new datasets than in the old datasets (by 8–
9 30% at 440 K and 9–30% at 560 K), as similarly found in the MMC strength in the NH in
10 DJF. The MMC shows a strengthening trend in all the datasets except for NCEP-NCAR and
11 ERA-Interim at 440 K (+1.4 to +6.0 %/decade in 1979–2012 and +8.5 %/decade in 1979–
12 2001 for ERA-40) and in all datasets except for ERA-Interim at 560 K (+0.6 to
13 +5.3 %/decade). The trend is negative only in ERA-Interim at both altitudes (-3.9 %/decade at
14 440 K and -6.7 %/decade at 560 K). The estimated trends are statistically significant for the
15 new datasets both at 440 K and 560 K.

16 Several recent studies have pointed out that AoA and BDC changes occur over timescales of
17 several years to decades (Aschmann et al., 2014; Mahieu et al., 2014; Ray et al., 2014; Abalos
18 et al., 2015; Poelger et al., 2015b). As summarized in Table 5, the upward mass flux at 440 K
19 shows a strengthening trend during the first 22 years (1979–2000, +1.3 to +9.6 %/decade), but
20 a weakening trend (or slowing strengthening trend) during the last 12 years (2001–2012, -6.3
21 to +0.9 %/decade) at 440 K in all datasets except for ERA-Interim. ERA-Interim shows a
22 larger negative trend during the last 12 years (-6.0 %/decade) than in the first 22 years (-
23 3.0 %/decade). A strong strengthening (weakening) trend is found at 560 K for NCEP-NCAR
24 and JRA-25 (ERA-Interim).

25 **3.3 Wave decomposition**

26 To gain a better understanding of the relative contributions of the wave phenomena on various
27 scales to isentropic mixing, the zonal eddy PV flux is decomposed into individual zonal
28 wavenumber components s using Fast Fourier Transform (FFT) and then normalized by the
29 background meridional PV gradient as follow:

$$1 \quad K_{yy(s)} = \left(\frac{\left[\overline{v'_s q'_s} \right]_l}{\left[\frac{\partial \bar{q}}{a \partial \varphi} \right]_l} \right)_\theta. \quad (11)$$

2 The eddy PV flux is separated into three groups: planetary waves (associated with the zonal
 3 wavenumbers 1–3), synoptic-scale waves (zonal wavenumbers 4–7), and small-scale waves
 4 (zonal wavenumbers 8 and higher). For this analysis, l is set to 1 day to include short-term
 5 variations in eddy mixing, although this may cause apparent diffusion in the K_{yy} estimates.

6 The relative contribution of these groups is summarized in Table 6. The relative contribution
 7 of the planetary-scale mixing to the eddy flux is larger in the new datasets than in the old
 8 datasets for all the datasets at 440 K in the NH mid-latitudes, whereas that of zonal
 9 disturbances at both synoptic and small scales are somewhat smaller. With conventional data
 10 assimilation techniques such as 3D-VAR, physical consistency cannot be maintained during
 11 the data assimilation analysis because of lack of flow-dependent background error
 12 information, and this probably together with the lower forecast model resolution may cause
 13 spurious disturbances and excessive mixing, especially at small scales in the old datasets. For
 14 planetary-scale and synoptic-scale waves, forecast model configurations may also be
 15 important for the obtained differences (e.g., Biagio et al., 2014).

16 Each wave group reveals different long-term variations. The planetary-scale disturbance
 17 shows increasing trends in the NH surf zone at both 440 K (+0.6 to +4.2 %/decade in 1979–
 18 2012 and +4.8 %/decade in 1979–2001 for ERA-40) and 560 K (+0.6 to +11.5 %/decade in
 19 1979–2012 and +11.1 %/decade in 1979–2001 for ERA-40). At 560 K, all the groups reveal
 20 positive trends in all the datasets, with smaller linear trends in the new datasets in most cases,
 21 consistent with the result of the K_{yy} trend analysis except for ERA (c.f., Section 3.1), although
 22 the estimated trends are not statistically significant. In the subtropical lower stratosphere (not
 23 shown in table), as commonly found in the NH surf zone, the planetary-scale disturbance is
 24 stronger, and the small-scale disturbance is weaker in the new datasets than in the old datasets.

25 **3.4 Relative importance of eddy meridional transport**

26 Atmospheric wave breaking drives both the mean and eddy transports, but in a different
 27 manner; it drives the MMC below the breaking level through downward control (Haynes et al.,
 28 1991), whereas it causes eddy mixing at the level of wave breaking (e.g., Waugh et al., 1994).

1 Therefore, changes in MMC and eddy mixing do not necessarily occur in the same way in
2 response to changes in atmospheric wave forcing. These wave-induced processes may be
3 represented differently among the reanalysis products.

4 We compare the relative ratio of the eddy and total meridional PV fluxes (Eq. 10) to
5 investigate the relative importance of these two transport processes. As shown in Fig. 8, the
6 mean meridional transport is dominant equator-ward of about 40° throughout the year, and in
7 the extratropics during the summer. The eddy transport is dominant in the extratropics of the
8 winter hemisphere. Air masses entering the stratosphere are thus dominantly transported by
9 the mean-meridional transport from low to mid-latitudes and then transported mainly by the
10 eddy mixing from the mid- to high latitudes in the winter hemisphere. The strong westerly
11 coincides with the large contribution of the eddy transport. These characteristics are found in
12 a consistent way in all the datasets. The mean vertical transport may be pronounced in some
13 regions (e.g., inside the polar vortex), but the relative importance of vertical transport is not
14 evaluated in our analysis.

15 The 34-year mean value of the transport ratio is larger (i.e., the contribution of eddy transport
16 is larger) in the new datasets than in the old datasets both in the extratropics (by 5–16 % at
17 440 K and by 11–27 % at 560 K) and subtropics (by 5–14 %) in the NH (Table 7 and Fig. 9).
18 In the NH surf zone at 440 hPa, the trend in the transport ratio is different among the datasets.
19 At 560 hPa, the transport ratio shows a negative trend (i.e., the contribution of the mean
20 transport becomes more important over time) in all the datasets. In the SH surf zone, the trend
21 in the transport ratio varies among the dataset in 1979–2012 (-10.8 to +0.4 %/decade at 440 K
22 and -3.2 to +2.6 %/decade at 560 K, except ERA-40). The standard deviation is generally
23 smaller in the new datasets than in the old datasets in both hemispheres.

24 In the NH subtropics at 440 K, the large mean value of the transport ratio in the new datasets
25 when compared to the old datasets suggests that eddy-induced recirculation of old air masses
26 from the extratropics to the tropics is more effective than the mean poleward transport of
27 fresh air masses from the tropics into the extratropics in the new datasets, which could have
28 led to older AoA as derived for example in the study by Monge-Sanz et al. (2012), as
29 discussed further in Section 4.2. The transport ratio in the NH subtropics shows a positive
30 trend in most datasets except for NCEP-NCAR and NCEP-CFSR. In the SH subtropics, the
31 trend is negative (the mean transport tends to become more important) in most datasets except

1 for ERA-Interim. The positive trend in ERA-Interim corresponds to the increasing trend in
2 K_{yy} that only appeared in ERA-Interim and NCEP-NCAR.

3 The transport ratio in the NH subtropics reveals a larger positive trend in the last 12 years
4 (+3.5 to +6.8 %/decade) than in the first 22 years (-2.8 to +1.8 %/decade) in ERA-Interim and
5 NCEP-NCAR (Table 8). This positive trend implies that the eddy transport has become more
6 important over time with this tendency becoming even more important during the later time
7 period considered. On the other hand, the trend changes from positive to negative between the
8 two periods for NCEP-CFSR, JRA-25, and JRA-55. In the SH subtropics, the mean transport
9 becomes more important in all datasets except ERA-Interim in the earlier time period. The
10 trend in the later time period is strongly positive in ERA-Interim (+8.3 %/decade) and JRA-25
11 (+21.9 %/decade).

12

13 **4 Discussion**

14 **4.1 Dynamical consistency**

15 The relationship between the Eliassen–Palm (E–P) flux divergence and the MMC is expressed
16 through the downward control principle for steady-state conditions. The eddy PV flux in K_{yy}
17 can also be related to the mass flux and E–P flux under the quasi-geostrophic assumption
18 (Andrews et al., 1987; Schneider, 2005). Conventional data assimilation may degrade the
19 wave mean flow and wave mixing relationships. To evaluate the short-term dynamical
20 balance in the reanalysis products, we computed the temporal correlation between the E–P
21 flux divergence and the mass stream function, and between the E–P flux divergence and K_{yy}
22 ($l=1$ day). The E–P flux was estimated based on the MIM zonal mean momentum equation, in
23 which forcings by sub-grid processes were not considered. Momentum changes through sub-
24 grid processes such as gravity wave drag (GWD) parameterization need to be taken into
25 consideration for strict momentum budget analysis. However, these data were not provided in
26 all the datasets, and hence could not be considered in the analysis. The wave-driven
27 adjustment time for the wave mean flow interactions is typically several weeks in the lower
28 stratosphere (Haynes et al., 1991); this effect was also not considered. Instantaneous analysis
29 fields with no time lag were used for the correlation calculation. We here discuss the
30 relationship between the E–P flux divergence at each vertical level and the MMC at 440 K and
31 between the E–P flux divergence at each vertical level and K_{yy} ($l=1$ day) at 560 K. The MMC

1 at 440 K and K_{yy} at 560 K are driven by wave forcings at mostly the same vertical levels
2 (approximately between 500 and 650 K), as discussed below.

3 As shown in Fig. 10 for ERA-Interim, the mass stream function at 440 K at NH mid-latitudes
4 shows a large correlation with the E–P flux divergence at higher levels, because of downward
5 control. The correlation coefficient averaged between 500 and 650 K at the NH mid-latitudes
6 is larger in the new datasets (approximately 0.5–0.6) than in the old datasets (approximately
7 0.4) except for JRA (Table 9). This result suggests that the physical balance associated with
8 the wave mean flow interaction is represented more strictly in the new datasets for the NCEP
9 and ERA datasets. JRA-25 and JRA-55 both reveal large correlation coefficients
10 (approximately 0.6).

11 K_{yy} ($l=1$ day) at 560 K shows strong correlations with the E–P flux divergence around 500–
12 650 K in ERA-Interim (Fig. 10). This confirms that the reanalysis products represent local
13 wave forcings that drive eddy mixing. The correlation coefficient averaged between 550 and
14 650 K in the NH mid latitudes is larger in the new datasets than in the old datasets by 0.04–
15 0.07 (i.e., by 24–37 %, Table 9). This result suggests that the wave-mixing relationship is also
16 more accurately represented in the new datasets. Note that not only the dynamic balance in
17 analysis increments, but also characteristics of atmospheric diffusivity in the forecast model
18 (e.g., associated with choice of transport scheme and numerical diffusion) could influence the
19 wave-mixing relationship in the reanalysis products. Also note, that an alternative diagnosis
20 of eddy mixing and its trends using effective diffusivity based on ERA-Interim and JRA-55
21 has recently become available by Abalos et al. (2016).

22 **4.2 Implications for AoA long-term variations**

23 Based on analyses of observational data, several observational studies (e.g., Engel et al.,
24 2009; Stiller et al., 2012; Ray et al., 2014; Hegglin et al., 2014) revealed a positive AoA trend
25 at the NH mid-latitudes from the middle to upper stratosphere, but for different period (1975–
26 2005 in Engel et al. (2009), 2002–2010 in Stiller et al. (2012), 1975–2012 in Ray et al. (2014),
27 and 1986–2010 in Hegglin et al. (2014)). Hegglin et al. (2014) also found a reversed trend in
28 the NH lower stratosphere in 1986–2010. Note that the analysis by Hegglin et al. (2014) was
29 based on observed changes in stratospheric water vapour and not the typical age-tracers used
30 in the other studies (SF_6 and CO_2), and therefore may only be seen as an approximation of
31 age-of-air.

1 Using ERA-Interim, Monge-Sanz (2012) and Diallo et al. (2013) have demonstrated an AoA
2 decreasing trend in the NH lower stratosphere and an increasing trend in the middle
3 stratosphere in 1990–2009 and 1989–2010, respectively, consistent with the available
4 observational estimates. As discussed by Garny et al. (2014) and Ploeger et al. (2015a, 2015b),
5 it is essential to take the effects of mixing along the transport pathway into consideration, on
6 top of the effects of the MMC for understanding AoA variations. Interpreting AoA variations
7 simply in terms of changes in the local MMC and eddy mixing effects can be misleading,
8 because the AoA of a given air parcel is the result of the integrated effect of the various
9 tendencies along the parcel trajectory. Although the local analysis given in this study does not
10 allow for the air parcel history and transport pathway variations to be analyzed, it can provide
11 useful information on the general structure of the transport intensity and its potential impacts
12 on AoA variations.

13 To investigate whether any reanalysis data have the potential to reveal useful implications of
14 long-term AoA variations, we summarize long-term variations in the three important transport
15 processes: the tropical upward mass flux, the relative contribution of eddy mixing in the
16 subtropics, and the mid-latitudes mean poleward motions ($\overline{v^*}$). We consider that, rather than
17 the mixing strength in the subtropics itself, the relative ratio of the mean and eddy transport is
18 essential to understanding AoA variations, since it determines the ratio of fresh air entering
19 and old air recirculating between the tropics and extratropics. Eddy mixing across the MMC
20 overturning may recirculate old air masses from the extratropics to the tropics, whereas mean
21 poleward flows carry fresh air masses from the tropics to the extratropics, and may return old
22 air masses that were recirculated from the extratropics by eddy transport. Here, we focus on
23 transport processes during winter to explain their possible influences on annual mean AoA
24 variations. In winter, atmospheric waves are active, and strongly induce both MMC and eddy
25 mixing. Abalos et al. (2015) showed that the DJF trends make a major contribution to the
26 overall structure of the annual mean trends of BDC in the NH. However, the influence of
27 transport processes in other seasons cannot be neglected (Konopka et al., 2015). Therefore,
28 the implication obtained from the local analysis of the wintertime transport processes in this
29 study is limited. Explicit calculations of AoA using the reanalysis datasets are required to
30 provide further insights into the role of each transport process on AoA variations.

31 As shown by Fig. 11, during the 34 years, from 1979 to 2012, the tropical upward mass flux
32 shows a positive trend from the lower to middle stratosphere, except for ERA-Interim in DJF

1 (consistent with the results of Seviour et al. (2011) and Abalos et al. (2015)), and NCEP-
2 NCAR and ERA-Interim in JJA (Fig. 11). The relative importance of the eddy transport
3 compared to the mean transport shows an increasing trend in ERA-Interim and JRA-55 from
4 the lower to middle stratosphere, which may act to increase AoA, especially in the BDC deep
5 branch. The value of $\overline{v^*}$ at the NH mid-latitudes shows a strengthening trend in the lower
6 stratosphere in most datasets, except for NCEP-CFSR below 440 K. The $\overline{v^*}$ trend varies
7 largely with height in ERA-Interim, showing a sharp positive trend peak around 400–430 K,
8 and a negative trend above 520 K. To summarize, AoA derived using the reanalysis products
9 in the NH middle stratosphere for the 34 years can be expected to increase in ERA-Interim,
10 and probably also in JRA-55, because of the large increase in the contribution of the eddy
11 mixing in the subtropics and the weakened mean poleward motion in the deep branch. AoA in
12 the NH lower stratosphere can be expected to decrease in all the datasets, except in NCEP-
13 CFSR.

14 In the SH, the relative importance of the eddy transport to the mean transport in the subtropics
15 shows an increasing trend only in ERA-Interim in the lower stratosphere, and in ERA-Interim
16 and NCEP-NCAR in the middle stratosphere (above 460 K). The value of $\overline{v^*}$ at the SH mid-
17 latitudes shows a strengthening trend in all the datasets, except for ERA-Interim above 500 K.
18 These changes suggest the possibility of an AoA decreasing trend for both the lower and
19 middle stratosphere in the SH in most datasets except for ERA-Interim during the 34 years.
20 The analysis of ERA-Interim may suggest the possibility of a weaker decreasing or a weak
21 positive AoA trend in the middle stratosphere.

22 The decadal scale variation in the transport processes will also cause changes in the AoA
23 trend on similar timescales. In the last 12 years (Fig. 12), the upward mass flux trend becomes
24 negative in the lower stratosphere in most datasets, and shows larger increasing trends in the
25 middle stratosphere in all the datasets in DJF. The contribution of the eddy transport in the
26 NH subtropics becomes even more important in the last 12 years in ERA-Interim. The value
27 of $\overline{v^*}$ at the NH mid-latitudes tends to weaken in most datasets, except for ERA-Interim
28 below 430 K, and all the datasets above that level in the last 12 years. Because of the larger
29 negative trend in $\overline{v^*}$ and the increased contribution of the eddy transport in the subtropics, a
30 larger increase in AoA in the NH is expected to be derived using ERA-Interim during the last
31 12 years than during the 34 years, as suggested by Ploeger et al. (2015b).

1 In the SH, ERA-Interim reveals a large decreasing contribution of the eddy transport in the
2 last 12 years, which may indicate a decrease in AoA in the SH stratosphere in the last 12
3 years, as consistently revealed by Stiller et al. (2012) for the 2002–2010 period. JRA-25
4 reveals substantial decadal scale changes in the transport processes in both hemispheres

5 **4.3 Implications for future developments of reanalyses**

6 Although the reanalysis systems have been individually updated at each operation center,
7 similar aspects were encountered regarding the effects of the system updates on the MMC and
8 eddy mixing, as follows:

- 9 1. Both the BDC shallow and deep branches reveal weaker MMC mean intensity (by 15–
10 60% in the NH) and smaller interannual variability (by 10–60% in the SH) in the new
11 datasets than in the old datasets. This tendency generally will tend to increase AoA
12 throughout the NH stratosphere.
- 13 2. The contribution of planetary-scale mixing to eddy mixing was stronger in the new
14 datasets than in the old datasets in the NH lower stratosphere, whereas that of small-scale
15 mixing was smaller. The weaker small-scale mixing in the new datasets may be a result of
16 reduced spurious eddies associated with analysis increments using the flow-dependent
17 analysis (but not in NCEP-CFSR) and the use of the higher forecast model resolution.
- 18 3. The relative importance of the eddy transport to the mean transport in the NH is larger in
19 the new datasets by 5–27% in the extratropics, and by 5–14% in the subtropics. The larger
20 eddy contribution in the subtropical lower stratosphere may cause an older AoA in the
21 entire stratosphere in the new datasets.
- 22 4. The wave mean flow relationship and the wave-mixing relationship in the NH surf zone
23 are more accurately represented in the new datasets for the NCEP and ERA datasets.

24 Updated systems can be expected to provide better representations of both the MMC and eddy
25 mixing, because of reduced systematic errors of the forecast model, the application of
26 improved bias correction algorithms, and advanced data assimilation techniques. Nevertheless,
27 it is not straightforward to identify which changes are mainly responsible for the differences
28 in the MMC and eddy mixing.

29 Monge-Sanz et al. (2007, 2012) investigated that ERA-Interim benefits using an omega-
30 equation balance operator in the background constraint, which is likely to have reduced

1 spurious propagation of eddy motion associated with analysis increments. They also
2 suggested other important factors, such as reduced stratospheric temperature biases of the
3 forecast model and improved bias corrections for satellite radiance data. In ERA-40 and JRA-
4 25, systematic analysis increments were introduced to compensate for model temperature
5 biases, which are thought to cause an overly strong BDC (Uppala et al., 2005). Kobayashi and
6 Iwasaki et al. (2015) found that in JRA-25, the temperature analysis increment evolves from
7 the 1990s to the 2000s, associated with changes in forecast model biases and lack of effective
8 bias corrections, either for the forecast model and assimilated measurements. As a result, the
9 MMC structure in JRA-25 was unrealistically distorted in a particular period corresponding to
10 the changes in external forcing.

11 The decadal scale variations of the transport processes in the reanalysis could be introduced
12 artificially because of measurement discontinuity. Stratospheric temperatures in ERA-Interim
13 are known to be affected by the introduction of AMSU-A data in 1998 and radio occultation
14 data at the end of 2006 (Dee et al., 2011). Improved bias correction schemes have been
15 implemented in recent reanalysis systems (e.g., variational bias correction in JRA-55
16 (Kobayashi et al., 2015) and ERA-Interim (Dee and Uppala, 2009)). However, it is still
17 difficult to distinguish between forecast models and measurements biases, and to correct them
18 properly.

19 In addition, the reanalysis quality could be strongly affected by the performance of the
20 forecast model. Any changes in wave propagation and breaking lead to changes in both the
21 MMC and mixing, but current models have large uncertainties in representing various waves,
22 including forcings from GWD parameterization (e.g., Sigmond and Shepherd, 2014). Further
23 efforts on model development are still important.

24 **5 Conclusions**

25 We compared the characteristics of the stratospheric MMC and eddy mixing in the six
26 reanalysis products for the period 1979–2012 based on mass-weighted isentropic zonal means.
27 Both of the mean and eddy transport processes play important roles in determining
28 distributions and variations of chemical tracers and AoA, thus it is important to clarify
29 whether there are any reanalysis data that can be used to investigate long-term BDC variations.

30 The mean K_{yy} value differs largely among the data products both in the extratropics and
31 subtropics. The contribution of planetary-scale mixing in the NH lower stratosphere is larger
32 in the new datasets, whereas that of small-scale mixing is smaller. The weaker small-scale

1 mixing in the new datasets is considered to be a result of reduced spurious eddies associated
2 with analysis increments in the 4D-VAR analysis. Isentropic mixing shows a strengthening
3 trend in the NH middle stratosphere surf zone in all datasets, associated with strengthened
4 large-scale mixing.

5 There are large differences in the MMC strength among the datasets, but are were some
6 similar characteristics. In the NH, the MMC is stronger in the new datasets (NCEP-CFSR,
7 ERA-Interim, and JRA-55) than in the old datasets (NCEP-NCAR, ERA-40, and JRA-25).
8 All the datasets show a strengthening trend in the BDC shallow branch in the winter
9 hemispheres, consistent with model simulations (e.g., Butchart et al., 2010). The MMC in the
10 BDC deep branch showed a weakening trend only in ERA-Interim at the NH mid-latitudes
11 and in ERA-Interim and JRA-25 in the SH mid-latitudes in winter. The vertical structure in
12 the changes as seen in ERA-Interim that would lead to a decrease in AoA in the lower
13 stratosphere and an increase in AoA in the middle (and upper) stratosphere are broadly
14 consistent with inferences made from changes observed in stratospheric water vapour
15 (Hegglin et al., 2014).

16 We also evaluated the relative importance of the mean and eddy transports. The contribution
17 of eddy mixing was generally stronger in the new datasets than in the old datasets in the NH.
18 In the subtropical lower stratosphere, the relative contribution of eddy transport showed an
19 increasing trend in datasets, except for NCEP-NCAR and NCEP-CFSR in the NH, and only in
20 ERA-Interim in the SH. These trends reveal changes in the relative effectiveness of eddy-
21 induced recirculation from the extratropics to the tropics, and the mean poleward transport of
22 fresh air into the extratropics; they are considered to be important in understanding AoA
23 variations.

24 The transport analysis suggests that the ERA-Interim provides a consistent transport picture,
25 with AoA trends derived based on observations in both the deep and shallow branches,
26 whereas other datasets would provide different implications. The increasing trend in AoA in
27 the NH middle stratosphere can be understood as a result of the weakening MMC in the deep
28 branch (found only in ERA-Interim), together with the increased contribution of eddy
29 transport compared with the mean transport in the subtropical lower and middle stratosphere
30 (found in ERA-Interim and JRA-55). The decreasing trend in AoA in the SH lower
31 stratosphere could be a result of the strengthening MMC trends in the shallow branch (large in
32 ERA-Interim and JRA-25). All the reanalysis datasets revealed decadal scale variations in the

1 strength of both MMC and eddy mixing, which may result in significant AoA trend variations.
2 For instance, the increased contribution of the eddy transport in the subtropics and the
3 weakened mean poleward motion in the middle stratosphere during the last 12 years (1979–
4 2000) compared to the first 22 years (2001–2012) in the NH suggests larger increasing trends
5 in AoA in ERA-Interim and NCEP-NCAR in the last 12 years.

6 Differences in data assimilation schemes and forecast models are thought to cause differences
7 in the expression of the MMC and eddy mixing in the BDC among reanalysis products. Our
8 analysis suggests that advanced reanalysis products are potentially useful for studying long-
9 term BDC variations, because of the flow-dependent background error covariance,
10 assimilation of the observations at the exact time, balance operators, and improved bias
11 correction algorithms used in these reanalyses. However, there still seem to be problems that
12 cause unrealistic atmospheric variations (e.g., large differences in long-term variations
13 between ERA-Interim and JRA-55) associated with discontinuities in the assimilated
14 measurements and large uncertainties in the forecast models. Further efforts are essential to
15 improve the reanalysis systems so that the reanalysis products become more consistent among
16 each other and that they can be used to study the details in BDC variations.

17 **Acknowledgements**

18 We would like to thank Felix Ploeger for his helpful comments on this study. We also would
19 like to thank the editor and two anonymous reviewers for their valuable comments. The work
20 was supported by Grant-in Aid for Scientific Research 15K05296 and 26287117 of MEXT,
21 Japan.

1 References

- 2 Abalos, M., Randel, W. J., Kinnison, D. E., and Serrano, E.: Quantifying tracer transport in
3 the tropical lower stratosphere using WACCM, *Atmos. Chem. Phys.*, 13, 10591-10607,
4 doi:10.5194/acp-13-10591-2013, 2013.
- 5 Abalos, M., B. Legras, F. Ploeger and W.J. Randel: Evaluating the advective Brewer-Dobson
6 circulation in three reanalyses for the period 1979-2012. *J. Geophys. Res.*, 120,
7 doi:10.1002/2015JD023182, 2015.
- 8 Abalos, M., Legras, B. and Shuckburgh, E., Interannual variability in effective diffusivity in
9 the upper troposphere/lower stratosphere from reanalysis data, *Q. J. R. Meteorol. Soc.*.
10 doi: 10.1002/qj.2779, 2016.
- 11 Andrews, D. G., and M. E. McIntyre: Planetary waves in horizontal and vertical shear: The
12 generalized Eliassen–Palm relation and mean zonal acceleration. *J. Atmos. Sci.*, 33, 2031–
13 2048, 1976.
- 14 Andrews, D.G. and M.E. McIntyre: An exact theory of nonlinear waves on a Lagrangian-
15 mean flow. *J. Fluid. Mech.*, 89, 609-646, 1978.
- 16 Andrews, D.G.: A finite-amplitude Eliassen-Palm theorem in isentropic coordinates. *J. Atmos.*
17 *Sci.*, 40, 1877-1883, 1983.
- 18 Andrews, D. G., J. R. Holton, and C. B. Leovy: *Middle Atmosphere Dynamics*. Academic
19 Press, Orlando, 489~pp., 1987.
- 20 Aschmann, J., Burrows, J. P., Gebhardt, C., Rozanov, A., Hommel, R., Weber, M., and
21 Thompson, A. M.: On the hiatus in the acceleration of tropical upwelling since the beginning
22 of the 21st century, *Atmos. Chem. Phys.*, 14, 12803-12814, doi:10.5194/acp-14-12803-2014,
23 2014.
- 24 Bartels, J., D. Peters, and G. Schmitz: Climatological Ertel’s potential-vorticity flux and mean
25 meridional circulation in the extratropical troposphere–lower stratosphere. *Ann. Geophys.*, 16,
26 250–265, 1998.
- 27 Birner, T. and H. Bönisch: Residual circulation trajectories and transit times into the
28 extratropical lowermost stratosphere. *Atmos. Chem. Phys.*, 11, 817–827, 2011.

1 Biagio, V., S. Calmanti, A. Dell'Aquila, and P. M. Ruti: Northern Hemisphere winter
2 midlatitude atmospheric variability in CMIP5 models, *Geophys. Res. Lett.*, 41, 1277–1282,
3 doi:10.1002/2013GL058928, 2014.

4 Brewer, A. W.: Evidence for a world circulation provided by measurements of helium and
5 water vapour distribution in the stratosphere, *Q. J. R. Meteorol. Soc.*, 75, 351–363,
6 doi:10.1002/qj.49707532603, 1949.

7 Butchart, N., Cionni, I., Eyring, V., Shepherd, T. G., Waugh, D. W., Akiyoshi, H., Austin, J.,
8 Bruhl, C., Chipperfield, M. P., Cordero, E., Dameris, M., Deckert, R., Dhomse, S., Frith, S.
9 M., Garcia, R. R., Gettelman, A., Giorgetta, M. A., Kinnison, D. E., Li, F., Mancini, E.,
10 McLandress, C., Pawson, S., Pitari, G., Plummer, D. A., Rozanov, E., Sassi, F., Scinocca, J.
11 F., Shibata, K., Steil, B., and Tian, W.: Chemistry-Climate Model simulations of twenty-first
12 century stratospheric climate and circulation changes, *J. Climate*, 23, 5349–5374,
13 doi:10.1175/2010JCLI3404.1, 2010.

14 Dee, D. P. and Uppala, S.: Variational bias correction of satellite radiance data in the ERA-
15 Interim reanalysis, *Q. J. Roy. Meteor. Soc.*, 135, 1830–1841, doi: 10.1002/qj.493, 2009.

16 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U.,
17 Balmaseda, M. A., Balsamo, G., Bauer, P. Bechtold, P., Beljaars, A. C. M., van de Berg, L.,
18 Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L.,
19 Healy, S. B., Hersbach, H., Ho'lm, E. V., Isaksen, L., Ka'llberg, P., Ko'hler, M., Matricardi,
20 M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.- K., Peubey, C., de
21 Rosnay, P., Tavolato, C., The'paut, J.-N., and Vitart, F.: The ERA-Interim reanalysis:
22 configuration and perfor- mance of the data assimilation system, *Q. J. Roy. Meteor. Soc.*, 137,
23 553–597, doi:10.1002/qj.828, 2011.

24 Diallo, M., B. Legras, and A. Chedin: Age of stratospheric air in the ERA-Interim, *Atmos.*
25 *Chem. Phys.*, 12, 12,133–12,154, doi:10.5194/ acp-12-12133-2012, 2012.

26 Dobson GMB, Harrison DN, Lawrence J.: Measurements of the amount of ozone in the
27 Earth's atmosphere and its relation to other geophysical conditions. *Proc. Roy. Soc., Ser. A*,
28 122, 456--486, 1929.

29 Dobson, G. M. B.: Origin and distribution of polyatomic molecules in the atmosphere, *Proc.*
30 *Roy. Soc., Ser. A*, 236, 187, 1956.

1 Engel, A., Möbius, T., Böning, H., Schmidt, U., Heinz, R., Levin, I., Atlas, E., Aoki, S.,
2 Nakazawa, T., Sugawara, S., Moore, F., Hurst, D., Elkins, J., Schauffler, S., Andrews, A., and
3 Boering, K.: Age of stratospheric air unchanged within un- certainties over the past 30 years,
4 *Nature Geosci.*, 2, 28–31, doi:10.1038/ngeo388, 2009.

5 Fueglistaler, S., Legras, B., Beljaars, A., Morcrette, J.-J., Sim- mons, A., Tompkins, A. M.,
6 and Uppala, S.: The diabatic heat budget of the upper troposphere and lower/mid stratosphere
7 in ECMWF reanalyses, *Q. J. Roy. Meteor. Soc.*, 135, 21–37, doi:10.1002/qj.361, 2009.

8 Garcia, R. R. and Randel, W. J.: Acceleration of Brewer-Dobson circulation due to the
9 increases in greenhouse gases, *J. Atmos. Sci.*, 65, 2731–2739, doi:10.1175/2008JAS2712.1,
10 2008.

11 Garny, H., T. Birner, H. Böning, and F. Bunzel: The effects of mixing on age of air, *J.*
12 *Geophys. Res.*, 119, 7015–7034, doi:10.1002/2013JD021417, 2014.

13 Haynes, P., and E. Shuckburgh: Effective diffusivity as a diagnostic of atmospheric transport:
14 1. Stratosphere, *J. Geophys. Res.*, 105, 22777–22794, doi:10.1029/2000JD900093, 2000.

15 Haynes, P., C. J. Marks, M. E. McIntyre, T. G. Shepherd, and K. P. Shine: On the
16 “downward control” of extratropical diabatic circulations by eddy-induced mean zonal
17 forces, *J. Atmos. Sci.*, 48, 651–678, 1991.

18 Hegglin, M. I., D. Plummer, J. Scinocca, T. G. Shepherd, J. Anderson, L. Froidevaux, B.
19 Funke, D. Hurst, A. Rozanov, J. Urban, T. v. Clarmann, K. A. Walker, R. Wang, S. Tegtmeier,
20 and K. Weigel, Variation of stratospheric water vapour trends with altitude from merged
21 satellite data, *Nat. Geosci.*, 7, 768–776, doi: 10.1038/NGEO2236, 2014.

22 Hoskins, B. J., M. E. McIntyre, and A. W. Robertson: On the use and significance of
23 isentropic potential vorticity maps, *Quart. J. Roy. Meteor. Soc.*, 111, 877–946, 1985.

24 Iwasaki, T.: A diagnostic formulation for wave–mean flow interactions and Lagrangian-mean
25 circulation with a hybrid vertical coordinate of pressure and isentropes, *J. Meteor. Soc. Japan*,
26 67, 293–312, 1989.

27 Iwasaki, T.: General circulation diagnosis in the pressure-isentrope hybrid vertical coordinate.
28 *J. Meteor. Soc. Japan*, 70, 673–687, 1992.

1 Iwasaki, T., Hamada, H., and Miyazaki, K.: Comparisons of Brewer-Dobson circulations
2 diagnosed from reanalyses, *J. Meteorol. Soc. Japan*, 87, 997–1006, doi:10.2151/jmsj.87.997,
3 2009.

4 Johnson, D.R.: A Generalized Transport Equation for Use with Meteorological Coordinate
5 Systems. *Mon. Wea. Rev.*, 108, 733–745, 1980.

6 Kalnay, E., M. Kanamitsu, R. Kistler, W. Collins, D. Deaven, L. Gandin, M. Iredell, S. Saha,
7 G. White, J. Woollen, Y. Zhu, A. Leetmaa, R. Reynolds, M. Chelliah, W. Ebisuzaki, W.
8 Higgins, J. Janowiak, K. C. Mo, C. Ropelewski, J. Wang, R. Jenne, D. Joseph: The
9 NCEP/NCAR 40-Year Reanalysis Project, *Bull. Am. Meteor. Soc.*, 77, 437–472, 1996.

10 Kawatani, Y., Hamilton, K., Miyazaki, K., Fujiwara, M., and Anstey, J.: Representation of the
11 Tropical Stratospheric Zonal Wind in Global Atmospheric Reanalyses, *Atmos. Chem. Phys.*
12 *Discuss.*, doi:10.5194/acp-2016-76, in review, 2016.

13 Kendall, M.G., *Rank Correlation Methods*, 4th edition, Charles Griffin, London, 1975.

14 Kim, Y.-H. and Chun, H.-Y.: Momentum forcing of the quasi-biennial oscillation by
15 equatorial waves in recent reanalyses, *Atmos. Chem. Phys.*, 15, 6577–6587, doi:10.5194/acp-
16 15-6577-2015, 2015.

17 Kobayashi, S., Y. Ota, Y. Harada, A. Ebita, M. Moriya, H. Onoda, K. Onogi, H. Kamahori, C.
18 Kobayashi, H. Endo, K. Miyaoka, and K. Takahashi: The JRA-55 Reanalysis: General
19 Specifications and Basic Characteristics, *J. Meteor. Soc. Japan*, 93, 5–48,
20 doi:10.2151/jmsj.2015-001, 2015.

21 Kobayashi, C. and T. Iwasaki, Brewer-Dobson circulation diagnosed from JRA-55, *J.*
22 *Geophys. Res.*, revised, 2015.

23 Konopka, P., Ploeger, F., Tao, M., Birner, T. and Riese, M.: Hemispheric asymmetries and
24 seasonality of mean age of air in the lower stratosphere: Deep versus shallow branch of the
25 Brewer-Dobson circulation, *J. Geophys. Res.*, 120, 2053–2066. doi: 10.1002/2014JD022429,
26 2015.

27 Lyjak, L. V., and V. A. Yudin: Diagnostics of the large-scale mixing properties from
28 stratospheric analyses, *J. Geophys. Res.*, 110, D17107, doi:10.1029/2004JD005577, 2005.

29 Mahieu, E., Chipperfield, M. P., Notholt, J., Reddmann, T., Anderson, J., Bernath, P. F.,
30 Blumenstock, T., Coffey, M. T., Dhomse, S. S., Feng, W., Franco, B., Froidevaux, L., Griffith,

1 D. W. T., Hannigan, J. W., Hase, F., Hossaini, R., Jones, N. B., Morino, I., Murata, I.,
2 Nakajima, H., Palm, M., Paton-Walsh, C., Russell III, J. M., Schneider, M., Servais, C.,
3 Smale, D. & Walker, K. A.: Recent Northern Hemisphere stratospheric HCl increase due to
4 atmospheric circulation changes, *Nature*, 515, 104–107, 2014.

5 Mann, H.B., Non-parametric tests against trend, *Econometrica* 13, 163–171, 1945.

6 McIntyre, M. E., and T. N. Palmer: Breaking planetary waves in the stratosphere, *Nature*, 305,
7 593–600, doi:10.1038/305593a0, 1983.

8 Miyazaki, K., and T. Iwasaki: Diagnosis of meridional ozone transport based on mass-
9 weighted isentropic zonal means, *J. Atmos. Sci.*, 63, 1192–1208, 2005.

10 Miyazaki, K., and T. Iwasaki: The gradient genesis of stratospheric trace species in the
11 subtropics and around the polar vortex, *J. Atmos. Sci.*, 65, 490–508, 2008.

12 Miyazaki, K., S. Watanabe, Y. Kawatani, Y. Tomikawa, M. Takahashi, and K. Sato:
13 Transport and mixing in the extratropical tropopause region in a high-vertical-resolution
14 GCM. Part I: Potential vorticity and heat budget analysis, *J. Atmos. Sci.*, 67, 1293–1314,
15 2010a.

16 Miyazaki, K., Sato, S. Watanabe, Y. Tomikawa, Y. Kawatani, and M. Takahashi: Transport
17 and mixing in the extratropical tropopause region in a high-vertical-resolution GCM. Part II:
18 Importance of small-scale disturbances, *J. Atmos. Sci.*, 67, 1315–1336, 2010b.

19 Monge-Sanz, B. M., Chipperfield, M. P., Simmons, A. J., and Uppala, S. M.: Mean age of air
20 and transport in a CTM: comparison of different ECMWF analyses, *Geophys. Res. Lett.*, 340,
21 L04801, doi:10.1029/2006GL028515, 2007.

22 Monge-Sanz, B. M., Chipperfield, M. P., Dee, D. P., Simmons, A. J., and Uppala, S. M.:
23 Improvements in the stratospheric transport achieved by a chemistry transport model with
24 ECMWF (re)analyses: identifying effects and remaining challenges, *Q. J. R. Meteorol. Soc.*,
25 139, 654–673, doi:10.1002/qj.1996, 2012.

26 Nakamura, N.: Two-dimensional mixing, edge formation, and permeability diagnosed in area
27 coordinates, *J. Atmos. Sci.*, 53, 1524–1537, 1996.

28 Nakamura, N.: Quantifying inhomogeneous, instantaneous, irreversible transport using
29 passive tracer field as a coordinate. *Transport and Mixing in Geophysical Flows*, J. B. Weiss
30 and A. Provenzale, Eds., *Lecture Notes in Physics*, Vol. 744, Springer-Verlag, 137–164, 2008

1 Newman, P. A., M. R. Schoeberl, R. A. Plumb, and J. E. Rosenfield: Mixing rates calculated
2 from potential vorticity, *J. Geophys. Res.*, 93, 5221–5240, 1988.

3 Onogi, K., J. Tsutsui, H. Koide, M. Sakamoto, S. Kobayashi, H. Hatsushika, T. Matsumoto, N.
4 Yamazaki, H. Kamahori, K. Takahashi, S. Kadokura, K. Wada, K. Kato, R. Oyama, T. Ose, N.
5 Mannoji and R. Taira: The JRA-25 Reanalysis, *J. Meteor. Soc. Japan*, 85, 369–432,
6 doi:10.2151/jmsj.85.369, 2007.

7 Pawson, S., and M. Fiorino: A comparison of reanalyses in the lower tropical stratosphere:
8 Part 2. The quasi-biennial oscillation, *Clim. Dyn.*, 14, 645–658, 1998.

9 Ploeger, F., M. Abalos, T. Birner, P. Konopka, B. Legras, R. Müller, and M. Riese:
10 Quantifying the effects of mixing and residual circulation on trends of stratospheric mean age
11 of air. *Geophys. Res. Lett.*, 42, 2047–2054. doi: 10.1002/2014GL062927, 2015a.

12 Ploeger, F., M. Riese, F. Haenel, P. Konopka, R. Müller, and G. Stiller: Variability of
13 stratospheric mean age of air and of the local effects of residual circulation and eddy mixing,
14 *J. Geophys. Res.*, 120, 716–733, doi:10.1002/2014JD022468, 2015b.

15 Plumb, R. A., A “tropical pipe” model of stratospheric transport, *J. Geophys. Res.*, 101,
16 3957–3972, 1996.

17 Randel, W. J., R. Garcia, and F. Wu: Dynamical balances and tropical stratospheric upwelling,
18 *J. Atmos. Sci.*, 65, 3584–3595, doi:10.1175/2008JAS2756.1, 2008.

19 Ray, E. A., Moore, F. L., Rosenlof, K. H., Davis, S. M., Boenisch, H., Morgenstern, O.,
20 Smale, D., Rozanov, E., Hegglin, M., Pitari, G., Mancini, E., Braesicke, P., Butchart, N.,
21 Hardiman, S., Li, F., Shibata, K., and Plummer, D. A.: Evidence for changes in stratospheric
22 transport and mixing over the past three decades based on multiple data sets and tropical
23 leaky pipe analysis, *J. Geophys. Res.*, 115, D21304, doi:10.1029/2010JD014206, 2010.

24 Ray, E. A., F. L. Moore, K. H. Rosenlof, S. M. Davis, C. Sweeney, P. Tans, T. Wang, J. W.
25 Elkins, H. Boenisch, A. Engel, S. Sugawara, T. Nakazawa, and S. Aoki: Improving
26 stratospheric transport trend analysis based on SF₆ and CO₂ measurements, *J. Geophys. Res.*,
27 119, 14,110–14,128, doi:10.1002/2014JD021802, 2014.

28 Saha, S., Moorthi, S., Wu, X., Wang, J., Nadiga, S., Tripp, P., Behringer, D., Hou, Y.-T.,
29 Chuang, H.-Y., Iredell, M., Ek, M., Meng, J., Yang, R., Peoa, M., van den Dool, M., Zhang,

1 Q., Wang, W., Chen, M., and Becker, M.: The NCEP climate forecast system version~2, *J.*
2 *Climate*, 27, 2185–2208, 2013.

3 Scheele, M. P., Siegmund, P. C., and Velthoven, P. F. J.: Stratospheric age of air computed
4 with trajectories based on various 3D-Var and 4D-Var data sets, *Atmos. Chem. Phys.*, 5, 1–7,
5 doi:10.5194/acp-5-1-2005, 2005.

6 Schneider, T.: Zonal momentum balance, potential vorticity dynamics, and mass fluxes on
7 near-surface isentropes, *J. Atmos. Sci.*, 62, 1884–1900, 2005.

8 Schoeberl, M. R., Douglass, A. R., Zhu, Z., and Pawson, S.: A comparison of the lower
9 stratospheric age spectra derived from a general circulation model and two data assimilation
10 systems, *J. Geophys. Res.*, 108, 4113, doi:10.1029/2002JD002652, 2003.

11 Seviour, W. J. M., Butchart, N., and Hardiman, S. C.: The Brewer-Dobson circulation
12 inferred from ERA-Interim, *Q. J. Roy. Meteor. Soc.*, 138, 878–888, doi:10.1002/qj.966, 2011.

13 Shepherd, T. G.: The middle atmosphere, *J. Atmos. Solar-Terr. Phys.*, 62, 1587-1601, doi:
14 10.1016/S1364-6826(00)00114-0, 2000.

15 Shepherd T. G., McLandress, C.: A robust mechanism for strengthening of the Brewer–
16 Dobson circulation in response to climate change: Critical-layer control of subtropical wave
17 breaking, *J. Atmos. Sci.*, 68, 784–797, 2011.

18 Sigmond, M., T. Shepherd: Compensation between Resolved Wave Driving and
19 Parameterized Orographic Gravity Wave Driving of the Brewer–Dobson Circulation and Its
20 Response to Climate Change, *J. Climate*, 27, 5601–5610, 2014.

21 Simmons, A., Uppala, S., Dee, D., Kobayashi, S.: ERA-Interim: New ECMWF reanalysis
22 products from 1989 onwards, *ECMWF newsletter*, 110, 25–35, 2007.

23 Stiller, G. P., von Clarmann, T., Haenel, F., Funke, B., Glatthor, N., Grabowski, U., Kellmann,
24 S., Kiefer, M., Linden, A., Lossow, S., and López-Puertas, M.: Observed temporal evolution
25 of global mean age of stratospheric air for the 2002 to 2010 period, *Atmos. Chem. Phys.*, 12,
26 3311–3331, doi:10.5194/acp-12-3311- 2012, 2012.

27 Tan, W., M. A. Geller, S. Pawson, and A. M. da Silva: A case study of excessive subtropical
28 transport in the stratosphere of a data assimilation system, *J. Geophys. Res.*, 109, D11102,
29 doi:10.1029/2003JD004057, 2004.

1 Tanaka, D., T. Iwasaki, S. Uno, M. Ujiie and K. Miyazaki, Eliassen-Palm flux diagnosis
2 based on isentropic representation, *J. Atmos. Sci.*, 61 2370-2383, 2004.

3 Tung, K.K.: On the two dimensional transport of stratospheric trace gases in isentropic
4 coordinates. *J. Atmos. Sci.*, 39, 2330-2355, 1982.

5 Tung, K.K.: On the relationship between the thermal structure of the stratosphere and the
6 seasonal distribution of ozone, *Geophys. Res. Lett.*, 13, 1308–1311, doi:
7 10.1029/GL013i012p01308, 1986.

8 Waugh, D., R. A. Plumb, R. J. Atkinson, M. R. Schoeberl, L. R. Lait, P. Newman, M.
9 Loewenstein, D. Toohey, L. Avallone, C. R. Webster, and R. May: Transport out of the lower
10 stratospheric Arctic vortex by Rossby wave breaking, *J. Geophys. Res.*, 99, 1071–1088,
11 doi:10.1029/93JD02556, 1994.

12 Waugh, D.W., and T.M. Hall: Age of stratospheric air: Theory, observations, and models,
13 *Rev. Geophys.* , 40, 4, doi:10.1029/2000RG000101, 2002.

14 Wohltmann, I. and Rex, M.: Improvement of vertical and residual velocities in pressure or
15 hybrid sigma-pressure coordinates in analysis data in the stratosphere, *Atmos. Chem. Phys.*, 8,
16 265-272, doi:10.5194/acp-8-265-2008, 2008.

17 Wright, J. S. and Fueglistaler, S.: Large differences in reanalyses of diabatic heating in the
18 tropical upper troposphere and lower stratosphere, *Atmos. Chem. Phys.*, 13, 9565-9576,
19 doi:10.5194/acp-13-9565-2013, 2013.

20 Uppala, S., Kallberg, P., Simmons, A., Andrae, U., da Costa Bechtold, V., Fiorino, M.,
21 Gibson, J., Haseler, J., Hernandez, A., Kelly, G., Li, X., Onogi, K., Saarinen, S., Sokka, N.,
22 Allan, R., Andersson, E., Arpe, K., Balmaseda, M., Beljaars, A., van de Berg, L., Bidlot, J.,
23 Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher, M., Fuentes, M.,
24 Hagemann, S., Holm, E., Hoskins, B., Isaksen, L., Janssen, P., Jenne, R., McNally, A.,
25 Mahfouf, J.-F., Morcrette, J.-J., Rayner, N., Saunders, R., Simon, P., Sterl, A., Trenberth, K.,
26 Untch, A., Vasiljevic, D., Viterbo, P., and Woollen, J.: The ERA-40 re-analysis, *Q. J. Roy.
27 Meteor. Soc.*, 131, 2961–3012, doi:10.1256/qj.04.176, 2005.

28

1 Table 1. Mean value (Mean) over 34 years (1979–2012) of K_{yy} ($l=1$ month) (in 10^6 m²/s)
2 averaged over 40–60N at 440 K and 560 K and over 15–25N at 440 K in DJF, and averaged
3 over 40–60S at 440 K and 560 K and over 15–25S at 440 K in JJA. The standard deviation
4 (SD) is shown in brackets. The linear trend slope (Trend, in %/decade) is also shown. The
5 linear trend with a confidence level greater than 95% is shown in bold.

		NCEP- NCAR	NCEP- CFSR	ERA-40	ERA- Interim	JRA-25	JRA-55
40N–60N	Mean (SD)	1.02 (15.0)	1.30 (11.6)	1.68 (11.6)	1.72 (13.1)	1.57 (10.8)	1.27 (12.3)
440K	Trend	-1.1	-2.9	+5.7	+7.0	-2.4	+3.3
40N–60N	Mean (SD)	1.02 (18.7)	1.48 (11.6)	2.25 (14.7)	2.27 (11.1)	1.60 (16.4)	1.55 (12.2)
560K	Trend	+5.2	+2.2	+0.2	+0.5	+4.9	+0.5
15N–25N	Mean (SD)	0.56 (6.0)	0.70 (7.3)	0.80 (8.9)	0.76 (9.0)	0.76 (8.8)	0.81 (9.6)
440K	Trend	+0.5	+4.1	+9.0	-0.8	-0.6	-1.7
40S–60S	Mean (SD)	0.45 (12.9)	0.57 (16.6)	0.79 (19.9)	0.66 (16.4)	0.73 (14.1)	0.48 (±16.4)
440K	Trend	-1.3	+4.2	-3.7	+5.5	+18.0	+16.1
40S–60S	Mean (SD)	0.38 (19.2)	0.60 (13.2)	0.55 (37.2)	0.87 (12.1)	0.53 (21.9)	0.60 (14.9)
560K	Trend	-3.6	+3.9	-30.8	+1.6	+13.9	+3.0
15S–25S	Mean (SD)	0.43 (7.9)	0.52 (10.1)	0.47 (12.8)	0.53 (10.9)	0.61 (9.1)	0.54 (11.4)
440K	Trend	+8.0	-10.0	-11.2	+1.2	-2.9	-1.8

6

1 Table 2. Linear trend slope of K_{yy} ($l=1$ month) (in %/decade) for 40N–60N in DJF and for
 2 40S–60S in JJA at 560 K during the first 22 years (1979–2000) and during the last 12 years
 3 (2001–2012). None of the linear trend has a confidence level of 95%.

	NCEP- NCAR	NCEP- CFSR	ERA-40	ERA- Interim	JRA-25	JRA-55
40N–60N 560K (1979– 2000)	-10.3	+4.0	-5.3	+0.0	+8.2	-1.8
40N–60N 560K (2001– 2012)	-31.5	-14.3		-16.8	-23.5	-5.7
40S–60S 560K (1979–2000)	+7.2	+8.1	-34.2	+7.6	+14.0	+4.4
40S–60S 560K (2001–2012)	-39.4	-24.4		-25.2	-40.4	-26.2

4

1 Table 3. Same as in Table 1, but for the mass stream function (in 10^{10} kg/s) averaged over 40–
 2 60N in DJF, and over 40–60S in JJA at 440 K and 560 K. The linear trend with a confidence
 3 level greater than 95% is shown in bold.

		NCEP- NCAR	NCEP- CFSR	ERA-40	ERA- Interim	JRA-25	JRA-55
40N–60N	Mean (SD)	0.71 (4.3)	0.45 (5.6)	0.56 (6.5)	0.44 (5.4)	0.61 (3.9)	0.48 (4.5)
440K	Trend	+2.1	+1.9	+3.4	+1.8	+0.9	+2.5
40N–60N	Mean (SD)	0.40 (5.1)	0.28 (5.4)	0.31 (6.6)	0.27 (4.9)	0.36 (4.7)	0.28 (5.0)
560K	Trend	+0.9	+2.2	-2.0	-3.3	+1.3	+4.0
40S–60S	Mean (SD)	-0.28 (9.4)	-0.22 (8.3)	-0.20 (9.1)	-0.18 (8.2)	-0.23 (7.2)	-0.25 (4.9)
440K	Trend	-5.4	-0.7	-22.3	+0.1	-8.4	-6.7
40S–60S	Mean (SD)	-0.16 (13.0)	-0.13 (9.5)	-0.10 (9.0)	-0.09 (3.5)	-0.14 (9.7)	0.14 (7.4)
560K	Trend	-6.4	+2.0	-18.2	+4.0	+1.1	-4.1

4

1 Table 4. Same as in Table 1, but for the annual mean total upward flux (in 10^{10} kg/s) and its
 2 linear trend slope (in %/decade) at 440 K and 560 K. The linear trend with a confidence level
 3 greater than 95% is shown in bold.

		NCEP- NCAR	NCEP- CFSR	ERA-40	ERA- Interim	JRA-25	JRA-55
440K	Mean (SD)	0.98 (2.7)	0.67 (3.3)	0.89 (4.7)	0.73 (2.2)	0.86 (1.8)	0.80 (1.3)
	Trend	-1.8	+6.0	+8.5	-3.9	+1.4	+1.7
560K	Mean (SD)	0.48 (4.3)	0.34 (4.8)	0.47 (3.4)	0.36 (3.2)	0.44 (3.4)	0.40 (1.9)
	Trend	+2.1	+5.3	+2.3	-6.7	+0.6	+2.1

4

1 Table 5. Linear trend of the annual mean tropical total mean upward flux (in %/decade)
 2 during the first 22 years (1979–2000), and during the last 12 years (2001–2012) at 440 K and
 3 560 K. The linear trend with a confidence level greater than 95% is shown in bold.

	NCEP- NCAR	NCEP- CFSR	ERA-40	ERA- Interim	JRA-25	JRA-55
1979–2000 440K	+1.3	+9.6	+9.3	-3.0	+3.8	2.7
2001–2012 440K	-6.3	-2.0		-6.0	+0.9	-1.6
1979–2000 560K	+1.2	+9.9	+3.5	-4.1	+2.1	+2.8
2001–2012 560K	+8.1	+5.9		-10.1	+20.0	-3.2

4

1 Table 6. Relative contribution of different wave components to the zonal eddy PV component
 2 (in %) and the linear trend slope (in %/decade in brackets) averaged over 40–60N at 440 K
 3 and 560 K in DJF in 1979–2012. None of the linear trend has a confidence level of 95%.

		NCEP- NCAR	NCEP- CFSR	ERA-40	ERA- Interim	JRA-25	JRA-55
40N–60N 440K	1-3	52.9 (+4.2)	56.9 (+3.5)	55.0 (+4.8)	56.1 (+0.6)	55.2 (+2.4)	56.1 (+3.2)
	4-7	36.7 (+1.6)	35.1 (+2.9)	36.4 (+0.5)	35.5 (-0.9)	36.5 (+0.2)	35.7 (+1.2)
	8-	10.4 (+1.2)	8.0 (+7.4)	8.7 (+8.9)	8.4 (+1.9)	8.4 (+3.3)	8.2 (+2.7)
40N–60N 560K	1-3	70.3 (+11.5)	72.3 (+4.6)	72.5 (+11.1)	71.1 (+0.6)	73.5 (+5.8)	72.3 (+3.6)
	4-7	23.4 (+1.9)	23.7 (+4.3)	23.3 (+11.3)	24.5 (+4.1)	23.2 (+7.8)	23.9 (+3.2)
	8-	6.2 (+17.3)	4.0 (+10.8)	4.2 (+23.8)	4.4 (+6.8)	3.4 (+11.3)	3.8 (+10.9)

4

1 Table 7. Same as in Table 1, but for the relative importance of the eddy transport to the total
2 meridional transport averaged over 40–60N at 440 and 560 K and 15–25N at 440 K in DJF,
3 and over 40–60S at 440 K and 560 K and 15S–25S at 440K in JJA, and its linear trend slope
4 (in %/decade). The linear trend with a confidence level greater than 95% is shown in bold.

		NCEP- NCAR	NCEP- CFSR	ERA-40	ERA- Interim	JRA-25	JRA-55
40N–60N	Mean (SD)	0.38 (5.8)	0.54 (4.0)	0.45 (7.5)	0.52 (4.1)	0.46 (6.8)	0.51 (4.4)
440K	Trend	-0.9	+0.7	-6.7	-0.4	+1.0	+3.0
40N–60N	Mean (SD)	0.29 (8.4)	0.56 (4.9)	0.47 (4.7)	0.58 (4.5)	0.36 (9.4)	0.55 (4.4)
560K	Trend	-0.3	-0.6	-7.6	-1.8	-0.5	-1.6
15N–25N	Mean (SD)	0.34 (3.3)	0.48 (3.0)	0.32 (4.8)	0.45 (2.4)	0.39 (3.6)	0.44 (2.0)
440K	Trend	-0.9	-1.4	+4.2	+2.4	+2.8	+2.6
40S–60S	Mean (SD)	0.51 (4.4)	0.56 (5.2)	0.66 (5.8)	0.58 (3.3)	0.55 (7.9)	0.49 (4.1)
440K	Trend	-5.8	+0.4	+4.8	-2.4	-10.8	-1.5
40S–60S	Mean (SD)	0.40 (6.7)	0.65 (4.5)	0.54 (9.6)	0.70 (2.8)	0.49 (6.7)	0.57 (3.5)
560K	Trend	-3.2	+2.6	-12.9	+0.8	+2.4	-1.8
15S–25S	Mean (SD)	0.40 (3.5)	0.45 (5.0)	0.32 (12.2)	0.40 (3.7)	0.58 (5.9)	0.36 (3.5)
440K	Trend	-1.4	-6.1	-24.5	+1.1	-17.2	-4.4

5

1 Table 8. Linear trend slope of the relative importance of the eddy transport to the total
 2 meridional transport (in %/decade) for 15N–25N in DJF, and for 15S–25S in JJA at 440 K
 3 during the first 22 years (1979–2000), and during the last 12 years (2001–2012). The linear
 4 trend with a confidence level greater than 95% is shown in bold.


	NCEP- NCAR	NCEP- CFSR	ERA-40	ERA- Interim	JRA-25	JRA-55
15N-25N 440K (1979-2000)	-2.8	+2.8	+3.8	+1.8	+1.4	+2.8
15N-25N 440K (2001-2012)	+3.5	-6.9		+6.8	-2.9	-1.0
15S-25S 440K (1979-2000)	-3.4	-7.7	-27.3	+0.6	-23.9	-11.1
15S-25S 440K (2001-2012)	-4.6	-1.8		+8.3	+21.9	-4.7

5

1 Table 9. Correlation coefficient between the 6-hourly mass stream function (Mean) at 50N,
 2 440 K, and the 6-hourly E–P flux divergence at each grid point averaged between 500 and
 3 650 hPa and 50N–60N, and between K_{yy} ($l=1$ day) at 50N, 560 K, and the 6-hourly E–P flux
 4 divergence at each grid point averaged between 550 and 650 hPa and 45N–55N in DJF in
 5 2001–2012 (1990–2001 for ERA-40).

	NCEP- NCAR	NCEP- CFSR	ERA-40	ERA- Interim	JRA-25	JRA-55
Mean	-0.42	-0.61	-0.39	-0.58	-0.61	-0.59
K_{yy} ($l=1$ day)	-0.17	-0.21	-0.19	-0.26	-0.16	-0.22


6



1

2 Figure 1. Latitude and potential temperature cross section of the isentropic diffusion
 3 coefficient K_{yy} ($l=1$ month) (in 10^6 m²/s) averaged for December to February (DJF, left
 4 panel) and June to August (JJA, right panel) over the time period 1979–2012 (1979–2002 for
 5 ERA-40.). The results are shown for the NCEP-NCAR reanalysis (top left), NCEP-CFSR (top
 6 right), ERA-40 (middle left), ERA-Interim (middle right), JRA-25 (bottom left), and JRA-55
 7 (bottom right) in each panel.

8




1

2 Figure 2. Temporal variations of the isentropic diffusion coefficient K_{yy} ($l=1$ month) (in 10^6
 3 m^2/s) averaged over 40–60N in DJF (left) and 40–60S in JJA (right) at 440 K (top) and 560 K
 4 (bottom) for 1978–2012. The linear trend slope is also shown for each dataset. The results are
 5 shown for the NCEP-NCAR reanalysis (purple), NCEP-CFSR (green), ERA-40 (light blue),
 6 ERA-Interim (blue), JRA-25 (orange), and JRA-55 (red).

7


1



2

3 Figure 3. Same as Fig. 1, but for the mass stream function (in 10^{10} kg/s).


4




1

2 Figure 4. Seasonal variation of the mass stream function (colour contours, in 10^{10} kg/s) and
 3 zonal mean zonal wind (black contour lines, with intervals of 10 m/s) at 560 K.

4




1
 2 Figure 5. Same as in Fig. 3, but for the linear trend slope (10^{10} kg/s/decade). Statistically
 3 significant trends at the 95% level are indicated by hatching.
 4



1


2 Figure 6. Same as Fig. 2, but for the mass stream function (in 10^{10} kg/s).

3




1

2 Figure 7. Temporal variations of the annual mean total mean upward mass flux (in 10^{10} kg/s)
 3 at (top) 440 K and (bottom) 560 K.




1
 2 Figure 8. Same as Fig. 1, but for the relative importance of the eddy transport to the total
 3 meridional transport. The value larger (smaller) than 0.5 indicates that the eddy transport
 4 (mean transport) is dominant in the meridional transport.




1

2 Figure 9. Same as Fig. 2, but for the ratio of the relative importance of the eddy transport to
 3 the total meridional transport.

4




1
 2 Figure 10. Latitude and potential temperature cross section of the correlation (left) between
 3 the 6-hourly mass stream function at 50N, 440 K, and the 6-hourly E–P flux divergence at
 4 each grid point, and (right) between K_{yy} ($l=1$ day) at 50N, 560 K, and the 6-hourly E–P flux
 5 divergence at each grid point in DJF in 2000–2012. Correlation coefficients at the 95 %
 6 confidence level (following t-test) are indicated by hatching.
 7



1

2

3 Figure 11. Vertical profiles of the linear trend of the tropical upward mass flux (left, in 10^{10}
 4 kg/s/decade), relative contribution of the eddy transport to the total meridional transport
 5 averaged over $15\text{--}25^\circ$ (2nd left, in 1/decade), and the mean meridional velocity $\overline{v^*}$ averaged
 6 over $30\text{--}50^\circ$ (m/s/decade) during DJF in the NH (upper panels), and during JJA in the SH
 7 (lower panels) for the 34 years, 1979–2012.



2 Figure 12. Same as Fig. 11, but for the last 12 years, 2001–2012.