

Climate modulation of the Tibetan Plateau on haze in China

X. Xu et al.

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Climate modulation of the Tibetan Plateau on haze in China

X. Xu¹, T. Zhao^{2,3}, F. Liu^{4,a}, S. L. Gong⁵, D. Kristovich⁴, C. Lu⁶, Y. Guo^{1,7}, X. Cheng², Y. Wang¹, and G. Ding¹

¹State Key Laboratory of Severe Weather, Chinese Academy of Meteorological Sciences, Beijing, 100081, China

²Collaborative Innovation Center on Forecast and Evaluation of Meteorological Disasters, Nanjing University of Information Science and Technology, Nanjing 210044, China

³Key Laboratory for Aerosol-Cloud-Precipitation of China Meteorological Administration, Nanjing University of Information Science & Technology, Jiangsu, 210044, China

⁴Climate and Atmospheric Science Section, Division of Illinois State Water Survey, Prairie Research Institute, University of Illinois at Urban-Champaign, Champaign, IL 61820, USA

⁵Institute of Atmospheric Composition/Key Laboratory of Atmospheric Chemistry of China Meteorological Administration, Chinese Academy of Meteorological Sciences, Beijing, 100081, China

⁶National Science Foundation, VA 22230, USA

⁷Tiannan Observatory, Tianjin Meteorological Bureau, Tianjin 200350, China

^anow at: Environmental Program Division, Maricopa Association of Governments, Phoenix, AZ 85003, USA

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Received: 2 October 2015 – Accepted: 9 October 2015 – Published: 26 October 2015

Correspondence to: T. Zhao (tlzhao@nuist.edu.cn)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ACPD

15, 28915–28937, 2015

**Climate modulation
of the Tibetan Plateau
on haze in China**

X. Xu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Rapid increases in pollutant emissions in conjunction with stagnant meteorological conditions result in haze pollution in China. Recent frequent haze in China has attracted worldwide attention. Here we show a relationship between the haze events and Tibetan Plateau (TP)'s environment and climate changes. Based on observational data taken over recent decades, we identify central-eastern China (CEC) as a climatological large-scale "susceptible region" of frequent haze, which is harbored by the TP with its impact on mid-latitude westerly winds. The observational and modeling studies demonstrate that the interannual variations in the thermal forcing of TP are positively correlated with the incidences of wintertime haze over CEC. Further analysis indicates that the TP-climate warming induced changes in atmospheric circulation driving frequent haze events in CEC. The frequent haze occurrences in CEC are consistent with decreasing winter monsoon winds, intensifying downward air flows and increasing atmospheric stability in the lower troposphere over the CEC in association with upstream plateau's thermal anomalies. Therefore, variations of haze in China are related to mechanical and thermal forcing by the TP. Our results also suggest that implications of the large TP-topography for environment and climate changes should be taken into account for air pollution mitigation policies in China.

1 Introduction

Haze in poor visibility with high particulate matter (PM) or aerosol levels is a pervasive air quality problem facing China, posing a major challenge for public health (Huang et al., 2014). The frequent haze pollution has been notable for hitting record high levels of PM pollution over central-eastern China (CEC) in history since 1961 (Ding and Liu, 2014). In January 2013, extremely severe and persistent haze events swept over much of CEC-region. A large area of CEC from the North China Plain, including Beijing, across the Lower Yangtze Valley Plain to the Sichuan Basin, was blanketed in thick

Climate modulation of the Tibetan Plateau on haze in China

X. Xu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Climate modulation of the Tibetan Plateau on haze in China

X. Xu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



haze and smog for almost one month. It is estimated by the Chinese government that this wintertime haze covered a quarter of the total land area in China with 600 million people, half of the Chinese population, exposed to the haze air pollution (Gu, 2013). China's National Meteorological Center released its first ever "haze" orange alert (CMA, 2010) in response to the air quality index frequently reaching hazardous levels for this regional haze event. The $PM_{2.5}$ (PM with an aerodynamic diameter less than $2.5\ \mu\text{m}$) concentrations at 33 cities in the CEC region were more than $300\ \mu\text{g m}^{-3}$ for longer than half a month, and some monitors reported hourly peak $PM_{2.5}$ levels of $900\ \mu\text{g m}^{-3}$, which is classified as "Beyond Index" (Gu, 2013). The suffering of those in China from haze and poor air quality has attracted worldwide attention (Wang et al., 2014; Pope III and Dockery, 2013; Chen et al., 2013; Kan et al., 2011; Park et al., 2013; Zhao et al., 2013).

China has been experiencing increased air pollution, commonly attributed to the large increases in pollutant emissions associated with the rapid economic development. The huge increases in CO_2 emissions from fossil fuel combustion with the certain attendant aerosol loading have a direct physical link to more frequent haze occurrences in situ in China (upper panel of Fig. 1). The regional emissions of air pollutants contribute largely to the haze pollution in CEC with the coefficient of determination, $R^2 = 0.9025$ between interannual variations of haze frequency and CO_2 emission in China (upper panel of Fig. 1). However, air quality is modulated by changes in meteorology and climate (Tagaris et al., 2009). In accompany with an unceasing increase in the Chinese pollutant emissions in recent decades, the significant interannual variations of haze occurrences in CEC have evolved with the trends of slow ascending from the 1960s to 70s, less changing during the 1980s–1990s and sharply rising with a trend reaching 13.0d/10a going into the 21st century (upper panel of Fig. 1), implying that climate change could also play an important role in the variations of haze events in CEC apart from the anthropogenic dimension of pollutant emission sources related to the rapid industrialization of China. A steady decline of East Asian monsoon winds is negatively correlated to haze occurrences in the CEC with the coefficient of deter-

2 Data and method

In this study, we used the observational records from 1961 to 2012 of haze and surface wind archived at the China Meteorological Administration (CMA) and the meteorological variables of air temperature, winds and relative humidity from the reanalysis data generated by the US National Center for Environmental Prediction–National Center for Atmospheric Research (NCEP/NCAR).

Following the studies of Yanai (1961), Yanai and Johnson (1993), Yanai and Tomita (1998), the apparent heat source (Q_1) and apparent moisture sink (Q_2) are calculated. Atmospheric heat sources and moisture sinks are respectively gauged with the Q_1 and Q_2 . As Q_1 includes Q_2 and radiative heating, here we concentrate only on the collective effect of apparent heating (Q_1) over the TP. The heat source column (in units of W m^{-2}) over the TP is obtained with both horizontal and vertical integration of Q_1 over the TP-area of $78\text{--}103^\circ\text{E}$ and $28\text{--}38^\circ\text{N}$ covering the most region with the altitude of higher than 3000 m (left panel of Fig. 2) to form a one-dimensional variable representing the TP-thermal forcing. The correlation coefficients between the TP-heat source column and the meteorological variables (U , V and W components of wind and air temperature) are calculated to build their horizontal and vertical distributions of correlations. Zonal, meridional and vertical components of the correlation vector are respectively derived through the correlation coefficients of the TP-heat source column to U , V and W components of vector of wind and air temperature, indicating the variations in wind and air temperature induced by the TP-thermal forcing.

In the modeling study, we used the global air quality model system GEM-AQ/EC, which is an integration of gas phase chemistry and aerosol modules in the meteorological model GEM (Global Environmental Multiscale weather prediction model of Environment Canada). Full details of the development and evaluation of GEM-AQ/EC are given by Gong et al. (2012). The validations of 10 year (1995–2004) GEM-AQ/EC modeling prove that GEM-AQ provided satisfactory simulations of the distribution and variation of global and regional aerosols (Gong et al., 2012; Zhao et al., 2012). Re-

Climate modulation of the Tibetan Plateau on haze in China

X. Xu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



gional variations of aerosols in East Asia are reasonably captured by the GEM-AQ/EC modeling compared to the observed aerosol concentrations and aerosol optical depth.

3 A climatological “susceptible region” for haze formation in China

Examination of ground-based observations of the frequency of haze events from 1961 to 2012 (CMA, 2010) reveals that the haze air pollution in China typically has the highest levels in the CEC region covering a vast area from the eastern edges of the TP and the Loess Plateau to China’s Pacific coast, and haze occurrences in CEC oscillate seasonally between the peak in winter and the low in summer (Fig. 2). Based upon these climate data, we could climatologically regard the CEC, with the lowlands harboured by the upstream plateaus of western China, as a large-scale “susceptible region” of frequent haze events in China (left panel of Fig. 2). Upper panel of Fig. 3 shows that low average wind speeds tend to be coincident with the centers of pollutant haze events over the CEC (left panel of Fig. 2), reflecting the climatological “susceptible region” of haze occurrences in connection with a stagnation area in the lower troposphere in China.

Due to the influence of the TP terrain on the typical westerly winds in this region, the air flowing from the windward plateaus descends in a north–south oriented zone between about 110 and 125° E (middle panel of Fig. 3). Accompanying this strong downward current are weak winds in the near-surface layers that lie in the lee side of the plateaus. These air flow and wind condition lead to development of a “harbor” that accumulates air pollutants in the CEC region. The weak wind and downward current areas coincide well with the centers of frequent haze events in China (middle and lower panels of Fig. 3). The “susceptible region” of haze events over the CEC region from the eastern edge of the plateaus to the lower flatlands is associated with the “harbor” effect of the unique TP topography under specific meteorological conditions that trap air pollutants.

Climate modulation of the Tibetan Plateau on haze in China

X. Xu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Because haze is climatologically mostly a winter phenomenon in the CEC (right panel of Fig. 2), the following analysis on the TP's climate effect on haze pollution in CEC and the related mechanisms is focused on the winter season.

4 Relationships between TP's thermal forcing and haze over CEC

As a vast elevated landmass, the TP acts thermodynamically as a synoptic-scale wintertime cooling source protruding into the free atmosphere (Qiu, 2008; Liu and Chen, 2000; Ruddiman and Kutzbach, 1989; Yeh et al., 1957). Aerosol transport and deposition have been increasingly dirtying and even melting the snow- and ice-dominated wintertime TP (Ramanathan and Carmichael, 2008; Xu et al., 2009). This process leads to decreases in the snow and ice albedos, which could be largely responsible for climate change in the TP region (Hansen and Nazarenko, 2004). Consequently, the TP region, as a wintertime cooling source (negative values of apparent heat source Q_1), has been experiencing a warming trend over recent decades, especially since 2001 (upper left panel of Fig. 4).

Against the backdrop of global climate change, the question may be posed: does the warming of the TP region cause changes in the atmospheric environment in China resulting in more frequent haze events in the CEC? The historical data analysis indicates that a significant correlation exists between the wintertime cooling source represented by the apparent heat source column Q_1 integrated over the TP and the number of haze days averaged regionally in the CEC over recent decades (upper panel of Fig. 4). It is also found in upper panel of Fig. 4 that the changes of wintertime Q_1 over the TP were reversed from cooling to warming in the late 1990s, which could be connected with the trends in haze occurrences with less changing over the 1980s–1990s and sharp increasing during the 21st century in China under the increases in pollutant emission levels (upper panel of Fig. 1). Based on the composite analysis on the haze frequencies in winter with positive and negative anomalies in wintertime cooling source of the TP, the haze increasing and decreasing incidences over the CEC are found in good agree-

ment with the positively and negatively anomalous TP-cooling sources (lower panel of Fig. 4). The frequency of haze events over the CEC region is positively correlated with climatic warming over the TP.

5 The TP-warming inducing favorable meteorology for CEC's haze

Further analyses provide information on the mechanisms relating climatic warming of the TP and enhancement in haze occurrence in the CEC. The favorable meteorology for haze occurrences is well known to be lower wind speeds, weaker vertical mixing, stronger subsidence, higher air humidity and more stable low-level stratification. We are still pondering the question whether climatic warming of the TP could strengthen the aforementioned meteorological conditions downstream for frequent haze events in the CEC to reveal the mechanism how thermal anomalies of the TP in climate change influence the incidence of haze over the CEC.

The East Asian winter monsoon, which climatologically prevails over the CEC, typically maintains near-surface northeastern winds (Ding, 1994). In upper panel of Fig. 5, two horizontal components of the correlation vector are derived through two correlation coefficients of Q_1 to U and V surface wind components, respectively, where the arrow length denotes the combined correlation with a longer arrow implying a better correlation, and the arrow direction means the direction of anomalous wind induced by the TP-thermal effect. The correlation vector over the CEC in upper panel of Fig. 5 indicates that the variations of thermal forcing over the TP could give rise to the weakening winter monsoon winds (southwest wind anomalies) induced by changes of Q_1 over the TP. Furthermore, the anomalous south wind components resulting from climate change in the TP (positive correlations of Q_1 to V wind components in upper panel of Fig. 5) can enhance transport of water vapor from the oceans to the CEC (Niu et al., 2010). By increasing the moisture in the lower troposphere driven by the strong vapor transport (Hung and Kao, 2010) (lower panel of Fig. 5), in addition to decline in

Climate modulation of the Tibetan Plateau on haze in China

X. Xu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the East Asian winter monsoon with weak advection of cold air, haze formation can be enhanced (Zhao et al., 2013; Xiao et al., 2011).

Upper panels of Fig. 6 present the results of composite analysis on vertical variations in air temperature in five winters respectively with the most positive and the most negative anomalies in the TP cooling source. The air temperature changes with upper warmer and lower cooler are found in the middle and lower troposphere over the CEC region in winter with positive Q_1 anomalies on the TP, and an inverse structure of the CEC air temperature changes in winter with negative Q_1 anomalies on the TP (upper panel of Fig. 6). The TP's warming and cooling anomalies could lead to a "warm shield" and a "cool shield" in the atmosphere over the CEC. The correlation analyses of observation data over 1981–2012 confirm that the vertical structure of anomalous air temperature similar to that by induced the TP's positive thermal effect (upper-left panel of Fig. 6) with a "warm shield" intensifying the subsidence in the lower troposphere is responsible for the frequent haze occurrences over the CEC (lower panel of Fig. 6). Associated with the warming TP, the vertical variations of air temperature with upper warmer and lower cooler could easily build an inversion layer in the atmosphere over the polluted CEC, which results in more stably stratified atmosphere in this region (Fig. 6). Heavy haze pollution processes in winter are highly related with the existence of atmospheric inversion layer (Xu et al., 2003).

The cumulative consequences of weakening winter monsoon winds, intensifying downward air flows, a more humid and more stable atmosphere as the favorable meteorological conditions for haze formation would be expected to strengthen the air pollutant "harbor" effect of the TP and increase the number and severity of haze events in the CEC. Therefore, the haze formation over CEC is significantly modulated by the TP's climate change under the increase and even without changes in the current levels of anthropogenic pollutant emissions.

Climate modulation of the Tibetan Plateau on haze in China

X. Xu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

6 A sensitivity simulation experiment

In order to more convincingly demonstrate the connection of the TP-warming to the haze frequency over CEC, a sensitivity simulation by employing the global air quality model GEM-AQ/EC is designed to isolate the emission influence on interannual variations of aerosols, where the monthly data of anthropogenic emissions by fossil fuel and biomass burning as well as the sulfate emissions compiled using EDGAR2.0 (Gong et al., 2012) are introduced without any interannual changes from 1995 to 2004. The sensitivity simulation experiment of GEM-AQ/EC was configured with 28 hybrid vertical levels with the model top at 10 hPa. The horizontal model grid was configured with a global uniform resolution of $1^\circ \times 1^\circ$. GEM-AQ/EC is run with the fully nudged variables of wind, temperature, pressure and water vapor of NCEP-reanalysis meteorology every 24 h from 1995 to 2004.

Haze and aerosol changes are controlled by both pollutant emission and meteorology, and the effects of meteorology are difficult to separate from aerosol observations. The 10-year GEM-AQ/EC simulation without interannual changes in the anthropogenic emissions provides a possibility to identify the meteorological effect on the interannual variations of aerosols. To investigate the implications of TP's climate change for interannual aerosol variations in CEC's haze, a composite analysis of surface aerosol concentrations over CEC (Fig. 7) were performed for two winters with lower TP's Q_1 (1996, 2002) and two winters with higher TP's Q_1 (1998, 2003) during the simulation period of 1995–2004 according to the interannual Q_1 changes over the TP (Fig. 4). Because the effect of emissions was singled out in the interannual aerosol variations modeled in the sensitivity simulation experiment, the simulated variations in aerosol concentrations over CEC could be purely attributed to the changes of meteorological drivers in the context of changing climate. The analysis results show that the TP heating anomalies could lead to enhances of 30–45% in wintertime surface aerosol concentrations over the CEC-region compared to the winters with the TP cooling anomalies (Fig. 7). Because changes of aerosol levels in the surface atmosphere determine haze

formation, this sensitivity simulation confirmed that the frequent haze in China with the significantly interannual variations is closely related to thermal forcing by the TP, and climate change of the TP could intensify pollutant haze in China even without increases in the current anthropogenic pollutant emissions.

7 Conclusions

Based on observational data over the recent decades, we identify the CEC region, the lower flatlands along the eastern plateau edges in China as a climatological large-scale “susceptible region” of pollutant haze in connected with downward currents and weak near-surface winds in consequence of the “harbor” impact of large TP topography on mid-latitude westerlies. The climate analysis reveals that the increasingly frequent haze in the CEC region is related with decreasing winter monsoon winds, intensifying descending air and increasing atmospheric stability in the lower-troposphere over the CEC in association with plateau’s thermal anomalies. Climate impact of the TP’s mechanical and thermal forcing driving changes in atmospheric circulation and meteorological conditions downstream is potentially contributing to the increasing trend in haze events in China. A sensitivity simulation also confirms that the frequent haze in CEC with the significantly interannual variations is closely connected with thermal forcing by the TP.

The Chinese government has been making great strides in reducing emissions and mitigating air pollution. However, the interplay of China’s unique landform distribution with climate change and its associated more extreme weather events could impair the effectiveness of air pollution control measures in China. With the influence of the TP climate change, the CEC region is facing a bigger challenge to realize air quality maintenance plan. The TP “harbor effect” and climate change should be considered in making decisions on the locations of new industrial facilities for development planning in China in order to preferentially reduce anthropogenic emissions in the “suscepti-

Climate modulation of the Tibetan Plateau on haze in China

X. Xu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ble region” of haze and in turn reduce the number and severity of haze events in the central-eastern region of China.

In this preliminary study based on long-term observational data and a sensitivity simulation experiment, we investigate a relationship between the haze pollution in China and TP’s environment and climate changes. The impacts of TP’s climate change on air quality in China could be further studied on the shifts in weather patterns, pollutant emissions and chemical reaction to more comprehensively understand the meteorological drivers of air quality in a changing climate and also to consider the ocean-related impacts of climate change.

Acknowledgements. This research is jointly supported by the Projects of Nature Science Fund of China (No. 41130960), Chinese National Science and Technology Project (2014BAC22B04), the projects of China Special Fund for Meteorological Research (GYHY201406001) and Environmental Protection (HY14093355; 201509001) in the Public Interest, Chinese Key Technologies R&D Program (2014BAC16B00), Science and Technology Support Program of Jiangsu Province (BE2012771) and the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD). We acknowledge the review of Beth Hall (ISWS).

References

- Chen, Y. Y., Ebenstein, A., Greenstone, M., and Li, H. B.: Evidence on the impact of sustained exposure to air pollution on life expectancy from China’s Huai River policy, *P. Natl. Acad. Sci. USA*, 110, 12936–12941, doi:10.1073/pnas.1300018110, 2013.
- CMA: China Meteorological Administration, National meteorological standard of the People’s Republic China: observation and forecasting levels of haze, QX-T 113–2010, 2010 (in Chinese).
- Ding, Y. H.: Monsoons over China, Kluwer Academic Publishers, Dordrecht, Boston, London, 1994.
- Ding, Y. H. and Liu, Y. J.: Analysis of long-term variations of fog and haze in China in recent 50 years and their relations with atmospheric humidity, *Science China: Earth Sciences*, 57, 36–46, doi:10.1007/s11430-013-4792-1, 2014.

Climate modulation of the Tibetan Plateau on haze in China

X. Xu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Climate modulation
of the Tibetan Plateau
on haze in China**

X. Xu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Gong, S. L., Lavoué, D., Zhao, T. L., Huang, P., and Kaminski, J. W.: GEM-AQ/EC, an on-line global multi-scale chemical weather modelling system: model development and evaluation of global aerosol climatology, *Atmos. Chem. Phys.*, 12, 8237–8256, doi:10.5194/acp-12-8237-2012, 2012.

5 Gu, L.: Half of Chinese live in haze: report, China News, available at: <http://www.ecns.cn/cns-wire/2013/07-12/72889.shtml> (last access: 12 July 2013) 2013.

Hansen, J. and Nazarenko, L.: Soot climate forcing via snow and ice albedos, *P. Natl. Acad. Sci. USA*, 101, 423–428, 2004.

10 Huang, R.-J., Zhang, Y., Bozzetti, C., Ho, K.-F., Cao, J.-J., Han, Y., Daellenbach, K. R., Slowik, J. G., Platt, S. M., Canonaco, F., Zotter, P., Wolf, R., Pieber, S. M., Bruns, E. A., Crippa, M., Ciarelli, G., Piazzalunga, A., Schwikowski, M., Abbaszade, G., Schnelle-Kreis, J., Zimmermann, R., An, Z., Szidat, S., Baltensperger, U., Haddad, I. E., and Prevot, A. S. H.: High secondary aerosol contribution to particulate pollution during haze events in China, *Nature*, 514, 218–222, doi:10.1038/nature13774, 2014.

15 Hung, C.-W. and Kao, P.-K.: Weakening of the winter monsoon and abrupt increase of winter rainfalls over northern taiwan and southern china in the early 1980s, *J. Climate*, 23, 2357–2367, doi:10.1175/2009JCLI3182.1, 2010.

Kan, H., Chen, R., and Tong, S.: Ambient air pollution, climate change, and population health in China, *Environ. Int.*, 42, 10–19, doi:10.1016/j.envint.2011.03.003, 2011.

20 Liu, X. and Chen, B.: Climatic warming in the Tibetan Plateau during recent decades, *Int. J. Climatol.*, 20, 1729–1742, 2000.

Niu, F., Li, Z., Li, C., Lee, K.-H., and Wang, M.: Increase of wintertime fog in China: potential impacts of weakening of the Eastern Asian monsoon circulation and increasing aerosol loading, *J. Geophys. Res.-Atmos.*, 115, D00K20, doi:10.1029/2009jd013484, 2010.

25 Oey, L.-Y., Chang, M.-C., Chang, Y.-L., Lin, Y.-C., and Xu, F.-H.: Decadal warming of coastal China Seas and coupling with winter monsoon and currents, *Geophys. Res. Lett.*, 40, 6288–6292, doi:10.1002/2013GL058202, 2013.

Park, S.-U., Cho, J.-H., and Park, M.-S.: Analyses of high aerosol concentration events (dense haze/mist) occurred in East Asia during 10–16 January 2013 using the data simulated by the Aerosol Modeling System, *International Journal of Chemistry*, 03, 10–26, 2013.

30 Pope III, C. A. and Dockery, D. W.: Air pollution and life expectancy in China and beyond, *P. Natl. Acad. Sci. USA*, 110, 12861–12862, doi:10.1073/pnas.1310925110, 2013.

Climate modulation of the Tibetan Plateau on haze in China

X. Xu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Qian, Y., Flanner, M. G., Leung, L. R., and Wang, W.: Sensitivity studies on the impacts of Tibetan Plateau snowpack pollution on the Asian hydrological cycle and monsoon climate, *Atmos. Chem. Phys.*, 11, 1929–1948, doi:10.5194/acp-11-1929-2011, 2011.
- Qiu, J.: China: the third pole, *Nature*, 454, 393–396, 2008.
- 5 Ramanathan, V. and Carmichael, G.: Global and regional climate changes due to black carbon, *Nature*, 1, 221–227, 2008.
- Ruddiman, W. F. and Kutzbach, J. E.: Forcing of late Cenozoic Northern Hemisphere climate by plateau uplift in southern Asia and the American west, *J. Geophys. Res.*, 94, 18409–18427, doi:10.1029/JD094iD15p18409, 1989.
- 10 Tagaris, E., Liao, K. J., Delucia, A. J., Deck, L., Amar, P., and Russell, A. G.: Potential impact of climate change on air pollution-related human health effects, *Environ. Sci. Technol.*, 43, 4979–4988, 2009.
- Wang, Y., Zhang, R. Y., and Saravanan, R.: Asian pollution climatically modulates midlatitude cyclones following hierarchical modeling and observational analysis, *Nature Commun.*, 5, 3098, doi:10.1038/ncomms4098, 2014.
- 15 Wu, G. X., Liu, Y., He, B., Q. Bao, Duan, A., and Jin, F.-F.: Thermal controls on the Asian summer monsoon, *Scientific Reports*, 2, 404, doi:10.1038/srep00404, 2012.
- Xiao, Z.-M., Zhang, Y.-F., Hong, S.-M., Bi, X.-H., Jiao, L., Feng, Y.-C., and Wang, Y.-Q.: Estimation of the main factors influencing haze, based on a long-term monitoring campaign in Hangzhou, China, *Aerosol Air Qual. Res.*, 11, 873–882, doi:10.4209/aaqr.2011.04.0052, 2011.
- 20 Xu, B., Cao, J. J., Hansen, J., Yao, T., Joswida, D. R., Wang, N., Wu, G., Wang, M., Zhao, H., Yang, W., Liu, X., and He, J.: Black soot and the survival of Tibetan glaciers, *P. Natl. Acad. Sci. USA*, 106, 22114–22118, 2009.
- 25 Xu, M., Chang, C.-P., Fu, C., Qi, Y., Robock, A., Robinson, D., and Zhang, H.: Steady decline of east Asian monsoon winds, 1969–2000: Evidence from direct ground measurements of wind speed, *J. Geophys. Res.*, 111, D24111, doi:10.1029/2006JD007337, 2006.
- Xu, X., Ding, G., Zhou, L., Zheng, X., Bian, L., Qiu, J., Yang, L., and Mao, J.: Localized 3D-structural features of dynamic-chemical processes of urban air pollution in Beijing winter, *Chinese Sci. Bull.*, 48, 819–825, 2003.
- 30 Xu, X., Lu, C., Shi, X., and Ding, Y.: Large-scale topography of China: A factor for the seasonal progression of the Meiyu rainband?, *J. Geophys. Res.*, 115, D02110, doi:10.1029/2009JD012444, 2010.

Climate modulation of the Tibetan Plateau on haze in China

X. Xu et al.




Figure 1. Interannual variations in the total CO₂ emissions in mainland China and the haze event frequency averaged in the CEC region over 1961–2012 with their relationship in the inserted figure (upper panel). The Chinese CO₂ emission data source: http://cdiac.ornl.gov/CO2_Emission/timeseries/national; two blue dotted lines separate the time series into three phases of the 1960s–1970s, the 1980s–1990s and the 21st century. Interannual variations in wind at 10 m (blue line) and the number of days with haze (red line) in the CEC over 1961–2012 (lower-left panel) and their scatter plot (lower-right panel).

Climate modulation
of the Tibetan Plateau
on haze in China

X. Xu et al.




Figure 2. Distribution of annual haze event frequency (days per year) averaged over 1961–2012 in China, and Chinese topography of the TP and the Loess Plateau with altitudes is shown with yellow shades (left panel). Monthly variation of haze frequency averaged from 1961–2012 over the CEC region (right panel).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Climate modulation of the Tibetan Plateau on haze in China

X. Xu et al.




Figure 3. Surface wind speed distribution (ms^{-1}) averaged over 1961–2012 in China with the red rectangle marking the region for cross sections in the middle and lower panels (upper panel). Cross sections of horizontal wind speed (ms^{-1} ; filled color contours) and vertical circulations illustrated by stream lines (middle panel) and zonal variations of annual haze event frequency (lower panel) at $27\text{--}41^\circ\text{N}$ averaged over 1961–2012. Note that near-surface vertical and horizontal winds are not illustrated well here due to north–south variations in the terrain and approximation of the location of the TP in this figure. All fields are for the annual-averages.

Climate modulation
of the Tibetan Plateau
on haze in China

X. Xu et al.




Figure 4. Interannual variability in the apparent heat source Q_1 (the negative values denote cooling) integrated vertically over the TP and haze event frequency averaged in the CEC in winter (December, January and February) over 1980–2012 and their correlation (upper panel). The differences of haze frequencies (days) averaged in five winters with most positive (lower left panel) and most negative Q_1 anomalies (lower right panel) on the TP relative to the mean haze frequency from 1980 to 2012.

Climate modulation of the Tibetan Plateau on haze in China

X. Xu et al.




Figure 5. The distributions of the interannual correlations of the apparent heat source Q_1 over the TP to the local V component of surface wind in winter over 1961–2012 (color shading). Arrows denote correlation vectors (showing both correlation coefficients of Q_1 to U and V surface wind components) in China. The correlation coefficients of 0.12 (–0.12), 0.14 (–0.14) and 0.19 (–0.19) respectively passing the significance levels of 90, 95 and 99%. A vertical section of the trends in vapour content (g kg⁻¹ per 10 years) in winter over 1961–2012 averaged along 27–41° N (lower panel).

Climate modulation
of the Tibetan Plateau
on haze in China

X. Xu et al.




Figure 6. Vertical sections of the anomalous air temperature ($^{\circ}\text{C}$) averaged along $27\text{--}41^{\circ}\text{N}$ in five winters with most positive (upper left panel) and most negative Q_1 anomalies (upper right panel) on the TP from 1980 to 2012, and vertical sections of the correlations of the number of haze days with air temperature (lower left panel) and vertical circulations (lower right panel) in winter from 1980 to 2012.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Climate modulation
of the Tibetan Plateau
on haze in China

X. Xu et al.




Figure 7. The percentages (%; contour lines) of differences of surface aerosol concentrations between winters of 1998 and 2003 with positive and winters of 1996 and 2002 with negative Q_1 anomalies on the TP relative to the surface aerosol levels averaged over winters of 1996, 1998, 2002 and 2003 ($\mu\text{g m}^{-3}$; color contours) modeled by the sensitivity simulation experiment with GEM-AQ/EC.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)