

Characterization of
polar organosulfates
in secondary organic
aerosol

M. S. Shalamzari et al.

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Characterization of polar organosulfates in secondary organic aerosol from the unsaturated aldehydes 2-*E*-pentenal, 2-*E*-hexenal, and 3-*Z*-hexenal

M. S. Shalamzari^{1,2}, R. Vermeylen¹, F. Blockhuys³, T. E. Kleindienst⁴,
M. Lewandowski⁴, R. Szmigielski⁵, K. J. Rudzinski⁵, G. Spólnik⁶,
W. Danikiewicz⁶, W. Maenhaut^{1,2}, and M. Claeys¹

¹Department of Pharmaceutical Sciences, University of Antwerp, 2610 Antwerp, Belgium

²Department of Analytical Chemistry, Ghent University, 9000 Gent, Belgium

³Department of Chemistry, University of Antwerp, 2020 Antwerp, Belgium

⁴National Exposure Research Laboratory, Office of Research and Development, United States Environmental Protection Agency, Research Triangle Park, North Carolina 27711, USA

⁵Institute of Physical Chemistry, Polish Academy of Sciences, 01-224 Warsaw, Poland

⁶Institute of Organic Chemistry, Polish Academy of Sciences, 01-224 Warsaw, Poland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Received: 4 August 2015 – Accepted: 30 September 2015 – Published: 29 October 2015

Correspondence to: M. Claeys (magda.claeys@uantwerpen.be)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ACPD

15, 29555–29590, 2015

Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

We show in the present study that the unsaturated aldehydes, 2-*E*-pental, 2-*E*-hexenal and 3-*Z*-hexenal, are biogenic volatile organic compound (BVOC) precursors for polar organosulfates with molecular weights (MWs) 230 and 214, which are also present in ambient fine aerosol from a forested site, i.e., K-pusztá, Hungary. These results complement those obtained in a previous study showing that the green leaf aldehyde 3-*Z*-hexenal serves as a precursor for MW 226 organosulfates. Thus, in addition to isoprene, the green leaf volatiles 2-*E*-hexenal and 3-*Z*-hexenal, emitted due to plant stress (mechanical wounding or insect attack), and 2-*E*-pental, a photolysis product of 3-*Z*-hexenal, should be taken into account for secondary organic aerosol and organosulfate formation. Polar organosulfates are of climatic relevance because of their hydrophilic properties and cloud effects. Extensive use was made of organic mass spectrometry (MS) and detailed interpretation of MS data (i.e., ion trap MS and accurate mass measurements) to elucidate the chemical structures of the MW 230, 214 and 170 organosulfates formed from 2-*E*-pental and indirectly from 2-*E*-hexenal and 3-*Z*-hexenal. In addition, quantum chemical calculations were performed to explain the different mass spectral behavior of 2,3-dihydroxypentanoic acid sulfate derivatives, where only the isomer with the sulfate group at C-3 results in the loss of SO₃. The MW 214 organosulfates formed from 2-*E*-pental are explained by epoxidation of the double bond in the gas phase and sulfation of the epoxy group with sulfuric acid in the particle phase through the same pathway as that proposed for 3-sulfoxy-2-hydroxy-2-methylpropanoic acid from the isoprene-related α,β -unsaturated aldehyde methacrolein in previous work (Lin et al., 2013). The MW 230 organosulfates formed from 2-*E*-pental are tentatively explained by a novel pathway, which bears features of the latter pathway but introduces an additional hydroxyl group at the C-4 position. Evidence is also presented that the MW 214 positional isomer, 2-sulfoxy-3-hydroxypentanoic acid is unstable and decarboxylates, giving rise to 1-sulfoxy-2-

Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



hydroxybutane, a MW 170 organosulfate. Furthermore, evidence is obtained that lactic acid sulfate is generated from 2-*E*-pentenal.

1 Introduction

Volatile organic compounds (VOCs) are ubiquitous in the troposphere, playing a key role as precursors for ozone and secondary organic aerosol (SOA). The formation and aging of SOA has received considerable attention during the last two decades because of its potential impact on climate and human health. Major classes of SOA precursors studied include alkanes, alkenes, aromatic hydrocarbons and oxygenated compounds, both from anthropogenic and biogenic origin. Among the biogenic VOCs that are precursors for SOA, isoprene and monoterpenes have received ample attention (for a review, see Hallquist et al., 2009), while other reactive VOCs such as green leaf volatiles (GLVs) have been far less examined. However, recent studies demonstrate that GLVs are also potential precursors for biogenic SOA (Hamilton et al., 2009; Harvey et al., 2014; Jain et al., 2014; Richards-Henderson et al., 2014; Shalamzari et al., 2014).

GLVs comprise unsaturated C₆ compounds that are released by plants when they are wounded (e.g., grass cutting, animal grazing, storms) or attacked by insects. They are named after their specific “green” odor, the fresh scent emitted by green plants, and play a crucial role in plant-plant and plant-insect communication (Hatanaka, 1993; Holopainen, 2004; Scala et al., 2013). GLVs are formed in plant leaves from the unsaturated fatty acids linoleic and α -linolenic acid, which are essential constituents of cell membrane lipids, by biochemical conversion with the enzymes lipoxygenase and hydroperoxide lyase (Hatanaka, 1993). One of the most abundant GLVs, 3-*Z*-hexenal, is formed by the cleavage of α -linolenic acid, and it partly isomerizes to 2-*E*-hexenal. Both alkenals can be further metabolized by an alcohol dehydrogenase and alcohol acyltransferase (D’Auria et al., 2007) to the corresponding alcohols and their esters (Matsui, 2006).

Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

GLVs also have a potential role in tropospheric chemistry as they may serve as precursors for ozone and SOA (Pinto et al., 2007; Holopainen, 2011; Hartikainen et al., 2012). Recently, the GLVs 3-Z-hexen-1-ol, 3-Z-hexenyl acetate, and methyl salicylate, methyl jasmonate and 2-methyl-3-butene-2-ol, have also been shown to undergo aqueous-phase oxidation with the hydroxyl radical and to result in considerable SOA yields ranging from 10 to 88 % (Richards-Henderson et al., 2014). SOA formation through photooxidation and ozonolysis was first studied at the molecular level for the GLVs 3-Z-hexen-1-ol and 3-Z-hexenyl acetate by Hamilton et al. (2009) with liquid chromatography combined with positive ion electrospray ionization mass spectrometry using lithium as a cationization agent. This study demonstrated that 3-hydroxypropanal, the primary oxidation product of 3-Z-hexen-1-ol, can hydrate and undergo further reactions with other aldehydes resulting in SOA dominated by higher-molecular weight (MW) oligomers, while, in contrast, considerably fewer higher MW species are detected in SOA produced from 3-Z-hexenylacetate. SOA formation was recently examined from turf grass volatiles (Jain et al., 2014), which comprise 3-Z-hexen-1-ol and 3-Z-hexenyl acetate, with near-infrared laser desorption/ionization aerosol mass spectrometry, and the ozonolysis SOA profile was found to closely resemble that of 3-Z-hexen-1-ol and to be dominated by low-volatility, higher-MW compounds such as oligoesters. In addition, there is recent evidence that 3-Z-hexenal results in polar m/z 225 (MW 226) organosulfates (OSs) through photooxidation and/or ozonolysis (Shalamzari et al., 2014), which occur at substantial concentrations in fine ambient aerosol ($PM_{2.5}$) from a forested site in Belgium (Gómez-González et al., 2012).

OSs are potential marker compounds for SOA formation occurring under acidic conditions by particle-phase reactions with sulfuric acid (Surratt et al., 2007b), formed by oxidation of sulfur dioxide, which is mainly from anthropogenic origin in continental regions of the globe (Smith et al., 2011). In addition to formation through reaction with sulfuric acid of epoxy-containing SOA compounds from the oxidation of β -pinene (Iinuma et al., 2007), isoprene (Surratt et al., 2010; Lin et al., 2013) and 2-methyl-3-buten-2-ol (Zhang et al., 2012), OSs may also result from the reactive uptake of

Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



unsaturated compounds in the particle phase and reaction with the sulfate anion radical (Rudzinski et al., 2009; Nozière et al., 2010; Schindelka et al., 2013; Szmigielski, 2013; Schöne et al., 2014). In addition, OSs may also be formed by uptake of tertiary organonitrates in the particle phase and nucleophilic substitution of the nitrate by a sulfate group, as shown in the case of 2-methyltetrol sulfates (Darer et al., 2011; Hu et al., 2011). Organosulfates are ubiquitous compounds in our environment, not only formed from BVOCs but also from anthropogenic VOCs such as aromatics (Kundu et al., 2013; Riva et al., 2015). It has been suggested that they are a significant component of fine ambient aerosol (e.g., Surratt et al., 2008; Lukács et al., 2009; Hawkins et al., 2010; Tolocka and Turpin, 2012). Using Fourier transform infrared measurements Hawkins et al. (2010) determined that 4 ± 8 % of the organic mass of continental outflow aerosols over the southeast Pacific Ocean during the VOCALS–Rex 2008 campaign was due to organosulfates during periods of high organic and sulfate concentrations. The annual average contributions of organosulfates to organic mass for twelve sites in the United States was found to be 5–10 % using a S-estimation method, and was higher during warm months when photochemical oxidation chemistry is most active (Tolocka and Turpin, 2012). As to K-puszta, Hungary, it was shown that organosulfates correspond to a substantial fraction of fine ambient summer aerosol, which was determined with S-estimation methods to be as high as 30 % of the total organic mass (Surratt et al., 2008) and to correspond to 6–12 % of the bulk sulfur mass (Lukács et al., 2009). The present study focuses on the chemical characterization of m/z 169, 213 and 229 OSs formed from the C_5 -unsaturated aldehyde, 2-*E*-pental, which is a known photolysis product of 3-*Z*-hexenal (O'Connor et al., 2006), as well as from the C_6 -unsaturated aldehydes 2-*E*-hexenal and 3-*Z*-hexenal. In a previous study (Gómez-González et al., 2008), it was shown that fine ambient aerosol ($PM_{2.5}$) contains polar m/z 213 OSs (i.e., sulfate esters of 2,3-dihydroxypentanoic acid), which could be related to 2-*E*-pental, have the same magnitude as the m/z 215 OSs that originate from isoprene (Surratt et al., 2007a, 2008) and are formed through sulfation of intermediary C_5 -epoxydiols (Surratt et al., 2010). Formation pathways for the m/z 213 and 229 OSs that originate from 2-*E*-

pentenal are proposed, thereby considering a known route leading to OSs, namely, the route established for 3-sulfooxy-2-hydroxy-2-methylpropanoic acid from methacrolein in the presence of NO_x ($\text{NO} + \text{NO}_2$), which, as 2-*E*-pentenal, is also a α, β -unsaturated aldehyde (Lin et al., 2013).

5 A first step in understanding the impact of an unknown organosulfate (or any organic compound) in the environment is its characterization at the molecular level, as this knowledge allows one to obtain information on its VOC precursor and its formation process (for a review, see Nozière et al., 2015). Only after its molecular structure has been established, ambient monitoring studies can be considered involving quantitative
10 measurements, which allow one to obtain further insights into its formation. In order to chemically characterize the unknown OSs from 2-*E*-pentenal in detail laboratory smog chamber experiments were conducted in the present study with 2-*E*-pentenal, 3-*Z*-hexenal and 2-*E*-hexenal, and the organosulfate profiles were compared with those of ambient fine ($\text{PM}_{2.5}$) aerosol collected from K-puszta, Hungary, a rural site with
15 mixed deciduous-coniferous vegetation. For the chemical analysis, liquid chromatography combined with negative ion electrospray ionization mass spectrometry (LC/(-)ESI-MS) was performed by resorting to reversed-phase LC using a trifunctionally-bonded C_{18} stationary phase (Shalamzari et al., 2013). As to mass spectrometric techniques, use was made of high-resolution MS to infer the elemental composition, while ion trap
20 MS was employed to obtain information about functional groups and their position in the molecules, a state-of-the-art analytical methodology which has proven its usefulness in previous studies on the molecular characterization of organosulfates (Gómez-González et al., 2008; Surratt et al., 2008; Shalamzari et al., 2013, 2014) and SOA (for a review, see Nozière et al., 2015). In addition, quantum chemical calculations
25 were performed to gain insight into the distinctive mass spectral behavior of positional isomers containing neighboring sulfate, hydroxyl, and carboxyl groups.

Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of the solution was adjusted to 3.1 with 0.1 M H₂SO₄, and the autoxidation of sulfite by the dissolved atmospheric oxygen was catalyzed by injecting a small aliquot of Fe₂(SO₄)₃ catalyst solution (2.5 × 10⁻⁵ M). The experiment was carried out at 25 ± 0.1 °C and the reaction time at which the 2-*E*-pentenoic acid reaction products were sampled was 5 min.

2.2.3 Ambient fine aerosol

Archived PM_{2.5} (particulate matter with an aerodynamic diameter ≤ 2.5 μm) aerosol samples collected from K-pusztá, Hungary, during the BIOSOL (Formation mechanisms, marker compounds, and source apportionment for biogenic atmospheric aerosols) campaign between 22 May and 29 June 2006, were used. Information about the site and the sampling campaign can be found in a previous article (Maenhaut et al., 2008). Information about the preparation of a pooled day- and night-time sample and the work-up of the sample, involving extraction with methanol, can be found in Shalamzari et al. (2013). Samples containing 570 μg of organic carbon were analyzed with LC/(-)ESI-MS using an aliquot of 5 μL.

2.3 Liquid chromatography/mass spectrometry

The LC/ESI-ITMS analysis was carried out using a Surveyor Plus system (pump and autosampler) (Thermo Scientific, San Jose, CA, USA) connected to an Atlantis T3 column (2.1 mm × 150 mm, 3 μm particle size, Waters, Milford, MA, USA), which provides retention for the organosulfate compounds under investigation. The mobile phases for reversed-phase (RP)-LC consisted of 50 mM ammonium formate buffer pH 3 and methanol. Information about the applied LC/MS gradient program and the operation conditions of the linear ion-trap mass spectrometer (LXQ, Thermo Scientific) can be found in Shalamzari et al. (2013). Accurate mass measurements were carried out using a Synapt G2-S HDMS instrument (Waters) equipped with an electrospray ion source and a quadrupole – time-of-flight type mass analyzer. The instrument was con-

diagnostic product ions formed by further fragmentation of m/z 185 (Fig. S2a) at m/z 167, 153, 141, and 123 (Scheme 1). Ions formed by further fragmentation of m/z 211 (Fig. S2b) to m/z 193, 167 and 97 are also explained in Scheme 1.

Furthermore, it is noted that the m/z 229 profiles obtained for K-pusztza fine aerosol and 2-*E*-pentenal (Fig. 2) are quite comparable, revealing a major compound eluting at 2.5 min, while additional later-eluting isomers could be observed for 3-*Z*-hexenal and 2-*E*-hexenal SOA (Fig. S1). Since only the m/z 229 peak eluting at 2.5 min is found in ambient fine aerosol, no attention is given to the latter later-eluting 3-*Z*-hexenal and 2-*E*-hexenal SOA isomers. As 2,3,4-trihydroxypentanoic acid has three chiral carbon atoms, several stereoisomeric forms (theoretically eight) are indeed possible. An early-eluting m/z 229 isomer (RT 2.2 min) is detected in fine ambient aerosol that is not formed from 2-*E*-pentenal, 3-*Z*-hexenal, and 2-*E*-hexenal. Selected MS data for the latter isomer are provided in Fig. S3 in the Supplement. It can be seen that the MS data only differ from those of the major peak at RT 2.5 min by the relative abundances of the product ions, consistent with stereoisomers.

3.3 Proposed formation pathway for m/z 229 organosulfates formed from 2-*E*-pentenal

As discussed above, m/z 229 compounds detected in 2-*E*-pentenal SOA could be assigned to stereoisomeric forms of 3-sulfooxy-2,4-dihydroxypentanoic acid. The detection of these compounds in 3-*Z*-hexenal SOA is in line with the fact that 2-*E*-pentenal is a photolysis product of 3-*Z*-hexenal (O'Connor et al., 2006). The formation of 2-*E*-pentenal through photolysis of 2-*E*-hexenal has not been reported so far although it is very likely that it may also be formed in this way.

A possible formation pathway leading to the structurally characterized isomeric m/z 229 organosulfates is proposed in Scheme 2, which bears features of the pathway proposed for the formation of 3-sulfooxy-2-hydroxy-2-methylpropanoic acid from methacrolein, which, as 2-*E*-pentenal, is also a α,β -unsaturated aldehyde (Lin et al., 2013). Reactions (1) are inspired by the latter pathway, involving abstraction of the

29567

Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Figure 5 shows selected LC/MS chromatographic data (m/z 169 EICs) obtained for ambient fine aerosol and 2-*E*-pentenal SOA, as well as selected MS data (MS² and MS³ product ion spectra). Three peaks can be distinguished in the m/z 169 EICs, of which the first peak co-elutes with the first-eluting m/z 213 OS (2.4 min), attributed to 2-sulfooxy-3-hydroxypentanoic acid. This co-elution behavior was also noted under improved chromatographic separation conditions where two Atlantis T3 columns were used in series (results not shown). It thus appears that 2-sulfooxy-3-hydroxypentanoic acid is unstable upon electrospray ionization. A possible fragmentation mechanism involving decarboxylation and resulting in m/z 169 is provided in Scheme 3a. The second peak (RT 2.7 min, measured mass, 168.9807; error: -2.9 mDa; elemental formula, C₃H₅O₆S) is attributed to lactic acid sulfate, a known organosulfate, which has been reported in ambient fine aerosol (Olson et al., 2011; Shalamzari et al., 2013; Hettiyadura et al., 2015). However, since no authentic standard was available, this assignment should be regarded as tentative. The third peak (RT 3.8 min, measured mass, 169.0177; error: $+0.6$ mDa; elemental formula, C₄H₉O₅S) reveals a fragmentation pattern that perfectly matches that of the first m/z 169 peak, which is due to decarboxylation. It is thus very likely that this compound results from decarboxylation of the corresponding m/z 213 OS; however, on the basis of this study it could not be determined where the degradation takes place (i.e., on the filter during sampling, during sample workup or during analysis). Furthermore, it was confirmed that the m/z 169 compounds that are present in 2-*E*-pentenal SOA are also formed from 3-*Z*-hexenal and 2-*E*-hexenal (Fig. S5). Based on MS data the product with RT 3.8 min in ambient fine aerosol (4.1 min in 2-*E*-pentenal SOA) is assigned to 1-sulfooxy-2-hydroxybutane (Scheme 3).

The instability of 2-sulfooxycarboxylic acids due to decarboxylation also provides an explanation why in the case of methacrolein only the formation of 3-sulfooxy-2-hydroxy-2-methylpropanoic acid has been reported (Gómez-González et al., 2008; Surratt et al., 2008; Shalamzari et al., 2013), and not that of its positional isomer 2-sulfooxy-3-hydroxy-2-methylpropanoic acid.

3.5 Proposed formation pathway for m/z 213 organosulfates formed from 2-*E*-pentenal

As discussed above, m/z 213 compounds detected in 2-*E*-pentenal SOA could be assigned to positional isomeric OSs of 2,3-dihydroxypentanoic acid, i.e., 2-sulfooxy-3-hydroxypentanoic acid and 3-sulfooxy-2-hydroxypentanoic acid. Their detection in 2-*E*-hexenal and 3-*Z*-hexenal SOA is logical as other organosulfates, i.e., the m/z 229 compounds (Sect. 3.2 and 3.3), could also be found in the latter SOA and the fact that 2-*E*-pentenal is a photolysis product of 3-*Z*-hexenal (O'Connor et al., 2006). A possible formation pathway leading to the structurally characterized isomeric m/z 213 organosulfates is proposed in Scheme 4. The suggested pathway is based on the mechanism reported for the formation of 3-sulfooxy-2-hydroxy-2-methylpropanoic acid from methacrolein, which, as 2-*E*-pentenal, is also a α,β -unsaturated aldehyde (Lin et al., 2013). It involves a sequence of reactions, comprising epoxidation of the double bond of 2-*E*-pentenal with the OH radical and reaction of the epoxy group with sulfuric acid in the particle phase. The results obtained in this study further suggest that the formation of the m/z 213 sulfate derivatives of 2,3-dihydroxypentanoic acid likely does not result from reactive uptake of 2-*E*-pentenal as only one positional isomer (3-sulfooxy-2-hydroxypentanoic acid) could be generated in the aqueous-phase sulfation of 2-*E*-pentenoic acid, whereas two positional isomers (the latter and 2-sulfooxy-3-hydroxypentanoic acid) are detected in both 2-*E*-pentenal SOA and fine ambient aerosol. The formation of 2-hydroxy-3-sulfooxypentanoic acid in the aqueous-phase reaction of 2-*E*-pentenoic acid with the sulfate radical anion (Scheme S1) is caused by stabilization of the resulting intermediate C-2 radical upon attack of the double bond with the sulfate radical anion.

Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

4 Conclusions and perspectives

We show in the present study that the unsaturated aldehydes, 2-*E*-pental, 2-*E*-hexenal and 3-*Z*-hexenal, can be converted into polar organosulfates with *m/z* 229 and 213. These results are in line with a previous study that the green leaf aldehyde 3-*Z*-hexenal serves as a precursor for *m/z* 225 organosulfates (Shalamzari et al., 2014). Thus, in addition to isoprene, the green leaf volatiles, 2-*E*-hexenal and 3-*Z*-hexenal, and 2-*E*-pental, a photolysis product of 3-*Z*-hexenal, should be taken into account for organosulfate formation. These results imply that biogenic volatile organic compounds such as the green leaf volatiles, which are emitted due to plant stress (mechanical wounding or insect attack), are a potential source of polar organosulfates and SOA.

Extensive use was made of organic mass spectrometry and detailed interpretation of mass spectral data to elucidate the chemical structures of the *m/z* 229, 213 and 169 organosulfates formed from 2-*E*-pental. In addition, quantum chemical calculations were performed to explain the different mass spectral behavior of 3-sulfoxy-2-hydroxypentanoic acid and 2-sulfoxy-3-hydroxypentanoic acid, where the isomer with the sulfate group at C-3 results in the loss of SO₃. The *m/z* 213 organosulfates formed from 2-*E*-pental are explained by the same route as that reported for 3-sulfoxy-2-hydroxy-2-methylpropanoic acid from methacrolein, which, as 2-*E*-pental, is also a α,β -unsaturated aldehyde. The pathway involves formation of an epoxide in the gas phase and sulfation of the epoxy group with sulfuric acid in the particle phase. The *m/z* 229 organosulfates formed from 2-*E*-pental are tentatively explained by a novel pathway that bears features of the latter pathway but introduces an additional hydroxyl group. Evidence is also presented that the *m/z* 213 OS, 2-sulfoxy-3-hydroxypentanoic acid, is unstable and decarboxylates, giving rise 1-sulfoxy-2-hydroxybutane, a *m/z* 169 organosulfate. The instability of 2-sulfoxy carboxylic acids due to decarboxylation also provides an explanation why in the case of the isoprene-related methacrolein only 3-sulfoxy-2-hydroxy-2-methylpropanoic acid has been detected in previous studies

Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and not its isomer 2-sulfooxy-3-hydroxy-2-methylpropanoic acid. Furthermore, it could be shown that lactic acid sulfate is generated from 2-*E*-pentenal. With regard to the proposed pathways for organosulfate formation from unsaturated aldehydes, further research is warranted to gain additional mechanistic insights.

5 **The Supplement related to this article is available online at
doi:10.5194/acpd-15-29555-2015-supplement.**

10 *Acknowledgements.* Research at the University of Antwerp was supported by the Belgian Federal Science Policy Office through the network project “Biogenic Influences on Oxidants and Secondary Organic Aerosol: theoretical, laboratory and modelling investigations (BIOSOA)” and by the Research Foundation – Flanders (FWO) through the project “Secondary Organic Aerosol formation from monoterpenes: GAPS in our current understanding (SOAGAPS)”. The Polish Academy of Sciences (PAS) collaborated in the study reported here through an academic exchange project supported by the PAS and FWO – Flanders. The US Environmental Protection Agency through its Office of Research and Development partially funded and collaborated in the research described here under contract EP-D-10-070 to Alion Science and Technology. The manuscript has been subjected to Agency review and has not been cleared for publication. Mention of trade names or commercial products does not constitute an endorsement or recommendation for use.

References

- 20 Boschan, R., Merrow, R. T., and Van Dolah, R. W.: The chemistry of nitrate esters, *Chem. Rev.*, 55, 485–510, 1955.
- Chen, X. H., Hulbert, D., and Shepson, P. B.: Measurement of the organic nitrate yield from OH reaction with isoprene, *J. Geophys. Res.-Atmos.*, 103, 25563–25568, 1998.
- Darer, A. I., Cole-Filipiak, N. C., O’Connor, A. E., and Elrod, M. J.: Formation and stability of atmospherically relevant isoprene-derived organosulfates and organonitrates, *Environ. Sci. Technol.*, 45, 1895–1902, 2011.

Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



D'Auria, J. C., Pichersky, E., Schaub, A., Hansel, A., and Gershenzon, J.: Characterization of a BAHD acyltransferase responsible for producing the green leaf volatile (Z)-3-hexen-1-yl acetate in *Arabidopsis thaliana*, *Plant J.*, 49, 194–207, 2007.

5 Frisch, M. J., Trucks, G. W., Schlegel, H. B., Scuseria, G. E., Robb, M. A., Cheeseman, J. R., Scalmani, G., Barone, V., Mennucci, B., Petersson, G. A., Nakatsuji, H., Caricato, M., Li, X., Hratchian, H. P., Izmaylov, A. F., Bloino, J., Zheng, G., Sonnenberg, J. L., Hada, M., Ehara, M., Toyota, K., Fukuda, R., Hasegawa, J., Ishida, M., Nakajima, T., Honda, Y., Kitao, O., Nakai, H., Vreven, T., Montgomery Jr., J. A., Peralta, J. E., Ogliaro, F., Bearpark, M., Heyd, J. J., Brothers, E., Kudin, K. N., Staroverov, V. N., Kobayashi, R., Normand, J., Raghavachari, K., Rendell, A., Burant, J. C., Iyengar, S. S., Tomasi, J., Cossi, M., Rega, N., Millam, J. M., Klene, M., Knox, J. E., Cross, J. B., Bakken, V., Adamo, C., Jaramillo, J., Gomperts, R., Stratmann, R. E., Yazyev, O., Austin, A. J., Cammi, R., Pomelli, C., Ochterski, J. W., Martin, R. L., Morokuma, K., Zakrzewski, V. G., Voth, G. A., Salvador, P., Dannenberg, J. J., Dapprich, S., Daniels, A. D., Farkas, O., Foresman, J. B., Ortiz, J. V., Cioslowski, J., and Fox, D. J.: Gaussian 09, Revision A.02, Gaussian Inc., Wallingford, CT, USA, 2009.

10 Gómez-González, Y., Surratt, J. D., Cuyckens, F., Szmigielski, R., Vermeylen, R., Jaoui, M., Lewandowski, M., Offenberg, J. H., Kleindienst, T. E., Edney, E. O., Blockhuys, F., Van Alsenoy, C., Maenhaut, W., and Claeys, M.: Characterization of organosulfates from the photooxidation of isoprene and unsaturated fatty acids in ambient aerosol using liquid chromatography/(–)electrospray ionization mass spectrometry, *J. Mass Spectrom.*, 43, 371–382, 2008.

15 Gómez-González, Y., Wang, W., Vermeylen, R., Chi, X., Neiryneck, J., Janssens, I. A., Maenhaut, W., and Claeys, M.: Chemical characterisation of atmospheric aerosols during a 2007 summer field campaign at Brasschaat, Belgium: sources and source processes of biogenic secondary organic aerosol, *Atmos. Chem. Phys.*, 12, 125–138, doi:10.5194/acp-12-125-2012, 2012.

20 Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, Th. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9, 5155–5236, doi:10.5194/acp-9-5155-2009, 2009.

Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Hamilton, J. F., Lewis, A. C., Carey, T. J., Wenger, J. C., Borrás i Garcia, E., and Muñoz, A.: Reactive oxidation products promote secondary organic aerosol formation from green leaf volatiles, *Atmos. Chem. Phys.*, 9, 3815–3823, doi:10.5194/acp-9-3815-2009, 2009.
- Hartikainen, K., Riikonen, J., Nerg, A. M., Kivimaenpää, M., Ahonen, V., Tervahauta, A., Karenlampi, S., Maenpää, M., Rousi, M., Kontunen-Soppela, S., Oksanen, E., and Holopainen, T.: Impact of elevated temperature and ozone on the emission of volatile organic compounds and gas exchange of silver birch (*Betula pendula Roth*), *Environ. Exp. Bot.*, 84, 33–43, 2012.
- Harvey, R. M., Zahardis, J., and Petrucci, G. A.: Establishing the contribution of lawn mowing to atmospheric aerosol levels in American suburbs, *Atmos. Chem. Phys.*, 14, 797–812, doi:10.5194/acp-14-797-2014, 2014.
- Hatanaka, A.: The biogenesis of green odour by green leaves, *Phytochemistry*, 34, 1201–1218, 1993.
- Hawkins, L. N., Russell, L. M., Covert, D. S., Quinn, P. K., and Bates, T. S.: Carboxylic acids, sulfates, and organosulfates in processed continental organic aerosol over the south-east Pacific Ocean during VOCALS-REx 2008, *J. Geophys. Res.-Atmos.*, 115, D13201, doi:10.1029/2009jd013276, 2010.
- Hettiyadura, A. P. S., Stone, E. A., Kundu, S., Baker, Z., Geddes, E., Richards, K., and Humphry, T.: Determination of atmospheric organosulfates using HILIC chromatography with MS detection, *Atmos. Meas. Tech.*, 8, 2347–2358, doi:10.5194/amt-8-2347-2015, 2015.
- Holopainen, J. K.: Can forest trees compensate for stress-generated growth losses by induced production of volatile compounds?, *Tree Physiol.*, 31, 1356–1377, 2011.
- Hu, K. S., Darer, A. I., and Elrod, M. J.: Thermodynamics and kinetics of the hydrolysis of atmospherically relevant organonitrates and organosulfates, *Atmos. Chem. Phys.*, 11, 8307–8320, doi:10.5194/acp-11-8307-2011, 2011.
- Iinuma, Y., Müller, C., Berndt, T., Böge, O., Claeys, M., and Herrmann, H.: Evidence for the existence of organosulfates from beta-pinene ozonolysis in ambient secondary organic aerosol, *Environ. Sci. Technol.*, 41, 6678–6683, 2007.
- Jain, S., Zahardis, J., and Petrucci, G. A.: Soft ionization chemical analysis of secondary organic aerosol from green leaf volatiles emitted by turf grass, *Environ. Sci. Technol.*, 48, 4835–4843, 2014.
- Kleindienst, T. E., Jaoui, M., Lewandowski, M., Offenberg, J. H., Lewis, C. W., Bhawe, V., and Edney, E. O.: Estimates of the contributions of biogenic and anthropogenic hydrocarbons to



**Characterization of
polar organosulfates
in secondary organic
aerosol**

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

secondary organic aerosol at a southeastern US location, Atmos. Environ., 41, 8288–8300, 2007.

Kundu, S., Quraishi, T. A., Yu, G., Suarez, C., Keutsch, F. N., and Stone, E. A.: Evidence and quantitation of aromatic organosulfates in ambient aerosols in Lahore, Pakistan, Atmos. Chem. Phys., 13, 4865–4875, doi:10.5194/acp-13-4865-2013, 2013.

Lin, Y. H., Zhang, H., Pye, H. O. T., Zhang, Z., Marth, W. J., Park, S., Arashiro, M., Cui, T., Budisulistiorini, S. H., Sexton, K. G., Vizuete, W., Xie, Y., Luecken, D. J., Piletic, I. R., Edney, E. O., Bartolotti, L. J., Gold, A., and Surratt, J. D.: Epoxide as a precursor to secondary organic aerosol formation from isoprene photooxidation in the presence of nitrogen oxides, P. Natl. Acad. Sci. USA, 110, 6718–6723, 2013.

Lukács, H., Gelencsér, A., Hoffer, A., Kiss, G., Horváth, K., and Hartyáni, Z.: Quantitative assessment of organosulfates in size-segregated rural fine aerosol, Atmos. Chem. Phys., 9, 231–238, doi:10.5194/acp-9-231-2009, 2009.

Maenhaut, W., Raes, N., Chi, X., Cafmeyer, J., and Wang, W.: Chemical composition and mass closure for $PM_{2.5}$ and PM_{10} aerosols at K-pusztá, Hungary, in summer 2006, X-Ray Spectrom., 37, 193–197, 2008.

Matsui, K.: Green leaf volatiles: hydroperoxide lyase pathway of oxylipin metabolism, Curr. Opin. Plant Biol., 9, 274–280, 2006.

Nozière, B., Ekström, S., Alsberg, T., and Holmström, S.: Radical-initiated formation of organosulfates and surfactants in atmospheric aerosols, Geophys. Res. Lett., 37, L05806, doi:10.1029/2009GL041683, 2010.

Nozière, B., Kalberer, M., Claeys, M., Allan, J., D'Anna, B., Decesari, S., Finessi, E., Glasius, M., Grgić, I., Hamilton, J. F., Hoffmann, T., Iinuma, Y., Jaoui, M., Kahnt, A., Kampf, C. J., Kourtchev, I., Maenhaut, W., Marsden, N., Saarikoski, S., Schnelle-Kreis, J., Surratt, J. D., Szidat, S., Szmigielski, R., and Wisthaler, A.: The Molecular identification of organic compounds in the atmosphere: state of the art and challenges, Chem. Rev., 115, 3919–3983, 2015.

O'Connor, M. P., Wenger, J. C., Mellouki, A., Wirtz, K., and Muñoz, A.: The atmospheric photolysis of E-2-hexenal, Z-3-hexenal and E,E-2,4-hexadienal, Phys. Chem. Chem. Phys., 8, 5236–5246, 2006.

Olson, C. N., Galloway, M. M., Yu, G., Hedman, C. J., Lockett, M. R., Yoon, T., Stone, E. A., Smith, L. M., and Keutsch, F. N.: Hydroxycarboxylic acid-derived organosulfates: synthesis, stability, and quantification in ambient aerosol, Environ. Sci. Technol., 45, 6468–6474, 2011.

Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Pinto, D. M., Nerg, A. M., and Holopainen, J. K.: The role of ozone-reactive compounds, terpenes, and green leaf volatiles (GLVs), in the orientation of *Cotesia plutellae*, J. Chem. Ecol., 33, 2218–2228, 2007.
- Richards-Henderson, N. K., Hansel, A. K., Valsaraj, K. T., and Anastasio, C.: Aqueous oxidation of green leaf volatiles by hydroxyl radical as a source of SOA: Kinetics and SOA yields, Atmos. Environ., 95, 105–112, 2014.
- Riva, M., Tomaz, S., Cui, T., Lin, Y.-H., Perraudin, E., Gold, A., Stone, E. A., Villenave, E., and Surratt, J. D.: Evidence for an unrecognized secondary anthropogenic source of organosulfates and sulfonates: Gas-phase oxidation of polycyclic aromatic hydrocarbons in the presence of sulfate aerosol, Environ. Sci. Technol., 49, 6654–6664, 2015.
- Romero, F. and Oehme, M.: Organosulfates – a new component of humic-like substances in atmospheric aerosols?, J. Atmos. Chem., 52, 283–294, 2005.
- Rudziński, K. J., Gmachowski, L., and Kuznietsova, I.: Reactions of isoprene and sulphoxy radical-anions – a possible source of atmospheric organosulphites and organosulphates, Atmos. Chem. Phys., 9, 2129–2140, doi:10.5194/acp-9-2129-2009, 2009.
- Sato, K.: Detection of nitrooxypolyols in secondary organic aerosol formed from the photooxidation of conjugated dienes under high-NO_x conditions, Atmos. Environ., 42, 6851–6861, 2008.
- Scala, A., Allmann, S., Mirabella, R., Haring, M. A., and Schuurink, R. C.: Green leaf volatiles: a plant’s multifunctional weapon against herbivores and pathogens, Int. J. Mol. Sci., 14, 17781–17811, 2013.
- Schindelka, J., Iinuma, Y., Hoffmann, D., and Herrmann, H.: Sulfate radical-initiated formation of isoprene-derived organosulfates in atmospheric aerosols, Faraday Discuss., 165, 237–259, 2013.
- Schöne, L., Schindelka, J., Szeremeta, E., Schaefer, T., Hoffmann, D., Rudzinski, K. J., Szmigielski, R., and Herrmann, H.: Atmospheric aqueous phase radical chemistry of the isoprene oxidation products methacrolein, methyl vinyl ketone, methacrylic acid and acrylic acid – kinetics and product studies, Phys. Chem. Chem. Phys., 16, 6257–6272, 2014.
- Shalamzari, M. S., Ryabtsova, O., Kahnt, A., Vermeylen, R., Herent, M. F., Quetin-Leclercq, J., Van der Veken, P., Maenhaut, W., and Claeys, M.: Mass spectrometric characterization of organosulfates related to secondary organic aerosol from isoprene, Rapid Commun. Mass Sp., 27, 784–794, 2013.

**Characterization of
polar organosulfates
in secondary organic
aerosol**

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Shalamzari, M. S., Kahnt, A., Vermeylen, R., Kleindienst, T. E., Lewandowski, M., Cuyckens, F., Maenhaut, W., and Claeys, M.: Characterization of polar organosulfates in secondary organic aerosol from the green leaf volatile 3-Z-hexenal, *Environ. Sci. Technol.*, 48, 12671–12678, 2014.

5 Smith, S. J., van Aardenne, J., Klimont, Z., Andres, R. J., Volke, A., and Delgado Arias, S.: Anthropogenic sulfur dioxide emissions: 1850–2005, *Atmos. Chem. Phys.*, 11, 1101–1116, doi:10.5194/acp-11-1101-2011, 2011.

Stephens, P. J., Devlin, F. J., Chabalowski, C. F., and Frisch, M. J.: Ab-initio calculation of vibrational absorption and circular-dichroism spectra using density-functional force-fields, *J. Phys. Chem.-US*, 98, 11623–11627, 1994.

10 Surratt, J. D., Kroll, J. H., Kleindienst, T. E., Edney, E. O., Claeys, M., Sorooshian, A., Ng, N. L., Offenberg, J. H., Lewandowski, M., Jaoui, M., Flagan, R. C., and Seinfeld, J. H.: Evidence for organosulfates in secondary organic aerosol, *Environ. Sci. Technol.*, 41, 517–527, 2007a.

15 Surratt, J. D., Lewandowski, M., Offenberg, J. H., Jaoui, M., Kleindienst, T. E., Edney, E. O., and Seinfeld, J. H.: Effect of acidity on secondary organic aerosol formation from isoprene, *Environ. Sci. Technol.*, 41, 5363–5369, 2007b.

Surratt, J. D., Gómez-González, Y., Chan, A. W. H., Vermeylen, R., Shahgholi, M., Kleindienst, T. E., Edney, E. O., Offenberg, J. H., Lewandowski, M., Jaoui, M., Maenhaut, W., Claeys, M., Flagan, R. C., and Seinfeld, J. H.: Organosulfate formation in biogenic secondary organic aerosol, *J. Phys. Chem. A*, 112, 8345–8378, 2008.

20 Surratt, J. D., Chan, A. W. H., Eddingsaas, N. C., Chan, M. N., Loza, C. L., Kwan, A. J., Hersey, S. P., Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Reactive intermediates revealed in secondary organic aerosol formation from isoprene, *P. Natl. Acad. Sci. USA*, 107, 6640–6645, 2010.

25 Szmigielski, R.: Chemistry of organic sulfates and nitrates in the urban atmosphere, in: Disposal of Dangerous Chemicals in Urban Areas and Mega Cities, Role of Oxides and Acids of Nitrogen in Atmospheric Chemistry, edited by: Barnes, I. and Rudzinski, K. J., Nato Science for Peace and Security Series – C: Environmental Chemistry, Springer, Dordrecht, the Netherlands, 211–226, 2013.

30 Tolocka, M. P. and Turpin, B.: Contribution of organosulfur compounds to organic aerosol mass, *Environ. Sci. Technol.*, 46, 7978–7983, 2012.

Yi, L., Dratter, J., Wang, C., Tunge, J. A., and Desire, H.: Identification of sulfation sites of metabolites and prediction of the compounds' biological effects, *Anal. Bioanal. Chem.*, 386, 666–674, 2006.

Zhang, H. F., Worton, D. R., Lewandowski, M., Ortega, J., Rubitschun, C. L., Park, J. H., Kristensen, K., Campuzano-Jost, P., Day, D. A., Jimenez, J. L., Jaoui, M., Offenberg, J. H., Kleindienst, T. E., Gilman, J., Kuster, W. C., de Gouw, J., Park, C., Schade, G. W., Frossard, A. A., Russell, L., Kaser, L., Jud, W., Hansel, A., Cappellin, L., Karl, T., Glasius, M., Guenther, A., Goldstein, A. H., Seinfeld, J. H., Gold, A., Kamens, R. M., and Surratt, J. D.: Organosulfates as tracers for secondary organic aerosol (SOA) formation from 2-methyl-3-buten-2-ol (MBO) in the atmosphere, *Environ. Sci. Technol.*, 46, 9437–9446, 2012.

Ziajka, J. and Rudzinski, K. J.: Autoxidation of S-IV inhibited by chlorophenols reacting with sulfate radicals, *Environ. Chem.*, 4, 355–363, 2007.

ACPD

15, 29555–29590, 2015

Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Chamber conditions for experiments used for the structural characterization of unknown organosulfates related to 2-*E*-pentenal, 2-*E*-hexenal, and 3-*E*-hexenal SOA. Abbreviations: SA, sulfuric acid; AS, ammonium sulfate; RH, relative humidity.

Sample code	Hydrocarbon		Hydrocarbon Conc.	NO _x Conc.	O ₃ Conc.	OC Conc.	Seed type Conc.	Other Conditions
ER625–GF5	2- <i>E</i> -pentenal	Initial:	4.8 ppm C	280 ppb	–	–	1/2 AS + 1/2 SA 28 μg m ⁻³	4 light banks 516 nmol m ⁻³ (H ⁺)air RH: 27 %
		Steady State:	1.1 ppm C	178 ppb	207 ppb	4.9 μg m ⁻³		
ER627–GF10	3- <i>Z</i> -hexenal	Initial:	2.5 ppm C	290 ppb	–	–	1/2 AS + 1/2 SA 30 μg m ⁻³	4 light banks 439 nmol m ⁻³ (H ⁺)air RH: 29 %
		Steady state:	0.3 ppm C	200 ppb	280 ppb	5 μg m ⁻³		
ER684–GF5	2- <i>E</i> -hexenal	Initial:	11.3 ppm C	400 ppb	–	–	1/2 AS + 1/2 SA 26 μg m ⁻³	4 light banks 587 nmol m ⁻³ (H ⁺)air RH: 30 %
		Steady State:	4.3 ppm C	195 ppb	207 ppb	20.1 μg m ⁻³		

* All the initial NO_x was NO.

Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close









Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Chemical structures of 2-*E*-pentenal, 2-*E*-hexenal, and 3-*Z*-hexenal, and of their identified organosulfates with mass spectral data (m/z , elemental formula, and m/z of main product ions).

VOC precursors		
		
2- <i>E</i> -pentenal	2- <i>E</i> -hexenal	3- <i>Z</i> -hexenal
Identified organosulfates	m/z (measured) (formula)	m/z product ions
	229.0021 (C ₅ H ₉ O ₈ S)	MS ² (229): 211, 195, 149, 97 MS ³ (229–149): 131, 103, 75 MS ³ (229–185): 167, 153, 141, 123 MS ³ (229–211): 193, 181, 167, 139, 97
3-sulfoxy-2,4-dihydroxypentanoic acid		
	213.0071 (C ₅ H ₉ O ₇ S)	MS ² (213): 195, 133, 97 MS ³ (213–133): 75
3-sulfoxy-2-hydroxypentanoic acid		
	213.0065 (C ₅ H ₉ O ₇ S)	MS ² (213): 195, 181, 169, 97 MS ³ (213–169): 151, 137, 97 MS ⁴ (213–169–137): 81, 73
2-sulfoxy-3-hydroxypentanoic acid		
	168.9807 (C ₃ H ₅ O ₆ S)	MS ² (169): 151, 125, 97
lactic acid sulfate		
	169.0177 (C ₄ H ₉ O ₅ S)	MS ² (169): 151, 137, 97 MS ³ (169–137): 81, 73
1-sulfoxy-2-hydroxybutane		




Figure 1. Selected LC/MS chromatographic data using regular RP-LC obtained for ambient fine aerosol, including a base peak chromatogram (BPC) and extracted ion chromatograms (EICs) at m/z 215, 169, 213 and 229, corresponding to organosulfates related to isoprene (m/z 215; 2-sulfoxy-1,3,4-trihydroxy-2-methylbutane) and OSs related to 2-*E*-pentenal and indirectly to 2-*E*-hexenal and 3-*Z*-hexenal (m/z 229, 213, and 169). For the molecular structures of the m/z 229, 213 and 169 organosulfates, see Table 2. Abbreviation: NL, normalization level.

Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.




Figure 2. Selected LC/MS chromatographic data (m/z 229 EICs) for **(a)** ambient fine aerosol and **(d)** SOA generated from 2-*E*-pentenal in the presence of acidic seed aerosol, as well as selected MS data (MS² or MS³ product ion spectra) for the peak eluting at 2.5 min in **(b and c)** ambient aerosol and **(e and f)** 2-*E*-pentenal SOA.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 3. Selected LC/MS chromatographic data (m/z 213 EICs) and selected MS data (MS², MS³ and MS⁴ product ion spectra) for ambient fine aerosol (a–j), 2-*E*-pentenal smog chamber SOA (k–p), and 2-*E*-pentenoic acid aqueous-phase reaction products (q–s). Abbreviation: NL, normalization level.

**Characterization of
polar organosulfates
in secondary organic
aerosol**

M. S. Shalamzari et al.




Figure 4. Optimized geometries in the gas-phase using ab-initio methods for **(a)** 3-sulfoxy-2-hydroxypentanoic acid and **(b)** 2-sulfoxy-3-hydroxypentanoic acid. After loss of SO_3 , the negative charge can be accommodated by the carboxyl group in the case of panel **(a)** (see also Scheme 3b).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.





Figure 5. Selected LC/MS chromatographic data (m/z 169 EICs) and selected MS data (MS^2 or MS^3 product ion spectra) for ambient fine aerosol (a–f) and 2-*E*-pentenal SOA (g–l). Abbreviation: NL, normalization level.

Characterization of polar organosulfates in secondary organic aerosol


M. S. Shalamzari et al.



Scheme 1. Proposed fragmentation routes for the m/z 229 organosulfate, related to 2-*E*-pentenal SOA and assigned to 3-sulfoxy-2,4-dihydroxypentanoic acid, on the basis of detailed interpretation of MS^2 , MS^3 , and accurate mass data.

Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.



Scheme 2. Proposed formation pathway for m/z 229 (MW 230) organosulfates related to 2-*E*-pentalenol (and indirectly to 3-*Z*-hexenal and 2-*E*-hexenal SOA), which occur in ambient fine aerosol, and are assigned to 3-sulfoxy-2,4-dihydroxypentanoic acid. The suggested route shares features of the mechanism reported for the formation of 3-sulfoxy-2-hydroxy-2-methylpropanoic acid from methacrolein (Lin et al., 2013).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Characterization of polar organosulfates in secondary organic aerosol


M. S. Shalamzari et al.



Scheme 3. Proposed fragmentation routes for m/z 213 organosulfates, related to 2-*E*-pentenal SOA and assigned to (a) 2-sulfoxy-3-hydroxypentanoic acid and (b) 3-sulfoxy-2-hydroxypentanoic acid, on the basis of detailed interpretation of MS^2 , MS^3 , MS^4 , and accurate mass data. For the fragmentation of the m/z 169 compound, formed by decomposition, see panel (a).

Characterization of polar organosulfates in secondary organic aerosol

M. S. Shalamzari et al.



Scheme 4. Proposed formation pathway for m/z 213 organosulfates related to 2-*E*-pentenal (and indirectly to 3-*Z*-hexenal and 2-*E*-hexenal SOA), which occur in ambient fine aerosol, and are assigned to 2-sulfoxy-3-hydroxypentanoic acid and 3-sulfoxy-2-hydroxypentanoic acid. The suggested mechanism is based on that reported for the formation of 3-sulfoxy-2-hydroxy-2-methylpropanoic acid from methacrolein (Lin et al., 2013).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

