Role of methyl group number on SOA formation from monocyclic aromatic hydrocarbons photooxidation under low NO_x conditions

L. Li^{1,2}, P. Tang^{1,2}, S. Nakao^{1,2,3}, C-L. Chen^{1,2,4} and D. R. Cocker III^{1,2}

5 [1] University of California, Riverside, Department of Chemical and Environmental Engineering, Riverside, CA 92507, USA

[2] College of Engineering-Center for Environmental Research and Technology (CE-CERT), Riverside, CA 92507, USA

[3] Currently at Clarkson University, Department of Chemical and BiomolecularEngineering, Potsdam, NY 13699, USA

[4] Currently at Scripps Institution of Oceanography, University of California, La Jolla, California, USA

Correspondence to: D. Cocker III (dcocker@engr.ucr.edu)

Abstract

- 15 Substitution of methyl groups onto the aromatic ring determines the SOA formation from the monocyclic aromatic hydrocarbon precursor (SOA yield and chemical composition). This study links the number of methyl groups on the aromatic ring to SOA formation from monocyclic aromatic hydrocarbons photooxidation under low NO_x conditions (HC/NO>10 ppbC:ppb). Monocyclic aromatic hydrocarbons with
- 20 increasing numbers of methyl groups are systematically studied. SOA formation from pentamethylbenzene and hexamethylbenzene are reported for the first time. A decreasing SOA yield with increasing number of methyl groups is observed. Linear trends are found in both f₄₄ vs. f₄₃ and O/C vs. H/C for SOA from monocyclic aromatic hydrocarbons with zero to six methyl groups. An SOA oxidation state
- 25 predictive method based on benzene is used to examine the effect of added methyl

groups on aromatic oxidation under low NO_x conditions. Further, the impact of methyl group number on density and volatility of SOA from monocyclic aromatic hydrocarbons is explored. Finally, a mechanism for methyl group impact on SOA formation is suggested. Overall, this work suggests as more methyl groups are

5 attached on the aromatic ring, SOA products from these monocyclic aromatic hydrocarbons become less oxidized per mass/carbon on a basis of SOA yield or chemical composition.

Key Words: Monocyclic aromatic hydrocarbons; Anthropogenic sources; SOA density; SOA composition; Aging; SOA yield

10 **1. Introduction**

Aromatic hydrocarbons are major anthropogenic SOA precursors (Kanakidou, et al., 2005; Henze, et al., 2008). Monocyclic aromatic hydrocarbons with less than four methyl groups are ubiquitous in the atmosphere (Singh et al., 1985; Singh et al., 1992; Fraser, et al., 1998; Pilling and Bartle, 1999; Holzinger, et al., 2001; Buczynska, et al.,

- 15 2009; Hu et al., 2015). Monocyclic aromatic hydrocarbons with more than 4 methyl groups are barely investigated in previous ambient studies, possibly due to a vapor pressure decrease with carbon number (Pankow and Asher, 2008; Table S1). However, a recent study observed that compounds with low vapor pressure are available to evaporate into the atmosphere (Vŏ and Morris, 2012). Monocyclic
- aromatic hydrocarbons with more than three methyl groups contributes to a large portion of components in products such as gasoline and crude oil (Diehl, et al., 2005; Darouich, et al., 2006). Moreover, hydrocarbon reactivity and OH reaction rate constant (k_{OH}) increase with methyl group number (k_{OH} Table S1; Glasson, et al., 1970; Calvert, et al, 2002; Atkinson and Arey, 2003; Aschmann, et al, 2013). OH-
- 25 initiated reactions, particularly OH addition to the aromatic ring, dominate aromatic photooxidation (Calvert, et al., 2002). Hence, photooxidation occurs rapidly once these low vapor pressure aromatic hydrocarbons evaporate into atmosphere. In addition, an increase in carbon number is associated with a decrease in vapor pressure

(Pankow and Asher, 2008). Higher carbon number products with a similar amount of functional groups have a higher tendency to participate in the particle phase. However, aging of organic aerosol is a combination of functionalization, fragmentation and oligomerization (Jimenez, et al., 2009; Kroll, et al., 2009).

5 Therefore, rapid aging does not necessarily lead to the highly oxidized compounds, which serve as an important source of SOA.

Recent studies have found that SOA yields from OH initiated alkane and alkene reactions increase with carbon chain length and decrease with the increase of branched structure (Lim, and Ziemann, 2009; Matsunaga, et al., 2009; Tkacik, et al.,

- 10 2012). However, SOA yield from monocyclic aromatics are found to decrease with carbon number by adding methyl groups to the aromatic ring (Odum, et al., 1997; Cocker, et al., 2001b; Sato, et al., 2012). This indicates that the role of methyl groups on the aromatic ring is different than for alkane and alkene hydrocarbons. Previous studies show that the relative methyl group position determines the alkoxyl radical
- (RO·) fragmentation ratio in alkane and alkene hydrocarbon oxidation (Atkinson, 2007; Ziemann, 2011). Therefore, it is necessary to explore the impact of methyl groups on SOA formation during monocyclic aromatic hydrocarbon oxidation.

Previous studies on SOA formation from monocyclic aromatic hydrocarbon in the presence of NO_x have been conducted at high NO_x levels (e.g., Odum, et al., 1997;

- 20 Cocker, et al., 2001b; Sato, et al., 2012). Ng, et al (2007) observed that SOA yield decreases with increasing carbon number under high NO_x conditions and no trends were observed for no NO_x conditions. Reaction mechanisms vary for different NO_x conditions (e.g., Song, et al., 2005; Kroll and Seinfeld, 2008) and thus impact SOA chemical composition. Therefore, it is necessary to investigate methyl group impact
- 25 on urban SOA formation from monocyclic aromatic hydrocarbon under more atmospherically relevant low NO_x conditions.

SOA budget underestimation of the urban environment is associated with mechanism uncertainty in aromatic hydrocarbon photooxidation and possibly missing aromatic

hydrocarbon precursors (Henze, et al., 2008; Hallquist, et al., 2009). Previous chamber studies seldom investigate SOA formation from monocyclic aromatic with more than 3 methyl groups (e.g. pentamethylbenzene and hexamethylbenzene). This study investigates SOA formation from the photooxidation of seven monocyclic

5 aromatic hydrocarbon (ranging from benzene to hexamethylbenzene) under the low NO_x (HC/NO>10 ppbC:ppb) condition. The impact of methyl group number on SOA yield, chemical composition and other physical properties are demonstrated. Possible methyl group impact on aromatic ring oxidation, decomposition and subsequent oligomerization are discussed.

10 **2. Method**

2.1 Environmental chamber

All experiments were conducted in the UC Riverside/CE-CERT indoor dual 90 m³ environmental chambers, which are described in detail elsewhere (Carter et al., 2005). All experiments were conducted at dry conditions (RH<0.1%), in the absence of

- 15 inorganic seed aerosol and with temperature controlled to 27±1°C. Two movable top frames were slowly lowered during each experiment to maintain a slight positive differential pressure (~0.02" H₂O) between the reactors and enclosure to minimize dilution and/or contamination of the reactors. 272 115 W Sylvania 350BL blacklights are used as light sources for photooxidation.
- 20 A known volume of high purity liquid hydrocarbon precursors (benzene Sigma-Aldrich, 99%; toluene Sigma-Aldrich, 99.5%; *m*-xylene Sigma-Aldrich, 99%; 1, 2, 4-trimethylbenzene Sigma-Aldrich, 98%) were injected through a heated glass injection manifold system and flushed into the chamber with pure N₂. A glass manifold packed with glass wool inside a temperature controlled oven (50-80 °C) is used to inject solid
- hydrocarbon precursors (1, 2, 4, 5-tetramethylbenzene Sigma-Aldrich, 98%;
 pentamethylbenzene Sigma-Aldrich, 98%; hexamethylbenzene Sigma-Aldrich, 99%).
 NO was introduced by flushing pure N₂ through a calibrated glass bulb filled to a

predetermined partial pressure of pure NO. All hydrocarbons and NO are injected and well mixed before the lights were turned on to commence the reaction.

2.2 Particle and Gas Measurement

Particle size distribution between 27 nm and 686 nm was monitored by dual custom

- 5 built Scanning Mobility Particle Sizers (SMPS) (Cocker et al., 2001a). Particle effective density was measured with a Kanomax Aerosol Particle Mass Analyzer (APM-SMPS) system (Malloy et al., 2009). Particle volatility was measured by a Volatility Tandem Differential Mobility Analyzer (VTDMA) (Rader and McMurry, 1986) with a Dekati® Thermodenuder controlled to 100°C and a 17 s heating zone
- 10 residence time (Qi, et al., 2010b). Volume fraction remaining (VFR) is calculated as $(D_{p, after TD}/D_{p, before TD})^3$.

Evolution of particle-phase chemical composition was measured by a High Resolution Time of Flight Aerosol Mass Spectrometer (HR-ToF-AMS; Aerodyne Research Inc.) (Canagaratna et al., 2007; DeCarlo et al., 2006). The sample was vaporized by a

- 15 600 °C oven followed by a 70 eV electron impact ionization. f_x in this study is calculated as the fraction of the organic signal at m/z=x. For example, f_{44} and f_{43} are the ratios of the organic signal at m/z 44 and 43 to the total organic signal, respectively (Chhabra et al., 2011; Duplissy et al., 2011). Elemental ratios for total organic mass, oxygen to carbon (O/C), and hydrogen to carbon (H/C) were
- determined using the elemental analysis (EA) technique (Aiken et al., 2007, 2008).
 Data was analyzed with ToF-AMS analysis toolkit squirrel 1.56D /PIKA 1.15D version.

The Agilent 6890 Gas Chromatograph – Flame Ionization Detector was used to measure aromatic hydrocarbon concentrations. A Thermal Environmental Instruments

25 Model 42C chemiluminescence NO analyzer was used to monitor NO, NO_y-NO and NO_y. The gas-phase reaction model SAPRC-11 developed by Carter and Heo (2012) was utilized to predict radical concentrations (·OH, HO₂·, RO₂· and NO₃·).

3. Results

3.1 SOA yield relationship with methyl group number

SOA yields from the photooxidation of seven monocyclic aromatic hydrocarbons are calculated as the mass based ratio of aerosol formed to hydrocarbon reacted (Odum, et

- al., 1996). The HC/NO ratio ranged from 12.6-110 ppbC:ppb for all experiments used in this study. Experiment conditions and SOA yield are listed from the current work (Table 1) along with additional *m*-xylene experiment conditions from previous studies (Table S2) (Song, et al, 2005) in the UCR CE-CERT chambers. SOA yield as a function of particle mass concentration (M₀) for all seven monocyclic aromatic
- 10 precursors (Fig.1) includes experiments listed in both Table 1 and Table S2. Each individual experiment is marked and colored by the number of methyl groups on each precursor aromatic ring. It is observed that SOA yield decreases as the number of methyl groups increases (Fig. 1). A similar yield trend is also observed in previous studies on SOA formation from monocyclic aromatic hydrocarbons, however,
- different absolute yield values are found, presumably due to higher NO_x levels
 (Odum, et al., 1997b; Kleindienst et al., 1999; Cocker et al., 2001b; Takekawa et al., 2003; Ng, et al., 2007; Sato, et al., 2012). SOA yields of benzene under comparable low NO_x conditions are higher than that in Sato, et al (2012), Borrás and Tortajada-Genaro (2012) and Martín-Reviejo and Wirtz (2005).

20 The two product semi-empirical model described by Odum, et al. (1996) is used to fit SOA yield as a function of M₀. Briefly, the two product model assumes that aerosol forming products can be lumped into lower and higher volatility groups whose mass fraction is defined by α_i and a partitioning parameter K_{om, i} (m³·µg⁻¹) described extensively in Odum, et al. (1996). Each monocyclic aromatic hydrocarbon is fitted

individually except for those with methyl group number greater than or equal to 4, which are grouped as C₁₀₊. The experimental fitting parameters (α₁, K_{om, 1}, α₂ and K_{om, 2} in Table 2) in the two product model were determined by minimizing the sum of the squared of the residuals. The higher-volatility partitioning parameter parameter

 $(K_{om,2})$ in all yield curve fitting are assigned to a fixed value by assuming similar high volatile compounds are formed during all monocyclic aromatic hydrocarbon photooxidation experiments. Benzene has much higher mass-based stoichiometric coefficients (α_2) than the other monocyclic aromatic compounds indicating that the

- 5 pathway leading to higher volatility products formation is favored. The lower volatility partitioning parameters (K_{om, 1}) vary widely for each monocyclic aromatic yield fitting curve. Benzene has the lowest K_{om, 1}, toluene has the highest K_{om, 1}, and the rest of monocyclic aromatics have similar mid-range K_{om, 1} values. The extremely low K_{om, 1} of benzene indicates that pathways associated with significant volatility
- 10 decrease occur far less during benzene photooxidation than for monocyclic aromatic compounds with methyl groups. Further, $K_{om, 1}$ is much lower in multi-methyl group monocyclic aromatic hydrocarbons (with the exception of toluene) while α_1 decreases with methyl group number. This suggests that increasing methyl group number on the aromatic ring suppresses formation of lower volatility products therefore lowering the
- 15 mass based aerosol yield. This suggests that monocyclic aromatics with more methyl groups are less oxidized per mass since the methyl group carbon is not well oxidized compared with the ring carbon.

The aromatic SOA growth curves (particle concentration M_0 vs. hydrocarbon consumption Δ HC) under similar HC/NO_x are shown in Fig. S1. The slope of the

- 20 growth curve is negatively correlated with the parent aromatics reaction rate (k_{OH}). This observation contrasts with a previous study that observed positive correlation between SOA formation rate and hydrocarbon reaction rate for systems where initial semivolatile products dominate gas-particle phase partitioning (Chan, et al., 2007). The reverse relationship observation in this study indicates that the effect of methyl
- 25 group number on SOA yield is greater than that of the increasing k_{OH} on SOA yield. There are two possibilities for the methyl group number effect: 1) the methyl group facilitates initial semivolatile products to react into more volatile compounds; 2) the methyl group prevents further generation semivolatile products formation by stereo-

hindrance. Therefore, the methyl group increases hydrocarbon mass consumption more than particle mass formation.

The relationship between radical levels and SOA yield was also analyzed. Table S3 lists modeled individual average radical concentrations throughout photooxidation

- 5 while Table S4 lists the correlation between SOA yields and individual average radical concentrations. None of the radical parameters (e.g. ·OH/HO₂·, HO₂·/ RO₂· etc.) is strongly correlated with SOA yield. Average OH radical concentration is the only parameter investigated with a statistically significant correlation (p<0.05), as k_{OH} varies with aromatic species and lower average OH concentrations are present
- with higher k_{OH}. Fig. S2 shows the time evolution of [·OH], [RO₂·] and [HO₂·] for different aromatic precursors under similar initial aromatic and NO_x loadings. Higher [·OH] is observed for aromatic precursors with lower k_{OH} while peroxide radicals ([RO₂·] and [HO₂·]), which depend on both k_{OH} and [·OH], are similar for all precursors. This suggests that SOA mass yield is determined by precursor structure
- 15 rather than gas-phase oxidation state since radical conditions for each aromatic hydrocarbon are comparable and $[RO_2 \cdot]$ and $[HO_2 \cdot]$ reactions are expected to determine SOA formation (Kroll and Seinfeld, 2008)

3.2 SOA chemical composition relationship with methyl group number

3.2.1 f44 vs f43

- Organic peaks at *m/z* 43 and *m/z* 44 are key fragments from AMS measurement toward characterization of oxygenated compounds in organic aerosol (Ng, et al., 2010; Ng, et al., 2011). A higher f₄₄ and a lower f₄₃ indicates a higher degree of oxidation (Ng et al., 2010, 2011). The f₄₄ and f₄₃ evolution during SOA formation from different monocyclic aromatic hydrocarbon photooxidation is shown for low NO_x
- conditions (Fig. 2). Each marker type represents an individual monocyclic aromatic hydrocarbon with the marker colored by photooxidation time (light to dark). The f_{44} and f_{43} range are comparable to previous chamber studies with slight shift due to

differences in initial conditions (e.g. NO_x etc.) (Ng, et al., 2010; Chhabra, et al., 2011; Loza, et al., 2012; Sato, et al., 2012). SOA compositions from monocyclic aromatic hydrocarbon photooxidation under low NO_x are in the LV-OOA and SV-OOA range of the f₄₄ vs f₄₃ triangle (Ng, et al., 2010) with those from benzene on the left side,

toluene inside and other monocyclic aromatics on the right side of the triangle confirming that laboratory SOA f₄₄ vs f₄₃ is precursor dependent (Chhabra, et al., 2011). Evolution of SOA composition (Heald, et al., 2009; Jimenez, et al., 2009) refers to SOA chemical composition changes with time and f₄₄ and f₄₃ evolution refers to the change of f₄₄ and f₄₃ with time. Significant f₄₄ and f₄₃ evolution is observed for
benzene and slightly for toluene, *m*-xylene and tetramethylbenzene.

In this work, average f_{44} and f_{43} are examined to demonstrate the methyl group impact on SOA chemical composition from monocyclic aromatic hydrocarbons. Average f_{44} vs f_{43} is marked with the aromatic compound name in Fig. 2. Generally decreasing f_{44} and increasing f_{43} are observed with increasing number of methyl groups on the

- 15 aromatic ring. Similar trends are also observed in previous studies (Ng, et al., 2010; Chhabra et al., 2011; Sato, et al., 2012). The f_{44} vs f_{43} trend is quantified by linear curve fitting (f_{44} = -0.58 f_{43} + 0.19, R₂ = 0.94). f_{28} is assumed to be equal to f_{44} in the AMS frag Table of Unit Resolution Analysis, which describes the mathematical formulation of the apportionment at each unit resolution sticks to aerosol species,
- 20 based on ambient studies (Zhang, et al., 2005; Takegawa et al., 2007) and $\text{CO}^+/\text{CO}_2^+$ ratio for SOA from aromatic oxidation is found around ~1 (0.9-1.3) (Chhabra et al., 2011). The slope of ~0.5 indicates that $2\Delta f_{44}$ =- Δf_{43} or $\Delta (f_{28}+f_{44})$ =- Δf_{43} in SOA formed from monocyclic aromatic hydrocarbons with different numbers of methyl groups. The CO₂⁺ fragment ion at *m/z* 44 and C₂H₃O⁺ fragment ion at *m/z* 43 are two
- 25 major AMS fragmentation ions from aromatic secondary organic aerosol. No significant C₃H₇⁺ is observed at *m/z* 43. CO₂⁺ represents oxidized aerosol and is associated with carboxylic acids (Alfarra, et al., 2004, Aiken, et al., 2007, Takegawa, et al., 2007; Canagaratna, et al., 2015) while C₂H₃O⁺ is associated with carbonyls (McLafferty and Turecek, 1993; Ng, et al., 2011). The CO⁺ fragment ion at *m/z* 28

can originate from carboxylic acid or alcohol (Canagaratna, et al., 2015). The $\Delta(f_{28}+f_{44}) = -\Delta f_{43}$ relationship observed in this study imply that adding the methyl group to the aromatic ring changes SOA from CO_2^+ to $C_2H_3O^+$ implying a less oxidized SOA chemical composition in AMS mass fragments. While bicyclic

hydrogen peroxides are considered to be the predominant species in aerosol phase from monocyclic aromatic photooxidation (Johnson, et al., 2004, 2005; Wyche, et al., 2009; Birdsall et al., 2010; Birdsall and Elrod, 2011; Nakao, et al., 2011), they are less likely to contribute to the CO₂⁺ ion fragment. Possible mechanisms to produce SOA products that form the CO₂⁺ fragments as well as produce C₂H₃O⁺ fragments by
adding methyl group are described in detail in Section 4.

3.2.2 H/C vs O/C

Elemental analysis (Aiken, et al., 2007, 2008) is used to elucidate SOA chemical composition and SOA formation mechanisms (Heald, et al., 2010; Chhabra, et al., 2011). Fig.3a shows the H/C and O/C time evolution of average SOA formed from

- 15 hydrocarbon photooxidation of various monocyclic aromatics under low NO_x conditions (marked and colored similarly to Fig.2). The H/C and O/C ranges are comparable to previous chamber studies with slight shift due to difference in initial conditions (e.g. NO_x etc.) (Chhabra, et al., 2011; Loza, et al., 2012; Sato, et al., 2012). All data points are located in between slope=-1 and slope=-2 (Fig. 3a, lower left
- 20 corner, zoom out panel). This suggests that SOA components from monocyclic aromatic photooxidation contain both carbonyl (ketone or aldehyde) and acid (carbonyl acid and hydroxycarbonyl) like functional groups. These elemental ratios also confirm that SOA formed from monocyclic aromatic hydrocarbon photooxidation under low NO_x are among the LV-OOA and SV-OOA regions (Ng, et al., 2011). The
- 25 change of elemental ratio (H/C and O/C) with time is referred as elemental ratio evolution. The elemental ratio evolution agrees with the f₄₄ vs f₄₃ evolution (significant evolution in benzene and slightly for toluene, *m*-xylene and 1,2,4,5-tetramethylbenzene). This study concentrates on average H/C and O/C in order to

demonstrate the methyl group impact on SOA chemical composition from monocyclic aromatic hydrocarbons.

Average H/C and O/C location is marked (Fig. 3a) for each aromatic compound by name. It is observed that H/C and O/C from SOA formed from *m*-xylene, 1, 2, 4-

- 5 trimethylbenzene and monocyclic aromatics with more than three methyl groups are similarly distributed in H/C vs O/C. A general decrease in O/C and an increase in H/C are noted as the number of methyl groups on the aromatic ring increase, which is consistent with other studies (Chhabra, et al., 2011; Sato, et al., 2012). The trend indicates that monocyclic aromatics are less oxidized per carbon as the number of
- 10 methyl groups increases, which can be attributed to less oxidation of the methyl groups compared to the aromatic ring carbons. The elemental ratio trends (O/C decreases and H/C increases as the number of methyl groups increases) are also consistent with the decreasing yield trends with increasing the number of methyl groups (Section 3.1), suggesting that SOA yield is dependent on SOA chemical
- 15 composition. It should also be noticed that the higher O/C and lower H/C observed in SOA formed from 1,2,4-trimethylbenzne (three methyl substitute aromatic hydrocarbon) than that from *m*-xylene (two methyl substitute aromatic hydrocarbon) is due to the isomer impact on SOA chemical composition, which is discussed in details by Li, et al (2015). Further, the yield and O/C ratio agrees with recent findings
- 20 that O/C ratio is well correlated to aerosol volatility (Section 3.3.2) (Cappa, et al., 2012, Yu, et al., 2014) thereby affecting the extent of gas to particle partitioning. The H/C vs O/C trend linear curve (H/C= -1.34 O/C + 2.00, $R^2 = 0.95$) shows an approximately -1 slope with a y-axis (H/C) intercept of 2. The H/C vs O/C trend slope observed in this work is similar to the toluene and *m*-xylene elemental ratio slope
- 25 observed under high NO_x and H_2O_2 only conditions observed in Chhabra et al (2011).

The Van Krevelen diagram can also be used to analyze the oxidation pathway from initial SOA precursor to final SOA chemical composition by comparing the initial H/C and O/C ratios from the precursor hydrocarbon to the final SOA H/C and O/C ratios. Fig. 3b shows the aromatic precursor location on the left (texted with aromatic

hydrocarbon name and colored by methyl group number) and average SOA chemical composition on the right. The SOA H/C increase in the final SOA chemical composition follows the initial aromatic precursor elemental ratio trend. A large O/C increase with a slight H/C increase is observed moving from precursor to SOA

- 5 composition. SOA formation from hydroperoxide bicyclic compounds contributes to O/C increases without loss of H. The slight H/C increase might result from hydrolysis of ring opened product oligomerization (Jang and Kamens, 2001; Jang et al., 2002; Kalberer, et al., 2004; Sato, et al., 2012). A slight H/C decrease rather than increase is observed in the hexamethylbenzene data suggesting that the six methyl groups
- 10 sterically inhibits certain reaction mechanism (eg. hydrolysis) to obtain H.

3.2.3 OSc and its prediction

O/C alone may not capture oxidative changes as a result of breaking and forming of bonds (Kroll, et al., 2009). Oxidation state of carbon (OS_c \approx 2O/C-H/C) was introduced into aerosol phase component analysis by Kroll et al. (2011). It is

- 15 considered to be a more accurate metric for describing oxidation in atmospheric organic aerosol (Ng et al., 2011; Canagaratna, et al., 2015; Lambe, et al., 2015) and therefore better correlated with gas-particle partitioning (Aumont, et al., 2012). Average SOA OS_c in this study ranges from -0.9 to 0.3 for monocyclic aromatic photooxidation under low NO_x conditions (Fig. 4b) and is comparable to previous
- studies (Kroll, et al., 2011 (toluene, *m*-xylene and trimethylbenzene); Sato, et al., 2012 (benzene and 1,3,5-trimethylbenzene)). OS_c observed is consistent with OS_c observed in field studies (Kroll, et al., 2011) especially in urban sites (e.g. $-1.6 \sim 0.1$, Mexico City) and supports the major role of monocyclic aromatic precursors in producing anthropogenic aerosol. Average SOA OS_c values are consistent with the LV-OOA and
- 25 SV-OOA regions (Ng, et al., 2011; Kroll, et al., 2011). OS_c only increases with oxidation time for benzene photooxidation $(0.2 \sim 0.4)$.

The methyl group substitute (-CH₃) affects O/C and H/C ratios by increasing both carbon and hydrogen number as they relate to SOA OS_c. It is hypothesized here that

the methyl group impacts remain similar in SOA elemental ratios as they do in the aromatic precursor (-CH₃ dilution effect). This would imply that the methyl group effect on SOA elemental ratio and OS_c from monocyclic aromatic hydrocarbons is predictable from benzene oxidation. Eq-1 and Eq-2 show the prediction formula for

5 O/C and H/C, respectively, where *i* represents the methyl group number on the monocyclic aromatic precursor, O/C_{benzene_SOA} and H/C_{benzene_SOA} are the measured O/C and H/C in SOA from benzene photooxidation experiments.

$$O/C_{pre,i_SOA} = \frac{6}{i+6} (O/C_{benzene_SOA})$$
 Eq-1

$$H/C_{pre,i_SOA} = \frac{2i}{i+6} + \frac{6}{i+6} (H/C_{benzene_SOA})$$
Eq-2

- Fig. 4a shows a comparison of measured (red) and predicted (green) H/C and O/C location marked with corresponding SOA precursor methyl groups. The difference between predicted and measured H/C and O/C ranges from -6.4~1.2% and -11.8~20.9%, respectively. However, the predicted H/C vs O/C line (Eq-1 & Eq-2) is H/C= -1.38O/C +2.00. This is comparable to a measured data fitting line (Section 3.2.2 H/C=
- 15 -1.34 O/C + 2.00, $R_2 = 0.95$). Predicted OS_c is then calculated based on the predicted H/C and O/C. Fig. 4b compares measured (red) and predicted (green) OS_c. The largest O/C and OS_c overestimation is observed in *m*-xylene (marked as 2 in Fig. 4a, bar 2 in Fig. 4b). This could be explained by the isomer selected for the two methyl group monocyclic aromatic hydrocarbon (*m*-xylene). A detailed analysis on isomer structure
- 20 impact on SOA chemical composition is found in Li, et al (2015). The largest O/C and OS_c underestimation is observed in hexamethylbenzene (marked as 6 in Fig. 4a, bar 6 in Fig. 4b). This suggests that the methyl groups attached to every aromatic carbon exert a steric inhibition effect on certain aromatic oxidation pathways, thus leading to increased importance of aerosol formation from other reaction pathways (possibly
- 25 fragmentation Kroll, et al., 2011, see Section 4) to form SOA. It is also noticed that O/C and OS_c is slightly overestimated in SOA formed from pentamethylbenzene. This indicates that the methyl group hindrance impact on aromatic hydrocarbon oxidation

should be explained by multiple pathways which have different impact on SOA formation.

The correlation between organic mass loading and chemical composition is also analyzed. Organic mass loading is well correlated (Pearson correlation) with chemical

- 5 composition parameters including f₄₄ (0.907), f₄₃ (-0.910), H/C (-0.890) and O/C (0.923) (Fig. S3). However, previous studies show that O/C and f₄₄ decrease as organic mass loading increases (Shilling et al., 2009; Ng, et al., 2010; Pfaffenberger; 2013). The findings of this study indicate that molecular species drive SOA chemical composition rather than organic mass. The positive trend between f₄₄ and organic
- 10 mass loading is driven by benzene and toluene experiments (Fig. S3) where the high mass loading results are concurrent with high f₄₄ results. However, the f₄₄ change with mass loading increase during benzene and toluene photooxidation is less significant compared with the f₄₄ difference caused by number of methyl groups on aromatic ring. Moreover, no significant correlation was found between mass loading and f₄₄ or O/C
- 15 when compared under similar mass loadings (including f₄₄ at low mass loading time point of toluene and benzene photoxidation). Organic nitrate accounts for less than 10% organic in SOA components in all monocyclic aromatic hydrocarbon photooxidation experiments in this work according to AMS measurement and will not be discussed.

20 **3.3 Physical property relationship with methyl group number**

3.3.1 SOA Density

SOA mass density is a fundamental parameter in understanding aerosol morphology, dynamics, phase and oxidation (De Carol, et al., 2004; Katrib, et al., 2005; Dinar, et al., 2006; Cross, et al., 2007). SOA density ranged from 1.24-1.44 g/cm³ for all

25 aromatic-NO_x photooxidation experiments in this study. The range is comparable to previous studies under similar conditions (Ng, et al; 2007; Sato, et al., 2010; Esther Borrás, et al., 2012). A general decreasing density trend is found with increasing

methyl group number on precursor aromatic rings (see Fig. 5a). Correlation between SOA density and chemical composition was statistically analyzed (Table S5). Besides the strong correlation with methyl group number (-0.943, Fig. 5a), SOA density was also well correlated with O/C ratio (0.873, Fig. 5b) and other measures of bulk

5 chemical composition (Table S5). Bahreini et al (2005) reported a density increase trend with f₄₄ in other compounds while Pang, et al. (2006) found that SOA density increases with O/C ratio. Kuwata, et al., 2011(Eq-3) and Nakao, et al., 2013 suggested a quantified relationship between SOA density and SOA elemental ratio. Eq-3 developed by

10
$$\rho = \frac{12 + H/C + 16 \times O/C}{7 + 5 \times H/C + 4.15 \times O/C}$$
 Eq-3

Kuwata, et al. (2011) is used in this work to predict density based on elemental ratio in order to explore the methyl group impact on SOA formation. Fig. 5c shows a good agreement between predicted and measured SOA densities (- $6.58\% \sim 10.42\%$). However, SOA density difference between prediction and measurement change from

- 15 positive (0&1 methyl group) to negative (2, 3, 4 or 5 methyl groups) with increasing methyl group number (except hexamethylbenzene) implying that the increase of methyl groups promotes mechanism(s) leading to changes in the ratio of several key organic fragments (e.g., m/z 28 : m/z 44) thereby challenging the applicability of the default fragment table for elemental ratio analysis. It is possible that CO⁺/CO₂⁺ and
- 20 H_2O^+/CO_2^+ ratios are different in SOA formed from different aromatic precursors. Nakao, et al (2013) shows that H_2O^+/CO_2^+ increases with methyl group number due to the constant H_2O^+ fraction and a decrease in CO_2^+ fraction. Canagaratna, et al (2015) demonstrated that CO^+/CO_2^+ and H_2O^+/CO_2^+ are underestimated in certain compounds (especially alcohols). Assuming that the major impact of methyl group on SOA
- 25 composition is to change –COOH to -COCH₃ (or other cyclic isomers), f_{CO2+} will decrease but H₂O⁺ and CO⁺ fraction might not change linearly. The alcohol contribution to CO⁺/CO₂⁺ and H₂O⁺/CO₂⁺ gradually grows as the methyl group prevents acid formation. Therefore, AMS measurements might underestimate O/C.

This is consistent with the density prediction from elemental ratios where a change of error from positive to negative is seen as the number of methyl group change from less than two to two or more than two, with the exception of hexamethylbenzene. This might relate to the difference in SOA formation pathways due to steric hindrance of the six methyl groups during hexamethylbenzene oxidation.

3.3.2 SOA Volatility

5

SOA volatility is a function of oxidation, fragmentation, oligomerization and SOA mass (Kalberer, et al., 2004; Salo, et al., 2011; Tritscher, et al., 2011; Yu, et al., 2014). Bulk SOA volatility can be described by the VFR after heating SOA to a fixed

- 10 temperature in a thermodenuder. VFRs for SOA formed early in the experiment are around 0.2 for all monocyclic aromatic precursors and then increase as the experiment progresses. Increasing VFR indicates the gas to particle partitioning of more oxidized products, which may include oligomerization products formed during aromatic photooxidation. The VFR trends and ranges are comparable to previous studies
- 15 (Kalberer et al., 2004; Qi et al., 2010a; Qi et al., 2010b; Nakao et al., 2012). Fig. 6a shows the relationship between SOA precursor methyl group number and SOA VFR at the end of the experiment (VFR_{end}). VFR shows a significant decreasing trend with increasing methyl group number from benzene to 1, 2, 4, 5-tetramethylbenzene. This implies that volatility of SOA-forming products increases as the number of methyl
- 20 groups on the aromatic ring increases. There is also a slight increase in VFR from 1, 2, 4, 5-tetramethylbenzene to hexamethylbenzene; however, VFR in SOA formed from all C₁₀₊ group aromatics is lower than that of 1, 2, 4-trimethylbenzene. The changing VFR trend suggests that chemical components contributing to SOA formation become different when more than four methyl groups are attached to a
- 25 single aromatic ring. A positive correlation (0.755, p=0.05) found between mass loading and VFR_{end} implies that the lower the volatility in the products formed from aromatic hydrocarbons, the higher the SOA mass concentration. An opposite correlation between mass loading and VFR is found in previous studies due to the

partitioning of more volatile compounds to the particle phase at high mass loading (Tritscher, et al., 2011;Salo, et al., 2011). Therefore, mass loading does not directly lead to the VFR trend in the current study, rather it is the methyl group number in the SOA precursor that affects the composition of SOA and therefore the monocyclic

- aromatic hydrocarbon yield (section 3.1) and volatility. The correlation between SOA volatility (VFR) and chemical composition is statistically analyzed (Table S5). O/C (0.937, p = 0.002) and OS_c (0.932, p = 0.02) have the highest correlation with VFR_{end}. Previous studies also observed that lower aerosol volatility is correlated to higher O/C ratio (Cappa, et al., 2012, Yu, et al., 2014) and OS_c (Aumont, et al., 2012; Hildebrandt
- Ruiz, et al., 2014). Fig. 6b and Fig. 6c illustrate the VFR_{end} and O/C or OS_c
 relationship among all the monocyclic aromatic precursors investigated in this study.
 Benzene and toluene are located on the right upper corner in both graphs suggesting that significantly more oxidized and less volatile components are formed from monocyclic aromatic precursors with less than two methyl groups. The VFR_{end} and
- 15 chemical components relationship becomes less significant when only monocyclic aromatic precursors with more than two methyl groups are considered.

4. Discussion

4. 1 SOA formation pathway from monocyclic aromatic hydrocarbon

- Bicyclic peroxide compounds are considered to be important SOA forming products
 from monocyclic aromatic photooxidation (Johnson, et al., 2004, 2005; Song, et al., 2005; Wyche, et al., 2009; Birdsall et al., 2010; Birdsall and Elrod, 2011; Nakao, et al., 2011). However, the significant CO₂⁺ fragment (f₄₄) observed for SOA by the AMS indicates a contribution of an additional pathway to SOA formation from monocyclic aromatic hydrocarbon photooxidation since it is unlikely that bicyclic
 peroxides could produce a CO₂⁺ in the AMS. Hydrogen abstraction from the methyl group is not further discussed here as it accounts for less than 10% monocyclic aromatic oxidation pathway (Calvert, et al., 2002). However, it is important to
 - consider the further reaction of bicyclic peroxide ring scission products, especially in

the presence of NO_x (Jang and Kamens, 2001; Atkinson and Arey, 2003; Song, et al., 2005; Hu, et al., 2007; Birdsall and Elrod, 2011; Carter, et al., 2013). First generation ring scission products include 1, 2-dicarbonyls (glyoxal and methylglyoxal) and unsaturated 1, 4-dicarbonyls (Forstner, et al., 1997; Jang and Kamens, 2001; Birdsall

- 5 and Elrod, 2011). These dicarbonyls are small volatile molecules that are unlikely to directly partition into the particle phase. However, these small molecules can potentially to grow into low volatility compounds through oligomerization. Previous studies suggest that oligomerization can be an important pathway for SOA formation from monocyclic aromatic precursors (Edney, et al., 2001; Baltensperger, et al., 2005;
- Hu, et al., 2007; Sato, et al., 2012). While Kalberer, et al (2004) proposed an oligomerization pathway of 1, 2-dicarbonyls, Arey, et al (2008) found that unsaturated 1, 4-dicarbonyls have a higher molar yield than 1, 2-dicarbonyls in OH radical-initiated reaction of monocyclic aromatic hydrocarbons. Further, OH radical reaction and photolysis rates are observed to be lower in 1, 2-dicarbonyls photolysis (Plum, et
- al., 1983; Chen, et al., 2000; Salter, et al., 2013; Lockhart, et al; 2013) than
 unsaturated 1, 4-dicarbonyls (Bierbach, et al., 1994; Xiang, et al; 2007). This suggests
 that secondary reaction of unsaturated 1, 4-dicarbonyls is more important than that of
 1, 2-dicarbonyls. Previous studies have found that unsaturated 1, 4-dicarbonyls react
 to form small cyclic furanone compounds (Jang, et al., 2001; Bloss, et al., 2005;
- 20 Aschmann, et al., 2011). Oligomerization is possible for these small cyclic compounds based on their similar molecular structure with glyoxal and methylglyoxal (Fig. 7 c-2-1 & c-2-2 pathway, Fig. S4). Products from further oligomerization of ring opening compounds can also partition into the aerosol phase and contribute to SOA formation. Hydrolysis is necessary in both oligomerization pathways (Fig. S4 and
- 25 Kalberer, et al., 2004), which is consistent with the slight H/C increase observed for most monocyclic aromatic hydrocarbon photooxidation results in this study. However, Nakao et al. (2012) showed that the glyoxal impact on SOA formation is majorly due to OH radical enhancement with glyoxal instead of oligomerization, especially under dry conditions. This indicates that oligomerization from small cyclic furanone is more

30 likely to contribute more to SOA formation than 1, 2- dicarbonyl in this work. Other

pathways reported in previous studies are also possible to contribute to SOA formation here (Edney, et al., 2001 (polyketone); Jang and Kamens, 2001 (aromatic ring retaining products; six and five member non-aromatic ring products; ring opening products); Bloss, et al., 2005 (benzoquinone, epoxide, phenol); Carter, 2013 (bicyclic

5 hydroperoxide)). Our work only addresses differences in the oligomerization pathway contribution to form SOA from monocyclic aromatic hydrocarbons.

A simplified monocyclic aromatic oxidation mechanism for low NO_x conditions is shown (Fig. 7 & Fig. 8; the figures only illustrate monocyclic aromatic oxidation related to particle formation). Fig. 7 illustrates the oxidation, fragmentation and

- 10 oligomerization after initial OH addition to aromatic ring, Fig. 8 shows the kinetic scheme for SOA formation from monocyclic aromatic hydrocarbons. S₁, S₂ and S₃ represent bicyclic hydroperoxide compounds, ring opening compounds and oligomerization products, respectively. Table S6 summarizes the predicted vapor pressures of the benzene photooxidation products using SIMPOL (Pankow and Asher,
- 15 2008). The bicyclic hydroperoxide (S₁, Fig. 8) is more volatile than the oligomers (S₃ in Fig. 8). The volatilities of the bicyclic hydroperoxides are sufficiently high to allow additional oxidation (e.g. add one more hydroperoxide functional group to form $C_6H_6O_8$). The further oxidized bicyclic hydroperoxide vapor pressure is predicted to be similar to oligomerization products from reaction of c-2-1 (Fig. S4) with glyoxal.
- 20 The higher vapor pressure of oligomer products from glyoxal as compared to oligomers from other products indicates that bicyclic hydroperoxides (S₁) contribute more to SOA formation in benzene than oligomerization products (S₃), especially at higher particle mass loadings, as compared with monocyclic aromatic hydrocarbons containing methyl groups according to the two product model fitting (Fig. 1 and
- 25 Table. 2).

4. 2 Methyl group number impact on SOA formation pathway from monocyclic aromatic hydrocarbon

It is observed that as the number of methyl groups on the monocyclic aromatic precursor increases, mass yield (Section 3.1), overall oxidation per carbon (Section

- 5 3.2), and SOA density all decrease and SOA volatility increases. The observed yield trend is attributed to the increasing methyl group number enhancing aromatic fragmentation and inhibiting oligomerization. First, the methyl group stabilizes the ring opening radical (Atkinson, 2007; Ziemann, 2011), thus favoring the ring opening pathway. Second, the methyl group hinders cyclic compound formation and
- 10 oligomerization (Fig. 7). Oligomerization is unlikely to occur directly from non-cyclic dicarbonyls (Kalberer, et al, 2004) or indirectly from cyclic compounds formed by unsaturated dicarbonyls (Fig. S5) with increasing methyl group number. Methyl groups both inhibit oligomerization (Fig.7 (c-1-3)) and prevent the formation of cyclic compounds from unsaturated dicarbonyls (Fig.7 (c-2-3)) when methyl groups are
- 15 attached to both ends of an unsaturated dicarbonyl. Oligomerization is possible for these ketones through reactions such as aldol condensation and hemiacetal formation (Jang et al., 2002) under acidic conditions. However, this is less favored for the current study in the absence of acidic seeds. Hence, less cyclic compounds are available for subsequent oligomerization, leading to more volatile products and a
- 20 decrease in SOA formation. Moreover, the SOA composition trend is well explained by a –CH₃ dilution effect. Previous studies on the different gas phase (Forstner, et al., 1997; Yu, et al., 1997) and particle phase (Hamilton, et al., 2005; Sato, et al., 2007; Sato, et al., 2012) products supports this methyl group dilution theory. A typical example is that more 3-methyl-2,5-furandione is observed in *m*-xylene than toluene
- and vice versa for 2,5-furandione. Sato et al. (2010) suggests that more low-reactive ketones are produced rather than aldehydes with increasing number of substituents.
 However, most ketones or aldehydes detected are so volatile that they mostly exist in the gas phase (Forstner, et al., 1997; Yu, et al., 1997; Cocker, et al., 2001b; Jang and

Kamens, 2001). Taken collectively, this implies the importance of oligomerization and methyl substitutes on SOA formation.

The observation of a slight H/C decrease from hexamethylbenzene to its SOA components in contrast with the increasing trend for monocyclic aromatic

- 5 photooxidation for zero to five methyl group substitutes (Section 3.2.2) suggests that hydrolysis followed by oligomerization might not be significant when all aromatic ring carbons have attached methyl groups. Besides, the higher O/C and lower H/C (or the higher OS_c) than predicted in Section 3.2.3 indicates that SOA components from hexamethylbenzene photooxidation are more oxidized per carbon due to oxidation of
- 10 the methyl groups, which is possibly related to the steric hindrance of the six methyl groups. Moreover, there is a slightly increasing trend in VFR from 1, 2, 4, 5- tetramethylbenzene to hexamethylbenzene (Section 3.3.2). Further studies (e.g. photooxidation using isotope labeled methyl group hexamethylbenzene) are required to probe the unique SOA aspects from hexamethylbenzene photooxidation.

15 5. Atmospheric Implication

20

The impact of the number of methyl group substituents on SOA formation has been comprehensively studied in this work by integrating SOA yield with SOA chemical composition and SOA physical properties. A generally decreasing trend is found in the SOA mass yield and the carbon number averaged oxidation level with increasing number of methyl groups. SOA physical properties agree with yield and oxidation results. Therefore, this study demonstrates that the addition of methyl group substitutes to monocyclic aromatic precursors decreases the oxidation of aromatic hydrocarbon to less volatile compounds. Offsetting the amount of CO_2^+ and $C_2H_3O^+$ suggests a methyl group dilution effect on SOA formation from monocyclic aromatic

25 hydrocarbons. The proposed methyl group dilution effect is then applied successfully to the predict SOA elemental ratio. Overall, this study clearly demonstrates the methyl group impact on SOA formation from monocyclic aromatic hydrocarbons. Benzene and toluene are evaluated as the most important monocyclic aromatic precursors to SOA formation among the six compounds studied due to their high SOA yields and highly oxidized components. Hexamethylbenzene is found to be significantly more oxidized than predicted based on other monocyclic aromatic

- 5 hydrocarbons studied here. This implies uniqueness in the methyl group behavior (no -H on aromatic ring) in hexamethylbenzene. Oligomerization is proposed to be an important pathway for SOA formation from monocyclic aromatic hydrocarbons. It is likely that oligomerization is even more valuable to SOA formation from monocyclic aromatic hydrocarbons under polluted areas (catalyzed effect, Jang, et al., 2002;
- Iinuma, et al., 2004; Noziere, et al., 2008) and ambient humidity (Liggio, et al., 2005a, b; Hastings, et al., 2005).

Acknowledgements

We acknowledge funding support from National Science Foundation (ATM 0901282) and W. M. Keck Foundation. Any opinions, findings, and conclusions expressed in
this material are those of the author(s) and do not necessarily reflect the views of the NSF.

References

20

Aiken, A. C., DeCarlo, P. F. and Jimenez, J. L.: Elemental analysis of organic species with electron ionization high-resolution mass spectrometry, Anal. Chem., 79(21), 8350-8358, doi:10.1021/ac071150w, 2007.

Aiken, A. C., DeCarlo, P. F., Kroll, J. H., Worsnop, D. R., Huffman, J. A., Docherty,
K. S., Ulbrich, I. M., Mohr, C., Kimmel, J. R., Sueper, D., Sun, Y., Zhang, Q.,
Trimborn, A., Northway, M., Ziemann, P. J., Canagaratna, M. R., Onasch, T. B.,
Alfarra, M. R., Prevot, A. S. H., Dommen, J., Duplissy, J., Metzger, A.,

25 Baltensperger, U. and Jimenez, J. H.: O/C and OM/OC ratios of primary, secondary, and ambient organic aerosols with high-resolution time-of-flight aerosol mass spectrometry, Environ. Sci. Technol., 42(12), 4478-4485, doi:10.1021/es703009q, 2008.

Alfarra, M. R., Coe, H., Allan, J. D., Bower, K. N., Boudries, H., Canagaratna, M. R.,
Jimenez, J. L., Jayne, J. T., Garforth, A. A., Li, S-M. and Worsnop, D. R.:

Characterization of urban and rural organic particulate in the lower Fraser valley using two aerodyne aerosol mass spectrometers, Atmos. Environ., 38(34), 5745-5758, doi:10.1016/j.atmosenv.2004.01.054, 2004.

Arey, J., Obermeyer, G., Aschmann, S. M., Chattopadhyay, S., Cusick, R. D. and
Atkinson, R.: Dicarbonyl products of the OH radical-initiated reaction of a series of aromatic hydrocarbons, Environ. Sci. Technol., 43(3), 683-689, doi:10.1021/es8019098, 2008.

Aschmann, S. M., Arey, J. and Atkinson, R.: Rate constants for the reactions of OH radicals with 1, 2, 4, 5-tetramethylbenzene, pentamethylbenzene, 2, 4, 5-

10 trimethylbenzaldehyde, 2, 4, 5-trimethylphenol, and 3-methyl-3-hexene-2, 5-dione and products of OH+ 1, 2, 4, 5-tetramethylbenzene, J. Phys. Chem. A., 117(12), 2556-2568, doi:10.1021/jp400323n, 2013.

Aschmann, S. M., Nishino, N., Arey, J. and Atkinson, R.: Kinetics of the Reactions of OH Radicals with 2-and 3-Methylfuran, 2, 3-and 2, 5-Dimethylfuran, and E-and Z-3-

Hexene-2, 5-dione, and Products of OH+ 2, 5-Dimethylfuran, Environ. Sci. Technol., 45(5), 1859-1865., doi:10.1021/es103207k, 2011.

Atkinson, R.: Rate constants for the atmospheric reactions of alkoxy radicals: An updated estimation method, Atmos. Environ., 41(38), 8468-8485, doi:10.1016/j.atmosenv.2007.07.002, 2007.

20 Atkinson, R. and Arey, J.: Atmospheric degradation of volatile organic compounds, Chem. Rev., 103(12), 4605-4638, doi:10.1021/cr0206420, 2003.

Aumont, B., Valorso, R., Mouchel-Vallon, C., Camredon, M., Lee-Taylor, J. and Madronich, S.: Modeling SOA formation from the oxidation of intermediate volatility n-alkanes, Atmos. Chem. Phys., 12(16), 7577-7589, doi:10.5194/acp-12-7577-2012, 2012.

Bahreini, R., Keywood, M. D., Ng, N. L., Varutbangkul, V., Gao, S., Flagan, R. C., Seinfeld, J. H., Worsnop, D. R. and Jimenez, J. L.: Measurements of secondary organic aerosol from oxidation of cycloalkenes, terpenes, and *m*-xylene using an Aerodyne aerosol mass spectrometer, Environ. Sci. Technol., 39(15), 5674-5688,

30 doi:10.1021/es048061a, 2005.

25

Baltensperger, U., Kalberer, M., Dommen, J., Paulsen, D., Alfarra, M. R., Coe, H.,Fisseha, R., Gascho, A., Gysel, M., Nyeki, S., Sax, M., Steinbacher, M., Prevot, A. S.H., Sjögren, S., Weingartnera, E. and Zenobib, R.: Secondary organic aerosols from

anthropogenic and biogenic precursors, Faraday. Discuss, 130, 265-278, doi:10.1039/b417367h, 2005.

Bierbach, A., Barnes, I., Becker, K. H. and Wiesen, E.: Atmospheric chemistry of unsaturated carbonyls: Butenedial, 4-oxo-2-pentenal, 3-hexene-2, 5-dione, maleic

5 anhydride, 3H-furan-2-one, and 5-methyl-3H-furan-2-one, Environ. Sci. Technol, 28(4), 715-729, doi:10.1021/es00053a028, 1994.

Birdsall, A. W., Andreoni, J. F. and Elrod, M. J.: Investigation of the role of bicyclic peroxy radicals in the oxidation mechanism of toluene, J. Phys. Chem. A., 114(39), 10655-10663, doi:10.1021/jp105467e, 2010.

Birdsall, A. W. and Elrod, M. J.: Comprehensive NO-dependent study of the products of the oxidation of atmospherically relevant aromatic compounds, J. Phys. Chem. A., 115(21), 5397-5407, doi:10.1021/jp2010327, 2011.

Bloss, C., Wagner, V., Jenkin, M. E., Volkamer, R., Bloss, W. J., Lee, J. D., Heard, D. E., Wirtz, K., Martin-Reviejo, M., Rea, G, Wenger, J. C., and Pilling M. J.:

15 Development of a detailed chemical mechanism (MCMv3. 1) for the atmospheric oxidation of aromatic hydrocarbons, Atmos. Chem. Phys., 5(3), 641-664, doi:10.5194/acp-5-641-2005, 2005.

Borrás, E. and Tortajada-Genaro, L. A.: Secondary organic aerosol formation from the photo-oxidation of benzene, Atmos. Environ., 47, 154-163,

20 doi:10.1016/j.atmosenv.2011.11.020, 2012.

Buczynska, A. J., Krata, A., Stranger, M., Godoi, A. F. L., Kontozova-Deutsch, V., Bencs, L., Naveau, I., Roekens, E. and Van Grieken, R.: Atmospheric BTEXconcentrations in an area with intensive street traffic, Atmos. Environ., 43(2), 311-318, doi:10.1016/j.atmosenv.2008.09.071, 2009.

Calvert, J. G., Atkinson, R., Becker, K. H., Kamens, R. M., Seinfeld, J. H.,
 Wallington, T. J. and Yarwood, G.: The mechanisms of atmospheric oxidation of aromatic hydrocarbons, Oxford University Press New York, 2002.

Canagaratna, M. R., Jayne, J. T., Jimenez, J. L., Allan, J. D., Alfarra, M. R., Zhang, Q., Onasch, T. B., Drewnick, F., Coe, H., Middlebrook, A., Delia, A., Williams, L. R.,

30 Trimborn, A. M., Northway, M. J., DeCarlo, P. F., Kolb, C. E., Davidovits, P. and Worsnop D. R.: Chemical and microphysical characterization of ambient aerosols with the aerodyne aerosol mass spectrometer, Mass. Spectrom. Rev., 26(2), 185-222, doi:10.1002/mas.20115, 2007. Canagaratna, M. R., Jimenez, J. L., Kroll, J. H., Chen, Q., Kessler, S. H., Massoli, P., Hildebrandt Ruiz, L., Fortner, E., Williams, L. R., Wilson, K. R., Surratt, J. D., Donahue, N. M., Jayne, J. T. and Worsnop, D. R.: Elemental ratio measurements of organic compounds using aerosol mass spectrometry: characterization, improved colibration, and implications. Atmos. Cham. Phys., 15(1), 252, 272, doi:10.5104/com

5 calibration, and implications, Atmos. Chem. Phys., 15(1), 253-272, doi:10.5194/acp-15-253-2015, 2015.

Cappa, C. D. and Wilson, K. R.: Multi-generation gas-phase oxidation, equilibrium partitioning, and the formation and evolution of secondary organic aerosol, Atmos. Chem. Phys., 12(20), 9505-9528, doi:10.5194/acp-12-9505-2012, 2012.

10 Carter, W. P. L. and Heo, G.: Development of revised SAPRC aromatics mechanisms, Atmos. Environ., 77, 404-414, doi:10.1016/j.atmosenv.2013.05.021, 2013.

Carter, W. P. L., Cocker III, D. R., Fitz, D. R., Malkina, I.L., Bumiller, K., Sauer, C.G., Pisano, J.T., Bufalino, C., Song, C.: A new environmental chamber for evaluation of gas-phase chemical mechanisms and secondary aerosol formation, Atmos Environ 20(40) 7768 7788 doi:10.1016/j.atmosony.2005.08.040.2005

15 Atmos. Environ., 39(40), 7768-7788, doi:10.1016/j.atmosenv.2005.08.040, 2005.

Carter, W. P. L. and Heo, G.: Development of Revised SAPRC Aromatics Mechanisms, California Air Resources Board, Sacramento, CA, USA, 2012.Chan, A. W. H., Kroll, J. H., Ng, N. L. and Seinfeld, J. H.: Kinetic modeling of secondary organic aerosol formation: effects of particle-and gas-phase reactions of semivolatile

20 products, Atmos. Chem. Phys., 7(15), 4135-4147, doi:10.5194/acp-7-4135-2007,2007.

Chen, Y., Wang, W. and Zhu, L.: Wavelength-dependent photolysis of methylglyoxal in the 290-440 nm region, J. Phys. Chem. A., 104(47), 11126-11131, doi:10.1021/jp002262t, 2000.

25 Chhabra, P. S., Ng, N. L., Canagaratna, M. R., Corrigan, A. L., Russell, L. M., Worsnop, D. R., Flagan, R. C. and Seinfeld, J. H.: Elemental composition and oxidation of chamber organic aerosol, Atmos. Chem. Phys., 11(17), 8827-8845, doi:10.5194/acp-11-8827-2011, 2011.

Cocker III, D. R., Flagan, R. C. and Seinfeld, J. H.: State-of-the-art chamber facility
for studying atmospheric aerosol chemistry, Environ. Sci. Technol., 35(12), 2594-2601, doi:10.1021/es0019169, 2001.

Cocker III, D. R., Mader, B. T., Kalberer, M., Flagan, R.C. and Seinfeld, J. H.: The effect of water on gas-particle partitioning of secondary organic aerosol: II. *m*-xylene

and 1, 3, 5-trimethylbenzene photooxidation systems, Atmos. Environ., 35(35), 6073-6085, doi:10.1016/S1352-2310(01)00405-8, 2001.

Cross, E. S., Slowik, J. G., Davidovits, P., Allan, J. D., Worsnop, D. R., Jayne, J. T., Lewis, D. K., Canagaratna, M. and Onasch, T. B.: Laboratory and ambient particle

 density determinations using light scattering in conjunction with aerosol mass spectrometry, Aerosol Sci. Tech., 41(4), 343-359., doi:10.1080/02786820701199736, 2007.

Darouich, T. AI., Behar, F. and Largeau., C.: Thermal cracking of the light aromatic fraction of Safaniya crude oil–experimental study and compositional modelling of

10 molecular classes. Org. Geochem., 37(9), 1130-1154, doi:.10.1016/j.orggeochem.2006.04.003, 2006

DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R. and Jimenez, J. L.: Field-deployable, high-resolution, time-of-flight aerosol mass spectrometer.

15 Anal. Chem., 78(24), 8281-8289, doi:10.1021/ac061249n, 2006.

DeCarlo, P. F., Slowik, J. G., Worsnop, D. R., Davidovits, P. and Jimenez, J. L.: Particle morphology and density characterization by combined mobility and aerodynamic diameter measurements. Part 1: Theory, Aerosol Sci. Tech., 38(12), 1185-1205, doi:10.1080/027868290903907, 2004.

Diehl, J. W. and Sanzo, F. P. Di.: Determination of aromatic hydrocarbons in gasolines by flow modulated comprehensive two-dimensional gas chromatography, J. Chromatogr. A., 1080(2): 157-165, doi:10.1016/j.chroma.2004.11.054, 2005.

Dinar, E., Mentel, T. and Rudich, Y.: The density of humic acids and humic like substances (HULIS) from fresh and aged wood burning and pollution aerosol

25 particles, Atmos. Chem. Phys., 6(12), 5213-5224, doi:10.5194/acp-6-5213-2006,
 2006.

Duplissy, J., DeCarlo, P. F., Dommen, J., Alfarra, M. R., Metzger, A., Barmpadimos, I., Prevot, A. S., Weingartner, E., Tritscher, T. and Gysel, M.: Relating hygroscopicity and composition of organic aerosol particulate matter, Atmos. Chem. Phys., 11(3),

Edney, E., Driscoll, D., Weathers, W., Kleindienst, T., Conver, T., McIver, C. and Li, W.: Formation of polyketones in irradiated toluene/propylene/NO_x/air mixtures, Aerosol Science & Technology, 35(6), 998-1008, doi:10.1080/027868201753306769, 2001.

^{30 1155-1165,} doi:10.5194/acp-11-1155-2011, 2011.

Forstner, H. J. L., Flagan, R. C. and Seinfeld, J. H.: Secondary organic aerosol from the photooxidation of aromatic hydrocarbons: Molecular composition, Environ. Sci. Technol., 31(5), 1345-1358, doi:10.1021/es9605376, 1997.

Fraser, M. P., Cass, G. R., Simoneit, B. R. and Rasmussen, R.: Air quality model
evaluation data for organics. 5. C₆-C₂₂ nonpolar and semipolar aromatic compounds, Environ. Sci. Technol., 32(12), 1760-1770, doi:10.1021/es970349v, 1998.

Glasson, W. A. and Tuesday, C. S.: Hydrocarbon reactivities in the atmospheric photooxidation of nitric oxide, Environ. Sci. Technol., 4(11), 916-924., doi:10.1021/es60046a002, 1970.

- Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, Th. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R. and Wildt, J.: The
- 15 formation, properties and impact of secondary organic aerosol: current and emerging issues, Atmos. Chem. Phys., 9(14), 5155-5236, doi:10.5194/acp-9-5155-2009, 2009.

Hamilton, J. F., Webb, P. J., Lewis, A. C. and Reviejo, M. M.: Quantifying small molecules in secondary organic aerosol formed during the photo-oxidation of toluene with hydroxyl radicals, Atmos. Environ., 39(38), 7263-7275,

20 doi:10.1016/j.atmosenv.2005.09.006, 2005.

Hastings, W. P., Koehler, C. A., Bailey, E. L. and De Haan, D. O.: Secondary organic aerosol formation by glyoxal hydration and oligomer formation: Humidity effects and equilibrium shifts during analysis, Environ. Sci. Technol., 39(22), 8728-8735, doi:10.1021/es0504461, 2005.

Heald, C. L., Kroll, J. H., Jimenez, J. L., Docherty, K. S., DeCarlo, P. F., Aiken, A. C., Chen, Q., Martin, S. T., Farmer, D. K. and Artaxo, P.: A simplified description of the evolution of organic aerosol composition in the atmosphere, Geophys. Res. Lett, 37(8), L08803, doi:10.1029/2010GL042737, 2010.

Henze, D. K., Seinfeld, J. H., Ng, N. L., Kroll, J. H., Fu, T-M., Jacob, D. J., Heald, C.

 L.: Global modeling of secondary organic aerosol formation from aromatic hydrocarbons: high-vs. low-yield pathways, Atmos. Chem. Phys., 8(9), 2405-2421, doi:10.5194/acp-8-2405-2008, 2008.

Hildebrandt Ruiz, L., Paciga, A., Cerully, K., Nenes, A., Donahue, N. M. and Pandis, S. N.: Aging of secondary organic aerosol from small aromatic VOCs: changes in

chemical composition, mass yield, volatility and hygroscopicity, Atmos. Chem. Phys. Disc., 14(22), 31441-31481, doi:10.5194/acpd-14-31441-2014, 2014.

Holzinger, R., Kleiss, B., Donoso, L. and Sanhueza, E.: Aromatic hydrocarbons at urban, sub-urban, rural (8° 52′ N; 67° 19′ W) and remote sites in Venezuela, Atmos. Environ., 35(29), 4917-4927, doi:10.1016/S1352-2310(01)00286-2, 2001.

5

20

Hu, D. and Kamens, R. M.: Evaluation of the UNC toluene-SOA mechanism with respect to other chamber studies and key model parameters, Atmos. Environ., 41(31), 6465-6477, doi:10.1016/j.atmosenv.2007.04.026, 2007.

Hu, D., Tolocka, M., Li, Q. and Kamens, R. M.: A kinetic mechanism for predicting
secondary organic aerosol formation from toluene oxidation in the presence of NOx and natural sunlight, Atmos. Environ., 41(31), 6478-6496, doi:10.1016/j.atmosenv.2007.04.025, 2007.

Hu, L., Millet, D. B., Baasandorj, M., Griffis, T. J., Travis, K. R., Tessum, C. W., Marshall, J. D., Reinhart, W. F., Mikoviny, T., Müller, M., Wisthaler, A., Graus, M.,

Warneke, C., and de Gouw, J.:Emissions of C₆–C₈ aromatic compounds in the United States: Constraints from tall tower and aircraft measurements, J. Geophys. Res.-Atmos., 120(2), 826-842, doi:10.1002/2014JD022627, 2015.

Iinuma, Y., Böge, O., Gnauk, T. and Herrmann, H.: Aerosol-chamber study of the α -pinene/O₃ reaction: influence of particle acidity on aerosol yields and products, Atmos. Environ., 38(5), 761-773, doi:10.1016/j.atmosenv.2003.10.015, 2004.

Jang, M., Czoschke, N. M., Lee, S., and Kamens, R. M.: Heterogeneous atmospheric aerosol production by acid-catalyzed particle-phase reactions, Science, 298(5594), 814-817, doi:10.1126/science.1075798, 2002.

Jang, M. and Kamens, R.M.: Characterization of secondary aerosol from the
photooxidation of toluene in the presence of NO_x and 1-propene, Environ. Sci.
Technol., 35(18), 3626-3639, doi:10.1021/es010676+, 2001.

Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q.,Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty,K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D.,

Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A.,
Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J.
M., Collins, D. R., Cubison, M. J., Dunlea1, E., J., Huffman, J. A., Onasch, T. B.,
Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F.,
Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R.,

Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M.,
Dzepina, K., Kimmel, J. R., Sueper, D., J. Jayne, T., Herndon, S. C., Trimborn, A.
M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb C. E., Baltensperger, U.
and Worsnop D. R.: Evolution of organic aerosols in the atmosphere, Science,
326(5959), 1525-1529, doi:10.1126/science.1180353, 2009.

Johnson, D., Jenkin, M. E., Wirtz, K., Martin-Reviejo, M.: Simulating the formation of secondary organic aerosol from the photooxidation of aromatic hydrocarbons,

Environ. Chem., 2(1), 35-48, doi:10.1071/EN04079, 2005.

5

15

Johnson, D., Jenkin, M. E., Wirtz, K., Martin-Reviejo, M.: Simulating the formation
of secondary organic aerosol from the photooxidation of toluene, Environ. Chem.,
1(3), 150-165, doi:10.1071/EN04069, 2004.

Kalberer, M., Paulsen, D., Sax, M., Steinbacher, M., Dommen, J., Prevot, A. S. H., Fisseha, R., Weingartner, E., Frankevich, V. and Zenobi, R.: Identification of polymers as major components of atmospheric organic aerosols, Science, 303(5664), 1659-1662, doi:10.1126/science.1092185, 2004.

Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J, Facchini, M.C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J.P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E.L., Tsigaridis, K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic aerosol and

20 global climate modelling: a review, Atmos. Chem. Phys., 5(4), 1053-1123, doi:10.5194/acp-5-1053-2005, 2005.

Katrib, Y., Martin, S. T., Rudich, Y., Davidovits, P., Jayne, J. T. and Worsnop, D. R.: Density changes of aerosol particles as a result of chemical reaction, Atmos. Chem. Phys., 5(1), 275-291, doi:10.5194/acp-5-275-2005, 2005.

25 Kleindienst, T. E., Smith, D. F., Li, W., Edney, E. O., Driscoll, D. J., Speer, R. E. and Weathers, W. S.: Secondary organic aerosol formation from the oxidation of aromatic hydrocarbons in the presence of dry submicron ammonium sulfate aerosol, Atmos. Environ., 33(22), 3669-3681, doi:10.1016/S1352-2310(99)00121-1, 1999.

Kroll, J. H., Donahue, N. M., Jimenez, J. L., Kessler, S. H., Canagaratna, M. R.,

Wilson, K. R., Altieri, K. E., Mazzoleni, L. R., Wozniak, A. S., Bluhm, H., Mysak, E.
 R., Smith, J. D., Kolb, C. E. and Worsnop D. R.: Carbon oxidation state as a metric for describing the chemistry of atmospheric organic aerosol, Nature Chemistry, 3(2), 133-139, doi:10.1038/nchem.948, 2011.

Kroll, J. H. and Seinfeld, J. H.: Chemistry of secondary organic aerosol: Formation and evolution of low-volatility organics in the atmosphere, Atmos. Environ., 42(16), 3593-3624, http://dx.doi.org/10.1016/j.atmosenv.2008.01.003, 2008.

Kroll, J. H., Smith, J. D., Che, D. L., Kessler, S. H., Worsnop, D. R. and Wilson,

K.R.: Measurement of fragmentation and functionalization pathways in the heterogeneous oxidation of oxidized organic aerosol, Phys. Chem. Chem. Phys., 11(36), 8005-8014, doi:10.1039/b905289e, 2009.

Kuwata, M., Zorn, S. R. and Martin, S. T.: Using elemental ratios to predict the density of organic material composed of carbon, hydrogen, and oxygen, Environ. Sci. Technol., 46(2), 787-794, doi:10.1021/es202525g, 2011.

Lambe, A. T., Chhabra, P. S., Onasch, T. B., Brune, W. H., Hunter, J. F., Kroll, J. H., Cummings, M. J., Brogan, J. F., Parmar, Y., Worsnop, D. R., Kolb, C. E., and Davidovits, P.: Effect of oxidant concentration, exposure time, and seed particles on secondary organic aerosol chemical composition and yield, Atmos. Chem. Phys., 15(6) 20(2) 2075. Li 10 5104/

15 15(6), 3063-3075, doi:10.5194/acp-15-3063-2015, 2015.

10

Liggio, J., Li, S.-M., McLaren, R.: Heterogeneous reactions of glyoxal on particulate matter: Identification of acetals and sulfate esters, Environ. Sci. Technol., 39(6), 1532-1541, doi:10.1021/es048375y, 2015a.

Liggio, J., Li, S.-M., McLaren, R.: Reactive uptake of glyoxal by particulate matter. J.
Geophys. Res.-Atmos., 110, D10304, doi:10.1029/2004JD005113, 2015b.

Li. L, Tang P., Nakao. S., and Cocker III, D. R..: Impact of Molecular Structure on Secondary Organic Aerosol Formation from Aromatic Hydrocarbon Photooxidation under Low NOx Conditions, Atmos. Chem. Phys. Discuss, 2015.

Lim, Y. B. and Ziemann, P. J.: Effects of molecular structure on aerosol yields from
OH radical-initiated reactions of linear, branched, and cyclic alkanes in the presence of NO_x, Environ. Sci. Technol., 43(7), 2328-2334, doi:10.1021/es803389s, 2009a.

Lockhart, J., Blitz, M., Heard, D., Seakins, P. and Shannon, R.: Kinetic study of the OH+ glyoxal reaction: experimental evidence and quantification of direct OH recycling, J. Phys. Chem. A., 117(43), 11027-11037, doi:10.1021/jp4076806, 2013.

Loza, C. L., Chhabra, P. S., Yee, L. D., Craven, J. S., Flagan, R. C. and Seinfeld, J.
H.: Chemical aging of *m*-xylene secondary organic aerosol: laboratory chamber study, Atmos. Chem. Phys., 12(1), 151-167, doi:10.5194/acp-12-151-2012, 2012. Malloy, Q. G., Nakao, S., Qi, L., Austin, R., Stothers, C., Hagino, H. and Cocker III,
D. R.: Real-Time Aerosol Density Determination Utilizing a Modified Scanning
Mobility Particle Sizer—Aerosol Particle Mass Analyzer System, Aerosol Sci. Tech.,
43(7), 673-678.doi:10.1080/02786820902832960, 2009.

5 Martín-Reviejo, M. and Wirtz, K.: Is benzene a precursor for secondary organic aerosol? Environ. Sci. Technol., 39(4), 1045-1054, doi:10.1021/es049802a, 2005.

Matsunaga, A., Docherty, K. S., Lim, Y. B. and Ziemann, P. J.: Composition and yields of secondary organic aerosol formed from OH radical-initiated reactions of linear alkenes in the presence of NO_x: Modeling and measurements, Atmos. Environ.,

10 43(6), 1349-1357, doi:10.1016/j.atmosenv.2008.12.004, 2009.

15

McLafferty, F.W. and Tureček, F.: Interpretation of mass spectra. Univ Science Books, Sausalito, CA, USA, 1993.

Nakao, S., Clark, C., Tang, P., Sato, K. and Cocker III, D. R.: Secondary organic aerosol formation from phenolic compounds in the absence of NO_x, Atmos. Chem. Phys, 11, 10649-10660, doi:10.5194/acp-11-10649-2011, 2011.

Nakao, S., Liu, Y., Tang, P., Chen, C.-L., Zhang, J. and Cocker III, D. R.: Chamber studies of SOA formation from aromatic hydrocarbons: observation of limited glyoxal uptake, Atmos. Chem. Phys., 12(9), 3927-3937, doi:10.5194/acp-12-3927-2012, 2012.

20 Nakao, S., Tang, P., Tang, X., Clark, C. H., Qi, L., Seo, E., Asa-Awuku, A. and Cocker III, D. R.: Density and elemental ratios of secondary organic aerosol: Application of a density prediction method, Atmos. Environ., 68, 273-277,doi:10.1016/j.atmosenv.2012.11.006, 2013.

Ng, N. L., Canagaratna, M. R., Jimenez, J. L., Chhabra, P. S., Seinfeld, J. H. and
Worsnop, D. R.: Changes in organic aerosol composition with aging inferred from aerosol mass spectra, Atmos. Chem. Phys, 11(13), 6465-6474, doi:10.5194/acp-11-6465-2011, 2011.

Ng, N. L., Canagaratna, M. R., Zhang, Q., Jimenez, J. L., Tian, J., Ulbrich, I. M., Kroll, J. H., Docherty, K. S., Chhabra, P. S., Bahreini, R., Murphy, S. M., Seinfeld, J.

30 H., Hildebrandt, L., Donahue, N. M., DeCarlo, P. F., Lanz, V. A., Prévôt, A. S. H., Dinar E., Rudich Y. and Worsnop D. R.: Organic aerosol components observed in Northern Hemispheric datasets from Aerosol Mass Spectrometry, Atmos. Chem. Phys., 10(10), 4625-4641, doi:10.5194/acp-10-4625-2010, 2010. Ng, N. L., Kroll, J. H., Chan, A. W. H., Chhabra, P. S., Flagan, R. C. and Seinfeld, J. H.: Secondary organic aerosol formation from *m*-xylene, toluene, and benzene, Atmos. Chem. Phys., 7(14), 3909-3922, doi: 10.5194/acp-7-3909-2007, 2007.

Noziere, B., Dziedzic, P. and Córdova, A.: Products and kinetics of the liquid-phase
reaction of glyoxal catalyzed by ammonium ions (NH4⁺), J. Phys. Chem. A., 113(1), 231-237, doi:10.1021/jp8078293, 2008.

Odum, J. R., Hoffmann, T., Bowman, F., Collins, D., Flagan, R. C. and Seinfeld, J. H.: Gas/particle partitioning and secondary organic aerosol yields, Environ. Sci. Technol., 30(8), 2580-2585, doi:10.1021/es950943+, 1996.

10 Odum, J. R., Jungkamp, T., Griffin, R., Flagan, R. C. and Seinfeld, J. H.: The atmospheric aerosol-forming potential of whole gasoline vapor, Science, 276(5309), 96-99, doi:10.1126/science.276.5309.96, 1997a.

Odum, J. R., Jungkamp, T., Griffin, R. J., Forstner, H., Flagan, R. C. and Seinfeld, J.H.: Aromatics, reformulated gasoline, and atmospheric organic aerosol formation, Environ. Sci. Technol., 31(7), 1890-1897, doi:10.1021/es9605351, 1997b.

Pang, Y., Turpin, B. and Gundel, L.: On the importance of organic oxygen for understanding organic aerosol particles, Aerosol Sci. Tech., 40(2), 128-133, doi:10.1080/02786820500423790, 2006.

Pankow, J. F. and Asher, W. E.: SIMPOL. 1: a simple group contribution method for
predicting vapor pressures and enthalpies of vaporization of multifunctional organic compounds, Atmos. Chem. Phys., 8(10), 2773-2796, doi:10.5194/acp-8-2773-2008, 2008.

Pfaffenberger, L., Barmet, P., Slowik, J. G., Praplan, A. P., Dommen, J., Prévôt, A. S. H. and Baltensperger, U. : The link between organic aerosol mass loading and degree

of oxygenation: an α-pinene photooxidation study, Atmos. Chem. Phys., 13(13),
 6493-6506, doi:10.5194/acp-13-6493-2013, 2013.

Hassoun, S., Pilling, M. J. and Bartle, K. D.: A catalogue of urban hydrocarbons for the city of Leeds: atmospheric monitoring of volatile organic compounds by thermal desorption-gas chromatography, J. Environ. Monitor., 1(5), 453-458,

30 doi:10.1039/a904879k, 1999.

15

Plum, C. N., Sanhueza, E., Atkinson, R., Carter, W. P. and Pitts, J. N.: Hydroxyl radical rate constants and photolysis rates of alpha-dicarbonyls, Environ. Sci. Technol., 17(8), 479-484, doi:10.1021/es00114a008, 1983.

Qi, L., Nakao, S., Malloy, Q., Warren, B. and Cocker, D. R.: Can secondary organic aerosol formed in an atmospheric simulation chamber continuously age? Atmos. Environ., 44(25), 2990-2996, doi:10.1016/j.atmosenv.2010.05.020, 2010a.

Qi, L., Nakao, S., Tang, P. and Cocker III, D. R.: Temperature effect on physical and 5 chemical properties of secondary organic aerosol from *m*-xylene photooxidation, Atmos. Chem. Phys., 10(8), 3847-3854, doi:10.5194/acp-10-3847-2010, 2010b.

Rader, D. J. and McMurry, P. H.: Application of the tandem differential mobility analyzer to studies of droplet growth or evaporation, J. Aerosol. Sci., 17(5), 771-787, doi:10.1016/0021-8502(86)90031-5, 1986.

- 10 Salo, K., Hallquist, M., Jonsson, A.M., Saathoff, H., Naumann, K.-H., Spindler, C., Tillmann, R., Fuchs, H., Bohn, B., Rubach, F., Mentel, T. F., Müller, L., Reinnig, M., Hoffmann, T., and Donahue, N. M.: Volatility of secondary organic aerosol during OH radical induced ageing, Atmos. Chem. Phys., 11(21), 11055-11067, doi:10.5194/acp-11-11055-2011, 2011.
- 15 Salter, R. J., Blitz, M. A., Heard, D. E., Kovács, T., Pilling, M. J., Rickard, A. R. and Seakins, P.W.: Quantum yields for the photolysis of glyoxal below 350 nm and parameterisations for its photolysis rate in the troposphere, Phys. Chem. Chem. Phys., 15(14), 4984-4994, doi:10.1039/c3cp43597k, 2013.
- Sato, K., Hatakeyama, S. and Imamura, T.: Secondary organic aerosol formation 20 during the photooxidation of toluene: NO_x dependence of chemical composition, J. Phys. Chem. A., 111(39), 9796-9808, doi:10.1021/jp071419f, 2007.

Sato, K., Takami, A., Isozaki, T., Hikida, T., Shimono, A. and Imamura, T.: 2010. Mass spectrometric study of secondary organic aerosol formed from the photooxidation of aromatic hydrocarbons, Atmos. Environ., 44(8), 1080-1087, doi:10.1016/j.atmosenv.2009.12.013, 2010.

Sato, K., Takami, A., Kato, Y., Seta, T., Fujitani, Y., Hikida, T., Shimono, A. and Imamura, T. 2012. AMS and LC/MS analyses of SOA from the photooxidation of benzene and 1, 3, 5-trimethylbenzene in the presence of NO_x: effects of chemical structure on SOA aging, Atmos. Chem. Phys, 12, 4667-4682, doi:10.5194/acp-12-4667-2012, 2012.

30

25

Shilling, J. E., Chen, Q., King, S. M., Rosenoern, T., Kroll, J. H., Worsnop, D. R., DeCarlo, P. F., Aiken, A. C., Sueper, D., Jimenez, J. L. and Martin, S. T.: Loadingdependent elemental composition of α -pinene SOA particles, Atmos. Chem. Phys., 9(3), 771-782, doi:10.5194/acp-9-771-2009, 2009.

Singh, H.B., Salas, L., Viezee, W., Sitton, B. and Ferek, R.: 1992. Measurement of volatile organic chemicals at selected sites in California, Atmos. Environ. A-Gen., 26(16), 2929-2946, doi:10.1016/0960-1686(92)90285-S, 1992.

Singh, H. B., Salas, L. J., Cantrell, B. K. and Redmond, R. M.: Distribution of
aromatic hydrocarbons in the ambient air, Atmos. Environ. (1967), 19(11), 19111919, doi:10.1016/0004-6981(85)90017-4, 1985.

Song, C., Na, K. and Cocker III, D. R.: Impact of the hydrocarbon to NO_x ratio on secondary organic aerosol formation, Environ. Sci. Technol., 39(9), 3143-3149, doi:10.1021/es0493244, 2005.

10 Takegawa, N., Miyakawa, T., Kawamura, K. and Kondo, Y.: Contribution of selected dicarboxylic and ω-oxocarboxylic acids in ambient aerosol to the m/z 44 signal of an Aerodyne aerosol mass spectrometer, Aerosol Sci. Tech., 41(4), 418-437, doi:10.1080/02786820701203215, 2007.

Takekawa, H., Minoura, H. and Yamazaki, S.: Temperature dependence of secondary
organic aerosol formation by photo-oxidation of hydrocarbons, Atmos. Environ.,
37(24), 3413-3424, doi:10.1016/S1352-2310(03)00359-5, 2003.

Tkacik, D. S., Presto, A. A., Donahue, N. M. and Robinson, A. L.: Secondary organic aerosol formation from intermediate-volatility organic compounds: cyclic, linear, and branched alkanes, Environ. Sci. Technol., 46(16), 8773-8781, doi:10.1021/es301112c, 2012.

20 201

Tritscher, T., Dommen, J., DeCarlo, P. F., Gysel, M., Barmet, P. B., Praplan, A. P., Weingartner, E., Prévôt, A. S. H., Riipinen, I., Donahue, N. M. and Baltensperger, U.: Volatility and hygroscopicity of aging secondary organic aerosol in a smog chamber, Atmos. Chem. Phys., 11(22), 11477-11496, doi:10.5194/acp-11-11477-2011, 2011.

25 Võ, U.-U.T. and Morris, M.P.: Nonvolatile, semivolatile, or volatile: Redefining volatile for volatile organic compounds. Journal of the Air & Waste Management Association, 64(6), 661-669, doi: 10.1080/10962247.2013.873746, 2014.

Wyche, K. P., Monks, P. S., Ellis, A. M., Cordell, R. L., Parker, A. E., Whyte, C., Metzger, A., Dommen, J., Duplissy, J., Prevot, A. S. H., Baltensperger, U., Rickard,

A. R., and Wulfert, F.: Gas phase precursors to anthropogenic secondary organic aerosol: detailed observations of 1, 3, 5-trimethylbenzene photooxidation, Atmos. Chem. Phys., 9(2), 635-665, doi:10.5194/acp-9-635-2009, 2009.

Xiang, B., Zhu, L. and Tang, Y.: Photolysis of 4-Oxo-2-pentenal in the 190-460 nm Region, J. Phys. Chem. A., 111(37), 9025-9033, doi:10.1021/jp0739972, 2007.

Yu, J., Jeffries, H. E. and Sexton, K. G. 1997. Atmospheric photooxidation of alkylbenzenes—I. Carbonyl product analyses, Atmos. Environ., 31(15), 2261-2280, doi:10.1016/S1352-2310(97)00011-3, 1997.

Yu, L., Smith, J., Laskin, A., Anastasio, C., Laskin, J., Zhang, Q.: Chemical
characterization of SOA formed from aqueous-phase reactions of phenols with the
triplet excited state of carbonyl and hydroxyl radical, Atmos. Chem. Phys., 14(24),
13801-13816, doi:10.5194/acp-14-13801-2014, 2014.

Zhang, Q., Alfarra, M. R., Worsnop, D. R., Allan, J. D., Coe, H., Canagaratna, M. R. and Jimenez, J. L.: Deconvolution and quantification of hydrocarbon-like and

10 oxygenated organic aerosols based on aerosol mass spectrometry, Environ. Sci.
 Technol., 39(13), 4938-4952, doi:10.1021/es0485681, 2005.

Ziemann, P.: Effects of molecular structure on the chemistry of aerosol formation from the OH-radical-initiated oxidation of alkanes and alkenes, International Reviews in Physical Chemistry, 30(2), 161-195, doi:10.1080/0144235X.2010.550728, 2011.

| Precursor | ID | HC/NO ^a | NO ^b | НС ^ь | ΔHC | M ₀ ^c | Yield |
|--------------------|----------------|--------------------|-----------------|-----------------|-----------------|-----------------------------|-------|
| Danzona | 1222 4 | 09.1 | 50.5 | 072 | 208 | 120 | 0.25 |
| Denzene | 1223A 1222D | 90.1 40.2 | 39.3 110 | 972 | 590 452 | 105 | 0.55 |
| | 12230 | 49.5 | 526 | 9/9 | 433 | 105 | 0.25 |
| | 1230A 1226D | 104 | 35.0 154 | 928 | 407 | 24.0 | 0.20 |
| | 12300 | 50.4 62.7 | 134 | 930 125 | 430 | 54.9 45.0 | 0.08 |
| | 1237A 1227D | 120 | 41.0 21.1 | 455 | 200 | 43.9 | 0.17 |
| | 123/D 1619A | 129 | 21.1 | 433 | 233 | 31.4 46.2 | 0.20 |
| Taluana | 1018A | 20.0 | 33.4 | 70.7 | 206 | 40.5 | 0.15 |
| Toluene | 1101A | 29.0 50.0 | 19.2 | 19.1 70.0 | 200 | 30.1 25.1 | 0.15 |
| | 1101B | 58.8 12.2 | 9.40 | /8.8 75 7 | 1/0 | 25.1 | 0.14 |
| | 1102A | 12.2 | 43.3 | /5./ | 223 | 21.8 | 0.10 |
| | 1102B | 1/.5 | 33.0 20.1 | 82.5 | 238 | 22.2 | 0.09 |
| | 1100A | 13.2 | 20.1 | 38.0 | 120 | 9.80 | 0.08 |
| | 1100B | 24.4 | 10.0 | 30.9 | | 12.4 | 0.11 |
| | 1408A | 20.1 | 04.1 | 239 | $\frac{60}{71}$ | 130 | 0.20 |
| | 1408B | 20.5 | 63.0 | 238 | 0/1 | 12/ | 0.19 |
| <i>m</i> -Xylene | 1191A | 12.0 | 52.2 | 82.1 | 298 | 15.2 | 0.05 |
| | 1191B | 14.0 | 45.7 | 83.6 | 340 | 14.0 | 0.04 |
| | 1193A | 15.5 | 36.8 | /1.1 | 239 | 13.0 | 0.06 |
| | 1193B | 15.2 | 36.5 | 69.5 92.1 | 236 | 11.2 | 0.05 |
| | 1191A | 12.0 | 52.2 | 82.1 | 298 | 15.2 | 0.05 |
| | 1191B | 14.0 | 45.7 | 83.6 | 340 | 14.0 | 0.04 |
| | 1516A | 27.8 | 26.7 | 92.9 | 357 | 48.7 | 0.14 |
| | 1950A | 14.1 | 45.5 | 80.0 | 327 | 26.3 | 0.08 |
| 1.0.4 | 1950B | 14.0 | 45.9 | 83.6 | 345 | 28.7 | 0.08 |
| 1,2,4- | 111/A | 69.8 | 10.3 | 80.0 | 335 | 16.8 | 0.05 |
| Trimethylbenzene | 111/B | 34.8 | 20.7 | 80.0 | 368 | 18.2 | 0.05 |
| | 1119A | 14.1 | 49.8 | 78.0 | 385 | 19.6 | 0.05 |
| | 1119B | 17.1 | 41.6 | 79.0 | 390 | 25.5 | 0.07 |
| | 1123A | 71.0 | 10.1 | 80.0 | 300 | 11.2 | 0.04 |
| | 1123B | 32.6 | 22.1 | 80.0 | 345 | 15.4 | 0.05 |
| | 1126A | 69.3 | 10.1 | 77.5 | 286 | 12.6 | 0.04 |
| | 1126B | 28.1 | 24.3 | 75.9 | 333 | 15.4 | 0.05 |
| | 1129B | 24.2 | 15.6 | 42.0 | 201 | 5.60 | 0.03 |
| 1,2,4,5- | 1531A | 72.0 | 25.0 | 180 | 752 | 17.9 | 0.02 |
| Tetramethylbenzene | 1603A | 109 | 11.2 | 122 | 469 | 3.12 | 0.01 |
| | 1603B | 110 | 11.1 | 123 | 464 | 2.54 | 0.01 |
| | 2085A | 60.6 | 33.4 | 202 | 862 | 29.2 | 0.03 |
| | 2085B | 136 | 12.9 | 175 | 502 | 8.20 | 0.02 |
| Pentamethylbenzene | 1521A | 68.8 | 23.5 | 147 | 893 | 32.7 | 0.04 |
| | 1627A | 77.9 | 20.0 | 142 | 769 | 20.6 | 0.03 |
| | 1627B | 26.6 | 50.0 | 121 | 719 | 24.8 | 0.03 |
| Hexamethylbenzene | 1557A | 72.0 | 28.0 | 168 | 999 | 23.4 | 0.02 |
| | 2083A | 78.4 | 11.6 | 76.0 | 442 | 15.2 | 0.03 |
| | 2083B | 41.3 | 22.0 | 76.0 | 483 | 14.0 | 0.03 |

1 Table 1. Experiment conditions*

 $1 \qquad \text{Note: a) Unit of HC/NO are ppbC:ppb; b) Unit of NO and HC are ppb; c) Unit of Δ HC and M_0 are μg·m-3, M_0 is a wall loss and M_0 are μg·m-3, M_0

2 density corrected particle mass concentration; * Only newly added data are listed here and published data are listed in Table S2.

| Yield Curve | α_1 | $K_{om,1}$ (m ³ /µg) | α2 | $K_{om,2}$ (m ³ /µg) |
|------------------------|------------|---------------------------------|-------|---------------------------------|
| Benzene | 0.082 | 0.017 | 0.617 | 0.005 |
| Toluene | 0.185 | 0.080 | 0.074 | 0.005 |
| <i>m</i> -Xylene | 0.148 | 0.047 | 0.079 | 0.005 |
| 1,2,4-Trimethylbenzene | 0.099 | 0.047 | 0.079 | 0.005 |
| C ₁₀₊ | 0.048 | 0.047 | 0.065 | 0.005 |
| | | | | |

3 Table 2. Two product yield curve fitting parameters

4

5



Fig. 1. Aromatic SOA yields as a function of M₀ *Note: Song, et al. (2005) *m*-xylene data are also
 included



2 Fig. 2. f₄₄ and f₄₃ evolution in SOA formed from photooxidation of different

3 monocyclic aromatic hydrocarbons under low NO_x (Benzene 1223A; Toluene 1468A;

4 *m*-Xylene 1950A; 1,2,4-Trimethylbenzene 1119A; 1,2,4,5-Tetramethylbenzene

5 2085A; Pentamethylbenzene 1627A; Hexamethylbenzene 2083A; colored solid circle

6 markers represent the location of average f_{44} and f_{43} value during photooxidation)



10

1



3 Fig. 3. a) H/C and O/C evolution; the inset graph shows the measured values relative 4 to the classic triangle plot (Ng et al., 2010). b) Average H/C and O/C in SOA formed from monocyclic aromatic hydrocarbon photooxidation under low NO_x (Benzene 5 1223A; Toluene 1468A; *m*-Xylene 1191A; 1,2,4-Trimethylbenzene 1119A; 1,2,4,5-6 7 Tetramethylbenzene 2085A; Pentamethylbenzene 1627A; Hexamethylbenzene 8 2083A).

9

1





^{7 (}OSc) (b) in SOA formation from monocyclic aromatic hydrocarbon photooxidation

10 1627A; Hexamethylbenzene 1557A)

⁸ under low NO_x (Benzene 1223A; Toluene 1468A; m-Xylene 1191A; 1, 2, 4-

⁹ Trimethylbenzene 1119A; 1,2,4,5-Tetramethylbenzene 1306A; Pentamethylbenzene



2 Fig. 5. Relationship between (a) SOA density and methyl group number; (b) SOA

3 density and O/C; (c) predicted and measured density from monocyclic aromatic

4 hydrocarbon photooxidation under low NO_x (number mark represents number of





Fig. 6. Relationship between a) SOA volatility and methyl group number; b) SOA
volatility and O/C; c) SOA volatility and oxidation state (OS_c) from monocyclic
aromatic hydrocarbon photooxidation under low NO_x (number mark represents
number of methyl groups on aromatic hydrocarbon ring).



2 Fig. 7. Monocyclic aromatic hydrocarbon oxidation pathways related to SOA

3 formation (methyl substitute on aromatic ring not shown)

1

| 4 | Aromatic Hydrocarbon + OH | \rightarrow S ¹ ^g | \rightarrow S ₂ ^g | \rightarrow S ₃ ^g |
|---|---------------------------|---|---|---|
| 5 | | ↓ ↑ | | ↓ ↑ |
| 6 | | $S_1{}^p$ | | $\mathbf{S_3}^p$ |

7 Fig. 8. Kinetic scheme for SOA formation from monocyclic aromatic hydrocarbon