

## Responses to Reviewers' Comments:

We thank the reviewers for the consideration and the constructive comments. The manuscript is revised based on the suggestions made and detailed responses to the reviewers are addressed as follows.

### Referee #1

The authors present a detailed study of the aqueous-phase oxidation of methyl ethyl ketone (MEK). Their most significant finding is that this species may be a precursor for methylglyoxal in the aqueous phase. The experiments were carefully performed and a detailed modeling analysis was used to determine the reaction mechanism and rate constants. The paper is suitable for publication in ACP after a few minor points are addressed.

1) My main concern about this paper is that I would like to see a discussion of the relative importance of this source of methylglyoxal as compared to uptake from the gas phase. The Henry's Law constant of MEK is very low, 2-3 orders of magnitude lower than that of methylglyoxal itself. Considering mass transfer limitations, how significant of a source of methylglyoxal, and by extension, aqueous SOA, is this pathway?

### Author's comment

In Table 1 the Henry constants of MEK and methylglyoxal as well as measured concentrations in the gas, particle and aqueous phase are summarised.

Table 1: Henry constant and concentrations of MEK and methylglyoxal in gas-, particle- and aqueous phase.

	Henry constant [M atm <sup>-1</sup> ]	Concentration gas phase [ppb]	Concentration particle phase [µg m <sup>-3</sup> ]	Concentration aqueous phase [µmol L <sup>-1</sup> ]
MEK	7.7 – 50 <sup>abcde</sup> fg	0.02 – 15 <sup>bijklmn</sup>	0 – 0.1 <sup>i</sup>	0 – 0.65(CW) <sup>uu</sup> ; < 0.0005(OBW) and 0.00228 (SML) <sup>v</sup>
Methyl- glyoxal	3.7 × 10 <sup>3</sup> <sup>h</sup> 3.2 × 10 <sup>4</sup> <sup>d</sup>	≤0.05 – 0.09 <sup>o</sup> 0.03 – 270.6 <sup>pqrs</sup>	0 – 0.075 <sup>p</sup>	0.3 – 18(CW) <sup>ou</sup>

CW cloud water, OBW ocean bulk water, SML surface microlayer

*a* Buttery et al., 1969; *b* Snider and Dawson, 1985; *c* Ashworth et al., 1988; *d* Zhou and Mopper, 1990; *e* Morillon et al., 1999; *f* Karl et al., 2002; *g* Schütze and Herrmann 2004; *h* Betterton and Hoffmann, 1988; *i* Grosjean et al., 1982, 1983, 2002; *j* Riemer et al., 1998; *k* Singh et al., 2004; *l* Goldan et al., 1995; *m* Müller et al., 2005; *n* Feng et al., 2004; *o* Munger et al., 1995; *p* Kawamura et al., 2013; *q* Oriiz et al., 2006; *r* Matsunaga et al., 2004; *s* Ieda et al., 2006; *t* Grosjean and Wright, 1983; *u* van Pinxteren et al., 2005; *v* Zhou and Mopper, 1997; *w* Collett et al., 1990

As it can be seen, MEK has a lower Henry constant with 50 M atm<sup>-1</sup> as the maximum available value (Schütze and Herrmann, 2004) compared to methylglyoxal ( $K_H = 3.7 \times 10^3$  M atm<sup>-1</sup>,

Betterton and Hoffmann, 1988) leading to the conclusion that the transfer from the gas in the aqueous phase is, according to the Henry constant, more important for methylglyoxal than for MEK. Noticeably, van Pinxteren et al. (2005) found higher concentrations of MEK in cloud water as it was expected according to the Henry constant. Thus, phase transfer could be more important than it is known at this moment. Besides its transfer from the gas phase, the formation of MEK in the aqueous phase of clouds and hygroscopic particles should be considered as an additional source for aqueous MEK as well. The so-formed aqueous MEK will react to form methylglyoxal with a molar yield of 9.5%.

Methylglyoxal present in the aqueous phase originates from i) uptake from the gas phase and ii) aqueous-phase reactions. The uptake of methylglyoxal from the gas phase in the aqueous phase is not fully understood. Kroll et al. (2005) investigated the uptake of methylglyoxal on inorganic seed particles under various relative humidities. It was found that the uptake was not relevant for methylglyoxal even under high relative humidity. Contrary, Zhao et al. (2006) measured an uptake coefficient of  $\gamma = 7.6 \times 10^{-3}$  on acidic solution. They found an irreversible uptake, which decreases with increasing acidity. Fu et al. (2008, 2009) determined an uptake coefficient on aqueous particles and cloud droplets  $\gamma = 2.9 \times 10^{-3}$ , which is in good agreement with the uptake coefficient measured by Zhao et al. (2006). Lin et al. (2014) modelled an uptake coefficient for several case studies with different multiphase process mechanisms. The uptake coefficient in deliquescent particles was determined to be  $\gamma = 1.47 - 2.92 \times 10^{-5}$ . Overall,  $\gamma$  was two orders of magnitude lower compared to the values determined by Zhao et al. (2006) and Fu et al. (2008, 2009) highlighting the discrepancies between the different literature studies.

As the uptake as well as the formation mechanisms of methylglyoxal are not fully understood, it cannot be excluded that the oxidation of MEK represents an important reaction channel for methylglyoxal formation.

Based on the Referee's comment the following paragraph was added to the manuscript:

Page 31907 Line 1 "*4. Atmospheric relevance*

*The sources of methylglyoxal in the aqueous phase are thus far not fully elucidated. Methylglyoxal can originate in the atmospheric aqueous phase through i) uptake from the gas phase, and/or ii) formation in the aqueous phase. The importance of the uptake from the gas into the aqueous phase is discussed in the literature but large discrepancies can be found. Kroll et al. (2005) investigated the uptake of methylglyoxal on inorganic seed particles under varying relative humidity. It was found that the uptake was not relevant for methylglyoxal even under*

high relative humidity. Contrary, Zhao et al. (2006) measured an uptake coefficient of  $\gamma = 7.6 \times 10^{-3}$  on acidic solution. They found an irreversible uptake, which decreases with increasing acidity. Fu et al. (2008, 2009) determined an uptake coefficient on aqueous particles and cloud droplets  $\gamma = 2.9 \times 10^{-3}$ , which is in good agreement with the uptake coefficient measured by Zhao et al. (2006). Lin et al. (2014) modelled an uptake coefficient for several case studies with different multiphase process mechanisms. The uptake coefficient in deliquescent particles was determined to be  $\gamma = 1.47 - 2.92 \times 10^{-5}$ . Overall,  $\gamma$  was two orders of magnitude lower compared to the values determined by Zhao et al. (2006) and Fu et al. (2008, 2009) showing the discrepancies between the different studies available in literature. The in-situ formation of methylglyoxal in the aqueous phase could be an important source as well (Blando and Turpin, 2000; Sempere and Kawamura 1994). Within the present study, MEK was found as a new precursor compound for methylglyoxal in the aqueous phase yielding methylglyoxal with a molar yield of 9.5%. Although the Henry constant of MEK (up to  $K_H = 50 \text{ M atm}^{-1}$ , Schütze and Herrmann, 2004) is lower compared to methylglyoxal ( $K_H = 3.7 \times 10^3 \text{ M atm}^{-1}$ , Betterton and Hoffmann, 1988), van Pinxteren et al. (2005) found higher concentrations in cloud water as it was expected. Thus, the phase transfer of MEK from the gas in the aqueous phase could be more important as currently derived from the available Henry constants. Besides the phase transfer, the in-situ formation of MEK in aqueous phase formation might also represents an important source. As, the oxidation of MEK yielding methylglyoxal has not been studied much before, it should be considered as a formation process of methylglyoxal.”

Minor comments:

1) Methylglyoxal is conventionally one word in English, please change throughout.

Author`s comment

The authors agree with the reviewers comment. The term “*methyl glyoxal*” was changed to “*methylglyoxal*” throughout the manuscript.

2) Was there any gas phase in the reactor? Was it in equilibrium with the aqueous phase?

Author`s comment

The bulk reactor was filled with 300 mL of the aqueous solution at the beginning of the experiment containing the precursor compound and H<sub>2</sub>O<sub>2</sub>. Over the course of the experiment

samples were taken in steps of 15 minutes (and for some cases 1 hour). Thus, the gas phase increased during the experiment. To take samples, the bulk reactor was opened. Even that the opening time was kept as small as possible the gas phase of the bulk reactor might be mixed with the laboratory air changing the chemical composition. Nevertheless, the Henry constant of methylglyoxal ( $K_H \approx 3.7 \times 10^3 \text{ M atm}^{-1}$ ; Betterton and Hoffmann, 1988) indicates that methylglyoxal will mainly stay in the aqueous phase even when the bulk reactor was opened. Contrary, the Henry constant of MEK was measured to be  $K_H = 7.7 - 50 \text{ M atm}^{-1}$ , thus a loss of MEK from the aqueous phase in the gas phase is possible. To proof the hypothesis, the results of blank experiment 2 can be used. As it was mentioned in the manuscript, no loss of MEK was observed. This indicates that MEK stays in the aqueous phase even during opening of the reactor.

3) p. 31898 line 7-8: please elaborate on how the relative photolysis rate constants 'confirm the experimental results'

Author`s comment

Fig. 1 B shows the molar yields of the oxidation products methylglyoxal, hydroxyacetone and 2,3-butanedione over a reaction period of 4 hours. As it can be seen, further reactions are taking place leading to lower molar yields of methylglyoxal, hydroxyacetone and 2,3-butanedione towards the end of the experiment.

The following rate constants for the OH radical reaction are reported in the literature:  $k_{OH} = 5.3 \times 10^8 - 1.1 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$  (Monod et al., 2005; Ervens et al., 2003; Herrmann et al., 2005; Tan et al., 2010) for methylglyoxal,  $k_{OH} = 0.8 - 1.2 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$  (Stefan and Bolton, 1999; Herrmann et al., 2005) for hydroxyacetone and  $k_{OH} = 1.4 - 1.8 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$  (Gligorovski and Herrmann, 2004, Lilie et al., 1968) for 2,3-butanedione. Additionally, photolysis rates were determined in the present study to be  $k_P = 3 \times 10^{-5} \text{ s}^{-1}$ ,  $2 \times 10^{-5} \text{ s}^{-1}$  and  $9 \times 10^{-6} \text{ s}^{-1}$  for methylglyoxal, hydroxyacetone and 2,3-butanedione, respectively.

According to the rate constants for the OH radical reaction reported in the literature and the measured photolysis rates, a faster consumption of methylglyoxal and hydroxyacetone can be expected than for 2,3-butanedione. Such a faster consumption can be clearly seen from Figure 1B because methylglyoxal and hydroxylacetone are completely consumed at the end of the experiment whereas 2,3-butanedione was still present. This might be caused by the smaller rate constant as well as by the lower photolysis rate.

According to the Referees' comment, the following sentence was added to the manuscript:  
Page 31898 Line 7-9 was changed to: *“This indicates a faster decomposition compared to 2,3-butanedione of methylglyoxal and hydroxyacetone, which showed a complete consumption at the end of the experiment (Fig. 1). 2,3-Butanedione was not completely consumed during the reaction time of 4 hours, which might be a result of the lower photolysis rate constants.”*

#### Referee #2

These authors offer an alternative pathway of methylglyoxal formation in the atmospheric aqueous phase. Namely, the authors have investigated the aqueous phase oxidation of methyl ethyl ketone (MEK) by highly reactive hydroxyl radicals (OH). Few reaction products of atmospheric relevance were detected and identified i.e. 2,3-butanedione, hydroxyacetone and methylglyoxal. A reaction mechanism has been proposed for the formation of methylglyoxal and evaluated with a model study. The experiments are very well done and the authors have carefully addressed all issues regarding the application of these results to atmospheric aqueous phase. The paper is concise and nicely written, with no excess information. ACP is an appropriate journal for the paper on this topic. It can therefore be published in ACP after minor revisions.

1) How relevant are the used concentrations of  $0.1 \text{ mmol l}^{-1}$  of MEK in these experiments compared to the concentrations of MEK in cloud water ranging between 70 and  $650 \text{ nmol l}^{-1}$  (van Pinxteren et al., 2005)? How the concentrations of MEK would impact the experiments?

#### Author's comment

The authors agree on the referees comment that the MEK concentration was much higher in the experiment than found in ambient measurements. Such a high MEK concentration of  $0.1 \text{ mmol L}^{-1}$  was necessary for the experiment to ensure the traceability of the reaction and in particular the formation of formed products.

With higher MEK concentrations, side reactions like photolysis of MEK will increase. In particular, more alkylperoxy radicals ( $\text{RO}_2$ ) will be present in the system and this will change the ratio of  $\text{RO}_2$  to hydroperoxy radicals ( $\text{HO}_2$ ). The  $[\text{RO}_2]/[\text{HO}_2]$  is a critical issue in terms of product distribution (e.g. Keywood et al., 2004; Docherty and Ziemann, 2003). As it can be seen in Figure 4, the reaction of  $\text{RO}_2$  radicals with  $\text{HO}_2$  radicals leads to the formation of organic peroxides (R14/R25), whereas the recombination of  $\text{RO}_2$  radicals leads to the formation of carbonyl compounds and alcohols. Thus, with higher  $\text{RO}_2$  radical concentration and a larger

$[RO_2]/[HO_2]$  the formation of organic peroxides might become of minor importance. In addition, if  $RO_2$  radical concentration is not sufficient side reactions like recombination of  $HO_2$  radicals (R5) or the reaction of  $HO_2$  radicals with OH radicals (R4) and  $H_2O_2$  (R6) get more importance. This will lead to additional sinks for OH and  $HO_2$  radicals.

Furthermore, if the concentrations of MEK or  $H_2O_2$  are too high the bulk reactor will not be completely penetrated by light. For that reason, a concentration of  $0.1 \text{ mmol L}^{-1}$  ensure a good balance to detect the formed products, to have a complete illuminated reactor and to avoid strong photolysis of MEK during the oxidation.

2) The authors did not precise at what temperature were performed the experiments? I suppose at ambient temperature (298K).

Author`s comment

The experiments were conducted at a temperature of 298 K.

Based on the referees` comment the following sentence was added:

Page 31895, Line 8: The sentence "*The experiments were conducted at a temperature of 298 K.*" was included in the manuscript.

How important is this pathway of methyl glyoxal formation in comparison to the other sources? Reactions of carbonyl compounds in tropospheric clouds produce organic aerosol mass through in-cloud oxidation and during post-cloud evaporation. To get a clearer picture about the impact of these processes on the SOA formation one has to compare the OH radical reactivity in the aqueous phase with the evaporation post-cloud processes at lower temperatures, at the range between 5 and 15 °C relevant for the tropospheric clouds. I would like to see discussion on this issue.

Author`s comment

Methylglyoxal can be originated in the aqueous phase through i) uptake from the gas phase, and ii) aqueous-phase reactions as it is discussed in comment 1 for referee #1. The uptake of methylglyoxal from the gas phase in the aqueous phase is not fully understood, thus the formation of methylglyoxal through aqueous phase reaction, in particular through oxidation of MEK might represent an important formation process for methylglyoxal. The temperature dependency was not the focus of this study and thus it was not investigated.

However, an increasing rate constant for the OH radical oxidation of MEK at higher temperature was found by Mezyk (1994). The rate constant increase from  $k = 0.490 \pm 0.005 \times 10^{-9} \text{ M}^{-1} \text{ s}^{-1}$  at 1.5 °C to  $k = 1.35 \pm 0.07 \times 10^{-9} \text{ M}^{-1} \text{ s}^{-1}$  at 66.5 °C (Table 2).

Table 2: Rate constants and Henry constants for MEK at different temperatures.

Henry constant [M atm <sup>-1</sup> ] <sup>a</sup>	Temperature [°C]	Rate constant × 10 <sup>-9</sup> [M <sup>-1</sup> s <sup>-1</sup> ] <sup>b</sup>	Temperature [°C]
49.0 ± 0.8	10	0.490 ± 0.005	1.5
19.8 ± 1.4	25	0.590 ± 0.001	13.5
14.1 ± 0.2	30	0.734 ± 0.008	24.4
10.9 ± 0.5	35	0.936 ± 0.009	41.5
		1.35 ± 0.07	66.5

a: Zhou and Mopper (1990), b: Mezyk (1994)

This indicates a faster OH radical oxidation at higher temperatures and hence a faster formation of methylglyoxal can be expected. On the other hand, it was found by Zhou and Mopper (1990), that the Henry constant decrease from  $K_H = 49 \pm 0.8 \text{ M atm}^{-1}$  at 0°C to  $K_H = 7.1 \pm 0.7 \text{ M atm}^{-1}$  at 45°C (Table 2), which shows a stronger partitioning of MEK from the gas phase in the aqueous phase with decreasing temperature. Thus, higher concentrations of MEK should be found in the aqueous phase at lower temperatures.


In summary, the influence of the temperature cannot be finally estimated because of the contrary dependency of the Henry constant and the rate constant for the OH radical oxidation.

Minor comments:

1) Perhaps a right axis can be added in Figure 1 with different scaling than the left axis in order to better present the molar yield of hydroxyacetone. In figure 3 you either describe MEK (black) and 2,3 butanedione (red) in the caption or in the legend. The same applies for Figure 6.

Author`s comment

A right axis was included in Figure 1 for the molar yield of hydroxyacetone.



In Figure 1, 3, 6, 7, S2 and S3 the description in the figure caption “black” for MEK, “green” for methylglyoxal, “blue” for hydroxyacetone and “red” for 2,3-butanedione was deleted.

### References:

- Ashworth, R. A., Howe, G. B., Mullins, M. E., and Rogers, T. N.: Air water partitioning coefficients of organics in dilute aqueous-solutions, *J. Hazard. Mater.*, 18, 25-36, 10.1016/0304-3894(88)85057-x, 1988.
- Betterton, E. A., and Hoffmann, M. R.: Henry’s law constants of some environmentally important aldehydes. *Environ. Sci.Technol.*, 22, 1415–1418, 1988.
- Blando, J. D., and Turpin B. J.: Secondary organic aerosol formation in cloud and fog droplets: a literature evaluation of plausibility, *Atmos. Environ.*, 34, 1623-1632, 2000.
- Buttery, R. G., Ling, L. C., and Guadagni, D. G.: Food volatiles - volatilities of aldehydes ketones and esters in dilute water solution, *J. Agric. Food Chem.*, 17, 385-&, 10.1021/jf60162a025, 1969.
- Collett, J.L., Daube, B.C., Gunz, D., and Hoffmann, M.R.: Intensive studies of Sierra Nevada cloud water chemistry and its relationship to precursor aerosol and gas concentrations. *Atmos. Environ.*, 7, 1741–1757, 1990.
- Docherty, K. S., and Ziemann, P. J.: Effects of Stabilized Criegee Intermediate and OH Radical Scavengers on Aerosol Formation from Reactions of  $\beta$ -Pinene with  $\text{O}_3$ , *Aero. Sci. Technol*, 37, 877-891, 2003.
- Ervens, B., Gligorovski, S., and Herrmann, H.: Temperature-dependent rate constants for hydroxyl radical reactions with organic compounds in aqueous solutions, *Phys. Chem. Chem. Phys.*, 5, 1811-1824, 10.1039/b300072a, 2003.
- Feng, Y. L., Wen, S., Wang, X. M., Sheng, G. Y., He, Q. S., Tang, J. H., and Fu, J. M.: Indoor and outdoor carbonyl compounds in the hotel ballrooms in Guangzhou, China, *Atmospheric Environment*, 38, 103-112, 10.1016/j.atmosenv.2003.09.061, 2004.
- Fu, T-M., Jacob, D. J., Wittrock, F., Burrows, J. P., Vrekoussis, M., and Henze, D. K.: Global budgets of atmospheric glyoxal and methylglyoxal, and implications for formation of secondary organic aerosols, *J. Geophys. Res.*, 113, 1-17, 2008.
- Fu, T-M., Jacob, D. J., and Heald, C. L.: Aqueous-phase reactive uptake of dicarbonyls as a source of organic aerosol over eastern North America, *Atmos. Environ.*, 43, 1814–1822, 2009.
- Gligorovski, S., and Herrmann, H.: Kinetics of reactions of OH with organic carbonyl compounds in aqueous solution, *Phys. Chem. Chem. Phys.*, 6, 4118-4126, 10.1039/b403070b, 2004.
- Goldan, P. D., Kuster, W. C., Fehsenfeld, F. C., and Montzka, S. A.: Hydrocarbon measurements in the southeastern United States: The Rural Oxidants in the Southern Environment (ROSE) program 1990, *Journal of Geophysical Research-Atmospheres*, 100, 25945-25963, 10.1029/95jd02607, 1995.
- Grosjean, D., and Wright, B.: Carbonyls in urban fog, ice fog, cloudwater and rainwater, *Atmospheric Environment*, 17, 2093-2096, 10.1016/0004-6981(83)90368-2, 1983.
- Grosjean, D.: Formaldehyde and other carbonyls in Los-Angeles ambient air, *Environmental Science & Technology*, 16, 254-262, 10.1021/es00099a005, 1982.
- Grosjean, D., Swanson, R. D., and Ellis, C.: Carbonyls in Los-Angeles air - contribution of direct emissions and photochemistry, *Sci. Total Environ.*, 29, 65-85, 10.1016/0048-9697(83)90034-7, 1983.



- Grosjean, D., Grosjean, E., and Moreira, L. F. R.: Speciated ambient carbonyls in Rio de Janeiro, Brazil, *Environmental Science & Technology*, 36, 1389-1395, 10.1021/es0111232, 2002.
- Herrmann, H., Tilgner, A., Barzagli, P., Majdik, Z., Gligorovski, S., Poulain, L., and Monod, A.: Towards a more detailed description of tropospheric aqueous phase organic chemistry: CAPRAM 3.0, *Atmospheric Environment*, 39, 4351-4363, 10.1016/j.atmosenv.2005.02.016, 2005.
- Ieda, T., Kitamori, Y., Mochida, M., Hirata, R., Hirano, T., Inukai, K., Fujinuma, Y., and Kawamura, K.: Diurnal variations and vertical gradients of biogenic volatile and semi-volatile organic compounds at the Tomakomai larch forest station in Japan, *Tellus B*, 58, 177-186, 2006.
- Karl, T., Chahan, Y., Jordan, A., and Lindinger, W.: Dynamic measurements of partition coefficients using proton-transfer-reaction mass spectrometry (PTR-MS), *International Journal of Mass Spectrometry*, 12269, 1-13, 2002.
- Kawamura, K., Okuzawa, K., Aggarwal, S. G., Irie, H., Kanaya, Y., and Wang, Z.: Determination of gaseous and particulate carbonyls (glycolaldehyde, hydroxyacetone, glyoxal, methylglyoxal, nonanal and decanal) in the atmosphere at Mt. Tai, *Atmos. Chem. Phys.*, 13, 5369 – 5380, 2013.
- Keywood, M. D., Kroll, J. H., Varutbangkul, V., Bahreini R., Flagan, R. C., and Seinfeld, J. H.: Secondary Organic Aerosol Formation from Cyclohexene Ozonolysis: Effect of OH Scavenger and the Role of Radical Chemistry, *Environ. Sci. Technol*, 38, 3343-3350, 2004.
- Kroll, J. H., Ng, N. L., Murphy, S. M., Varutbangkul, V., Flagan, R. C., and Seinfeld, J. H.: Chamber studies of secondary organic aerosol growth by reactive uptake of simple carbonyl compounds, *J. Geophys. Res.*, 110, 1-10, 2005.
- Lilie, J., Beck, G., and Henglein, A.: Pulse radiolysis of acetoin radicals and diacetyl anions in aqueous solution, *Berichte Der Bunsen-Gesellschaft Fur Physikalische Chemie*, 72, 529-&, 1968.
- Lin, G., Sillman, S., Penner, J. E., and Ito, A.: Global modeling of SOA: the use of different mechanisms for aqueous-phase formation, *Atmos. Chem. Phys.*, 14, 5451-5475, 2014.
- Matsunaga, S., Mochida, M., and Kawamura, K.: Variation on the atmospheric concentrations of biogenic carbonyl compounds and their removal processes in the northern forest at Moshiri, Hokkaido Island in Japan, *J. Geophys. Res.*, 109, , doi:10.1029/2003JD004100, 2004.
- Mezyk, S. P.: Rate constant and activation energy determination for reaction of e<sup>-</sup>(aq) and OH with 2-butanone and propanal, *Can. J. Chem.*, 72, 116 – 1119, 1994.
- Monod, A., Poulain, L., Grubert, S., Voisin, D., and Wortham, H.: Kinetics of OH-initiated oxidation of oxygenated organic compounds in the aqueous phase: new rate constants, structure-activity relationships and atmospheric implications, *Atmospheric Environment*, 39, 7667-7688, 10.1016/j.atmosenv.2005.03.019, 2005.
- Morillon, V., Debeaufort, F., Jose, J., Tharrault, J. F., Capelle, M., Blond, G., and Voilley, A.: Water vapour pressure above saturated salt solutions at low temperatures, *Fluid Phase Equilib.*, 155, 297-309, 10.1016/s0378-3812(99)00009-6, 1999.
- Müller, K., van Pinxteren, D., Plewka, A., Svrčina, B., Kramberger, H., Hofmann, D., Bachmann, K., and Herrmann, H.: Aerosol characterisation at the FEBUKO upwind station Goldlauter (II): Detailed organic chemical characterisation, *Atmospheric Environment*, 39, 4219-4231, 10.1016/j.atmosenv.2005.02.008, 2005.
- Munger, J. W., Jacob, D. J., Daube, B. C., Horowitz, L. W., Keene, W. C., and Heikes, B. G.: Formaldehyde, glyoxal, and methylglyoxal in air and cloud water at a rural mountain site in central Virginia, *J. Geophys. Res.*, 100, 9325 – 9333, 1995.
- Ortiz, R., Hagino, H., Sekiguchi, K., Wang, Q., and Sakamoto, K.: Ambient air measurements of six bifunctional carbonyls in a suburban area, *Atmos. Res.*, 82, 709-718, 2006.
- Riemer, D., Pos, W., Milne, P., Farmer, C., Zika, R., Apel, E., Olszyna, K., Kliendienst, T., Lonneman, W., Bertman, S., Shepson, P., and Starn, T.: Observations of nonmethane hydrocarbons and oxygenated volatile organic compounds at a rural site in the southeastern United States, *Journal of Geophysical Research-Atmospheres*, 103, 28111-28128, 10.1029/98jd02677, 1998.
- Schütze, M., and Herrmann, H.: Uptake of acetone, 2-butanone, 2,3-butanedione and 2-oxopropanal on a water surface, *Phys. Chem. Chem. Phys.*, 6, 965-971, 2004.
- Sempere, R., and Kawamura K.: Comparative distribution of dicarboxylic acids and related polar compounds in snow, rain and aerosols from urban, *Atmos. Environ.*, 28, 449-459, 1994.
- Singh, H. B., Salas, L. J., Chatfield, R. B., Czech, E., Fried, A., Walega, J., Evans, M. J., Field, B. D., Jacob, D. J., Blake, D., Heikes, B., Talbot, R., Sachse, G., Crawford, J. H., Avery, M. A., Sandholm, S., and Fuelberg, H.: Analysis of the atmospheric distribution, sources, and sinks of oxygenated volatile organic chemicals based on measurements over the Pacific during TRACE-P, *Journal of Geophysical Research-Atmospheres*, 109, 20, 10.1029/2003jd003883, 2004.
- Snider, J. R., and Dawson, G. A.: Tropospheric light alcohols, carbonyls and acetonitrile - concentrations in southwestern united-states and henry law data, *Journal of Geophysical Research-Atmospheres*, 90, 3797-3805, 10.1029/JD090iD02p03797, 1985.

- Stefan, M. I., and Bolton, J. R.: Reinvestigation of the acetone degradation mechanism in dilute aqueous solution by the UV/H<sub>2</sub>O<sub>2</sub> process, *Environmental Science & Technology*, 33, 870-873, 10.1021/es9808548, 1999.
- Tan, Y., Carlton, A. G., Seitzinger, S. P., and Turpin, B. J.: SOA from methylglyoxal in clouds and wet aerosols: Measurement and prediction of key products, *Atmospheric Environment*, 44, 5218-5226, 10.1016/j.atmosenv.2010.08.045, 2010.
- van Pinxteren, D., Plewka, A., Hofmann, D., Müller, K., Kramberger, H., Svrčina, B., Bachmann, K., Jaeschke, W., Mertes, S., Collett, J. L., and Herrmann, H.: Schmucke hill cap cloud and valley stations aerosol characterisation during FEBUKO (II): organic compounds, *Atmos. Environ.*, 39, 4305–4320, 2005.
- Zhao, J., Levitt, N. P., Zhang, R., and Chen, J.: Heterogeneous Reactions of Methylglyoxal in Acidic Media: Implications for Secondary Organic Aerosol Formation, *Environ. Sci. Technol.*, 40, 7682-7687, 2006.
- Zhou, X. L., and Mopper, K.: Apparent partition-coefficients of 15 carbonyl-compounds between air and seawater and between air and fresh-water – implications for air sea exchange, *Environ. Sci. Technol.*, 24, 1864–1869, 1990.
- Zhou, X. L., and Mopper, K.: Photochemical production of low-molecular-weight carbonyl compounds in seawater and surface microlayer and their air-sea exchange, *Mar. Chem.*, 56, 201-213, 10.1016/s0304-4203(96)00076-x, 1997.

### The following changes were made to the manuscript

The term “*methyl glyoxal*” was changed to “*methylglyoxal*” in the manuscript text as well as in Fig. 1, 4, 6, 7 and Table 1 and 2.

Page 31895 Line 8: The sentence “*The experiments were conducted at a temperature of 298 K.*” was included in the manuscript.

Page 31898 Line 7-9: “*This indicates a faster decomposition of methyl glyoxal and hydroxyacetone and confirms the experimental results.*” was changed to “*This indicates a faster decomposition compared to 2,3-butanedione of methylglyoxal and hydroxyacetone, which showed a complete consumption at the end of the experiment (Fig. 1). 2,3-Butanedione was not completely consumed during the reaction time of 4 hours, which might be a result of the lower photolysis rate constants.*”

Page 31898 Line 24-25: “*This is in good agreement with the experimental results obtained in the the present study.*” was changed to “*This is in good agreement with the experimental results obtained in the present study.*”

Page 31906 Line 1: The paragraph “*4. Atmospheric relevance*

*The sources of methylglyoxal in the aqueous phase are thus far not fully elucidated. Methylglyoxal can originate in the atmospheric aqueous phase through i) uptake from the gas phase, and/or ii) formation in the aqueous phase. The importance of the uptake from the gas into the aqueous phase is discussed in the literature but large discrepancies can be found. Kroll et al. (2005) investigated the uptake of methylglyoxal on inorganic seed particles under varying relative humidity. It was found that the uptake was not relevant for methylglyoxal even under*

high relative humidity. Contrary, Zhao et al. (2006) measured an uptake coefficient of  $\gamma = 7.6 \times 10^{-3}$  on acidic solution. They found an irreversible uptake, which decreases with increasing acidity. Fu et al. (2008, 2009) determined an uptake coefficient on aqueous particles and cloud droplets  $\gamma = 2.9 \times 10^{-3}$ , which is in good agreement with the uptake coefficient measured by Zhao et al. (2006). Lin et al. (2014) modelled an uptake coefficient for several case studies with different multiphase process mechanisms. The uptake coefficient in deliquescent particles was determined to be  $\gamma = 1.47 - 2.92 \times 10^{-5}$ . Overall,  $\gamma$  was two orders of magnitude lower compared to the values determined by Zhao et al. (2006) and Fu et al. (2008, 2009) showing the discrepancies between the different studies available in literature. The in-situ formation of methylglyoxal in the aqueous phase could be an important source as well (Blando and Turpin, 2000; Sempere and Kawamura 1994). Within the present study, MEK was found as a new precursor compound for methylglyoxal in the aqueous phase yielding methylglyoxal with a molar yield of 9.5%. Although the Henry constant of MEK (up to  $K_H = 50 \text{ M atm}^{-1}$ , Schütze and Herrmann, 2004) is lower compared to methylglyoxal ( $K_H = 3.7 \times 10^3 \text{ M atm}^{-1}$ , Betterton and Hoffmann, 1988), van Pinxteren et al. (2005) found higher concentrations in cloud water as it was expected. Thus, the phase transfer of MEK from the gas in the aqueous phase could be more important as currently derived from the available Henry constants. Besides the phase transfer, the in-situ formation of MEK in aqueous phase formation might also represents an important source.

As, the oxidation of MEK yielding methylglyoxal has not been studied much before, it should be considered as a formation process of methylglyoxal.” was included in the manuscript.

Page 31906 Line 1: “4. Summary” was changed to “5. Summary”

Page 31908 Line 13: The reference “Blando, J. D., and Turpin B. J.: Secondary organic aerosol formation in cloud and fog droplets: a literature evaluation of plausibility, *Atmos. Environ.*, 34, 1623-1632, 2000.” was included in the manuscript.

Page 31909 Line 28: The references “Fu, T-M., Jacob, D. J., Wittrock, F., Burrows, J. P., Vrekoussis, M., and Henze, D. K., Global budgets of atmospheric glyoxal and methylglyoxal, and implications for formation of secondary organic aerosols, *J. Geophys. Res.*, 113, 1-17, 2008.” and “Fu, T-M., Jacob, D. J., and Heald, C. L., Aqueous-phase reactive uptake of dicarbonyls as a source of organic aerosol over eastern North America, *Atmos. Environ.*, 43, 1814–1822, 2009.” were included in the manuscript.

Page 31911 Line 16: The reference “Kroll, J. H., Ng, N. L., Murphy, S. M., Varutbangkul, V., Flagan, R. C., and Seinfeld, J. H., *Chamber studies of secondary organic aerosol growth by reactive uptake of simple carbonyl compounds*, *J. Geophys. Res.*, 110, 1-10, 2005.” was added to the manuscript.

Page 31911 Line 18: The reference “Lin, G., Sillman, S., Penner, J. E., and Ito, A., *Global modeling of SOA: the use of different mechanisms for aqueous-phase formation*, *Atmos. Chem. Phys.*, 14, 5451–5475, 2014.” was included in the manuscript.

Page 31912 Line 31: The reference “Sempere, R., and Kawamura K.: *Comparative distribution of dicarboxylic acids and related polar compounds in snow, rain and aerosols from urban*, *Atmos. Environ.*, 28, 449-459, 1994.” was included in the manuscript.

Page 31914 Line 1: The reference: “Zhao, J., Levitt, N. P., Zhang, R., and Chen, J., *Heterogeneous Reactions of Methylglyoxal in Acidic Media: Implications for Secondary Organic Aerosol Formation*, *Environ. Sci. Technol.*, 40, 7682-7687, 2006.” was added to the manuscript.

Page 31918: The figure caption “*Figure 1: Consumption of MEK (black; A) during the oxidation with OH radicals and time-resolved formation of the products methylglyoxal (green; B), 2,3-butanedione (red; B), and hydroxyacetone (blue; B)*” was changed to “*Figure 1: Consumption of MEK (A) during the oxidation with OH radicals and time-resolved formation of the products methylglyoxal (B), 2,3-butanedione (B), and hydroxyacetone (B).*”

Page 31920: The figure caption “*Figure 3: Photolysis of MEK (black) and time-resolved formation of 2,3-butanedione (red).*” was changed to “*Figure 3: Photolysis of MEK and time-resolved formation of 2,3-butanedione.*”

Page 31923: The figure caption “*Figure 6: Comparison of the molar yields of 2,3-butanedione (A, red) and methylglyoxal (B, green) for the model and experimental results using different branching ratios of the primary/secondary H-atom abstraction varied between 60/40 up to 10/90.*” was changed to “*Figure 6: Comparison of the molar yields of 2,3-butanedione (A) and methylglyoxal (B) for the model and experimental results using different branching ratios of the primary/secondary H-atom abstraction varied between 60/40 up to 10/90.*”

Page 31924: The figure caption “*Figure 7: Comparison of the model and experimental results for MEK (A, black), 2,3-butanedione (B, red), and methylglyoxal (C, green).*” was changed to

*“Figure 7: Comparison of the model and experimental results for MEK (A), 2,3-butanedione (B), and methylglyoxal (C).”*

### General changes made to the supplement

The term “*methyl glyoxal*” was changed to “*methylglyoxal*” in the supplement text and Fig, S1, S2 and S3.

Page 2: The figure caption “*Figure S2: Formation of methylglyoxal (green) due to the photolysis of hydroxyacetone (blue) and 2,3-butanedione (red).*” was changed to “*Figure S2: Formation of methylglyoxal due to the photolysis of hydroxyacetone and 2,3-butanedione.*”

Page 3: The figure caption “*Figure S3: Oxidation of hydroxyacetone (blue) and formation of methylglyoxal (green).*” was changed to “*Figure S3: Oxidation of hydroxyacetone and formation of methylglyoxal.*”

# A new source of methylglyoxal in the aqueous phase

Maria Rodigast, Anke Mutzel, Janine Schindelka and Hartmut Herrmann\*

*Leibniz Institute for Tropospheric Research (TROPOS), Atmospheric Chemistry Dept. (ACD), Permoserstr. 15,  
D-04318 Leipzig, Germany*

\*Corresponding author. Tel: +49-341-2717-7024; fax: +49-341-271799-7024.

*E-mail address:* herrmann@tropos.de

For submission to: Atmospheric Chemistry and Physics

First submitted on 27.10.2015

---

## 1 Abstract

2 Carbonyl compounds are ubiquitous in atmospheric multiphase system participating in gas, particle, and aqueous-  
3 phase chemistry. One important compound is methyl ethyl ketone (MEK), as it is detected in significant amounts  
4 in the gas phase as well as in cloud water, ice, and rain. Consequently, it can be expected that MEK influences the  
5 liquid phase chemistry. Therefore, the oxidation of MEK and the formation of corresponding oxidation products  
6 were investigated in the aqueous phase. Several oxidation products were identified from the oxidation with OH  
7 radicals, including 2,3-butanedione, hydroxyacetone, and methylglyoxal. The molar yields were 29.5% for  
8 2,3-butanedione, 3.0% for hydroxyacetone, and 9.5% for methylglyoxal. Since methylglyoxal is often related to  
9 the formation of organics in the aqueous phase, MEK should be considered for the formation of aqueous secondary  
10 organic aerosol (aqSOA). Based on the experimentally obtained data, a reaction mechanism for the formation of  
11 methylglyoxal has been developed and evaluated with a model study. Besides known rate constants, the model  
12 contains measured photolysis rate constants for MEK ( $k_p = 5 \times 10^{-5} \text{ s}^{-1}$ ), 2,3-butanedione ( $k_p = 9 \times 10^{-6} \text{ s}^{-1}$ ),  
13 methylglyoxal ( $k_p = 3 \times 10^{-5} \text{ s}^{-1}$ ), and hydroxyacetone ( $k_p = 2 \times 10^{-5} \text{ s}^{-1}$ ). From the model predictions, a branching  
14 ratio of 60/40 for primary/secondary H-atom abstraction at the MEK skeleton was found. This branching ratio  
15 reproduces the experiment results very well, especially the methylglyoxal formation, which showed excellent  
16 agreement. Overall, this study demonstrates MEK as a methylglyoxal precursor compound for the first time.  
17

## 18 1. Introduction

19 In the last decades, carbonyl compounds have been a subject of intense research due to their ubiquitous abundance  
20 and their effect on atmospheric chemistry and human health. They are emitted directly from biogenic and  
21 anthropogenic sources or formed through the oxidation of hydrocarbons (e.g., Atkinson, 1997; Matthews and  
22 Howell, 1981; Lipari et al., 1984; Ciccioli et al., 1993; Mopper and Stahovec, 1986; Carlier et al., 1986; Hallquist  
23 et al., 2009). One carbonyl compound that is emitted from numerous and mainly biological sources is methyl ethyl  
24 ketone (MEK). It is released from grass, clover (Kirstine et al., 1998; de Gouw et al., 1999), different types of  
25 forests, and biomass burning processes (Khalil and Rasmussen, 1992; Warneke et al., 1999; Isidorov et al., 1985).  
26 Anthropogenic emissions are also important MEK sources, such as artificial biomass burning (Andreae and  
27 Merlet, 2001; Akagi et al., 2011; Yokelson et al., 2013; Brilli et al., 2014) and tobacco smoke (Buyske et al.,  
28 1956; Yokelson et al., 2013). In addition, MEK is emitted into the atmosphere through the application as solvent  
29 for the production of glue, resins, cellulose, rubber, paraffin wax and lacquer (Ware, 1988).

30 Tropospheric MEK gas-phase concentration was found to be in the range of 0.02 – 15 ppbv, depending  
31 on the region (Grosjean et al., 2002; Riemer et al., 1998; Singh et al., 2004; Snider and Dawson, 1985; Goldan et  
32 al., 1995; Grosjean et al., 1983; Grosjean, 1982; Müller et al., 2005; Feng et al., 2004). Singh et al. (2004)  
33 measured a concentration in a remote region of 0.02 ppbv, whereas Grosjean et al. (1983) observed a MEK  
34 concentration of 11.3 ppbv in Los Angeles. Brown et al. (1994) concluded that MEK is one of the major volatile  
35 organic compounds (VOCs) in indoor air.

36 In addition to the gas-phase measurements, the concentrations measured in bulk water samples collected  
37 at an open station near the Bahamas reached a concentration of  $< 0.5 \text{ nmol L}^{-1}$  (Zhou and Mopper, 1997).  
38 Furthermore, an enrichment of MEK in the surface micro layer was found with concentrations up to  $2.28 \text{ nmol L}^{-1}$   
39 (Zhou and Mopper, 1997). MEK was also investigated in ice, fog, and rain samples (Grosjean and Wright, 1983).  
40 It was not found in fog but there were traces in rain water. In cloud water, a concentration of up to  $650 \text{ nmol L}^{-1}$   
41 was measured. This is supported by van Pinxteren et al. (2005), who measured a concentration of  
42 70 to  $300 \text{ nmol L}^{-1}$  in cloud water. These studies concluded that the liquid-phase fraction of MEK is higher than  
43 the expected fraction calculated according to the Henry constant.

44 The Henry constants at a temperature of  $25 \text{ }^{\circ}\text{C}$  were found to vary between  $7.7$  and  $21 \text{ M atm}^{-1}$  in  
45 numerous studies (Buttery et al., 1969; Snider and Dawson, 1985; Ashworth et al., 1988; Zhou and Mopper, 1990;  
46 Morillon et al., 1999; Karl et al., 2002). However, Schütze and Herrmann (2004) estimated the Henry constant to  
47 be between  $23$  and  $50 \text{ M atm}^{-1}$ , which is higher than the previous measured values found in the literature. This  
48 higher Henry constant supports the conclusion from van Pinxteren et al. (2005) and tends to support the  
49 investigation of MEK in the liquid phase as aqSOA precursor compound. AqSOA is formed through the oxidation  
50 of organic compounds in the aqueous particle phase and is often related to missing SOA sources. These missing  
51 sources are most likely responsible for the huge discrepancies between measured and calculated SOA burden. As  
52 model results usually underestimate the SOA burden (Kanakidou et al., 2005; Goldstein and Galbally, 2007),  
53 missing SOA sources have to be considered in such models to close this gap (Ervens et al., 2011; Herrmann et al.,  
54 1999; Herrmann et al., 2015).

55 In the present study, the reaction of MEK with OH radicals in water was investigated. Based on the  
56 experimentally obtained data, a reaction mechanism was developed to explain methylglyoxal formation. The

57 mechanism was included in a COPASI (Complex Pathway Simulator) model and evaluated by comparing the  
58 experimentally obtained data and the model results.

## 59 2. Experimental

### 60 2.1 Chemicals and standards

61 Cyclohexanone-2,2,6,6-d<sub>4</sub> (98%), hydrochloric acid, and catalase from bovine liver (40000-60000 units mg<sup>-1</sup>  
62 protein) were obtained from Sigma-Aldrich (Hamburg, Germany). *O*-(2,3,4,5,6-  
63 pentafluorobenzyl)hydroxylamine hydrochloride ( $\geq 99\%$ ), 2,3-butanedione (99%), hydroxyacetone (90%), and  
64 methylglyoxal (40% in water) were purchased from Fluka (Hamburg, Germany). Dichloromethane (Chromasolv  
65 99.8%) and methyl ethyl ketone (99.7%) were obtained from Riedel-de Haen (Seelze, Germany), and hydrogen  
66 peroxide (30% Suprapur<sup>®</sup>) was obtained from Merck KGaA (Darmstadt, Germany). Ultrapure water was used to  
67 prepare the reaction solutions for the bulk reactor experiments and the stock solutions of the authentic standard  
68 compounds (Milli-Q gradient A 10, 18.2 M $\Omega$  cm<sup>-1</sup>, 3 ppb TOC, Millipore, USA).

### 69 2.2 Bulk reactor experiments

70 The aqueous-phase oxidation of MEK was conducted in a 300 mL batch reactor using the photolysis of hydrogen  
71 peroxide (H<sub>2</sub>O<sub>2</sub>) as an OH radical source (Set 1). The experiments were conducted at a temperature of 298 K. For  
72 the experiments, 0.1 mmol L<sup>-1</sup> of the precursor compound was mixed with 2 mmol L<sup>-1</sup> of H<sub>2</sub>O<sub>2</sub>. The solution was  
73 then irradiated at  $\lambda = 254$  nm for 4 hours to continuously generate OH radicals (500 W xenon-mercury lamp,  
74 Andover Corporation Optical bandpass filter: L.O.T.- Oriel GmbH & Co. KG, Darmstadt, Germany). To obtain  
75 time-resolved data, samples were taken once per hour (0 – 4 hours; number of repetitions  $n = 3$ ) or in steps of  
76 15 minutes ( $n = 1$ ) and analyzed after derivatization using GC/MS. To avoid further reactions of the organics  
77 present in the sample with remaining H<sub>2</sub>O<sub>2</sub>, 100  $\mu$ L of catalase (4 mg mL<sup>-1</sup> in water) were added to each sample.  
78 Furthermore, a set of blank experiments was conducted to exclude that (a) MEK reacts with H<sub>2</sub>O<sub>2</sub> (Set 2),  
79 (b) identified oxidation products originate from photolysis (Set 3), (c) the photolysis of contaminants in the H<sub>2</sub>O<sub>2</sub>  
80 solution results in the formation of organics (Set 4), and (d) oxidation of 2,3-butanedione or hydroxyacetone also  
81 forms methylglyoxal (Set 5/6). To determine the photolysis rate constants of 2,3-butanedione, methylglyoxal, and  
82 hydroxyacetone, the photolysis of these products was investigated as well with the setup used (Set 7). A complete  
83 overview of the experiments conducted is given in Table 1.

### 84 2.3 Sample preparation

85 Two types of samples were taken over a period of 4 hours. For the first type of samples, 60  $\mu$ L of the reaction  
86 mixture were diluted with 2940  $\mu$ L of water to avoid saturation of the GC/MS detector during the quantification  
87 of MEK, 2,3-butanedione or hydroxyacetone. For the second type of samples, 3 mL of the reaction mixture were  
88 taken and injected without any dilution to enable the identification and quantification of the formed carbonyl  
89 compounds.

90 Samples of all sets were derivatized with 300  $\mu$ L of *o*-(2,3,4,5,6-pentafluorobenzyl)hydroxylamine  
91 hydrochloride (PFBHA, 5 mg mL<sup>-1</sup>) at room temperature (Rodigast et al., 2015). Cyclohexanone-2,2,6,6-d<sub>4</sub> was  
92 used as an internal standard (150  $\mu$ L, 100  $\mu$ mol L<sup>-1</sup>), and after 24 hours, a pH value of 1 was adjusted by adding  
93 hydrochloric acid (37%) to the reaction mixture. The target compounds were extracted for 30 minutes with 250  $\mu$ L  
94 of dichloromethane using an orbital shaker (1500 rpm, revolutions per minutes). Finally, 1  $\mu$ L of the organic phase



95 was used for GC/MS analysis. A 5-point calibration was performed for each chromatographic run using a series  
96 of the standard solutions (MEK, methylglyoxal, 2,3-butanedione, and hydroxyacetone) with concentrations  
97 ranging from 1 to 50  $\mu\text{mol L}^{-1}$ .

#### 98 2.4 Instrumentation

99 Derivatized carbonyl compounds were analyzed using a GC System (6890 Series Agilent Technologies, Frankfurt,  
100 Germany) coupled with an electron ionization quadrupole mass spectrometer in splitless mode at a temperature  
101 of 250 °C (Agilent 5973 Network mass selective detector, Frankfurt, Germany). They were separated with an HP-  
102 5MS UI column (Agilent J & W GC columns, 30 m  $\times$  0.25 mm  $\times$  0.25  $\mu\text{m}$ ) using the following temperature  
103 program: 50 °C isothermal for 2 minutes and elevated to 230 °C with 10 °C  $\text{minute}^{-1}$ . The temperature of 230 °C  
104 was held for 1 minute, and the temperature gradient ended with 320 °C, which was held constant for 10 minutes.

#### 105 2.4 Instrumentation

106 Derivatized carbonyl compounds were analysed using a GC System (6890 Series Agilent Technologies, Frankfurt,  
107 Germany) coupled with an electron ionization quadrupole mass spectrometer (Agilent 5973 Network mass  
108 selective detector, Frankfurt, Germany) in the splitless mode at a temperature of 250 °C. They were separated  
109 with a HP-5MS UI column (Agilent J & W GC columns, 30 m  $\times$  0.25 mm  $\times$  0.25  $\mu\text{m}$ ) using the following  
110 temperature program: 50 °C isothermal for 2 minutes and elevated to 230 °C with 10 °C  $\text{minute}^{-1}$ . The temperature  
111 of 230 °C was held for 1 minute and the temperature gradient ended with 320 °C which was held constant for  
112 10 minutes.

### 113 3. Results

#### 114 3.1 Experimental results from bulk reactor

115 MEK was oxidized with OH radicals, and the decay of MEK was monitored by GC/MS. Fig. 1A shows the  
116 consumption of the precursor compound MEK. As can be seen, MEK was almost consumed after 180 minutes of  
117 reaction time. From the analysis of the collected samples, 2,3-butanedione, hydroxyacetone, and methylglyoxal  
118 were observed as the most dominating oxidation products. The formation of methylglyoxal from the oxidation of  
119 MEK was unexpected as it has not been reported in the literature before. Due to the relevance of methylglyoxal  
120 for aqSOA formation, its formation was comprehensively characterized in the present study.

121 Nevertheless, in Fig. 1B, 2,3-butanedione was found as the main oxidation product of MEK reaching a  
122 maximum yield of  $\approx 29.5 \pm 6.0\%$  after 60 minutes. The small variations of the molar yields over the course of the  
123 experiment might result from slight temperature changes during the experiment despite the reactor temperature  
124 being controlled. However, the curve shape shows that the concentration of 2,3-butanedione starts to decrease  
125 after 60 minutes, indicating further reactions of 2,3-butanedione. This decrease results in a molar yield of  
126  $18.9 \pm 3.9\%$  at the end of the experiment (240 minutes).

127 A similar trend was observed for hydroxyacetone and methylglyoxal, as the concentration of  
128 methylglyoxal was the highest after 15 minutes ( $\approx 9.5\%$ ) and started to decrease afterwards. In comparison,  
129 hydroxyacetone reached the highest concentration after 60 minutes with a molar yield of  $3.0 \pm 2.6\%$ . The standard  
130 deviation of the determined molar yields of hydroxyacetone was high, and therefore, only after reaction times of  
131 60 and 120 minutes could a molar yield of hydroxyacetone be determined.

132 **Methylglyoxal** and hydroxyacetone were completely consumed at the end of the experiment. The strong  
133 decrease of the concentrations of the detected carbonyl compounds might result from the reaction with OH radicals  
134 and/or from photolysis. Both mechanisms are most likely, as it has been demonstrated that the detected carbonyl  
135 compounds react quickly with OH radicals (Lilie et al., 1968; Gligorovski and Herrmann, 2004; Doussin and  
136 Monod, 2013; Monod et al., 2005; Ervens et al., 2003; Herrmann et al., 2005; Tan et al., 2010; Stefan and Bolton,  
137 1999) and they are prone to photolysis (Faust et al., 1997; Tan et al., 2010). The photolysis rate constants of the  
138 detected carbonyl compounds were determined in the present study because of the dependency on the setup used  
139 (Set 7; see supplement S2 for more details). **Methylglyoxal** and hydroxyacetone showed higher photolysis rate  
140 constants of  $k_p = 3 \times 10^{-5} \text{ s}^{-1}$  and  $k_p = 2 \times 10^{-5} \text{ s}^{-1}$  in comparison to 2,3-butanedione ( $k_p = 9 \times 10^{-6} \text{ s}^{-1}$ ). **This**  
141 **indicates a faster decomposition compared to 2,3-butanedione of methylglyoxal and hydroxyacetone, which**  
142 **showed a complete consumption at the end of the experiment (Fig. 1). 2,3-Butanedione was not completely**  
143 **consumed during the reaction time of 4 hours, which might be a result of the lower photolysis rate constants.**  
144 Furthermore, during the photolysis of 2,3-butanedione and hydroxyacetone, **methylglyoxal** was found with molar  
145 yields of  $\approx 17\%$  and  $\approx 19\%$ , respectively (see Fig. S2, supplement S2). Due to the low molar yield of  
146 hydroxyacetone during the oxidation of MEK ( $\approx 3\%$ ) and the slow photolysis rate constant of 2,3-butanedione,  
147 these processes are of minor importance for the **methylglyoxal** yield.

148 For 2,3-butanedione, a huge discrepancy of the rate constants for the OH radical oxidation can be found  
149 between the different literature studies. They vary by one order of magnitude in a range of  $k = 1.4 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$   
150 (Gligorovski and Herrmann, 2004) up to  $k = 1.86 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$  (Doussin and Monod, 2013). Rate constants were  
151 determined between  $k = 5.3 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$  and  $k = 1.1 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$  for **methylglyoxal** (Monod et al., 2005; Ervens  
152 et al., 2003; Herrmann et al., 2005; Tan et al., 2010) and between  $k = 0.8 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$  and  $k = 1.2 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$   
153 for hydroxyacetone (Stefan and Bolton, 1999; Herrmann et al., 2005). In consideration of the similar rate constant  
154 for 2,3-butanedione determined by Lilie et al. (1968) ( $k = 1.8 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ ) and Gligorovski and Herrmann (2004)  
155 ( $k = 1.4 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ ), 2,3-butanedione revealed a slower OH radical reaction than **methylglyoxal** and  
156 hydroxyacetone. **This is in good agreement with the experimental results obtained in the present study.**

157 Since 2,3-butanedione is the main oxidation product, it was necessary to investigate the contribution of  
158 2,3-butanedione to the product distribution, especially for the formation of **methylglyoxal**. In the oxidation of 2,3-  
159 butanedione (Set 5), no **methylglyoxal** was detected in the GC/MS chromatogram over a reaction period of  
160 240 min (Fig. 2). Consequently, a contribution of 2,3-butanedione to the **methylglyoxal** formation could be  
161 excluded.

162 Despite the low molar yield of hydroxyacetone during MEK oxidation, the oxidation of hydroxyacetone  
163 was investigated for **methylglyoxal** formation as well (Herrmann et al., 2005; Schaefer et al., 2012; Set 6). During  
164 the oxidation, a molar yield of 100% was found after a reaction time of 60 minutes (see supplement S3, Fig. S3).  
165 After 60 minutes of reaction time, the molar yield of **methylglyoxal** decreases through further reactions, as was  
166 observed during MEK oxidation. However, due to the low molar yield of hydroxyacetone (3.0%), the oxidation  
167 has only minor importance for the observed molar yield of **methylglyoxal**.

168 To ensure **methylglyoxal** was only formed during the oxidation of MEK, an experiment was conducted  
169 to investigate the non-radical reaction of MEK with  $\text{H}_2\text{O}_2$  in the dark (Set 2). During this experiment, no  
170 decomposition of MEK was observed, excluding the non-radical reaction of MEK with  $\text{H}_2\text{O}_2$  as a source of  
171 **methylglyoxal**. Furthermore, the photolysis of MEK has to be considered as a source for **methylglyoxal**. Hence,

172 the photolysis of MEK was studied at  $\lambda = 254$  nm (Set 3). As can be seen in Fig. 3, the photolysis of MEK leads  
173 to the formation of 2,3-butanedione with a molar yield of  $\approx 2.2\%$  indicating that the photolysis of MEK is an  
174 additional source. No further carbonyl compounds were detected from the photolysis of MEK, and thus, the  
175 photolysis of MEK can be excluded as a methylglyoxal source. The blank experiment (Set 4) indicated the absence  
176 of methylglyoxal as well.

177 In summary, the oxidation of MEK constitutes a source for methylglyoxal, and due to the high  
178 concentration of MEK in cloud water ( $70 - 650$  nmol L<sup>-1</sup>; Grosjean and Wright, 1983; van Pinxteren et al., 2005),  
179 this source is of atmospheric relevance. Due to the molar yields of 9.5% for methylglyoxal, 29.5% for 2,3-  
180 butanedione, and 3.0% for hydroxyacetone,  $\approx 42\%$  of the oxidation products of MEK could be elucidated in the  
181 present study with only these carbonyl compounds (Table 2). This highlights the importance of carbonyl  
182 compounds for the aqueous phase chemistry. Based on the experimental findings, a reaction mechanism was  
183 developed to describe the formation of methylglyoxal (Fig. 4).

### 184 3.2 Oxidation mechanism and model description

185 According to the structure of MEK, the OH radical attack can proceed at three different positions (Fig. 4; H-atoms  
186 at carbons 1, 3, and 4). For the present study, only the attack at carbons 3 and 4 is considered because these  
187 processes lead to the formation of the observed products (Fig. 4). Note that the abstraction of a hydrogen atom at  
188 carbon 3 leads to a secondary alkyl radical (A), whereas at the terminal carbon, a primary alkyl radical is formed  
189 (B). The branching ratios for the formation of the primary and secondary alkyl radicals will be discussed in detail  
190 in the next section (Sect. 3.2.1).

191 The primary and secondary alkyl radicals react rapidly with oxygen to form alkylperoxy radicals. The alkylperoxy  
192 radical recombines to a tetroxide and reacts further in three different ways, including the formation of an carbonyl  
193 compound and an alcohol (i), the formation of two carbonyl compounds and H<sub>2</sub>O<sub>2</sub> (ii), and the decomposition into  
194 an alkoxy radical (iii) (von Sonntag and Schuchmann, 1991). The decomposition of the tetroxide into a peroxide  
195 and oxygen was not considered further due to the minor importance of this process (von Sonntag and Schuchmann,  
196 1997). The secondary and primary alkylperoxy radicals can react with HO<sub>2</sub>, forming organic hydroperoxides. The  
197 organic peroxides react with OH radicals or photolyze, resulting in the formation of an alkoxy radical or tetroxide  
198 that can react further, as described before through pathways i–iii. The described mechanism was included in a  
199 COPASI model to examine the developed oxidation mechanism, the decomposition of the precursor compound,  
200 and the formation of the observed products. Table 3 shows the considered reactions, the rate constants, and their  
201 references. Only the reactions leading to the formation of the products identified are discussed in detail.  
202 Surprisingly, the products 3-oxobutanal and hydroxybutanone were not observed during the experiments. Since  
203 3-oxobutanal contains two carbonyl groups, two derivatives can be formed, including one (mass to charge ratio  
204  $m/z$  281; M<sup>+</sup>) or two ( $m/z$  476; M<sup>+</sup>) derivatized groups. Hydroxybutanone has one carbonyl group with a  
205 derivatized  $m/z$  283 (M<sup>+</sup>). As can be seen in Fig. 5,  $m/z$  283 was not found in the extracted ion chromatogram  
206 (EIC). Furthermore,  $m/z$  281 and 476 can be detected in the EIC of the bulk reactor samples, but they can also be  
207 found in the EIC of the authentic standard compounds. The sample of the authentic standard compounds does not  
208 contain hydroxybutanone and 3-oxobutanal. According to the comparison of the EIC of the bulk reactor samples  
209 and the authentic standard, it can be assumed that  $m/z$  281 and 476 are fragments of the internal standard and 2,3-  
210 butandione. No additional signals in the EIC of the samples were detected at  $m/z$  281, 283, and 476, and thus, 3-

211 oxobutanal and hydroxybutanone are not formed during the experiment. Hence, their formation pathways were  
212 excluded from the model.

### 213 3.2.1 HO<sub>x</sub> chemistry and OH radical attack

214 The relevant reactions for HO<sub>x</sub> chemistry according to R1-R6 are included in the mechanism. OH radicals were  
215 formed through the photolysis of H<sub>2</sub>O<sub>2</sub> at  $\lambda = 254$  nm with measured photolysis rate constants of  $k_p = 7.6 \times 10^{-6} \text{ s}^{-1}$   
216 (R1, see supplement S1). The formed OH radicals react further with MEK in a first oxidation step, leading to the  
217 formation of a primary (R7) and secondary alkyl radical (R8). The rate constants were postulated by  
218 Herrmann et al. (2005) to be  $k = 1.17 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$  and  $k = 1.3 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$  for the formation of the primary and  
219 secondary alkyl radicals, respectively. This is in good agreement with the model study by Sebbar et al., who  
220 postulated that the H-atom abstraction most likely proceeds at the secondary carbon of MEK (Sebbar et al., 2014;  
221 Sebbar et al., 2011) due to the lower C-H bond dissociation energy ( $\approx 377 \text{ kJ mol}^{-1}$ , C4) in comparison to the  
222 primary carbon atom ( $\approx 423 \text{ kJ mol}^{-1}$ , C3). In the present study, a rate constant  $k = 1.5 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$  (Gligorovski  
223 and Herrmann, 2004) was used, and the branching ratio was varied from 60/40 to 10/90 for the primary/secondary  
224 H-atom abstraction.

225 The results are shown in Fig. 6 and discussed based on the molar yields of the products. As can be seen  
226 for 2,3-butanedione (Fig. 6A), a branching ratio of 60% for the primary H-atom abstraction and 40% for the  
227 secondary H-atom abstraction leads to lower molar yields, whereas the molar yields start to increase with an  
228 increasing fraction of secondary H-atom abstraction. According to the mechanism (Fig. 4), 2,3-butanedione is  
229 only formed via secondary H-atom abstraction, and thus, it is feasible to reach higher molar yields with a higher  
230 fraction of secondary H-atom abstraction. However, with an increasing secondary H-atom abstraction, the  
231 experimentally determined concentration was increasingly overestimated, especially at the beginning of the  
232 experiment. After 60 minutes of reaction time, the highest experimentally determined molar yield ( $29.5 \pm 6.0\%$ )  
233 was overestimated by a factor of  $\approx 2$  with a ratio of 10/90 for primary/secondary H-atom abstraction (molar yield  
234  $\approx 49.9\%$ ), whereas a ratio of 60/40 resulted in reasonably good agreement (molar yield  $\approx 23.7\%$ ). Based on this,  
235 a branching ratio of 60/40 for primary/secondary H-atom abstraction was used in the present model.

236 In contrast, methylglyoxal molar yields were increasingly underestimated with an increasing fraction of  
237 secondary H-atom abstraction (Fig. 6B). Thus, after 15 minutes of reaction time, the experimental molar yield  
238 ( $\approx 9.5\%$ ) was underestimated by a factor of  $\approx 5$  with a higher fraction of the secondary H-atom abstraction (molar  
239 yield  $\approx 2\%$ ). In comparison with a ratio of 60/40 for primary/secondary H-atom abstraction, a molar yield of  
240  $\approx 11.4\%$  was observed, which is in good agreement with the experiment. Overall, the branching ratio of 60/40  
241 (primary/secondary H-atom abstraction) resulted in reasonable agreement for 2,3-butanedione and in an excellent  
242 conformity to the methylglyoxal molar yields. This shows the importance of the primary H-atom abstraction as  
243 the main decomposition pathway of MEK and thus for methylglyoxal formation.

244 The primary and secondary alkyl radicals react further with oxygen (R9/R21) with a rate constant of  
245  $k = 3.1 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$  (Zegota et al., 1986; Glowa et al., 2000), which was reported for the formation of  
246 acetylperoxy radicals. The rate constants of the acetylperoxy radical were used for the formation of the  
247 primary alkylperoxy radical (R9) and further reactions of the alkylperoxy radical because of their structural  
248 similarity. They were also applied by Glowa et al. (2000) for the formation of the secondary alkylperoxy radical  
249 of MEK. Further reactions of the primary and secondary alkylperoxy radicals resulting in different oxidation  
250 products will be discussed in detail in the following section (Sect. 3.2.2 and 3.2.3).

251 *3.2.2 Oxidation of MEK leading to the formation of methylglyoxal and hydroxyacetone*

252 Only the formation of the alkoxy radical (iii; R10) leads to the formation of methylglyoxal. The alkoxy radical  
253 further reacts rapidly with oxygen into an acetylperoxy radical under elimination of formaldehyde. The  
254 acetylperoxy radicals can recombine again to form a tetroxide (Schaefer et al., 2012). The latter is able to  
255 decompose through pathways i–iii, which are illustrated in R11–R13. Consequently, the decomposition of the  
256 tetroxide can explain the formation of hydroxyacetone and methylglyoxal (R11;  $k = 2 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ ; Zegota et al.,  
257 1986), methylglyoxal (R12;  $k = 4 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ ; Zegota et al., 1986), and methylglyoxal and hydrogen peroxide  
258 (R13;  $k = 4 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ ; Zegota et al., 1986; Schaefer et al., 2012).

259 In addition to the discussed pathway, the primary alkylperoxy radical has the opportunity to react with  
260  $\text{HO}_2$  instead of the recombination, forming an organic hydroperoxide (R14). The rate constant was reported by  
261 von Sonntag and Schuchmann (1991) to be from  $10^7 \text{ M}^{-1} \text{ s}^{-1}$  up to  $10^9 \text{ M}^{-1} \text{ s}^{-1}$ . For the primary alkylperoxy radical,  
262  $k = 1 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$  was used as it led to the best agreement of the experimental data and the model results. The  
263 formed peroxide can react with OH radicals (R15) or photolyze (R16). The same rate constant  
264 ( $k = 2.7 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ ) as that for the OH radical oxidation of  $\text{H}_2\text{O}_2$  was used for R15 due to a lack of literature  
265 data. According to the assumption by Monod et al. (2007), the rate constant for the photolysis of the formed  
266 peroxide (R16) was defined to be 1/10 of the photolysis rate constants of hydrogen peroxide. Thus, a photolysis  
267 rate constant of  $k_p = 7.6 \times 10^{-7} \text{ s}^{-1}$  was included in the model. This was used as a first approximation due to a lack  
268 of literature data for the formed hydroperoxides. The photolysis of the organic peroxide leads to the formation of  
269 formaldehyde, OH radicals, and an acetyl radical, which reacts further with oxygen (R17) to form an  
270 acetylperoxy radical ( $k = 3.1 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ ; Zegota et al., 1986; R19) and subsequently methylglyoxal  
271 (R18 – 20).

272 *3.2.3 Oxidation of MEK leading to the formation of methylglyoxal and 2,3-butanedione*

273 As described for the primary alkylperoxy radical, the secondary alkylperoxy radical recombines and forms a  
274 tetroxide. This reacts to form either (i) 2,3-butanedione and acetoin (R22) or (ii) 2,3-butanedione and hydrogen  
275 peroxide (R23) and is considered with rate constants of  $k = 2.5 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$  and  $k = 4.5 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$  (Glowa et  
276 al., 2000). The tetroxide can decompose into an alkoxy radical as well (iii) and react further to form 2,3-  
277 butanedione, methylglyoxal, a methyl radical, and  $\text{HO}_2$  (R24;  $k = 5 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ ; Glowa et al., 2000). The rate  
278 constants postulated by Glowa et al. (2000) are derived from the branching ratios determined by Zegota et al.  
279 (1986) for acetone. Glowa et al. (2000) simulated the concentration profiles of MEK and the corresponding  
280 products and postulated the rate coefficient used in the present model study. The secondary alkylperoxy radical  
281 also has the opportunity to react with  $\text{HO}_2$  to form an organic peroxide (R25), which photolyzes (R26;  
282  $k = 7.6 \times 10^{-7} \text{ s}^{-1}$ ) or reacts further with OH radicals (R28;  $k = 2.7 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ ). Through the photolysis, an  
283 alkoxy radical was formed that leads to methylglyoxal, 2,3-butanedione, a methyl group, and  $\text{HO}_2$  radicals  
284 (R27;  $k = 4 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ ).

285 *3.2.4 Further reactions and photolysis of formed oxidation products and MEK*

286 The products 2,3-butanedione, methylglyoxal, and hydroxyacetone positively identified by GC/MS analysis might  
287 also react further, forming a variety of oxidation products. The rate constant of methylglyoxal with OH radicals  
288 is given in the range of  $k = 5.3 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$  to  $k = 1.1 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$  (Monod et al., 2005; Ervens et al., 2003;  
289 Herrmann et al., 2005; Tan et al., 2010), whereas a rate constant of  $k = 5.3 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$  (R29; Monod et al., 2005)

290 leads to the best agreement between the experimental data and the model results. In comparison, the rate constants  
291 determined by Ervens et al. (2003), Herrmann et al. (2005), and Tan et al. (2010) resulted in an underestimation  
292 of the molar yield. **Methylglyoxal** also has the opportunity to photolyze. The photolysis is included with  
293  $k_p = 3 \times 10^{-5} \text{ s}^{-1}$  (R30) in the present model.

294 2,3-Butanedione is also prone to OH radical oxidation and photolysis. As discussed, a huge discrepancy  
295 exists in the rate constants for the reaction of OH radicals with 2,3-butanedione (Lilie et al., 1968; Doussin and  
296 Monod 2013; Gligorovski and Herrmann, 2004). In the present study, the value of  $k = 1.4 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$  (R31)  
297 determined by Gligorovski and Herrmann (2004) was used. The higher rate constants reported by Lilie et al.  
298 (1968) and Doussin and Monod (2013) resulted in consumption that was too fast. The photolysis was included  
299 with a rate constant of  $k_p = 9 \times 10^{-6} \text{ s}^{-1}$  in the COPASI model (R32). It should be mentioned that **methylglyoxal**  
300 was formed during the photolysis of 2,3-butanedione. However, the photolysis was too small to contribute  
301 significantly to **methylglyoxal** formation. For more details, see supplement S2.

302 The oxidation of hydroxyacetone with OH radicals was also considered in the model study with a rate  
303 constant of  $k = 8 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$  (R33; Stefan and Bolton, 1999). During the experiment, **methylglyoxal** was formed  
304 with 100% molar yield. Thus, the reaction of hydroxyacetone to **methylglyoxal** was included in the model study.  
305 The photolysis of hydroxyacetone was measured with a rate of  $k_p = 2 \times 10^{-5} \text{ s}^{-1}$ , leading to **methylglyoxal** with a  
306 molar yield of  $\approx 19\%$  (R34). The photolysis rate constant of MEK was measured as  $k_p = 5 \times 10^{-5} \text{ s}^{-1}$  (see  
307 supplement S2, Fig. S1). During the experiment, 2,3-butanedione was found with a molar yield of 2.2%. Thus,  
308 the reaction of MEK leading to 2,3-butanedione was included in the model study (R35).

### 309 **3.3 Model results and comparison with the experimental dataset**

310 The described reactions are included in a model, and the decomposition of MEK and molar yields of  
311 formed products were compared to the experimentally obtained data (Fig. 7). The model was not validated with  
312 the time course of hydroxyacetone due to the high standard deviation of the experimental results. The comparison  
313 of the model study and the experiment showed very good agreement for the consumption of MEK (Fig. 7A).  
314 There is also good agreement for the formation of **methylglyoxal** and limited agreement for the molar yields of  
315 2,3-butanedione. The initial high molar yield of 2,3-butanedione is reflected well (Fig. 7B). Thus, after 60 minutes  
316 of reaction time, molar yields of 23.7% in the model study and  $29.5 \pm 6.0\%$  in the experiment were reached. Under  
317 consideration of the standard deviation, this is in good agreement with the COPASI model.

318 The determined molar yields up to a reaction time of 120 minutes showed very good conformity with the  
319 experiment (18.2% and  $22.1 \pm 9.0\%$ ). The temporal behavior shows that the determined molar yield in the model  
320 study is somewhat lower towards the end of the experiment. Hence, it is possible that there are other reaction  
321 pathways that lead to higher molar yields of 2,3-butanedione at the end of the experiment, which are not  
322 considered in the model study as of now. Good agreement between the model results and experiment was observed  
323 for **methylglyoxal** (Fig. 7C). The curve shapes are very similar, and hence, fast formation and decomposition of  
324 **methylglyoxal** were found in the model study and in the experiment. After 15 minutes of reaction time, a molar  
325 yield of 9.5% was found in the experiment. In comparison, the model study resulted in a molar yield of 11.4%,  
326 which is in good agreement with the experiment and validates the COPASI model.

#### 327 **4. Atmospheric relevance**

328 The sources of methylglyoxal in the aqueous phase are thus-far not fully elucidated. Methylglyoxal can originate  
329 in the atmospheric aqueous phase through i) uptake from the gas phase, and/or ii) formation in the aqueous phase.  
330 The importance of the uptake from the gas into the aqueous phase is discussed in the literature but large  
331 discrepancies can be found. Kroll et al. (2005) investigated the uptake of methylglyoxal on inorganic seed particles  
332 under varying relative humidity. It was found that the uptake was not relevant for methylglyoxal even under high  
333 relative humidity. Contrary, Zhao et al. (2006) measured an uptake coefficient of  $\gamma = 7.6 \times 10^{-3}$  on acidic solution.  
334 They found an irreversible uptake, which decreases with increasing acidity. Fu et al. (2008, 2009) determined an  
335 uptake coefficient on aqueous particles and cloud droplets  $\gamma = 2.9 \times 10^{-3}$ , which is in good agreement with the  
336 uptake coefficient measured by Zhao et al. (2006). Lin et al. (2014) modelled an uptake coefficient for several  
337 case studies with different multiphase process mechanisms. The uptake coefficient in deliquescent particles was  
338 determined to be  $\gamma = 1.47 - 2.92 \times 10^{-5}$ . Overall,  $\gamma$  was two orders of magnitude lower compared to the values  
339 determined by Zhao et al. (2006) and Fu et al. (2008, 2009) showing the discrepancies between the different  
340 studies available in literature.

341 The in-situ formation of methylglyoxal in the aqueous phase could be an important source as well (Blando and  
342 Turpin, 2000; Sempere and Kawamura 1994). Within the present study, MEK was found as a new precursor  
343 compound for methylglyoxal in the aqueous phase yielding methylglyoxal with a molar yield of 9.5%. Although  
344 the Henry constant of MEK (up to  $K_H = 50 \text{ M atm}^{-1}$ , Schütze and Herrmann, 2004) is lower compared to  
345 methylglyoxal ( $K_H = 3.7 \times 10^3 \text{ M atm}^{-1}$ , Betterton and Hoffmann, 1988), van Pinxteren et al. (2005) found higher  
346 concentrations in cloud water as it was expected. Thus, the phase transfer of MEK from the gas in the aqueous  
347 phase could be more important as currently derived from the available Henry constants. Besides the phase transfer,  
348 the in-situ formation of MEK in aqueous phase formation might also represents an important source.

349 As, the oxidation of MEK yielding methylglyoxal has not been studied much before, it should be considered as a  
350 formation process of methylglyoxal.

351

#### 352 **5. Summary**

353 In the present study, MEK was identified as a new source for methylglyoxal in the aqueous phase. It was  
354 demonstrated that methylglyoxal originates directly from MEK oxidation and not from side reactions such as  
355 photolysis or non-radical reactions. A molar yield of  $\approx 9.5\%$  was determined during the oxidation. Based on the  
356 experimental results, a reaction mechanism could be developed. The calculations with a COPASI model supported  
357 the experimental results and confirm MEK as a precursor compound for methylglyoxal in aqueous medium.

358 Further carbonyl compounds could be identified and quantified. 2,3-Butanedione was found as the main  
359 oxidation product (molar yield  $\approx 29.5\%$ ) and was formed during the photolysis of MEK as well. As a further  
360 oxidation product, hydroxyacetone was identified and was formed with a molar yield of  $\approx 3.0\%$  during the  
361 oxidation of MEK.

362 The oxidation mechanism of MEK in aqueous solution was elucidated, and MEK was demonstrated to  
363 be as a precursor compound for methylglyoxal in the aqueous phase. Regarding the important role of  
364 methylglyoxal for the aqSOA formation, MEK has to be considered for aqSOA as well, which could be a next  
365 step in reducing the underestimation of the SOA burden by model studies.

366 **5. Acknowledgements**

367 This study was supported by the Scholarship program of the German Federal Environmental Foundation  
368 (Deutsche Bundesstiftung Umwelt, DBU; grant number 20013/244).

369 **References**

- 370 Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crounse, J. D., and Wennberg,  
371 P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmos. Chem.*  
372 *Phys.*, 11, 4039-4072, 10.5194/acp-11-4039-2011, 2011.
- 373 Andreae, M. O., and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Glob. Biogeochem.*  
374 *Cycle*, 15, 955-966, 10.1029/2000gb001382, 2001.
- 375 Ashworth, R. A., Howe, G. B., Mullins, M. E., and Rogers, T. N.: Air water partitioning coefficients of organics  
376 in dilute aqueous-solutions, *J. Hazard. Mater.*, 18, 25-36, 10.1016/0304-3894(88)85057-x, 1988.
- 377 Atkinson, R.: Gas-phase tropospheric chemistry of volatile organic compounds .I. Alkanes and alkenes, *J. Phys.*  
378 *Chem. Ref. Data*, 26, 215-290, 1997.
- 379 Blando, J. D., and Turpin B. J.: Secondary organic aerosol formation in cloud and fog droplets: a literature  
380 evaluation of plausibility, *Atmos. Environ.*, 34, 1623-1632, 2000.
- 381 Brilli, F., Gioli, B., Ciccioli, P., Zona, D., Loreto, F., Janssens, I. A., and Ceulemans, R.: Proton Transfer Reaction  
382 Time-of-Flight Mass Spectrometric (PTR-TOF-MS) determination of volatile organic compounds (VOCs)  
383 emitted from a biomass fire developed under stable nocturnal conditions, *Atmospheric Environment*, 97, 54-  
384 67, 10.1016/j.atmosenv.2014.08.007, 2014.
- 385 Brown, S. K., Sim, M. R., Abramson, M. J., and Gray, C. N.: Concentrations of volatile organic-compounds in  
386 indoor air - a review, *Indoor Air-Int. J. Indoor Air Qual. Clim.*, 4, 123-134, 10.1111/j.1600-0668.1994.t01-2-  
387 00007.x, 1994.
- 388 Buttery, R. G., Ling, L. C., and Guadagni, D. G.: Food volatiles - volatilities of aldehydes ketones and esters in  
389 dilute water solution, *J. Agric. Food Chem.*, 17, 385-&, 10.1021/jf60162a025, 1969.
- 390 Buyske, D. A., Owen, L. H., Wilder, P., and Hobbs, M. E.: Chromatography of the 2,4-dinitrophenylhydrazones  
391 of some aldehydes and ketones in tobacco smoke, *Anal. Chem.*, 28, 910-913, 10.1021/ac60113a040, 1956.
- 392 Carlier, P., Hannachi, H., and Mouvier, G.: The chemistry of carbonyl-compounds in the atmosphere - a review,  
393 *Atmospheric Environment*, 20, 2079-2099, 10.1016/0004-6981(86)90304-5, 1986.
- 394 Christensen, H., Sehested, K., and Corfitzen, H.: Reactions of hydroxyl radicals with hydrogen-peroxide at  
395 ambient and elevated-temperatures, *J. Phys. Chem.*, 86, 1588-1590, 10.1021/j100206a023, 1982.
- 396 Christensen, H., and Sehested, K.: HO<sub>2</sub> and O<sub>2</sub>- radicals at elevated-temperatures, *J. Phys. Chem.*, 92, 3007-3011,  
397 10.1021/j100321a060, 1988.
- 398 Ciccioli, P., Brancaleoni, E., Frattoni, M., Cecinato, A., and Brachetti, A.: Ubiquitous occurrence of semivolatile  
399 carbonyl-compounds in tropospheric samples and their possible sources, *Atmospheric Environment Part a-*  
400 *General Topics*, 27, 1891-1901, 10.1016/0960-1686(93)90294-9, 1993.
- 401 de Gouw, J. A., Howard, C. J., Custer, T. G., and Fall, R.: Emissions of volatile organic compounds from cut  
402 grasses and clover are enhanced during the drying process, *Geophysical Research Letters*, 26, 811-814, 1999.
- 403 Doussin, J. F., and Monod, A.: Structure-activity relationship for the estimation of OH-oxidation rate constants  
404 of carbonyl compounds in the aqueous phase, *Atmos. Chem. Phys.*, 13, 11625-11641, 10.5194/acp-13-11625-  
405 2013, 2013.
- 406 Elliot, A. J., and Buxton, G. V.: Temperature-dependence of the reactions OH + O<sub>2</sub><sup>-</sup> and OH + HO<sub>2</sub> in water up  
407 to 200 °C, *J. Chem. Soc.-Faraday Trans.*, 88, 2465-2470, 10.1039/ft9928802465, 1992.
- 408 Ervens, B., Gligorovski, S., and Herrmann, H.: Temperature-dependent rate constants for hydroxyl radical  
409 reactions with organic compounds in aqueous solutions, *Phys. Chem. Chem. Phys.*, 5, 1811-1824,  
410 10.1039/b300072a, 2003.
- 411 Ervens, B., Turpin, B. J., and Weber, R. J.: Secondary organic aerosol formation in cloud droplets and aqueous  
412 particles (aqSOA): a review of laboratory, field and model studies, *Atmos. Chem. Phys.*, 11, 11069-11102,  
413 10.5194/acp-11-11069-2011, 2011.
- 414 Faust, B. C., Powell, K., Rao, C. J., and Anastasio, C.: Aqueous-phase photolysis of biacetyl (An α-dicarbonyl  
415 compound): A sink for biacetyl, and a source of acetic acid, peroxyacetic acid, hydrogen peroxide, and the  
416 highly oxidizing acetylperoxyl radical in aqueous aerosols, fogs, and clouds, *Atmospheric Environment*, 31,  
417 497-510, 1997.
- 418 Feng, Y. L., Wen, S., Wang, X. M., Sheng, G. Y., He, Q. S., Tang, J. H., and Fu, J. M.: Indoor and outdoor  
419 carbonyl compounds in the hotel ballrooms in Guangzhou, China, *Atmospheric Environment*, 38, 103-112,  
420 10.1016/j.atmosenv.2003.09.061, 2004.



421 Fu, T-M., Jacob, D. J., Wittrock, F., Burrows, J. P., Vrekoussis, M., and Henze, D. K., Global budgets of  
422 atmospheric glyoxal and methylglyoxal, and implications for formation of secondary organic aerosols, *J.*  
423 *Geophys. Res.*, 113, 1-17, 2008.

424 Fu, T-M., Jacob, D. J., and Heald, C. L., Aqueous-phase reactive uptake of dicarbonyls as a source of organic  
425 aerosol over eastern North America, *Atmos. Environ.*, 43, 1814-1822, 2009.

426 Gligorovski, S., and Herrmann, H.: Kinetics of reactions of OH with organic carbonyl compounds in aqueous  
427 solution, *Phys. Chem. Chem. Phys.*, 6, 4118-4126, 10.1039/b403070b, 2004.

428 Glowa, G., Driver, P., and Wren, J. C.: Irradiation of MEK - II: A detailed kinetic model for the degradation under  
429 of 2-butanone in aerated aqueous solutions steady-state gamma-radiolysis conditions, *Radiat. Phys. Chem.*,  
430 58, 49-68, 10.1016/s0969-806x(99)00360-6, 2000.

431 Goldan, P. D., Kuster, W. C., Fehsenfeld, F. C., and Montzka, S. A.: Hydrocarbon measurements in the  
432 southeastern United States: The Rural Oxidants in the Southern Environment (ROSE) program 1990, *Journal*  
433 *of Geophysical Research-Atmospheres*, 100, 25945-25963, 10.1029/95jd02607, 1995.

434 Goldstein, A. H., and Galbally, I. E.: Known and unexplored organic constituents in the earth's atmosphere,  
435 *Environmental Science & Technology*, 41, 1514-1521, 10.1021/es072476p, 2007.

436 Grosjean, D.: Formaldehyde and other carbonyls in Los-Angeles ambient air, *Environmental Science &*  
437 *Technology*, 16, 254-262, 10.1021/es00099a005, 1982.

438 Grosjean, D., Swanson, R. D., and Ellis, C.: Carbonyls in Los-Angeles air - contribution of direct emissions and  
439 photochemistry, *Sci. Total Environ.*, 29, 65-85, 10.1016/0048-9697(83)90034-7, 1983.

440 Grosjean, D., and Wright, B.: Carbonyls in urban fog, ice fog, cloudwater and rainwater, *Atmospheric*  
441 *Environment*, 17, 2093-2096, 10.1016/0004-6981(83)90368-2, 1983.

442 Grosjean, D., Grosjean, E., and Moreira, L. F. R.: Speciated ambient carbonyls in Rio de Janeiro, Brazil,  
443 *Environmental Science & Technology*, 36, 1389-1395, 10.1021/es0111232, 2002.

444 Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N.  
445 M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin,  
446 M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, T. F., Monod, A., Prevot,  
447 A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of  
448 secondary organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9, 5155-5236, 2009.

449 Herrmann, H., Ervens, B., Nowacki, P., Wolke, R., and Zellner, R.: A chemical aqueous phase radical mechanism  
450 for tropospheric chemistry, *Chemosphere*, 38, 1223-1232, 10.1016/s0045-6535(98)00520-7, 1999.

451 Herrmann, H., Tilgner, A., Barzagli, P., Majdik, Z., Gligorovski, S., Poulain, L., and Monod, A.: Towards a more  
452 detailed description of tropospheric aqueous phase organic chemistry: CAPRAM 3.0, *Atmospheric*  
453 *Environment*, 39, 4351-4363, 10.1016/j.atmosenv.2005.02.016, 2005.

454 Herrmann, H., Schaefer, T., Tilgner, A., Styler, S. A., Weller, C., Teich, M., and Otto, T.: Tropospheric  
455 Aqueous-Phase Chemistry: Kinetics, Mechanisms, and Its Coupling to a Changing Gas Phase, *Chem. Rev.*,  
456 115, 4259-4334, 10.1021/cr500447k, 2015.

457 Isidorov, V. A., Zenkevich, I. G., and Ioffe, B. V.: Volatile organic-compounds in the atmosphere of forests,  
458 *Atmospheric Environment*, 19, 1-8, 10.1016/0004-6981(85)90131-3, 1985.

459 Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R.,  
460 Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat,  
461 G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic  
462 aerosol and global climate modelling: a review, *Atmos. Chem. Phys.*, 5, 1053-1123, 2005.

463 Karl, T., Chahan, Y., Jordan, A., and Lindinger, W.: Dynamic measurements of partition coefficients using proton-  
464 transfer-reaction mass spectrometry (PTR-MS), *International Journal of Mass Spectrometry*, 12269, 1-13,  
465 2002.

466 Khalil, M. A. K., and Rasmussen, R. A.: Forest hydrocarbon emissions - relationships between fluxes and ambient  
467 concentrations, *J. Air Waste Manage. Assoc.*, 42, 810-813, 1992.

468 Kirstine, W., Galbally, I., Ye, Y. R., and Hooper, M.: Emissions of volatile organic compounds (primarily  
469 oxygenated species) from pasture, *Journal of Geophysical Research-Atmospheres*, 103, 10605-10619,  
470 10.1029/97jd03753, 1998.

471 Kroll, J. H., Ng, N. L., Murphy, S. M., Varutbangkul, V., Flagan, R. C., and Seinfeld, J. H., Chamber studies of  
472 secondary organic aerosol growth by reactive uptake of simple carbonyl compounds, *J. Geophys. Res.*, 110, 1-  
473 10, 2005.

474 Lilie, J., Beck, G., and Henglein, A.: Pulse radiolysis of acetoin radicals and diacetyl anions in aqueous solution,  
475 *Berichte Der Bunsen-Gesellschaft Fur Physikalische Chemie*, 72, 529-&, 1968.

476 Lin, G., Sillman, S., Penner, J. E., and Ito, A., Global modeling of SOA: the use of different mechanisms for  
477 aqueous-phase formation, *Atmos. Chem. Phys.*, 14, 5451-5475, 2014.

478 Lipari, F., Dasch, J. M., and Scruggs, W. F.: Aldehyde emissions from wood-burning fireplaces, *Environmental*  
479 *Science & Technology*, 18, 326-330, 10.1021/es00123a007, 1984.

480 Matthews, T. G., and Howell, T. C.: Visual colorimetric formaldehyde screening analysis for indoor air, *Journal*  
481 *of the Air Pollution Control Association*, 31, 1181-1184, 1981.

482 Monod, A., Poulain, L., Grubert, S., Voisin, D., and Wortham, H.: Kinetics of OH-initiated oxidation of  
483 oxygenated organic compounds in the aqueous phase: new rate constants, structure-activity relationships and  
484 atmospheric implications, *Atmospheric Environment*, 39, 7667-7688, 10.1016/j.atmosenv.2005.03.019, 2005.

485 Monod, A., Chevallier, E., Jolibois, R. D., Doussin, J. F., Picquet-Varrault, B., and Carlier, P.: Photooxidation of  
486 methylhydroperoxide and ethylhydroperoxide in the aqueous phase under simulated cloud droplet conditions,  
487 *Atmospheric Environment*, 41, 2412-2426, 10.1016/j.atmosenv.2006.10.006, 2007.

488 Mopper, K., and Stahovec, W. L.: Sources and sinks of low-molecular-weight organic carbonyl-compounds in  
489 seawater, *Mar. Chem.*, 19, 305-321, 10.1016/0304-4203(86)90052-6, 1986.

490 Morillon, V., Debeaufort, F., Jose, J., Tharrault, J. F., Capelle, M., Blond, G., and Voilley, A.: Water vapour  
491 pressure above saturated salt solutions at low temperatures, *Fluid Phase Equilib.*, 155, 297-309,  
492 10.1016/s0378-3812(99)00009-6, 1999.

493 Müller, K., van Pinxteren, D., Plewka, A., Svrčina, B., Kramberger, H., Hofmann, D., Bachmann, K., and  
494 Herrmann, H.: Aerosol characterisation at the FEBUKO upwind station Goldlauter (II): Detailed organic  
495 chemical characterisation, *Atmospheric Environment*, 39, 4219-4231, 10.1016/j.atmosenv.2005.02.008, 2005.

496 Pastina, B., and LaVerne, J. A.: Effect of molecular hydrogen on hydrogen peroxide in water radiolysis, *J. Phys.*  
497 *Chem. A*, 105, 9316-9322, 10.1021/jp012245j, 2001.

498 Riemer, D., Pos, W., Milne, P., Farmer, C., Zika, R., Apel, E., Olszyna, K., Kliendienst, T., Lonneman, W.,  
499 Bertman, S., Shepson, P., and Starn, T.: Observations of nonmethane hydrocarbons and oxygenated volatile  
500 organic compounds at a rural site in the southeastern United States, *Journal of Geophysical Research-*  
501 *Atmospheres*, 103, 28111-28128, 10.1029/98jd02677, 1998.

502 Rodigast, M., Mutzel, A., Iinuma, Y., Haferkorn, S., and Herrmann, H.: Characterisation and optimisation of a  
503 sample preparation method for the detection and quantification of atmospherically relevant carbonyl  
504 compounds in aqueous medium, *Atmos. Meas. Tech.*, 8, 2409-2416, 10.5194/amt-8-2409-2015, 2015.

505 Schaefer, T., Schindelka, J., Hoffmann, D., and Herrmann, H.: Laboratory Kinetic and Mechanistic Studies on  
506 the OH-Initiated Oxidation of Acetone in Aqueous Solution, *J. Phys. Chem. A*, 116, 6317-6326,  
507 10.1021/jp2120753, 2012.

508 Schütze, M., and Herrmann, H.: Uptake of acetone, 2-butanone, 2,3-butanedione and 2-oxopropanal on a water  
509 surface, *Phys. Chem. Chem. Phys.*, 6, 965-971, 10.1039/b313474a, 2004.

510 Sebbar, N., Bozzelli, J. W., and Bockhorn, H.: Thermochemistry and Kinetics for 2-Butanone-1-yl Radical (CH<sub>2</sub>  
511 center dot C(=O)CH<sub>2</sub>CH<sub>3</sub>) Reactions with O<sub>2</sub>, *J. Phys. Chem. A*, 118, 21-37, 10.1021/jp408708u, 2014.

512 Sebbar, N., Bozzelli, J. W., and Bockhorn, H.: Reactivity, thermochemistry and kinetics of 2-butanone radicals:  
513 CH<sub>2</sub>C(=O)CHCH<sub>3</sub>, CH<sub>3</sub>C(=O)CHCH<sub>3</sub> and CH<sub>3</sub>C(=O)CH<sub>2</sub>CH<sub>2</sub>, 2011.

514 [Sempere, R., and Kawamura K.: Comparative distribution of dicarboxylic acids and related polar compounds in](#)  
515 [snow, rain and aerosols from urban, \*Atmos. Environ.\*, 28, 449-459, 1994.](#)

516 Singh, H. B., Salas, L. J., Chatfield, R. B., Czech, E., Fried, A., Walega, J., Evans, M. J., Field, B. D., Jacob, D.  
517 J., Blake, D., Heikes, B., Talbot, R., Sachse, G., Crawford, J. H., Avery, M. A., Sandholm, S., and Fuelberg,  
518 H.: Analysis of the atmospheric distribution, sources, and sinks of oxygenated volatile organic chemicals based  
519 on measurements over the Pacific during TRACE-P, *Journal of Geophysical Research-Atmospheres*, 109, 20,  
520 10.1029/2003jd003883, 2004.

521 Snider, J. R., and Dawson, G. A.: Tropospheric light alcohols, carbonyls and acetonitrile - concentrations in  
522 southwestern united-states and henry law data, *Journal of Geophysical Research-Atmospheres*, 90, 3797-3805,  
523 10.1029/JD090iD02p03797, 1985.

524 Stefan, M. I., and Bolton, J. R.: Reinvestigation of the acetone degradation mechanism in dilute aqueous solution  
525 by the UV/H<sub>2</sub>O<sub>2</sub> process, *Environmental Science & Technology*, 33, 870-873, 10.1021/es9808548, 1999.

526 Tan, Y., Carlton, A. G., Seitzinger, S. P., and Turpin, B. J.: SOA from methylglyoxal in clouds and wet aerosols:  
527 Measurement and prediction of key products, *Atmospheric Environment*, 44, 5218-5226,  
528 10.1016/j.atmosenv.2010.08.045, 2010.

529 van Pinxteren, D., Plewka, A., Hofmann, D., Müller, K., Kramberger, H., Svrčina, B., Bachmann, K., Jaeschke,  
530 W., Mertes, S., Collett, J. L., and Herrmann, H.: Schmucke hill cap cloud and valley stations aerosol  
531 characterisation during FEBUKO (II): Organic compounds, *Atmospheric Environment*, 39, 4305-4320,  
532 10.1016/j.atmosenv.2005.02.014, 2005.

533 von Sonntag, C., and Schuchmann, H. P.: Aufklärung von Peroxyl-Radikalreaktionen in wäßriger Lösung mit  
534 strahlenchemischen Techniken, *Angew. Chem.*, 130, 1255-1279, 1991.

535 von Sonntag, C., and Schuchmann, H. P.: Bimolecular Decay of peroxy radicals, in: *Peroxy radicals*, edited by:  
536 Alfassi, Z. B., John Wiley & sons, 199-207, 1997.

537 Ware, G. W.: Methyl ethyl ketone, *Reviews of Environmental Contamination and Toxicology*, 106, 165-174,  
538 1988.

539 Warneke, C., Karl, T., Judmaier, H., Hansel, A., Jordan, A., Lindinger, W., and Crutzen, P. J.: Acetone, methanol,  
540 and other partially oxidized volatile organic emissions from dead plant matter by abiological processes:  
541 Significance for atmospheric HO<sub>x</sub> chemistry, *Glob. Biogeochem. Cycle*, 13, 9-17, 10.1029/98gb02428, 1999.  
542 Yokelson, R. J., Burling, I. R., Gilman, J. B., Warneke, C., Stockwell, C. E., de Gouw, J., Akagi, S. K., Urbanski,  
543 S. P., Veres, P., Roberts, J. M., Kuster, W. C., Reardon, J., Griffith, D. W. T., Johnson, T. J., Hosseini, S.,  
544 Miller, J. W., Cocker, D. R., Jung, H., and Weise, D. R.: Coupling field and laboratory measurements to  
545 estimate the emission factors of identified and unidentified trace gases for prescribed fires, *Atmos. Chem.*  
546 *Phys.*, 13, 89-116, 10.5194/acp-13-89-2013, 2013.  
547 Zegota, H., Schuchmann, M. N., Schulz, D., and Vonsonntag, C.: Acetylperoxy radicals, CH<sub>3</sub>COCH<sub>2</sub>O<sub>2</sub> - a  
548 study on the gamma-radiolysis and pulse-radiolysis of acetone in oxygenated aqueous-solutions,  
549 *Z.Naturforsch.(B)*, 41, 1015-1022, 1986.  
550 [Zhao, J., Levitt, N. P., Zhang, R., and Chen, J., Heterogeneous Reactions of Methylglyoxal in Acidic Media:  
551 Implications for Secondary Organic Aerosol Formation, \*Environ. Sci. Technol.\*, 40, 7682-7687, 2006.](#)  
552 Zhou, X. L., and Mopper, K.: Apparent partition-coefficients of 15 carbonyl-compounds between air and seawater  
553 and between air and fresh-water - implications for air sea exchange, *Environmental Science & Technology*,  
554 24, 1864-1869, 10.1021/es00082a013, 1990.  
555 Zhou, X. L., and Mopper, K.: Photochemical production of low-molecular-weight carbonyl compounds in  
556 seawater and surface microlayer and their air-sea exchange, *Mar. Chem.*, 56, 201-213, 10.1016/s0304-  
557 4203(96)00076-x, 1997.

558

Table 1: Conducted experiments in the bulk reactor.

Number of experiment	Type of experiment	Concentration precursor compound [mmol L <sup>-1</sup> ]	Concentration H <sub>2</sub> O <sub>2</sub> [mmol L <sup>-1</sup> ]	UV light (λ = 254 nm)	Reaction time [hours]	Number of repetition
1	Oxidation of MEK	0.1	2	✓	4	3
2	Reaction of H <sub>2</sub> O <sub>2</sub> with MEK	0.1	2	-	4	1
3	Photolysis of MEK	0.1	-	✓	4	1
4	Photolysis of H <sub>2</sub> O <sub>2</sub>	-	2	✓	4	1
5	Oxidation of 2,3-butanedione	0.1	2	✓	4	1
6	Oxidation of hydroxyacetone	0.1	2	✓	4	1
7	Photolysis of 2,3-butanedione, hydroxyacetone, methylglyoxal	0.1	-	✓	4	1

MEK: Methyl ethyl ketone




Figure 1: Consumption of MEK (A) during the oxidation with OH radicals and time-resolved formation of the products methylglyoxal (B), 2,3-butanedione (B), and hydroxyacetone (B).




Figure 2: GC/MS chromatogram of oxidation of 2,3-butanedione (a,b,c) and the authentic standard compounds 2,3-butanedione (a,b,c) and methylglyoxal (d,e).




Figure 3: Photolysis of MEK and time-resolved formation of 2,3-butanedione.

Table 2: Maximal molar yields of the oxidation products.

Oxidation product	Maximal molar yield (± standard deviation) in %	Reaction time in minutes
2,3-Butanedione	29.5 ± 6.0	60
Methylglyoxal	9.5	15
Hydroxyacetone	3.0 ± 2.6	60






Figure 4: Recommendation of an oxidation mechanism of MEK for the formation of 2,3-butanedione, hydroxyacetone and methylglyoxal.

2 Table 3: Reaction mechanism and rate constants for the modelling of the experiment with COPASI.

	Reaction	Rate constant $k$	Comment
R1	$\text{H}_2\text{O}_2 + h\nu \rightarrow 2 \text{OH}$	$7.6 \times 10^{-6} \text{ s}^{-1}$	measured
R2	$2 \text{OH} \rightarrow \text{H}_2\text{O}_2$	$3.6 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$	Elliot and Buxton, 1992
R3	$\text{OH} + \text{H}_2\text{O}_2 \rightarrow \text{HO}_2 + \text{H}_2\text{O}$	$2.7 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$	Christensen et al., 1982
R4	$\text{OH} + \text{HO}_2 \rightarrow \text{H}_2\text{O} + \text{O}_2$	$6 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$	Elliot and Buxton, 1992
R5	$2 \text{HO}_2 \rightarrow \text{H}_2\text{O}_2 + \text{O}_2$	$9.8 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$	Christensen and Sehested, 1988
R6	$\text{HO}_2 + \text{H}_2\text{O}_2 \rightarrow \text{OH} + \text{O}_2 + \text{H}_2\text{O}$	$0.5 \text{ M}^{-1} \text{ s}^{-1}$	Pastina and LaVerne, 2001
R7	$\text{CH}_3\text{C}(\text{O})\text{CH}_2\text{CH}_3 + \text{OH} \rightarrow \text{CH}_3\text{C}(\text{O})\text{CH}_2\text{CH}_2 + \text{H}_2\text{O}$	$9 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$	changed after Gligorovski and Herrmann, 2004
R8	$\text{CH}_3\text{C}(\text{O})\text{CH}_2\text{CH}_3 + \text{OH} \rightarrow \text{CH}_3\text{C}(\text{O})\text{CHCH}_3 + \text{H}_2\text{O}$	$6 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$	changed after Gligorovski and Herrmann, 2004
R9	$\text{CH}_3\text{C}(\text{O})\text{CH}_2\text{CH}_2 + \text{O}_2 \rightarrow \text{CH}_3\text{C}(\text{O})\text{CH}_2\text{CH}_2\text{O}_2$	$3.1 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$	Zegota et al., 1986
R10	$2 \text{CH}_3\text{C}(\text{O})\text{CH}_2\text{CH}_2\text{O}_2 + \text{O}_2 \rightarrow \text{CH}_3\text{C}(\text{O})\text{CH}_2\text{O}_2 + 2 \text{HCHO}$	$4 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$	Zegota et al., 1986
R11	$2 \text{CH}_3\text{C}(\text{O})\text{CH}_2\text{O}_2 \rightarrow \text{CH}_3\text{C}(\text{O})\text{CH}_2\text{OH} + \text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{H} + \text{O}_2$	$2 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$	Zegota et al., 1986
R12	$2 \text{CH}_3\text{C}(\text{O})\text{CH}_2\text{O}_2 \rightarrow 2 \text{HO}_2 + 2 \text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{H}$	$4 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$	Zegota et al., 1986
R13	$2 \text{CH}_3\text{C}(\text{O})\text{CH}_2\text{O}_2 \rightarrow 2 \text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{H} + \text{H}_2\text{O}_2$	$4 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$	Zegota et al., 1986
R14	$\text{CH}_3\text{C}(\text{O})\text{CH}_2\text{CH}_2\text{O}_2 + \text{HO}_2 \rightarrow \text{CH}_3\text{C}(\text{O})\text{CH}_2\text{CH}_2\text{OOH} + \text{O}_2$	$1 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$	Von Sonntag and Schuchmann, 1991
R15	$\text{CH}_3\text{C}(\text{O})\text{CH}_2\text{CH}_2\text{OOH} + \text{OH} \rightarrow \text{Product}$	$2.7 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$	Christensen et al., 1982
R16	$\text{CH}_3\text{C}(\text{O})\text{CH}_2\text{CH}_2\text{OOH} \rightarrow \text{OH} + \text{HCHO} + \text{CH}_3\text{C}(\text{O})\text{CH}_2$	$7.6 \times 10^{-7} \text{ s}^{-1}$	after Monod et al., 2007
R17	$\text{CH}_3\text{C}(\text{O})\text{CH}_2 + \text{O}_2 \rightarrow \text{CH}_3\text{C}(\text{O})\text{CH}_2\text{O}_2$	$3.1 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$	Zegota et al., 1986
R18	$2 \text{CH}_3\text{C}(\text{O})\text{CH}_2\text{O}_2 \rightarrow \text{CH}_3\text{C}(\text{O})\text{CH}_2\text{OH} + \text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{H} + \text{O}_2$	$2 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$	Zegota et al., 1986
R19	$2 \text{CH}_3\text{C}(\text{O})\text{CH}_2\text{O}_2 \rightarrow \text{H}_2\text{O}_2 + 2 \text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{H}$	$4 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$	Zegota et al., 1986
R20	$2 \text{CH}_3\text{C}(\text{O})\text{CH}_2\text{O}_2 \rightarrow 2 \text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{H} + \text{HO}_2$	$4 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$	Zegota et al., 1986
R21	$\text{CH}_3\text{C}(\text{O})\text{CHCH}_3 + \text{O}_2 \rightarrow \text{CH}_3\text{C}(\text{O})\text{CHO}_2\text{CH}_3$	$3.1 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$	Glowa et al., 2000
R22	$2 \text{CH}_3\text{C}(\text{O})\text{CHO}_2\text{CH}_3 \rightarrow \text{O}_2 + \text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{CH}_3 + \text{CH}_3\text{C}(\text{O})\text{CHCH}_3\text{OH}$	$2.5 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$	Glowa et al., 2000
R23	$2 \text{CH}_3\text{C}(\text{O})\text{CHO}_2\text{CH}_3 \rightarrow \text{H}_2\text{O}_2 + 2 \text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{CH}_3$	$4.5 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$	Glowa et al., 2000
R24	$2 \text{CH}_3\text{C}(\text{O})\text{CHO}_2\text{CH}_3 \rightarrow \text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{CH}_3 + \text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{H} + \text{CH}_3 + \text{HO}_2$	$5 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$	Glowa et al., 2000
R25	$\text{CH}_3\text{C}(\text{O})\text{CHO}_2\text{CH}_3 + \text{HO}_2 \rightarrow \text{CH}_3\text{C}(\text{O})\text{CHOOHCH}_3 + \text{O}_2$	$1 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$	Von Sonntag and Schuchmann, 1991
R26	$\text{CH}_3\text{C}(\text{O})\text{CHOOHCH}_3 \rightarrow \text{CH}_3\text{C}(\text{O})\text{CHOCH}_3 + \text{OH}$	$7.6 \times 10^{-7} \text{ s}^{-1}$	after Monod et al., 2007
R27	$2 \text{CH}_3\text{C}(\text{O})\text{CHOCH}_3 \rightarrow \text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{H} + \text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{CH}_3 + \text{CH}_3 + \text{HO}_2$	$4 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$	Zegota et al., 1986
R28	$\text{CH}_3\text{C}(\text{O})\text{CHOOHCH}_3 + \text{OH} \rightarrow \text{Product}$	$2.7 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$	Christensen et al., 1982
R29	$\text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{H} + \text{OH} \rightarrow \text{Product}$	$5.3 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$	Monod et al., 2005
R30	$\text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{H} \rightarrow \text{Product}$	$3 \times 10^{-5} \text{ s}^{-1}$	measured
R31	$\text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{CH}_3 + \text{OH} \rightarrow \text{Product}$	$1.4 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$	Gligorovski and Herrmann, 2004
R32	$\text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{CH}_3 \rightarrow \text{Product} + 0.17 \text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{H}$	$9 \times 10^{-6} \text{ s}^{-1}$	measured
R33	$\text{CH}_3\text{C}(\text{O})\text{CH}_2\text{OH} + \text{OH} \rightarrow \text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{H}$	$8 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$	Stefan and Bolton, 1999
R34	$\text{CH}_3\text{C}(\text{O})\text{CH}_2\text{OH} \rightarrow \text{Product} + 0.19 \text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{H}$	$2 \times 10^{-5} \text{ s}^{-1}$	measured
R35	$\text{CH}_3\text{C}(\text{O})\text{CH}_2\text{CH}_3 \rightarrow \text{Product} + 0.02 \text{CH}_3\text{C}(\text{O})\text{C}(\text{O})\text{CH}_3$	$5 \times 10^{-5} \text{ s}^{-1}$	measured




3  
4 Figure 5: Extracted ion chromatogram (EIC) of  $m/z$  281  $M^+$ ,  $m/z$  476  $M^+$  and  $m/z$  283  $M^+$  during the oxidation  
5 of MEK and EIC of the authentic standard compounds.



6  
7 Figure 6: Comparison of the molar yields of 2,3-butanedione (A) and methylglyoxal (B) for the model and  
8 experimental results using different branching ratios of the primary/secondary H-atom abstraction varied between  
9 60/40 up to 10/90.





10

11 Figure 7: Comparison of the model and experimental results for MEK (A), 2,3-butanedione (B), and

12 methylglyoxal (C).

## 13 Supplement

### 14 **S 1 Determination of photon flux of Xe/Hg lamp and the photolysis rate constants of H<sub>2</sub>O<sub>2</sub>**

15 The photon flux of the 500-W Xe/Hg lamp used in the experimental setup was determined with ferrioxalate  
16 actinometry (Hatchard and Parker, 1956). For this method, two types of solutions were prepared. Solution 1  
17 contained 0.006 mol L<sup>-1</sup> of K<sub>3</sub>Fe(C<sub>2</sub>O<sub>4</sub>)<sub>3</sub> in 0.05 mol L<sup>-1</sup> of H<sub>2</sub>SO<sub>4</sub> and was made under red light in the absence of  
18 oxygen to avoid Fe<sup>2+</sup> formation. Solution 2 contained 7.5 mL of phenanthroline (5 × 10<sup>-3</sup> mol L<sup>-1</sup>) and 1 mL of  
19 acetate solution. The acetate solution was made of 0.6 mol L<sup>-1</sup> of sodium acetate trihydrate and 0.17 mol L<sup>-1</sup> of  
20 H<sub>2</sub>SO<sub>4</sub> in water. To determine the photon flux, 300 mL of solution 1 was filled in the bulk reactor and illuminated  
21 (λ = 254 nm). In steps of 5 minutes, 2 mL of the solution were collected from the bulk reactor and mixed with  
22 solution 2. This mixture was allowed to react for 30 minutes. The absorption was measured with UV/Vis  
23 spectroscopy at λ = 510 nm. Based on the absorption, the photon flux was calculated with q = 4.94 × 10<sup>-9</sup> mol s<sup>-1</sup>  
24 (q = 2.96 molecules s<sup>-1</sup>). Using the photon flux, the decomposition rate of H<sub>2</sub>O<sub>2</sub> was calculated according to  
25 Eq. (S1):

$$26 \quad \frac{d[H_2O_2]}{dt} = \frac{q \times \theta}{V \times N_A} \times (1 - 10^{\varepsilon \times c \times d}) \quad \text{Eq. (1)}$$

27	q	Photon flux [molecules s <sup>-1</sup> ]
28	θ	Quantum yield H <sub>2</sub> O <sub>2</sub> (Kwon and Kwon, 2010)
29	V	Reaction volume [L]
30	N <sub>A</sub>	Avogadro constant [molecules mol <sup>-1</sup> ]
31	ε	Extinction coefficient H <sub>2</sub> O <sub>2</sub> [L mol <sup>-1</sup> cm <sup>-1</sup> ]
32	c	Concentration H <sub>2</sub> O <sub>2</sub> [mol L <sup>-1</sup> ]
33	d	Optical path length [cm]

34 Based on the decay of H<sub>2</sub>O<sub>2</sub>, a photolysis rate constant of k<sub>pH<sub>2</sub>O<sub>2</sub></sub> = 7.6 × 10<sup>-6</sup> s<sup>-1</sup> was calculated according to  
35 Eq. (S2):

$$36 \quad k_{pH_2O_2} = \frac{\frac{d[H_2O_2]}{dt}}{c} \quad \text{Eq. (2)}$$

## S 2 Photolysis rate constants of MEK, 2,3-butanedione, hydroxyacetone and methylglyoxal

Photolysis rate constants were determined using the decay of the respective precursor compound (MEK, 2,3-butanedione, hydroxyacetone, and methylglyoxal). The photolysis rate constants were determined by plotting the logarithmic concentration of the precursor compound against the reaction time in seconds. According to the linear regression ( $y = mx + n$ ), the photolysis rate constants correspond to the slope of the linear fit ( $m$ ). The results are illustrated in Fig. S1.




Figure S1: Photolysis rate constants of MEK, methylglyoxal, hydroxyacetone, and 2,3-butanedione determined through the slope  $m$  of the linear fit.

For MEK, 2,3-butanedione, methylglyoxal, and hydroxyacetone, the photolysis rate constants were determined with  $k_{p\text{MEK}} = 5 \times 10^{-5} \text{ s}^{-1}$ ,  $k_{p\text{Methylglyoxal}} = 3 \times 10^{-5} \text{ s}^{-1}$ ,  $k_{p\text{Hydroxyacetone}} = 2 \times 10^{-5} \text{ s}^{-1}$ , and  $k_{p2,3\text{-Butanedione}} = 9 \times 10^{-6} \text{ s}^{-1}$ .

During the photolysis of 2,3-butanedione and hydroxyacetone, methylglyoxal was formed with molar yields of  $\approx 17.0\%$  and  $19.5\%$  (Fig. S2).




Figure S2: Formation of methylglyoxal due to the photolysis of hydroxyacetone and 2,3-butanedione.

These two additional methylglyoxal sources were included in the model as well.

### S 3 Oxidation of hydroxyacetone

The oxidation of hydroxyacetone was investigated to determine its contribution to the formation of methylglyoxal during the oxidation of MEK (Fig. S3). The contribution of the oxidation of 2,3-butanedione to the methylglyoxal formation is discussed in the manuscript.




Figure S3: Oxidation of hydroxyacetone and formation of methylglyoxal.

During the oxidation, hydroxyacetone was completely consumed after a reaction time of 240 minutes. After 60 minutes, methylglyoxal was formed with a molar yield of  $\approx 100\%$ . Afterwards, the concentration of methylglyoxal starts to decrease and resulted in complete consumption at the end of the experiment. It can be concluded that hydroxyacetone is an important source of methylglyoxal. Notably, hydroxyacetone was formed in the present study with a molar yield of  $\approx 3.0\%$  ( $1.9 \mu\text{mol L}^{-1}$ ). Such a low concentration cannot explain the huge amount of methylglyoxal formed from the oxidation of MEK. Based on this, it can be stated that the oxidation of hydroxyacetone has only a small contribution to the methylglyoxal formation.

## References

Hatchard, C. G., and Parker, C. A.: A new sensitive chemical actinometer .2. potassium ferrioxalate as standard chemical actinometer, *Proceedings of the Royal Society of London Series a-Mathematical and Physical Sciences*, 235, 518-536, 10.1098/rspa.1956.0102, 1956.

Kwon, B. G., and Kwon, J. H.: Measurement of the hydroxyl radical formation from H<sub>2</sub>O<sub>2</sub>, NO<sub>3</sub><sup>-</sup>, and Fe(III) using a continuous flow injection analysis, *J. Ind. Eng. Chem.*, 16, 193-199, 10.1016/j.jiec.2009.10.007, 2010.