

Answer to Anonymous Referee 1 Received and published: 10 December 2015

The paper is well written. It needs only minor technical corrections. p2. L25-35: Refer to more papers on BB in Europe, adding more geographical areas and more recent papers.

Three other papers: Sciare et al. (2008), Bougiatioti et al. (2013), Edler et al. (2011) are now
5 referenced in the new version to extend the biomass burning (BB) characterization to the Eastern
Mediterranean basin and to emphasize the use of chemical characterization of the aerosol to identify
BB episode.

*p2. L37, p. 5. L132, etc.: In all manuscript (text, fig, tables) refer ONLY to the well-established
lidar ratio $[(LR)=\alpha(aer)/\beta(aer)]$, and not to BER, as defined in French manuscripts.*

10 We agree. This has been corrected.

*p3. In section 2.2, the authors should mention the uncertainties of the parameters retrieved by
each instrument (lidar, MODIS etc) and cite the corresponding papers. In addition, how the depo-
larization ratio is calibrated and retrieved.*

We thank the reviewer for this useful comment. We fully agree that references to the uncertainties
15 were sometimes missing, especially for the depolarization ratio. Three sentences have been added
in section 2.2 and 3.1 to describe how the depolarization calibration is performed and the related
error calculated for airborne and ground based lidar (p. 4 L 105 to 111) and CALIOP (p. 6 L 151-
153.). The error bar is given for CALIOP data presented in section 3 and 4 by using the error of the
backscatter signal from the CALIOP data products. The error on the airborne and ground based lidar
20 depolarization ratio is also estimated assuming a 20% uncertainty on the calibration by the molecular
signal. Therefore the error values on the depolarization ratios are now provided in section 3 and 4
(for example see Table 3 for the ground based lidar, p7 L 184 to 187 for CALIOP and p.19 L 323
for the airborne lidar). A reference for the error on the MODIS AOD daily product is also added on
p.5 L 125.

25 *p6. L165: change from "June 17 to June 25" to the correct "17 to 25 June 2013" Also, in all
manuscript put the dates in European format "dd mm yyyy" (e.g see p.16, Fig. 9, legend)*

done

*p.7, legend Fig. 2.: Again, do not duplicate the month when you refer to the same month period.
The correct wording should be "17 to 25 June 2013"*

30 done

*p.9, Fig.4. legend: again, in all legends add the year as well. p.10-Fig.5, p12-Fig. 6, p13-Fig.7.,
p14-Fig.8. p21-Fig11, p22-Fig.12, : legend: again, in all legends add the year as well*

Year is now always mentioned in the figure captions.

p17., L293, replace "J.Pelon and co-workers "by Pelon et al. (to be submitted)".

35 We kept the initial wording to avoid listing a non-published paper in the reference list. The name
of the co-worker may also change.

p24., L383, replace "12% while" by "12%, while".

done

*Also, in the References add full information about the forthcoming paper, as follows: "Pelon J.,
40, title, Journal (to be submitted)*

See answer to the p17., L293 comment.

Anonymous Referee 2 Received and published: 18 December 2015

*In the presented work gives an extensive examination of the aerosol situation in summer 2013 in
the Western Mediterranean area. The paper is well written and interesting to read, and I have only
45 minor comments which should be considered before publication in ACP.*

*Introduction: You define BER as extinction to backscatter ratio. In Section 3.1 you define the lidar
ratio (LR) as BER-1. To my knowledge the general definition of the LR is extinction-to-backscatter
ratio.*

Yes the reviewer is right. It was a mistake we meant backscatter to extinction ratio. The word BER
50 has now been removed from the paper as the most popular parameter is the lidar ratio.

Section 2.1: Please give references for this information.

done

Section 3.1: Can you give a reference for the forward Klett inversion? Please give more information on the exponent k (order, differences depending on aerosol type).

55 In fact for a forward inversion scheme it is known as the Fernald forward inversion. The name has been corrected and a reference to Fernald (1984) is added p. 6 L 147. The range for the k value goes from 0 to 2 according to the contribution of the fine mode. This has been added p. 6 L 158.

Section 3.3: What do you mean by 'Atlantic dust sources'?

60 Yes we agree the word is misleading. There is of course no dust source over the Atlantic. It has been replaced by the dust aerosol layers over the Atlantic ocean used for the initialization of the FLEXPART Lagrangian model.

Figure 1: Please give the time of the CALIOP tracks and the FALCON 20 flight in the figure caption.

65 The legend in Figure 1 has been changed and date and time of the aircraft flight is given in the caption.

Figure 2: Panels are not top and bottom. Would it be possible to use the same scale for both panels?

Done.

70 *Figure 3: How do you explain the very inhomogeneous structure in Depol and CR (especially on 21 June) which seems not really to correspond with the layering shown in R532-plot? Also on 22 June the intensive optical properties seem to be very variable within the aerosol plume. I would not expect such large differences.*

75 The CALIOP figures (3 and 7) have been reprocessed to remove the noisy data in the depolarization and color ratio vertical cross sections. Data are not provided for backscatter ratio less than 2 because the ratio of two noisy values is not reliable. Some inhomogeneity remains because the aerosol mixing and composition change near the boundaries of the layers. Data provided in the paper are averages over the aerosol layers identified in the backscatter ratio vertical cross sections.

Figure 8: Please indicate the source of these plots (FLEXPART?) in the figure caption.

Done

80 **List of author's changes in the manuscript**

See attached pdf to identify in red the modifications of the text.

Although it was not requested by the reviewers, a new table (Table 4) is added in section 4.3 to summarize the aerosol optical properties of the 4 different aerosol types described in the paper.

Fig. 1, 2, 3 and 7 have been updated. Error bars are given for depolarization ratios in Table 3.

Manuscript prepared for Atmos. Chem. Phys.

with version 2014/09/16 7.15 Copernicus papers of the L^AT_EX class copernicus.cls.

Date: 21 January 2016

Long range transport and mixing of aerosol sources during the 2013 North American biomass burning episode: analysis of multiple lidar observations in the Western Mediterranean basin

Gerard Ancellet¹, Jacques Pelon¹, Julien Totems², Patrick Chazette², Ariane Bazureau¹, Michaël Sicard³, Tatiana Di Iorio⁴, Francois Dulac², and Marc Mallet⁵

¹Sorbonne Université, UPMC; Université Versailles St-Quentin; CNRS/INSU; LATMOS, Paris, France

²LSCE, Laboratoire des sciences du Climat et de l'Environnement; CEA; Université Versailles St-Quentin; CNRS/INSU; Gif sur Yvette, France

³RSLab/CTE-CRAE-IEEC, Universitat Politècnica de Catalunya, Barcelona, Spain

⁴ENEA, Agenzia nazionale per le nuove tecnologie, l'energia e lo sviluppo economico sostenibile, Roma, Italy

⁵Laboratoire d'Aérodynamique, Université Paul Sabatier; CNRS/INSU, Toulouse, France

Correspondence to: Gerard Ancellet: gerard.ancellet@upmc.fr

85 **Abstract.** Long range transport of biomass burning (BB) aerosols between North America and the Mediterranean region took place in June 2013. A large number of ground based and airborne lidar measurements were deployed in the Western Mediterranean during the Chemistry-AeRosol Mediterranean EXperiment (ChArMEX) intensive observation period. A detailed analysis of the potential North American aerosol sources is conducted including the assessment of their transport to Europe
90 using forward simulations of the FLEXPART Lagrangian particle dispersion model initialized using satellite observations by MODIS and CALIOP. The three dimensional structure of the aerosol distribution in the ChArMEX domain observed by the ground-based lidars (Menorca, Barcelona and Lampedusa), a Falcon-20 aircraft flight and three CALIOP tracks, agree very well with the model simulation of the three major sources considered in this work: Canadian and Colorado fires, a dust storm from Western US and the contribution of Saharan dust streamers advected from the North
95 Atlantic trade wind region into the Westerlies region. Four aerosol types were identified using the optical properties of the observed aerosol layers (aerosol depolarization ratio, lidar ratio) and the transport model analysis of the contribution of each aerosol source: (I) pure BB layer, (II) weakly dusty BB, (III) significant mixture of BB and dust transported from the trade wind region (IV) the
100 outflow of Saharan dust by the subtropical jet and not mixed with BB aerosol. The contribution of the Canadian fires is the major aerosol source during this episode while mixing of dust and BB is only significant at altitude above 5 km. The mixing corresponds to a 20%-30% dust contribution in the total aerosol backscatter. The comparison with the MODIS AOD horizontal distribution during this episode over the Western Mediterranean sea shows that the Canadian fires contribution were as
105 large as the direct northward dust outflow from Sahara.

1 Introduction

Forest fires are a significant source of tropospheric aerosol particles at northern latitudes in Spring and Summer (Generoso et al., 2003; Warneke et al., 2009) and many studies project higher temperatures and longer growing season e.g. (Flannigan et al., 2009; Liu et al., 2014). The focus of biomass burning emission impact on the atmospheric composition is often on the effect of these fires on the aerosol distribution in North America and Siberia (Eck et al., 2009; Warneke et al., 2010). Long range transport of biomass burning plumes has been also recognized as a significant source of aerosol in the mid-latitude free troposphere over Europe (Müller et al., 2005; Fiebig et al., 2003; Sciare et al., 2008; Adler et al., 2011). Air mass aging related to long range transport also leads to aerosol optical and chemical properties different from results obtained when looking at observations close to the fire region (Lioussé et al., 1995; Müller et al., 2007; Bougiatioti et al., 2014). As an example, the absorbing efficiency in the visible spectral range is known to significantly increase in case of internally mixed BC (coating with secondary compounds) compared to externally mixed BC (Schnaiter et al., 2005). So far little attention has been paid to the frequent mixing of dust and biomass burning (BB) aerosol occurring during their transatlantic long range transport while lidar data analysis has shown that such a mixing will likely modify the extinction to backscatter ratio often called Lidar Ratio (LR) and then the aerosol optical depth (AOD) (Cattrall et al., 2005; Gross et al., 2011). Results of Paris et al. (2010) also show that the solubility of iron is enhanced by the mixing with biomass burning aerosols, while aerosol deposition may influence the rate of nitrogen fixation by microorganisms, and subsequently the global carbon cycle (Guieu et al., 2014). Although episodic, such long-range transport of smoke aerosols over the Mediterranean can also impact the regional energy budget by changing the distribution of solar energy. Indeed, for an aged BB plume, Formenti et al. (2002) report a net shortwave radiative forcing over the sea (daytime average) up to -64 W m^{-2} , at the surface and up to -22 W m^{-2} , at the top of the atmosphere (for an AOD of 0.40 at 550 nm). The large concentration of absorbing material (BC particles) within smoke plumes leads to significant absorption of solar radiations within the atmospheric layer where smoke resides, that could perturb the relative humidity and temperature vertical profiles. In the framework of the Chemistry-Aerosol Mediterranean Experiment/Aerosol Direct Radiative Impact in the Mediterranean (ChArMEx/ADRIMED) experimental campaign, many aerosol lidar and aircraft measurements have been made in June-July 2013 in the Mediterranean region during a case of intense biomass burning transport from North America to Europe (Mallet et al., 2016; Chazette et al., 2015; Pelon et al., 2015). Only a few studies report such long-range transport observations from North America to Europe (Forster et al., 2001; Petzold et al., 2007) or even the eastern Mediterranean (Formenti et al., 2002).

The purpose of this paper is to analyze the transatlantic long range transport of BB and dust aerosol sources from North America during this period. The context of our study is described in section 2 by describing the main characteristics of the summer 2013 BB episode in North America and the observation network considered for the analysis of the aerosol distribution in the Mediterranean

region. The aerosol sources are identified using satellite observations and the transport of dust or BB plumes is calculated with the FLEXPART Lagrangian model (see section 3). The aerosol lidar
145 observations are discussed in section 4, where the contribution of the different aerosol sources is assessed using the comparison of the spatial distribution of the layers with the FLEXPART model simulations (forward from the sources region and backward to calculate the potential emission sensitivity for each observed aerosol layers). The mixing between dust and BB plumes is mainly derived from the analysis of the aerosol layer optical properties. The Menorca and aircraft lidar observations
150 during ChArMEx are thoroughly described in a companion paper (Chazette et al., 2015) submitted with this paper and in a paper in preparation by J.Pelon and coworkers.

2 Context

2.1 The 2013 North American biomass burning period

June 2013 was on the drier side in the USA High Plains Region with most areas receiving less
155 than 70% of normal precipitation. It was especially dry for most of Colorado and Wyoming which received less than 50% of normal precipitation and many locations in the western areas of those states received little to no precipitation. As a consequence many fires took place in North America. Fire started in Colorado State on 10 June and lasted until 22 June 2013 (Colorado HSEM, 2013). Two large fires burning in Southern Colorado even produced pyrocumulonimbus clouds and very
160 large smoke plumes on 19 and 20 June 2013 in the West Fork Complex, and in the East Park.

In Canada there have been also many fires (334) during the period 13 to 26 June 2013 burning 632,000 ha. The seasonal fire occurrence was below average while the area burned was more than twice the 10-year average, due to large fires burning in Quebec. The majority of fires were spread between Manitoba, Alberta, Yukon and North Western Territories and Quebec, while 75% of the
165 area burned was in Quebec and 20% in Manitoba (CIFFC, 2013). The total amount of area burned was around 500,000 ha for the period 12 to 25 June 2013, i.e. more than twice the 10-year summer average for the same period. The East Canadian fires at -80°W and -100°W took place during 4-6 days between 18 and 24 June while the fires west from -120°W took place during 2-3 days starting on 17 June in Alaska and 22 June in the Mackenzie mountains.

170 2.2 The 2013 Mediterranean lidar observation network

During ChArMEx an intensive observation period took place in Western Mediterranean region from 11 June to 5 July 2013 (SOP-1a) when airborne measurements were made by two aircraft (ATR42 and F20) and ground based observations at 4 sites in Lampedusa, Corsica, Barcelona and Menorca (Mallet et al., 2016). During ChArMEx, aerosol backscatter vertical profiles were made by airborne
175 and ground based lidar systems which provide a very good opportunity to characterize the vertical distribution of the North American BB plume over the Mediterranean region. The map of the

ChArMEx lidar observation network is shown in Fig.1. The Falcon 20 aircraft was equipped with an airborne lidar LNG (Pelon et al., 2002) providing attenuated backscatter vertical profiles at three wavelength (1064, 532 and 355 nm). It was based in Cagliari, Sardinia. The LNG lidar has been
180 mainly used with a downward looking mode. Two tracks have been made in late June during the passage of the BB plume over the Western Mediterranean: a transect between Cagliari and Menorca on 27 June 2013 and a loop around Sicily on 28 June. Only the loop on 28 June is considered in this work because the 27 June data will be discussed in a future paper by J. Pelon and co-workers on the airborne observations during ChArMEx. The ground based lidar are located in Menorca (40°N,
185 4°E), Barcelona (41.4°N, 2°E) and Lampedusa (35.5°N, 12.5°E). The Menorca lidar works at 355 nm, while the Barcelona and Lampedusa lidar measure the atmospheric backscatter signal at 532 nm. The ground based lidar systems are respectively described in Chazette et al. (2014), Kumar et al. (2011), Di Iorio et al. (2009). The airborne lidar LNG was also run every morning in Cagliari (39°N, 9°W) from 24 to 30 June 2013 pointing upward from the surface. All the lidars can record
190 also the depolarization ratio between the signal polarized parallel and perpendicular to the plane of the outgoing beam. For the ground based lidar, the uncertainty on the aerosol depolarization ratio is of the order 1-2% as explained in Chazette et al. (2012). For the airborne lidar LNG, the depolarization ratio is measured at 355 nm and it is calibrated on molecular scattering using a value of $1.5 \pm 0.3\%$ for clean air, corresponding also to 1-2% error on the aerosol depolarization ratio.

195 In addition to ground based and airborne lidar, the observations of the spaceborne Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) are known to be very useful to track aerosol plumes (Winker et al., 2009). Three CALIOP night-time tracks shown in Fig.1 on 27 and 28 June 2013 are ideally located above the ChArMEx area when the BB plume is expected over Europe.

3 Aerosol sources and transport

200 3.1 Methodology

Satellite remote sensing were considered for the BB aerosol sources identifications: both Moderate Resolution Imaging Spectroradiometers (MODIS) on Terra and Aqua platforms, and CALIOP. The distribution of the fires was taken from the NASA Fire Information for Resource Management System (FIRMS) which provides the analysis of the MODIS hot spots in terms of fire radiative power
205 (FRP) given in MW. Only fire areas with $FRP > 0.8$ GW are included in this analysis. The MODIS $0.5 \mu\text{m}$ AOD daily product is also considered to estimate the horizontal extent of BB plume when large $AOD > 0.3$ is found near the spots with elevated FRP. Both MODIS instruments on Aqua and Terra are considered to derive the daily mean. The mean error on the MODIS AOD daily product is 0.03 with a root mean square error of 0.14 according to Ruiz-Arias et al. (2013). When a CALIOP
210 overpass is found near the MODIS BB plume, the lidar vertical cross section is used to specify the vertical extent of the MODIS BB plume.

For the dust aerosol sources, two main information sources were considered: (i) North American dust storms identified in the NAAPS (Navy Aerosol Analysis and Prediction System) Global Aerosol Model simulations and (ii) $0.5 \mu\text{m}$ AOD anomalies from the MODIS daily products. AOD streamers transported from the tropical Atlantic belt of elevated $0.5 \mu\text{m}$ AOD to the mid-latitudes are related to the transport of Saharan dust across the Atlantic. CALIOP overpasses near the AOD anomalies again provide the vertical extent of dust aerosol layers.

In this work we use the new CALIOP level-1 (L1) version 4.0 attenuated backscatter coefficients β_{1064} and β_{532} because they correspond to a better calibration of the lidar data. They are averaged using a 10 km horizontal resolution and a 60 m vertical resolution (Vaughan et al., 2012). Before making horizontal or vertical averaging, the initial 333 m horizontal resolution (1 km above the altitude 8.2 km) are filtered to remove the cloud layer contribution (Winker et al., 2009). This cloud mask makes use of the Version 3 level-2 (L2) cloud layer data products (Vaughan et al., 2009). Our scheme for distinguishing cloud and aerosol is described in Ancellet et al. (2014). Although the lidar ratio (LR) is available from the CALIOP Version 3 L2 aerosol layer data products, it is often based on an aerosol classification algorithm (Omar et al., 2009). In our work the lidar ratio (LR) is recalculated by using the aerosol layer transmittance and the integrated attenuated backscatter in




Figure 1. Map of the ChArMEx lidar observations. The colored vertical line are the positions of the nighttime CALIOP tracks on 27 and 28 June 2013. The red thick line shows the loop followed by the Falcon 20 aircraft on 28 June 2013 from 13 UT to 15 UT, while the blue crosses are for the Menorca, Barcelona and Lampedusa ground based lidar.

the aerosol layer following the method described in Young (1995). To reduce the error when using high horizontal resolution CALIOP profiles, β_{532} is averaged over 80 km to compute the plume transmittance whenever it is possible. **The attenuated backscatter is then corrected for the molecular and aerosol attenuation using a forward Fernald inversion (Fernald, 1984)** before calculating the backscatter ratio $R(z) = (\beta_a + \beta_R)/\beta_R$ at 532 nm and 1064 nm using the CALIOP atmospheric density model to calculate the β_R Rayleigh backscatter vertical profiles. The aerosol depolarization ratio δ_{532} is also calculated using the perpendicular- to the parallel plus perpendicular polarized aerosol backscatter coefficient. **The calibration of the relative ratio between the two 532 nm channels is based on regular use of a pseudo depolarizer located ahead of the beamsplitter which separates the signal polarized parallel and perpendicular to the plane of the outgoing beam (Winker et al., 2009).** We have also derived the color ratio defined as the ratio of the aerosol backscatter coefficients at 1064 and 532 nm ($C_a(z) = \beta_{a1064}/\beta_{a532} = (R_{1064}(z) - 1)/[16(R_{532}(z) - 1)]$). The aerosol color ratio can be also written as $C_a(z) = 2^{-k}$, where k is an exponent depending on the aerosol microphysical properties (Cattrall et al., 2005). **The exponent k varies from 0 to 2 when increasing the fine mode aerosol contribution.** These two ratios are provided only for $R(z) > 1.3$ because the uncertainty on the depolarization and color ratios are large for weak aerosol layers. Whenever it is possible, the use of nighttime overpasses are preferred to improve the signal-to-noise ratio (SNR).

The transport of the aerosol sources is analyzed using the FLEXPART model version 8.23 (Stohl et al., 2002) driven by 6-hourly ECMWF analysis (T213L91) interleaved with operational forecasts every 3 hours. The model is run using a forward simulation with a tracer released within a volume estimated from the satellite observations. The release time period ranges from 1-3 days according to the MODIS AOD observations. The total mass of the tracer emitted is estimated using the aerosol concentration given in the NAAPS Global Aerosol Model simulations and FLEXPART calculates the gridded tracer concentration in $\text{ng}\cdot\text{m}^{-3}$. Considering the uncertainty in the estimate of the emitted tracer mass, the tracer distribution in the ChArMEx domain is analyzed using a relative mass fraction between the emitted mass and the calculated mass within the model grid cell. A factor is applied to calculate this ratio in order to take into account on one hand the difference between the emission volume ($\approx 5 \cdot 10^5 \text{ km}^3$) and the grid cell volume of the tracer concentration field ($\approx 2 \cdot 10^3 \text{ km}^3$), and on the other hand, the time difference between the emission period (1-3 days) and the integration time (6 h) used for the calculation of the tracer gridded concentration. The relative mass fraction is 100% when the air mass is advected above the $0.5^\circ \times 0.5^\circ$ grid cell chosen for the gridded concentration calculation, without dilution ($< 100\%$) or concentration ($> 100\%$) of the tracer.

260 **3.2 North American biomass burning aerosol**

The MODIS FRP distributions are plotted in Fig.2 from 17 to 25 June 2013 showing the 6 main fire regions over Canada and Colorado. The map of the $0.5 \mu\text{m}$ daily AOD MODIS also show aerosol plumes on 22 June near Hudson Bay, Colorado and over the Atlantic ocean where the AOD is > 0.4 .

The white area on the daily mean MODIS map often correspond to the cloud distribution which was high over Quebec explaining the lack of large AOD daily mean values near this strong BB source. Nearby CALIOP tracks on 21 June over Colorado and 22 June over Canada show the vertical extent of the aerosol layers related to the fires (Fig.3). The aerosol layers reach 8 km over Colorado, while they remain below 4 km over Canada. The aerosol depolarization ratio is less than $7\pm 3\%$ for the layers over Canada, while it is near $9\pm 3\%$ in the mid troposphere over Colorado. The uncertainty on the CALIOP aerosol depolarization ratio averaged over the two layers are calculated using the error on the 532 nm backscatter signals. Notice also the high depolarization ratio ($>15\%$) over Colorado below 3 km showing that the BB plume overlays dust layers in the lower troposphere. The 6 areas shown in blue in Fig.2 are considered for a forward run of FLEXPART in order to study the long range transport of the Canadian and Colorado biomass burning tracer. The depth of the volume is set according to the CALIOP vertical distributions shown in Fig.3. The parameters of the different BB sources considered in the FLEXPART simulations are given in Table 1.

The map of the biomass burning tracer plume over the ChArMEx domain on 27 and 28 June is shown in Fig.4 using the relative fraction between the emitted mass and the simulated mass in the grid cell of the tracer field as explained before. Two different maps are given for the Canadian and Colorado fire contribution respectively. The Canadian plume has crossed the whole Western Mediterranean basin being over Menorca already on 27 June 06 UT and passing over Sicily on 28 June during the day. The Colorado fires do not play a major role in the aerosol layers observed on 27 June, but according to the transport model they could be observed on 28 June mainly over Spain and also in a 200- km wide strip parallel to a line from Gibraltar to Messina. The vertical cross sections (Fig.5) show that the front edge of the Canadian fires are above 4 km on 27 June while the tails bring aerosol at lower altitudes in the 1-4 km altitude range on June 28. The Colorado fires can be only detected above 5 km. The relative mass fraction is larger than 30% in the Canadian fire plume showing that a significant part is indeed advected above the Mediterranean while the remaining part is transported to Central Europe as observed by the EARLINET lidar network in Germany Gross et al. (2015). The relative fraction for the Colorado fires remains in the range 20-30% because the major part of the plume remains over Spain and the Atlantic ocean.




Figure 2. (Left panel) MODIS Fire Radiative Power from 17 to 25 June 2013. Areas with red dots are considered as significant fires. The blue boxes correspond to area chosen for the release of particles in the FLEXPART forward simulation. (Right panel) Daily AOD $0.5 \mu\text{m}$ measured by MODIS on 22 June 2013. The CALIOP tracks used to estimate the height of layer influenced by the fires are shown in red on 22 June near Hudson Bay fires and 21 June 2013 near Colorado fires




Figure 3. CALIOP vertical cross section of backscatter ratio (top), aerosol depolarization ratio (middle) and aerosol color ratio (bottom) for the two tracks shown in Fig.2 on 21 (left) and 22 (right) June 2013. Depolarization and color ratios are only reported for backscatter ratio >2 . Aerosol layers on 21 and 22 June 2013, are near the Colorado and Canadian fires, respectively.




Figure 4. Map of the relative fraction of the FLEXPART biomass burning tracer plume in % for the Canadian (top) and Colorado (bottom) fires on 27 June 2013 06 UT (left) and 28 June 18 UT (right). The altitude range corresponds to the vertical levels included in the calculation of the tracer relative fraction.




Figure 5. Vertical cross section of the relative fraction of the FLEXPART biomass burning tracer in % for the Canadian fires on 27 June 2013 06 UT (top), 28 June 18 UT (middle) and the Colorado fires on 28 June 18 UT (bottom).

3.3 North American dust layers

Modeling and satellite observations suggest that the Western USA is a significant contributor to the global mineral dust aerosol budget (Ginoux et al., 2001) and mineral dust emissions from this source region may have increased during the last 20 years (Brahney et al., 2013). Several dust blows hit Utah, Colorado and Wyoming in June 2013 due to the very dry conditions and strong winds which were also the cause of the Colorado forest fires (Hahnenberger and Nicoll, 2012). The NAAPS aerosol transport model simulations indicate elevated surface dust concentrations ($> 300 \mu\text{g}\cdot\text{m}^{-3}$) from 19 to 22 June 2013 in a region almost similar to the large MODIS AOD area related to the Colorado fires. It also explains the aerosol layers with large depolarization seen by CALIOP on 21 June 2013 at 41 N at 3 km below the Colorado fires (Fig.3). In addition to the local sources coming from Western USA, the MODIS maps on 20 and 21 June also show that dust streamers are transported at latitudes north of 30°N from the large scale Saharan dust plume, crossing the Atlantic because of the trade winds. Three streamers are shown in Fig.6 over the Atlantic ocean where the 0.5 μm AOD is enhanced with values > 0.3 . Nearby CALIOP tracks on 20 and 21 June show that the AOD enhancement are indeed related to the contribution of aerosol layers with large depolarization $> 20\%$ (Fig.7). **The uncertainty on the average depolarization ratio for the dust layers is of the order of 5%.** The 3 areas shown in blue in Fig.6 are considered in our study in order to analyze the role of dust layers over the Atlantic in the aerosol distribution over the Mediterranean Sea. According to the CALIOP vertical cross sections, the northern layer at 42°N was already uplifted in the altitude range 3-5 km while the dust plumes near 30°N remains below 3 km. Four areas are then selected for a FLEXPART forward run of dust tracers (see Table 2). The emission volume is set according to the MODIS AOD anomalies horizontal extent and the CALIOP vertical distribution of the dust layers. The emission period is chosen between 20 and 22 June for the dust layers over the Atlantic when the AOD anomalies are observed with MODIS, while the time frame for the High Plain region dust source is set according to the NAAPS model simulations.

The amount of tracer related to the High Plains dust sources was found to be negligible over the ChArMEx area during the period 27 and 28 June (mass fraction $< 10\%$) and it will not be considered any further. It may have been however mixed with the lower boundary of the Colorado fire plume seen at higher altitudes as shown in the previous section. The maps of the Atlantic dust tracer plume over the ChArMEx campaign domain is shown on 27 and 28 June in Fig.8. The values of the mass fraction is significant ($> 30\%$) showing that the contribution of long range transport of dust cannot be neglected even during the event of biomass burning aerosol transport to Europe. A first plume of dust was advected across the Western Mediterranean basin already before 27 June and a second crossed the basin on 28 June. The tail of the first one is at relatively low altitude (< 4 km) on 27 June while the second one is above 5 km on 28 June.




Figure 6. Daily AOD $0.5 \mu\text{m}$ measured by MODIS on 20 June 2013. The CALIOP tracks used to estimate the heights of the dust layers over the Atlantic ocean are shown in red on 20 June at 42°N and 21 June at 30°N . The blue boxes correspond to areas chosen for the release of particles in the FLEXPART forward simulation.




Figure 7. Same as Figure 3 for the two tracks shown in Fig. 6 on 20 (left) and 21 (right) June 2013. Dust layers are seen above the Atlantic ocean in the altitude range 1-4 km, near 32° and 42°N, on 20 and 21 June 2013, respectively.




Figure 8. Map (top) and vertical cross section (bottom) of the relative fraction of the FLEXPART Atlantic dust tracer in % on 27 June 2013 06 UT (left) and 28 June 18 UT (right). The altitude ranges in the top figures correspond to the vertical levels included in the calculation of the tracer relative fraction.

Table 1. Characteristics of the biomass burning tracer emission used for the forward FLEXPART simulation. The emitted mass is only a rough estimate explaining the use of relative mass fraction in the simulation analysis. Altitude is given above ground level.

Aerosol Source	Release Time	Horizontal domain	Vertical range,km	Emitted mass, kg
Quebec BB	18-24 June 2013	-80W/-70W, 51N/54N	0-3	$3 \cdot 10^7$
Manitoba BB	20-24 June 2013	-102W/-95W, 57N/61N	0-3	$2.5 \cdot 10^7$
NWT BB	22-24 June 2013	-128W/-121W, 61N/65N	0-5	$2 \cdot 10^7$
N.Alaska BB	17-19 June 2013	-160W/-154W, 60N/64N	0-5	$1.9 \cdot 10^7$
S.Alaska BB	19-22 June 2013	-154W/-148W, 65N/69N	0-5	$3.6 \cdot 10^7$
Colorado BB	19-22 June 2013	-105W/-96W, 37N/41N	0-6	$5 \cdot 10^7$

Table 2. Same as Table 1 for the dust tracer emission

Aerosol Source	Release Time	Horizontal domain	Vertical range,km	Emitted mass, kg
Dust High Plains	19-22 June 2013	-105W/-99W, 37N/40N	0-3	$5 \cdot 10^7$
Dust over Atlantic	20-21 June 2013	-60W/-50W, 37N/43N	1-5	$5 \cdot 10^7$
Dust over Atlantic	20-21 June 2013	-69W/-59W, 25N/33N	1-4	$5 \cdot 10^7$
Dust over Atlantic	20-21 June 2013	-48W/-38W, 25N/33N	1-4	$5 \cdot 10^7$

3.4 Saharan dust

Although the synoptic wind conditions (northwesterly flow) from 25 to 29 June 2013 were not favorable for the export of Saharan dust to the basin as explained in Mallet et al. (2016), it is important to set the northern limit of the area influenced by the northward transport of Saharan dust. The characteristics of the dust emissions were estimated using the Multi-angle Imaging SpectroRadiometer (MISR) AOD maps for the period 22 to 28 June (not shown) because the multiangle observations are better suited to distinguish surface and dust contribution to the solar reflection. The depth of the Saharan dust layer has been estimated looking at several CALIOP overpasses above North Africa during the same period. A FLEXPART forward run with a Saharan dust tracer was made for a wide area over Northern Africa in the box (24°-34.5°N, 0°-10°E, 0-6 km) from 23 to 28 June 2013.

The vertical layering of the Saharan dust tracer over the ChArMEx domain is shown on 28 June in Fig.9. As expected for a nearby source, the relative mass fraction is very large ($> 100\%$). Although the dust outflow from Sahara is transported above Lampedusa, it remains south of 36.6°N between Lampedusa and Cagliari. No Saharan dust is expected above Menorca. The altitude of the dust plume is between 2.5 and 4.5 km because the uplifting in the westerly flow is very limited.

4 Aerosol observations in the Mediterranean basin

In this section, the ChArMEx aircraft or ground based lidar observations and the CALIOP vertical cross sections on 27 and 28 June 2013 are compared with the expected contributions of the different
345 aerosol sources transported across the Atlantic.

4.1 Spatial distribution of the aerosol layers

Three nighttime CALIOP overpasses are suitable for a comparison with the different BB plumes:
27 June at -10°W and at 10°E , and 28 June at 0°W . The backscatter ratio $R(z)$ and the aerosol
depolarization ratio δ_{532} are shown in Fig.10 in the latitude range where cloud free sky made possible
350 the observations of aerosol layers. The $R(z)$ values are larger than 3 in these layers. On 27 June,
the layers are in the altitude range 5 to 7.5 km at 10°E while it is between 2 and 5 km at 10°W . In
both cases low δ_{532} values ($<10\%$) are found, showing that the plumes are not mixed with significant
amount of dust (except at 10°W where δ_{532} may reach 10% in some layers). **The uncertainty for δ_{532}
is of the order of 3%.** These results are in good agreement with the characteristics of the Canadian
355 fire plumes discussed in section 3. Indeed it was found that the front edge of the plume was at 10°E
on 27 June with an altitude range 4-7 km, while the tail is at -10°W in the altitude range 2-5 km
(see vertical cross section on 28 June in Fig.5). Although the Colorado fires may be present in the 27
June CALIOP cross section at -10°W according to the FLEXPART simulations, the altitude range
of the observed aerosol layers is not consistent with the influence of the Colorado BB plume which
360 is expected at altitude above 5 km. On 28 June at 0°W , the CALIOP observations show also aerosol
layers in the 3-5 km altitude range, with slightly higher depolarization ratio ($\approx 10\pm 3\%$), but still in




Figure 9. Same as figure 8 for the FLEXPART Saharan tracer on 28 June 2013 18 UT.

the range expected for biomass burning aerosol (Nisantzi et al., 2014). The altitude range is again in good agreement with a major role of the tail of the Canadian fire plume.

Several ground based lidar observations have also identified aerosol plumes possibly related to the transatlantic transport. The characteristics of the aerosol layers are summarized in Table 3. The Menorca lidar data are discussed in a companion paper by Chazette et al. (2015). An aerosol layer between 3-5 km seen in Menorca is quite similar to the CALIOP observations on 28 June. A second layer between 5-7 km is also seen in Menorca with a noticeable depolarization ($\delta_{355} > 12 \pm 1\%$). The upper layer is not seen by CALIOP because it is expected at latitudes higher than 40°N and is masked by overlaying clouds. In Menorca the vertical profiles of the water vapor mixing ratio were also measured during the night (Chazette et al., 2014) showing elevated mixing ratio > 1 g/kg above 5 km and values near 0.5 g/kg in the aerosol layer observed around 4 km. The time series of the Menorca lidar is also useful to estimate the horizontal range of BB plume. The plume is observed for 24 hours from 27 June 00 UT to 28 June 12 UT and the wind speed at 4 km is between 30-40 km/h, therefore the plume zonal extent is of the order of 1200 km. It is very similar to the size of the Canadian tracer plume obtained in the FLEXPART simulations (15° longitude difference between the front edge and the tail).

As expected the Barcelona lidar detects similar features: a strong layer between 5-7 km with $\delta_{532} \approx 10\%$ and an optically thin layer between 3-5 km with $\delta_{532} < 10\%$. The spectral variation of the aerosol depolarization ratio between Barcelona and Menorca cannot be accurately estimated but is less than 1.5. It is consistent with a small influence of urban aerosol (Burton et al., 2012). When looking at the Lampedusa lidar data at 35°N, a layer is seen between 2 and 4 km on 28 June which is influenced by the Saharan dust outflow discussed in section 3 since $\delta_{532} \geq 30\%$, i.e. a value similar to other dust layers observed over Menorca during ChArMEx (Chazette et al., 2015).

The Lampedusa lidar measures aerosol layers in the 2-4 km altitude range on both days, but with very different optical characteristics. A dust layer with $\delta_{532} > 30\%$ on 28 June 12 UT while a mixture of dust and BB aerosol is seen on 27 June from 8 UT to 16 UT. The aerosol layer seen by CALIOP on 27 June 01 UT near 36°N has optical characteristics close to the layer observed in Lampedusa on 27 June (Fig.10), i.e. a depolarization between 10%-15% and LR between 50-55 sr.

The LNG airborne lidar data obtained during ChArMEx will be thoroughly discussed in a forthcoming paper by J.Pelon and co-workers. Here we will only consider the vertical structure of the aerosol layers observed on 28 June 2013 along the loop shown in Fig.1. The 3 corresponding vertical cross sections of attenuated $R(z)$ at 532 nm are shown in Fig.11. Three interesting regions can be identified:

- (A) the 38.2°N layer at 2-4 km on the Cagliari-Lampedusa section and at 11-14°E on the return section between Messina and Cagliari,
- (B) the upper altitude layer in the 4-6 km altitude range covering a southwest (36°N, 12°E)-northeast (39°N, 15°E) band, the width of which is of the order of 100 km,

Table 3. Characteristics of the aerosol layers observed in the free troposphere by the ground based lidars listed in Fig.1 on 27 and 28 June 2013.

Lidar	Wavelength nm	Layer altitude	Time period	Scattering ratio	Depolarization ratio	Lidar ratio
Menorca	355	5-7 km	27/6 00UT to 28/6 00UT	1.87 ± 0.03	$10\pm 1\%$	42 ± 5 sr
Menorca	355	3-5 km	27/6 00UT to 28/6 12UT	1.22 ± 0.02	$4\pm 1\%$	59 ± 5 sr
Barcelona	532	5-7 km	26/6 20UT to 27/6 16UT cloudy after	4.3	$\approx 10\pm 2\%$	
Barcelona	532	3-5 km	26/6 12UT to 27/6 16UT cloudy after	1.4	$7\pm 2\%$	
Lampedusa	532	2-4 km	27/6 08UT to 27/6 16UT	1.5 ± 0.03	$12\pm 2\%$	51 ± 9 sr
Lampedusa	532	2-4 km	28/6 08UT to 28/6 14UT	2.8 ± 0.04	$30\pm 2\%$	30 ± 6 sr

400 – (C) a low altitude layer between 2-4 km south of 36 N which corresponds to the layer seen by the Lampedusa lidar.

The spatial distribution of the aerosol layers seen by the LNG lidar corresponds quite well with the position of North American BB plumes and the expected latitudinal extent of the Saharan dust calculated with the FLEXPART simulation in section 3. Indeed the layer A is related to the tail of the Canadian BB plume. The layer B is also in the latitude range of the Canadian BB plume, possibly mixed with the Colorado BB plume present between Gibraltar and Messina (see Fig.4). Layer A and B seen by the LNG airborne lidar on 28 June are also consistent with the superposition of two different aerosol layers seen above Menorca 24 hours before. In layer B, $\delta_{355} \approx 10\pm 1\%$, i.e. higher than the low values found in layer A ($\delta_{355} \leq 5\pm 2\%$).

4.2 Aerosol source attribution

410 Although the comparison with the position of the FLEXPART tracer plumes can already help to attribute a specific source to the observed layers in the ChArMEx area, it can be further checked by calculating the potential emission sensitivity (PES) values by running the FLEXPART model in the backward mode for 10-11 days to identify the area where surface emissions may influence the observed aerosol structure seen by CALIOP, the Menorca ground based lidar and the LNG airborne instrument. The PES is given in s unit in order to be multiplied by model surface fluxes to produce




Figure 10. CALIOP vertical cross section of backscatter ratio (left panel), aerosol depolarization ratio (right panel) for the 3 tracks shown in Fig.1 on 27 June 2013 03 UT at -10° W (top panel), on 28 June 2013 2 UT at 0° W (middle panel), on 27 June 2013 01 UT at 10° E (bottom panel).

concentrations at the receptor location. The PES is calculated using 6 hours averages on a three dimensional $1^\circ \times 1^\circ \times 1$ km grid. The results are shown for the CALIOP and Menorca observations on 27 June 2013 (Fig.12). Similar calculations were also made for layer A, B and C seen by the airborne lidar on 28 June (Fig.13). The simulations for the layers seen by CALIOP on 28 June 02
420 UT are not shown because they are very similar to the results obtain for the Menorca lidar on 27 June at 12 UT or for layer B seen by LNG.

The aerosol layers observed by CALIOP along the two cross sections at -10°W and 10°E are indeed mainly related to aerosol sources over Canada and Alaska, but the retro-plume altitude and latitude at 60°W are quite different when reaching the Atlantic Ocean. The probability of dust and
425 biomass mixing is higher for the CALIOP layers at -10°W which is located in the 40° - 50°N latitude band at lower altitude (5-7 km) than for the CALIOP layers at 10°E . This may explain the slight depolarization difference for the two CALIOP tracks since there are more layers with $\delta_{532} \approx 10\%$ at -10°W than at 10°E . The mixing of dust layers over the Atlantic and Canadian BB aerosol is even more explicit for the Menorca layer at 6 km since two branches of elevated PES are seen over the two
430 aerosol source regions identified for this layer. It explains the relatively higher aerosol depolarization ratio (up to 12%) at 6 km than at 4 km in Menorca during this episode. Such a transport pathway is also consistent with the water vapor mixing ratio maximum > 1 g/kg seen by the Menorca water vapor lidar near 6 km since uplifting of air masses from the lower troposphere above the Atlantic Ocean is likely to increase the humidity in the mid-troposphere.

435 When considering the PES related to the airborne lidar layers, the layer A PES is similar to the 10°E plume showing a strong influence of the Canadian aerosol BB source, while the layer B PES distribution resembles the results obtained with the Menorca layer seen one day earlier. For the dusty layer C seen both by the aircraft and at the Lampedusa station, the PES distribution shows that there is no transatlantic transport for the period 17 to 28 June while the aerosol sources are mainly
440 located above North Africa and Western Europe at low altitude (< 3 km). Although air masses are still advected from Western Europe, Saharan dust emission remains the major aerosol source since Western Europe air masses were heavily influenced by Saharan dust layers during the period 16 to 20 June (Mallet et al., 2016). It is consistent with the large depolarization seen both above Lampedusa ($\delta_{532} \approx 30\%$).

445 The ATR42 aircraft also flew between Cagliari and Lampedusa on 28 June around 12 UT to sample the aerosol layers with in-situ measurements (Pelon et al., 2015). The analysis of the CO and BC in-situ measurements made on-board the ATR42 shows that the layer A and B correspond to a CO excess above background of the order of 100 ppbv while ΔCO is less than 20 ppbv for layer C (not shown). The BC variability shows also the same pattern. This is in very good agreement with
450 the conclusions derived from the lidar data analysis coupled with the Lagrangian transport model simulations.




Figure 11. Airborne lidar vertical cross sections of attenuated backscatter ratio at 532 nm on 28 June 2013 along the loop shown in Fig.1: (top) Lampedusa/Cagliari around 13 UT (middle) Lampedusa/Messina around 14 UT (bottom) Cagliari/Messina around 14:40 UT.




Figure 12. (Left panel) FLEXPART Potential emission sensitivity (PES) in s for 3 aerosol layers identified by CALIOP and Menorca lidar: 27 June 2013 02 UT, -10°W , 43°N (top) 27 June 12 UT, in Menorca (middle) 27 June 01 UT, 10°E , 39°N (bottom). The PES Vertical cross section are along the red line following the North American East Coast (right panel).




Figure 13. Same as figure 12 for the 3 aerosol layers identified by the Falcon 20 lidar on 28 June 2013: layer A (top) layer B (middle) layer C (bottom).

4.3 Aerosol optical properties

In this section, we will summarize the results about the aerosol layer optical properties and the aerosol source attribution. The analysis conducted in the previous sections leads to the identification
455 of 4 different aerosol layers during the passage of the BB plume over the ChArMEx area:

- (I) pure BB layer at 10°E above 4 km on 27 June 2013 01 UT (CALIOP),
- (II) weakly dusty BB layer below 5 km observed between -10°W and 10°E on 27 and 28 June
02 UT by CALIOP, the Menorca lidar and layer A seen by LNG flying around Sicily on 28
June.
- 460 – (III) significant mixture of BB and dust transported across the Atlantic above 5 km at Menorca
on 27 June 12 UT and layer B seen by LNG on 28 June 14 UT.
- (IV) the outflow of Saharan dust above the sea at latitudes South of 36°N on 28 June 13 UT
(Lampedusa lidar and layer C seen by LNG).

For layer (I) δ_{532} is $< 5\%$, while the LR at 532 nm is 60 ± 20 sr when using the aerosol layer
465 transmission from the averaged L1 CALIOP attenuated backscatter, and it is 65 sr in the level-2 (L2)
CALIOP operational aerosol data products. The color ratio is between 0.2 and 0.4. Both LR and C_a
are in the range expected for a pure BB layer (Burton et al., 2012) in agreement with our source
identification.

For layer (II) δ_{532} and δ_{355} are respectively in the range 5-10% and $< 8\%$ with the lowest values
470 on 27 June at 45°N along the -10°W CALIOP overpass. The LR calculated from the ground based
and airborne lidars are more accurate and they are 59 ± 5 sr at 355 nm and 60 ± 5 sr at 532 nm ,
respectively. The CALIOP 27 June (-10°W) and 28 June LR at 532 nm estimated to be 60 ± 20 sr
and 50 ± 20 sr, respectively, using the L1 data analysis and are of the order of 60 sr for both layers
using the L2 operational products. It also gives confidence in the LR retrieval to see the largest
475 LR is obtained where δ_{532} is minimum. The CALIOP C_a is in the range 0.4-0.5. These aerosol
optical parameters are still in the range expected for a BB layer. However differences with the optical
parameters found for layer (I) (higher depolarization and C_a , slightly lower LR) are consistent with
a BB mixed with a small amount of dust or an increase in relative humidity. According to the small
values (< 0.5 g/kg) of water vapor recorded by the Menorca lidar observations (Chazette et al., 2015),
480 the mixing with a small amount of dust is more likely.

For layer (III) mainly seen by the Menorca ground based and airborne lidar, δ_{355} values are in the
range 8-12%, while LR are 45 ± 5 sr and 42 ± 5 sr at 532 and 355 nm, respectively. The LR of the
layer B seen by LNG is calculated by including also the contribution of the underlying layer between
3-4 km to get a better molecular reference. Following the methodology proposed by Tesche et al.
485 (2009), the aerosol depolarization value for layer (III) is consistent with a contribution of 20%-30%
of dust and 80%-70% of BB aerosol in the total aerosol backscatter in layer (III), if we assume that

Table 4. Optical properties of the four types of aerosol encountered during the passage of the BB plume: depolarization ratios (δ_{532} , δ_{355}), lidar ratios (LR_{532} , LR_{355}), color ratios (CR_a 1064/532, CR_a 532/355)

Aerosol type	δ_{532}	δ_{355}	LR_{532} , sr	LR_{355} , sr	CR_a 1064/532	CR_a 532/355
Pure BB (I)	<5%	-	60±20	-	0.2-0.4	-
Dusty BB (II)	5-10%	5-8%	60±5*, 51±9**	59±5	0.4-0.5	0.35
BB/dust mixture (III)	≈10±2%	8-12%	45±5	42±5	-	0.74
Saharan dust (IV)	30±2%	-	48±5*, 30±5**	-	-	-

* airborne lidar ** Lampedusa lidar

pure dust and pure BB aerosol types have δ_{355} of 25% and 5%, respectively and LR at 355 nm of 45±10 sr and 60±10 sr, respectively. It is also interesting to calculate C_a between 355 and 532 nm using the Barcelona and Menorca observations assuming that $R(z)$ is stationary during the advection of the aerosol layers between Barcelona and Menorca. The C_a value is 0.74 for layer (III) while it is only 0.35 for the layer (II). It is consistent with a larger contribution of the accumulation mode when BB is mixed with dust, but also with a larger water vapor mixing ratio (1 g/kg) for layer (III) than layer (II).

For layer (IV) larger depolarization up to 30% are seen by the Lampedusa lidar at 532 nm. The lidar ratio calculated by the LNG lidar and the ground based lidar at 532 nm are respectively 48±5 sr and 30±10 sr. The layer optical parameters are consistent with a dust plume with a large depolarization, while a large variability is observed for LR. The large depolarization ratio and the low LR value at 355 nm is quite similar to previous observation by Di Iorio et al. (2009) in fresh dust exported over the Mediterranean sea. The strong variation in the LR values between the layer C of the LNG lidar at 13.5 °E and the Lampedusa observations at 12.5°E suggests an increase of the mixing between the northward African dust outflow and the BB plume as the aircraft moved across the boundary between layer (IV) and (II) between Lampedusa and the southern cape of Sicily.

The aerosol properties and spatial distribution of the four aerosol types are summarized in, Table 4 and Fig.14 respectively. The spatial distribution of the MODIS AOD at 0.5 μ m is also shown in Fig.14, where the largest AOD values are seen before the plumes dispersion above Northern Spain. For the type (II) aerosol, i.e. an aged BB plume seen below 5 km and mixed with a small amount of dust mainly from continental origin, two areas are distinguished for the Colorado and Canadian fires using the results of the FLEXPART forward simulations. The Canadian fires significantly contributes to the AOD observed by MODIS over the Mediterranean sea. The additional contribution of the upper aerosol layers of type (III) where the BB is mixed with dust also explains the significant AOD increase over the Western Mediterranean region. The BB contribution to AOD is as large as the North African dust contribution (type IV) that dominates the southern part of the domain with AOD values in the same range of 0.3-0.4 over Northeastern Algeria and Tunisia.

5 Conclusions

515 A very interesting event of long range transport of biomass burning (BB) aerosols between North
America and the Western Mediterranean region that took place in late June 2013 has been docu-
mented during the ChArMEX/ADRIMED campaign. Although the occurrence of such events has
been discussed in previous publications, the contribution of this work is to take advantage of a large
number of ground based and airborne lidar measurements used in conjunction with spaceborne li-
520 dar observations by CALIOP during this period. A detailed analysis of the biomass burning North
American sources was conducted including the assessment of their transport to Europe using for-
ward simulations with the FLEXPART model initialized using satellite observations. The specific
question of mixing between dust and BB particles was addressed by considering the possible dust
sources transported along the same transport pathway. The role of mixing was quantified by consid-
525 ering the optical properties of the different aerosol layers observed during two days of the ChArMEX
campaign (27 and 28 June 2013) when the biomass burning aerosol load was at its maximum over
the Western Mediterranean. The three dimensional structure of the aerosol distribution revealed by
the lidar network and the airborne lidar flight provides a detailed assessment of the different aerosol
source contributions when it is coupled with the results of the Lagrangian FLEXPART transport
530 model. Four aerosol types were identified using the depolarization ratio and the three dimensional
structure of the aerosol plume: (I) pure BB layer, (II) weakly dusty BB, (III) a significant mixture
of BB and dust transported from the North Atlantic trade wind region (IV) the direct northward out-
flow by the subtropical jet of Saharan dust not mixed with BB aerosol. Mixing of dust and BB can
correspond to a 20%-30% dust contribution in the total aerosol backscatter. The comparison with
535 the MODIS AOD distribution during this episode over the Western Mediterranean sea shows that
the Canadian fires contribution were surprisingly as large as the direct northward dust outflow from
Sahara. An additional contribution from a mid-tropospheric aerosol layer due a mixture of dust and
BB aerosol was found in the region of higher AOD seen by MODIS. The next step will now concern
the use of all presented and analyzed data for evaluating 3-D regional models to simulate this specific
540 event, in terms of optical properties, possible mixing and vertical extent of mineral dust and forest
fire aerosol layers.

Acknowledgements. This work was funded by the French MISTRALS program funded by CNRS/INSU, ADEME,
CEA, Météo-France and CNES for aerosol and cloud satellite missions validation. The NILU team lead by A.
Stohl is gratefully acknowledged for distributing the FLEXPART model. The SAFIRE team, INSU DT and
545 D. Bruneau from LATMOS are gratefully acknowledged for the aircraft flight operation and the LNG lidar
operation. The lidar measurements in Barcelona were supported by the 7th Framework Programme project
Aerosols, Clouds, and Trace Gases Research Infrastructure Network (ACTRIS) (grant agreement no. 262254)
and by the Spanish Ministry of Science and Innovation and FEDER funds under the projects TEC2012-34575,
UNPC10-4E-442 and CGL2011-13580-E/CLI.




Figure 14. MODIS AOD horizontal distribution on 27 and 28 June 2013 over the Mediterranean region. The area corresponding to the aerosol types identified during our analysis of the BB plume passage are delimited by the red lines. The black crosses are for the Menorca and Lampedusa stations.

550 References

- Adler, G., Flores, J. M., Abo Riziq, A., Borrmann, S., and Rudich, Y.: Chemical, physical, and optical evolution of biomass burning aerosols: a case study, *Atmospheric Chemistry and Physics*, 11, 1491–1503, doi:10.5194/acp-11-1491-2011, <http://www.atmos-chem-phys.net/11/1491/2011/>, 2011.
- 555 Ancellet, G., Pelon, J., Blanchard, Y., Quennehen, B., Bazureau, A., Law, K. S., and Schwarzenboeck, A.: Transport of aerosol to the Arctic: analysis of CALIOP and French aircraft data during the spring 2008 POLARCAT campaign, *Atmospheric Chemistry and Physics*, 14, 8235–8254, doi:10.5194/acp-14-8235-2014, <http://www.atmos-chem-phys.net/14/8235/2014/>, 2014.
- Bougiatioti, A., Stavroulas, I., Kostenidou, E., Zampas, P., Theodosi, C., Kouvarakis, G., Canonaco, F., Prévôt, A. S. H., Nenes, A., Pandis, S. N., and Mihalopoulos, N.: Processing of biomass-burning aerosol in the eastern Mediterranean during summertime, *Atmospheric Chemistry and Physics*, 14, 4793–4807, doi:10.5194/acp-14-4793-2014, <http://www.atmos-chem-phys.net/14/4793/2014/>, 2014.
- 560 Brahney, J., Ballantyne, A., Sievers, C., and Neff, J.: Increasing Ca²⁺ deposition in the western US: The role of mineral aerosols, *Aeolian Research*, 10, 77 – 87, doi:<http://dx.doi.org/10.1016/j.aeolia.2013.04.003>, <http://www.sciencedirect.com/science/article/pii/S1875963713000281>, 2013.
- 565 Burton, S. P., Ferrare, R. A., Hostetler, C. A., Hair, J. W., Rogers, R. R., Obland, M. D., Butler, C. F., Cook, A. L., Harper, D. B., and Froyd, K. D.: Aerosol classification using airborne High Spectral Resolution Lidar measurements - methodology and examples, *Atmospheric Measurement Techniques*, 5, 73–98, doi:10.5194/amt-5-73-2012, <http://www.atmos-meas-tech.net/5/73/2012/>, 2012.
- Cattrell, C., Reagan, J., Thome, K., and Dubovik, O.: Variability of aerosol and spectral lidar and backscatter and extinction ratios of key aerosol types derived from selected Aerosol Robotic Network locations, *J. Geophys. Res.*, 110, D10S11, doi:10.1029/2004JD005124, 2005.
- 570 Chazette, P., Dabas, A., Sanak, J., Lardier, M., and Royer, P.: French airborne lidar measurements for Eyjafjallajökull ash plume survey, *Atmospheric Chemistry and Physics*, 12, 7059–7072, doi:10.5194/acp-12-7059-2012, <http://www.atmos-chem-phys.net/12/7059/2012/>, 2012.
- 575 Chazette, P., Marnas, F., and Totems, J.: The mobile Water vapor Aerosol Raman Lidar and its implication in the framework of the HyMeX and ChArMEx programs: application to a dust transport process, *Atmospheric Measurement Techniques*, 7, 1629–1647, doi:10.5194/amt-7-1629-2014, <http://www.atmos-meas-tech.net/7/1629/2014/>, 2014.
- Chazette, P., Totems, J., Ancellet, G., Pelon, J., and Sicard, M.: Temporal consistency of lidar observables during aerosol transport events in the framework of the ChArMEx/ADRIMED campaign at Menorca Island in June 2013, *Atmospheric Chemistry and Physics Discussions*, 15, 32 723–32 757, doi:10.5194/acpd-15-32723-2015, <http://www.atmos-chem-phys-discuss.net/15/32723/2015/>, 2015.
- 580 CIFFC: Canadian Interagency Forest Fire Center Inc. Report 2013, Tech. rep., <http://cwfis.cfs.nrcan.gc.ca/report/archives?year=2013&month=06&day=26&process=Submit>, 2013.
- 585 Colorado HSEM: Homeland Security and Emergency Management: Colorado Wildfire Report - June 26 2013, Tech. rep., <http://www.coemergency.com/2013/06/colorado-wildfire-report-june-26-cofire.html>, 2013.
- Di Iorio, T., di Sarra, A., Sferlazzo, D. M., Cacciani, M., Meloni, D., Monteleone, F., Fuá, D., and Fiocco, G.: Seasonal evolution of the tropospheric aerosol vertical profile in the central Mediterranean and role

of desert dust, *Journal of Geophysical Research: Atmospheres*, 114, D02 201, doi:10.1029/2008JD010593,
590 <http://dx.doi.org/10.1029/2008JD010593>, 2009.

Eck, T. F., Holben, B. N., Reid, J. S., Sinyuk, A., Hyer, E. J., O'Neill, N. T., Shaw, G. E., Vande Castle,
J. R., Chapin, F. S., Dubovik, O., Smirnov, A., Vermote, E., Schafer, J. S., Giles, D., Slutsker, I., Sorokine,
M., and Newcomb, W. W.: Optical properties of boreal region biomass burning aerosols in central Alaska
and seasonal variation of aerosol optical depth at an Arctic coastal site, *Journal of Geophysical Research:*
595 *Atmospheres*, 114, n/a–n/a, doi:10.1029/2008JD010870, <http://dx.doi.org/10.1029/2008JD010870>, 2009.

Fernald, F. G.: Analysis of atmospheric lidar observations: some comments, *Appl. Opt.*, 23, 652–653,
doi:10.1364/AO.23.000652, <http://ao.osa.org/abstract.cfm?URI=ao-23-5-652>, 1984.

Fiebig, M., Stohl, A., Wendisch, M., Eckhardt, S., and Petzold, A.: Dependence of solar radiative forcing
of forest fire aerosol on ageing and state of mixture, *Atmospheric Chemistry and Physics*, 3, 881–891,
600 doi:10.5194/acp-3-881-2003, <http://www.atmos-chem-phys.net/3/881/2003/>, 2003.

Flannigan, M., Stocks, B., Turetsky, M., and Wotton, M.: Impacts of climate change on fire activity and
fire management in the circumboreal forest, *Global Change Biology*, 15, 549–560, doi:10.1111/j.1365-
2486.2008.01660.x, <http://dx.doi.org/10.1111/j.1365-2486.2008.01660.x>, 2009.

Formenti, P., Boucher, O., Reiner, T., Sprung, D., Andreae, M. O., Wendisch, M., Wex, H., Kindred, D.,
605 Tzortziou, M., Vasaras, A., and Zerefos, C.: STAAARTE-MED 1998 summer airborne measurements over
the Aegean Sea 2. Aerosol scattering and absorption, and radiative calculations, *Journal of Geophysical
Research: Atmospheres*, 107, 4451, doi:10.1029/2001JD001536, <http://dx.doi.org/10.1029/2001JD001536>,
2002.

Forster, C., Wandler, U., Wotawa, G., James, P., Mattis, I., Althausen, D., Simmonds, P., O'Doherty, S.,
610 Jennings, S. G., Kleefeld, C., Schneider, J., Trickl, T., Kreipl, S., Jäger, H., and Stohl, A.: Transport of
boreal forest fire emissions from Canada to Europe, *Journal of Geophysical Research: Atmospheres*, 106,
22 887–22 906, doi:10.1029/2001JD900115, <http://dx.doi.org/10.1029/2001JD900115>, 2001.

Generoso, S., Bréon, F.-M., Balkanski, Y., Boucher, O., and Schulz, M.: Improving the seasonal cycle and
interannual variations of biomass burning aerosol sources, *Atmospheric Chemistry and Physics*, 3, 1211–
615 1222, doi:10.5194/acp-3-1211-2003, <http://www.atmos-chem-phys.net/3/1211/2003/>, 2003.

Ginoux, P., Chin, M., Tegen, I., Prospero, J. M., Holben, B., Dubovik, O., and Lin, S.-J.: Sources and distribu-
tions of dust aerosols simulated with the GOCART model, *Journal of Geophysical Research: Atmospheres*,
106, 20 255–20 273, doi:10.1029/2000JD000053, <http://dx.doi.org/10.1029/2000JD000053>, 2001.

Gross, S., Tesche, M., Freudenthaler, V., Toledano, C., Wiegner, M., Ansmann, A., Althausen, D., and Seefeld-
620 ner, M.: Characterization of Saharan dust, marine aerosols and mixtures of biomass-burning aerosols and dust
by means of multi-wavelength depolarization and Raman lidar measurements during SAMUM 2, *Tellus B*,
63, 706–724, doi:10.1111/j.1600-0889.2011.00556.x, <http://dx.doi.org/10.1111/j.1600-0889.2011.00556.x>,
2011.

Gross, S., Geiss, A., Heimerl, K., Gasteiger, J., Freudenthaler, V., Weinzier, B., and Wiegner, M.: Characteriza-
625 tion of long-range transported Canadian biomass burning over Central Europe - A case study, *EGU General
Assembly 2015*, 2015.

- Guieu, C., Ridame, C., Pulido-Villena, E., Bressac, M., Desboeufs, K., and Dulac, F.: Impact of dust deposition on carbon budget: a tentative assessment from a mesocosm approach, *Biogeosciences*, 11, 5621–5635, doi:10.5194/bg-11-5621-2014, <http://www.biogeosciences.net/11/5621/2014/>, 2014.
- 630 Hahnenberger, M. and Nicoll, K.: Meteorological characteristics of dust storm events in the eastern Great Basin of Utah, U.S.A., *Atmospheric Environment*, 60, 601 – 612, doi:<http://dx.doi.org/10.1016/j.atmosenv.2012.06.029>, <http://www.sciencedirect.com/science/article/pii/S1352231012005808>, 2012.
- Kumar, D., Rocadenbosch, F., Sicard, M., Comeron, C., Munoz, C., Lange, D., Tomas, S., and Gregorio, E.:
635 Six-channel polychromator design and implementation for the UPC elasticRaman LIDAR, in: *SPIE Remote Sensing*, p. 81820, International Society for Optics and Photonics, Prague, Czech Republic, 2011.
- Lioussé, C., Devaux, C., Dulac, F., and Cachier, H.: Aging of savanna biomass burning aerosols: Consequences on their optical properties, *Journal of Atmospheric Chemistry*, 22, 1–17, doi:10.1007/BF00708178, <http://dx.doi.org/10.1007/BF00708178>, 1995.
- 640 Liu, Y., Goodrick, S., and Heilman, W.: Wildland fire emissions, carbon, and climate: Wildfire–climate interactions, *Forest Ecology and Management*, 317, 80 – 96, doi:<http://dx.doi.org/10.1016/j.foreco.2013.02.020>, <http://www.sciencedirect.com/science/article/pii/S037811271300114X>, 2014.
- Mallet, M., Dulac, F., Formenti, P., Nabat, P., Sciare, J., Roberts, G., Pelon, J., Ancellet, G., Tanré, D., Parol, F., Denjean, C., Brogniez, G., Di Sarra, A., Alados-Arboledas, L., Arndt, J., Auriol, F., Blarel, L., Bourriane,
645 T., Chazette, P., Chevaillier, S., Claeys, M., D’Anna, B., Derimian, Y., Desboeufs, K., Di Iorio, T., Doussin, J.-F., Durand, P., Féron, A., Freney, E., Gaimoz, C., Goloub, P., Gómez-Amo, J. L., Granados-Muñoz, M. J., Grand, N., Hamonou, E., Jankowiak, I., Jeannot, M., Léon, J.-F., Maillé, M., Mailler, S., Meloni, D., Menut, L., Momboisse, G., Nicolas, J., Podvin, T., Pont, V., Rea, G., Renard, J.-B., Roblou, L., Schepanski, K., Schwarzenboeck, A., Sellegri, K., Sicard, M., Solmon, F., Somot, S., Torres, B., Totems, J., Triquet, S.,
650 Verdier, N., Verwaerde, C., Waquet, F., Wenger, J., and Zapf, P.: Overview of the Chemistry-Aerosol Mediterranean Experiment/Aerosol Direct Radiative Forcing on the Mediterranean Climate (ChArMEx/ADRIMED) summer 2013 campaign, *Atmospheric Chemistry and Physics*, 16, 455–504, doi:10.5194/acp-16-455-2016, <http://www.atmos-chem-phys.net/16/455/2016/>, 2016.
- Müller, D., Mattis, I., Wandinger, U., Ansmann, A., Althausen, D., and Stohl, A.: Raman lidar observations of
655 aged Siberian and Canadian forest fire smoke in the free troposphere over Germany in 2003: Microphysical particle characterization, *Journal of Geophysical Research: Atmospheres*, 110, doi:10.1029/2004JD005756, <http://dx.doi.org/10.1029/2004JD005756>, 2005.
- Müller, D., Mattis, I., Ansmann, A., Wandinger, U., Ritter, C., and Kaiser, D.: Multiwavelength Raman lidar observations of particle growth during long-range transport of forest-fire smoke in the free tropo-
660 sphere, *Geophysical Research Letters*, 34, L05 803, doi:10.1029/2006GL027936, <http://dx.doi.org/10.1029/2006GL027936>, 2007.
- Nisantzi, A., Mamouri, R. E., Ansmann, A., and Hadjimitsis, D.: Injection of mineral dust into the free troposphere during fire events observed with polarization lidar at Limassol, Cyprus, *Atmospheric Chemistry and Physics*, 14, 12 155–12 165, doi:10.5194/acp-14-12155-2014, <http://www.atmos-chem-phys.net/14/12155/2014/>, 2014.
665

- Omar, A., Winker, D., Kittaka, C., Vaughan, M., Liu, Z., Hu, Y., Treppe, C., Rogers, R., Ferrare, R., Lee, K., Kuehn, R., and Hostetler, C.: The CALIPSO Automated Aerosol Classification and Lidar Ratio Selection Algorithm, *J. Atmos. Ocean. Tech.*, 26, 1994–2014, doi:10.1175/2009JTECHA1231.1, 2009.
- 670 Paris, R., Desboeufs, K. V., Formenti, P., Nava, S., and Chou, C.: Chemical characterisation of iron in dust and biomass burning aerosols during AMMA-SOP0/DABEX: implication for iron solubility, *Atmospheric Chemistry and Physics*, 10, 4273–4282, doi:10.5194/acp-10-4273-2010, <http://www.atmos-chem-phys.net/10/4273/2010/>, 2010.
- 675 Pelon, J., Flamant, C., Chazette, P., Leon, J.-F., Tanre, D., Sicard, M., and Satheesh, S. K.: Characterization of aerosol spatial distribution and optical properties over the Indian Ocean from airborne LIDAR and radiometry during INDOEX'99, *Journal of Geophysical Research: Atmospheres*, 107, 8029, doi:10.1029/2001JD000402, <http://dx.doi.org/10.1029/2001JD000402>, 2002.
- 680 Pelon, J., Ancellet, G., Chazette, P., Totems, J., Sicard, M., Dulac, F., Di Ioro, T., Formenti, P., and Mallet, M.: Lagrangian analysis of forest fire aerosol emissions from North America to Western Mediterranean basin during the CHARMEX 2013 summer campaign, *EGU General Assembly 2015*, <https://hal-insu.archives-ouvertes.fr/insu-01142730>, 2015.
- 685 Petzold, A., Weinzierl, B., Huntrieser, H., Stohl, A., Real, E., Cozic, J., Fiebig, M., Hendricks, J., Lauer, A., Law, K., Roiger, A., Schlager, H., and Weingartner, E.: Perturbation of the European free troposphere aerosol by North American forest fire plumes during the ICARTT-ITOP experiment in summer 2004, *Atmospheric Chemistry and Physics*, 7, 5105–5127, doi:10.5194/acp-7-5105-2007, <http://www.atmos-chem-phys.net/7/5105/2007/>, 2007.
- Ruiz-Arias, J. A., Dudhia, J., Gueymard, C. A., and Pozo-Vázquez, D.: Assessment of the Level-3 MODIS daily aerosol optical depth in the context of surface solar radiation and numerical weather modeling, *Atmospheric Chemistry and Physics*, 13, 675–692, doi:10.5194/acp-13-675-2013, <http://www.atmos-chem-phys.net/13/675/2013/>, 2013.
- 690 Schnaiter, M., Linke, C., Müller, O., Naumann, K.-H., Saathoff, H., Wagner, R., Schurath, U., and Wehner, B.: Absorption amplification of black carbon internally mixed with secondary organic aerosol, *Journal of Geophysical Research: Atmospheres*, 110, D19 204, doi:10.1029/2005JD006046, <http://dx.doi.org/10.1029/2005JD006046>, 2005.
- 695 Sciare, J., Oikonomou, K., Favez, O., Liakakou, E., Markaki, Z., Cachier, H., and Mihalopoulos, N.: Long-term measurements of carbonaceous aerosols in the Eastern Mediterranean: evidence of long-range transport of biomass burning, *Atmospheric Chemistry and Physics*, 8, 5551–5563, doi:10.5194/acp-8-5551-2008, <http://www.atmos-chem-phys.net/8/5551/2008/>, 2008.
- 700 Stohl, A., Eckhardt, S., Forster, C., James, P., Spichtinger, N., and Seibert, P.: A replacement for simple back trajectory calculations in the interpretation of atmospheric trace substance measurements, *Atmospheric Environment*, 36, 4635 – 4648, doi:10.1016/S1352-2310(02)00416-8, <http://www.sciencedirect.com/science/article/B6VH3-46PBJBX-8/2/7d8c7b6557524176d31e8d96169cd1df>, 2002.
- 705 Tesche, M., Ansmann, A., Müller, D., Althausen, D., Engelmann, R., Freudenthaler, V., and Gross, S.: Vertically resolved separation of dust and smoke over Cape Verde using multiwavelength Raman and polarization lidars during Saharan Mineral Dust Experiment 2008, *Journal of Geophysical Research: Atmospheres*, 114, D13 202, doi:10.1029/2009JD011862, <http://dx.doi.org/10.1029/2009JD011862>, 2009.

- Vaughan, M. A., Powell, K. A., Winker, D. M., Hostetler, C. A., Kuehn, R. E., Hunt, W. H., Getzewich, B. J., Young, S. A., Liu, Z., and McGill, M. J.: Fully Automated Detection of Cloud and Aerosol Layers in the CALIPSO Lidar Measurements, *J. Atmos. Oceanic Technol.*, 26, 2034–2050, doi:10.1175/2009JTECHA1228.1, <http://dx.doi.org/10.1175/2009JTECHA1228.1>, 2009.
- 710 Vaughan, M. A., Garnier, A., Liu, Z., Josset, D., Hu, Y., Lee, K.-P., Hunt, W., Vernier, J.-P., Rodier, S., Pelon, J., and Winker, D.: Chaos, consternation and CALIPSO calibration: new strategies for calibrating the CALIOP 1064 nm Channel, in: *Proceedings of the 26th Int. Laser Radar Conf.*, Porto Heli, Greece, pp. 39–55, Alexandros Papayannis, University of Athens, Greece, 2012.
- 715 Warneke, C., Bahreini, R., Brioude, J., Brock, C., de Gouw, J., Fahey, D., Froyd, K., Holloway, J., Middlebrook, A., Miller, L., Montzka, S., Murphy, D., Peischl, J., Ryerson, T., Schwarz, J., Spackman, J., and Veres, P.: Biomass burning in Siberia and Kazakhstan as an important source for haze over the Alaskan Arctic in April 2008, *Geophys. Res. Lett.*, 36, L02 813, doi:10.1029/2008GL036194, 2009.
- 720 Warneke, C., Froyd, K. D., Brioude, J., Bahreini, R., Brock, C. A., Cozic, J., de Gouw, J. A., Fahey, D. W., Ferrare, R., Holloway, J. S., Middlebrook, A. M., Miller, L., Montzka, S., Schwarz, J. P., Sodemann, H., Spackman, J. R., and Stohl, A.: An important contribution to springtime Arctic aerosol from biomass burning in Russia, *Geophys. Res. Lett.*, 37, L01 801, doi:10.1029/2009GL041816, <http://dx.doi.org/10.1029/2009GL041816>, 2010.
- 725 Winker, D. M., Vaughan, M. A., Omar, A., Hu, Y., Powell, K. A., Liu, Z., Hunt, W. H., and Young, S. A.: Overview of the CALIPSO Mission and CALIOP Data Processing Algorithms, *J. Atmos. Oceanic Technol.*, 26, 2310–2323, doi:10.1175/2009JTECHA1281.1, <http://dx.doi.org/10.1175/2009JTECHA1281.1>, 2009.
- Young, S. A.: Analysis of lidar backscatter profiles in optically thin clouds, *Appl. Opt.*, 34, 7019–7031, doi:10.1364/AO.34.007019, <http://ao.osa.org/abstract.cfm?URI=ao-34-30-7019>, 1995.