



Black carbon emissions from Russian diesel sources

M. Evans et al.

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Black carbon emissions from Russian diesel sources: case study of Murmansk

M. Evans¹, N. Kholod¹, V. Malyshev², S. Tretyakova³, E. Gusev², S. Yu¹, and A. Barinov²

¹Joint Global Change Research Institute, Pacific Northwest National Laboratory, 5825 University Research Court, Suite 3500, College Park, MD 20740, USA

²Department of Transport and Energy, Murmansk State Technical University, Murmansk, Russian Federation

³Department of Environment, Murmansk State Technical University, Murmansk, Russian Federation

Received: 25 November 2014 – Accepted: 12 January 2015 – Published: 3 February 2015

Correspondence to: M. Evans (m.evans@pnnl.gov)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Black carbon (BC) is a potent pollutant because of its effects on climate change, ecosystems and human health. Black carbon has a particularly pronounced impact as a climate forcer in the Arctic because of its effect on snow albedo and cloud formation. We have estimated BC emissions from diesel sources in Murmansk Region and Murmansk City, the largest city in the world above the Arctic Circle. In this study we developed a detailed inventory of diesel sources including on-road vehicles, off-road transport (mining, locomotives, construction and agriculture), fishing and diesel generators. For on-road transport, we conducted several surveys to understand the vehicle fleet and driving patterns, and, for all sources, we also relied on publicly available local data sets and analysis. We calculated that BC emission in Murmansk Region were 0.40 Gg in 2012. The mining industry is the largest source of BC emissions in the region, emitting 70 % of all BC emissions because of its large diesel consumption and absence of emissions controls. On-road vehicles are the second largest source emitting about 12 % of emissions. Old heavy duty trucks are the major source of emissions. Emission controls on new vehicles limit total emissions from on-road transportation. Vehicle traffic and fleet surveys show that many of the older cars on the registry are lightly or never used. We also estimated that total BC emissions from diesel sources in Russia were 56.7 Gg in 2010, and on-road transport contributed 55 % of diesel BC emissions. Agricultural machinery is also a significant source Russia-wide, in part because of the lack of controls on off-road vehicles.

1 Introduction

Black carbon (BC) is a potent pollutant, with a global warming potential 680 times that of CO₂ (on a 100 year basis). It also contributes to adverse impacts on human health, ecosystems and air visibility. In particular, it is associated with respiratory and cardiovascular effects, as well as premature death. BC is the product of incomplete combus-

Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tion, resulting in small, light-absorbing particles of 2.5 microns or less (or to state this in another way, BC is a major component of $PM_{2.5}$). Diesel and biomass combustion are both important global sources of BC and $PM_{2.5}$ emissions. Black carbon has a particularly pronounced impact as a climate forcer in the Arctic because of its effect on snow albedo and cloud formation (EPA, 2012).

This article provides a detailed inventory of BC emissions from diesel sources in Russia's Murmansk Region. Murmansk City is the largest city in the world above the Arctic Circle. Russian BC emissions are poorly understood in general (Stohl, 2013); this represents an important gap in our understanding of BC emissions and global BC forcing because Russia is by far the largest Arctic state in terms of territory. Bond et al. (2004 and 2013) provide an overview of global emissions of black carbon and their forcing (Bond et al., 2004, 2013). The US Department of Agriculture estimates BC emissions from agricultural burning in Russia (USDA, 2012). McCarty et al. (2012) estimate the range of average annual BC emissions from cropland burning in Russia at 8.90 Gg, based on agricultural statistics. Cheng (2014) estimates the likely geographic distribution of Russian black carbon emission sources.

Diesel is an important source of emissions globally, for example, the US EPA Report to Congress on Black Carbon indicates that nearly 50% of BC emissions in the United States came from mobile diesel engines in 2005 (EPA, 2012). Russia has several trends that affect its diesel consumption and emissions in the transport sector. Diesel is growing as a transportation fuel. Road traffic has grown rapidly in Russia in the past decade, linked to economic growth and growing demand for cars. The popularity of diesel light-duty vehicles has grown: many higher class or sports utility vehicles that perform well in snow rely on diesel. In Murmansk, we found that 12% of light-duty passenger vehicles used diesel, which is somewhat higher than older estimates Russia wide. Freight transport has also been growing in Russia. At the same time, Russia has European standards for limiting particulate emissions from on-road vehicles: currently, new or imported vehicles must be at least Euro 4. Euro 4 vehicle regulations require emissions that are 20–30 times lower than vehicles with no controls (e.g., Euro 0). In

the past year, two of the largest bus companies in Murmansk Region began to upgrade their bus fleets, retiring old Euro 0 buses and replacing them with Euro IV and Euro V buses; our inventory base year (2012) predates this change.

Russia has also adopted European standards for fuel quality, which is important because emissions controls will not operate properly when diesel has high sulfur content. Russia has not introduced fuel quality standards as rapidly as its vehicle standards, so currently, three types of diesel are available on the market in Murmansk: Euro 3, 4 and 5. In 2013, Euro 5, with a maximum sulfur content of 10 ppm, accounted for 52 % of Russian diesel production for the domestic market while the share of Euro 4 was 18 % and Euro 3 was 26 % (Novak, 2014).

Russia has no requirements for emission controls on off-road vehicles, so off-road vehicles, particularly in open-pit mines in Murmansk Region, represent a major source of black carbon emissions. While Russia has considered adopting European standards for off-road vehicles, it has not yet done so. At the same time, as with on-road transportation, we found evidence that some off-road vehicles in Russia exceed current requirements.

Regarding rail emissions, most Murmansk rail operates on electricity. Diesel locomotives operate in freight depots and within industrial facilities. Diesel locomotives in Murmansk do not appear to have controls. Likewise, we did not find evidence that diesel generators typically have controls, and there are no regulations requiring such controls.

We also assessed emissions from the large Murmansk fishing fleet. Despite the size of the fleet, it does not account for a large share of emissions in Murmansk Region. Most of the large fishing vessels registered in Murmansk rarely if ever call in to Murmansk Port, based on port registries.

By design and because of sensitivities and data availability, we did not include military consumption or consumption from commercial shipping in our analysis. The military likely represents an important source of consumption; commercial shipping, on the

Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



other hand, primarily relies on heavy fuel oil, not diesel, and most of the ships quickly leave Russian territorial waters.

The impact of regulations in reducing emissions is quite clear based on our analysis in Murmansk. Without regulation of vehicles and fuel, emissions would be substantially higher. Likewise, off-road vehicles and other sources would be significantly lower if emission controls were obligatory. For example, EPA calculates the effect of emission regulations of off-road vehicles in the US and estimates that BC emissions will decrease by 92 % between 2005 and 2030 as a result of emission regulations (EPA, 2012).

2 Methodology

Our approach to estimating BC emissions involved combining fuel consumption and activity data with emission factors, which is consistent with the literature (Bond et al., 2004; Klimont et al., 2002; EPA, 2012; EEA, 2009, 2013; Streets et al., 2004). Since measured BC emission factors from Russian diesel sources are not available, we estimated BC emissions from PM emissions and then apply a speciation ratio to estimate BC emissions. We used similar methods to estimate organic carbon (OC) emissions.

We applied different methodologies to different fuel combustion technologies. Wherever possible, we used Russian methodologies or emission factors for PM; for example, we used both Russian and European emission factors to estimate emissions from on-road vehicles; the Russian methodologies included emission factors for the typical vehicle fleet on Russian roads (NIIAT, 2008a, b), though by international comparison, some of the Russian cold start emission factors seemed quite low. For most other sources, we used emission factors from the European Environment Agency (EEA, 2013). Additional detail on our methodology can be found in (Evans et al., 2012).

We collected detailed bottom up activity data from several sources, depending on the needs of the emission calculation methodology. We collected extensive primary data on road traffic in Murmansk (see Table 2 for details).

Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Regarding off-road vehicles, we used statistical data as well as public information from annual corporate reports and other public sources. For power generators, we received a detailed list of the largest off-grid diesel generators in Murmansk Region, and supplemented this with analysis comparing population centers with the power grid and statistics on fuel use. We also relied on regional statistical data about non-transport diesel consumption by different sectors of the economy. Regarding the fishing fleet, we used public data from Russian ship registries and port calls. We only counted the fraction of fishing vessel emissions corresponding to the time these vessels spent in Russian territorial waters.

3 Analysis of fuel consumption in Murmansk Region

We reviewed the official statistical data on diesel consumption in Murmansk Region, which include annual summary data on consumption and stock changes by broad categories, and a breakdown of enterprise consumption for transport and non-transport needs organized by economic activity. The summary data and the more detailed data appear to have some methodological differences, and the summary data appear to include different categories across different years, causing major swings in the total reported fuel use. Because of these factors, we also estimated consumption by sector using bottom up calculations where possible. Except in the case of mines, statistical data were significantly different from our bottom-up estimates.

In Table 1 below, we provide a consolidated estimate of diesel use in Murmansk Region in 2012.

The Supplement provides more details: Table S1 provides additional details on our bottom-up fuel calculation for on-road transport; Table S9 highlights these calculations for mines, and Tables S16 and S17 estimate fuel use for fishing and diesel generators, respectively.

Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.2 Emissions estimates

We used several methodologies to estimate emissions in the city and the region. We reviewed several Russian methodologies, including two prepared by the Scientific Research Institute for Automobile Transport (NIIAT, 2008a, b), as well as the European Environmental Agency methodology, COPERT (Emisia, 2011). The NIIAT methodologies use Russian-specific emission factors for PM_{2.5} based on the average fleet of vehicles of each ecological class on Russian roads. At the same time, the Russian methodologies have much lower emission factors for cold starts in small vehicles than other international methodologies. While some Russian drivers warm their cars before they begin driving, which reduces emissions from cold starts, without survey data measuring cold start emissions more precisely, we decided it would be more consistent with inventories elsewhere to use European emission factors for cold starts, particularly given the cold Russian climate. Thus, we used COPERT with Russian emission factors for the hot operation stage to reflect the Russian vehicle fleet. Figure 1 summarizes our emission estimates by vehicle type using COPERT with Russian emission factors.

As a cross-check, we also calculated emissions with the Russian methodologies. We used the NIIAT methodology for large cities. We also used the NIIAT universal methodology, which factors in low usage of registered vehicles in Russia in its formulas.

Table 3 presents a summary of total vehicle emissions in the city using each of the methodologies. Additional details on the emissions are available in Table S5.

The results in Table 3 clearly show that one should be very careful in using registry data for emission estimates. The difference between estimated emissions from the observed fleet is 4.7 times smaller than the potential emissions from the fleet of all registered vehicles, as the Russian vehicle registries likely contain many vehicles not actually in use.

We also calculated total road transport emissions in Murmansk Region using the NIIAT universal methodology (NIIAT, 2008b). This methodology is simpler and designed for use with limited vehicle activity data; estimating emissions at the regional level pro-

Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



operate a wide range of machinery, including excavators, bulldozers, loaders, drilling equipment and other machinery. On average, excavators and bulldozers operate 7270 and 6660 h per year, respectively. The mines also use supplementary, smaller on-road trucks with payloads from 13 to 45 t.

Statistical data in the region indicate that mining companies consumed 139 013 t of diesel fuel in 2012. We also cross-checked this data through bottom-up estimates of fuel use in the largest mines. The results of cross-checking showed that the statistical data and bottom-up calculations match closely (with a difference of less than 1 %).

Russia does not have emission regulations for off-road vehicles but often uses foreign-made, off-road vehicles and equipment. Thus, we have used both US EPA and European Environment Agency information about emission requirements for off-road vehicles. Table S7 shows PM emission requirements in the US and Europe.

The extent of controls is one of the important uncertainties regarding emission estimates from the mining sector. Since there are no emission control requirements, the mining vehicles may not meet even Tier 1 requirements. Based on information from Cummins, 88 % of the large, Cummins-powered, BELAZ mining trucks have no controls on their engine exhaust and the remaining 12 % meet EPA Tier 1 requirements (Mueller, 2014). A smaller population of Caterpillar and Komatsu trucks meets Tier 1 or Tier 2 requirements (Table S11).

The $PM_{2.5}$ emission factor for off-road, industrial mobile sources and machinery without emission controls is 3.551 g kg^{-1} fuel and the emission factor for equipment with some controls is 0.967 g kg^{-1} fuel (EEA, 2013).

We estimated that $PM_{2.5}$ emissions in the mining industry in Murmansk Region are 450.5 t per year. The speciation BC/PM ratio is 0.62. Total BC and OC emissions in the mining industry in Murmansk Region estimated to be 279.3 and 83.8 t per year, respectively.

Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5.2 Locomotives

Diesel locomotives are only in limited use in Murmansk Region because all the main railroads are electrified. According to data from the Murmansk statistical office, diesel locomotives at the Murmansk branch of Russian Railways consumed 21 200 t of diesel in 2012 (GKS, 2012).

Diesel locomotives in Russia do not have any emission controls. Some of the locomotives in Murmansk Region are more than 30 years old. Since we have limited information on the activity of the small line haul and switch locomotives, the only way to estimate BC emissions is to use the fuel consumption method.

The emission factor for $PM_{2.5}$ of switch locomotives is 1.44 grams per kilogram of fuel. The speciation ratio for BC/ $PM_{2.5}$ for locomotives is 0.73. Thus, locomotives in Murmansk Region emitted 30.5 t of $PM_{2.5}$, including 22.3 t of BC and 4.5 t of OC.

5.3 Construction and road management

This sector includes building construction and road management. According to official statistics, the building construction industry used 3205 t and road management companies used 865 t of diesel fuel for off-road vehicles, machinery and equipment in 2012.

Building construction is stagnant in Murmansk Region. The region's population is declining and the formerly powerful construction industry is deteriorating. The vast majority of equipment in the construction industry is very old. There are over 1800 pieces of equipment and more than 50 % of equipment and machinery need replacement (see Table S12 for details). We assume that 90 % of equipment has no emission controls and 10 % has some controls.

We used EMEP-EEA emission factors (EEA, 2013) for off-road vehicles in the construction industry, e.g. $4.038 \text{ g } PM_{2.5} \text{ kg}^{-1}$ fuel for vehicles without controls and $0.967 \text{ g } \text{kg}^{-1}$ fuel for equipment with some controls. Hence, off-road building construction vehicles in Murmansk Region emitted 12.7 t of $PM_{2.5}$, 9.8 t BC and 2.0 t OC.

Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The road management sector includes minor road reconstruction and snow removal. Murmansk City is located on the shore of the Barents Sea and the level of precipitation is quite high. On average, there is snow on the ground 180–200 days per year. The snow removal fleet was significantly updated recently with Russian-made, multifunctional vehicles and off-road vehicles, including new tractors and graders, do not have any emission controls.

Similarly to construction, we have to exclude on-road vehicles from the emission calculations. The emission factor for off-road machinery without emission controls in this sector is $3.551 \text{ gPM}_{2.5} \text{ kg}^{-1} \text{ fuel}$ (EEA, 2013). Off-road vehicles in this sector in Murmansk Region emitted 2.8 t of $\text{PM}_{2.5}$, 2.2 t BC and 0.4 t OC.

Total emissions from off-road vehicles and equipment in building construction and road management sector were 15.6 t of $\text{PM}_{2.5}$, 12.0 t BC and 2.4 t OC.

5.4 Agriculture

Over 90 % of Murmansk Region lies above the Arctic Circle and agriculture is not well developed. The agricultural machinery in the region is Russian-made with a small fraction of foreign-made equipment; 62 % of agricultural machinery is older than 10 years.

According to regional statistics agricultural enterprises in Murmansk Region consumed 1344 t of diesel in 2012. The emission factor for agricultural equipment without emission controls is $3.755 \text{ gPM}_{2.5} \text{ kg}^{-1} \text{ fuel}$ assuming no controls (EEA, 2013). We thus estimated total PM emissions at 5.1 t $\text{PM}_{2.5}$. Using EPA speciation ratio (EPA, 2012), emissions from agricultural equipment in Murmansk region in 2012 estimated to be 3.9 t of BC and 0.8 t of OC.

6 Fishing

Fishing is an important part of Murmansk's economy. The fishing industry in Murmansk Region provides 16 % of Russia's total fish catch. Fishing companies in Murmansk Re-

Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



plied an emission factor of 0.35 g kWh^{-1} (Corbett, 2010) to calculate BC emissions within Russian territorial waters from fishing vessels. We assumed that all fishing vessels use diesel. (According to the EEA emissions inventory guidebook, only 3.8 % of fishing vessels use both diesel and bunker fuel oil). We estimate that these large and medium fishing vessels emitted 4.3 t of BC and 0.9 t of OC in 2012.

In addition, there are about 100 small fishing ships. Detailed registries and other data about installed engine capacity and hours of operation are not available, so in consultation with local fishing and marine experts, we assumed that the average engine capacity is 50 kW, engine load is 60 %, the boats sail 800 h per year. The total BC emissions by small fishing boats were 840 kg per year. Total BC emissions from all types of fishing vessels in Murmansk Region territorial waters were 5117 kg.

We also prepared bottom-up estimates of fuel use, based on information about rated engine power, hours of operation and specific fuel consumption (g fuel kWh^{-1}). The specific fuel consumption is $203 \text{ g diesel kWh}^{-1}$ (EEA, 2013). The fuel consumption by large and medium ships during their travel within Russian territorial waters is 2481 t per year. The fuel consumption by small boats is 487 t year^{-1} .

7 Diesel generators

We found several types of diesel generators and heaters in Murmansk Region. The largest category in terms of fuel use and emissions is generators and heaters that small market shops and service providers operate in settled areas. The next largest category includes off-grid generators that operate for a large portion of the year, typically up to 12 h a day.

We found the least data for the very small generators and heaters used in commerce and services – the government does not appear to regulate or keep statistics on these small generators. The data quality regarding diesel generators is very low and the uncertainty is very high. In total, government statistics show that non-transport diesel use from these sectors was 7100 t in 2012. We also verified the existence of such

generators by looking at the number of dealers selling diesel generators in Murmansk. With an emission factor for diesel generators of 4.0 g PM kg^{-1} fuel use and a BC/PM ratio of 0.74 for this category, we assumed that such small generators and heaters emitted 21.0 t of BC and 4.2 t OC in 2012.

Regarding off-grid generators, it is important to note that the majority of Murmansk Region's urban and rural energy consumers receive their power from the Kola Power Grid. Several dozen settlements in the region lack access to centralized electricity supply, due to their remote locations; instead they rely on diesel generators (Minin, 2012). The largest villages without centralized electricity supply receive diesel subsidies. Table S17 shows the capacity of these subsidized diesel generators and their annual fuel consumption. In total, according to the Development Strategy for Energy Savings in Murmansk Region, there were 80 settlements without centralized electricity supply in 2009. About 150 village diesel generators with a total capacity of 3.8 MW provided electricity to these settlements (Government of Murmansk Region, 2009). We used information about fuel consumption and power capacity of generators with subsidized fuel and proportionally calculated the possible total fuel consumption by this category of generators. Using bottom-up calculations, we estimated that off-grid generators consume 1700 t of diesel per year. We further estimate that off-grid generators in Murmansk Region emitted 5.2 t of BC and 1.0 t of OC in 2012.

The total BC emissions from diesel generators in the region estimated to be 26.3 t and OC emissions were 5.3 t.

8 Uncertainty analysis

Uncertainties exist in emission factors, activity data and emission controls; we used multiple approaches to estimate and reduce uncertainties of the BC emissions inventory. This could help us validate the inventory estimates, choose appropriate methodological approaches and improve the accuracy of the results (IPCC, 2006). This could also help peer reviewers understand the reliability of our inventory estimates. We used

Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



five methods to assess and minimize uncertainties (EEA, 2013; IPCC, 2006, 2000), including:

- multiple approaches to collecting and validating activity data;
- literature and other documented data for cross-checks;
- cross-checks of bottom-up activity data and fuel allocation;
- error propagation; and
- expert judgment.

We derived aggregate uncertainties of the emissions inventory based on the error propagation method. We combined uncertainties of emission factor and activity data by source category, and then combined uncertainties by source category to estimate overall uncertainty of the inventory (IPCC, 2006). For emission factors, we use uncertainties and confidence intervals reported by previous studies (e.g. emission factors from EPA, EEA, Russian methodologies and journal articles). Uncertainties in activity data are primarily assessed based on expert judgment.

For major sources of BC emissions, we also used cross-checks to assess sectoral uncertainties. For on-road emissions, we checked our results against multiple methodologies, and there is 19 % difference between estimated emissions from COPERT with NIAT emission factors and COPERT with COPERT emission factors.

The largest uncertainty in mining lies in assumptions on emission controls and fuel use (Table S19 and S20). Uncertainty in emissions from mining vehicles appears to be the greatest. Uncertainty about Tier distribution could significantly change the results of our emissions calculation given the significant fuel consumption in the mining industry.

9 Simple estimate of Russian diesel emissions

According to IEA data, Russia consumed 23.3 million t of diesel in 2010 (IEA, 2012). On-road transport accounted for 12.7 million t of diesel, while agriculture and forestry

Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



consumed an additional 2.8 million t and industry 2.6 million t of diesel. All other sectors combined consumed an additional 2.9 million t of fuel.

Since on-road transport is the largest consumer of diesel we conducted a more detailed analysis of BC emissions by on-road vehicles. We simply applied fuel-based emission factors to all other sectors.

According to the Federal State Statistics Service of the Russian Federation, there were 5 181 200 diesel vehicles in Russia in 2010. NIIAT conducted bottom-up calculations of fuel consumption by on-road vehicles in Russia and estimated it at 17.3 million t per year. Table S20 shows fuels consumption by different types of vehicles and Table S21 shows diesel fleet distribution by ecological class. We used these estimates to calculate BC emissions from on-road transport; for all other sectors we used IEA data.

We calculated PM emissions by using NIIAT fuel-based emissions factors (NIIAT, 2008) and applied the BC/PM speciation ratio (EPA, 2012) to determine BC emissions. Table 4 shows the results of the BC emissions calculations from on-road diesel vehicles in Russia in 2012.

We estimated total BC emissions from on-road diesel vehicles in Russia in 2010 at 31 117 t, and total OC emissions at 4588 t. The vast majority of BC emissions (68 %) came from Euro 0 trucks.

We cross-checked this result with the EEA methodology using bulk emissions factors (EEA, 2013). The total BC emissions are 34 226 t (the difference is 9 %). These results are similar to those presented in the EPA Report to Congress on Black Carbon (EPA, 2012). According to EPA estimates, BC emissions from transport (including aircrafts and marine shipping) in Russia were 32 Gg in 2000.

BC emissions from diesel sources in agriculture and forestry were 8180 t, industrial emissions were 5610 t (including 2536 t from mining) and while emissions from other sectors combined were 11 818 t. We estimated total BC emissions in Russia from diesel combustion in 2010 at 56 726 t (56.7 Gg).

Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



10 Conclusions

We conducted a detailed, bottom-up assessment of emissions from diesel combustion in Murmansk Region, based on surveys of vehicles, traffic and data collection regarding other significant sources (see Table 5).

We also conducted an initial estimate of Russian emissions from diesel combustion. In both Murmansk and Russia, on-road transportation is a large source of BC emissions. Within this category, Euro 0 trucks make up the vast majority of emissions. This reflects the fact that Russia now has requirements for emission controls on new vehicles, resulting in comparatively low emissions for cars and most new trucks and buses. We also found that many registered vehicles, particularly older vehicles, are driven infrequently based on parking lot and traffic video surveys, which is consistent with the literature. Surprisingly, we found that regional statistic on fuel use for on-road transportation indicate significantly lower consumption than our bottom-up estimates of fuel use in this category. In Murmansk Region, the largest category of emissions is off-road vehicles, in particular mining. In Russia as a whole, agriculture represents the second largest diesel BC source. In both these cases, the high emissions are linked to the absence of control technologies and the lack of emission standards for off-road vehicles. Off-road vehicles represent an important opportunity for reducing emissions, for example, with emission standards for new vehicles and engines.

The Supplement related to this article is available online at doi:10.5194/acpd-15-3257-2015-supplement.

Acknowledgements. The authors are grateful for research support provided by the US Environmental Protection Agency, Office of International and Tribal Affairs (grant no. X4-83527901) and the US Department of State. Battelle Memorial Institute operates the Pacific Northwest National Laboratory for the US Department of Energy under contract DE-AC05-76RL01831. We thank

ACPD

15, 3257–3284, 2015

Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the members of the Technical Steering Group for their helpful comments and suggestions. The views and opinions expressed in this paper are those of the authors alone.

References

- Bond, T. C., Streets, D. G., Yarber, K. F., Nelson, S. M., Woo, J.-H., and Klimont, Z.:
5 A technology-based global inventory of black and organic carbon emissions from combustion, *J. Geophys. Res.-Atmos.*, 109, D14203, doi:10.1029/2003jd003697, 2004.
- Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C.,
10 Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: a scientific assessment, *J. Geophys. Res.-Atmos.*, 118, 5380–5552, doi:10.1002/jgrd.50171, 2013.
- Cheng, M.-D.: Geolocating Russian sources for Arctic black carbon, *Atmos. Environ.*, 92, 398–
15 410, doi:10.1016/j.atmosenv.2014.04.031, 2014.
- Committee for the Fishery Complex of Murmansk Region: The state program of Murmansk Region “The Development of Fishery Complex of Murmansk region for 2014–2020” (in Russian), Murmansk, Russia, 2013.
- Corbett, J. J., Lack, D. A., Winebrake, J. J., Harder, S., Silberman, J. A., and Gold, M.: Arctic shipping emissions inventories and future scenarios, *Atmos. Chem. Phys.*, 10, 9689–9704,
20 doi:10.5194/acp-10-9689-2010, 2010.
- EEA: EMEP/EEA Air Pollutant Emission Inventory Guidebook – 2009, European Environment Agency, Copenhagen, Denmark, 2009.
- EEA: EMEP/EEA Air Pollutant Emission Inventory Guidebook – 2013, European Environment Agency, Copenhagen, Denmark, 2013.
- Emisia: COPERT 4 (Computer programme to calculate emissions from road transport), Prepared for the European Environment Agency (EEA), available at: <http://www.emisia.com/content/copert-download> (last access: 15 July 2014), 2011.
- EPA: Report to Congress on Black Carbon, US Environmental Protection Agency, Washington
25 DCEPA-450/R-12-001, 2012.
- 30

Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Black carbon
emissions from
Russian diesel
sources**

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Evans, M., Kholod, N., Yu, S., Tretyakova, S., Gusev, E., and Malyshev, V.: Understanding black carbon from diesel sources in Russia: methodology for preparing an emissions inventory in Murmansk Region, Battelle Memorial Institute, 2012.

GKS: Information about fuel use in Murmansk Region, Federal State Statistical Service Moscow, 2012.

Government of Murmansk Region: Strategy of Energy Saving in Murmansk Region, Murmansk, Russia, 2009.

IEA: Fuel balances of non-OECD countries, International Energy Agency, Paris, 2012.

IPCC: IPCC good practice guidance and uncertainty management in national greenhouse gas inventories, Institute for Global Environmental Strategies, Hayama, Japan, 4-88788-000-6, 2000.

IPCC: 2006 IPCC guidelines for national greenhouse gas inventories, Institute for Global Environmental Strategies, Hayama, Japan, 4-88788-032-4, 2006.

Klimont, Z., Cofala, J., Bertok, I., Amann, M., Heyes, C., and Gyarmas, F.: Modeling particulate emissions in Europe. A framework to estimate reduction potential and control costs, International Institute for Applied Systems Analysis (IIASA), Laxenburg, Australia, 2002.

McCarty, J. L., Ellicott, E. A., Romanenkov, V., Rukhovitch, D., and Koroleva, P.: Multi-year black carbon emissions from cropland burning in the Russian Federation, *Atmos. Environ.*, 63, 223–238, doi:10.1016/j.atmosenv.2012.08.053, 2012.

Minin, V.: Economic aspects of small-scale renewable energy development in remote settlements of the Kola Peninsula, Bellona Murmansk, Russia, 2012.

Mining Magazine: BELAZ at Apatity, Mining Magazine, October, 47, 2007.

Mueller, R.: Personal correspondence with Ralf Mueller, Territory Manager – Mining Business, Europe, Middle East and CIS; e-mail response to technical questions about Cummins products, 2014.

Murmansk Fishing Port: Information System on State Port Control, Murmansk, 2014.

NIAT: Calculation instruction (methodology) for emission inventory from vehicles on the territory of the largest cities (in Russian), Scientific Research Institute of Automobiles and Transportation, Moscow, Russia, 2008a.

NIAT: Calculation instruction (methodology) for emission inventory from vehicles into the air (in Russian), Scientific Research Institute of Automobiles and Transportation, Moscow, Russia, 2008b.

Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Novak, A. V.: The results of activity of the fuel and energy complex of Russia, The Ministry of Energy of the Russian Federation, Moscow, 2014.

Petrovich, A. A., Belyavskiy, D. A., and Garavskiy, A. O.: BELAZ trucks at open pit mines in Russia (in Russian), Mining J., 1, 75–77, 2013.

5 Stohl, A., Klimont, Z., Eckhardt, S., Kupiainen, K., Shevchenko, V. P., Kopeikin, V. M., and Novigatsky, A. N.: Black carbon in the Arctic: the underestimated role of gas flaring and residential combustion emissions, Atmos. Chem. Phys., 13, 8833–8855, doi:10.5194/acp-13-8833-2013, 2013.

10 Streets, D. G., Bond, T. C., Lee, T., and Jang, C.: On the future of carbonaceous aerosol emissions, J. Geophys. Res.-Atmos., 109, D24212, doi:10.1029/2004jd004902, 2004.

The Russian Maritime Register of Shipping: The Register Book, On-line database, available at: <http://info.rs-head.spb.ru/webFS/regbook/regbookVessel?In=en> (last access: 29 July 2014), 2014.

15 USDA: Developing Options for Avoiding, Reducing or Mitigating Agricultural Burning that Contributes to Black Carbon Deposition in the Arctic, United States Department of Agriculture, Washington, D. C., 2012.

Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Estimated diesel consumption by sector in Murmansk Region, 2012.

Activity	Diesel use (t)
On-road transport*	65 100
Mines	139 000
Locomotives	21 200
Construction	4100
Agriculture	1300
Diesel generators, including:	8800
Small generators for commerce and services*	7100
Off-grid generators*	1700
Fishing (in Russian territorial waters), including:	3000
large and medium vessels*	2500
Small boats*	500
Total	242 500

* – Bottom up calculations. The other numbers come from regional statistics. This table does not consider marine shipping and military fuel use.

Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Main data sources on vehicle fleet and activity.

Type of data	Description	Notes
Vehicle fleet	Basic registry information on each vehicle registered in Murmansk Region from Avtostat	We categorized the vehicles by make, model and age, and then assessed diesel use and ecological class based on manufacturer data of the models.
Passenger cars in use	Parking lot surveys at several locations throughout central and suburban Murmansk City	The surveys provided data on the vehicle models actually in use. We assessed the models for age and ecological class as we did with the Avtostat data.
Passenger cars in use and odometer readings	Database of vehicle inspection station on MSTU campus	This provided additional data on vehicles on the roads as well as their age and odometer readings (average km travelled per year).
Traffic intensity	Video surveys	MSTU conducted video surveys to count total traffic by vehicle type (cars, light-duty vehicles, buses and trucks) on different road categories in both central and suburban Murmansk City. Surveys covered different hours of the day.
Road categories and length	Municipal data on road categories and lengths	We used this data to help select road segments for the video surveys and to correlate the video survey data with the rest of the city roads by category.
Road speed and grade	GPS logger data	We used specialized GPS data loggers to track road speed by road type at different times of day. The loggers also provided data on road grade. In addition, we used data from the Yandex traffic service to assess road speed.

The Supplement provides additional details on several of these data sets.

Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. BC emissions in Murmansk City from on-road transport, different methodologies, t yr^{-1} .

	COPERT with NIIAT EFs	NIIAT universal	NIIAT for large cities	COPERT with NIIAT EFs (full registry)
Cars	3.9	2.5	3.0	6.1
LDV	1.7	1.1	1.1	14.4
Trucks	3.9	3.9	2.7	28.7
Buses	2.2	2.2	1.0	5.7
Total	11.7	9.7	7.8	54.9

Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 4. BC emissions from on-road diesel vehicles in Russia in 2010, t.

Ecological class/ vehicle type	Euro 0	Euro 1	Euro 2	Euro 3 and higher
Cars	203	0	62	365
Trucks	21 203	1029	2287	1871
Buses	2973	245	511	368

Source: Authors' calculations.

Black carbon emissions from Russian diesel sources

M. Evans et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 5. PM_{2.5}, BC and OC emissions in Murmansk Region, 2012 (t).

Activity	PM _{2.5}	BC	OC
On-road transport in Murmansk Region	98.9	53.7	36.2
Mines	450.5	279.3	83.8
Locomotives	30.5	22.3	4.5
Construction	15.6	12.0	2.4
Agriculture	5.0	3.9	0.8
Diesel generators	35.2	27.1	5.4
Fishing (in Russian waters)	16.5	5.1	1.0
Total	652.3	403.4	134.1




Figure 1. Cold start and hot stage BC emissions in Murmansk City by vehicle type, 2012 (in t).

Black carbon emissions from Russian diesel sources

M. Evans et al.

[Title Page](#)

[Abstract](#) | [Introduction](#)

[Conclusions](#) | [References](#)

[Tables](#) | [Figures](#)

[◀](#) | [▶](#)

[◀](#) | [▶](#)

[Back](#) | [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Black carbon emissions from Russian diesel sources

M. Evans et al.




Figure 2. BC emissions in Murmansk City by ecological class and vehicle type.