

Comparison of eddy covariance and modified Bowen ratio methods

D. J. Bolinius et al.

Comparison of eddy covariance and modified Bowen ratio methods for measuring gas fluxes and implications for measuring fluxes of persistent organic pollutants

D. J. Bolinius¹, A. Jahnke^{1,2}, and M. MacLeod¹

¹Department of Environmental Science and Analytical Chemistry (ACES), Stockholm University, Svante Arrhenius väg 8, 106 91 Stockholm, Sweden

²Department Cell Toxicology, Helmholtz Centre for Environmental Research (UFZ), Permoserstr. 15, 04318 Leipzig, Germany

Received: 7 September 2015 – Accepted: 4 November 2015 – Published: 20 November 2015

Correspondence to: D. J. Bolinius (damien.bolinus@aces.su.se)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Semi-volatile persistent organic pollutants (POPs) cycle between the atmosphere and terrestrial surfaces, however measuring fluxes of POPs between the atmosphere and other media is challenging. Sampling times of hours to days are required to accurately measure trace concentrations of POPs in the atmosphere, which rules out the use of eddy covariance techniques that are used to measure gas fluxes of major air pollutants. An alternative, the modified Bowen ratio (MBR) method, has been used instead. In this study we used data from FLUXNET for CO₂ and water vapor (H₂O) to compare fluxes measured by eddy covariance to fluxes measured with the MBR method using vertical concentration gradients in air derived from averaged data that simulates the long sampling times typically required to measure POPs. When concentration gradients are strong and fluxes are unidirectional, the MBR method and the eddy covariance method agree within a factor of 3 for CO₂, and within a factor of 10 for H₂O. To remain within the range of applicability of the MBR method field, studies should be carried out under conditions such that the direction of net flux does not change during the sampling period. If that condition is met then the performance of the MBR method is not strongly affected by the length of sample duration nor the use of a fixed value for the transfer coefficient.

1 Introduction

Despite the more than decade-old global ban on the production and use of persistent organic pollutants (POPs) such as polychlorinated biphenyls (PCBs), hexachlorobenzene and several organochlorine pesticides, these chemicals are still present in the environment and continue to raise concerns due to their persistence, bioaccumulation, toxicity and potential for long-range atmospheric transport (Stockholm Convention, 2010). As the production and use of POPs continues to decline, large cities, old stocks and re-volatilization from soil are expected to become more important sources to the

ACPD

15, 32759–32777, 2015

Comparison of eddy covariance and modified Bowen ratio methods

D. J. Bolinius et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



atmosphere (Nizzetto et al., 2010). Studying the sources and fate of organic pollutants in the environment is an important prerequisite to exposure and risk assessment, and environmental fate models that calculate fluxes of pollutants between air, water, soil, vegetation and other media have proven to be a valuable tool in this respect (McKone and MacLeod, 2003). Measurements of fluxes of POPs emanating from source areas and between the atmosphere and other environmental media are needed to parameterize and evaluate the chemical fate models that are used as scientific support for international conventions on POPs (Gusev et al., 2012).

The preferred approach to measure the flux of major air pollutants between the earth's surface and the atmosphere is the eddy covariance (EC) technique (Baldocchi et al., 1988). It is based on measuring the covariance of the concentration of a pollutant and the vertical wind velocity, using data from very fast measurements (e.g. 5–10 Hz). This approach works well for compounds such as CO₂, methane, ozone and more recently also for mercury (Pierce et al., 2015), since concentrations can be measured at a high temporal resolution. However, it cannot be applied directly when studying trace-level organic micropollutants that require sampling times of at minimum several hours when using active high-volume sampling, or even several weeks when using passive samplers (Hung et al., 2013) to result in reliable, quantifiable data.

One way to estimate chemical fluxes from measurements based on sampling times of hours to days is to use the modified Bowen ratio (MBR) method (Businger, 1986). The MBR method is based on the assumption that turbulent atmospheric transport occurs indiscriminately for chemicals, heat and other scalar quantities that can be described entirely by their magnitude without reference to direction. It can be used to measure the flux of a chemical pollutant (x) from measurements of its concentration at two heights and the measured transfer coefficient of another scalar such as heat (y) over the same height interval (Meyers et al., 1996; Eq. 1):

$$F_x = -K_y \cdot \frac{\Delta C_x}{\Delta Z} \quad (1)$$

Comparison of eddy covariance and modified Bowen ratio methods

D. J. Bolinius et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Comparison of eddy covariance and modified Bowen ratio methods

D. J. Bolinius et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Where F_x ($\text{ng m}^{-2} \text{h}^{-1}$) is the flux of the chemical x of interest, K_y ($\text{m}^2 \text{h}^{-1}$) is the measured eddy diffusion coefficient for a scalar y over the height interval ΔZ (m) and ΔC_x (ng m^{-3}) is the measured concentration gradient of x over the height interval. The negative sign on the right hand side of the equation enforces the convention that downward fluxes have a negative sign, and upward fluxes a positive sign.

Among other applications, the MBR method has been used to measure volatilization fluxes of pesticides applied to agricultural fields (Majewski, 1999), to estimate PCB fluxes from Lake Superior to the overlying air phase (Rowe and Perlinger, 2012) and fluxes of polycyclic aromatic hydrocarbons (PAHs) above a forest canopy in Canada (Choi et al., 2008). In the study of Choi et al., air was sampled for 24 h every 3 days at different heights for a period of one month while leaves in the forest canopy were developing. The samples were analyzed for PAHs and the data was combined with the eddy diffusivity of heat (K_{heat}) determined from eddy covariance measurements from the FLUXNET network to estimate vertical PAH fluxes using the MBR method.

Our goal in this study was to test the limits of applicability of the MBR method and to evaluate its accuracy relative to the “standard” EC technique. We used data from the FLUXNET network to calculate fluxes of CO_2 and water vapor (H_2O) with the MBR method under different sampling duration scenarios and different assumptions about data availability for the eddy diffusion coefficient K_y . Thus, we took advantage of the high-frequency measurement data for CO_2 and H_2O , and used them as proxies for organic micropollutants in order to analyze the performance of the MBR method compared to the EC method. By averaging the FLUXNET data over periods that ranged from 1 h to 1 week, we simulated sampling times that are typically required to measure POPs and other organic micropollutants in air. Our approach is similar to the one used by Majewski (see Majewski, 1999), who simulated 24 h sampling periods from higher frequency data to characterize the potential for long sampling times to introduce errors to the aerodynamic profiling method.

2 Methods

2.1 Datasets

All data used in this study can be accessed freely on the web via the FLUXNET homepage (<http://fluxnet.ornl.gov/>). A figure showing the flux tower and associated instruments can be found in the Supplement Fig. S1. The dataset used in this study contains eddy flux parameters and micrometeorological measurements for the year 2009 taken at the Borden mixed deciduous forest site in Ontario, Canada (FLUXNET site code: CA-Cbo). A list of all parameters is given in the Table S1 in the Supplement. We selected measurements taken at heights of 33.3 and 40.7 m. Air sampling in the study by Choi et al. (2008) was conducted at the same site in 2003 at heights of 29.1 and 44.4 m.

Prior to any calculations, we filtered the data to remove about 25 % of the observations that were flagged as unreliable for CO₂ or H₂O. Details on the criteria for the flags can be found on the FLUXNET homepage. Common reasons for flagging are instrument malfunctions, calibration problems and outliers. All flagged data was filtered out simultaneously, such that our analysis only includes data points collected at times when data were not flagged for either CO₂ or H₂O.

On inspection of the distribution of the CO₂ and H₂O concentration gradients, it was apparent that a few outliers that had not been flagged could significantly alter the average gradient when pooling the data to simulate long sampling times. These outliers in some cases led to net flux estimations based on the MBR method that were in the opposite direction compared to the EC method. To exclude such outliers and reduce the influence of extreme values of measured parameters, the highest and lowest 2.5 % of values of the CO₂ gradient and the H₂O gradient were removed from the dataset prior to further calculations.

Comparison of eddy covariance and modified Bowen ratio methods

D. J. Bolinius et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2 Modified Bowen ratio

We used K_{heat} derived from EC flux measurements and measurements of the scalar temperature at two heights, as described in the paper by Choi et al. (2008) to specify the eddy diffusivity in the MBR method (K_y in Eq. 1). Specifically, K_{heat} was inferred from the dataset by first calculating the vertical turbulent flux of the sonic anemometer temperature $W'T'$ (K m s^{-1} , Eq. 2) from the turbulent sensible heat flux (Q in W m^{-2} or $\text{J s}^{-1} \text{m}^{-2}$), using the air density (σ_{air} in kg m^{-3}) and the gravimetric heat capacity (c_p) of air measured at 33 m height ($1005.7 \text{ J kg}^{-1} \text{K}^{-1}$). Spurious K_{heat} values less than or equal to zero were removed from the dataset as these would indicate a heat flux against the measured temperature gradient.

$$\overline{W'T'} = \frac{Q}{c_p \cdot \sigma_{\text{air}}} \quad (2)$$

K_{heat} ($\text{m}^2 \text{s}^{-1}$) was then calculated based on temperatures measured at 40.7 and 33.3 m according to Eq. (3):

$$K_{\text{heat}} = -\overline{W'T'} \frac{(40.7 \text{ m} - 33.3 \text{ m})}{(T_{40.7} - T_{33.3})} \quad (3)$$

Finally, vertical turbulent fluxes of CO_2 and H_2O were calculated using the MBR method and measured concentrations at 33 and 41.5 m averaged over different time intervals selected to represent sampling times typical for organic micropollutants, as described below. Fluxes calculated with the MBR method were compared with those measured by the EC method available from the FLUXNET dataset.

2.3 Data-analysis

To simulate sampling times typical for organic micropollutants, concentrations of CO_2 and H_2O reported as 30 min averages in the database were pooled and averaged over

periods of 1, 2, 4, 8, 24 h, 3 days (72 h), and 1 week (168 h). Fluxes during four two-month periods selected to represent each of the four seasons were then calculated from median values of these pooled data points during the entire period. Thus for example, fluxes calculated from 1 h simulated sampling times are based on the median of average vertical concentration gradients in 1 h pools measured at the same time each day over the entire 2 month period (see Fig. S2 for a visual representation). Medians were used instead of geometric means because of the presence of negative flux values. January and February represented winter, April and May spring, July and August summer and October and November fall.

We tested two approaches to specify K_{heat} in the MBR method calculations. In the first approach hourly average K_{heat} values were calculated from 30 min averages of temperature measurements reported in the database. In the second approach a geometric mean of K_{heat} was calculated for all time points across the entire period corresponding to the simulated sampling time. The first approach takes advantage of the availability of high temporal resolution information about K_{heat} at the FLUXNET site, but the second approach is likely to be common when applying the MBR since high frequency meteorological data is not always available.

The direction of flux for CO_2 can change on a diurnal basis (see Figs. 1 and S3). During the day the flux of CO_2 is often negative (i.e., downward) due to photosynthesis while during the night, plant respiration produces CO_2 and fluxes are positive (i.e. to the atmosphere). In addition, atmospheric conditions during the night are typically much more stable than during the day, resulting in a lack of large turbulent eddies and a higher contribution of additional transport mechanisms, such as horizontal advection, to the total flux. The result is that fluxes measured using EC during the night are often underestimated (Aubinet, 2008).

To understand the impact of changing directions of flux and to investigate potential underestimation of flux at night, fluxes during the day and during the night calculated with the MBR method were evaluated against EC measurements separately. Nighttime data was set to cover from 9 p.m. to 5 a.m. LT across all seasons, and daytime data was

Comparison of eddy covariance and modified Bowen ratio methods

D. J. Bolinius et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



set from 9 a.m. to 5 p.m. LT. The nighttime/daytime divisions were selected based on the shortest interval between sunrise and sunset at the site. The 8 h periods representing day and night allowed us to construct 24 h, 3 day and 1 week sampling periods by averaging a whole number of 8 h periods taken at the same time of day over multiple days. In addition to the nighttime/daytime split data, we also examined the performance of the MBR method relative to the EC method when using continuous data that ignored day/night differences.

3 Results

10 Our calculated K_{heat} values (Fig. 1) are in good agreement with values for the same site during the same time of year in 2003 (Choi et al., 2008; shown in Fig. S4). Values are close to 0 during the night and in the range of 0.0026 to 35.8 m^2s^{-1} during the day over the summer period, with 95 % of the values between 0.029 and 22.11 m^2s^{-1} .

The fluxes calculated with the MBR method are proportional to the product of K_{heat} and the concentration gradient of either CO_2 or H_2O (Eq. 1). The raw data at 30 min time resolution that was pooled and used to calculate the fluxes with the MBR method is visualized in Fig. 1.

3.2 Performance of the MBR method on continuous time series

3.3 Performance of the MBR method with hourly-resolved and fixed values of

K_{heat}

The use of either hourly-resolved data for K_{heat} or a fixed value did not significantly affect the the MBR method. A student t test comparing the similarity of the 2 datasets resulted in a P value below 0.0001.

Results using a fixed value for K_{heat} are shown in Table 1; those using hourly-resolved data for K_{heat} are given in the Supplement (see Table S2).

3.4 Performance of the MBR method on day/night split data

Fluxes of CO_2 during the nighttime only, measured using the MBR method in combination with simulated sampling times ranging from 1 h to 1 week are on average a factor of 1.7 and up to a factor of 2.1 larger than those derived with the EC method (see Table 1). Fluxes of CO_2 during the daytime only, measured using the MBR method have, in some cases, the opposite sign of the fluxes reported using the EC technique. Specifically, the MBR method produces daytime fluxes with the opposite sign compared to the EC method during daytime in the spring and for the 1 week duration simulated sampling scenario in the winter (values marked with (!) in Table 1). In those cases where the direction of flux calculated with the MBR does not agree with the EC method, the disagreement is attributable to the median value of ΔCO_2 (between 41.5 and 33 m) selected to represent the sampling period having a sign that implies a flux in the opposite direction of the flux measured with the EC method.

In cases where the direction of daytime flux measured using the MBR method agreed with the EC method, the ratio of the two fluxes ranged from 0.32 to 1.4, implying that the two methods differed by factors that range from 1.4 to 3.0 and that the MBR method may either underestimate or overestimate fluxes relative to the EC method.

For H_2O , the fluxes measured with the MBR method are in the same direction as those measured by the EC method during spring, summer, and during the day in the fall. When the two methods agree about the direction of flux, the ratio of fluxes mea-

ACPD

15, 32759–32777, 2015

Comparison of eddy covariance and modified Bowen ratio methods

D. J. Bolinius et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sured by the MBR method to the EC method is between 0.9 and 0.052, implying differences between the methods by factors of 1.1 to 20, and that fluxes of H₂O are underestimated if the MBR method is applied as compared to the EC method. Fluxes measured during the winter season with the MBR method are negative (i.e., from the atmosphere to the surface), while the EC method indicates fluxes are positive (i.e., from the surface to the atmosphere). Fluxes measured by both methods during the night in the fall are small, with a upward flux measured using the EC method and downward fluxes measured by the MBR method in all cases except the longest 1 week simulated sampling period.

In general, the fluxes measured using the two different methods are in better agreement for CO₂ than for H₂O (Table 1). It is interesting to note that the MBR method generally overestimates the flux of CO₂ relative to the EC method, while it in most cases underestimates fluxes of H₂O.

4 Discussion

The MBR method fails for CO₂ when using continuous time series in cases where the simulated sampling period encompasses the shift between night and day and there is also a shift from upward fluxes (dominated by respiration) at night to downward fluxes (dominated by photosynthesis) during daytime. As shown in Eq. (3), the flux estimated using the MBR method only depends on K_{heat} and the vertical concentration gradient for the compound of interest. The error arises because the average values of K_{heat} are dominated by high values that occur during the day, while average values of ΔCO_2 are dominated by extreme values that occur at night (see especially the summer season for CO₂ in Fig. 1 and data for CO₂ in the summer visualized in Fig. S5).

The CO₂ fluxes determined using the MBR method during the daytime for the spring season and when simulating a sampling time of 1 week during daytime for the winter were in the opposite direction relative to the fluxes determined using the EC method. During spring this could be caused by a reversal of the direction of flux due to the

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Comparison of eddy covariance and modified Bowen ratio methods

D. J. Bolinius et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



development of leaves and the start of photosynthesis taking place halfway across the season, which would produce a shift from a continuous net flux of CO_2 out of the canopy to a diel cycle of CO_2 uptake and release. Furthermore, it is possible that the simulated 1 week sampling duration during daytime in the winter might have encompassed periods when the net direction of flux of CO_2 changed due to the movement of air masses with variable CO_2 concentrations across the region. Thus, all of the cases of disagreement between the MBR method and the EC method about the direction of flux of CO_2 that are shown in Table 2 might be attributable to applying the MBR method using simulated sampling times that encompass a change of direction of the net flux.

In general, the duration of simulated sampling does not have a strong influence on the fluxes measured with the MBR method. Exceptions are the longest simulated sampling times in during the daytime in winter for CO_2 and during nighttime in the fall for H_2O . We simulated sampling times of 24 h, 3 days and 1 week by combining data that were measured over non-consecutive 8 h periods during 2 month time windows selected to represent the four seasons. For the 3 day and 1 week simulated sampling times there are just 3 to 4 data-points per season, depending on the data availability, which may introduce higher uncertainty in the median value used in our MBR method calculations compared to the shorter simulated sampling times.

For H_2O , which nearly always has a net flux upwards from the canopy during both the day and the night, pooling of the data over longer time intervals and application of the MBR method also led to estimations of the direction of flux that were opposite the EC method in the winter and fall seasons. In winter, fluxes of H_2O measured by the EC method were small and upwards, while those calculated with the MBR method were small and downwards (Table 2). A recent study focusing on drainage basins in Canada, reported low but positive fluxes of water vapour during the winter season (Wang et al., 2014), which is consistent with measurements using the EC method. In this case we traced the origin of the different directions of flux calculated with the MBR method relative to the EC method to differences in H_2O concentration gradients across different height intervals. Reported fluxes of H_2O using the EC method were measured at 33.3 m

Comparison of eddy covariance and modified Bowen ratio methods

D. J. Bolinius et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



while for the MBR method we have used a gradient of concentrations measured at 41.5 and 33 m. During the winter for H₂O there was a clear discrepancy between gradients measured at these heights and between 33 and 25.7 m, with the latter being more consistent with the EC measurements (Fig. S6). The cold and low humidity during the winter in Canada might play a role here as discrepancies between the concentration gradients of H₂O at different heights were only observed in the winter and to a lesser extent in the fall.

It is clear from our analysis that a requirement for the MBR method to give accurate results for prolonged sampling times is to only sample during a time period when the chemicals of interest are expected to have a unidirectional flux. The occurrence of a day/night regime has implications for designing sampling campaigns for organic pollutants that require sampling times longer than the 8 h intervals with stable conditions chosen as daytime or nighttime above, and which may exhibit changes in the direction of flux between daytime and nighttime periods. This can be the case for many POPs, and the direction of fluxes can be estimated using appropriately parameterized dynamic chemical multimedia fate models (see, for example, Gasic et al., 2010). When longer sampling times are needed, samples could be pooled by sampling at the same time of day during consecutive 24 h periods as was simulated here by combining data from selected 8 h periods.

When sampling over time periods of several hours to several days, as simulated in this study, it is very likely that steady-state conditions during the sampling period are not achieved. Our results indicate however that when fluxes were unidirectional, measurements using the MBR method were usually within 1 order of magnitude of those from the EC method, and that in most cases the difference was less than a factor of 4. The summer period, in which K_{heat} is low and the gradients of CO₂ and H₂O are large due to stable atmospheric conditions, showed the best agreement between the MBR and EC methods. In the study of Choi et al. (2008), they estimated their uncertainty in fluxes of PAHs derived from the MBR method was an order of magnitude,

which is within the range of agreement we obtained between the MBR method and the EC method.

Fluxes measured with the MBR method corresponded better with the EC data for CO₂ than for H₂O, the reason of which is unknown. However our investigation of the discrepancy in direction of flux between the two methods for H₂O in the winter suggests that the concentration gradient of H₂O is more variable over height than that of CO₂. Therefore the gradient that we selected between 41.5 and 33 m may be more representative of the flux measured at 33.3 m with the EC method for CO₂ than for H₂O.

In general, the variation between fluxes determined using the MBR method for different sampling frequencies was small, suggesting that longer sampling times did not introduce a higher bias relative to EC measurements. Using a single value for K_{heat} , in this case the geometric mean across the sampling period, was found to be a good substitute for hourly K_{heat} values, indicating that it is possible to use the MBR method when there is no high frequency data available for the transfer coefficient.

Some studies have shown that additional transport mechanisms aside from eddy diffusion, such as advection, can become more important during night, thereby violating the conditions needed for EC measurements to take place, and resulting in underestimations of the night time fluxes (Aubinet, 2008). In this study, there is no clear difference in the performance of the two methods relative to one another during either day or night. We note however that the MBR method relies on K_{heat} determined under the assumption that only turbulent atmospheric processes occur, so the effect of additional transport mechanisms might not be evident in our analysis.

Based on the findings in this study it is clear that field studies that use the MBR method to measure gas fluxes of POPs and other organic micropollutants should be designed such that the direction of the flux does not change during the sampling period. If this condition is fulfilled and the concentration gradients are large enough to be measured accurately, then we find no strong evidence that the duration of sample collection affects the performance of the MBR method. Furthermore, using a fixed

Comparison of eddy covariance and modified Bowen ratio methods

D. J. Bolinius et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Comparison of eddy covariance and modified Bowen ratio methods

D. J. Bolinius et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



value for the transfer coefficient instead of hourly data is a good alternative that should not introduce a significant bias when there is no high frequency data available. The MBR method could be used to measure fluxes of POPs in depositional areas such as forests as shown by the study of Choi et al. (2008) or in source areas such as large cities where recent studies of vertical concentration gradients of POPs did not lead to quantitative flux estimates (Li et al., 2009; Moreau-Guigon et al., 2007).

**The Supplement related to this article is available online at
doi:10.5194/acpd-15-32759-2015-supplement.**

Author contributions. M. MacLeod had the idea for this study, D. J. Bolinius and M. MacLeod designed the study and D. J. Bolinius gathered the data, performed the analysis and prepared the manuscript with contributions from A. Jahnke and M. MacLeod.

Acknowledgements. We thank Frank Wania for constructive suggestions during the early phases of this work. This research was funded by the Swedish Research Council Vetenskap-srådet (VR), project number 2011-3890, “Investigating thermodynamic controls on the cycling of persistent organic chemicals in forest systems”.

References

- Aubinet, M.: Eddy covariance CO₂ flux measurements in nocturnal conditions: an analysis of the problem, *Ecol. Appl.*, 18, 1368–1378, doi:10.1890/06-1336.1, 2008.
- Baldocchi, D. D., Hincks, B. B., and Meyers, T. P.: Measuring biosphere–atmosphere ex- changes of biologically related gases with micrometeorological methods, *Ecology*, 69, 1331, doi:10.2307/1941631, 1988.
- Businger, J. A.: Evaluation of the accuracy with which dry deposition can be measured with current micrometeorological techniques, *J. Clim. Appl. Meteorol.*, 25, 1100–1124, doi:10.1175/1520-0450(1986)025<1100:EOTAWW>2.0.CO;2, 1986.

Comparison of eddy covariance and modified Bowen ratio methods

D. J. Bolinius et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Choi, S.-D., Staebler, R. M., Li, H., Su, Y., Gevaio, B., Harner, T., and Wania, F.: Depletion of gaseous polycyclic aromatic hydrocarbons by a forest canopy, *Atmos. Chem. Phys.*, 8, 4105–4113, doi:10.5194/acp-8-4105-2008, 2008.

Gasic, B., MacLeod, M., Scheringer, M., and Hungerbuhler, K.: Assessing the impact of weather events at mid-latitudes on the atmospheric transport of chemical pollutants using a 2-dimensional multimedia meteorological model, *Atmos. Environ.*, 44, 4489–4496, doi:10.1016/j.atmosenv.2010.07.016, 2010.

Gusev, A., MacLeod, M., and Bartlett, P.: Intercontinental transport of persistent organic pollutants: a review of key findings and recommendations of the task force on hemispheric transport of air pollutants and directions for future research, *Atmospheric Pollut. Res.*, 3, 463–465, doi:10.5094/APR.2012.053, 2012.

Hung, H., MacLeod, M., Guardans, R., Scheringer, M., Barra, R., Harner, T., and Zhang, G.: Toward the next generation of air quality monitoring: persistent organic pollutants, *Atmos. Environ.*, 80, 591–598, doi:10.1016/j.atmosenv.2013.05.067, 2013.

Li, Y., Zhang, Q., Ji, D., Wang, T., Wang, Y., Wang, P., Ding, L., and Jiang, G.: Levels and vertical distributions of PCBs, PBDEs, and OCPs in the atmospheric boundary layer: observation from the Beijing 325-m meteorological tower, *Environ. Sci. Technol.*, 43, 1030–1035, 2009.

Majewski, M. S.: Micrometeorologic methods for measuring the post-application volatilization of pesticides, *Water Air. Soil Pollut.*, 115, 83–113, doi:10.1023/A:1005297121445, 1999.

McKone, T. E. and MacLeod, M.: Tracking multiple pathways of human exposure to persistent multimedia pollutants: regional, continental and global-scale models, *Annu. Rev. Env. Resour.*, 28, 463–492, doi:10.1146/annurev.energy.28.050302.105623, 2003.

Meyers, T. P., Hall, M. E., Lindberg, S. E., and Kim, K.: Use of the modified bowen-ratio technique to measure fluxes of trace gases, *Atmos. Environ.*, 30, 3321–3329, doi:10.1016/1352-2310(96)00082-9, 1996.

Moreau-Guigon, E., Motelay-Massei, A., Harner, T., Pozo, K., Diamond, M., Chevreuil, M., and Blanchoud, H.: Vertical and temporal distribution of persistent organic pollutants in Toronto. 1. Organochlorine pesticides, *Environ. Sci. Technol.*, 41, 2172–2177, doi:10.1021/es062705s, 2007.

Nizzetto, L., Macleod, M., Borgå, K., Cabrerizo, A., Dachs, J., Guardo, A. D., Ghirardello, D., Hansen, K. M., Jarvis, A., Lindroth, A., Ludwig, B., Monteith, D., Perlinger, J. A., Scheringer, M., Schwendenmann, L., Semple, K. T., Wick, L. Y., Zhang, G., and Jones, K. C.: Past,

present, and future controls on levels of persistent organic pollutants in the global environment, Environ. Sci. Technol., 44, 6526–6531, doi:10.1021/es100178f, 2010.

Pierce, A. M., Moore, C. W., Wohlfahrt, G., Hörtnagl, L., Kljun, N., and Obrist, D.: Eddy covariance flux measurements of gaseous elemental mercury using Cavity Ring-Down Spectroscopy, Environ. Sci. Technol., 49, 1559–1568, doi:10.1021/es505080z, 2015.

Stockholm Convention: Stockholm convention on Persistent Organic Pollutants (POPs) as amended in 2009, available at: <http://chm.pops.int/TheConvention/Overview/TextoftheConvention/tabid/2232/Default.aspx> (last access: November 2015), 2010.

Wang, S., Huang, J., Li, J., Rivera, A., McKenney, D. W., and Sheffield, J.: Assessment of water budget for sixteen large drainage basins in Canada, J. Hydrol., 512, 1–15, doi:10.1016/j.jhydrol.2014.02.058, 2014.

ACPD

15, 32759–32777, 2015

Comparison of eddy covariance and modified Bowen ratio methods

D. J. Bolinius et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Comparison of eddy covariance and modified Bowen ratio methods

D. J. Bolinius et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Cumulative fluxes for 8 h periods representing day and night across the 2 month periods representing spring, summer, fall and winter. Fluxes measured by the MBR method that are in the opposite direction than those measured by the EC method are marked with “!”. Positive fluxes are defined as fluxes moving upwards from the canopy. The ratio of MBR results over EC results is based on the geometric mean of the MBR results divided by the EC result. The MBR fluxes for the 1 week sampling period were left out in the calculation of the geometric mean during the day in winter for CO₂ and during the night in fall for H₂O. This table shows fluxes calculated with a constant value for K_{heat} .

CO ₂ (PPM m)		spring		summer		fall		winter	
		Day	Night	Day	Night	Day	Night	Day	Night
eddy covariance		−0.332	0.295	−3.360	1.071	0.074	0.227	0.189	0.112
modified Bowen ratio constant K_{heat}	1/h	0.275 (!)	0.408	−1.202	1.529	0.073	0.390	0.163	0.224
	1/2 h	0.313 (!)	0.434	−1.124	1.703	0.113	0.421	0.187	0.196
	1/4 h	0.319 (!)	0.413	−1.107	1.544	0.067	0.396	0.163	0.199
	1/8 h	0.250 (!)	0.441	−1.000	1.697	0.093	0.392	0.159	0.159
	1/day	0.353 (!)	0.466	−1.064	2.602	0.145	0.527	0.176	0.185
	1/3 days	0.335 (!)	0.658	−0.991	2.258	0.138	0.618	0.096	0.185
	1/week	0.409 (!)	0.459	−1.114	2.576	0.136	0.621	−0.019 (!)	0.032
MBR/EC method		−0.960 (!)	1.566	0.323	1.811	1.425	2.071	0.893	1.317
H ₂ O (PPT m)		spring		summer		fall		winter	
		Day	Night	Day	Night	Day	Night	Day	Night
eddy covariance		0.420	0.016	1.118	0.038	0.180	0.005	0.049	0.001
modified Bowen ratio constant K_{heat}	1/h	0.048	0.006	0.243	0.027	0.011	−0.001 (!)	−0.007 (!)	−0.009 (!)
	1/2 h	0.052	0.006	0.236	0.030	0.009	−0.001 (!)	−0.007 (!)	−0.008 (!)
	1/4 h	0.040	0.007	0.265	0.032	0.008	−0.001 (!)	−0.006 (!)	−0.007 (!)
	1/8 h	0.033	0.009	0.239	0.030	0.008	−0.001 (!)	−0.008 (!)	−0.008 (!)
	1/day	0.029	0.009	0.269	0.044	0.007	−0.001 (!)	−0.004 (!)	−0.011 (!)
	1/3 days	0.046	0.012	0.332	0.050	0.010	−0.003 (!)	−0.015 (!)	−0.015 (!)
	1/week	0.061	0.007	0.323	0.038	0.014	0.001	−0.009 (!)	−0.014 (!)
MBR/EC method		0.102	0.464	0.241	0.915	0.052	−0.169 (!)	−0.149 (!)	−9.305 (!)

Comparison of eddy covariance and modified Bowen ratio methods

D. J. Bolinius et al.




Figure 1. Half-hourly averages of K_{heat} and the concentration gradients of CO_2 and H_2O across the different seasons (concentration at 40.7 m – concentration at 33.3 m). The dotted line indicates 0 in all plots, the grey areas indicate the 8 h periods representing day and night.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Comparison of eddy covariance and modified Bowen ratio methods

D. J. Bolinius et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 2. A comparison of measured fluxes using the modified Bowen ratio (MBR) method (1 and 24 h pooled data using hourly K_{heat} values) with the eddy covariance (EC) measurements. The simulated 24 h sampling time includes a change in the direction of the CO_2 flux, which results in a measured flux with the MBR method that has the wrong direction and magnitude.