

Quantification of black carbon mixing state from traffic: implications for aerosol optical properties

Megan D. Willis¹, Robert M. Healy^{2,3,a}, Nicole Riemer⁴, Matthew West⁵, Jon M. Wang², Cheol-Heon Jeong², John C. Wenger³, Greg J. Evans², Jonathan P. D. Abbatt¹, and Alex K. Y. Lee¹

¹Department of Chemistry, University of Toronto, Ontario, Canada

²Southern Ontario Centre for Atmospheric Aerosol Research, Department of Chemical Engineering and Applied Chemistry, University of Toronto, Ontario, Canada

³Department of Chemistry and Environmental Research Institute, University College Cork, Ireland

⁴Department of Atmospheric Sciences, University of Illinois at Urbana-Champaign, Urbana, USA

⁵Department of Mechanical Science and Engineering, University of Illinois at Urbana-Champaign, Urbana, USA

^aNow at Air Monitoring and Transboundary Air Sciences Section, Environmental Monitoring and Reporting Branch, Ontario Ministry of the Environment and Climate Change, Ontario, Canada

Correspondence to: M.D. Willis (megan.willis@mail.utoronto.ca) and A.K.Y. Lee (alexky.lee@utoronto.ca)

Abstract. The climatic impacts of black carbon (BC) aerosol, an important absorber of solar radiation in the atmosphere, remain poorly constrained and are intimately related to its particle-scale physical and chemical properties. Using particle-resolved modelling informed by quantitative measurements from a soot-particle aerosol mass spectrometer, we confirm that the mixing state (the distribution of co-emitted aerosol amongst fresh BC-containing particles) at the time of emission significantly affects BC-aerosol optical properties even after a day of atmospheric processing. Both single particle and ensemble aerosol mass spectrometry observations indicate that BC near the point of emission co-exists with hydrocarbon-like organic aerosol (HOA) in two distinct particle types: HOA-rich and BC-rich particles. The average mass fraction of black carbon in HOA-rich and BC-rich particle classes was <0.1 and 0.8 , respectively. Notably, approximately 90 % of BC mass resides in BC-rich particles. This new measurement capability provides quantitative insight into the physical and chemical nature of BC-containing particles and is used to drive a particle-resolved aerosol box model. Significant differences in calculated single scattering albedo (an increase of 0.1) arise from accurate treatment of initial particle mixing state as compared to the assumption of uniform aerosol composition at the point of BC injection into the atmosphere.

1 Introduction

Incomplete combustion emits teragram quantities of black carbon (BC) aerosol to the troposphere each year, resulting in a significant warming effect on climate that may be second only to carbon

dioxide (Bond et al., 2013; Ramanathan and Carmichael, 2008; Jacobson, 2001). BC influences climate directly, by absorbing solar radiation, and indirectly, by changing cloud properties and altering snow and ice melt. BC impacts on the global scale remain poorly constrained and are intimately related to its particle-scale physical and chemical properties (Bond et al., 2013). The majority of BC emissions in North America, Europe and Latin America are derived from traffic-related sources, though the specific physical and chemical properties of BC-containing particles at emission depend greatly on the source (Bond et al., 2013). BC-containing particles are generally hydrophobic near emission and become mixed over time with hydrophilic species through condensation and coagulation (Johnson et al., 2005; Moteki et al., 2007; Moffet and Prather, 2009), with resulting impacts on particle hygroscopicity (McMeeking et al., 2011b; Liu et al., 2013a; Laborde et al., 2013) and optical properties (Zhang et al., 2008; Cappa et al., 2012; Lack et al., 2012; Knox et al., 2009; Liu et al., 2015). Despite extensive previous work, our understanding of the role of mixing state in influencing the climate impacts of BC remains incomplete, in part because of instrumental challenges in particle characterization.

Studies assessing both the composition and amount of non-BC species in BC-containing particles are rare. A large body of evidence from urban, tunnel and engine emission studies has shown that individual combustion particles are mixtures of BC, inorganic species, metals, and hydrocarbon-like organic species at the time of emission (e.g., Johnson et al., 2005; Toner et al., 2006; Schneider et al., 2006; Tritscher et al., 2011; Chirico et al., 2010; Zhang et al., 2013; Massoli et al., 2012; Dallmann et al., 2014). Among these studies, single particle mass spectrometry has directly demonstrated mixing of BC and non-BC species in traffic-derived particles. For example, Toner et al. 2006 observed that emissions from heavy duty diesel engines were dominated by particles containing BC, organic species, calcium and phosphate, with one particle type dominated by BC and another with higher levels of organic species. In contrast, Healy et al. 2012 observed BC-dominated particles from traffic emissions in a European city. Other approaches to measure BC mixing state, including hygroscopicity and volatility differential mobility techniques, have highlighted the presence of an external mixture in terms of particle volatility with lower volatility, BC-containing aerosol being less hygroscopic (Kuwata et al., 2009; McMeeking et al., 2011a). Further, single particle soot photometer measurements of coating thickness in urban areas have shown that traffic-related BC-containing particles are largely uncoated or very thinly coated (Shiraiwa et al., 2008; McMeeking et al., 2011b; Laborde et al., 2013; Liu et al., 2014). In addition, microscopy studies have illustrated the dominance of bare or thinly coated BC-containing particles in traffic emissions (China et al., 2014) and the occurrence of coating, embedding and compaction of BC as it is aged (Adachi and Buseck, 2013; Adachi et al., 2014). While previous work has provided valuable insight into BC mixing state, the majority of past approaches have not allowed simultaneous quantification of BC mixing state on a mass basis and chemical characterization of non-BC species.

55 Considerable attention has been paid to BC-containing aerosol because of the potential for short-
term climate mitigation through emissions reduction (Shindell et al., 2012). Therefore, a quantitative
understanding of BC mixing state is crucial for three reasons: first, to assess rates of BC processing
and removal in the atmosphere; second, to assess the role that BC-containing particles play as cloud
nuclei; and third, to assess direct effects on solar radiation. Aerosol optical properties are central
60 parameters required to evaluate direct radiative forcing (DRF). One of the largest contributors to
uncertainty in DRF calculations is the single scattering albedo (SSA) (McComiskey et al., 2008),
defined as the ratio of aerosol scattering to total light extinction. Previous work has clearly demon-
strated that calculations of aerosol optical properties depend upon assumptions about particle mixing
state (e.g., Jacobson, 2000), with assumptions of uniform internal mixing producing overestimates
65 of absorption efficiency and underestimates of single scattering albedo (Oshima et al., 2009; Zaveri
et al., 2010; Matsui et al., 2013). The role of BC-aerosol mixing state at emission in affecting its
mixing state in the atmosphere has also been highlighted in modelling studies explicitly treating
detailed aerosol micro-physical and chemical processes (e.g., Matsui et al., 2013). However, model
assessments driven by quantitative measurements of aerosol mixing state remain rare.

70 In this work, we use a soot-particle aerosol mass spectrometer, equipped with a light-scattering
module, to determine the mixing state of BC-containing particles from traffic-dominated sources
in an urban environment. These measurements provide quantitative insight into the physical and
chemical nature of BC-containing particles near emission, and are used to drive a particle-resolved
aerosol box model to assess the effect of accurately representing BC mixing on aerosol optical
75 properties.

2 Methods

2.1 The soot-particle aerosol mass spectrometer (SP-AMS)

2.1.1 Two urban studies: locations and SP-AMS configuration

A soot-particle aerosol mass spectrometer (SP-AMS, Aerodyne Research Inc., Billerica, MA, USA),
80 equipped with a light scattering module, was deployed in two urban studies to assess the mixing state
of refractory black carbon (rBC) containing particles derived from vehicle emissions (Lee et al.,
2015a). The first study was conducted in downtown Toronto away from major roadways (referred
to as the “non-roadside” study), and the second was performed at ground level, near a busy road in
downtown Toronto (referred to as the “roadside” study), to investigate fresh vehicle emissions. The
85 non-roadside site is described in detail in Lee et al 2015a. The roadside study took place from May
31 to June 24, 2013 at the Southern Ontario Centre for Atmospheric Aerosol Research (SOCAAR)
facility in downtown Toronto, Canada, located at ground level and adjacent to a road with traffic
volumes ranging from 16000 to 25000 vehicles per day (Sabaliauskas et al., 2014). Ambient air was

sampled at 170 L min^{-1} through a 10 cm (inner diameter) stainless steel tube fitted with a $2.5 \mu\text{m}$ cut-off inlet, located 15 m from the roadside at a height of 3 m a.g.l.

Two main modes of operation are possible with the SP-AMS, depending on the vaporizer configuration. The instrument was operated such that both rBC-containing particles and non-rBC-containing particles were detected in the non-roadside study (i.e., a dual-vaporizer configuration), whereas exclusively rBC-containing particles were detected in the roadside study (i.e., a laser-only configuration) (Lee et al., 2015a; Onasch et al., 2012).

2.1.2 SP-AMS operation and calibration

Details of the SP-AMS (and SP-AMS with light scattering) have been described elsewhere (Lee et al., 2015a; Onasch et al., 2012). The SP-AMS detects black carbon, which evaporates at $\sim 4000 \text{ K}$, as C_x^+ fragments (predominantly m/z 12, 24, 36, 48 and 60) and is referred to as refractory black carbon (rBC) (Onasch et al., 2012). We use the term “black carbon” (BC) when referring generally to the concept of “soot”-type aerosol, while rBC is used when referring to quantities measured by the SP-AMS. In the SP-AMS, rBC and associated species are volatilized in an infrared laser beam (1064 nm) and are ionized using 70 eV electron impact ionization followed by detection in a time-of-flight mass spectrometer (ToF-MS) operated in “V-mode.” For the majority of the study the SP-AMS was operated at one-minute time resolution alternating between bulk mass spectrum (MS), particle time-of-flight (pToF) and single particle modes. The SP-AMS was operated at high-time resolution (1 Hz) in MS mode for a total of five days during the roadside study. SP-AMS data was analysed using the Igor Pro-based analysis tool PIKA (Seuper, 2010). The single particle categorization procedure and k -means clustering algorithm that were used to analyse the single particle data followed the description in Lee et al. (2015a). Positive matrix factorization (PMF) analysis of ensemble data was performed to identify the sources of rBC and organics, and to speciate different types of organic species (i.e., oxygenated organic aerosol (OOA), biomass burning organic aerosol (BBOA), and hydrocarbon-like organic aerosol (HOA)), based on procedures described previously (Paatero and Tapper, 1994; Zhang et al., 2011; Ulbrich et al., 2009).

Without the tungsten vaporizer, direct calibrations of the ionization efficiency for NH_4NO_3 (IE_{NO_3}) are not possible. Therefore, size-selected (300 nm) Regal Black (Regal 400R Pigment, Cabot Corp.) particles were used to determine the mass based ionization efficiency of rBC (mIE_{rBC}) (Onasch et al., 2012; Willis et al., 2014). The relative ionization efficiency for rBC ($\text{RIE}_{\text{rBC}} = \text{mIE}_{\text{rBC}}/\text{mIE}_{\text{NO}_3}$) was 0.2 ± 0.05 , as experimentally determined before removal of the tungsten vaporizer. Assuming that RIE_{rBC} is constant, IE_{NO_3} could be calculated based on known values of mIE_{rBC} and RIE_{rBC} . The average mIE_{rBC} was $189 \pm 20 \text{ ions pg}^{-1}$ in the roadside study. The calculated IE_{NO_3} was then used with recommended relative ionization efficiencies to quantify other aerosol species associated with rBC (Jimenez et al., 2003). Collection efficiency for rBC particles was determined in the roadside study using beam width probe (BWP) measurements described in (Willis et al., 2014). Ambient

125 rBC-containing particles had an average beam width $\sigma = 0.46 \pm 0.03$ mm, which is close to, but
wider than that of 300 nm Regal Black particles ($\sigma = 0.40 \pm 0.08$) (Willis et al., 2014). Therefore,
a collection efficiency (CE) of 0.6 was applied for absolute quantification of rBC and associated
species. A time-varying collection efficiency was not possible with BWP measurements available
here; the assumption of a constant CE over periods of local and long-range transport influence in
130 this study provided a good linear correlation with photoacoustic soot spectrometer (PASS) absorp-
tion measurements (405 and 781 nm) and SP-AMS rBC (Healy et al., 2015, Fig. 4). Note that the CE
applied will not impact calculations of the mass fraction of rBC (mf_{rBC}). However, two additional
uncertainties in SP-AMS measurements may affect calculation of mf_{rBC} . First, there are uncertainties
in the recommended RIE for organic species evaporating from rBC in the SP-AMS of up to ~ 50 %
135 (Lee et al., 2015a; Willis et al., 2014), which could cause an overestimation in the mass of coating
material and a corresponding underestimation in mf_{rBC} . Second, it is possible for rBC-containing
particles to pass through the edges of the laser vaporizer, thus producing a heating effect sufficient
to evaporate some fraction of the coating materials but not evaporate the rBC itself. This effect may
also lead to an underestimation in mf_{rBC} . SP-AMS CE and quantification are discussed in further
140 detail in Supplement Sect. 1.

2.2 Photoacoustic soot spectrometer (PASS)

A photoacoustic soot spectrometer (PASS-3, Droplet Measurement Techniques, Boulder, CO) was
used to measure aerosol absorption (b_{abs}) and scattering (b_{sca}) coefficients ($M m^{-1}$) at 405 and
781 nm. A 532 nm laser is not installed in this particular unit. The PASS determines aerosol absorp-
145 tion ($M m^{-1}$) in a cavity which acts as an acoustic resonator. The absorption of incoming radiation
heats the particles, which in turn heat the surrounding air in the cavity (Arnott et al., 1999). The
aerosol-laden air thus expands, resulting in a pressure disturbance. By modulating the laser power
at the resonance frequency of the cavity, the pressure disturbance is amplified and the resulting
acoustic wave is measured using a microphone. Light scattering at both wavelengths is concurrently
150 measured using reciprocal nephelometry (Moosmüller et al., 2009; Flowers et al., 2010; Chan et al.,
2011). Scattering signals were not corrected for truncation; however, since mass-based particle size
distributions at the roadside site generally peaked near or below 100 nm a negative bias in the scat-
tering measurement is unlikely. Since a 532 nm laser was not present in this unit an NO_2 calibration
was not possible, and the instrument was calibrated using a propane soot generator (miniCAST,
155 6203A, Jing). PASS measurements of the bulk single scattering albedo at 405 nm (selected due to
superior signal-to-noise ratio for scattering relative to the 781 nm channel) are used here only to
illustrate differences in optical properties in vehicle plumes with varying composition.

2.3 Particle-resolved box model simulations

We used the stochastic particle-resolved box model PartMC-MOSAIC (Particle Monte Carlo - Model
160 for Simulating Aerosol Interactions and Chemistry) (Riemer et al., 2009; Zaveri et al., 2008) to quan-
tify the importance of the observed mixing state information for the calculation of CCN and aerosol
optical properties. PartMC-MOSAIC is suited for this task, as it explicitly tracks the composition
of individual aerosol particles (in our case about 10^5 particles) in a population of different particle
types within a well-mixed computational volume. Particle emissions, dilution with the background,
165 and Brownian coagulation are simulated stochastically with PartMC, by generating a realization of
a Poisson process using weighted particles in the sense of DeVile et al. (DeVile et al., 2011) and an
accelerated binned sampling strategy (Michelotti et al., 2013). Coupled to PartMC is the MOSAIC
chemistry code, which simulates gas chemistry (Zaveri and Peters, 1999), particle phase thermody-
namics (Zaveri et al., 2005a, b), and dynamic gas-particle mass transfer (Zaveri et al., 2008) in a
170 deterministic manner.

Here we compare two simulations that differ in the way the mixing state of the aerosol initial
conditions and the aerosol emissions are prescribed. The set-up of the particle-resolved simulations
is similar to the urban plume scenarios in Riemer et al. (2009) and Zaveri et al. (2010). Common
to both simulations are the following specifics: The number of computational particles used in the
175 simulation is about 10^5 . The total simulation time is 24 h, with a simulation start of 06:00 local
standard time (LST). The initial gas phase concentrations are the same as in Zaveri et al. (2010).
In contrast to Zaveri et al. (2010), we prescribe a constant mixing height (400 m), relative humidity
(70 %) and temperature (298.15 K) to simplify the interpretation of the results. Since the entrainment
of background aerosol can modify the aerosol mixing state substantially (Liu et al., 2011), we include
180 constant dilution with the background at a rate of $1.5 \times 10^{-5} \text{ s}^{-1}$. The background aerosol is non-
absorbing and consists of ammonium sulfate mixed with biogenic secondary organic aerosol. This
dilution rate corresponds to about 75 % of the aerosol being replaced in a 24 h period, comparable
to a diurnal mixing height increase of 500 m to 2000 m (e.g., Liu et al., 2011), although imposed
uniformly over the day for simplicity. We also reduced the gas phase emissions by a factor of two
185 compared to Zaveri et al. (2010) to create more moderately polluted conditions. The initial aerosol
population was sampled from log-normal size distributions fitted from the measurement data, as
shown in Table S1 in the Supplement. The derivation of model inputs from measurement data is
described in Supplement Sect. 3.

To fully exploit the information supplied by the observations, we introduced a new method of
190 sampling the aerosol composition for the aerosol initial conditions and emissions. We now allow
for stochastic composition variation at each diameter, with a prescribed standard deviation around a
mean value. The new composition-sampling method proceeds as follows. For each aerosol particle,
the diameter is first sampled, and then the composition is sampled as a vector of mass fractions
 $\mathbf{w} = [w_{\text{BC}}, w_{\text{HOA}}]^T$. Although we only sample two species for the simulations in this paper, we

195 specify the sampling algorithm for any number of species. We use bold symbols to denote vectors, so \mathbf{mf} is the mean mass fraction vector from Table S1 in the Supplement. We write Σ for the diagonal covariance matrix with diagonal entries $(\sigma_i)^2$, $\mathbf{1}$ for the vector of ones, and \mathbf{x}^\top for the transpose of a vector \mathbf{x} . We sample the composition vector \mathbf{w} from a multivariate normal distribution on the affine hyperplane of vectors that sum to one, truncated to the positive closed orthant (and thus to
 200 the probability or standard simplex). This is done as shown in Algorithm 1 using an accept-reject procedure.

Algorithm 1 Composition sampling

```

1: repeat
2:    $\mathbf{x} \leftarrow \mathcal{N}(\mathbf{mf}, \Sigma)$ 
3:    $\mathbf{w} \leftarrow \mathbf{x} + (1 - \mathbf{1}^\top \mathbf{x})\mathbf{mf}$ 
4: until  $w_i \geq 0$  for all  $i$ 

```

Algorithm 1 has five important properties, as follows. (1) It is clear that if the algorithm terminates then \mathbf{w} lies in the positive closed orthant, so every species has a non-negative mass fraction. (2) The sum of the sampled mass fractions \mathbf{w} is one, as can be seen from $\mathbf{1}^\top \mathbf{w} = \mathbf{1}^\top \mathbf{x} + (1 - \mathbf{1}^\top \mathbf{x})\mathbf{1}^\top \mathbf{mf} =$
 205 $\mathbf{1}^\top \mathbf{x} + 1 - \mathbf{1}^\top \mathbf{x} = 1$, assuming that the mean mass fractions sum to one and so $\mathbf{1}^\top \mathbf{mf} = 1$. (3) The distribution of \mathbf{w} is a truncated multivariate normal, which follows from the fact that \mathbf{x} has a multivariate normal distribution, \mathbf{w} is an affine function of \mathbf{x} , and the accept-reject procedure samples from the truncation of the per-iteration distribution. Note, however, that the resulting marginal distributions for each single-species mass fraction will not in general have mean and standard deviation
 210 given by \mathbf{mf}_i and σ_i , respectively. (4) Adding additional species with zero mean mass fraction and zero standard deviation will not change the distribution of the other sampled mass fractions, as the additional species will have zero components in both \mathbf{x} and \mathbf{w} and will not alter the calculation of any other components. (5) Finally, the algorithm will terminate with probability one, under the conditions that that the mean mass fractions sum to one and each σ_i is non-negative and zero only if the
 215 corresponding \mathbf{mf}_i is non-negative. This last property can be seen by first considering the set A of \mathbf{x} that will result in a \mathbf{w} in the positive closed orthant, terminating the algorithm. This set A is exactly the direct sum of the probability simplex with the span of $\{\mathbf{mf}\}$, as can be seen by checking that $\mathbf{x} = \mathbf{p} + \alpha\mathbf{mf}$ for \mathbf{p} in the probability simplex and any scalar α results in $\mathbf{w} = \mathbf{p}$, so that the entire probability simplex is thus obtained from A . The conditions on \mathbf{mf} and the σ_i imply that \mathbf{x} has non-
 220 zero probability of lying in A , and hence the algorithm has non-zero probability of terminating on each iteration. Because each iteration is an i.i.d. Bernoulli trial, the algorithm thus terminates with probability one.

The two simulations differ in the assigned mixing state of the initial aerosol populations and the aerosol emissions. For the measurement-constrained case we distinguish between rBC-rich and
 225 HOA-rich particle classes, as the observations indicate. Both classes consist of two modes, which

amounts to four modes in total, as listed in Table S1 in the Supplement. Note that the modes BC1 and HOA1 have the same geometric mean diameter and geometric standard deviation, but differ in their abundance and their emission rate, respectively. The same applies to modes BC2 and HOA2. This specific choice is guided by the observations as shown in Table S1 in the Supplement.

230 For the uniform case only two modes are prescribed, with the same geometric mean diameter and standard deviation as modes BC1/HOA1 and BC2/HOA2, respectively (Table S2 in the Supplement). The mass fractions of BC and HOA in these two modes are identical, and they are equal to the mass fractions of the bulk concentrations for the measurement constrained case. Hence the uniform case represents conditions for which the bulk mass concentrations and the size distribution are the same
235 as for the measurement-constrained case, but for which the detailed mixing state information is not available. A comparison of the two cases therefore quantifies the importance of mixing state information at emission for properties of interest, such as optical properties and CCN activation properties.

Figure S10 in the Supplement shows the temporal evolution of the bulk aerosol species over
240 the course of the 24 h simulation. The primary species black carbon and organic carbon increase initially owing to emissions, and then decrease after emissions are discontinued at $t = 12$ h as a result of dilution with the background. The model simulations also show the production of inorganic and organic secondary aerosol species, which condense on the primary particles, continuously modifying the composition of each particle in the aerosol population. Not shown on this figure is the aerosol
245 water content. While the particles start out dry, they take up water around $t = 6$ h when nitrate formation begins.

We calculate the optical properties for ambient conditions (including aerosol water content) at a wavelength of 550 nm and the critical supersaturations of each particle in a post processing step according to Zaveri et al. (2010). For the optical properties we assume spherical particles with a
250 “core-and-shell” configuration in which BC forms the particle core and the other substances compose the shell. We then use Mie calculations (Ackerman and Toon, 1981) to determine the extinction and scattering cross sections for each particle in the population. From these values, the volume extinction, scattering, and absorption coefficients can be reconstructed. Note that we assume all primary and secondary organic substances to be non-absorbing, hence we do not consider any effects due to
255 brown carbon.

To determine the critical supersaturations, we employ κ -Köhler theory (see Zaveri et al. (2010) for details). Based on the critical supersaturation values of each particle in the population, we construct CCN spectra, as shown in Figure S11 in the Supplement for different times throughout the simulation. As the simulation progresses, the CCN/CN fraction at low values of critical supersatura-
260 tion increases, since the the particles become more hygroscopic due to the condensation of secondary aerosol material. Since both primary organic carbon ($\kappa = 0.001$) and black carbon ($\kappa = 0$) are model species with similar and low hygroscopicity parameters, the differences between the uniform case

and the measurement-constrained case are expected to be small, and the two CCN spectra are almost indistinguishable, as we will show in Section 3.3.

265 3 Results and discussion

3.1 Two classes of fresh black carbon particles

Based on single particle SP-AMS measurements in both roadside and non-roadside studies, we identified two types of particles originating from vehicle exhaust, which are primarily composed of refractory black carbon (rBC) and hydrocarbon-like organic aerosol (Fig. 1a – d). One type is dominated by HOA mass (referred to as the “HOA-rich” particle class) and the other is dominated by rBC (“rBC-rich” particle class). The mass spectra of HOA-rich particles (Fig. 1a and c) exhibit fragmentation patterns associated with hydrocarbon structures (e.g., m/z 43 ($C_3H_7^+$), 57 ($C_4H_9^+$), 71 ($C_5H_{11}^+$)), and are consistent with previous aerosol mass spectrometer observations of gasoline/diesel vehicle exhaust and unburned lubricating oil (Massoli et al., 2012; Canagaratna et al., 2004; Mohr et al., 2009). In conjunction with differences in mass spectra, clear differences in particle size distributions (Fig. 2c and d) demonstrate that the division of traffic-related rBC-containing particles into HOA-rich and rBC-rich classes is physically meaningful.

The average mass fractions of rBC (mf_{rBC}) in HOA-rich particles during the non-roadside and roadside studies are low: 0.03 and 0.05, respectively. The narrow distribution of mf_{rBC} (Fig. 3a) suggests that most of the HOA-rich particles are either mixed with a small amount of rBC or do not contain detectable rBC mass, while a very small number of particles contain larger mf_{rBC} (Fig. 3, split axis). Mass spectra of rBC-rich particles are dominated by C_x^+ fragments (i.e., m/z 12, 24, 36, 48 and 60 in Fig. 1b and d) arising from black carbon. Smaller signals associated with HOA-like material in these particles are similar to the primary organic materials derived from diesel engine exhaust observed in laboratory studies (Sage et al., 2008). In contrast to the HOA-rich particles, Figure 3c illustrates a wider range of mf_{rBC} values in rBC-rich particles. In the non-roadside and roadside environments the average mf_{rBC} values (± 1 standard deviation) in this particle class are 0.72 (± 0.18) and 0.86 (± 0.14), respectively, highlighting the possibility that condensation and/or coagulation of HOA material on rBC particles might occur within a short time of emission.

The mass fraction of rBC derived from SP-AMS measurements may be best regarded as a lower limit. For the reasons detailed in Sect. 2.1.2, including the potential for incomplete vaporisation and uncertainty in SP-AMS sensitivity to rBC coating materials, mf_{rBC} values presented here may be underestimated. A 50% overestimation in the mass of HOA would increase mf_{rBC} in rBC-rich particles to 0.85 (± 0.16) and 0.92 (± 0.10) in the non-roadside and roadside studies, respectively (Fig. 3c, dashed lines). In addition to these uncertainties, in the roadside study only rBC-containing particles are detected by the SP-AMS, while in the non-roadside study it is possible that the HOA-rich class includes HOA particles that do not contain rBC (i.e., “externally mixed” HOA).

3.1.1 Differences in particle coating thickness

With single particle detection capability, SP-AMS observations can be used to quantify the thickness of non-refractory particulate matter on individual rBC-containing particles. This calculation focuses
300 only on determining the thickness of HOA coating derived from traffic emissions; however, oxygenated organic aerosol (OOA), biomass burning organic aerosol (BBOA) and secondary inorganic species were mixed with rBC in more aged accumulation mode particles in these urban studies and a similar calculation could be carried out for these particle classes (Lee et al., 2015a, b).

305 We assume a core-shell structure in order to determine the thickness of HOA coating on fresh rBC-containing particles. In addition, we assume a spherical rBC core, a uniform thickness of coating, and an HOA density of 0.9 g cm^{-3} . To simulate the effect of the fractal structure of ambient rBC, the effective density of rBC ($\rho_{\text{eff,rBC}}$) was varied between 0.3 and 1.3 g cm^{-3} ; values which have been observed for laboratory soot standards and engine exhaust (Maricq and Ning, 2004; Gysel et al.,
310 2011). Figure 4 presents the two-dimensional histograms ($m_{\text{f,rBC}}$ vs. particle aerodynamic diameter, d_{va}) of rBC-rich and HOA-rich particles identified by cluster analysis in the two urban studies, with calculated coating thickness curves overlaid for comparison (dashed lines). The coating thickness is relatively insensitive to variations in $\rho_{\text{eff,rBC}}$ (Fig. 4, dashed lines).

These observations demonstrate that rBC-rich particles are only very thinly coated with HOA-
315 material, as compared to HOA-rich particles that have only very small rBC inclusions (Fig. 4). Here we use two specific examples to illustrate that the coating thickness on HOA-rich and rBC-rich particles can vary significantly. First, a 200 nm (d_{va}) rBC-rich particle (i.e., physical particle diameter $\sim 245 \text{ nm}$, assuming $\rho_{\text{eff,rBC}} = 0.8 \text{ g cm}^{-3}$) with $72 - 86 \%$ rBC by mass (the average $m_{\text{f,rBC}}$ of rBC-rich particles) would be covered by a $5 - 11 \text{ nm}$ HOA coating. Second, a 300 nm (d_{va}) HOA-
320 rich particle (i.e., physical particle diameter $\sim 335 \text{ nm}$, assuming $\rho_{\text{eff,rBC}} = 0.8 \text{ g cm}^{-3}$) that co-exists with a small amount of rBC mass ($3 - 5 \%$ by mass, the average $m_{\text{f,rBC}}$ of HOA-rich particles) has an rBC core size of $108 - 119 \text{ nm}$ and a thick HOA coating ($108 - 113 \text{ nm}$).

3.2 Black carbon mass is dominated by black carbon rich particles

Single particle measurements provide direct insight into mixing state at an individual particle level;
325 however, single particle detection has an inherent bias towards large particle sizes because light scattering triggers particle detection (Lee et al., 2015a; Cross et al., 2009; Freutel et al., 2013; Liu et al., 2013b). Ensemble size distributions of rBC peaked at $\sim 100 \text{ nm}$ in both roadside and non-roadside studies indicating that fresh rBC-containing particles emitted from vehicle exhaust are small (Fig. 2a and b). Smaller rBC-containing particles fall below the lower size limit of single particle detection
330 (Fig. 2c and d). Therefore, single particle observations do not necessarily quantify mixing state at the population level. To provide a complementary view of rBC mixing state we turn to ensemble measurements from the roadside study where only rBC-containing particles were detected in the

SP-AMS. Ensemble measurements better represent mixing state on a population basis compared to single particle observations, because they cover a wider range of particle size (i.e., 80 – 1000 nm vacuum aerodynamic diameter, d_{va}).

By measuring very close to a busy road, ensemble SP-AMS observations of rBC and organic aerosol enabled the analysis of more than one hundred vehicle exhaust plumes over the course of the roadside study (Fig. 5a and b). Positive matrix factorization (PMF) (Zhang et al., 2011; Ulbrich et al., 2009) indicates three major sources of rBC-containing aerosol in the roadside environment: transported biomass burning (BBOA), regional background (OOA), and traffic emissions comprised of two PMF factors (HOA-rich and rBC-rich factors, Fig. 5c – f). Inorganic species evident in the bulk aerosol size distributions (Fig. 2a and b) are largely associated with OOA and BBOA factors, while the traffic-related factors contain the majority of rBC and HOA (Lee et al., 2015a, b). Mass spectra of all four PMF factors and a discussion of selection of the number of factors is presented in Supplement Sect. 2. Not previously observed with an AMS, the HOA-rich and rBC-rich factors represent two types of vehicle exhaust plumes with different amounts of rBC and HOA (Fig. 1e and f). The mf_{rBC} of HOA-rich and rBC-rich factors are 0.16 and 0.75, respectively. High-time resolution SP-AMS measurements (Fig. 6) also demonstrated varying organic and rBC levels in vehicle plumes (Fig. 1g and h), with corresponding differences in optical properties (SSA at 405 nm) indicating that rBC-rich plumes were more highly absorbing. The origin of different plume types at this site has been characterized through long-term measurements, and is related to a variety of engine types, operating conditions and pollution control features (Wang et al., 2015).

Mass spectra of HOA-rich and rBC-rich PMF factors (Fig. 1e and f) are strikingly similar to the mass spectra of HOA-rich and rBC-rich particle classes identified in the single particle measurements (Fig. 1c and d), validating the division of traffic emissions into the two factors observed here. The single particle and ensemble mass spectra show some differences. First, the HOA-rich factor has a larger rBC content compared to the HOA-rich particle class; second, the rBC-rich factor contains higher CO^+ and CO_2^+ signals compared to the rBC-rich particle class (inserts of Figs. 1d, e and f). The latter difference arises because highly oxygenated organic fragments (m/z 28 (CO^+) and 44 (CO_2^+)) were excluded in the single particle analysis due to interference of air signals in UMR spectra (Lee et al., 2015a). Differences in mf_{rBC} between single particle and ensemble measurements, especially for HOA-rich particles that contain a small amount of rBC, could be due to the probability of rBC detection in a single particle such that single particle mf_{rBC} may be underestimated relative to ensemble mf_{rBC} Lee et al. (2015a).

With support of direct measurements of single particle mixing state, this work presents the first interpretation of ensemble AMS results in terms of rBC and HOA mixing state. Using the intensity of the two traffic-related factors in plumes (Fig. 5e and f), we estimate that rBC-rich particles account for ~90 % of the observed total traffic-related rBC mass (Fig. 3d). In a similar manner, we estimate that ~60 % of the total HOA mass is due to HOA-rich particles (Fig. 3b). Purple dashed lines shown

370 in Figure 3b and d represent the inclusion of all CO_x^+ fragments (due to surface functionality of ambient rBC (Corbin et al., 2014)) in the calculation of rBC mass (see Supplement Sect. 1), providing similar results. Dashed lines in Figure 3a, b and c represent the effect of a 50 % overestimation of HOA mass (see Supplement Sect. 1), again providing similar results.

3.3 Black carbon mixing state at emission impacts modelled optical properties

375 These novel single particle and ensemble observations of rBC mixing state were used to initialize the particle-resolved aerosol box model PartMC-MOSAIC (see Section 2.3) (Zaveri et al., 2008; Riemer et al., 2009), which simulates the evolution of aerosol due to condensation and coagulation in an idealized urban setting. As described in Section 2.3, we simulate two cases to isolate the impact of mixing state of rBC emissions on aerosol properties. First, a uniform mixing state case in which
380 all particles are assigned identical composition at emission (measured average $m_{f_{\text{rBC}}}$), and second, a measurement-constrained mixing state case where $m_{f_{\text{rBC}}}$ is prescribed directly from our mixing state observations. The evolution of dry $m_{f_{\text{rBC}}}$ over a 24 h period is illustrated in Figure 7a and b for the two cases. As ageing proceeds $m_{f_{\text{rBC}}}$ shifts to lower values, but in the uniform initial composition case no particles ever have $m_{f_{\text{rBC}}}$ larger than $\sim 45\%$. Note that the bulk aerosol composition
385 and number size distributions are identical in the two cases (Fig. S10 in the Supplement), and any differences in optical properties arise only due to the distribution of aerosol components amongst the particles.

Optical properties are determined for each particle in the population using Mie calculations and assuming a core-shell structure. We acknowledge that the underlying assumptions of Mie calculations may not be appropriate for rBC-containing aerosol in the real atmosphere (e.g., Adachi et al.,
390 2011; Scarnato et al., 2013), which is often not spherical and may not exhibit a core-shell configuration (e.g., China et al., 2015). In particular, the enhancement of rBC absorption due to non-absorbing coatings may be overestimated (Cappa et al., 2012; Healy et al., 2015), except in situations where rBC sources and ageing promote extensive coating (Liu et al., 2015; China et al., 2015). However,
395 Mie calculations are commonly applied in regional and global models (Chung and Seinfeld, 2005; Zhao et al., 2013; Fast et al., 2006) to estimate optical properties, and we therefore include this comparison to illustrate the sensitivity to mixing state.

Consistent with previous studies (e.g., Zaveri et al., 2010; Matsui et al., 2013), we observe a difference in volume absorption and scattering coefficients (B_{abs} and B_{scat} , Fig. 7c), and a resulting
400 increase of 0.1 in SSA from the uniform mixing state to the measurement-constrained case that is present during the emission period in the simulation and, notably, persists throughout ageing (Fig. 7d). Note that a direct comparison between measured and modelled SSA is not valid since we measure SSA of the bulk aerosol population and model it only for a subset of this population (i.e., the rBC containing particles). No significant differences in calculated cloud condensation nuclei
405 activity were observed in these simulations (Fig. S11 in the Supplement), owing to the very similar

hygroscopicity of rBC and HOA species. Calculations of aerosol DRF are very sensitive to changes in SSA (McComiskey et al., 2008) (e.g., uncertainties in SSA on the order of only 0.02 can result in a DRF uncertainty of 1 W m^{-2} for a particular particle type (Chin et al., 2009)), illustrating the importance of accurately measuring and simulating mixing state for calculating climatologically relevant aerosol properties.

4 Conclusions

We present mass-based measurements of the mixing state of black carbon-containing aerosol, using a soot-particle aerosol mass spectrometer, from traffic emissions in an urban environment. Observations from single particle mass spectrometry indicate that rBC co-exists with hydrocarbon-like organic aerosol (HOA) in two distinct particle types; those containing a larger mass fraction of rBC, and those containing a larger mass fraction of HOA. Source apportionment of ensemble mass spectral observations using positive matrix factorization also indicates two types of rBC-containing aerosol related to traffic: rBC-rich and HOA-rich aerosol, validated by single particle observations. Ensemble measurements expand the particle size range over which mixing state can be investigated, providing a better insight into the mixing state of the particle population and indicating that approximately 90 % of rBC mass resides in rBC-rich particles. These measurements were used to drive the particle-resolved aerosol box model, PartMC-MOSAIC. Our results indicate an increase in SSA of ~ 0.1 when mixing state at the point of emission is treated accurately in the model, compared to the assumption of uniform mixing state. The approach described here for quantitative assessment of black carbon mixing state from traffic can also be used to assess mixing state from other sources, and to explore the evolution of mixing state during atmospheric processing. Such measurements will be crucial to drive accurate model assessment of black carbon climate impacts on a broader scale.

Acknowledgements. This work was financially supported by Natural Sciences and Engineering Research Council (NSERC) of Canada, Environment Canada, the Canada Foundation for Innovation and the Marie Curie Action FP7-PEOPLE-IOF-2011 (Project: CHEMBC, No. 299755). N.R. and M.W. acknowledge funding from the Department of Energy under Grant DOE DE-SC0011771.

References

- Ackerman, T. and Toon, O.: Absorption of visible radiation in atmospheres containing mixtures of absorbing and nonabsorbing particles, *Applied Optics*, 20, 3661–3668, doi:10.1364/AO.20.003661, 1981.
- 435 Adachi, K. and Buseck, P. R.: Changes of ns-soot mixing states and shapes in an urban area during CalNex, *Journal of Geophysical Research-Atmospheres*, 118, 3723–3730, 2013.
- Adachi, K., Freney, E. J., and Buseck, P. R.: Shapes of internally mixed hygroscopic aerosol particles after deliquescence, and their effect on light scattering, *Geophysical Research Letters*, 38, L13804, doi:10.1029/2011GL047540, L13804, 2011.
- 440 Adachi, K., Zaizen, Y., Kajino, M., and Igarashi, Y.: Mixing state of regionally transported soot particles and the coating effect on their size and shape at a mountain site in Japan, *Journal of Geophysical Research-Atmospheres*, 119, 5386–5396, doi:10.1002/2013JD020880, 2014.
- Arnott, W. P., Moosmüller, H., Rogers, C. F., Jin, T., and Bruch, R.: Photoacoustic spectrometer for measuring light absorption by aerosol: instrument description, *Atmospheric Environment*, 33, 2845 –
- 445 2852, doi:http://dx.doi.org/10.1016/S1352-2310(98)00361-6, http://www.sciencedirect.com/science/article/pii/S1352231098003616, 1999.
- Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z.,
- 450 Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, *Journal of Geophysical Research-Atmospheres*, 118, 5380–5552, 2013.
- Canagaratna, M. R., Jayne, J. T., Ghertner, D. A., Herndon, S., Shi, Q., Jimenez, J. L., Silva, P. J., Williams, P., Lanni, T., Drewnick, F., Demerjian, K. L., Kolb, C. E., and Worsnop, D. R.: Chase studies of particulate
- 455 emissions from in-use New York City vehicles, *Aerosol Science and Technology*, 38, 555–573, 2004.
- Cappa, C. D., Onasch, T. B., Massoli, P., Worsnop, D. R., Bates, T. S., Cross, E. S., Davidovits, P., Hakala, J., Hayden, K. L., Jobson, B. T., Kolesar, K. R., Lack, D. A., Lerner, B. M., Li, S. M., Mellon, D., Nuaaman, I., Olfert, J. S., Petaja, T., Quinn, P. K., Song, C., Subramanian, R., Williams, E. J., and Zaveri, R. A.: Radiative Absorption Enhancements Due to the Mixing State of Atmospheric Black Carbon, *Science*, 337, 1078–1081,
- 460 2012.
- Chan, T. W., Brook, J. R., Smallwood, G. J., and Lu, G.: Time-resolved measurements of black carbon light absorption enhancement in urban and near-urban locations of southern Ontario, Canada, *Atmospheric Chemistry and Physics*, 11, 10407–10432, doi:10.5194/acp-11-10407-2011, http://www.atmos-chem-phys.net/11/10407/2011/, 2011.
- 465 Chin, M., Kahn, R., Rind, D., Feingold, G., Schwartz, S., and DeCola, P.: Atmospheric Aerosol Properties and Climate Impacts, A Report by the U.S. Climate Change Science Program and the Subcommittee on Global Change Research, Tech. Rep. SAP 2.3, NASA, Washington, D.C., 2009.
- China, S., Salvadori, N., and Mazzoleni, C.: Effect of Traffic and Driving Characteristics on Morphology of Atmospheric Soot Particles at Freeway On-Ramps, *Environmental Science and Technology*, 48, 3128–3135,
- 470 2014.

- China, S., Scarnato, B., Owen, R., Zhang, B., Ampadu, M., Kumar, S., Dzepina, K., Dziobak, M., Fialho, P., Perlinger, J., Hueber, J., D. Helmig, Mazzoleni, L., and Mazzoleni, C.: Morphology and mixing state of aged soot particles at a remote marine free troposphere site: Implications for optical properties, *Geophysical Research Letters*, 42, 1243–1250, 2015.
- 475 Chirico, R., DeCarlo, P. F., Heringa, M. F., Tritscher, T., Richter, R., Prevot, A. S. H., Dommen, J., Weingartner, E., Wehrle, G., Gysel, M., Laborde, M., and Baltensperger, U.: Impact of aftertreatment devices on primary emissions and secondary organic aerosol formation potential from in-use diesel vehicles: results from smog chamber experiments, *Atmospheric Chemistry and Physics*, 10, 11 545–11 563, 2010.
- 480 Chung, S. H. and Seinfeld, J. H.: Climate response of direct radiative forcing of anthropogenic black carbon, *Journal of Geophysical Research: Atmospheres*, 110, doi:10.1029/2004JD005441, 2005.
- Corbin, J. C., Sierau, B., Gysel, M., Laborde, M., Keller, A., Kim, J., Petzold, A., Onasch, T. B., Lohmann, U., and Mensah, A.: Mass spectrometry of refractory black carbon particles from six sources: carbon-cluster and oxygenated ions, *Atmospheric Chemistry and Physics*, 14, 2591–2603, 2014.
- 485 Cross, E. S., Onasch, T. B., Canagaratna, M., Jayne, J. T., Kimmel, J., Yu, X. Y., Alexander, M. L., Worsnop, D. R., and Davidovits, P.: Single particle characterization using a light scattering module coupled to a time-of-flight aerosol mass spectrometer, *Atmospheric Chemistry and Physics*, 9, 7769–7793, 2009.
- Dallmann, T., Onasch, T. B., Kirchstetter, T. W., Worton, D., Fortner, E. C., Herndon, S. C., Wood, E. C., Franklin, J. P., Worsnop, D. R., Goldstein, A., and Harley, R.: Characterization of particulate matter emission from on-road gasoline and diesel vehicles using a soot particle aerosol mass spectrometer, *Atmospheric*
- 490 *Chemistry and Physics*, 14, 7585–7599, 2014.
- DeVile, R., Riemer, N., and West, M.: Weighted Flow Algorithms (WFA) for stochastic particle coagulation, *Journal of Computational Physics*, 230, 8427–8451, 2011.
- Fast, J. D., Gustafson, W. I., Easter, R. C., Zaveri, R. A., Barnard, J. C., Chapman, E. G., Grell, G. A., and Peckham, S. E.: Evolution of ozone, particulates, and aerosol direct radiative forcing in the vicinity of Houston using a fully coupled meteorology-chemistry-aerosol model, *Journal of Geophysical Research: Atmospheres*,
- 495 111, doi:10.1029/2005JD006721, 2006.
- Flowers, B. A., Dubey, M. K., Mazzoleni, C., Stone, E. A., Schauer, J. J., Kim, S.-W., and Yoon, S. C.: Optical-chemical-microphysical relationships and closure studies for mixed carbonaceous aerosols observed at Jeju Island; 3-laser photoacoustic spectrometer, particle sizing, and filter analysis, *Atmospheric Chemistry and*
- 500 *Physics*, 10, 10 387–10 398, doi:10.5194/acp-10-10387-2010, <http://www.atmos-chem-phys.net/10/10387/2010/>, 2010.
- Freutel, F., Drewnick, F., Schneider, J., Klimach, T., and Borrmann, S.: Quantitative single-particle analysis with the Aerodyne aerosol mass spectrometer: development of a new classification algorithm and its application to field data, *Atmospheric Measurement Techniques*, 6, 3131–3145, 2013.
- 505 Gysel, M., Laborde, M., Olfert, J. S., Subramanian, R., and Groehn, A. J.: Effective density of Aquadag and fullerene soot black carbon reference materials used for SP2 calibration, *Atmospheric Measurement Techniques*, 4, 2851–2858, 2011.
- Healy, R. M., Sciare, J., Poulain, L., Kamili, K., Merkel, M., Mueller, T., Wiedensohler, A., Eckhardt, S., Stohl, A., Sarda-Estevé, R., McGillicuddy, E., O'Connor, I. P., Sodeau, J. R., and Wenger, J. C.: Sources and mixing

- 510 state of size-resolved elemental carbon particles in a European megacity: Paris, *Atmospheric Chemistry and Physics*, 12, 1681–1700, 2012.
- Healy, R. M., Wang, J. M., Jeong, C.-H., Lee, A. K. Y., Willis, M. D., Jaroudi, E., Zimmerman, N., Hilker, N., Murphy, M., Eckhardt, S., Stohl, A., Abbatt, J. P. D., Wenger, J. C., and Evans, G. J.: Light-absorbing properties of ambient black carbon and brown carbon from fossil fuel and biomass burning sources, *Journal of Geophysical Research: Atmospheres*, 120, 6619–6633, doi:10.1002/2015JD023382, 2015.
- 515 Jacobson, M. Z.: A physically-based treatment of elemental carbon optics: Implications for global direct forcing of aerosols, *Geophysical Research Letters*, 27, 217–220, doi:10.1029/1999GL010968, 2000.
- Jacobson, M. Z.: Strong radiative heating due to the mixing state of black carbon in atmospheric aerosols, *Nature*, 409, 695–697, 2001.
- 520 Jimenez, J. L., Jayne, J. T., Shi, Q., Kolb, C. E., Worsnop, D. R., Yourshaw, I., Seinfeld, J. H., Flagan, R., Zhang, X., Smith, K., Morris, J., and Davidovits, P.: Ambient aerosol sampling using the Aerodyne Aerosol Mass Spectrometer, *Journal of Geophysical Research-Atmospheres*, 108, doi:10.1029/2001JD001213, 2003.
- Johnson, K. S., Zuberi, B., Molina, L. T., Molina, M. J., Iedema, M. J., Cowin, J. P., Gaspar, D. J., Wang, C., and Laskin, A.: Processing of soot in an urban environment: case study from the Mexico City Metropolitan Area, *Atmospheric Chemistry and Physics*, 5, 3033–3043, 2005.
- 525 Knox, A., Evans, G. J., Brook, J. R., Yao, X., Jeong, C. H., Godri, K. J., Sabaliauskas, K., and Slowik, J. G.: Mass Absorption Cross-Section of Ambient Black Carbon Aerosol in Relation to Chemical Age, *Aerosol Science and Technology*, 43, 522–532, 2009.
- Kuwata, M., Kondo, Y., and Takegawa, N.: Critical condensed mass for activation of black carbon as cloud condensation nuclei in Tokyo, *Journal of Geophysical Research-Atmospheres*, 114, doi:10.1029/2009JD012086, 2009.
- 530 Laborde, M., Crippa, M., Tritscher, T., Juranyi, Z., Decarlo, P. F., Temime-Roussel, B., Marchand, N., Eckhardt, S., Stohl, A., Baltensperger, U., Prevot, A. S. H., Weingartner, E., and Gysel, M.: Black carbon physical properties and mixing state in the European megacity Paris, *Atmospheric Chemistry and Physics*, 13, 5831–5856, 2013.
- 535 Lack, D. A., Langridge, J. M., Bahreini, R., Cappa, C. D., Middlebrook, A. M., and Schwarz, J. P.: Brown carbon and internal mixing in biomass burning particles, *Proceedings of the National Academy of Sciences of the United States of America*, 109, 14 802–14 807, 2012.
- Lee, A. K. Y., Willis, M. D., Healy, R. M., Onasch, T. B., and Abbatt, J. P. D.: Mixing state of carbonaceous aerosol in an urban environment: Single particle characterization using a soot particle aerosol mass spectrometer (SP-AMS), *Atmospheric Chemistry and Physics*, 15, 1823–1841, 2015a.
- 540 Lee, A. K. Y., Willis, M. D., Healy, R. M., Wang, J. M., Jeong, C.-H., Wenger, J. C., Evans, G. J., and Abbatt, J. P. D.: Single particle characterization of biomass burning organic aerosol (BBOA): evidence for non-uniform mixing of high molecular weight organics and potassium, *Atmospheric Chemistry and Physics Discussions*, 15, 32 157–32 183, doi:10.5194/acpd-15-32157-2015, <http://www.atmos-chem-phys-discuss.net/15/32157/2015/>, 2015b.
- 545 Liu, D., Allan, J., Whitehead, J., Young, D., Flynn, M., Coe, H., McFiggans, G., Fleming, Z. L., and Bandy, B.: Ambient black carbon particle hygroscopic properties controlled by mixing state and composition, *Atmospheric Chemistry and Physics*, 13, 2015–2029, 2013a.

- 550 Liu, D., Allan, J. D., Young, D. E., Coe, H., Beddows, D., Fleming, Z. L., Flynn, M. J., Gallagher, M. W.,
Harrison, R. M., Lee, J., Prevot, A. S. H., Taylor, J. W., Yin, J., Williams, P. I., and Zotter, P.: Size distribution,
mixing state and source apportionment of black carbon aerosol in London during wintertime, *Atmospheric
Chemistry and Physics*, 14, 10061–10084, 2014.
- Liu, P. F., Zhao, C. S., Göbel, T., Hallbauer, E., Nowak, A., Ran, L., Xu, W. Y., Deng, Z. Z., Ma, N., Mildener-
555 berger, K., Henning, S., Stratmann, F., and Wiedensohler, A.: Hygroscopic properties of aerosol particles
at high relative humidity and their diurnal variations in the North China Plain, *Atmospheric Chemistry and
Physics*, 11, 3479–3494, doi:10.5194/acp-11-3479-2011, <http://www.atmos-chem-phys.net/11/3479/2011/>,
2011.
- Liu, S., Russell, L. M., Sueper, D. T., and Onasch, T. B.: Organic particle types by single-particle measure-
560 ments using a time-of-flight aerosol mass spectrometer coupled with a light scattering module, *Atmospheric
Measurement Techniques*, 6, 187–197, 2013b.
- Liu, S., Aiken, A. C., Gorkowski, K., Dubey, M., Cappa, C. D., Williams, L. R., Herndon, S. C., Massoli, P.,
Fortner, E. C., Chhabra, P., Brooks, W., Onasch, T. B., Jayne, J. T., Worsnop, D., China, S., Sharma, N.,
Mazzoleni, C., Xu, L., Ng, N., Liu, D., Allan, J. D., Lee, J., Fleming, Z. L., Mohr, C., Zotter, P., Szidat, P., and
565 Prevot, A. S. H.: Enhanced light absorption by mixed source black and brown carbon particles in UK winter,
Nature Communications, 6, doi:10.1038/ncomms9435, 2015.
- Maricq, M. M. and Ning, X.: The effective density and fractal dimension of soot particles from premixed flames
and motor vehicle exhaust, *Journal of Aerosol Science*, 35, 1251–1274, 2004.
- Massoli, P., Fortner, E. C., Canagaratna, M. R., Williams, L. R., Zhang, Q., Sun, Y., Schwab, J. J., Trimborn,
570 A., Onasch, T. B., Demerjian, K. L., Kolb, C. E., Worsnop, D. R., and Jayne, J. T.: Pollution Gradients and
Chemical Characterization of Particulate Matter from Vehicular Traffic near Major Roadways: Results from
the 2009 Queens College Air Quality Study in NYC, *Aerosol Science and Technology*, 46, 1201–1218, 2012.
- Matsui, H., Koike, M., Kondo, Y., Moteki, N., Fast, J. D., and Zaveri, R. A.: Development and validation of a
black carbon mixing state resolved three-dimensional model: Aging processes and radiative impact, *Journal
575 of Geophysical Research: Atmospheres*, 118, 2304–2326, doi:10.1029/2012JD018446, 2013.
- McComiskey, A., Schwartz, S., Schmid, B., Guan, H., Lewis, E., Ricchiazzi, P., and Ogren, J.: Direct aerosol
forcing: Calculation from observables and sensitivities to inputs, *Journal of Geophysical Research*, 113,
doi:10.1029/2007JD009170, 2008.
- McMeeking, G. R., Good, N., Petters, M. D., McFiggans, G., and Coe, H.: Influences on the fraction of hy-
580 drophobic and hydrophilic black carbon in the atmosphere, *Atmospheric Chemistry and Physics*, 11, 5099–
5112, 2011a.
- McMeeking, G. R., Morgan, W. T., Flynn, M., Highwood, E. J., Turnbull, K., Haywood, J., and Coe, H.:
Black carbon aerosol mixing state, organic aerosols and aerosol optical properties over the United Kingdom,
Atmospheric Chemistry and Physics, 11, 9037–9052, 2011b.
- 585 Michelotti, M., Heath, M., and West, M.: Binning for efficient stochastic multiscale particle simulations, *Mul-
tiscale Model Simulations*, 11, 1071–1096, 2013.
- Moffet, R. C. and Prather, K. A.: In-situ measurements of the mixing state and optical properties of soot with
implications for radiative forcing estimates, *Proceedings of the National Academy of Sciences of the United
States of America*, 106, 11872–11877, 2009.

- 590 Mohr, C., Huffman, J. A., Cubison, M. J., Aiken, A. C., Docherty, K. S., Kimmel, J. R., Ulbrich, I. M., Han-
nigan, M., and Jimenez, J. L.: Characterization of Primary Organic Aerosol Emissions from Meat Cooking,
Trash Burning, and Motor Vehicles with High-Resolution Aerosol Mass Spectrometry and Comparison with
Ambient and Chamber Observations, *Environmental Science and Technology*, 43, 2443–2449, 2009.
- Moosmüller, H., Chakrabarty, R., and Arnott, W.: Aerosol light absorption and its measure-
595 ment: A review, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 110, 844 – 878,
doi:<http://dx.doi.org/10.1016/j.jqsrt.2009.02.035>, [http://www.sciencedirect.com/science/article/pii/
S0022407309000879](http://www.sciencedirect.com/science/article/pii/S0022407309000879), light Scattering: Mie and More Commemorating 100 years of Mie’s 1908 pub-
lication, 2009.
- Moteki, N., Kondo, Y., Miyazaki, Y., Takegawa, N., Komazaki, Y., Kurata, G., Shirai, T., Blake, D. R.,
600 Miyakawa, T., and Koike, M.: Evolution of mixing state of black carbon particles: Aircraft measurements
over the western Pacific in March 2004, *Geophysical Research Letters*, 34, doi:10.1029/2006GL028943,
2007.
- Onasch, T. B., Trimborn, A., Fortner, E. C., Jayne, J. T., Kok, G. L., Williams, L. R., Davidovits, P., and
Worsnop, D. R.: Soot Particle Aerosol Mass Spectrometer: Development, Validation, and Initial Application,
605 *Aerosol Science and Technology*, 46, 804–817, 2012.
- Oshima, N., Koike, M., Zhang, Y., and Kondo, Y.: Aging of black carbon in outflow from anthropogenic sources
using a mixing state resolved model: 2. Aerosol optical properties and cloud condensation nuclei activities,
Journal of Geophysical Research: Atmospheres, 114, doi:10.1029/2008JD011681, [http://dx.doi.org/10.1029/
2008JD011681](http://dx.doi.org/10.1029/
2008JD011681), d18202, 2009.
- 610 Paatero, P. and Tapper, U.: Positive matrix factorization: A non-negative factor model with optimal utilization
of error estimate of data values, *Environmetrics*, 5, 111–126, 1994.
- Ramanathan, V. and Carmichael, G.: Global and regional climate changes due to black carbon, *Nature Geo-
science*, 1, 221–227, 2008.
- Riemer, N., West, M., Zaveri, R. A., and Easter, R. C.: Simulating the evolution of soot mix-
615 ing state with a particle-resolved aerosol model, *Journal of Geophysical Research-Atmospheres*, 114,
doi:10.1029/2008JD011073, 2009.
- Sabaliauskas, K., Jeong, C.-H., Yao, X., and Evans, G. J.: The application of wavelet decomposition to quan-
tify the local and regional sources of ultrafine particles in cities, *Atmospheric Environment*, 95, 249–257,
doi:<http://dx.doi.org/10.1016/j.atmosenv.2014.05.035>, 2014.
- 620 Sage, A. M., Weitkamp, E. A., Robinson, A. L., and Donahue, N. M.: Evolving mass spectra of the oxidized
component of organic aerosol: results from aerosol mass spectrometer analyses of aged diesel emissions,
Atmospheric Chemistry and Physics, 8, 1139–1152, 2008.
- Scarnato, B. V., Vahidinia, S., Richard, D. T., and Kirchstetter, T. W.: Effects of internal mixing and aggregate
morphology on optical properties of black carbon using a discrete dipole approximation model, *Atmospheric
625 Chemistry and Physics*, 13, 5089–5101, doi:10.5194/acp-13-5089-2013, 2013.
- Schneider, J., Weimer, S., Drewnick, F., Borrmann, S., Helas, G., Gwaze, P., Schmid, O., Andreae, M. O., and
Kirchner, U.: Mass spectrometric analysis and aerodynamic properties of various types of combustion-related
aerosol particles, *International Journal of Mass Spectrometry*, 258, 37–49, 2006.

Seuper, D.: ToF-AMS analysis software, <http://cires.colorado.edu/jimenez-group/ToFAMSResources/ToFSoftware/index.html>, 2010.

630 Shindell, D., Kuylensstierna, J. C. I., Vignati, E., van Dingenen, R., Amann, M., Klimont, Z., Anenberg, S. C., Muller, N., Janssens-Maenhout, G., Raes, F., Schwartz, J., Faluvegi, G., Pozzoli, L., Kupiainen, K., Hoglund-Isaksson, L., Emberson, L., Streets, D., Ramanathan, V., Hicks, K., Oanh, N. T. K., Milly, G., Williams, M., Demkine, V., and Fowler, D.: Simultaneously Mitigating Near-Term Climate Change and Improving Human

635 Health and Food Security, *Science*, 335, 183–189, 2012.

Shiraiwa, M., Kondo, Y., Moteki, N., Takegawa, N., Sahu, L. K., Takami, A., Hatakeyama, S., Yonemura, S., and Blake, D. R.: Radiative impact of mixing state of black carbon aerosol in Asian outflow, *Journal of Geophysical Research-Atmospheres*, 113, doi:10.1029/2008JD010546, 2008.

Toner, S. M., Sodeman, D. A., and Prather, K. A.: Single particle characterization of ultrafine and accumulation

640 mode particles from heavy duty diesel vehicles using aerosol time-of-flight mass spectrometry, *Environmental Science and Technology*, 40, 3912–3921, 2006.

Tritscher, T., Juranyi, Z., Martin, M., Chirico, R., Gysel, M., Heringa, M. F., DeCarlo, P. F., Sierau, B., Prevot, A. S. H., Weingartner, E., and Baltensperger, U.: Changes of hygroscopicity and morphology during ageing of diesel soot, *Environmental Research Letters*, 6, doi:10.1088/1748-9326/6/3/034026, 2011.

645 Ulbrich, I. M., Canagaratna, M. R., Zhang, Q., Worsnop, D. R., and Jimenez, J. L.: Interpretation of organic components from Positive Matrix Factorization of aerosol mass spectrometric data, *Atmospheric Chemistry and Physics*, 9, 2891–2918, 2009.

Wang, J., Jeong, C.-H., Zimmerman, N., Healy, R. M., Wang, D., Ke, F., and Evans, G.: Plume-based analysis of vehicle fleet air pollutant emissions and the contribution from high emitters, *Atmospheric Measurement*

650 *Techniques*, 8, 3263–3275, doi:10.5194/amt-8-3263-2015, 2015.

Willis, M. D., Lee, A. K. Y., Onasch, T. B., Fortner, E. C., Williams, L. R., Lambe, A. T., Worsnop, D. R., and Abbatt, J.: Collection efficiency of the soot-particle aerosol mass spectrometer (SP-AMS) for internally mixed particulate black carbon, *Atmospheric Measurement Techniques*, 7, 4507–4516, 2014.

Zaveri, R., Barnard, J., Easter, R., Riemer, N., and West, M.: Particle-resolved simulation of aerosol size, composition, mixing state, and the associated optical and cloud condensation nuclei activation properties in an

655 evolving urban plume, *Journal of Geophysical Research*, 115, doi:10.1029/2009JD013616, 2010.

Zaveri, R. A. and Peters, L. K.: A new lumped structure photochemical mechanism for large-scale applications, *Journal of Geophysical Research: Atmospheres*, 104, 30 387–30 415, doi:10.1029/1999JD900876, 1999.

Zaveri, R. A., Easter, R. C., and Peters, L. K.: A computationally efficient multicomponent equilibrium solver

660 for aerosols (MESA), *Journal of Geophysical Research: Atmospheres*, 110, doi:10.1029/2004JD005618, 2005a.

Zaveri, R. A., Easter, R. C., and Wexler, A. S.: A new method for multicomponent activity coefficients of electrolytes in aqueous atmospheric aerosols, *Journal of Geophysical Research: Atmospheres*, 110, doi:10.1029/2004JD004681, 2005b.

665 Zaveri, R. A., Easter, R. C., Fast, J. D., and L.K., P.: Model for Simulating Aerosol Interactions and Chemistry (MOSAIC), *Journal of Geophysical Research*, 113, doi:10.1029/2007JD008782, 2008.

- Zhang, G., Bi, X., Li, L., Chan, L. Y., Li, M., Wang, X., Sheng, G., Fu, J., and Zhou, Z.: Mixing state of individual submicron carbon-containing particles during spring and fall seasons in urban Guangzhou, China: a case study, *Atmospheric Chemistry and Physics*, 13, 4723–4735, 2013.
- 670 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Ulbrich, I. M., Ng, N., Worsnop, D., and Sun, Y.: Understanding atmospheric organic aerosols via factor analysis of aerosol mass spectrometry: A review, *Anal. Bioanal. Chem.*, 401, 3045–3067, doi:10.1007/s00216-011-5355-y, 2011.
- Zhang, R. Y., Khalizov, A. F., Pagels, J., Zhang, D., Xue, H. X., and McMurry, P. H.: Variability in morphology, hygroscopicity, and optical properties of soot aerosols during atmospheric processing, *Proceedings of the*
675 *National Academy of Sciences of the United States of America*, 105, 10 291–10 296, 2008.
- Zhao, C., Liu, X., Qian, Y., Yoon, J., Hou, Z., Lin, G., McFarlane, S., Wang, H., Yang, B., Ma, P.-L., Yan, H., and Bao, J.: A sensitivity study of radiative fluxes at the top of atmosphere to cloud-microphysics and aerosol parameters in the community atmosphere model CAM5, *Atmospheric Chemistry and Physics*, 13, 10 969–10 987, 2013.




Figure 1. Unit mass resolution (UMR) spectra of HOA-rich (a) and rBC-rich (b) particle classes identified by *k*-means cluster analysis of single particle SP-AMS data acquired in the non-roadside environment. UMR spectra of HOA-rich (c) and rBC-rich (d) particle classes identified from single particle data in the roadside environment. High-resolution mass spectra of HOA-rich (e) and rBC-rich (f) PMF factors from ensemble SP-AMS data from the roadside study (i.e., for rBC-containing particles only). Examples of UMR spectra collected at high-time resolution during HOA-rich (g) and rBC-rich (h) plumes (corresponding time series are shown in Figure 6) during the roadside study. Insets in the lower panels illustrate the UMR spectrum of organic species (without CO_x^+ fragments) in the rBC-rich mass spectra. Black and colored sticks represent the fragments of rBC (C_x^+) and organic species (C_xH_y^+ , $\text{C}_x\text{H}_y\text{O}_w^+$), respectively.




Figure 2. Particle size distributions from the non-roadside and roadside studies. **(a)** Mass-based ensemble size distribution of refractory black carbon (rBC, black), organic species (Org, green), sulphate (SO_4 , red) and nitrate (NO_3 , blue) for the non-roadside study. **(b)** Mass-based ensemble size distribution from the roadside study. **(c)** Ion-signal based single particle size distributions for HOA-rich particle classes from the roadside (filled green) and non-roadside (blue) studies. **(d)** Ion-signal based single particle size distributions for rBC-rich particle classes from the roadside (filled green) and non-roadside (blue) studies. The SP-AMS measured rBC-containing particles only during the roadside study, while both rBC-containing and non-rBC-containing particles were measured in the non-roadside study.




Figure 3. (a) & (c) The distribution of mf_{rBC} (presented as a normalized histogram) in the HOA-rich (a) and rBC-rich (c) particle classes in roadside (green and black) and non-roadside (blue) environments. Bars and solid lines represent mf_{rBC} calculated from rBC fragments (i.e. C_x^+), and dashed lines represent the impact of a 50% decrease in HOA mass loading, to illustrate the impact of uncertainty in HOA quantification. (b) & (d) Normalized histograms of mf_{Org} in the HOA-rich factor (b) and mf_{rBC} in the rBC-rich factor (d), i.e., frequency distribution showing the fraction of total rBC (HOA) contributed by the rBC-rich (HOA-rich) factor. Bars represent mf_{rBC} calculated by rBC fragments (i.e. C_x^+), whereas purple dashed lines represent the mf_{rBC} calculated including both CO_x^+ and C_x^+ fragments, yielding similar results (see Supplement Sect. 1). Red dashed lines represent the impact of a 50% decrease in HOA mass loading, to illustrate the impact of uncertainty in HOA quantification.




Figure 4. Two-dimensional histograms (mf_{rBC} vs. particle aerodynamic diameter) of rBC-rich (a) and HOA-rich (b) particle classes. The color scale represents the number of particles in each particle class. The dashed lines represent the physical thickness (nm) of organic coating on rBC-containing particles determined by modelling the particles as a core-shell structure. Note that the d_{va} of model outputs are calculated by the physical diameter (d_{p}) and the density (ρ_{p}) of particles (i.e., $d_{\text{va}} = d_{\text{p}} \cdot \rho_{\text{p}}$ where ρ_{p} is the linear combination of mass-weighted HOA and rBC density). To examine the potential effect of fractal structure of ambient rBC particles on predicted coating thickness, the effective density of rBC was varied (green: 0.3 g cm^{-3} , white: 0.8 g cm^{-3} , and red: 1.3 g cm^{-3}).




Figure 5. Time series of (a) ensemble organic aerosol and (b) refractory black carbon mass loadings during the roadside study. Positive matrix factorization (PMF) results for ensemble data (c) to (f): (c) regionally sourced rBC mixed with oxygenated organic aerosol, (d) biomass-burning organic aerosol mixed with rBC, and traffic-related rBC in an rBC-rich and HOA-rich factor (e) and (f). Mass spectra of all PMF factors are shown in Fig. S1 in the Supplement.




Figure 6. Examples of rBC-dominated (a) and organic-dominated (b) plumes observed using the SP-AMS with high-time resolution (1 Hz) sampling, with measurements of single scattering albedo (SSA) at 405 nm from a photoacoustic soot spectrometer (PASS-3). Gaps in SP-AMS data correspond to periods used for the subtraction of gas-phase contributions from particle signals. Mass spectra shown in Figure 1(g) and (h) correspond to plumes in (b) and (a), respectively.




Figure 7. Evolution of dry mf_{rBC} for the measurement-constrained (a) and uniform initial (b) mixing state cases. See Methods and Supplement Sect. 3 for a description of the model input parameters. Evolution of volume absorption (B_{abs}), scattering (B_{scat}) and extinction (B_{ext}) coefficients at 550 nm (c), and single scattering albedo (SSA) at 550 nm over the 24h simulation period (d) for measurement constrained (solid lines) and uniform initial mixing state (dashed lines) cases.