

We thank the reviewer for taking the time to examine our manuscript. Below, the reviewer's comments are reproduced in *green*, and our replies are in black.

Did the authors compare the E-AIM predictions for NH₃ closure with those of another thermodynamic model, such as ISORROPIA?

We have not compared the results of E-AIM with other models such as ISORROPIA, and consider it well beyond the scope of the present manuscript to perform such a comparison, which would be a study in itself. An intercomparison has been done previously for several systems with each model (including AIM-2, a precursor of E-AIM), albeit some time ago, for example in Zhang et al. (2000), who found that the various models are generally comparable, with some differences arising due to the different equilibrium constants used, and the different equilibria considered in the models. E-AIM's strength is in its accuracy; ISORROPIA's strength is in its speed and also its good accuracy. Since speed was not a concern here, we picked the thermodynamics code with the best accuracy. E-AIM has been described as a benchmark model (Zaveri et al., 2008) since it has been written with a minimum of assumptions and simplifications, without compromises to increase computational efficiency. Therefore we consider E-AIM to be a suitable model to perform these calculations with.

We have included the following text in the manuscript:

"E-AIM has been written with the aim of reproducing the thermodynamics of the aerosol system as precisely as possible, without making compromises for the sake of computational efficiency. It is therefore considered a benchmark model (e.g. Zaveri et al., 2008)"

This manuscript contains an overload of data and information. My recommendation is that technical figures that go towards characterizing the chamber rather than experimental data, such as figures 1 and 2, be moved to an online supporting information document. Tables 4 and 5 likewise contain far too much information and could be moved to the SI. A shorter table could be presented that highlights only specific experiments which are discussed in the modeling section.

We believe that the figures mentioned here are an important part of describing the experimental process, and are necessary for understanding what was done. We would like to keep them in the main text. Likewise for the tables mentioned, removing them to a SI section will not improve the readability of the manuscript, as the casual reader can easily skip them. On the other hand, it is frustrating to have to download the manuscript and SI separately and combine them manually if one wants to have the complete paper. Therefore we have not created a SI section.

Finally, the glyoxal experiments presented here are an afterthought, serving more as a proof of concept rather than providing new insight. It is my assessment that they should be removed from this manuscript, expanded upon, and published separately.

We appreciate the suggestion to publish a paper focusing on the glyoxal data, and we are certainly working in that direction, as these experiments are very interesting in their own right. We wish to retain the current plot in this paper however, as it illustrates an important point about the exchange between the gas phase and the chamber walls.

References

Zaveri, R. A., Easter, R. C., Fast, J. D., and Peters, L. K.: Model for Simulating Aerosol Interactions and Chemistry (MOSAIC), *J. Geophys. Res.-Atmos.*, 113, D13204, doi:10.1029/2007jd008782, 2008.

Zhang et al., A comparative review of inorganic aerosol thermodynamic equilibrium modules: similarities, differences, and their likely causes, *Atmospheric Environment* 34 (2000) 117-137

We would like to thank the reviewer for taking the time to evaluate our manuscript. In the response below, the reviewer's questions are reproduced in *green* and our replies are shown in black.

1) Throughout the manuscript: I think the authors confuse the term 'reaction rate' and '(reaction) rate constant (or 'rate coefficient'). A reaction rate dc/dt is the product of the rate constant (k) and reactant concentrations (e.g., $[A]$ and $[B]$), i.e. $dc/dt = k [A] [B]$ (cf e.g. Seinfeld and Pandis, 2006) Therefore, a model includes rate constants that are based on laboratory studies. Reaction rates could only be compared between different studies if the same solute concentrations were present. The text should be carefully checked and revised for the correct use of 'reaction rates' and 'rate constants'.

We have now corrected this throughout the manuscript.

2) p. 33850, l. 11: Did any ice form in any of the experiments? If so, might that be differentiated from liquid droplets? Related to this comment: p. 33852, l. 16: Does 'phase transition' only refer here to gas-to-liquid transition or is any ice phase involved?

During the CLOUD8 campaign, ice was formed at the end of the expansions, when the inlet gas lines to the chamber were re-opened. This issue was fixed before CLOUD9. The presence of an ice phase was detected in the Particle Phase Discriminator (PPD-2K) measurements. The onset of ice formation was also seen in the change of SIMONE forward scattering and depolarization signals. The growth of the aerosol droplets always ended well before the ice formation occurred, and no further growth was observed during or after the ice formation.

"Phase transition" refers to liquid/ice.

We have added the following text to the paper, in section 4.3: "During the CLOUD8 experiments at -10°C , the presence of an ice phase was detected from the PPD-2K measurements. The onset of ice formation was also seen in the change of SIMONE forward scattering and depolarization signals. This ice formation was always detected after the pressure decrease had ended, by which time the aerosol growth had also ceased. No further growth of the aerosol was observed during or after the formation of ice. An upgrade in the chamber expansion system between CLOUD8 and CLOUD9 prevented formation of ice during the CLOUD9 experiments."

We added the following text to the instrument description section:

"In addition, the Particle Phase Discriminator mark 2, Karlsruhe edition (PPD-2K) was used to monitor the phase of particles in the size range of 6 to $60\mu\text{m}$, and thus to detect ice formation. Ice particles and water droplets were differentiated by the PPD-2K based on their forward scattering signals (Vochezer et al. 2016).

3) p. 33857, l. 21: Would it be possible to use E-AIM at pressures other than 1 atm? How much difference in the results could be expected if the variable pressure as in the chamber were considered?

Activities and activity coefficients are relatively weak functions of pressure, so for the modest pressure ranges explored in the CLOUD chamber, variations with pressure are not significant.

4) p. 33858/9: Add units to J_c , D_g , J_k in the equations or text before.

The units for D_g are given on page 33858, line 16 (cm^2s^{-1}). The units for J_c are given on page 33858, line 13 (moles per second). We have added the units for J_k (moles per second) to line 2 of page 33859, immediately before equation 3.

5) What are the assumptions of the mass accommodation coefficients for H₂SO₄ and NH₃? What is the limit between 'high' and 'low' accommodation coefficient?

In the paper of McMurry and Grosjean 1985, they consider the limit to be an accommodation coefficient of 6×10^{-6} . For gases with lower accommodation coefficients, gas phase diffusion does not play a role, as the uptake is slow (kinetic solution). For gases with higher accommodation coefficients, the uptake on the chamber walls is fast enough that the limiting process is diffusion of the gas to the walls (diffusion limited solution). The rate of diffusion depends also on the coefficient of eddy diffusion, accounting for mixing within the chamber. They consider ammonia to have an accommodation coefficient of the order of $1-5 \times 10^{-8}$ for their chamber walls, however, in terms of exchange between the gas phase and the chamber walls, the fastest possible rate is that obtained for the diffusion limited approach. As we find that the diffusion limited rate is far too slow to replenish the gas phase, the accommodation coefficient is less relevant, as values less than 1.0 would only lead to an even slower exchange between the walls and the gas phase.

In the CLOUD chamber, during decompression, the rate of diffusion may increase (an increase in the eddy diffusion coefficient), leading to a more rapid exchange between the walls and the gas phase. We have changed the discussion of diffusion in the paper, with the following text replacing the discussion between page 33868, line 6 and line 22:

"The uptake of gases on chamber walls was investigated by McMurry and Grosjean (1985), who pointed out that when the accommodation coefficient of the gas on the chamber walls is high (in their case higher than approximately 6×10^{-6}), transport is diffusion-limited, with the rate of diffusion also depending on the turbulence in the chamber (parameterised by an eddy diffusion coefficient). For their Teflon chamber, they find an accommodation coefficient of $1.2-4.8 \times 10^{-8}$ for ammonia, suggesting that exchange with the walls occurs even slower than the rate of diffusion. For the CLOUD chamber, previous measurements have shown wall loss rates for sulphuric acid of $1.7 \times 10^{-3} \text{ s}^{-1}$ corresponding to a lifetime of approximately 10 min. Assuming ammonia to behave in a similar way to sulphuric acid in the CLOUD chamber, this suggests that the transport between the walls and the gas phase is orders of magnitude too slow to maintain a constant gas phase ammonia mixing ratio during the cloud formation."

We have also changed the text on page 33869 line 25 to the following:

"It is possible that the pressure change and increased turbulence during the decompression of the chamber lead to a better ventilation of the chamber walls. This would be characterised by a higher eddy diffusion coefficient, increasing the rate of exchange between the gas phase and the walls above the value that is observed during non-decompression periods."

6) p. 33869, l. 2ff: Is the increased evaporation of ammonia from unactivated aerosol greater because of the larger curvature effect or because of differences in water activity? Does E-AIM take into account both of these effects?

This calculation was only performed accounting for the differences in water content of the aerosol/droplets. EAIM does not account for curvature effects.

7) p. 33869: The section about glyoxal gets a bit lost in this section and should be rather made to a separate section. A few more details should be given in terms of the comparison to NH₃, e.g. how comparable are the solubilities (Henry's law constants) and vapor pressure of these gases? Which parameter determines their uptake/evaporation behavior?

We have put the glyoxal discussion in a separate subsection. We have included information on the effective Henry's law coefficients of glyoxal and ammonia. We would prefer not to speculate on the exact mechanism behind the mobilisation of the glyoxal from the chamber walls at this stage, especially as we have no measurements of ammonia that can be

compared to the glyoxal data. Therefore we did not include a discussion of which parameter determines the uptake or evaporation of the gases. We merely suggest that based on the behaviour of glyoxal, there appears to be a rapid mobilisation of gases from the chamber walls to the gas phase, and this may also be the case for ammonia.

8) Table 1: It should be added which equilibria K_1 , K_2 , K_3 refer to. Done

p. 33845, l. 3: CLOUD and CERN should be spelled out here: Done

p. 33858, l. 17: P_{vap} should have lower case p: Done

p. 33864, l. 19: either 'all aerosol is activated' or 'all aerosols are activated': changed to "all aerosol particles are activated"

p. 33867, l. 27/8: 'concentration' misspelled: Fixed

p. 33873, l. 13: Spell out CIGAR: Done

p. 33874, l. 29: received: Fixed

All references end with numbers (Page numbers?). I assume that this happened during copy editing but should be removed. : We shall tell the typesetters to fix this – the numbers are not in the latex file that we are editing, so they must be added by some processing software during typesetting.

Table 3: Is there no T dependence of k_0 ? If so, the minus sign should be removed and '0' added instead.: Correct, there is no temperature dependence. Dash removed, "0" inserted.

Figure 1 (and some other figures) 'shows that' can be omitted: Done.

Aqueous phase oxidation of sulphur dioxide by ozone in cloud droplets

**C. R. Hoyle^{1,2}, C. Fuchs¹, E. Järvinen³, H. Saathoff³, A. Dias⁴, I. El Haddad¹,
M. Gysel¹, S. C. Coburn⁵, J. Tröstl¹, A.-K. Bernhammer^{6,16}, F. Bianchi¹,
M. Breitenlechner⁶, J. C. Corbin¹, J. Craven^{7,a}, N. M. Donahue⁸, J. Duplissy⁹,
S. Ehrhart⁴, C. Frege¹, H. Gordon⁴, N. Höppel³, M. Heinritzi¹⁰, T. B. Kristensen¹¹,
U. Molteni¹, L. Nichman¹², T. Pinterich¹³, A. S. H. Prévôt¹, M. Simon¹⁰, J. G. Slowik¹,
G. Steiner^{9,6,13}, A. Tomé¹⁴, A. L. Vogel⁴, R. Volkamer⁵, A. C. Wagner¹⁰, R. Wagner⁹,
A. S. Wexler¹⁵, C. Williamson^{10,b,c}, P. M. Winkler¹³, C. Yan⁹, A. Amorim¹⁴,
J. Dommen¹, J. Curtius¹⁰, M. W. Gallagher^{12,18}, R. C. Flagan⁷, A. Hansel^{6,16},
J. Kirkby^{4,10}, M. Kulmala⁹, O. Möhler³, F. Stratmann¹¹, D. R. Worsnop^{9,17}, and
U. Baltensperger¹**

¹Laboratory of Atmospheric Chemistry, Paul Scherrer Institute, Villigen, Switzerland

²WSL Institute for Snow and Avalanche Research SLF Davos, Switzerland

³Karlsruhe Institute of Technology, Institute for Meteorology and Climate Research, P.O. Box 3640,
76021 Karlsruhe, Germany

⁴CERN, 1211 Geneva, Switzerland

⁵Department of Chemistry and Biochemistry & CIRES, University of Colorado, Boulder, CO, USA

⁶University of Innsbruck, Institute for Ion Physics and Applied Physics, Technikerstrasse 25, 6020
Innsbruck, Austria

⁷California Institute of Technology, Department of Chemical Engineering, Pasadena, CA, 91125,
USA

⁸Carnegie Mellon University Center for Atmospheric Particle Studies, 5000 Forbes Ave, Pittsburgh,
PA 15213, USA

Abstract

The growth of aerosol due to the aqueous phase oxidation of SO_2 by O_3 was measured in laboratory generated clouds created in the CLOUD chamber at CERN. Experiments were performed at 10 and -10°C , on acidic (sulphuric acid) and on partially to fully neutralised (ammonium sulphate) seed aerosol. Clouds were generated by performing an adiabatic expansion – pressurising the chamber to 220 hPa above atmospheric pressure, and then rapidly releasing the excess pressure, resulting in a cooling, condensation of water on the aerosol and a cloud lifetime of approximately 6 min. A model was developed to compare the observed aerosol growth with that predicted by oxidation rates previously measured in bulk solutions. The model captured the measured aerosol growth very well for experiments performed at 10 and -10°C , indicating that, in contrast to some previous studies, the oxidation rates of SO_2 in a dispersed aqueous system are well represented by accepted rates, based on bulk measurements. To the best of our knowledge, these are the first laboratory based measurements of aqueous phase oxidation in a dispersed, super-cooled population of droplets. The measurements are therefore important in confirming that the extrapolation of currently accepted reaction rates to temperatures below 0°C is correct.

1 Introduction

Sulphur dioxide is an important tropospheric species, influencing air quality as well as the acidity of precipitation (and therefore that of soil, lakes and rivers). It also influences climate directly and indirectly through its oxidation to sulphate and subsequent role in atmospheric new particle formation (e.g. Kulmala, 2003; Riccobono et al., 2014), and the growth of aerosol particles (e.g. Hoppel et al., 1994a) and thus its effect on their cloud condensation nuclei (CCN) properties (e.g. Hobbs, 1971).

Global anthropogenic emissions of SO_2 around the year 1990 were estimated to be approximately 73 Tg S yr^{-1} (Rodhe, 1999), more than twice the total sulphur emissions from natural sources. Similar values were given by Faloon (2009) in a review of modelled SO_2

emissions. In heavily industrialised regions, the ratio of anthropogenic to natural emissions can be higher than 10. Although air quality legislation in Europe and the USA has led to a significant decline of industrial emissions of SO_2 in the last couple of decades, emissions from Asia and developing countries in other locations are increasing (Forster et al., 2007).

The major sink of atmospheric SO_2 is oxidation to sulphate, estimated at approximately 51 Tg Syr^{-1} (Faloona, 2009), and anthropogenic emissions of SO_2 account for approximately 72 % of sulphate aerosol in the atmosphere (Forster et al., 2007).

SO_2 is either oxidised in the gas phase (by reaction with the OH radical), or it can be taken up by cloud droplets and undergo aqueous phase oxidation. In the aqueous phase, oxidation is primarily by reaction with H_2O_2 or O_3 , with the O_3 reaction becoming larger than typical gas phase reaction rates at pH higher than approximately 4 (Seinfeld and Pandis, 2006). Model studies suggest that aqueous oxidation comprises a large majority of the global sulphate production, perhaps 80 % or more (Faloona, 2009; Barth et al., 2000), however the range of model estimates is rather large, reflecting the difficulty in reproducing cloud processes in large scale models.

The aqueous phase oxidation of SO_2 by O_3 proceeds by the absorption of SO_2 into the cloud droplet, and the establishment of equilibrium between $\text{SO}_2 \cdot \text{H}_2\text{O}$, HSO_3^- , and SO_3^{2-} . Dissolved O_3 then oxidises the latter three species, forming SO_4^{2-} . The Henry's law coefficients for dissolution of O_3 and SO_2 , as well as the equilibrium constants for the hydration of SO_2 and the reaction rates for the subsequent reaction with O_3 have all been well studied in bulk solutions in the past (predominantly at temperatures of approximately 20°C or higher, see Sect. 3.2). The oxidation rates recommended by Hoffmann (1986) are commonly adopted in models simulating cloud chemistry (for example, all of the 7 models simulating SO_2 oxidation in clouds, which were compared by Kreidenweis et al., 2003, used these rates). However, to the best of our knowledge, no studies have been performed at temperatures below 0°C , thus values for sub-zero temperatures are based on extrapolations of the temperature dependence at higher temperatures. The temperature dependence recommended by Hoffmann (1986), and adopted in most modelling studies, is that of Erickson et al. (1977), based on measurements at 25 and 16°C . In that study, experiments were

only performed up to a pH of 4.02, where the vast majority (i.e. 0.994) of dissolved S(IV) is present as HSO_3^- . At higher pH, as encountered in cloud droplets, more of the S(IV) is present as SO_3^{2-} , which reacts with O_3 at a rate approximately 4 orders of magnitude faster than that of HSO_3^- . The temperature dependence of this reaction is therefore very important. Measurements performed by Maahs (1983), at 25 and 10 °C, using buffered solutions at pH up to 6.2 suggest a slightly weaker temperature dependence, and therefore greater oxidation rates at low temperatures than suggested by Erickson et al. (1977).

A few previous cloud chamber studies have examined the aqueous phase oxidation of SO_2 in cloud droplets, hydrated aerosol, and fogs, with the aim of reconciling reaction rates measured in bulk solutions with those inferred from S(VI) production in a dispersed aqueous system. As discussed below, the results are mixed, and it is not possible to exclude the influence of un-measured contaminants on the rate of S(IV) oxidation.

The 6.7 m³, Desert Research Institute Dynamic Cloud Chamber was described by Steele et al. (1981), who performed experiments with the oxidation of SO_2 in droplets formed on a range of CCN, without the addition of additional oxidants such as O_3 . The same chamber was then used by Miller et al. (1986) to perform experiments with the addition of O_3 . They used a Na_2SO_4 seed aerosol, and formed clouds by performing adiabatic expansions, which produced temperature decreases from approximately 21 to 10 °C, and cloud lifetimes of 440 to 490 s. Although the chamber could maintain a minimum wall temperature of −40 °C, they only performed experiments beginning around 20 °C, representative of warm clouds. Similarly to the present study, they used O_3 mixing ratios of 120 to 150 ppbv, with a wider range of SO_2 mixing ratios (23 to 310 ppbv). They found good agreement between the measured (dual channel ion chromatograph) increase in S(VI) in the cloud water, and the oxidation rates measured by Erickson et al. (1977) in bulk solutions, suggesting that such rates are indeed applicable to the chemistry occurring in cloud droplets, at warm temperatures.

The aqueous phase oxidation of SO_2 by H_2O_2 and by O_3 was also investigated in the Calspan 590 m³ environmental chamber by Hoppel et al. (1994b). The chamber was filled with outside air that had been filtered through activated charcoal and aerosol filters and experiments were performed at ambient temperature. After humidifying the air and injecting O_3

and SO_2 , they achieved 4 min cloud periods by performing adiabatic expansions from 15 mb over atmospheric pressure to 5 mb below atmospheric pressure. Aqueous phase oxidation rates determined from the resulting aerosol growth were found to be much higher than the bulk measurement based rates recommended by Hoffmann (1986). This experiment was repeated by Caffrey et al. (2001), with improved instrumentation and an improved model, but the measured oxidation rates were still found to be a factor of 5 higher than those of Hoffmann (1986). Small amounts of un-measured ammonia contamination (increasing the droplet pH and raising the reaction rate) were given as a possible explanation.

The CLOUD (Cosmics Leaving Outdoor Droplets) chamber at CERN provides an essentially contaminant free, and precisely controlled environment in which to perform experimental observations of aqueous phase reactions occurring in cloud droplets. In this study, the aqueous phase oxidation rate of SO_2 by O_3 , in cloud droplets is examined. This study is based on measurements performed during two experimental campaigns, in 2013 (CLOUD8) and 2014 (CLOUD9). Experiments were carried out at temperatures of 10 and -10°C , with acidic (H_2SO_4) and partially to fully neutral (ammonium sulphate) aerosol as CCN, and a chemical-microphysical model of the chamber was written to compare commonly accepted reaction rates with the formation of sulphate observed in the cloud chamber droplets.

In the next section, the CLOUD chamber and associated instrumentation is described. In Sect. 3, the details of the model are given, and in Sect. 4 we present a discussion of the results of the experiments and a comparison with the modelled values. Conclusions are drawn in Sect. 5.

2 Experimental setup

The experiments were conducted in the CLOUD chamber at CERN. The chamber consists of a 3 m diameter electro-polished stainless steel cylinder, with a volume of 26.1 m^3 . The temperature in the chamber can be accurately controlled at any point between 183 and 300 K, by regulating the temperature of air flowing between the chamber wall and its thermal

insulation housing (Kirkby et al., 2011). Experiments are performed in a well-mixed flow chamber mode, with the sample air drawn off by the instruments continually being replaced, and the mixing ratio of any added gas phase species being held approximately constant. In CLOUD8, the sample flow, and therefore the addition of gases to the chamber totalled 250 L min^{-1} , leading to a dilution lifetime of 105 min. In CLOUD9, the sample flow demands of the instruments were lower, leading to a flow of 150 L min^{-1} and thus a dilution lifetime of 174 min. As species such as O_3 and SO_2 were continually added to the chamber to maintain approximately constant mixing ratios, the dilution lifetime only applies to the concentration of the aerosol particles.

Gases in the chamber were mixed by two stainless steel fans, mounted at the top and bottom of the chamber, and magnetically coupled to their gearboxes, which are mounted outside the chamber. Between CLOUD8 and CLOUD9, the gearboxes were upgraded to allow a fourfold increase in fan speed, but, for the experiments presented here, the fans were set to the same speed in CLOUD8 and CLOUD9. At the beginning of a series of experiments, the chamber is cleaned by heating to 373 K, flushing with ultra-pure water, and drying with a pure air flow, resulting in extremely low levels of contaminants. The pure air added to the chamber is created by the evaporation of liquid N_2 and liquid O_2 , at a ratio of 79:21. The desired relative humidity (RH) in the chamber is achieved by passing the necessary fraction of the inflow air through a Nafion humidifier, using water which was purified by recirculation through a bank of Millipore Super-Q filters and irradiated with ultra-violet radiation to suppress biological activity (this treatment process is also used to create the water used to clean the chamber). SO_2 is added to the chamber from a gas cylinder ($998.2 \text{ ppmv} \pm 2\%$ in N_2), as is NH_3 for the neutral seed experiments ($1\% \text{ NH}_3$ in N_2). O_3 is created by irradiating a pure air flow at wavelengths below 200 nm, in an external O_3 generator, from which it is piped into the chamber. All gases have dedicated lines for injection into the chamber to avoid contamination and reactions occurring in the gas lines. Fittings and gas lines are made from stainless steel to avoid contamination. More details of the CLOUD chamber are given in (Duplissy et al., 2016).

2.1 Expansion system

By increasing the input flow of air beyond the sample flow drawn off by the instruments, the chamber can be pressurised up to 220 hPa above ambient pressure. This overpressure can be released through an exhaust valve, resulting in an adiabatic cooling, and, at sufficiently high initial RH, the activation of aerosol particles to form cloud droplets. At lower temperatures, ice particles may form. The pressure-release valve is computer controlled, and can be programmed to follow a linear decrease in pressure over a given time period, or to follow any other prescribed pressure profile, such as an initial rapid pressure decrease until a cloud is formed, followed by a slower pressure reduction to maintain the cloud as long as possible. A vacuum pump is mounted in the exhaust line, to ensure that the rate of change of pressure does not decrease with the difference between the chamber and ambient pressure. During and after the adiabatic cooling, the air in the chamber is continually heated by the chamber walls, as the temperature control system is maintained at the pre-expansion temperature, causing the evaporation of the cloud after approximately 4–6 min.

2.2 Temperature and pressure measurement

The temperature and pressure measurements in the CLOUD chamber are described in detail by Dias et al., 2015 (manuscript in preparation). The temperature inside the chamber was measured with a string of 6 thermocouples (TC, type K) which were mounted horizontally between the chamber wall and the centre of the chamber at distances of 100, 170, 270, 400, 650, and 950 mm from the chamber wall. The line of thermocouples was located midway between the top and bottom of the chamber. The TC have a precision of below 0.1 °C, and were calibrated by an in-situ measurement with a string of well calibrated Pt100 sensors (one at each TC position). The temperature measured before, during and after an expansion is shown in Fig. 1. During normal operation, there was no systematic horizontal gradient across the chamber. During expansions, a small oscillation in the temperature was observed. This is caused by the mixing of air within the chamber, with the variation between

the temperature sensors being typically lower than approximately 0.5 °C. In the modelling work here, the temperature was taken as the mean of the 3 innermost TCs (TC4–TC6).

The pressure in the chamber was measured by a Vegabar 51 pressure transmitter with a precision of 0.1 bar.

2.3 Trace gas and aerosol measurement configuration

The gas phase concentrations of SO₂ and O₃ were measured by trace gas monitors (Enhanced Trace Level SO₂ Analyser, Model 43i-TLE, Thermo Scientific and O₃ Analyser, Thermo Environmental Instruments Inc., Model 49C, respectively). A proton transfer reaction time of flight mass spectrometer (PTR-TOF-MS) with a mass resolving power of 4000–5000 ($m/\Delta m$, FWHM) and a mass accuracy within 10 ppm (Graus et al., 2010) was present, however the gas phase ammonia was typically below the detection limit during the experiments. The aerosol particle number size distributions were simultaneously measured by three different scanning mobility particle sizer (SMPS) systems, consisting of a differential mobility sizer (DMA) and a condensation particle counter (CPC, TSI model 3010). The first of these was attached to a total sampling line that allowed the measurement of all aerosols in the chamber. The second was attached to a cyclone, with a cut off of 2 µm, which enabled the measurement of the interstitial (i.e. non-activated) aerosol during the cloud periods. During the subsaturated periods the SMPS attached to the total and the interstitial lines measure essentially the same size distributions. The third SMPS was attached to an additional sampling line, on which a Pumped Counterflow Virtual Impactor (PCVI) was installed with the PCVI flow rate set for a cutoff of 5 µm, thus sampling only the aerosol contained in cloud droplets. The three SMPS systems scanned the size range between 17 and 415 nm approximately every minute. Unfortunately, the PCVI and cyclone appear to have been influenced by pressure changes during the expansions. It is, therefore, not possible to clearly identify the activated fraction of aerosol from these measurements.

The total humidity in the chamber was measured by MBW dew point mirror instruments (model MBW973 during CLOUD9 and MBW373LX during CLOUD8) attached to a heated sampling line. During CLOUD9 a tuneable diode laser (TDL) hygrometer, comparable to

the APicT instrument as described by Fahey et al. (2014), was used to measure the water vapour content in-situ with 1 Hz time resolution using a single optical path once across the middle plane of the CLOUD chamber. From the difference between total water and water vapour, the liquid water content (LWC) or ice water content (IWC) could be calculated. An optical particle sizer (WELAS Promo 2000, Palas GmbH) measured the droplet size distributions during the cloud periods. An in-situ particle backscatter instrument (the SIMONE, Schnaiter et al., 2012) detected the presence of droplets and phase transitions within the chamber. In addition, the Particle Phase Discriminator mark 2, Karlsruhe edition (PPD-2K) was used to monitor the phase of particles in the size range of 6 to 60 μm , and thus to detect ice formation. Ice particles and water droplets were differentiated by the PPD-2K based on their forward scattering signals (Vochezer et al., 2016).

The droplet size distributions were also measured during the cloud periods with the cloud and aerosol spectrometer with polarisation detection (CASPOL), which measures the forward and backward scattering from single particles in the diameter range of 0.6 to 50 μm (Glen and Brooks, 2013).

During both CLOUD8 and CLOUD9, the saturation relative humidity (100 % with respect to liquid water) was determined from the chamber temperature at the time that the SIMONE first detected an increased forward scattering signal, indicating the beginning of droplet growth. As the SIMONE measured in-situ, and was not influenced by transmission effects or temperature changes in the sampling line, it is expected that this provides the most accurate way of determining the dew point and thus the LWC during the cloud periods. From the dew point temperature, the water vapour mixing ratio was calculated. A high and a low value of the dew point was also determined from the SIMONE data, based on the uncertainty in the detection time of the cloud formation with the scattering signal. The total water content of the air in the chamber was assumed to be constant over the time scale of the experiments modelled here. As aerosol growth was only observed during the cloud periods, and the dew point was determined at the beginning of each 4–6 min cloud period, the assumption of constant total water content is not expected to influence the results. This approach was validated during CLOUD9 by comparison with the TDL data. In Fig. 2,

the SIMONE derived and measured peak condensed water during CLOUD9 expansions is plotted. The SIMONE derived values are calculated from the dew point determined as described above, the measured condensed water is found by taking the difference of the TDL (which only measures gas phase water) and the MBW (which measures total water after the droplets have been evaporated). From this comparison, it can be seen that the dew point and thus the total condensed water can be accurately determined using the combination of SIMONE and temperature data.

During both CLOUD8 and CLOUD9, aerosol composition (ammonium to sulphate ratio) was measured with an Aerodyne high-resolution time-of-flight aerosol mass spectrometer (HR-ToF-AMS, DeCarlo et al., 2006). To dry the sample flow, a Nafion dryer (PermaPure) was attached to the sampling line prior to the HR-ToF-AMS, and a pressure controlled inlet (PCI), as described in Bahreini et al. (2008), was used to eliminate the effects of pressure variations in the sampling line on the measured concentrations.

The ionic composition of the aerosol formed is derived from HR-ToF-AMS data using the PIKA software package. A critical parameter required in the model is the ammonium to sulphate ratio (ASR) in the aerosol phase, which influences the pH of the cloud droplets and consequently SO_2 reaction rates. The determination of this ratio heavily depends on the relative ionisation efficiencies (RIE) of ammonium and sulphate. Ammonium RIE, equal to 3.9 ± 0.2 , was determined during the IE calibration by nebulising NH_4NO_3 , whereas the sulphate RIE, equal to 1.38 ± 0.07 , was determined by nebulising $(\text{NH}_4)_2\text{SO}_4$. Another technical limitation that might affect the determination of ammonium concentrations is related to the possible interference of water and oxygen fragments: O^+ and HO^+ . Figure 3 displays the intensities of the main ammonium ions normalised by that of NH_2^+ . The clustering of these ratios around a single value ($\text{NH}_3^+/\text{NH}_2^+ = 1.23 \pm 0.05$ and $\text{NH}^+/\text{NH}_2^+ = 0.072 \pm 0.008$), irrespective of NH_4^+ concentration indicate that the interference from O^+ and HO^+ is negligible in our case.

Measurements of glyoxal (CHOCHO) were performed with the Light Emitting Diode Cavity Enhanced DOAS (LED-CE-DOAS). This instrument is a multispectral sensor that selectively and simultaneously measures glyoxal, oxygen collision complexes ($\text{O}_2\text{-O}_2$), methyl

glyoxal (CH_3COCHO), nitrogen dioxide (NO_2) and other gases utilizing an Ocean Optics QE65000 spectrometer (Thalman and Volkamer, 2010; Coburn et al., 2014). The detection limit of this instrument during the CLOUD9 experiment was determined to be 15–20 pptv at the native one minute measurement resolution employed for this study. The instrument has been extensively compared to gravitational, UV-vis and IR absorption, phosphorescence and mass spectrometric measurement techniques for alpha-dicarbonyls and NO_2 (Thalman et al., 2015) and remote-sensing techniques (Volkamer et al., 2015).

2.4 Seed aerosol formation

Two kinds of seed aerosol were used in these experiments, pure H_2SO_4 , and partially to fully neutralised ammonium sulphate aerosol. The pure H_2SO_4 aerosol was formed in an external CCN generator, which comprised a temperature controlled stainless steel vessel holding a ceramic crucible filled with concentrated H_2SO_4 . After heating the vessel to between 150 and 180 °C, depending on the desired characteristics of the aerosol population, a flow of N_2 was passed through the vessel, above the crucible to transport the hot H_2SO_4 vapour into the chamber. In addition, during CLOUD9, a humidified flow of N_2 was added to the aerosol injection line immediately downstream of the H_2SO_4 vessel, to create more reproducible size distributions. As the vapour cooled in the injection line, H_2SO_4 droplets formed. The partially or fully neutralised aerosol was formed by using the same aerosol generator, and injecting NH_3 directly into the chamber, where it was taken up by the acidic aerosol. The mode diameter of the aerosol distribution produced by this method was approximately 65–75 nm, with a full width at half maximum (FWHM) of approximately 50–70 nm.

2.5 Experimental procedure

A typical experiment began by pressurising the chamber to 220 hPa above ambient pressure, and injecting SO_2 and O_3 so as to reach the desired mixing ratios. Seed particles were then added to the chamber, and observed for approximately 40 min to detect any growth under subsaturated conditions, before the pressure in the chamber was rapidly reduced to

form the cloud. After a 15–30 min waiting time, during which the chamber temperature re-stabilised, the pressure was increased again, and further expansions were performed. This was repeated until the seed aerosol numbers decreased below a few hundred per cubic centimetre. With initial particle number densities of approximately $6000\text{--}8000\text{ cm}^{-3}$, two to four cloud cycles could usually be performed. The properties of the aerosol, cloud droplets, and gas phase species were continually measured during all stages of the experiment.

In order to better illustrate the experimental procedure, it is useful to examine some basic measurements. The SMPS measurements of the aerosol size distribution during one experiment are shown in Fig. 4. At approximately 13:00 UTC, sulphuric acid aerosol was injected into the chamber (not shown). The conditions in the chamber were held constant for approximately two hours, during which time the number of aerosol particles reduced due to dilution. At approximately 15:10 UTC, the first expansion was performed, and a cloud formed (marked in purple in the figure). The resulting aerosol growth can clearly be seen, both in the distribution, and from the mode diameter (line of white dots), which increases from 70 to 90 nm. The chamber was re-pressurised and a second expansion was performed shortly after 17:00 UTC, also resulting in substantial aerosol growth. No growth was observed during sub-saturated periods.

The chamber conditions for the second expansion in Fig. 4, which were used as input for the model, are shown in Fig. 5. Before the expansion, the temperature and pressure were approximately constant at 283 K and 1160 hPa respectively, and the experiment was performed with approximately 18–20 ppbv SO_2 and 120 ppbv O_3 . At approximately 600 s the expansion begins, and the pressure dropped, leading to an adiabatic cooling. The vertical line denotes where the SIMONE first detected the presence of water droplets, indicating that the dew point was approximately 1 K below the chamber temperature before the expansion started. During this expansion, the temperature reached approximately 1.5 K below the dew point, and the liquid water content of the air (LWC) reached a total of 0.7 g m^{-3} . In panel d, it can be seen that the total aerosol number concentration (the integral of the total SMPS size distribution) decreased due to the approximately 20 % pressure reduction during the expansion, and due to the sedimentation and deposition of cloud droplets containing

aerosol. However, before and after the expansion, the measured values remained relatively stable, with a slight decrease due to dilution. In the bottom panel of Fig. 5 the diameter of the 25th and 75th percentile of the particles, as well as the median diameter are plotted as a function of time. The growth in the dry aerosol diameter resulting from aqueous phase chemistry during the cloud is clearly visible. The aerosol growth occurred rapidly as the LWC increases, with growth ending soon after the LWC reached its peak.

3 Model description

A microphysical and chemical box model was constructed to simulate the experiments in the CLOUD chamber. The model was initialised approximately 10 min before each expansion with the dry aerosol size distribution measured by the SMPS attached to the total sampling line. The aerosol sizes measured by the SMPS were not adjusted to account for any remaining water present at the measurement RH (approximately 3 % during CLOUD8 and approximately 20 % during CLOUD9). This will lead to a slight overestimate of the non-water volume of aerosol during CLOUD9, however this effect is similar to the magnitude of the measurement uncertainty of the SMPS instruments (approximately 14 % in volume for particles larger than 50 nm diameter and approximately 22 % in volume for particles smaller than 50 nm in diameter). Subsequently, the model was integrated along a time series of data measured during the chamber experiments. Temperature and pressure, as well as the mixing ratios of SO₂ and O₃ were input at 1 s resolution, and while the actual resolution of the aerosol NH₄⁺ fraction derived from the AMS data was approximately 10 s, this was also interpolated to a 1 s resolution for model input. The water vapour mixing ratio for each experiment was set at the beginning of the simulation (calculated from the dew point temperature, as described in Sect. 2.3), and the sum of the gas and liquid phase water was held constant. The gas phase ammonia mixing ratio was determined for the starting time of the simulation and held constant throughout. As the gas phase ammonia mixing ratio was below the detection limit of the PTR-TOF-MS, this mixing ratio was determined using the Extended Aerosol Inorganic Model (E-AIM) (Clegg et al., 1998; Wexler and Clegg, 2002),

by summing the total amount of sulphate and ammonium present in the aerosol population, and calculating the corresponding equilibrium gas phase ammonia mixing ratio under the RH and temperature conditions of the chamber. The validity and limitations of this approach will be discussed further in Sect. 4.2. As described below, E-AIM is also used to calculate the water activity and vapour pressure over the droplets. E-AIM has been written with the aim of reproducing the thermodynamics of the aerosol system as precisely as possible, without making compromises for the sake of computational efficiency. It is therefore considered a benchmark model (e.g. Zaveri2008).

3.1 Aerosol water content and droplet growth

At sub-saturated conditions, the RH in the chamber during the simulated time periods was typically above 90 %. At RH below 93 %, the hydrated aerosol was assumed to be in equilibrium with the gas phase, and the water activity was calculated, at the maximum model time step of 1 s, using E-AIM. This calculation took into account the amount of SO_4^{2-} and NH_4^+ in the aerosol, as well as the temperature, while the pressure was held constant at 1 atm.

Once the RH reached or exceeded 93 %, a full kinetic calculation of the water uptake into the aerosol or droplets was performed. The Kelvin effect was accounted for in both the equilibrium and the kinetic calculations. At $\text{RH} \geq 93 \%$, the equilibrium vapour pressure of water over the aerosol or droplets, as well as the surface tension (the latter calculated in E-AIM following Dutcher et al., 2010) and the solution density were determined based on interpolation of a pre-calculated lookup table. The lookup table was created using E-AIM, and provides the equilibrium vapour pressure as a function of temperature, $\text{NH}_4^+ : \text{SO}_4^{2-}$ ratio and $\text{H}_2\text{O} : \text{SO}_4^{2-}$ ratio. The lookup table covers the range of $\text{H}_2\text{O} : \text{SO}_4^{2-}$ ratios from 12 to pure water, temperature from 250 to 300 K and $\text{NH}_4^+ : \text{SO}_4^{2-}$ ratio from 0 to 2. The vapour pressure determined from the lookup table was then adjusted to account for the Kelvin effect. The equilibrium vapour pressure over very dilute $\text{NH}_4^+ / \text{SO}_4^{2-} / \text{H}_2\text{O}$ solution droplets (assumed to be pure water) was calculated using the approach of Lowe and Ficke (1974), who parameterised the equation of Goff and Gratch (1946).

The physics of water uptake by aerosol and cloud droplets is well discussed in standard atmospheric physics textbooks, however, as this is the first time the model is described, we believe it beneficial to provide as detailed a description as possible.

When a droplet is large compared to the mean free path of water in air (continuum regime), the total flux of water to a droplet, and thus the uptake (in moles per second), is given by:

$$J_c = \frac{4\pi R_p D_g (p_{\text{H}_2\text{O}} - p_{\text{vap}}) \times 10^{-4}}{RT}, \quad (1)$$

where R_p is the droplet radius in cm, D_g the diffusivity of water in air (in $\text{cm}^2 \text{s}^{-1}$), $p_{\text{H}_2\text{O}}$ the partial pressure of water vapour (in hPa), P_{vap} the vapour pressure of water at the droplet surface (in hPa, calculated from the product of the equilibrium vapour pressure of water at a given temperature (Lowe and Ficke, 1974) and the activity of water in the droplet), R is the gas constant ($8.314 \text{ J mol}^{-1} \text{ K}^{-1}$), and T is the temperature (K). D_g is calculated as

$$D_g = \frac{0.211 \times 1013.0}{P \left(\frac{T}{273.15} \right)^{1.94}}, \quad (2)$$

where P is the pressure in hPa. On the other hand, small particles find themselves in the kinetic regime, with the flux (in moles per second) described by:

$$J_k = \frac{\pi R_p^2 \nu \alpha (p_{\text{H}_2\text{O}} - p_{\text{vap}}) \times 10^{-4}}{RT}, \quad (3)$$

where α is the mass accommodation coefficient for water (assumed to be 1.0). The mean speed of the water molecules (in cm s^{-1}) is given by:

$$\nu = \left(\frac{8 \times 10^{-7} RT}{\pi M_{\text{H}_2\text{O}}} \right)^{\frac{1}{2}}, \quad (4)$$

where $M_{\text{H}_2\text{O}}$ is the molar mass of water (in g mol^{-1}).

The transition from the kinetic to continuum regime is accounted for in the model by the flux matching approach of Fuchs and Sutugin (1971):

$$\frac{J}{J_c} = \frac{1 + Kn}{1 + 1.71Kn + 1.33Kn^2}, \quad Kn = \frac{\lambda}{R_p}, \quad \lambda = \frac{3D_g}{\nu}, \quad (5)$$

where Kn is the Knudsen number (dimensionless) and λ is the mean free path (cm).

The Kelvin effect is accounted for by multiplying p_{vap} by a correction factor, C :

$$C = \exp \frac{2M_{\text{H}_2\text{O}}\sigma}{1 \times 10^7 RT \rho R_p}, \quad (6)$$

where σ is the surface tension and ρ is the density of the $\text{H}_2\text{O}/\text{H}_2\text{SO}_4$ solution at the temperature of interest.

The time step for the calculation of the water flux to or from the droplets was calculated according to the ratio between the flux and the total droplet water content, so that the droplet water content could change by no more than 2 % in a single time step. The maximum time step was restricted to 1 s.

The aerosol and the water droplets were assumed to be in thermal equilibrium with the gas phase at all times. The aerosol concentration varies linearly with the pressure in the chamber, with concentrations decreasing by around 20 % during an expansion due only to the pressure change. During some expansions, particularly when the number of seed aerosol was low, the observed change in particle number concentration was larger than the modelled change, indicating that processes other than the pressure change influenced the particle loss. Corrections for both these effects are applied in the analysis.

3.2 Chemistry

The partitioning of SO_2 and O_3 to the cloud droplets, as well as the subsequent aqueous phase chemistry was calculated on a time step 100 times smaller than that for the water partitioning. Performing the analysis described by Schwartz (1988) showed that, under the

conditions of the experiments presented here (even at -10°C), the partitioning of SO_2 to the cloud droplets can be represented with an effective Henry's law approach, neglecting mass transport limitations. Thus the total amount of S(IV) in the droplet is given by

$$[\text{S(IV)}] = [\text{HSO}_3^-] + [\text{SO}_3^{2-}] + [\text{SO}_2 \cdot \text{H}_2\text{O}] \quad (7)$$

with

$$[\text{HSO}_3^-] = \frac{H_{\text{SO}_2} K_1 p_{\text{SO}_2}}{[\text{H}^+]}, \quad (8)$$

$$[\text{SO}_3^{2-}] = \frac{H_{\text{SO}_2} K_1 K_2 p_{\text{SO}_2}}{[\text{H}^+]^2} \quad (9)$$

and

$$[\text{SO}_2 \cdot \text{H}_2\text{O}] = H_{\text{SO}_2} p_{\text{SO}_2}, \quad (10)$$

where p_{SO_2} is the gas phase partial pressure of SO_2 , K_1 and K_2 are equilibrium constants (Table 1), and H_{SO_2} is the Henry's law coefficient for the dissolution of SO_2 in water (Table 2). H_2SO_4 completely dissociates to HSO_4^- and H^+ . The equilibrium constants given in Table 1 are taken from Seinfeld and Pandis (2006), who, in turn, reported the values recommended by Smith and Martell (1976), based on an evaluation of experimental studies conducted between 1910 and 1974. The heat of dissolution for H_{SO_2} is based on measurements between 25 and 50°C .

The pseudo first order rate coefficient for the aqueous phase oxidation of S(IV) by O_3 is

$$k_{\text{S(IV)}} = (k_0 \times [\text{SO}_2 \cdot \text{H}_2\text{O}]) + (k_1 \times [\text{HSO}_3^-]) + (k_2 \times [\text{SO}_3^{2-}]), \quad (11)$$

with k_0 , k_1 and k_2 given in Table 3. Ozone, on the other hand, is subject to mass transport limitations under the experimental conditions described here, therefore, similarly to Caffrey et al. (2001), we follow the approach of Schwartz (1988), with the change in aqueous phase

O₃ concentration being given by

$$\frac{dC_{\text{aq}}}{dt} = \frac{k_{\text{mt}}}{RT} p_{\text{O}_3} - \frac{k_{\text{mt}}}{H_{\text{O}_3} RT} C_{\text{aq}} - Q R_p, \quad (12)$$

with

$$k_{\text{mt}} = \left[\frac{R_p^2 RT}{3D_g} + \frac{R_p (2\pi M_{\text{O}_3} RT)^{(1/2)}}{3\alpha} \right]^{-1}, \quad (13)$$

$$Q = 3 \left(\frac{\coth q}{q} - \frac{1}{q^2} \right), \quad (14)$$

and

$$q = R_p \sqrt{\frac{k_{\text{S(IV)}}}{D_{\text{aq}}}}. \quad (15)$$

Here, p_{O_3} is the partial pressure of O₃, H_{O_3} is the Henry's law coefficient for O₃, measured by Kosak-Channing and Helz (1983) at temperatures between 5 and 30 °C (Table 2) and C_{aq} is the aqueous phase concentration at the surface of the droplet. The coefficient k_{mt} accounts for the gas and interfacial mass transport limitations. R_p is the radius of the droplet or aerosol particle, M_{O_3} is the molar mass of O₃ and α is the mass accommodation coefficient (4.0×10^{-2} , Müller and Heal, 2002). In Eq. 12, Q , given by Eq. (14), is a correction factor to account for the lower aqueous phase concentrations of O₃ caused by diffusion limited transport within the droplet. D_{aq} is the aqueous phase diffusion coefficient of O₃, taken as a generally representative value of 1×10^{-5} (Schwartz, 1988), and assumed to be temperature independent. D_g is the gas phase diffusion coefficient of O₃, where the typical value of $0.1 \text{ cm}^2 \text{ s}^{-1}$, for a temperature of 298 K, given by Schwartz (1986) is used as a starting point, and scaled for a particular temperature and pressure as follows. According to Poling et al. (2001), the value of the gas phase diffusion coefficient of a gas A in a second gas B

can be calculated as:

$$D_{AB} = \frac{0.00266 T^{3/2}}{P_{\text{bar}} M_{AB}^{1/2} \sigma_{AB}^2 \Omega_D}, \quad (16)$$

where P_{bar} is the pressure in bar, σ_{AB}^2 is a characteristic length in Å, and Ω_D is a dimensionless diffusion collision integral. Knowing the value of D_g at a particular temperature and pressure (298 K, 1 bar) permits the calculation of a constant to represent all terms in Eq. (16), except T and P_{bar} , enabling scaling to other temperatures and pressures:

$$D_g(T_f, P_f) = 1.94 \times 10^{-05} \frac{T_f^{3/2}}{P_f}, \quad (17)$$

Where T_f and P_f are the temperature and pressure of interest.

At the beginning of the chemistry time step, the partial pressures of O_3 and SO_2 , as well as the temperature and pressure and the AMS derived NH_4 mass fraction of the aerosol from the input file were interpolated to the model time. The total NH_4 mass in each aerosol particle or droplet was calculated relative to the S(VI) mass, and the NH_4 and S(VI) concentrations were calculated using the water volume calculated for a particle or droplet in that size bin. Subsequently, the concentration of H^+ in each droplet, and thus the concentration of the other ions was calculated by iteratively solving the electroneutrality equation,

$$[\text{H}^+] + [\text{NH}_4^+]_{\text{AMS}} + [\text{NH}_4^+]_{\text{part}} = [\text{OH}^-] + [\text{HSO}_3^-] + 2[\text{SO}_3^{2-}] + 2[\text{SO}_4^{2-}] + [\text{HSO}_4^-], \quad (18)$$

where $[\text{NH}_4^+]_{\text{AMS}}$ and $[\text{NH}_4^+]_{\text{part}}$ are the NH_4^+ determined from the AMS mass fraction and from the partitioning of gas phase NH_3 into the cloud droplet, respectively. The latter is given by:

$$[\text{NH}_4^+]_{\text{part}} = H_{\text{NH}_3} \times p_{\text{NH}_3} \times K_{\text{NH}_3} \times [\text{H}^+] / K_{\text{H}_2\text{O}}, \quad (19)$$

with p_{NH_3} being the gas phase partial pressure of NH_3 , and the remaining constants defined in Tables 1 and 2.

In Eq. 18, $[\text{HSO}_3^-]$ is given by Eq. (8), $[\text{SO}_3^{2-}]$ by Eq. (9), $[\text{SO}_4^{2-}]$ by

$$[\text{SO}_4^{2-}] = \frac{K_3 [\text{S(VI)}]}{[\text{H}^+] + K_3} \quad (20)$$

and $[\text{HSO}_4^-]$ by

$$[\text{HSO}_4^-] = \frac{[\text{H}^+] [\text{S(VI)}]}{[\text{H}^+] + K_3}. \quad (21)$$

Once the concentrations of the ions are known, the production of S(VI) is calculated from

$$\Delta \text{S(VI)} = Q k_{\text{S(IV)}} [\text{O}_3]_{\text{aq}} dt. \quad (22)$$

The new aqueous phase O_3 concentration is calculated using a backward Eulerian approach to solve Eq. (12), so that a relatively long time step can be used without large fluctuations in $[\text{O}_3]_{\text{aq}}$ which could lead to values that are negative, or exceed the maximum concentration determined by Henry's law.

4 Results

Data from a total of 31 expansions were analysed, 15 at 10°C , and 16 at -10°C . Table 4 lists the most important data describing the different experiments and model runs. SO_2 mixing ratios ranged from 0.9 to 26.4 ppbv, and O_3 from 63.5 to 137.2 ppbv.

4.1 Aerosol activation and cloud droplets

Peak LWCs of up to 1.5 g m^{-3} were seen during the 10°C experiments, and the largest size reached by the median droplets in the modelled size distribution was $17 \mu\text{m}$, with the smallest being $4.9 \mu\text{m}$. Most of the aerosol were activated during the expansions, with modelled activated fractions generally being around 0.9. In one case, however, only 34 % of the aerosol were activated.

The number of modelled cloud droplets (defined as the number of droplets larger than $1\text{ }\mu\text{m}$ in diameter; orange symbols) is compared with the number measured by the WELAS, in Fig. 6, for several of the CLOUD9 experiments. WELAS data from two of the CLOUD9 experiments, and from the CLOUD8 experiments were not available. At aerosol numbers below 1000 cm^{-3} , all aerosol are activated, and the modelled number of droplets matches the number of droplets detected by the WELAS. At higher aerosol concentrations, the modelled and measured numbers of droplets diverge, with the modelled number being higher than that measured by the WELAS. The LWC calculated from the WELAS data is sometimes lower than the LWC derived from the SIMONE, TDL and MBW data, which may explain part of the discrepancy between modelled and WELAS-measured droplet numbers. Further, cooling in the chamber is unlikely to be totally homogeneous, possibly resulting in slightly differing activated fractions of aerosol in different regions of the chamber in the initial stages of the expansion. Much of the clear relationship between the modelled droplet number and that measured by the WELAS can likely be explained by the fact that both depend strongly on the number of aerosol in the chamber (black diamonds in Fig. 6), nevertheless it is encouraging that the modelled and measured droplet numbers are well correlated.

4.2 Ammonia in the CLOUD chamber

At this point, it is useful to examine the assumptions made with regard to the gas phase ammonia, and the amount of ammonium in the aerosol and the cloud droplets in greater detail. In Sect. 3, it was stated that the ASR of the aerosol were measured using an AMS, that these ratios were then used to determine the gas phase ammonia in the chamber, using E-AIM, and that the uptake of this ammonia to cloud droplets was calculated assuming that the droplets were in an effective Henry's law equilibrium with a constant gas phase ammonia concentration. For this approach to be accurate, a number of conditions must be met, each of which are discussed in the subsections below. We find that the pre-expansion determination of the gas phase ammonia mixing ratio is likely to be correct, as is the assumption that the ammonium concentration in the droplets is in equilibrium with the gas

phase ammonia. However, the gas phase ammonia mixing ratio during the cloud periods cannot generally be assumed to be constant.

4.2.1 Accuracy of AMS measurement of ammonia in hydrated aerosol

Firstly, the ammonium to sulphate ratio in the particles must be accurately measured by the AMS. Some technical aspects of this measurement were already discussed in Sect. 2.3. The $\text{NH}_4^+ : \text{SO}_4^{2-}$ ratio used to determine the gas phase ammonia mixing ratio was measured during the subsaturated period immediately preceding each expansion, when the RH in the chamber was approximately 95 %. The calculation of the gas phase ammonia was carried out with E-AIM, using the AMS determined $\text{NH}_4^+ : \text{SO}_4^{2-}$ ratio, at a pressure of one atmosphere and the temperature and RH conditions measured in the chamber. Any reduction in $\text{NH}_4^+ : \text{SO}_4^{2-}$ ratio due to processes such as the drying of the air on the way to the AMS would lead to a lower calculated gas phase ammonia mixing ratio (Fig. 7). Calculations with E-AIM show that under the assumption of a metastable aqueous phase, without the formation of solids, the $\text{NH}_4^+ : \text{SO}_4^{2-}$ ratio can be reduced from 2, to 1.5 during a change in RH from 95 % to 35 %. If solid phases are allowed to form however, the $\text{NH}_4^+ : \text{SO}_4^{2-}$ ratio is much less sensitive to changes in RH. It has been observed that ammonium sulphate particles bounce from the AMS vapouriser, therefore we expect that ammonium sulphate is in a solid state and that there are minimal ammonium losses during the drying process.

In a recent study, nebulised mixtures of ammonium sulphate and ammonium nitrate were measured with an AMS (nitrate activity coefficients of 0, 0.3, 0.5, 0.7 and 1). It was found that the AMS reliably measures the predicted ammonium content of the mixtures of these internally mixed ammonium nitrate and ammonium sulphate particles (Xu et al., manuscript in preparation). We are not aware of any analogous study for aerosol containing only varying ratios of ammonium and sulphate, however if ammonia is not lost from the mixtures described above during the measurement process, we do not expect that it will be lost from aerosol formed from pure ammonium and sulphate solution. Therefore it is likely that the AMS determined ammonium to sulphate ratios are correct.

4.2.2 Uptake of ammonia by a droplet

Since it is possible to estimate the gas phase ammonia before the expansion and cloud formation from the AMS measurements with a reasonable degree of accuracy, the next question is whether or not our assumption of an effective Henry's law equilibrium between droplet and gas phase ammonia is accurate. Using an experiment with a relatively high gas phase ammonia concentration as an example (CLD8_20_11_1b), the uptake of ammonia by a 7 μm diameter water droplet was calculated, under the assumption that the droplet initially contained no ammonium, and was exposed to a 22 pptv ammonia gas phase. The development of the droplet ammonium concentration with time is plotted in Fig. 8, for three different solution pH. At the lowest, $\text{pH} = 4$, equilibrium is reached after approximately 600 s, which is comparable with the time scale of the clouds formed in the chamber (400–600 s). At higher pH, the timescale becomes shorter very rapidly. As shown in Table 4, for most of the experiments, the pH lay between 4 and 5. We therefore expect that the ammonium concentration in the droplets will be essentially equal to the values predicted by assuming an effective Henry's law equilibrium with the gas phase.

4.2.3 Assumption of constant gas phase ammonia

The assumption that the gas phase ammonia is constant during the cloud formation is more difficult to constrain. As shown in Table 5, under the assumption of a constant gas phase, and droplet equilibrium with the gas phase, the amount of ammonia that was taken up into the cloud droplets suspended in a cubic centimeter of air was between approximately 12 and 2500 times the gas phase amount in that same air volume. The experiments with the highest values of this ratio were the ones performed under essentially acidic conditions, such that ammonia was almost absent from the gas phase. The results of these experiments are therefore not sensitive to uncertainties in the gas phase ammonia concentration.

For the experiments with near neutral seed aerosol (for example CLD8_20_11_2a), the droplets would have needed to take up around 500 times the gas phase ammonia in order for our assumption of droplet equilibrium with a constant gas phase to hold. The only possi-

ble sources of ammonia in the chamber during the cloud formation were the chamber walls, any water in the bottom of the chamber or the particles themselves. The total wall area of the chamber is approximately 42 m^2 , which is comparable to the total surface area of the droplets (8.5 m^2 for $10 \text{ }\mu\text{m}$ diameter droplets at a concentration of 1000 cm^{-3} , 34 m^2 for $20 \text{ }\mu\text{m}$ diameter droplets at a concentration of 1000 cm^{-3}). Before the expansion, the walls were in equilibrium with the gas phase and the particles (no changes in the $\text{NH}_4^+ : \text{SO}_4^{2-}$ ratio of the particles were observed before the expansion, unless further ammonia was injected into the chamber), being coated in a mixture of ammonium, sulphuric acid and water. During the expansion, the walls maintained a constant temperature, while the temperature of the gas in the chamber decreased.

The uptake of gases on chamber walls was investigated by McMurry and Grosjean (1985), who pointed out that when the accommodation coefficient of the gas on the chamber walls is high (in their case higher than approximately 6×10^{-6}), transport is diffusion-limited, with the rate of diffusion also depending on the turbulence in the chamber (parameterised by an eddy diffusion coefficient). For their Teflon chamber, they find an accommodation coefficient of $1.2 - 4.8 \times 10^{-8}$ for ammonia, suggesting that exchange with the walls occurs even more slowly than the rate of diffusion. For the CLOUD chamber, previous measurements have shown wall loss rates for sulphuric acid of $1.7 \times 10^{-3} \text{ s}^{-1}$, corresponding to a lifetime of approximately 10 min. Assuming ammonia to behave in a similar way to sulphuric acid in the CLOUD chamber, this suggests that the transport between the walls and the gas phase is orders of magnitude too slow to maintain a constant gas phase ammonia mixing ratio during the cloud formation.

A further possible source of ammonia is the un-activated particles. Once the cloud forms, the ammonia vapour pressure over these deliquesced aerosol will be higher than over the droplets, and ammonia may be transferred via the gas phase from the aerosol to the droplets. However, from the data shown in Table 5, it can be seen that the ammonia contained in the particles was only 2–4 times greater than the additional ammonia required for the droplets to reach their equilibrium value. Using CLD8_20_11_1b as an example again, the maximum ammonium loss from a non-activated aerosol can be calculated, such that the ammonia

vapour pressure over that aerosol reaches the vapour pressure over a droplet. In this case, only approximately 16% of the ammonia can be removed from a non-activated aerosol, making it necessary that as little as 30% of the aerosol activate if the remaining aerosol were to supply the missing ammonia to the droplets. The aerosol in CLD8_20_11_1b had an $\text{NH}_4^+:\text{SO}_4^{2-}$ ratio of 1.67. As shown in Fig. 7, the vapour pressure over the aerosol is highly sensitive to the $\text{NH}_4^+:\text{SO}_4^{2-}$ ratio, particularly as this approaches 2. This means that in other experiments where the $\text{NH}_4^+:\text{SO}_4^{2-}$ ratio was higher, an even smaller fraction of the ammonia in the unactivated aerosol can be given up. As shown in Fig. 6, the model likely overestimates the activation, particularly at larger aerosol numbers, however in order to provide enough ammonia, only a minority of the aerosol could have been activated. A visual inspection of the aerosol diameters measured with the SMPS showed that in all experiments, the majority of the aerosol grew during the cloud periods, and must therefore have activated to form cloud droplets. To summarise, the unactivated aerosol likely released some ammonia during the cloud periods, however this would not have been sufficient to supply the missing ammonia to the cloud droplets.

4.2.4 Glyoxal as an indicator of exchange between chamber walls and the gas phase

During the analysis of other experiments being carried out in the CLOUD chamber (ozone initiated oxidation of isoprene), high precision measurements of gas phase glyoxal were performed. During these measurements, it was found that as soon as the pressure decrease associated with an expansion began, the gas phase glyoxal increased rapidly from around 50 pptv to almost 500 pptv (Fig. 9). Glyoxal is relatively soluble, with an effective Henry's law coefficient of approximately $4 \times 10^5 \text{ M atm}^{-1}$ at 298 K in pure water (Ip et al., 2009), a value that increases by orders of magnitude for solutions containing sulphate (Ip et al., 2009; Kampf et al., 2013). Therefore this gas phase increase occurs in spite of the simultaneous uptake by the droplets. Ammonia is comparably soluble, with the effective Henry's law coefficient ranging between approximately $1 \times 10^4 \text{ M atm}^{-1}$ for a solution pH of 7, and $1 \times 10^7 \text{ M atm}^{-1}$ when the pH is 4 (Seinfeld and Pandis, 2006). It is possible that the pres-

sure change and increased turbulence during the decompression of the chamber lead to a better ventilation of the chamber walls. This would be characterised by a higher eddy diffusion coefficient, increasing the rate of exchange between the gas phase and the walls above the value that is observed during non-decompression periods.

Although high precision ammonia measurements are not available for the CLOUD8 and CLOUD9 experiments, we expect any soluble gas adsorbed onto the chamber walls to respond to the pressure change in a similar way as glyoxal has been shown to behave. In the experiments with neutral or semi-neutral seed aerosol, a large amount of ammonia was injected into the chamber, which was partly taken up by the initially acidic seed, and partly deposited on the chamber walls. A re-mobilisation of ammonia from the chamber walls similar to that observed for glyoxal would lead to a large amount of ammonia in the gas phase, which could be taken up by the droplets.

In summary, it is not possible to support the assumption of a constant gas phase ammonia mixing ratio. It is likely that the droplets take up a large amount of ammonia from the gas phase, due to a rapid mobilisation of ammonia that was previously deposited on the chamber walls. However, it is not possible to quantify the size of this source. It is also likely that a certain amount of ammonia is transferred from the unactivated aerosol to the cloud droplets during the cloud formation.

As mentioned above, the gas phase ammonia is very small when the $\text{NH}_4^+ : \text{SO}_4^{2-}$ ratio is low (below approximately 1.0). Below, we show that the more acidic experiments are uninfluenced by the uncertainty in gas phase ammonia, as for these experiments it was only present at trace levels. We believe it is still interesting to include the more neutralised experiments in the analysis to investigate the sensitivity of the results to ammonium in the droplets and aerosol. In order to perform the model simulations, we maintain the assumption of a constant gas phase, as a base case, and discuss the implications further below.

4.3 Aerosol growth and the effect of ammonia

In this section, the modelled and measured growth of the dry aerosol volume before and after the expansion is discussed.

The pH of droplets formed during the expansions was seldom above 5, due to the uptake of SO_2 and rapid production of SO_4^{2-} at higher pH, and the resulting reduction in pH. The only exceptions were experiments such as CLD8_05_12_1a and CLD8_05_12_1b where there were comparatively high gas phase NH_3 mixing ratios (approximately 21 and 24 pptv respectively, determined with the E-AIM model, as described in Sect. 3). A total of 12 of the experiments were performed at $\text{NH}_4^+ : \text{SO}_4^{2-}$ ratios of greater than 1.

The modelled and measured change in total volume of the aerosol population was calculated by integrating the dry aerosol size distribution (as measured by the SMPS attached to the total sampling line), and subtracting the total volume before the cloud period from that afterwards. Both modelled and measured data were corrected for particle losses during the expansion, by multiplying the volume after the expansion by the ratio of the number density of aerosol before the expansion to that after the expansion.

The dependence of the aerosol growth on the seed aerosol composition is shown in Fig. 10a and b, for the measured and modelled growth respectively. The aerosol growth during the supersaturated periods clearly depends on the aerosol seed composition, with the strength of the effect increasing rapidly at $\text{NH}_4^+ : \text{SO}_4^{2-}$ ratios above approximately 1. This is due to the fact that the more neutral seeds are in equilibrium with a higher gas phase NH_3 amount, which will partition into the cloud droplets when they form. This leads to a higher pH during the cloud phase and accelerated SO_2 oxidation. For the aerosols with an $\text{NH}_4^+ : \text{SO}_4^{2-}$ ratio above 1.2, the additional amount of ammonium which would partition to the droplets under the constant gas phase assumption would have more than balanced the sulphate in the droplets, leading to a very large increase in pH, and correspondingly, a very large increase in aerosol growth. In contrast, at $\text{NH}_4^+ : \text{SO}_4^{2-}$ ratios below approximately 1, there is little ammonia in the gas phase, and it does not influence the aerosol growth. The modelled dependence of aerosol growth on seed composition is very similar to the measured dependence, however Fig. 10b shows that the model systematically predicts greater aerosol growth at 10°C than at -10°C , an effect that is not seen in the measured data (in the experimental data it appears that the lower reaction rates at lower temperatures are balanced by the greater solubility of the gases). This may point to a slightly too large

temperature dependence of the reaction rates, and indeed, using the temperature dependence of Maahs (1983) leads to a slight closing of the gap between the -10°C and the 10°C modelled points (not shown). However, as the rates given in the studies of Erickson et al. (1977) and Maahs (1983) were based on measurements at 25°C , weakening the temperature dependence also increases the modelled growth of aerosol at 10°C . During the CLOUD8 experiments at -10°C , the presence of an ice phase was detected from the PPD-2K measurements. The onset of ice formation was also seen in the change of SIMONE forward scattering and depolarization signals. This ice formation was always detected after the pressure decrease had ended, by which time the aerosol growth had also ceased. No further growth of the aerosol was observed during or after the formation of ice. An upgrade in the chamber expansion system between CLOUD8 and CLOUD9 prevented formation of ice during the CLOUD9 experiments.

A comparison of the modelled and measured aerosol volume growth is shown in Fig. 11a. In this panel, the gas phase ammonia ratio is calculated with E-AIM, as described above, and is held constant during the cloud period. In general, the modelled volume growth matches the measured values to within a factor of two (dot dashed lines in Fig. 11). This indicates that observed aerosol growth in the CLOUD chamber matches the growth predicted by the reaction rates specified by Hoffmann (1986), at both 10 and -10°C . The model appears to slightly overestimate the aerosol growth at 10°C , in the case of the more acidic seed aerosols, while the -10°C points are more evenly distributed about the $1:1$ line. Applying the temperature dependence of Maahs (1983) has the effect of increasing the modelled growth of both the -10°C and the 10°C points (the latter to a slightly lesser degree), and rather decreases the agreement between the modelled and measured values (not shown).

It should once again be noted that the assumption of a constant gas phase ammonia cannot be supported. While the walls almost certainly provide a large source of ammonia, this source cannot be quantified and the relatively good agreement of modelled and measured growth during the 12 partially to fully neutralised experiments must be ascribed to coincidence in each case. As the modelled results of the more neutralised experiments de-

pend on the gas phase NH_3 mixing ratio, two further model runs were performed, the first with doubled gas phase NH_3 (Fig. 11b), the second with no gas phase NH_3 (Fig. 11c), to assess the sensitivity. From panel c, it is clear that the gas phase NH_3 is necessary for the model to reproduce the observed growth of the more neutralised seed aerosol. Particularly in the 10°C experiments at $\text{NH}_4^+ : \text{SO}_4^{2-}$ ratios of close to 2, the modelled aerosol growth is far too low if gas phase NH_3 is neglected. On the other hand, doubling the gas phase NH_3 with respect to the original values determined with E-AIM leads to an overestimation of the aerosol growth. In general however, it can be seen that although the modelled growth is clearly influenced by the gas phase NH_3 , small variations do not greatly impact aerosol growth. It is also clear that for the more acidic aerosol, the results are essentially identical if one assumes no gas phase ammonia, or a constant gas phase ammonia mixing ratio in equilibrium with the aerosol, as the latter is almost negligible.

4.4 Influence of ions on aqueous phase oxidation of SO_2

One of the features of the CLOUD chamber is the ability to perform experiments under conditions with varying concentrations of gas phase ions. By applying a 30 kV electric field across the chamber, ions can effectively be swept from the chamber, providing a neutral environment. In the absence of the clearing field, natural radiation sources such as galactic cosmic rays (GCR) lead to the ionisation of gases, creating ion pair concentrations of approximately 650 cm^{-3} , representative of the number concentrations found in the atmospheric boundary layer. Higher ion-pair concentrations can be achieved by using a pion beam from the CERN proton synchrotron, however for the time period in which the experiments described here were conducted, this was not available. Instead, the CIGAR was used to create higher ion concentrations. The CIGAR is a stainless steel wire, 100 mm in length and $100\text{ }\mu\text{m}$ in diameter, situated in an argon surrounding atmosphere in direct contact with the air in the CLOUD chamber. A voltage of 3.5 kV is applied across the wire, which is just below the voltage necessary to produce a corona discharge. The CIGAR produced ion pair concentrations of approximately 1.5×10^5 to $2.0 \times 10^5\text{ cm}^{-3}$.

SO₂ mixing ratios of 20 ppbv correspond to approximately 5×10^{11} molecules cm⁻³, at 1013.25 hPa and as the aqueous phase oxidation proceeds rapidly, one would not necessarily expect an influence of, at most, a few hundred thousand ions cm⁻³ on the aerosol growth. Nevertheless, a few experiments were performed to confirm that this was the case, and that no unexpected surface charge effects could influence the results. The modelled and measured aerosol growth are again plotted in Fig. 12, with the colours indicating under which conditions the experiments were performed. All except five of the experiments were performed under natural GCR conditions, however the CIGAR and neutral points show no sign of any systematic bias, confirming that ions do not have a measurable influence on SO₂ uptake and oxidation.

5 Conclusions

Experiments have been performed to investigate the aqueous phase oxidation of SO₂ by O₃ in cloud droplets. The observed aerosol growth was compared with the growth predicted by a model using the reaction rates recommended by Hoffmann (1986), which are widely used in the modelling of these reactions. It was shown that the modelled growth under acidic seed aerosol conditions generally agrees with the observed growth, suggesting that the rates of Hoffmann (1986), which were measured in bulk solutions, do indeed accurately represent the chemistry occurring in dispersed aqueous systems. This contrasts with the findings of at least two previous studies (Hoppel et al., 1994b; Caffrey et al., 2001), and seems to confirm the suggestion made by those authors that the observed disagreement resulted from the presence of contaminants such as ammonia. Furthermore, we have performed what we believe to be the first laboratory based measurements of the aqueous phase oxidation of SO₂ in a population of super-cooled cloud droplets, and confirmed that the generally accepted temperature dependence of the oxidation reactions, measured by Erickson et al. (1977), is consistent with our experimental results at temperatures of -10 °C.

Experiments were also performed for partially to essentially fully neutralised aerosol (ammonium sulphate), however as there is likely to be a large, unquantifiable source of ammo-

nia from the chamber walls during the cloud formation, these experiments can only be used to illustrate the sensitivity of modelled aerosol growth to ammonia.

The agreement between the model and the experimental data presented here illustrates that chamber experiments, performed under well defined conditions, may be used to determine aqueous phase reaction rates. Additionally, such experiments may be used to determine reaction rates in super-cooled droplets, which are important for atmospheric applications but impossible to measure in bulk solutions.

Acknowledgements. We thank Martin Schnaiter for his assistance with the SIMONE and PPD-2K data interpretation. The PPD-2K was made available by funding from the Deutsche Forschungsgemeinschaft under grant SCHN 1140/2-1. C. R. Hoyle was supported by the Swiss National Science Foundation (SNSF) (grant number 200021_140663). T. B. Kristensen gratefully acknowledges funding from the German Federal Ministry of Education and Research (BMBF) through the CLOUD12 project. J. Craven received funding through the Dreyfus Award EP-11-117. N. M. Donahue received funding through US National Science Foundation Grants AGS-1447056 and AGS-1439551. This research has received funding from the EC Seventh Framework Programme (Marie Curie Initial Training Network “CLOUD-TRAIN” grant no. 316662, and the German Federal Ministry of Education and Research (project no. 01LK1222A and B).

References

- Barth, M. C., Rasch, P. J., Kiehl, J. T., Benkovitz, C. M., and Schwartz, S. E.: Sulfur chemistry in the National Center for Atmospheric Research Community Climate Model: description, evaluation, features, and sensitivity to aqueous chemistry, *J. Geophys. Res.*, 105, 1387–1415, doi:10.1029/1999JD900773, 2000.
- Caffrey, P., Hoppel, W., Frick, G., Pasternack, L., Fitzgerald, J., Hegg, D., Gao, S., Leaitch, R., Shantz, N., Albrechtski, T., and Ambrusko, J.: In-cloud oxidation of SO₂ by O₃ and H₂O₂: cloud chamber measurements and modeling of particle growth, *J. Geophys. Res.*, 106, 27587–27601, doi:10.1029/2000JD900844, 2001.
- Clegg, S., Brimblecombe, P., and Wexler, A.: Thermodynamic model of the system H⁺-NH₄⁺-SO₄²⁻-NO₃-H₂O at tropospheric temperatures, *J. Phys. Chem.-US*, 102, 2137–2154, doi:10.1021/jp973042r, 1998.

- Coburn, S., Ortega, I., Thalman, R., Blomquist, B., Fairall, C. W., and Volkamer, R.: Measurements of diurnal variations and eddy covariance (EC) fluxes of glyoxal in the tropical marine boundary layer: description of the Fast LED-CE-DOAS instrument, *Atmos. Meas. Tech.*, 7, 3579–3595, doi:10.5194/amt-7-3579-2014, 2014.
- DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R., and Jimenez, J. L.: Field-deployable, high-resolution, time-of-flight aerosol mass spectrometer, *Anal. Chem.*, 78, 8281–8289, doi:10.1021/ac061249n, 2006.
- Duplissy, J., Merikanto, J., Franchin, A., Tsagkogeorgas, G., Kangasluoma, J., Wimmer, D., Vuollekoski, H., Schobesberger, S., Lehtipalo, K., Flagan, R., Brus, D., Donahue, N., Vehkämäki, H., Almeida, J., Amorim, A., Barmet, P., Bianchi, F., Breitenlechner, M., Dunne, E., Guida, R., Henschel, H., Junninen, H., Kirkby, J., Kürten, A., Kupc, A., Määttänen, A., Makhmutov, V., Mathot, S., Nieminen, T., Onnela, A., Praplan, A., Riccobono, F., Rondo, L., Steiner, G., Tome, A., Walther, H., Baltensperger, U., Carslaw, K., Dommen, J., Hansel, A., Petäjä, T., Sipilä, M., Stratmann, F., Vrtala, A., Wagner, P., Worsnop, D., Curtius, J., and Kulmala, M.: Effect of ions on sulfuric acid-water binary particle formation II: Experimental data and comparison with QC-normalized classical nucleation theory, *Journal of Geophysical Research: Atmospheres*, doi:10.1002/2015JD023539, 2016.
- Dutcher, C. S., Wexler, A. S., and Clegg, S. L.: Surface tensions of inorganic multicomponent aqueous electrolyte solutions and melts, *J. Phys. Chem.-US*, 114, 12216–12230, doi:10.1021/jp105191z, 2010.
- Erickson, R., Yates, L., Clark, R., and Mcewen, D.: Reaction of sulfur-dioxide with ozone in water and its possible atmospheric significance, *Atmos. Environ.*, 11, 813–817, doi:10.1016/0004-6981(77)90043-9, 1977.
- Fahey, D. W., Gao, R.-S., Möhler, O., Saathoff, H., Schiller, C., Ebert, V., Krämer, M., Peter, T., Amarouche, N., Avallone, L. M., Bauer, R., Bozóki, Z., Christensen, L. E., Davis, S. M., Durr, G., Dyroff, C., Herman, R. L., Hunsmann, S., Khaykin, S. M., Mackrodt, P., Meyer, J., Smith, J. B., Spelten, N., Troy, R. F., Vömel, H., Wagner, S., and Wienhold, F. G.: The AquaVIT-1 intercomparison of atmospheric water vapor measurement techniques, *Atmos. Meas. Tech.*, 7, 3177–3213, doi:10.5194/amt-7-3177-2014, 2014.
- Faloona, I.: Sulfur processing in the marine atmospheric boundary layer: a review and critical assessment of modeling uncertainties, *Atmos. Environ.*, 43, 2841–2854, doi:10.1016/j.atmosenv.2009.02.043, 2009.

- Forster, P., Ramaswamy, V., Artaxo, P., Berntsen, T., Betts, R., Fahey, D., Haywood, J., Lean, J., Lowe, D., Myhre, G., Nganga, J., Prinn, R., Raga, G., Schulz, M., and Van Dorland, R.: Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, chap. 2, Changes in Atmospheric Constituents and in Radiative Forcing, Cambridge University Press, Cambridge, UK and New York, NY, USA, 2007.
- Fuchs, N. A. and Sutugin, A. G.: High dispersed aerosols, in: Topics in Current Aerosol Research, edited by: Hidy, G. M., and Brock, J. R., Pergamon, New York, USA, 1–60, 1971.
- Glen, A. and Brooks, S. D.: A new method for measuring optical scattering properties of atmospherically relevant dusts using the Cloud and Aerosol Spectrometer with Polarization (CASPOL), Atmos. Chem. Phys., 13, 1345–1356, doi:10.5194/acp-13-1345-2013, 2013.
- Goff, J. A. and Gratch, S.: Low-pressure properties of water from -160 to 212°F , T. Am. Soc. Heat. Vent. Eng., 52, 95–121, 1946.
- Graus, M., Mueller, M., and Hansel, A.: High resolution PTR-TOF: quantification and formula confirmation of VOC in real time, J. Am. Soc. Mass Spectr., 21, 1037–1044, doi:10.1016/j.jasms.2010.02.006, 2010.
- Hobbs, P. V.: Simultaneous airborne measurements of cloud condensation nuclei and sodium-containing particles over the ocean, Q. J. Roy. Meteorol. Soc., 97, 263–271, doi:10.1002/qj.49709741302, 1971.
- Hoffmann, M.: On the kinetics and mechanism of oxidation of aquated sulfur-dioxide by ozone, Atmos. Environ., 20, 1145–1154, doi:10.1016/0004-6981(86)90147-2, 1986.
- Hoppel, W., Frick, G., Fitzgerald, J., and Larson, R.: Marine boundary-layer measurements of new particle formation and the effects nonprecipitating clouds have on aerosol-size distribution, J. Geophys. Res., 99, 14443–14459, doi:10.1029/94JD00797, 1994a.
- Hoppel, W., Frick, G., Fitzgerald, J., and Wattle, B.: A cloud chamber study of the effect that nonprecipitating water clouds have on the aerosol-size distribution, Aerosol Sci. Tech., 20, 1–30, doi:10.1080/02786829408959660, 1994b.
- Ip, H. S. S., Huang, X. H. H., and Yu, J. Z.: Effective Henry's law constants of glyoxal, glyoxylic acid, and glycolic acid, , 36, doi:10.1029/2008GL036212, 2009.
- Kampf, C. J., Waxman, E. M., Slowik, J. G., Dommen, J., Pfaffenberger, L., Praplan, A. P., Prevot, A. S. H., Baltensperger, U., Hoffmann, T., and Volkamer, R.: Effective Henry's law partitioning and the salting constant of Glyoxal in Aerosols Containing Sulfate, Environmental Science & Technology, 47, 4236–4244, doi:10.1021/es400083d, 2013.

- Kirkby, J., Curtius, J., Almeida, J., Dunne, E., Duplissy, J., Ehrhart, S., Franchin, A., Gagne, S., Ickes, L., Kuerten, A., Kupc, A., Metzger, A., Riccobono, F., Rondo, L., Schobesberger, S., Tsagko-georgas, G., Wimmer, D., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Downard, A., Ehn, M., Flagan, R. C., Haider, S., Hansel, A., Hauser, D., Jud, W., Junninen, H., Kreissl, F., Kvashin, A., Laaksonen, A., Lehtipalo, K., Lima, J., Lovejoy, E. R., Makhmutov, V., Mathot, S., Mikkila, J., Minginette, P., Mogo, S., Nieminen, T., Onnela, A., Pereira, P., Petaja, T., Schnitzhofer, R., Seinfeld, J. H., Sipila, M., Stozhkov, Y., Stratmann, F., Tome, A., Vanhanen, J., Viisanen, Y., Vrtala, A., Wagner, P. E., Walther, H., Weingartner, E., Wex, H., Winkler, P. M., Carslaw, K. S., Worsnop, D. R., Baltensperger, U., and Kulmala, M.: Role of sulphuric acid, ammonia and galactic cosmic rays in atmospheric aerosol nucleation, *Nature*, 476, 429–433, doi:10.1038/nature10343, 2011.
- Kosak-Channing, L. and Helz, G.: Solubility of ozone in aqueous-solutions of 0–0.6 M ionic-strength at 5–30 °C, *Environ. Sci. Technol.*, 17, 145–149, doi:10.1021/es00109a005, 1983.
- Kreidenweis, S., Walcek, C., Feingold, G., Gong, W., Jacobson, M., Kim, C., Liu, X., Penner, J., Nenes, A., and Seinfeld, J.: Modification of aerosol mass and size distribution due to aqueous-phase SO₂ oxidation in clouds: Comparisons of several models, *J. Geophys. Res.*, 108, 4213, doi:10.1029/2002JD002697, 2003.
- Kulmala, M.: How particles nucleate and grow, *Science*, 302, 1000–1001, doi:10.1126/science.1090848, 2003.
- Lowe, P. R. and Ficke, J. M.: The Computation of Saturation Vapor Pressure, Tech. Rep., Environmental Prediction Research Facility (Navy), Monterey, CA, USA, 1974.
- Maahs, H.: Kinetics and mechanism of the oxidation of S(IV) by ozone in aqueous-solution with particular reference to SO₂ conversion in nonurban tropospheric clouds, *J. Geophys. Res.*, 88, 721–732, doi:10.1029/JC088iC15p10721, 1983.
- McMurry, P. H. and Grosjean, D.: Gas and aerosol wall losses in Teflon film smog chambers, *Environ. Sci. Technol.*, 19, 1176–1182, doi:10.1021/es00142a006, 1985.
- Miller, D., Gertler, A., Whitbeck, M., and Lamb, D.: Ozone-induced oxidation of SO₂ in simulated clouds, *J. Geophys. Res.*, 91, 14439–14444, doi:10.1029/JD091iD13p14439, 1986.
- Müller, B., and Heal, M.: The mass accommodation coefficient of ozone on an aqueous surface, *Phys. Chem. Chem. Phys.*, 4, 3365–3369, doi:10.1039/b202491h, 2002.
- Pandis, S. and Seinfeld, J.: Sensitivity analysis of a chemical mechanism for aqueous-phase atmospheric chemistry, *J. Geophys. Res.*, 94, 1105–1126, doi:10.1029/JD094iD01p01105, 1989.

- Poling, B. E., Prausnitz, J. M., and O'Connell, J. P.: The Properties of Gases and Liquids, McGraw-Hill, New York, 2001.
- Riccobono, F., Schobesberger, S., Scott, C. E., Dommen, J., Ortega, I. K., Rondo, L., Almeida, J., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Downard, A., Dunne, E. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, A., Hansel, A., Junninen, H., Kajos, M., Keskinen, H., Kupc, A., Kuerten, A., Kvashin, A. N., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Nieminen, T., Onnela, A., Petaja, T., Praplan, A. P., Santos, F. D., Schallhart, S., Seinfeld, J. H., Sipila, M., Spracklen, D. V., Stozhkov, Y., Stratmann, F., Tome, A., Tsagkogeorgas, G., Vaattovaara, P., Viisanen, Y., Vrtala, A., Wagner, P. E., Weingartner, E., Wex, H., Wimmer, D., Carslaw, K. S., Curtius, J., Donahue, N. M., Kirkby, J., Kulmala, M., Worsnop, D. R., and Baltensperger, U.: Oxidation products of biogenic emissions contribute to nucleation of atmospheric particles, *Science*, 344, 717–721, doi:10.1126/science.1243527, 2014.
- Rodhe, H.: Human impact on the atmospheric sulfur balance, *Tellus A*, 51, 110–122, 1999.
- Schnaiter, M., Büttner, S., Möhler, O., Skrotzki, J., Vragel, M., and Wagner, R.: Influence of particle size and shape on the backscattering linear depolarisation ratio of small ice crystals – cloud chamber measurements in the context of contrail and cirrus microphysics, *Atmos. Chem. Phys.*, 12, 10465–10484, doi:10.5194/acp-12-10465-2012, 2012.
- Schwartz, S. E.: Mass transport considerations pertinent to aqueous phase reactions of gases in liquid water clouds, in: *Chemistry of Multiphase Atmospheric Systems*, edited by: Jaeschke, W., NATO ASI series Series G: Ecological Sciences, 6, 415–471, Springer-Verlag Berlin Heidelberg New York Tokyo, 1986.
- Schwartz, S.: Mass-transport limitation to the rate of in-cloud oxidation of SO₂ – re-examination in the light of new data, *Atmos. Environ.*, 22, 2491–2499, doi:10.1016/0004-6981(88)90480-5, 1988.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics, From Air Pollution to Climate Change*, Wiley, New York, USA, 2006.
- Smith, R. M. and Martell, A. E.: *Critical Stability Constants*, vol. 4: Inorganic Complexes, Plenum, New York, 1976.
- Steele, R., Gertler, A., Katz, U., Lamb, D., and Miller, D.: Cloud chamber studies of dark transformations of sulfur-dioxide in cloud droplets, *Atmos. Environ.*, 15, 2341–2352, 1981.
- Thalman, R. and Volkamer, R.: Inherent calibration of a blue LED-CE-DOAS instrument to measure iodine oxide, glyoxal, methyl glyoxal, nitrogen dioxide, water vapour and aerosol extinction in open cavity mode, *Atmos. Meas. Tech.*, 3, 1797–1814, doi:10.5194/amt-3-1797-2010, 2010.

- Thalman, R., Baeza-Romero, M. T., Ball, S. M., Borrás, E., Daniels, M. J. S., Goodall, I. C. A., Henry, S. B., Karl, T., Keutsch, F. N., Kim, S., Mak, J., Monks, P. S., Muñoz, A., Orlando, J., Peppe, S., Rickard, A. R., Ródenas, M., Sánchez, P., Seco, R., Su, L., Tyndall, G., Vázquez, M., Vera, T., Waxman, E., and Volkamer, R.: Instrument intercomparison of glyoxal, methyl glyoxal and NO₂ under simulated atmospheric conditions, *Atmos. Meas. Tech.*, 8, 1835–1862, doi:10.5194/amt-8-1835-2015, 2015.
- Vochezer, P., Järvinen, E., Wagner, R., Kupiszewski, P., Leisner, T., and Schnaiter, M.: In situ characterization of mixed phase clouds using the Small Ice Detector and the Particle Phase Discriminator, *Atmospheric Measurement Techniques*, 9, 159–177, doi:10.5194/amt-9-159-2016, <http://www.atmos-meas-tech.net/9/159/2016/>, 2016.
- Volkamer, R., Baidar, S., Campos, T. L., Coburn, S., DiGangi, J. P., Dix, B., Eloranta, E. W., Koenig, T. K., Morley, B., Ortega, I., Pierce, B. R., Reeves, M., Sinreich, R., Wang, S., Zondlo, M. A., and Romashkin, P. A.: Aircraft measurements of BrO, IO, glyoxal, NO₂, H₂O, O₂–O₂ and aerosol extinction profiles in the tropics: comparison with aircraft-/ship-based in situ and lidar measurements, *Atmos. Meas. Tech.*, 8, 2121–2148, doi:10.5194/amt-8-2121-2015, 2015.
- Wexler, A. and Clegg, S.: Atmospheric aerosol models for systems including the ions H⁺, NH₄⁺, Na⁺, SO₄²⁻, NO₃⁻, Cl⁻, Br⁻, and H₂O, *J. Geophys. Res.*, 107, 4207, doi:10.1029/2001JD000451, 2002.
- Zaveri, R. A., Easter, R. C., Fast, J. D., and Peters, L. K.: Model for Simulating Aerosol Interactions and Chemistry (MOSAIC), *J. Geophys. Res.*, 113, D13204, doi:10.1029/2007JD008782, 2008.

Table 1. Equilibrium constants used in the model. The temperature dependence of the equilibrium constants is given by $K = K_{298} \exp \left[-\frac{\Delta H}{R} \left(\frac{1}{T} - \frac{1}{298} \right) \right]$, where K is the equilibrium constant at temperature T in Kelvin. K_1 refers to the equilibrium $\text{SO}_2 \cdot \text{H}_2\text{O} \rightleftharpoons \text{HSO}_3^- + \text{H}^+$, K_2 refers to the equilibrium $\text{HSO}_3^- \rightleftharpoons \text{SO}_3^{2-} + \text{H}^+$ and K_3 to the equilibrium $\text{HSO}_4^- \rightleftharpoons \text{SO}_4^{2-} + \text{H}^+$.

	K_{298} [M]	$-\Delta H/R$ [K]	Reference
K_1	1.3×10^{-2}	1960	Seinfeld and Pandis (2006); Smith and Martell (1976)
K_2	6.6×10^{-8}	1500	Seinfeld and Pandis (2006); Smith and Martell (1976)
K_3	1.02×10^{-2}	2720	Seinfeld and Pandis (2006); Smith and Martell (1976)
K_{NH_3}	1.7×10^{-5}	−4353.09	Seinfeld and Pandis (2006)
$K_{\text{H}_2\text{O}}$	1.0×10^{-14}	−6710	Seinfeld and Pandis (2006); Smith and Martell (1976)

Table 2. Henry's law coefficients and heats of dissolution. The temperature dependence of the coefficients is given by $H_A(T_2) = H_A(T_1) \exp \left[\frac{\Delta H_A}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right) \right]$.

	H [M atm ⁻¹ at 298 K]	ΔH_A [kcal mol ⁻¹ at 298 K]	Reference
H_{O_3}	1.1×10^{-2}	-5.04	Seinfeld and Pandis (2006) Kosak-Channing and Helz (1983)
H_{SO_2}	1.23	-6.25	Seinfeld and Pandis (2006) Pandis and Seinfeld (1989) Smith and Martell (1976)

Table 3. Rate constants for the aqueous phase oxidation of S(IV) with dissolved ozone, used in Eq. (11). The temperature dependence of the rate constants is given by $k(T) = \frac{k_{298}}{\exp\left[-\frac{E}{R}\left(\frac{1}{298} - \frac{1}{T}\right)\right]}$. The activation energies given for the k_1 and k_2 values are those of Erickson et al. (1977), based on measurements at 16 and 25 °C.

	$K_{298} [\text{M}^{-1} \text{s}^{-1}]$	$-E/R [\text{K}]$	Reference
k_0	2.4×10^4	–	Seinfeld and Pandis (2006); Hoffmann (1986)
k_1	3.7×10^5	–5530	Seinfeld and Pandis (2006); Hoffmann (1986)
k_2	1.5×10^9	–5280	Seinfeld and Pandis (2006); Hoffmann (1986)

Table 4. A list of the measured and modelled conditions during each experiment. T_i is the temperature just before the expansion, and T_{\min} is the minimum temperature measured during the expansion. Peak LWC is the maximum condensable water during the expansion, derived from the dew point temperature (T_{dew}) and T_{\min} . The median drop diameter (modelled) is the largest size reached by the median drop during the expansion and the median dry diameter is taken from the model output just prior to the expansion. The maximum activated fraction is the maximum modelled number of droplets larger than $1 \mu\text{m}$ in diameter during the expansion, divided by the total number of aerosol. The change in measured volume divided by the change in modelled volume is the numerical value for the points plotted in Fig. (11). The maximum pH is the maximum pH reached by the median sized droplet during the expansion (modelled), the NH_4/SO_4 molar ratio is calculated from the AMS data just prior to the expansion. The gas phase ammonia mixing ratio was calculated from the AMS data immediately prior to the expansion, as described in Sect. 3 and the number of aerosol is the modelled number of aerosol just prior to the expansion.

Run	Stage number	SO_2 [ppbv]	O_3 [ppbv]	T_i [K]	T_{\min} [K]	T_{dew} [K]	Peak LWC [g m^{-3}]	Median drop Diameter [μm]	Median dry Diameter [nm]	max. activated fraction	Δ Meas. vol/ Δ Mod. vol	max pH	$\text{NH}_4^+/\text{SO}_4^{2-}$ molar ratio	$\text{NH}_3(\text{g})$ pptv	N_{aerosol}
CLD8_17_11_1a	1243.04	17.4	82.3	282.3	277.6	281.0	0.9	9.8	66.1	0.92	0.84	4.37	0.03	0.0142	2100.35
CLD8_17_11_1b	1243.06	18.3	125.8	282.4	278.4	281.2	0.8	13.4	79.6	0.92	0.53	4.28	0.03	0.0123	732.06
CLD8_18_11_1b	1245.03	19.9	126.5	282.5	278.5	280.8	0.5	8.6	66.5	0.73	0.69	4.24	0.02	0.0377	2288.45
CLD8_20_11_1a	1249.03	1.3	136.3	282.3	278.0	281.3	0.9	8.7	76.4	0.87	2.33	5.55	1.58	13.0602	3041.92
CLD8_20_11_1b	1249.09	2.3	137.2	282.3	278.5	281.7	0.9	6.9	88.2	0.95	1.25	5.55	1.67	21.5740	6323.98
CLD8_20_11_2a	1250.11	23.7	136.8	262.7	258.4	261.8	0.3	5.1	71.0	0.95	1.75	4.49	1.87	1.4404	5628.41
CLD8_20_11_2b	1251.04	26.6	128.9	262.3	258.5	261.5	0.3	4.9	91.4	0.90	0.40	4.57	1.88	2.5927	6309.59
CLD8_20_11_2c	1251.09	21.0	126.9	262.4	258.5	261.4	0.3	7.0	113.3	0.90	1.01	4.47	1.85	1.1498	1874.42
CLD8_21_11_1a	1252.04	18.3	115.3	262.3	258.3	261.6	0.3	6.0	81.9	0.96	0.66	4.48	1.87	1.0562	3213.07
CLD8_21_11_1b	1252.12	18.5	115.4	262.3	258.5	261.7	0.3	5.0	82.1	0.98	0.56	4.52	1.80	1.3962	4322.27
CLD8_21_11_1c	1252.17	19.8	118.8	262.5	258.5	261.5	0.3	7.2	101.8	0.96	0.83	4.42	1.69	0.7903	4322.27
CLD8_05_12_1a	1306.20	1.1	92.4	282.7	278.8	281.6	0.6	9.8	53.2	0.34	0.94	5.79	1.75	21.1771	3920.10
CLD8_05_12_1b	1306.27	0.9	131.0	282.8	278.9	281.6	0.6	11.6	71.1	0.59	0.65	5.70	1.74	24.2970	1341.82
CLD8_07_12_1a	1310.14	19.6	107.1	262.6	259.0	261.5	0.2	5.5	71.2	0.92	6.15	4.25	1.24	0.2614	2672.03
CLD8_07_12_1b	1310.17	10.3	178.1	262.6	259.2	261.6	0.2	7.3	109.6	0.96	3.62	4.23	1.21	0.1665	1086.43
CLD8_07_12_1c	1310.23	10.2	203.9	262.5	259.0	261.5	0.2	6.8	88.5	0.90	2.66	4.21	0.85	0.1085	1440.50
CLD9_24_09_2b	1418.11	18.5	112.8	283.4	279.2	282.4	0.8	8.1	66.0	0.86	0.41	4.40	0.61	0.5036	3492.84
CLD9_24_09_2c	1418.15	19.4	115.6	283.4	279.1	282.9	1.0	10.6	76.2	0.90	0.25	4.44	0.84	0.6745	1794.43
CLD9_25_09_1a	1422.05	29.2	90.8	283.7	278.3	282.2	0.9	10.1	98.8	0.98	0.39	4.41	0.92	1.7512	1856.36
CLD9_25_09_1b	1422.08	26.4	85.8	284.0	278.7	282.3	1.0	12.1	112.0	0.97	0.66	4.36	0.89	0.5860	998.07
CLD9_25_09_1c	1422.12	25.4	73.9	284.8	278.7	282.6	1.1	17.0	136.4	1.00	0.63	4.32	0.55	0.2159	410.01
CLD9_25_09_2a	1422.15	25.0	63.5	283.8	278.6	283.1	1.4	8.6	75.7	0.86	0.43	4.35	0.77	0.6497	5028.45
CLD9_25_09_2b	1422.17	25.1	89.2	284.2	277.9	282.9	1.5	10.7	91.6	0.89	0.39	4.35	0.58	0.7761	2756.07
CLD9_25_09_2c	1422.20	25.0	84.9	284.1	277.8	282.9	1.4	12.7	113.0	0.92	0.42	4.42	1.01	1.3842	1476.15
CLD9_28_09_1a	1434.11	17.5	85.3	262.3	257.7	260.8	0.3	5.7	82.2	0.94	2.06	3.91	0.28	0.0098	3385.48
CLD9_28_09_1b	1434.13	20.4	92.8	263.3	257.9	261.7	0.4	6.8	88.1	0.97	1.64	3.97	0.32	0.0093	1970.34
CLD9_28_09_1c	1434.16	22.9	109.4	263.3	258.0	261.9	0.4	8.7	99.1	0.98	1.31	4.02	0.43	0.0124	1149.79
CLD9_28_09_1d	1434.19	21.8	112.2	263.4	258.1	261.9	0.4	10.5	113.6	0.99	1.27	4.04	0.63	0.0174	595.93
CLD9_28_09_2a	1435.04	15.5	107.2	263.5	258.1	261.8	0.3	8.3	49.7	0.92	0.56	4.17	0.10	0.0081	1445.49
CLD9_28_09_2b	1435.06	7.6	109.1	263.1	257.9	261.8	0.4	9.4	73.9	0.96	0.31	4.29	0.92	0.0176	782.26
CLD9_28_09_2c	1435.09	4.0	111.9	263.2	257.7	261.7	0.4	11.9	88.1	0.95	0.40	4.37	0.71	0.0077	373.06

Table 5. Data from the model simulations, showing the gas phase concentration of ammonia, the total particle phase ammonium contained in all particles in one cm^3 of air before cloud formation, and the maximum additional amount of ammonium taken up into all droplets in one cm^3 of air during the cloud formation. Values in brackets indicate the ratio of the number of moles of ammonium contained in the particles or droplets to the gas phase ammonia concentration. The first three columns are reproduced from Table 4, for convenience. Note that the aqueous phase ammonium is in addition to the particle phase ammonium, which is also contained in any droplets that form.

Run	Stage number	$\text{NH}_4^+/\text{SO}_4^{2-}$ [molar ratio]	NH_3 (g) [pptv]	NH_3 (g) [mol cm^{-3}]	NH_4^+ (particle) [mol cm^{-3}], (ratio)	NH_4^+ (aq, peak) [mol cm^{-3}], (ratio)
CLD8_17_11_1a	1243.04	0.03	0.0142	7.05×10^{-19}	5.86×10^{-16} (830.32)	3.86×10^{-16} (547.13)
CLD8_17_11_1b	1243.06	0.03	0.0123	6.11×10^{-19}	2.62×10^{-16} (428.41)	2.63×10^{-16} (430.42)
CLD8_18_11_1b	1245.03	0.02	0.0377	1.88×10^{-18}	9.32×10^{-16} (496.51)	9.83×10^{-16} (523.54)
CLD8_20_11_1a	1249.03	1.58	13.0602	6.50×10^{-16}	4.14×10^{-14} (63.70)	1.65×10^{-14} (25.37)
CLD8_20_11_1b	1249.09	1.67	21.5740	1.07×10^{-15}	1.41×10^{-13} (131.59)	3.26×10^{-14} (30.38)
CLD8_20_11_2a	1250.11	1.87	1.4404	7.71×10^{-17}	7.23×10^{-14} (937.81)	3.89×10^{-14} (504.72)
CLD8_20_11_2b	1251.04	1.88	2.5927	1.39×10^{-16}	1.87×10^{-13} (1342.01)	6.33×10^{-14} (455.78)
CLD8_20_11_2c	1251.09	1.85	1.1498	6.16×10^{-17}	7.20×10^{-14} (1167.93)	3.18×10^{-14} (515.48)
CLD8_21_11_1a	1252.04	1.87	1.0562	5.66×10^{-17}	7.22×10^{-14} (1274.75)	3.20×10^{-14} (564.51)
CLD8_21_11_1b	1252.12	1.80	1.3962	7.48×10^{-17}	7.55×10^{-14} (1008.44)	2.93×10^{-14} (391.31)
CLD8_21_11_1c	1252.17	1.69	0.7903	4.23×10^{-17}	4.04×10^{-14} (953.98)	2.43×10^{-14} (574.69)
CLD8_05_12_1a	1306.20	1.75	21.1771	1.07×10^{-15}	4.85×10^{-14} (45.31)	1.33×10^{-14} (12.38)
CLD8_05_12_1b	1306.27	1.74	24.2970	1.23×10^{-15}	5.58×10^{-14} (45.46)	1.43×10^{-14} (11.67)
CLD8_07_12_1a	1310.14	1.24	0.2614	1.42×10^{-17}	4.03×10^{-14} (2829.00)	1.19×10^{-14} (839.10)
CLD8_07_12_1b	1310.17	1.21	0.1665	9.07×10^{-18}	2.88×10^{-14} (3172.37)	6.87×10^{-15} (757.97)
CLD8_07_12_1c	1310.23	0.85	0.1085	5.91×10^{-18}	2.64×10^{-14} (4464.63)	6.56×10^{-15} (1109.05)
CLD9_24_09_2b	1418.11	0.61	0.5036	2.55×10^{-17}	1.15×10^{-14} (450.25)	1.76×10^{-15} (68.79)
CLD9_24_09_2c	1418.15	0.84	0.6745	3.42×10^{-17}	8.48×10^{-15} (247.82)	9.17×10^{-15} (268.19)
CLD9_25_09_1a	1422.05	0.92	1.7512	8.87×10^{-17}	2.44×10^{-14} (275.10)	2.40×10^{-14} (270.27)
CLD9_25_09_1b	1422.08	0.89	0.5860	2.97×10^{-17}	1.20×10^{-14} (403.80)	1.31×10^{-15} (44.29)
CLD9_25_09_1c	1422.12	0.55	0.2159	1.09×10^{-17}	5.05×10^{-15} (463.43)	6.33×10^{-16} (58.07)
CLD9_25_09_2a	1422.15	0.77	0.6497	3.29×10^{-17}	2.66×10^{-14} (808.86)	3.81×10^{-15} (115.77)
CLD9_25_09_2b	1422.17	0.58	0.7761	3.92×10^{-17}	2.00×10^{-14} (509.26)	1.73×10^{-14} (442.08)
CLD9_25_09_2c	1422.20	1.01	1.3842	7.00×10^{-17}	2.32×10^{-14} (331.90)	1.47×10^{-14} (210.51)
CLD9_28_09_1a	1434.11	0.28	0.0098	5.35×10^{-19}	9.05×10^{-15} (16911.79)	1.37×10^{-15} (2555.99)
CLD9_28_09_1b	1434.13	0.32	0.0093	5.08×10^{-19}	5.19×10^{-15} (10223.71)	1.05×10^{-15} (2064.01)
CLD9_28_09_1c	1434.16	0.43	0.0124	6.76×10^{-19}	6.35×10^{-15} (9391.70)	2.66×10^{-16} (393.23)
CLD9_28_09_1d	1434.19	0.63	0.0174	9.49×10^{-19}	4.92×10^{-15} (5184.55)	1.44×10^{-15} (1519.16)
CLD9_28_09_2a	1435.04	0.10	0.0081	4.40×10^{-19}	1.17×10^{-15} (2664.48)	7.30×10^{-16} (1657.98)
CLD9_28_09_2b	1435.06	0.92	0.0176	9.60×10^{-19}	4.21×10^{-15} (4390.07)	1.00×10^{-15} (1045.57)
CLD9_28_09_2c	1435.09	0.71	0.0077	4.19×10^{-19}	2.02×10^{-15} (4814.28)	1.40×10^{-16} (333.88)




Figure 1. (a) shows the evolution of temperature measured by the 6 TC before, during and after an expansion. Pressure above atmospheric pressure is also shown (dashed line, right hand axis). (b) shows the deviation of the temperature measured by each individual TC from the mean of all TC.




Figure 2. A comparison of the SIMONE-derived and measured (MBW total water minus TDL gas phase water) peak condensed water during CLOUD9 expansions.




Figure 3. AMS measurements of SO_4 (red line) concentrations and $\text{NH}_4^+/\text{SO}_4$ molar ratios (blue markers for raw data and black line for smoothed data). Also shown are the ratios between the main ammonium fragments, to inspect a possible interference from O^+ and HO^+ on the determination of NH_4 concentrations.




Figure 4. The aerosol size distribution measured by the SMPS attached to the total sampling line, for experiments performed on the 17 November 2013. The white line of points shows the mode diameter. Aerosol growth is clearly observed during the cloud periods (marked by the purple vertical bars).




Figure 5. Measured parameters during the second expansion performed on 17 November 2013. (a) shows the time series of pressure and temperature, with the expansion occurring at approximately 600 s. The resulting temperature decrease below the dew point and increase in liquid water content are shown in (b). Vertical lines in (a) and (b) indicate the time at which the SIMONE first detected enhanced forward scattering, indicating the presence of droplets. The temperature measured at this time was taken as the dew point and thus the total water content could be calculated. (c) and (d) show gas mixing ratios and the total density of particles, while (e) shows the diameter of the 25th and 75th percentile of particles, as well as the median diameter (dashed, dot-dashed and solid lines respectively).




Figure 6. The number of cloud droplets formed in the model (orange stars) and the number of aerosol immediately before the expansion (black diamonds), plotted against the maximum number of droplets detected by WELAS during the presence of the cloud. The error bars on the modelled droplet numbers correspond to simulations using the high and low values of the dew point calculated from the SIMONE data. Data are shown for all CLOUD9 experiments where data were available.




Figure 7. The equilibrium gas phase ammonia mixing ratio, calculated with E-AIM, as a function of the ammonium to sulphate ratio of the aerosol. Data shown are for -10 and 10°C , at $\text{RH} = 95\%$.




Figure 8. The aqueous phase concentration of NH_4^+ in a 7 μm diameter water droplet, exposed to a 22 pptv NH_3 gas phase, similar to the conditions of experiment CLD8_20_11_1b.




Figure 9. The change in glyoxal mixing ratio during an expansion performed in the CLOUD chamber as part of a series of isoprene oxidation experiments. Pressure and temperature are also shown, for reference.




Figure 10. (a) The measured growth in the total aerosol volume as a function of $\text{NH}_4^+/\text{SO}_4^{2-}$ ratio in the seed aerosol, at 10 and -10°C . The growth is determined as the difference in aerosol volume before and after the cloud. The error bars on the measured growth stem from the uncertainty in diameter in the SMPS measurements (which translates to approximately 14 % in volume for particles with a diameter greater than 50 nm and 22 % for those with a diameter less than 50 nm). **(b)** As in **(a)**, except here the modelled data is presented. The error bars stem from simulations carried out using the high and low limits of the dew point, as determined from the SIMONE data.




Figure 11. (a) The modelled vs. observed total increase in aerosol volume due to the aqueous phase oxidation of SO₂ to SO₄²⁻ at 10 and -10 °C. Note that the masses are corrected to account for any particle loss during the expansion (see text). The dashed lines indicate the range of a factor of two higher or lower than the observed values. The error bars are as for Fig. 10. **(b)** As for **(a)**, but with doubled gas phase NH₃. **(c)** As for **(a)**, but with no gas phase NH₃.




Figure 12. As for Fig. 11a, except that the colours indicate the ion conditions under which the experiments were performed: ion free (neutral), natural ion concentrations (GCR) and enhanced ion concentrations (CIGAR).