

Author Comments

1) Comment 1: I would think this could be answered by looking some cases and perhaps design some experiments to test where the inconsistency comes from.

(1) Retrieval failures in the tropics

We further looked into the statistic of parameters that are obtained during the inversion process. The coefficient of correlation between EOF 1 and EOF 2 scores is centered at -0.8 in the tropics and strongly skewed, whereas it is centered at zero and nearly symmetric in the extratropics. This is an indication that the tropics measurements do not provide the directional distribution of polarized radiance that allows independent retrieval of EOF 1 and EOF 2 scores. Also, the retrieved cloud top height in the tropics contains significant error, compared to the collocated CALIOP dataset.

These are presumably because the observations are aligned near the backscattering direction with low zenith angles in the tropics. The measurements in limited scattering angle provide insufficient directionality information to the inversion system, and low zenith angles lead to low airmass, that makes the spectral contrast of polarization signal weak.

(2) Inconsistencies in the extratropics

A sensitivity study is conducted with different particle shapes, and the result shows that the selection of particle shape has a significant impact on the alignment between forward model and the observation density. Also, the CALIOP collocated dataset indicates that the removal of multilayer clouds and aerosol above clouds do not have a major impact on the retrieved distribution of EOF scores. Other possibility to look into is to design an experiment that compares the effect of viewing zenith angles, but the time was limited to address this issue in this revision.

2) Comment 2: I think that reviewer 2's statement is fair and I don't quite understand your explanations. Could you please elaborate on this?

We revised a part explaining our EOF technique to make it clear that EOF analysis of $-P_{12}$ theoretically allows mixing of particles, and the interpolation in the polarized reflectivity based on them is accurate enough for our purpose.

3) Comment 4: I am bothered by the extrapolations of the linear regression relationship between EOF1 score and roughness, because when a regression is performed, the relationship should be valid only in the range used in the regression process. As pointed out by reviewer 2, we don't really know if this relationship remains the same outside the range. However, I agree with you that it seems less useful if any roughness outside the range is unphysical anyway. If the wrong choice of ice particle model could be the main reason for the unphysical EOF1 score/roughness, could you please at least show one

example, demonstrating how sensitive EOF1 score is to the choice of the ice particle model? Since it may be inevitable to apply an inappropriate ice particle model for actual observations, what would be appropriate criteria to remove unphysical retrieval? In short, a couple of paragraphs that detail when/what factors will lead to unphysical EOF1 score/roughness and how sensitive retrievals are to those factors will be helpful for readers to apply the proposed method and to understand what those retrievals represent.

We added a new figure to elaborate on the sensitivity of the retrieval to assumed particle shapes. As the roughness retrieval is quite sensitive to the assumed particle shapes, the shape must be known in advance to eliminate the unexpectedly large roughness values if the roughness is retrieved independently. However, there is a kernel space in the shape and roughness retrieval (e.g. van Diedenhoven et al. 2012), so it may be possible to constrain the combination of shape and roughness. Our EOF analysis is proposed in a hope of making this process more theoretically robust. We cannot fully address this question even at best of our knowledge, but our hope is to do so in the near future.

4) Comment 5–6: More detailed analysis and supporting evidence are needed.

New evidences from correlation histogram and the cloud top height comparison indicates that the difference between the tropics and extratropics are probably due to the viewing geometry, as the reviewer 2 speculated.

List of relevant changes

- Section 2.2.1 is revised to respond to the comment (2).
- A new section is inserted before the 3.2: Inference Failure in Tropics to expand upon the discrepancy in the extratropical retrievals to respond other comments.
- Two new paragraphs are added to Section 3.2 (new number 3.3)
- Three new figures are included, and one old figure is deleted.
- Minor corrections to equations and notations

1 For submission to Atmospheric Chemistry and Physics

2

3 Degree of ice particle surface roughness in- 4 ferred from polarimetric observations

5 S. Hioki¹, P. Yang¹, B. A. Baum², S. Platnick³, K. G. Meyer⁴, M. D. King⁵, J.
6 Riedi⁶

7 [1]{Department of Atmospheric Sciences, Texas A&M University, College Station,
8 Texas}

9 [2]{Space Science and Engineering Center, University of Wisconsin-Madison, Madi-
10 son, Wisconsin}

11 [3]{Earth Science Division, NASA Goddard Space Flight Center, Greenbelt, Maryland}

12 [4]{Goddard Earth Sciences Technology and Research, Universities Space Research
13 Association, Columbia, Maryland}

14 [5]{Laboratory for Atmospheric & Space Physics, University of Colorado, Boulder,
15 CO}

16 [6]{Laboratoire d'Optique Atmosphérique, Université de Lille - Sciences et Technol-
17 ogies, Villeneuve d'Ascq, France}

18

19

20 Correspondence to S. Hioki (s.hioki@tamu.edu)

21 **Abstract**

22 The degree of surface roughness of ice particles within thick, cold ice clouds is in-
23 ferred from multi-directional, multi-spectral satellite polarimetric observations over
24 oceans, assuming a column-aggregate particle habit. An improved roughness infer-
25 ence scheme is employed that provides a more noise-resilient roughness estimate
26 than the conventional best-fit approach. The improvements include the introduction

1 of a quantitative roughness parameter based on empirical orthogonal function anal-
2 ysis and proper treatment of polarization due to atmospheric scattering above
3 clouds. A global one-month data sample supports the use of a severely roughened
4 ice habit to simulate the polarized reflectivity associated with ice clouds over ocean.
5 The density distribution of the roughness parameter inferred from the global one-
6 month data sample and further analyses of a few case studies demonstrate the sig-
7 nificant variability of ice cloud single-scattering properties. However, the present
8 theoretical results do not agree with observations in the tropics. In the extratropics,
9 the roughness parameter is inferred but 74% of the sample is out of the expected
10 parameter range. Potential improvements are discussed to enhance the depiction of
11 the natural variability on a global scale.

12

13 **1. Introduction**

14 Satellite observations at visible and infrared wavelengths can characterize
15 global cloud microphysical parameters and radiative properties. Numerous
16 techniques have been developed to retrieve ice cloud optical and microphysical
17 properties from radiometric measurements (e.g., Inoue, 1987; Nakajima and King,
18 1990; Minnis et al., 1993) and have been adopted in operational retrieval efforts
19 (Rolland et al., 2000; Platnick et al., 2003; Minnis et al., 2011). A synergetic
20 combination of satellite and in-situ observations (e.g., Heymsfield et al., 2002, 2013)
21 serves as a constraint for the parameterization of bulk ice cloud optical properties
22 for remote sensing implementations as well as general circulation models (GCMs).

23 The accuracy of these retrieval techniques and the validity of downstream
24 applications including GCM radiation parameterization hinges on steady
25 improvements in the single-scattering calculations involving ice crystals. As ice
26 clouds consist of nonspherical particles with characteristic sizes much larger than

1 the wavelengths of interest, the single-scattering properties depend on the size,
2 shape, and microscopic morphology of the particles (Macke et al., 1996; Yang et al.,
3 2008a; Xie et al., 2009; Baum et al., 2010; Um and McFarquhar, 2007, 2009;
4 Ulanowski et al., 2006, 2014). In the solar shortwave spectrum, particle shape,
5 surface texture, and crystal imperfections have a substantial influence on the single-
6 scattering properties. Recent improvements in scattering calculation techniques are
7 being incorporated into models that represent diverse ice particle populations in
8 clouds.

9 However, it is challenging to quantify some of these influential microphysical
10 parameters, given current satellite sensors. As a result, little information about their
11 variability is available. The discrepancies among climate models (Waliser et al.,
12 2009) in terms of ice water path (IWP) indicate that the GCM parameterizations
13 need more reliable constraints on IWP. Recent work by Sourdeval et al. (2015),
14 which includes direct retrieval of IWP, is a novel approach to this problem.

15 The ability of passive and lidar sensors to correctly infer IWP requires
16 knowledge of ice cloud radiative properties. Application of an unrealistic ice model,
17 e.g., with only smooth (unroughened) surfaces, results in an overall global bias
18 (Macke and Mishchenko 1996; Yang et al., 2007, 2008b; Holz et al. 2015) as well as
19 seasonal biases (Zhang et al., 2009) in cloud property retrievals. The overarching
20 goal of this paper is to gain a better understanding of the constraints in the
21 microphysical parameters of global ice clouds using angular polarimetric
22 observations and state-of-the-art light-scattering computational capabilities.

23 Multidirectional polarimetric observations can constrain the representative
24 particle shape and surface texture condition (specifically, the degree of surface
25 roughness), owing to the sensitivity of the polarization state of reflected light to
26 small-scale particle structures. These measurements have been used to infer both
27 particle habit (Chepfer, 1998; C.-Labonnote et al., 2001; Masuda et al., 2002; Knap et

Souichiro Hioki 5/12/16 7:50 PM
Deleted: overarching

1 al., 2005; Baran and C.-Labonnote, 2007) and surface roughness (Baran and C.-
2 Labonnote, 2006; Cole et al., 2013, 2014). Since polarized reflectivity saturates at
3 relatively small optical thickness (generally about $\tau = 5$, Masuda and Takashima,
4 1992), the conventional “best-fit” approach to this problem computes the residual
5 sum of squares (RSS) from the multi-angle observations of polarized reflectivity and
6 reflectivity simulations, and selects the ice particle model that minimizes the RSS
7 when $\tau > 5$.

8 The previous studies imply that the use of roughened particles is necessary to
9 achieve maximum consistency between observations and numerical scattering
10 calculations. Furthermore, the spectral consistency of visible/near-infrared and
11 thermal infrared retrievals (Baran and Francis, 2004) was recently investigated by
12 Liu et al. (2014) and Holz et al. (2015), who report that retrieved ice cloud optical
13 thicknesses are more consistent when particles are roughened.

14 The treatment of particle surface roughness here is not a rigorous approach.
15 Rather, it is an approximation of the effects of roughened surface texture (Neshyba
16 et al., 2013) and other kinds of imperfections present in natural ice cloud particles.
17 The scattering properties calculated by this approximate method are in reasonable
18 agreement with those calculated by rigorous ray-tracing methods (Yang et al.,
19 2008a). Although previous studies suggest that some degree of roughness is
20 desirable, the issue remains as to the amount of roughness that should be adopted
21 for global satellite-based retrievals, or used in numerical models.

22 Recent work by van Diedenhoven et al. (2012, 2014) simultaneously infers
23 both the aspect ratio and the degree of roughness from a combination of
24 polarimetric and intensity observations over a virtually continuous parameter
25 space, assuming that simple hexagonal ice particles can explain observations. The
26 ability to infer a representative ice cloud particle aspect ratio adds yet another
27 dimension to the problem. Such exploration into the variability of ice particle

Souichiro Hioki 5/12/16 7:50 PM

Deleted: 2015

Souichiro Hioki 5/12/16 7:50 PM

Deleted: a particle is

Souichiro Hioki 5/12/16 7:50 PM

Deleted: property

Souichiro Hioki 5/12/16 7:50 PM

Deleted: is

Souichiro Hioki 5/12/16 7:50 PM

Deleted: that

1 microphysical properties can lead to a more reliable satellite climatology of ice
2 clouds. This study focuses on the quantitative inference of ice particle roughness
3 parameter for a specific particle habit, and will not include a detailed investigation
4 of aspect ratio.

5 While a conventional “best-fit” approach can constrain the range of the
6 average roughness parameter at the global scale, it is not suitable for pixel-by-pixel
7 inferences. This is because the signal-to-noise ratio for particle roughness is low,
8 and in the conventional “best-fit” approach, even random observational errors can
9 modify the inferred histogram significantly when it is applied to individual pixels.
10 Figure 1 illustrates how such a modification takes place if the method is applied to a
11 synthetic signal with random noise. To produce Fig. 1, viewing geometries are
12 extracted from one month (September 2005) of cloud observations by the
13 POLarization and Directionality of the Earth’s Reflectance (POLDER) sensor
14 (Deschamps et al., 1994) onboard the Polarization and Anisotropy of Reflectances
15 for Atmospheric Sciences coupled with Observation from Lidar (PARASOL) satellite
16 (Fougnie et al., 2007). The “best-fit” inference is applied to synthetic multi-angle
17 cloud polarized reflectivities (L_p , defined in Section 2.1) with and without random
18 noise. In synthesizing the signal, a column aggregate particle shape (e.g. Yang et al.,
19 2013) is assumed with a roughness parameter of $\sigma^2 = 0.15$ (variance of the slope of
20 random facet tilts, see Yang et al., 2008b for details), and the random error has a
21 normal distribution with variance equivalent to the POLDER observational error,
22 which is estimated in Section 2.1. The hatched bar is the histogram with noise and
23 the gray bar is that without noise. Note that the distinct peak at $\sigma^2 = 0.15$ is no
24 longer apparent when instrumental noise is included, indicating the necessity of
25 appropriate treatment of the error distribution in the analysis.

26 This paper demonstrates how a continuous parameter space for the roughness
27 retrieval is constructed and how it can be used to infer the particle roughness of

Souichiro Hioki 5/12/16 7:50 PM

Deleted: the

1 optically thick ice clouds on a pixel-by-pixel basis. Section 2 provides the details of
2 the data and inversion method we employ, and the result of the application to one
3 month of global data is described in Section 3. Concluding remarks are given in
4 Section 4.

Souichiro Hioki 5/12/16 7:50 PM

Deleted: employed

6 **2 Methodologies**

7 To establish a method resilient to observational error, we first examine
8 random errors in POLDER data and select pixels based on the MODIS Collection 6
9 cloud product, as given in Section 2.1. Then, a continuous parameter space for
10 inferring roughness is constructed by using an empirical orthogonal function
11 analysis, and used in the retrieval scheme with the maximum likelihood method.
12 The construction of the parameter space, and the design and the performance of the
13 forward model, are discussed in Section 2.2.

Souichiro Hioki 5/12/16 7:50 PM

Deleted: scrutinize

15 **2.1 Observations**

16 **2.1.1 Reflectivity from POLDER**

17 The POLDER sensor onboard the PARASOL satellite provides multispectral
18 polarimetric observations at up to 16 viewing geometries for a single overpass
19 (Fougnie et al., 2007). The PARASOL satellite was in the A-train satellite
20 constellation from 2004 to 2009 and continued operation in a separate orbit until
21 late 2013, providing a total of nine years of global polarimetric observation data.
22 The design of the instrument is inherited from previous POLDER sensors on the
23 ADEOS (ADvanced Earth Observing Satellite) platforms. POLDER sensors provide
24 the first three elements of the Stokes vector from three images taken successively
25 with linear polarization filters (Deschamps et al., 1994).

1 This study uses the single-pixel data set in [the](#) PARASOL Level 1B product. [The](#)
2 approximate resolution is $6 \text{ km} \times 6 \text{ km}$.

3 PARASOL products report the intensity of reflection in terms of normalized
4 radiance L_n , which is equal to the reflectivity R of the surface-atmosphere system
5 multiplied by the factor $\mu_0 = \cos \theta_0$ (cosine of solar zenith angle).

$$L_n(\mu, \mu_0, \varphi, \varphi_0) = \mu_0 R(\mu, \mu_0, \varphi, \varphi_0) \quad (1)$$

6 The reflectivity R is defined as

$$R(\mu, \mu_0, \varphi, \varphi_0) = \frac{\pi I(\mu, \mu_0, \varphi, \varphi_0)}{E_0 \mu_0} \quad (2)$$

7 where $I(\mu, \mu_0, \varphi, \varphi_0)$ is the radiance and $E_0 \mu_0$ is the irradiance of incoming unpolar-
8 ized light (i.e., solar irradiance; E_0 is the beam flux).

9 In a similar manner, the polarized reflectivity is reported in terms of
10 normalized radiance, so (L_n, Q, U) become the first three Stokes parameters. In
11 other words, the normalized polarized radiance $L_{np} = \sqrt{Q^2 + U^2}$ is equal to the
12 polarized reflectivity R_p multiplied by μ_0 .

$$L_{np}(\mu, \mu_0, \varphi, \varphi_0) = \mu_0 R_p(\mu, \mu_0, \varphi, \varphi_0) = \frac{\pi \sqrt{Q_i^2 + U_i^2}}{E_0}, \quad (3)$$

13 where Q_i and U_i are defined to form the first three Stokes parameters in terms of
14 radiance (I, Q_i, U_i) . It is worth noting the similarity between Eqs. (1) and (3). We
15 conduct the analysis in terms of $L_{np} = \mu_0 R_p$ defined in Eq. (3) to simplify the error
16 estimate.

17 The distribution of random errors in L_{np} observed with POLDER is estimated
18 in the following procedure. A reflection property of an optically thick ice cloud is
19 that the modified polarized reflectivity $L_{nmp} = \eta(\mu + \mu_0) L_{np} / \mu_0$ (where $\eta = \pm 1$, C.-
20 Labonnote et al., 2001) crosses zero at scattering angle $\Theta \approx 170^\circ$ as shown in Fig. 2.

Souichiro Hioki 5/12/16 7:50 PM

Deleted: (i.e. the

Souichiro Hioki 5/12/16 7:50 PM

Deleted:).

Souichiro Hioki 5/12/16 7:50 PM

$$\frac{\pi \sqrt{Q_i^2 + U_i^2}}{E_0 \mu_0}$$

Souichiro Hioki 5/12/16 7:50 PM

Deleted: L_p

1 This implies that the polarization signal at $\theta \approx 170^\circ$ is primarily due to the
2 observational noise with additional contributions from the variation of cloud
3 particle scattering properties. We utilize this reflection property to estimate the
4 magnitude of observational noise from the POLDER data at scattering angles
5 between 168° and 172° , and further estimate the noise level at other angles with a
6 typical polarization state of cloud reflection.

7 The POLDER observational noise consists of radiometric noise and
8 misregistration noise. The misregistration noise is inherent in the POLDER sensor's
9 design that extracts polarimetric information from three images successively taken
10 with different polarizers. The co-registration process of these three images is an
11 inevitable source of error. As the distribution of misregistration noise is unknown,
12 our instrument model attempts to explain both noise components with a
13 radiometric noise model in the following analysis.

14 We define a random variable L_{np} that serves as a statistical model of observed
15 L_{np} as follows.

$$L_{np} = \sqrt{X_1^2 + X_2^2 + X_3^2 - X_1X_2 - X_2X_3 - X_3X_1} \quad (4)$$

16 where random variables X_1 , X_2 , and X_3 represent the radiances of a pixel in the orig-
17 inal three images with different polarizers (not available in a product). With the sta-
18 tistical model outlined in Eq. (4), we first assume that X_1 , X_2 , and X_3 follow the same
19 normal distribution centered at 0.5 with variance s^2 (i.e., $X_i \sim N(0.5, s^2)$) because the
20 expectation of polarized radiance L_{np} is assumed to be zero at scattering angles be-
21 tween 168° and 172° . With this assumption, we apply the parametric bootstrap
22 method (e.g., Evans and Rosenthal, 2010) to obtain the distribution of L_{np} as a func-
23 tion of variance s^2 . The observational distribution of L_{np} at $0.865 \mu\text{m}$ in the scatter-
24 ing angles between 168° and 172° (within the rectangular box in Fig. 2) is shown in

Souichiro Hioki 5/12/16 7:50 PM

Deleted: misregistra-tion

Souichiro Hioki 5/12/16 7:50 PM

Deleted: channel

1 the bar chart of Fig. 3, and compared with the theoretical distribution with
2 $s = 0.00095$ (solid line). Figure 4 justifies our selection of $s = 0.00095$ by showing
3 that the sum of squared errors of the density in each bin of the histogram (Fig. 3) is
4 minimized when $s = 0.00095$. Therefore, we take $s = 0.00095$ as the standard er-
5 ror for X_1, X_2 , and X_3 . In Fig. 3, the distribution from observations is slightly more
6 skewed than the distribution from bootstrapping, but their agreement justifies the
7 use of the simple statistical model formulated in Eq. (4) to quantify the magnitude of
8 measurement errors.

9 To obtain the approximate magnitude of the L_{np} error at other scattering
10 angles, the same parametric bootstrap method is applied with the degree of linear
11 polarization fixed at 5%, which is the upper limit for typical ice cloud reflection. This
12 selection does not significantly affect the following analysis. When the signal is
13 polarized, random variables X_1, X_2 , and X_3 do not follow the same distribution, but
14 we assume that the standard errors for X_1, X_2 , and X_3 still stay the same. Figure 5
15 shows the estimated magnitude of error (variance) as a function of normalized
16 radiance L_n . The variance of L_{np} asymptotes to a near-constant value once L_n
17 reaches $L_n = 0.2$. As shown in insets, the distribution becomes closer to a normal
18 distribution with increasing L_n . Based on the discussion above, we conclude that the
19 error distribution of L_{np} approximately follows a normal distribution with variance
20 $\text{var}(L_{np}) = 1.35 \times 10^{-6}$ for a reflective target ($L_n \geq 0.2$). This estimate of error is
21 about the same magnitude as the value by Fougnie et al. (2007). Note that we
22 assume that the error is purely from observational noise, neglecting any natural
23 cloud variability. Therefore, the actual radiometric noise level should be somewhat
24 smaller than our estimate. We estimate the magnitude of error using the 0.865 μm
25 channel because the channel is likely to be the least contaminated by other sources
26 of uncertainty such as ozone absorption (0.67 μm) and Rayleigh scattering (0.49 μm ,

Souichiro Hioki 5/12/16 7:50 PM
Deleted: 856

1 0.67 μm). We apply the same variance to all three POLDER channels used in the
2 analysis (0.865, 0.67, and 0.49 μm).

3 **2.1.2 Ancillary Data from MODIS and AIRS**

4 The MODerate resolution Imaging Spectroradiometer (MODIS) instruments
5 onboard the Aqua and Terra satellites measure radiance at multiple visible and
6 infrared wavelengths, providing various products (King et al., 2003) that are
7 complementary to those from PARASOL. Of interest here is the Collection 6 Level 2
8 cloud product (MYD06) from Aqua MODIS, with which the PARASOL satellite was
9 flying in formation until 2009. Cloud top temperature and thermodynamic phase are
10 extracted from MYD06 and are collocated to POLDER data to be used in the analysis
11 described later in this section. In addition, Level 3 monthly mean ozone
12 concentration from the Atmospheric Infrared Sounder (AIRS) on the Aqua satellite
13 is also used, in particular to account for absorption by ozone that attenuates
14 reflected radiation in the visible range.

15 **2.1.3 Collocation and Selection**

16 The PARASOL Level 1 radiometric data is first collocated with the MODIS Level
17 2 cloud product (Platnick et al., 2015) to select pixels containing ice clouds. Only
18 PARASOL pixels that have corresponding MODIS observations are selected, and
19 filtered by the criteria summarized in Table 1. The intent of the filtering process is to
20 avoid cloud edge contamination, to avoid supercooled water droplets, and to select
21 pixels where clouds are optically thick. The selection criterion of 208 K is a
22 threshold used to identify convective precipitation in the tropics (Mapes and Houze,
23 1993). The analysis is applied only over oceans so the influence of surface reflection
24 is minimal.

25 A “pixel” in the PARASOL Level 1 product contains reflectivity data observed
26 from up to 16 viewing angles. An individual reflectivity value stored in a pixel is

Souichiro Hioki 5/12/16 7:50 PM

Deleted: multiple

Souichiro Hioki 5/12/16 7:50 PM

Deleted: geometries. Individual

Souichiro Hioki 5/12/16 7:50 PM

Deleted: data

1 called a “view”, and we select valid views using criteria relating to scattering angle
2 and sunglint angle (see Table 1). When five or more valid views are contained in a
3 pixel that satisfy all pixel criteria previously mentioned, the pixel is marked as valid,
4 and the roughness inference is attempted.

5 2.2 Inversion Methods

6 2.2.1 Selection of Retrieval Parameters

7 To overcome the problem of the conventional “best-fit” approach that uses a
8 discrete set of roughness parameters, we construct a continuous parameter space
9 for the particle roughness with empirical orthogonal functions (EOF). The goal of
10 the EOF analysis is to find the parameter space that describes the variation of the
11 $-P_{12}$ element of the phase matrix when varying the particle roughness. An ideal
12 approach would be to use a collection of $-P_{12}$ values from observations (Rodgers,
13 2000), but such a dataset is unavailable. For this reason, we apply EOF analysis to
14 the $-P_{12}$ simulated with light scattering calculations. The first and second EOFs
15 together explain 99.3% of the entire variation of $-P_{12}$ in the scattering angle range
16 from 60° to 160° . This implies that following approximation is valid in the scattering
17 angle range $60^\circ < \theta < 160^\circ$.

$$-P_{12}(\theta) = x_1(\sigma^2)Q_1(\theta) + x_2(\sigma^2)Q_2(\theta) \quad (5)$$

18 where θ is scattering angle, $Q_1(\theta)$ and $Q_2(\theta)$ are the first and second EOFs, and
19 $x_1(\sigma^2)$ and $x_2(\sigma^2)$ are weights for EOFs (EOF Scores). Note that the set of EOFs and
20 EOF scores obtained in this way depends on the selection of particle shapes and the
21 degree of roughness. In our EOF analysis, ten prescribed roughness parameter (σ^2)
22 values are used: 0, 0.03, 0.05, 0.1, 0.15, 0.2, 0.3, 0.4, 0.5, and 0.7. These roughness
23 parameters are selected to outline the variation of $-P_{12}$ over the course of rough-
24 ness changes, including the roughness parameter used in MODIS Collection 6

Souichiro Hioki 5/12/16 7:50 PM

Deleted: °, so $-P_{12}$ for a particle of any roughness can be reconstructed almost exactly from the first and second EOFs by a simple linear combination. Also, $-P_{12}$ for a mixture of particles with different roughness parameters can be reconstructed by the same linear combination.

Souichiro Hioki 5/12/16 7:50 PM

Formatted: No widow/orphan control, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Souichiro Hioki 5/12/16 7:50 PM

Moved (insertion) [1]

1 ($\sigma^2 = 0.5$). The EOF scores are shown in Fig. 6. The EOF 1 primarily describes the
2 degree of roughness, and the EOF 2 score has sensitivity to large roughness param-
3 eters. Therefore, the EOF 1 and EOF 2 scores are selected as retrieval parameters re-
4 lated to particle roughness.
5 Since the phase matrix follows a linear mixing rule, $-P_{12}$ of a mixture containing
6 multiple degrees of roughness is also approximated by Eq. (5). For example, a mix-
7 ture of MODIS Collection 6 particle ($\sigma^2 = 0.5$) and moderately roughened particle
8 ($\sigma^2 = 0.03$) produces EOF scores (x_1, x_2) on a straight line between
9 $(x_1(0.5), x_2(0.5))$ and $(x_1(0.03), x_2(0.03))$. Constructing a continuous parameter
10 space using EOF scores (x_1, x_2) is very powerful because the method guarantees
11 that the parameter space contains any mixture of prescribed shapes or degree of
12 roughness. In Section 2.2.1, we show that these EOF scores can be used to accurately
13 parameterize the normalized polarized reflectance L_{np} for a given direction.

14 The details of the particle model are as follows. The EOF analysis is applied to
15 $-P_{12}$ elements of the phase matrices calculated by the method described by Yang et
16 al. (2013), which is a combination of the Improved Geometric Optics Method (IGOM,
17 Yang and Liou, 1996) and the Amsterdam Discrete Dipole Approximation method
18 (ADDA, Yurkin et al., 2007). Surface roughness is applied only in the IGOM
19 computation ($D_{max} > 10 \mu\text{m}$). The column aggregate shape is chosen because the
20 most extensive previous study on a global scale (Cole et al., 2014) implies that this
21 habit produces the most consistent agreement with observations. In addition, this
22 habit is used in the retrieval scheme for the operational MODIS Collection 6 cloud
23 products. This particle shape is an aggregate of eight column elements that are solid
24 hexagonal particles with slightly different particle aspect ratios (originally defined
25 in Yang and Liou 1996, see Yang et al., 2013 for geometric parameters). A gamma
26 particle size distribution with an effective size (diameter) of $60 \mu\text{m}$ and an effective

Souichiro Hioki 5/12/16 7:50 PM
Moved (insertion) [2]

Souichiro Hioki 5/12/16 7:50 PM
Deleted:), and ten prescribed roughness parameter (σ^2) values are used in calculations: 0, 0.03, 0.05, 0.1, 0.15, 0.2, 0.3, 0.4, 0.5, and 0.7.

Souichiro Hioki 5/12/16 7:50 PM
Moved up [1]: over the course of roughness changes, including the roughness parameter used in MODIS Collection 6 ($\sigma^2 = 0.5$).

Souichiro Hioki 5/12/16 7:50 PM
Deleted: The choice of roughness parameter is somewhat arbitrary, but our intent is to outline the variation of $-P_{12}$

1 variance of 0.1 is used in this study since we expect little impact on our analysis due
2 to this size distribution selection (Cole et al., 2014).

3 As the EOF 1 score is a monotonic function of the roughness parameter and
4 explains most of the $-P_{12}$ variation (85.6%), it can be considered as an effective
5 roughness parameter for this shape. The relation between EOF 1 scores and the
6 natural logarithm of roughness parameters is nearly linear (Fig. 7), indicating that
7 the roughness parameter can be subsequently inferred after the inference of the
8 EOF 1 score. The straight line in Fig. 7 is the regression line defined in the form

$$\sigma^2 = \exp[-115.755x_1 - 2.3543]. \quad (6)$$

9 As the roughness parameter computed from Eq. (6) does not account for the varia-
10 tion of EOF 2 score, it is inaccurate for the mixture of particles containing multiple
11 degrees of roughness. Equation (6) is introduced to compare our retrievals to the
12 conventional discrete parameter space.

13 Another factor that can impact the roughness retrieval is atmospheric
14 Rayleigh scattering above the cloud. Above-cloud Rayleigh scattering has previously
15 been used to infer cloud top pressure from polarimetric measurements (e.g., Buriez
16 et al., 1997), with results comparable to those from O₂ A band retrievals and ISCCP
17 (Parol et al., 1999). With the POLDER instruments, Rayleigh scattering is primarily
18 detected as a spectral and directional difference of polarized reflectivities. Figure 8
19 shows the change of L_{np} at 0.865 μm , as a function of scattering angle, in response to
20 a 300 hPa change in cloud top pressure (i.e., from 200 to 500 hPa, the red line) and a
21 change of similar effect in roughness parameter (from $\sigma^2 = 0.15$ to 0.5, the dashed
22 green line). The effects of cloud top pressure and roughness parameter changes on
23 L_{np} have different directional patterns but comparable magnitudes. The variation of

Souichiro Hioki 5/12/16 7:50 PM

Deleted: The weights for the EOFs (EOF scores) are displayed in Fig. 6, and some original and reconstructed $-P_{12}$ curves are plotted in Fig. 7. Figure 6 shows that

Souichiro Hioki 5/12/16 7:50 PM

Deleted: . Since the first EOF

Souichiro Hioki 5/12/16 7:50 PM

Deleted: 8

Souichiro Hioki 5/12/16 7:50 PM

Deleted: 8

Souichiro Hioki 5/12/16 7:50 PM

Formatted: Font:+Theme Body

Souichiro Hioki 5/12/16 7:50 PM

$[-115.755 \times (\text{EOF 1 score}) - 2.3543]$.

Souichiro Hioki 5/12/16 7:50 PM

Deleted: 5

Souichiro Hioki 5/12/16 7:50 PM

Deleted: In addition, the EOF 2 score provides constraints on the retrieval especially for large roughness parameters, while the EOF 1 score is less sensitive to the roughness variation.

Souichiro Hioki 5/12/16 7:50 PM

Moved up [2]: Therefore, the EOF 1 and EOF 2 scores are selected as retrieval parameters related to particle roughness. .

Souichiro Hioki 5/12/16 7:50 PM

Deleted: 9

Souichiro Hioki 5/12/16 7:50 PM

Deleted:

1 the cloud top height must therefore be well constrained or retrieved simultaneously
2 when attempting to infer the roughness parameter.

3

4 2.2.2 Construction of Forward Model

5 Once the inverse problem is formulated, the next step is to construct a forward
6 model that is fast enough to be embedded in the inversion algorithm. From the
7 discussion in Section 2.1 and 2.2.1, the inverse problem is formalized as follows: (1)
8 the parameters to be inverted are the EOF 1 and EOF 2 scores, and cloud top
9 pressure; and (2) observations are MODIS-AIRS-collocated L_{np} from POLDER at
10 central channel wavelengths 0.865 μm , 0.67 μm , and 0.49 μm . To satisfy the
11 requirements for numerical efficiency, the present forward model is based on a look
12 up table. The adding-doubling radiative transfer program is used to compute L_{np} for
13 every phase matrix with seven atmospheric scattering optical thicknesses above the
14 cloud: 0, 0.02, 0.05, 0.1, 0.15, 0.2, and 0.3. The result at a specific viewing geometry
15 (denoted by subscript i) and a optical thickness (denoted by subscript j) is
16 parameterized by a simple linear regression model defined as:

$$L_{np,(i,j)} = a_{(i,j)} + b_{(i,j)}x_1 + c_{(i,j)}x_2 \quad (7)$$

17 where x_1 and x_2 are EOF scores obtained in Sect. 2.2.1, and $a_{(i,j)}$, $b_{(i,j)}$, and $c_{(i,j)}$ are
18 regression coefficients determined by the polarized reflectivities for multiple phase
19 matrices. The viewing geometry is gridded as follows: solar zenith angles from 0° to
20 81° , viewing zenith angles from 0° to 75° , and relative azimuth angles from 0° to
21 180° , with an interval of 3° for each. The regression is repeated for seven atmos-
22 pheric scattering optical thicknesses above the cloud and more than 40,000 viewing
23 geometries. With this fast forward model, once cloud top height and EOF scores are
24 given, L_{np} can be obtained for each specific viewing geometry and wavelength. Note

- Souichiro Hioki 5/12/16 7:50 PM
Deleted: function
- Souichiro Hioki 5/12/16 7:50 PM
Deleted: thickness
- Souichiro Hioki 5/12/16 7:50 PM
$$L_{np,i} = a_i + b_i x_1 + c_i x_2$$
- Souichiro Hioki 5/12/16 7:50 PM
Deleted: 6
- Souichiro Hioki 5/12/16 7:50 PM
Deleted: a_i ,
- Souichiro Hioki 5/12/16 7:50 PM
Deleted: b_i ,
- Souichiro Hioki 5/12/16 7:50 PM
Deleted: c_i
- Souichiro Hioki 5/12/16 7:50 PM
Deleted: for each viewing geometry, which is denoted
- Souichiro Hioki 5/12/16 7:50 PM
Deleted: subscript i . The linear relation between L_{np} and $-P_{12}$ in
- Souichiro Hioki 5/12/16 7:50 PM
Deleted: single scattering approximation underscores the appropriateness of this formulation
- Souichiro Hioki 5/12/16 7:50 PM
Deleted: stored in a file so that one can apply appropriate interpolations. In this way

1 that when a set of EOF scores (x_1, x_2) is not exactly at the values corresponding to
2 the ten prescribed phase matrices, the forward model linearly interpolates the po-
3 larized reflectivity. We confirmed that the interpolation usually produces a reliable
4 polarized reflectivity simulation for a phase matrix of intermediate roughness and a
5 mixture of phase matrices.

6 The fast model constructed in this way is accurate enough to solve our inverse
7 problem. A typical difference between an exact calculation and our forward model is
8 shown in Fig. 9. The overall accuracy is within 1×10^{-4} in terms of L_{np} and the peak-
9 to-peak variation is 5×10^{-4} even in the worst case ($\sigma^2 = 0.03$). The overall error of
10 1×10^{-4} implies that the model bias is less than 10% of the observation error given
11 by ($\sqrt{\text{var}(L_{np})} = \sqrt{1.35 \times 10^{-6}} = 1.16 \times 10^{-3}$). The bias may be detected in the
12 residual of the inversion, but the influence on the roughness inference is negligible.

13 In calculating cloud reflectivity, a single-layer homogeneous cloud is assumed,
14 and the cloud optical thickness is set to 5 (roughly the saturation point of polarized
15 reflectance). No aerosol is assumed to be present above and below clouds. As
16 optically thick cold ice clouds occur in the upper troposphere, the radiometric
17 contribution from lower tropospheric aerosols is neglected. For the same reason,
18 the surface is assumed to be dark. There may be an influence from aerosols above
19 the cloud layer, such as transported mineral dust and stratospheric sulfates, but we
20 disregard them to be in line with previous studies. The influence of such aerosol
21 layers on inferences of cloud properties is beyond the scope of this paper but should
22 be investigated in the future.

23 The adding-doubling radiative transfer program formulated by de Haan et al.
24 (1987) with significant improvements by Huang et al. (2015) is used in the
25 calculation. The first-order scattering is calculated analytically and combined with
26 the multiple scattering results from the adding-doubling model, following the TMS
27 method (Nakajima and Tanaka, 1988). Further, the cloud reflectivity is multiplied by

Souichiro Hioki 5/12/16 7:50 PM
Deleted: 10

Souichiro Hioki 5/12/16 7:50 PM
Deleted: 1991

1 the transmissivity that changes due to ozone absorption; the transmissivity is
2 calculated from the monthly mean AIRS ozone concentration.

3 **2.2.3 Maximum Likelihood Estimation**

4 Once the inverse problem is formulated and the forward model is built, the last
5 step is to find the set of parameters for each pixel based on observations. The simple
6 but powerful maximum likelihood method with a normal error distribution is
7 appropriate for our problem because we have little knowledge about the
8 distribution of parameters (EOF scores). As each pixel typically contains five to eight
9 valid views (Section 2.1.2) at three wavelengths, the number of observations in each
10 pixel ranges from 15 to 24.

11 The standard deviation (SD) and correlation (Corr) of inferred parameters are
12 calculated in the framework of maximum likelihood estimation, and used to avoid
13 under-constrained inferences. The pixel is rejected if $SD(\text{EOF 1 Score}) > 0.02$,
14 $SD(\text{EOF 2 Score}) > 0.02$, or $\text{Corr}(\text{EOF 1 Score}, \text{EOF 2 Score}) > 0.3$. The standard
15 deviation and the correlation depend strongly on the observation geometry and
16 particle model and are almost independent of the observed polarized reflectivity.
17 Therefore, this rejection process can be interpreted as the refinement of pixels
18 based on the information content to achieve a reliable inference.

19 The error distribution is confirmed to be normal (see Section 2.1.1), so if the
20 problem is not strongly nonlinear, the parameters' error distributions are expected
21 to be normal as well (Rodgers, 2000). As expected, Fig. 10 demonstrates that the
22 application of the maximum likelihood method with synthetic L_{np} data results in a
23 symmetric distribution about the EOF 1 score corresponding to the true roughness
24 parameter $\sigma^2 = 0.15$. The distribution is not strictly normal because the number of
25 observations in each pixel varies, but the error distribution of each pixel is
26 theoretically derivable, as well as the confidence interval.

Souichiro Hioki 5/12/16 7:50 PM

Deleted: .

Souichiro Hioki 5/12/16 7:50 PM

Deleted: 11

1 For the synthetic retrieval in Fig. 10, the median of the inverted EOF 1 score is
2 -0.00336 and the corresponding roughness parameter is $\sigma^2 = 0.14$. The
3 interquartile range of the EOF 1 score distribution is [-0.01146:0.00476], which
4 corresponds to the roughness parameter range of [0.05:0.36]. The result indicates
5 that our approach has a practical skill in estimating the particle roughness
6 parameter out of observations superimposed with noise. This is a remarkable
7 contrast with the traditional “best-fit” approach (cf. Fig. 1).

Souichiro Hioki 5/12/16 7:50 PM
Deleted: 11

8 The distribution of the χ^2 values for the synthetic retrieval is presented in Fig.
9 11. The χ^2 value is a variance-normalized residual squared sum that is defined for
10 each pixel, and follows the χ^2 distribution with degrees of freedom N_d if the
11 inversion is successful, where N_d is the observational degrees of freedom
12 (approximately, the number of observations in a pixel). As the χ^2 distribution of N_d
13 degrees of freedom has a peak about N_d , the distribution of the χ^2 value indicates
14 whether the inversion is successful. If the location of the peak of a distribution of χ^2
15 values is smaller than N_d , the observation error may be overestimated, and if the
16 location of the peak is larger than N_d , the observation error is underestimated, or
17 the forward model does not represent reality (Rodgers, 2000). The distribution in
18 Fig. 11 has a peak at about 12, and very few pixels have a χ^2 value over 40. This is a
19 reasonable distribution because the number of observations ($\approx N_d$) is about 15 to
20 24 for most pixels. Because the 95th percentile for the χ^2 distribution with 24
21 degrees of freedom is 36.42, it is no surprise that very few pixels have a χ^2 value
22 over 40.

Souichiro Hioki 5/12/16 7:50 PM
Deleted: 12

23 Figures 10 and 11 demonstrate the validity of our inference framework under
24 an idealized situation, where the error distribution and the true roughness
25 parameter are constant. In application to actual satellite data, however, the true
26 roughness parameter varies from pixel to pixel while the error distribution stays the
27 same. Therefore, the distribution of the EOF 1 score must be more spread out as a

Souichiro Hioki 5/12/16 7:50 PM
Deleted: 12

Souichiro Hioki 5/12/16 7:50 PM
Deleted: 11
Souichiro Hioki 5/12/16 7:50 PM
Deleted: 12

1 result of convolution of the error distribution and the true roughness parameter
2 distribution. In contrast, the χ^2 distribution is expected to be about the same. The
3 result of the application to actual data is given in the next section.

Souichiro Hioki 5/12/16 7:50 PM

Deleted: the

5 3. Results and Discussion

6 3.1 Roughness Parameter of Cold Ice Cloud over Oceans

7 With the cloud selection criteria listed in Table 1, 79192 pixels based on one
8 month of collocated PARASOL/MODIS data over oceans during September 2005
9 were selected for inversion. The information content was sufficient for full analysis
10 of 23359 pixels, for which results are presented in this section.

Souichiro Hioki 5/12/16 7:50 PM

Deleted: 79132 pixels

11 The histogram of the inferred EOF 1 score is presented in Fig. 12 for the
12 extratropical (latitude > 30°) oceans. The width of the histogram in Fig. 12 is
13 broader than the monodispersive roughness case (Fig. 10), indicating significant
14 variability in the microphysical properties of clouds. The median of the distribution
15 is -0.0293 , corresponding to a surface roughness parameter of 2.82. The
16 interquartile range of the EOF 1 score is $[-0.0429; -0.0165]$, implying 50% of the
17 data is within the roughness parameter (σ^2) range of $[0.65; 13.6]$. The result
18 supports the use of the roughened particle model in extratropical ice cloud
19 retrievals as suggested by previous studies. While our analysis is limited to very
20 cold ice clouds over ocean, the validity of using roughened crystals in the MODIS
21 Collection 6 ice model is supported, although further explorations into warmer and
22 optically thinner clouds are desirable. In general, cloud particles become more
23 complex as the cloud temperature increases (Heymsfield, 2002), thus we expect
24 more roughened particles in warmer clouds that are not included in our analysis.

Souichiro Hioki 5/12/16 7:50 PM

Deleted: 13 based on one month of collocated PARASOL/MODIS data over

Souichiro Hioki 5/12/16 7:50 PM

Deleted: during September 2005.

Souichiro Hioki 5/12/16 7:50 PM

Deleted: 13

Souichiro Hioki 5/12/16 7:50 PM

Deleted: 11

Souichiro Hioki 5/12/16 7:50 PM

Moved down [3]: As the roughness parameter of 2.82 lies outside of our prescribed roughness parameter range (0 to 0.7), it is an estimate by extrapolation. Yet, this projection of roughness parameter implies that the conventional degree of roughness may not be sufficient to represent actual cloud particles with the aggregate column model. The proportion of pixels that contains inferred roughness parameter $\sigma^2 > 0.7$ is 74%, which also indicates the limit of this particle shape. As the accuracy of roughness approximation for such a large roughness parameter is questionable, a particle shape that can fit observations with less intense roughening may be suitable for the representation of natural clouds.

25 The distributions of the χ^2 value in the tropics and extratropics are separately
26 presented in Fig. 13. As discussed in the previous section, the distribution of χ^2

Souichiro Hioki 5/12/16 7:50 PM

Deleted: 14

Souichiro Hioki 5/12/16 7:50 PM

Deleted: a

1 values indicates the validity of the inversion. While the distribution of the χ^2 values
2 in the extratropics shows reasonable behavior (Fig. 13a), the distribution of the χ^2
3 values in the tropics has a very long tail with the mean χ^2 being 59.7, which is
4 unacceptably large (Fig. 13b). This long tail implies that our forward model does not
5 properly reproduce the observed L_{np} field in the tropics, presumably because some
6 underlying assumptions are not appropriate or the information content is not
7 enough. Some possibilities that violate our underlying assumptions include sub-
8 pixel scale cloud heterogeneity, the presence of ice particles with other habits or
9 aspect ratios, their vertical heterogeneity, cloud 3-D effects, and the effect of
10 aerosols.

3.2 Unexpectedly Large Roughness Values in the Extratropics

As the roughness parameter of 2.82 lies outside of our prescribed roughness
parameter range (0 to 0.7), it is an estimate by extrapolation. Yet, this projection of
roughness parameter implies that the conventional degree of roughness may not be
sufficient to represent actual cloud particles with the aggregate column model. The
proportion of pixels that contains inferred roughness parameter $\sigma^2 > 0.7$ is 74%,
which also indicates the limit of this particle shape. As the accuracy of roughness
approximation for such a large roughness parameter is questionable, a particle
shape that can fit observations with less intense roughening may be suitable for the
representation of natural clouds.

To attribute the cause of unphysically large roughness value in the
extratropics, the same retrieval process is repeated assuming three additional
particle shapes. Figure 14a shows the original inference with aggregate of columns
shape, in which the observation density peaks away from the line connecting 10
points that corresponds to prescribed roughness values. The aggregate of plates
(Fig. 14d) performs worst among the tested particles, and the solid bullet rosette

- Souichiro Hioki 5/12/16 7:50 PM
Deleted: a
- Souichiro Hioki 5/12/16 7:50 PM
Deleted: 14a
- Souichiro Hioki 5/12/16 7:50 PM
Deleted: 14b

- Souichiro Hioki 5/12/16 7:50 PM
Deleted: .

- Souichiro Hioki 5/12/16 7:50 PM
Deleted: and
- Souichiro Hioki 5/12/16 7:50 PM
Deleted: . The extension
- Souichiro Hioki 5/12/16 7:50 PM
Deleted: the parameter space to include other habits will be investigated in future work.
- Souichiro Hioki 5/12/16 7:50 PM
Deleted: 3.2 Possible Causes of
- Souichiro Hioki 5/12/16 7:50 PM
Moved (insertion) [3]

1 [shape \(Fig. 14c\) shows the largest overlap of parameter space and observation](#)
2 [density. These results indicate that the roughness retrieval is sensitive to an](#)
3 [assumed particle shape.](#)

4 [We also investigated the contamination by multi-layer clouds and aerosol](#)
5 [above clouds by collocating Cloud-Aerosol Lidar with Orthogonal Polarization](#)
6 [\(CALIOP\) vertical feature mask product and cloud layer product. The distribution of](#)
7 [the retrieved EOF scores is approximately the same even when assuring the absence](#)
8 [of aerosol above cloud and limiting the analysis to single-layer clouds \(Fig. 15\).](#)
9 [However, the removal of the multi-layer clouds helps to reduce the number of pixels](#)
10 [with very large \$\chi^2\$ values.](#)

11 **3.3 Inference Failure in Tropics**

12 To gain a better insight into the cause of the long tail in the tropics, a case
13 study is conducted for two cloudy scenes: a typical extratropical scene and a tropical
14 cloud scene with systematically large χ^2 values. Figure [16](#) displays true color
15 composites from PARASOL with markers indicating the locations of detailed
16 analysis. A green circle is shown where the χ^2 value is less than the 95th percentile
17 of the χ^2 distribution (reasonable deviation from the forward model), and a
18 magenta cross is shown where the χ^2 value is more than the 95th percentile (too far
19 from the forward model). The locations of the magenta crosses in Fig. [16a](#) (typical
20 extratropics) are somewhat systematic; they appear at cloud boundaries or at
21 isolated locations. This may suggest that cloud heterogeneity and cloud 3-D effects
22 cause a small number of inference failures in the extratropics.

23 In contrast to the typical extratropical scene, magenta crosses are prevalent
24 throughout the tropical scene in Fig. [16b](#). Since the cloud reflectivity is comparable
25 to the typical extratropical scene, it is not likely that the inference failures are due to
26 contamination by surface reflection. Also failures cannot be fully explained by 3-D

Souichiro Hioki 5/12/16 7:50 PM

Deleted: 15

Souichiro Hioki 5/12/16 7:50 PM

Deleted: 15a

Souichiro Hioki 5/12/16 7:50 PM

Deleted: 15b

1 effects of clouds as a few green circles appear randomly. Flaws in the assumptions
2 that depend little on the relative location in a cloud, such as cloud particle shape and
3 cloud heterogeneity (e.g. Oreopoulos, 2009), or the lack of information content due
4 to the limited scattering angle range are therefore suspected as causes of the
5 inference failure in the tropics.

6 A close investigation into the correlation of EOF 1 and EOF 2 scores supports
7 the hypothesis that the information content is limited. Figure 17 shows the
8 coefficient of correlation between retrieved EOF 1 and EOF 2 scores. While the
9 distribution is centered at 0 in the extratropics, it peaks at -0.8 in the tropics,
10 indicating limited information content to constrain the parameter space. Also, a
11 validation of retrieved cloud top height using the CALIOP data indicated that the
12 cloud top heights are not properly retrieved in the tropics.

13 The insufficient information content for roughness and cloud top height
14 retrievals is presumably caused by that sampled scattering angles are concentrated
15 near the backscattering direction and zenith angles are small. The directional
16 distribution of the polarized reflectivity is not well captured to constrain the degree
17 of roughness and the spectral contrast of the Rayleigh scattering signal is too weak
18 to infer cloud top height accurately.

19 **3.4 Comparison with Scattering Properties in the MODIS Retrieval Scheme**

20 By definition of our EOF scores, the inverted EOF scores translate into $-P_{12}$ on
21 a pixel-by-pixel basis. The reconstructed $-P_{12}$ reflects a variation due to observation
22 error and natural variability. To accurately interpret the result, we reconstructed
23 $-P_{12}$ from extratropical data with a precise EOF 2 score ($SD(EOF2) < 0.01$). The
24 area shaded with gray in Fig. 18 shows the interquartile range of the reconstructed
25 $-P_{12}$ which indicates that 50% of our extratropical observations fall within the
26 shade at a given scattering angle. The blue line is $-P_{12}$ for the particle shape used in

Souichiro Hioki 5/12/16 7:50 PM

Deleted: A separate validation using the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) data indicated that our cloud top height retrieval is not in agreement with CALIOP cloud top height data, possibly indicating the limited information content.

Souichiro Hioki 5/12/16 7:50 PM

Deleted: 3

Souichiro Hioki 5/12/16 7:50 PM

Deleted: 16

1 MODIS Collection 6, and the green line is that for the shape in MODIS Collection 5.
2 Both particle models assume a gamma distribution with effective particle size of 60
3 μm and effective variance of 0.1. The blue line (Collection 6) is closer to our
4 reconstruction, while the green line (Collection 5) significantly deviates from our
5 reconstruction. This result indicates that the particle habit adopted for MODIS
6 Collection 6 is more consistent with polarimetric observations than the habit
7 mixture used for MODIS Collection 5, for which only one of the habits **includes** a
8 limited degree of roughness.

Souichiro Hioki 5/12/16 7:50 PM

Deleted: includes

9 The reconstructed $-P_{12}$ shows stronger side scattering between 80° and 120° than
10 the MODIS Collection 6 particle model. As the increasing roughness enhances side
11 scattering, weak side scattering of the column aggregate shape may be responsible
12 for the unphysically large roughness parameter in the extratropical inferences. By
13 using a shape that has stronger side scattering, it is likely that the degree of rough-
14 ness that is needed to explain the observations becomes smaller. An example of such
15 a habit mixture is shown by the thick magenta line in Fig. 18. A mixture of two habits
16 (70% column aggregate particles with roughness parameter of $\sigma^2 = 0.8$ and 30%
17 severely roughened hollow bullet rosette particles with $\sigma^2 = 0.5$) included in the
18 scattering property library by Yang et al. (2013), results in a phase function with
19 strong side scattering.

Souichiro Hioki 5/12/16 7:50 PM

Deleted: 16

20 **4. Summary and Future Directions**

21 In this study, the particle roughness parameter of very cold ice clouds over
22 ocean is inferred by employing a new framework that is resilient to the
23 observational error. The distinct feature of the framework is the continuous
24 parameter space that is constructed with an empirical orthogonal function (EOF)
25 analysis. Two EOFs are found to be sufficient to explain the variation of $-P_{12}$ with a

1 changing particle roughness parameter, substantially reducing the number of
2 parameters for the forward model.

3 From unpolarized cloud reflection at a scattering angle of 170° , the
4 observational error of the PARASOL data is empirically estimated. Supported by the
5 error analysis with parametric bootstrapping, the maximum likelihood method is
6 applied to the inverse problem. The method provides error estimates and
7 correlations for inverted parameters, which are unavailable with the “best-fit”
8 approach used in the previous studies. To correctly incorporate the effect of
9 atmospheric Rayleigh scattering, the cloud-top height is inferred simultaneously.

10 The application of the present method to cold ice clouds over extratropical
11 oceans results in a roughness parameter of 2.82, implying that the use of the
12 roughened particle model is suitable for cloud property retrievals. By contrasting
13 the distribution of χ^2 values in the tropics and extratropics, we find that the
14 performance of our method needs to be enhanced in the tropics. Possible future
15 technical improvements may be an extension of parameter space to include multiple
16 particle shapes, application to optically thin clouds, and integration with
17 unpolarized radiance observations.

18 The reconstructed $-P_{12}$ curve shows better consistency with $-P_{12}$ from the
19 particle shape model used in MODIS Collection 6 than $-P_{12}$ from MODIS Collection
20 5. The addition of roughness and a hollow bullet rosette particle shape to the MODIS
21 Collection 6 model further improves the consistency.

22 Since its launch in 2004, the PARASOL satellite observed global polarimetric
23 reflectivity nearly simultaneously with MODIS for five years until leaving the A-train
24 constellation in 2009. A large amount of PARASOL data is available to apply the
25 framework described in this paper. Local variations of the roughness parameter,
26 correlation of the roughness parameter to other meteorological data, and the impact
27 of cloud heterogeneity are to be investigated in our future study.

1 **Acknowledgements**

2 The authors thank the ICARE Data and Service Center for providing
3 POLDER/PARASOL data, the NASA LAADS system for providing MODIS atmosphere
4 products, and the NASA GSFC GES DAAC for providing AIRS data. The ice particle
5 scattering calculations are conducted at the Texas A&M University Supercomputing
6 Facility. The authors are grateful to Dr. Lei Bi for help in light scattering
7 computations. This work was funded by NASA Grants NNX11AR06G and
8 NNX15A12H, and the authors are grateful for continued support from Dr. Hal
9 Maring.

10

1 **References**

- 2 Baran, A. J. and Francis, P. N.: On the radiative properties of cirrus cloud at solar and
3 thermal wavelengths: A test of model consistency using high-resolution airborne
4 radiance measurements. *Quart. J. Roy. Meteor. Soc.*, 130, 763–778, doi:
5 | 10.1256/qj.03.151, 2004.
- 6 Baran, A. and C-Labonnote, L.: On the reflection and polarisation properties of ice
7 cloud, *J. Quant. Spectrosc. Ra.*, 100, 41–54, 2006.
- 8 Baran, A. J. and C-Labonnote, L.: A self-consistent scattering model for cirrus. I: The
9 solar region, *Quart. J. Roy. Meteor. Soc.*, 133, 1899–1912, 2007.
- 10 Baum, B. A., Yang, P., Hu, Y. X., and Feng, Q.: The impact of ice particle roughness on
11 the scattering phase matrix, *J. Quant. Spectrosc. Ra.*, 111, 2534–2549, 2010.
- 12 Buriez, J. C., Vanbauce, C., Parol, F., Goloub, P., Herman, M., Bonnel, B., Fouquart, Y.,
13 Couvert, P., and Seze, G.: Cloud detection and derivation of cloud properties from
14 POLDER, *Int. J. Remote Sens.*, 18, 2785-2813, doi: 10.1080/014311697217332,
15 1997.
- 16 Chepfer, H., Brogniez, G., and Fouquart, Y.: Cirrus clouds' microphysical properties
17 deduced from POLDER observations, *J. Quant. Spectrosc. Ra.*, 60, 375–390, 1998.
- 18 Cole, B. H., Yang, P., Baum, B. A., Riedi, J., and C-Labonnote, L.: Ice particle habit and
19 surface roughness derived from PARASOL polarization measurements, *Atmos.*
20 *Chem. Phys.*, 14, 3739–3750, 2014.
- 21 Cole, B. H., Yang, P., Baum, B. A., Riedi, J., C-Labonnote, L., Thieuleux, F., and Platnick,
22 S.: Comparison of PARASOL observations with polarized reflectances simulated
23 using different ice habit mixtures, *J. Appl. Meteor. Climatol.*, 52, 186–196, 2013.

1 Deschamps, P., Bréon, F., Leroy, M., Podaire, A., Bricaud, A., Buriez, J. C., and Sèze, G.:
2 The POLDER mission – instrument characteristics and scientific objectives, IEEE
3 T. Geosci. Remote, 32, 598–615, 1994.

4 de Haan, J. F., Bosma, P. B., and Hovenier, J. W.: The adding method for multiple scat-
5 tering calculations of polarized light, *Astron. Astrophys.* 183, 371–391, 1987.

6 Evans, M. J. and Rosenthal, J. S., *Probability and statistics: The science of uncertainty*,
7 2nd ed., W. H. Freeman and Company, 2010.

8 Fougnie, B., Bracco, G., Lafrance, B., Ruffel, C., Hagolle, O., and Tinell, C.: PARASOL in-
9 flight calibration and performance, *Appl. Opt.*, 46, 5435–5451, 2007.

10 Heymsfield, A. J., Bansemer, A., Field, P. R., Durden, S. L., Stith, J. L., Dye, J. E., Hall, W.,
11 and Grainger, C. A.: Observations and parameterizations of particle size distribu-
12 tions in deep tropical cirrus and stratiform precipitating clouds: Results from in
13 situ observations in TRMM field campaigns, *J. Atmos. Sci.*, 59, 3457–3491, 2002.

14 Heymsfield, A. J., Schmitt, C., and Bansemer, A.: Ice cloud particle size distributions
15 and pressure-dependent terminal velocities from in situ observations at temper-
16 atures from 0° to -86°C, *J. Atmos. Sci.*, 70, 4123–4154, 2013.

17 Holz, R. E., Platnick, S., Meyer, K., Vaughan, M., Wind, G., Dutcher, S., Ackerman, S.,
18 Heidinger, A., Amarasinghe, N., Wang, C., and Yang, P.: Resolving cirrus optical
19 depth biases between CALIOP and MODIS using IR retrievals. *Atmos. Chem. Phys.*,
20 submitted Sept. 2015.

21 Huang, X., Yang, P., Kattawar, G., and Liou, K. N.: Effect of mineral dust aerosol aspect
22 ratio on polarized reflectance, *J. Quant. Spectrosc. Ra.*, 151, 97–109, 2015.

23 Inoue, T.: A Cloud type classification with NOAA 7 split-window measurements, *J.*
24 *Geophys. Res.*, 92, 3991–4000, 1987.

- 1 King, M. D., Menzel, W. P., Kaufman, Y. J., Tanré, D., Gao, B.-C., Platnick, S., Ackerman,
2 S. A., Remer, L. A., Pincus, R., and Hubanks, P. A.: Cloud and aerosol properties,
3 precipitable water, and profiles of temperature and water vapor from MODIS,
4 IEEE T. Geosci. Remote, 41, 442–458, 2003.
- 5 Knap, W. H., C.-Labonnote, L., Brogniez, G., and Stammes, P.: Modeling total and po-
6 larized reflectances of ice clouds: evaluation by means of POLDER and ATSR-2
7 measurements, Appl. Opt., 44, 4060–4073, 2005.
- 8 C.-Labonnote, L., Brogniez, G., Buriez, J.-C., Doutriaux-Boucher, M., Gayet, J., and
9 Macke, A.: Polarized light scattering by inhomogeneous hexagonal monocrystals:
10 Validation with ADEOS-POLDER measurements, J. Geophys. Res., 106, 12139–
11 12153, 2001.
- 12 Liu, C., Yang, P., Minnis, P., Loeb, N., Kato, S., Heymsfield, A., and Schmitt, C.: A two-
13 habit model for the microphysical and optical properties of ice clouds, Atmos.
14 Chem. Phys., 14, 13719–13737, 2014.
- 15 Macke, A., and Mishchenko, M. I.: Applicability of regular particle shapes in light
16 scattering calculations for atmospheric ice particles. Appl. Opt., 35, 4291-4296,
17 doi:10.1364/AO.35.004291, 1996.
- 18 Macke, A., Mueller, J., and Raschke, E.: Single scattering properties of atmospheric ice
19 crystals, J. Atmos. Sci., 53, 2813–2825, 1996.
- 20 Mapes, B. E., and Houze, R. A.: Cloud clusters and superclusters over the oceanic
21 warm pool, Mon. Weather Rev., 121, 1398–1415, 1993.
- 22 Masuda, K., Ishimoto, H., and Takashima, T.: Retrieval of cirrus optical thickness and
23 ice-shape information using total and polarized reflectance from satellite meas-
24 urements, J. Quant. Spectrosc. Ra., 75, 39–51, 2002.

- 1 Masuda, K. and Takashima, T.: Feasibility study of derivation of cirrus information
2 using polarimetric measurements from satellite, *Remote Sens. Environ.*, 39, 45–
3 59, 1992.
- 4 Minnis, P., Heck, P. W., and Young, D. F.: Inference of cirrus cloud properties using
5 satellite-observed visible and infrared radiances. Part II: Verification of theoretical
6 cirrus radiative properties, *J. Atmos. Sci.*, 50, 1305–1322, 1993.
- 7 Minnis, P., Sun-Mack, S., Young, D. F., Heck, P. W., Garber, D. P., Chen, Y., Spangenberg,
8 D. A., Arduini, R. F., Trepte, Q. Z., Smith, Jr., W. L., Ayers, J. K., Gibson, S. C.,
9 Miller, W. F., Hong, G., Chakrapani, V., Takano, Y., Liou, K.-N., Xie, Y., and Yang, P.:
10 CERES Edition-2 cloud property retrievals using TRMM VIRS and Terra and Aqua
11 MODIS data - Part I: Algorithms, *IEEE T. Geosci. Remote*, 49, 4374–4400, 2011.
- 12 Nakajima, T. and King, M. D.: Determination of the optical thickness and effective
13 particle radius of clouds from reflected solar radiation measurements. Part I:
14 Theory, *J. Atmos. Sci.*, 47, 1878–1893, 1990.
- 15 Nakajima, T. and Tanaka, M.: Algorithms for radiative intensity calculations in moderately
16 thick atmospheres using a truncation approximation, *J. Quant. Spectrosc.*
17 *Ra.*, 40, 51-69, 1988.
- 18 Neshyba, S. P., Lowen, B., Benning, M., Lawson, A., and Rowe, P. M.: Roughness metrics
19 of prismatic facets of ice, *J. Geophys. Res.*, 118, 3309–3318, 2013.
- 20 Oreopoulos, L., Platnick, S., Hong, G., Yang, P., and Cahalan, R. F.: The shortwave radiative
21 forcing bias of liquid and ice clouds from MODIS observations, *Atmos. Chem.*
22 *Phys.*, 9, 5865–5875, 2009.
- 23 Parol, F., Buriez, J.-C., Vanbauce, C., Couvert, P., Sèze, G., Goloub, P., and Cheinet, S.:
24 First results of the POLDER “Earth Radiation Budget and Clouds” operational algorithm,
25 *IEEE T. Geosci. Remote*, 37, 1597–1612, 1999.

- 1 Platnick, S., Ackerman, S., King, M. D., et al.: MODIS atmosphere L2 cloud product
2 (06_L2), NASA MODIS Adaptive Processing System, Goddard Space Flight Center,
3 doi:dx.doi.org/10.5067/MODIS/MYD06_L2.006, 2015.
- 4 Platnick, S., King, M. D., Ackerman, S. A., Menzel, W. P., Baum, B. A., Riedi, J. C., and
5 Frey, R. A.: The MODIS cloud products: Algorithms and examples from Terra,
6 IEEE T. Geosci. Remote, 41, 459–473, 2003.
- 7 Rodgers, C. D., Inverse methods for atmospheric sounding: Theory and practice,
8 World Scientific Publishing, 2000.
- 9 Rolland, P., Liou, K.-N., King, M. D., Tsay, S.-C., and McFarquhar, G. M.: Remote sens-
10 ing of optical and microphysical properties of cirrus clouds using Moderate-
11 Resolution Imaging Spectroradiometer channels: Methodology and sensitivity to
12 physical assumptions, J. Geophys. Res., 105, 11721–11738, 2000.
- 13 Sourdeval, O., C-Labonnote, L., Baran, A. J. and Brogniez, G.: A methodology for sim-
14 ultaneous retrieval of ice and liquid water cloud properties. Part I: Information
15 content and case study. Quart. J. Roy. Meteor. Soc., 141, 870–882, doi:
16 10.1002/qj.2405, 2015.
- 17 Ulanowski, Z., Hesse, E., Kaye, P. H., and Baran, A. J.: Light scattering by complex ice-
18 analogue crystals, J. Quant. Spectrosc. Ra., 100, 382–392,
19 doi:10.1016/j.jqsrt.2005.11.052, 2006.
- 20 Ulanowski, Z., Kaye, P. H., Hirst, E., Greenaway, R. S., Cotton, R. J., Hesse, E., and Colli-
21 er, C. T.: Incidence of rough and irregular atmospheric ice particles from Small Ice
22 Detector 3 measurements, Atmos. Chem. Phys., 14, 1649–1662, doi:10.5194/acp-
23 14-1649-2014, 2014.
- 24 Um, J. and McFarquhar, G. M.: Single-scattering properties of aggregates of bullet
25 rosettes in cirrus, J. Appl. Meteorol. Climatol., 46, 757-775, 2007.

- 1 Um, J. and McFarquhar, G. M.: Single-scattering properties of aggregates of plates,
2 Quart. J. Roy. Meteor. Soc., 135, 291-304, 2009.
- 3 van Diedenhoven, B., Cairns, B., Geogdzhayev, I. V., Fridlind, A. M., Ackerman, A. S.,
4 Yang, P., and Baum, B. A.: Remote sensing of ice crystal asymmetry parameter us-
5 ing multi-directional polarization measurements - Part 1: Methodology and eval-
6 uation with simulated measurements, Atmos. Meas. Tech., 5, 2361-2374, 2012.
- 7 van Diedenhoven, B., Fridlind, A. M., Cairns, B., and Ackerman, A. S.: Variation of ice
8 crystal size, shape, and asymmetry parameter in tops of tropical deep convective
9 clouds, J. Geophys. Res., 119, 11809-11825, 2014.
- 10 Waliser, D. E., Li, J.-L. F., Woods, C. P., Austin, R. T., Bacmeister, J., Chern, J., Del Genio,
11 A., Jiang, J. H., Kuang, Z., Meng, H., Minnis, P., Platnick, S., Rossow, W. B., Stephens,
12 G. L., Sun-Mack, S., Tao, W.-K., Tompkins, A. M., Vane, D. G., Walker, C., and Wu, D.:
13 Cloud ice: A climate model challenge with signs and expectations of progress, J.
14 Geophys. Res., 114, D00A21, doi:10.1029/2008JD010015, 2009.
- 15 Xie, Y., Yang, P., Kattawar, G. W., Minnis, P., and Hu, Y. X.: Effect of the inhomogeneity
16 of ice crystals on retrieving ice cloud optical thickness and effective particle size,
17 J. Geophys. Res., 114, D11203, doi:10.1029/2008JD011216, 2009.
- 18 Yang, P., Bi, L., Baum, B. A., Liou, K.-N., Kattawar, G. W., Mishchenko, M. I., and Cole,
19 B.: Spectrally consistent scattering, absorption, and polarization properties of
20 atmospheric ice crystals at wavelengths from 0.2 to 100 μm , J. Atmos. Sci., 70,
21 330-347, 2013.
- 22 Yang, P., Hong, G., Kattawar, G. W., Minnis, P., and Hu, Y.: Uncertainties associated
23 with the surface texture of ice particles in satellite-based retrieval of cirrus
24 clouds: Part II - Effect of particle surface roughness on retrieved cloud optical

1 thickness and effective particle size, IEEE T. Geosci. Remote, 46, 1948–1957,
2 2008b.

3 Yang, P., Kattawar, G. W., Hong, G., Minnis, P., and Hu, Y.: Uncertainties associated
4 with the surface texture of ice particles in satellite-based retrieval of cirrus
5 clouds - Part I: Single-scattering properties of ice crystals with surface roughness,
6 IEEE T. Geosci. Remote, 46, 1940–1947, 2008a.

7 Yang, P., and Liou, K. N.: Geometric-optics-integral-equation method for light
8 scattering by nonspherical ice crystals, Appl. Opt., 35, 6568–6584, 1996.

9 Yang, P., Zhang, L., Hong, G., Nasiri, S. L., Baum, B. A., Huang, H.-L., King, M. D., and
10 Platnick, S.: Differences between collection 4 and 5 MODIS ice cloud opti-
11 cal/microphysical products and their impact on radiative forcing simulations,
12 IEEE T. Geosci. Remote, 45, 2886–2899, 2007.

13 Yurkin, M. A., Maltsev, V. P., and Hoekstra, A. G.: The discrete dipole approximation
14 for simulation of light scattering by particles much larger than the wavelength, J.
15 Quant. Spectrosc. Ra., 106, 546–557, 2007.

16 Zhang, Z., Yang, P., Kattawar, G., Riedi, J. C., C.-Labonnote, L., Baum, B. A., Platnick, S.,
17 and Huang, H.-L.: Influence of ice particle model on satellite ice cloud retrieval:
18 lessons learned from MODIS and POLDER cloud product comparison, Atmos.
19 Chem. Phys., 9, 7115–7129, 2009.

20

1 **Tables**


2

3 **Table 1.** PARASOL pixel and view selection criteria.

Parameters	Criteria	Applied to
MODIS Brightness Temperature at 11 μ m	Median is less than 208 K	Pixel
MODIS Infrared Cloud Phase	Ice	Pixel
PARASOL Ocean/Land Flag	Ocean	Pixel
Number of Valid Views	At least 5	Pixel
Scattering Angle	60° to 160°	View
Sunlint Angle	Greater than 30°	View


4

1 **Figures**




2


3 **Fig. 1.** The response of the conventional “best-fit” approach to a synthetic signal
4 with and without random measurement noise. The addition of noise to the synthetic
5 signal results in a distribution of the roughness parameter (hatched bars), from
6 which the true roughness cannot be inferred. This figure is to be compared to Fig.
7 11.
8




1
 2 **Fig. 2.** Observation density of modified polarized reflectivity (L_{nmp}) over the West-
 3 ern Pacific during September 2005. L_{nmp} crosses zero at a scattering angle of ap-
 4 proximately 170° . The data in the rectangular box is used to derive the histogram in
 5 Fig. 3.
 6




1
2 **Fig. 3.** Histogram of observed normalized polarized radiance (L_{np}) from the data in
3 the rectangular box in Fig. 2. The solid line is the simulated error using a parametric
4 bootstrapping method with $s = 0.00095$. The agreement is sufficient for estimating
5 the noise level.
6



1
2 **Fig. 4.** Sum of squared error as a function of standard error (s) of the original sensor
3 noise. The minimum error is achieved when $s = 0.00095$.
4



1
 2 **Fig. 5.** The simulated variance of L_{np} as a function of L_n . The variance of L_{np} increases
 3 as the normalized radiance L_n (brightness of a pixel) increases, becoming nearly
 4 constant at $\text{var}(L_{np}) = 1.35 \times 10^{-6}$ once L_n reaches $L_n = 0.2$. Insets show that the
 5 distribution of L_{np} tends to a normal distribution, justifying the use of a normal dis-
 6 tribution as an error distribution of L_{np} for a reflective cloudy pixel.
 7



1
 2 **Fig. 6.** The pairs of EOF scores needed to reconstruct the original $-P_{12}$. The EOF 1
 3 score is a monotonic function of particle roughness parameter σ^2 . The EOF 2 score
 4 reaches a minimum at particle roughness parameter of $\sigma^2 = 0.1$.
 5

1
2
3
4
5
6
7






Fig. 7. The relationship between the particle roughness parameter and the EOF1 score. The natural logarithm of the particle roughness parameter is nearly linear to the EOF 1 score. This implies that the particle roughness can be directly inferred from the EOF 1 score.

Souichiro Hioki 5/12/16 7:50 PM




Deleted:
Souichiro Hioki 5/12/16 7:50 PM
Deleted: Comparison of the reconstructed and the original $-P_{12}$ elements of the phase matrix. Reconstructed $-P_{12}$ (colored solid lines) agrees with original $-P_{12}$ (black dashed lines). Only the first and second EOFs are used in the reconstruction. This confirms that two EOFs explain almost the entire variation of $-P_{12}$ due to the particle roughness change. - [...](#) [1]




1
 2 **Fig. 8.** The impact of particle roughness parameter change ($\sigma^2 = 0.15 \rightarrow 0.5$) and
 3 cloud top pressure change (200 \rightarrow 500 hPa). The magnitudes of the differences are
 4 comparable while the directional patterns are different. In this plot, the solar zenith
 5 angle is 54° and the viewing zenith angle is 30° .
 6

Souichiro Hioki 5/12/16 7:50 PM
 Deleted: 9



1
 2 | **Fig. 9.** Difference in L_{np} between exact radiative transfer calculations and our sim-
 3 | plified forward model. At almost all angles, the difference is less than 1×10^{-4} . The
 4 | polar plot shows the distribution of bias when the particle roughness parameter is
 5 | $\sigma^2 = 0.15$. The bias is a function of scattering angle. However, the magnitude of er-
 6 | ror is acceptably small compared to the random observational error.
 7


Souichiro Hioki 5/12/16 7:50 PM
 Deleted: 10



1
2
3
4
5
6

Fig. 10. The distribution of inferred EOF 1 scores for synthetic data with and without noise. The distribution for the noise-added synthetic data is symmetric about the EOF 1 score corresponding to the true roughness. The median of EOF 1 score is -0.00336, corresponding to roughness parameter of $\sigma^2 = 0.14$.


Souichiro Hioki 5/12/16 7:50 PM
Deleted: 11



1
 2 | **Fig. 11.** Frequency distribution of the χ^2 values (variance-normalized residual
 3 square sum). The distribution has a peak at about 12, tapering to nearly zero at ap-
 4 proximately 40. This is a reasonable distribution because most pixels contain 15 to
 5 24 observations.
 6


Souichiro Hioki 5/12/16 7:50 PM

Deleted: 12



1
 2 **Fig. 12.** The distribution of EOF 1 scores obtained from cold ice clouds over extra-
 3 tropical oceans during September 2005. The median of the EOF 1 score is -0.0293,
 4 corresponding to a roughness parameter of 2.82. Consistent with previous studies,
 5 roughened particles better simulate the measured polarized reflectivity.
 6

Souichiro Hioki 5/12/16 7:50 PM
 Deleted: 13
 Souichiro Hioki 5/12/16 7:50 PM
 Deleted: ocean



1
 2 | **Fig. 13.** Distributions of χ^2 values in the tropics and extratropics. The distribution of
 3 the χ^2 value in the tropics (b) implies that the forward model is not correctly simu-
 4 lating the reflectivity in the tropics, while the distribution of the χ^2 value in the ex-
 5 tratropics (a) indicates successful inversion.


6 |

Souichiro Hioki 5/12/16 7:50 PM

Deleted: 14


Souichiro Hioki 5/12/16 7:50 PM


Formatted: Left, Line spacing: single




1
2
3
4
5
6
7

Fig. 14. Distributions of EOF 1 and EOF 2 scores with different particle shapes. The observation frequency is shaded with color, and the solid line connects the EOF scores for 10 prescribed roughness values (circles). (a) The result of inference with aggregate of columns, (b) hollow column, (c) solid bullet rosette, and (d) aggregate of plates.






1
 2 Fig. 15. The distribution of retrieved EOF 1 and EOF 2 scores when filtering out the
 3 multilayer clouds and cloud with aerosol loading on top. The observation frequency
 4 is color shaded, and the EOF scores for aggregate of column models (circles) are
 5 connected by a line. This analysis is conducted on a different EOF space from Fig. 6.
 6 The minimum degree of roughness is $\sigma^2 = 0.03$ and the maximum is $\sigma^2 = 1.0$.
 7




1
 2 **Fig. 16.** Comparison of (a) a typical cloud scene in the extratropics and (b) a cloud
 3 scene in the tropics where the χ^2 values are much larger than expected. Green circles
 4 are inference locations where the χ^2 value is less than the 95th percentile of the
 5 χ^2 distribution, whereas magenta crosses are inference locations where the χ^2 val-
 6 ue exceeds the 95th percentile. These figures indicate that the causes of a large χ^2
 7 value may be different in the extratropics and tropics.
 8

Souichiro Hioki 5/12/16 7:50 PM
 Formatted: Left, Line spacing: single



1
2
3
4
5
6
7

Fig. 17. Histograms of the coefficient of correlation between EOF 1 and EOF 2 scores. Out of 79192 total inferred pixels, 49902 pixels are selected by the condition $SD(\text{EOF 1 Score}) < 0.02, SD(\text{EOF 2 Score}) < 0.02$. The results in Fig. 12-14 are based on the data within the center six bins in these histograms.



2
3 **Fig. 18.** Comparison of inferred $-P_{12}$ and the counterparts used in MODIS Collec-
4 tions 5 and 6. The $-P_{12}$ of MODIS Collection 6 (blue line) is more consistent with the
5 reconstructed $-P_{12}$ (black thick line) than the $-P_{12}$ of MODIS Collection 5 (green
6 line). However, better consistency is obtained with a two-habit model (thick magenta
7 line), by increasing the roughness to $\sigma^2 = 0.8$ and adding 30% of hollow bullet
8 rosette particles.

Souichiro Hioki 5/12/16 7:50 PM

Deleted: 16