

1 **Short- and long-term variability of spectral solar UV**
2 **irradiance at Thessaloniki, Greece: effects of changes in**
3 **aerosols, total ozone and clouds**

4

5 **Ilias Fountoulakis¹, Alkiviadis F. Bais¹, Kostantinos Fragkos¹, Charickleia**
6 **Meleti¹, Kleareti Tourpali¹, Melina Maria Zempila^{1,2}**

7 [1]{Aristotle University of Thessaloniki, Laboratory of Atmospheric Physics, Thessaloniki,
8 Greece}

9 [2]{Now to: Natural Resource Ecology Laboratory, Colorado State University, Fort Collins,
10 USA}

11

12 Correspondence to: I. Fountoulakis (iliasn@auth.gr)

13

14 **Abstract**

15 In this study, we discuss the short- and the long-term variability of spectral UV irradiance at
16 Thessaloniki, Greece using a long, quality-controlled data set from two Brewer
17 spectrophotometers. Long-term changes in spectral UV irradiance at 307.5, 324 and 350 nm
18 for the period 1994 – 2014 are presented for different solar zenith angles and discussed in
19 association to changes in total ozone column (TOC), aerosol optical depth (AOD) and
20 cloudiness observed in the same period. Positive changes in annual mean anomalies of UV
21 irradiance, ranging from 2% to 6% per decade, have been detected both for clear- and all-sky
22 conditions. The changes are generally greater for larger solar zenith angles and for shorter
23 wavelengths. For clear skies, these changes are, in most cases, statistically significant at the
24 95% confidence limit. Decreases in the aerosol load and weakening of the attenuation by
25 clouds lead to increases in UV irradiance in the summer, of 7 – 9% per decade for 64° solar
26 zenith angle. The increasing TOC in winter counteracts the effect of decreasing AOD for this
27 particular season, leading to small, statistically insignificant, negative long-term changes in
28 irradiance at 307.5 nm. Annual mean UV irradiance levels are increasing from 1994 to 2006
29 and remain relatively stable thereafter, possibly due to the combined changes in the amount

1 and optical properties of aerosols. However, no statistically significant corresponding turning
2 point has been detected in the long-term changes of AOD. The absence of signatures of
3 changes in AOD in the short-term variability of irradiance in the UV-A may have been caused
4 by changes in the single scattering albedo of aerosols, which may counteract the effects of
5 changes in AOD on irradiance. The anti-correlation between the year-to-year variability of the
6 irradiance at 307.5nm and TOC is clear and becomes clearer as the AOD decreases.

7

8 **1 Introduction**

9 Although ultraviolet (UV) radiation is only a small fraction (<10%) of the total solar radiation
10 that reaches the earth's surface, it is vital for the life on earth (Asta et al., 2011; Häder et al.,
11 2015; Lucas et al., 2015; Madronich et al., 2015; UNEP, 2010; Williamson et al., 2014). The
12 amount of solar UV radiation reaching the atmosphere is changing periodically due to
13 changes in the earth-sun distance and the solar activity. Solar radiation with wavelengths
14 shorter than 290nm is entirely blocked by the atmosphere, while for longer wavelengths the
15 fraction that penetrates to the surface depends mainly on the solar zenith angle, the
16 composition of the atmosphere and the characteristics of the surface (Kerr and Fioletov,
17 2008). The interaction between the UV radiation, the atmospheric constituents and the
18 characteristics of the surface is complicated and not yet fully understood (Bernhard et al.,
19 2007; Kerr and Fioletov, 2008; Meinander et al., 2009). The geophysical parameters that
20 mainly affect the levels of the surface UV irradiance are: ozone, clouds, surface reflectivity
21 and aerosols (Arola et al., 2003; Bais et al., 1993; Bernhard et al., 2007; WMO, 2007).

22 From the early 1980s until the mid-1990s, the sharp decline of the stratospheric ozone over
23 the mid and high latitudes led to important increases in the surface UV irradiance (Kerr and
24 McElroy, 1993; Madronich et al., 1998). The successful implementation of the Montreal
25 protocol decelerated the weakening of the ozone layer (Egorova et al., 2013; Mäder et al.,
26 2010; Newman and McKenzie, 2011) and many recent studies indicate the first signs of
27 recovery over the northern hemisphere (Kuttippurath et al., 2013; McLinden and Fioletov,
28 2011; Newchurch et al., 2003; Smedley et al., 2012). Signs from the onset of ozone recovery
29 since the late 1990s on surface UV-B irradiance have been mainly detected over the northern
30 high latitudes; recent studies indicate that UV-B has been declining during the last two
31 decades (Bernhard, 2011; Eleftheratos et al., 2014). However, during the same period, both
32 the UV-B and the UV-A irradiance are increasing over many locations in the northern

1 hemisphere mid-latitudes, mainly due to the negative trends in the amount of clouds and
2 aerosols (De Bock et al., 2014; Fitzka et al., 2012; Román et al., 2014; Smedley et al., 2012;
3 Zerefos et al., 2012). In a recent study, Fragkos et al. (2015) showed that in Thessaloniki,
4 even under extreme high (low) TOC conditions, the erythemal irradiance can be lower
5 (higher) than its climatological values due to the dominant effect of aerosols. Zerefos et al.
6 (2012) suggest that over Canada, Europe and Japan there is a statistically significant evidence
7 of a slowdown or even a turning point in the upward UV-B trends after 2006, which is an
8 indication that since then the negative trends of aerosols are no more the main driver of the
9 long-term changes in UV-B irradiance.

10 In the next decades, ozone, aerosols, clouds and surface reflectivity are projected to undergo
11 important changes (IPCC, 2013). Changes in these factors may alter the levels of the surface
12 UV irradiance (Bais et al., 2015; Bais et al., 2011; Hegglin and Shepherd, 2009; Tourpali et
13 al., 2009; Watanabe et al., 2011) with important impacts on the human health and the balance
14 of the ecosystems (UNEP, 2010; Williamson et al., 2014). However, the uncertainties in the
15 spatial and temporal variability, the magnitude, and the direction of the projected changes of
16 surface UV irradiance are still high (Bais et al., 2015). Thus, good quality measurements of
17 the spectral UV irradiance and the main factors controlling its levels at the earth's surface are
18 of great importance for achieving better understanding and more accurate modelling of the
19 interactions among UV radiation, ozone, aerosols, clouds and surface reflectivity (García et
20 al., 2015; Kreuter et al., 2014; Mayer and Kylling, 2005; Schwander et al., 1997).

21 Accurate knowledge of the levels of spectral surface UV irradiance is necessary in order to
22 quantify effects on the health of humans (Kazantzidis et al., 2015; Webb et al., 2010) and
23 ecosystems (Ballare et al., 2011; Häder et al., 2011), and prevent potential impacts from over-
24 or under-exposure to UV radiation (Lucas et al., 2015). Additionally, reliable estimations of
25 the trends of spectral surface UV irradiance provide useful information for assessing these
26 impacts and for adopting proper measures (Morgenstern et al., 2008; Newman and McKenzie,
27 2011; van Dijk et al., 2013). Climatologies and trends of surface UV irradiance (spectral or
28 broadband) can be derived either directly from ground based measurements (Fitzka et al.,
29 2012; Glandorf et al., 2005; Zerefos, 2002), or indirectly from measurements of surface
30 reflectivity, ozone, aerosols and cloudiness derived either from satellites (Damiani et al.,
31 2014; Fioletov et al., 2004; Li et al., 2000), or from ground based instruments (Antón et al.,
32 2011; Román et al., 2014; Walker, 2009). The uncertainties of these parameters and the

1 applied methodologies increase the uncertainty of the indirectly derived UV irradiance, when
2 compared to measurements (Cordero et al., 2013; Weihs and Webb, 1997). Thus, long records
3 of good quality measurements of UV irradiance lead to more reliable estimations of its short-
4 term and long-term changes (Arola et al., 2003; Weatherhead et al., 1998).

5 The present study aims at the quantification of the long-term changes in surface UV
6 irradiance using spectral measurements which are recorded since 1990 at Thessaloniki (Bais
7 et al., 2001; Garane et al., 2006; Gröbner et al., 2006), one of the longest time series globally
8 (Glandorf et al., 2005). An important aspect is also the attribution of the trends and variability
9 of UV irradiance to changes in the total ozone column (TOC), the aerosol optical depth
10 (AOD) and cloudiness during the same period. Special emphasis is given to the reported
11 slowdown of the positive trends of UV irradiance (Zerefos et al., 2012) and their correlation
12 with the reported aerosol decline over the northern hemisphere (Turnock et al., 2015).

13

14 **2 Instrumentation and data**

15 In the 1980s, the increased concern for the stratospheric ozone depletion (Farman et al., 1985;
16 Solomon et al., 1986) and its effect on the levels of UV radiation at the Earth's surface (Kerr
17 and McElroy, 1993; Madronich et al., 1995; Zerefos, 2002), led to increased deployment of
18 ground-based instruments worldwide (Fioletov et al., 1999), to monitor the TOC and the
19 surface UV irradiance. Among these instruments, several Brewer spectrophotometers
20 (Brewer, 1973; Kerr et al., 1985) were deployed at different locations including Thessaloniki,
21 Greece, where the first commercially available single-monochromator Brewer with serial
22 number 005 (B005) was installed in 1982. Since then B005 performs continuous
23 measurements of the TOC and the columnar SO_2 (Bais et al., 1993; Bais et al., 1985; Meleti et
24 al., 2012; Zerefos, 1984). These measurements are also used to derive the aerosol optical
25 depth at specific UV-B wavelengths (Kazadzis et al., 2007). Monitoring of spectral UV
26 irradiance with B005 started in 1990 (Bais et al., 1996; Bais et al., 1993; Garane et al., 2006).
27 Since 1993, a second, double-monochromator, Brewer spectrophotometer with serial number
28 086 (B086) is also operating at Thessaloniki for continuous monitoring of the spectral UV
29 irradiance (Bais et al., 1996). Both instruments are located at the facilities of the Laboratory
30 of Atmospheric Physics (latitude 40.634° N, longitude 22.956° N, altitude 60 m above sea
31 level).

1 The spectral measurements of B005 cover the wavelength range 290-325nm in steps of 0.5
2 nm and spectral resolution of about 0.55 nm (FWHM). The corresponding spectral range for
3 B086 is 290-363nm, with the same wavelength step and very similar spectral resolution. The
4 UV dataset of both instruments was quality checked and re-evaluated up to the end of 2005
5 (Garane et al., 2006) and has been used in different studies (Kazadzis et al., 2009; Kazantzidis
6 et al., 2006; Kazantzidis et al., 2009; Meleti et al., 2009). Garane et al., (2006) presented a
7 comprehensive analysis of the uncertainties from all known sources that may affect the
8 calibration procedures and the spectral measurements of B005 and B086. In this study the
9 overall 1σ uncertainty of the measured irradiance was estimated to 5% for B086 and from
10 6.5% near 305nm to 5% near 320nm for B005. Recently, the quality control and the re-
11 evaluation of the post-2005 dataset have been completed and the time series is now extended
12 to the end of 2014, comprising about 170.000 spectra for B005 and 140.000 spectra for B086.

13 Direct spectral irradiance measurements performed with B005 at 306.3, 310.0, 313.5, 316.8
14 and 320.1 nm are used to derive the TOC (Kerr et al. 1981) and the AOD (Gröbner and
15 Meleti, 2004; Meleti and Cappellani, 2000). The uncertainty of the TOC measurements is
16 estimated to about 1% or less (Kerr et al., 1985), while for the AOD the uncertainty is of the
17 order of 0.04 at 320 nm for air mass 1.4 (Kazadzis et al., 2007). Comparisons with AOD data
18 for the period 2005 – 2014 provided from a collocated Cimel sun-photometer which is part of
19 AERONET (<http://aeronet.gsfc.nasa.gov/>) revealed an overall agreement to within 0.1 for air
20 mass values up to 3.2.

21 For the trend analysis, which will be discussed later, data for the 11-year solar cycle and the
22 Quasi-Biennial Oscillation (QBO) of the winds in the equatorial stratosphere have been used.
23 Monthly means for the solar flux at 10.7 cm were downloaded from the NOAA national
24 geophysical data centre (<http://www.ngdc.noaa.gov/ftp/space-weather/solar-data/solar-features/solar-radio/noontime-flux/penticton/>), while for the QBO wind data were
25 downloaded from the Freie Universität Berlin (<http://www.geo.fu-berlin.de/en/met/ag/strat/produkte/qbo/index.html>).
26

27 The spectral irradiances used in this study are averages of five measurements (within ± 1 nm
28 about the nominal wavelength) and the analysis is performed for 307.5, 324 and 350 nm. The
29 irradiance for the former two wavelengths is derived from both instruments while for 350 nm
30 it is derived only from B086. The solar irradiance at 307.5nm is strongly absorbed by ozone
31 in contrast to the other two wavelengths where the ozone absorption is either weak (324 nm)
32

1 or negligible (350 nm). In many studies, the irradiance at 305 nm has been used to estimate the
2 effect of TOC on UVB irradiance. However, for large solar zenith angles and/or under cloudy
3 conditions the effects from the dark signal and the stray light are very important for such low
4 wavelength, especially for the single monochromator Brewers (Karppinen et al., 2014).
5 Therefore, in the present study the irradiance at 307.5 nm has been chosen because it is
6 stronger than at 305 nm, with significantly higher signal-to-noise ratio required for more
7 accurate determination of trends, while the effect of ozone absorption remains very strong.
8 Additionally, the changes in irradiance for 307.5 nm are more representative of the changes in
9 the erythema irradiance, which is an important, human health-related metric. Finally, the
10 effect of SO₂ absorption in relation to the effect of ozone absorption at 307.5 nm is the
11 weakest in the range 306 - 309 nm. The changes in irradiance at 324 nm and 350 nm for clear
12 skies are mainly determined by the changes in the amount and optical properties of the
13 aerosols. Thus the effect of TOC on the long-term trends can be estimated by comparing the
14 trend in irradiance at 307.5 nm with the trend at these longer wavelengths. In order to detect
15 the presence of clouds during measurements and separate the data under clear skies, data from
16 a collocated pyranometer (type Kipp & Zonen CM-21) recorded at high temporal resolution
17 (once per minute) have been used (Bais et al., 2013).

18 The long-term variability of the spectral UV irradiance was investigated for specific solar
19 zenith angles, in order to minimize interferences from the different paths of radiation in the
20 atmosphere. Specifically, averages of measurements corresponding to SZAs within $\pm 1^\circ$ about
21 the nominal SZA were used. In order to eliminate remaining biases induced by these slightly
22 different SZAs, correction factors were derived with the radiative transfer model UVSPEC,
23 which is included in version 1.7 of the libRadtran package (Mayer and Kylling, 2005). The
24 simulations were made for a range of TOC and AOD values within the expected range of
25 variability over Thessaloniki: 250 – 550 DU for TOC and 0 – 1.4 for the AOD at 320 nm,
26 using the US standard atmospheric profile (Anderson et al., 1986), the aerosol profile
27 suggested by Shettle (1989), and typical values of the surface reflectivity, the single scattering
28 albedo and the asymmetry factor of 0.05, 0.85 and 0.7 respectively (Bais et al., 2005). The
29 simulations revealed that while for small SZAs and long wavelengths the differences in clear-
30 sky UV irradiance are small, at 305 nm the differences escalate to 60% for a change in SZA
31 from 69° to 71° . The following empirical relationship has been derived to correct the
32 measured irradiance at SZAs different than the nominal:

$$\frac{I_0}{I_\theta} = 1 + \alpha(\lambda, \theta_0) \cdot \left(\frac{1}{\cos \theta} - \frac{1}{\cos \theta_0} \right) \quad (1)$$

2 Where θ_0 is the nominal SZA, θ is the actual SZA, I_0 is the irradiance for θ_0 , I_θ is the
 3 irradiance for θ and $\alpha(\lambda, \theta_0)$ is the correction factor which depends on wavelength λ and θ_0 .
 4 After applying the correction, the differences between irradiances for SZAs which differ by
 5 up to 2° do not exceed 10% ($\pm 5\%$ about the mean) for wavelengths ranging from 290 to 400
 6 nm and SZAs from 15° to 80° . For the wavelengths above 310 nm and SZAs smaller than 70°
 7 the remaining discrepancies are generally below 2%, while for the same SZAs and
 8 wavelengths between 305 and 310 nm the remaining discrepancies range between 1% and
 9 5%. Thus for 307.5 nm, the remaining uncertainties due to differences from the nominal SZA
 10 range from 1 to 5% for SZAs between 30° and 70° . For the same range of SZAs the
 11 corresponding uncertainties are lower than 2% and 1% for 324 and 350 nm, respectively.

12 The monthly mean values of the irradiance at 307.5 nm and at 324 nm, the TOC, and the
 13 AOD derived from B005 since 1990 are presented in Figure 1. The period from June 1991
 14 until December 1993 has been shaded to highlight the low TOC values due to the Mt.
 15 Pinatubo volcanic eruption in June of 1991 (Hofmann et al., 1994; Randel et al., 1995). The
 16 annual cycle of the irradiance at 307.5 nm is clearly anti-correlated with the annual cycle of
 17 TOC while for the irradiance at 324 nm the annual variability is mainly caused by changes in
 18 cloudiness. There is an indication of increasing tendency in irradiance, for both wavelengths
 19 since the beginning of the record, which will be further discussed in the following sections.
 20 The monthly mean TOC generally ranges between about 280 and 400 DU with no obvious
 21 long-term trend. The high aerosol load over Thessaloniki is depicted in the monthly mean
 22 values of AOD that range between about 0.3 and 0.9. However, during the last two decades,
 23 the mean levels of the AOD are decreasing and its inter-annual variability becomes weaker.
 24 As will be shown in the following, changes in aerosols play a key role in both the short- and
 25 the long-term variability of the UV irradiance over Thessaloniki.

26

27 **3 Data Analysis and Results**

28 **3.1 Methodology**

29 The present study has two main objectives: First, to quantify and discuss the long- and short-
 30 term changes in UV irradiance, and second, to investigate whether a turning point exists in the

1 long-term variability of the UV time-series, during the period 1994-2014. Data before 1994
2 are not used in the analysis, to eliminate the effect on total ozone from the volcanic eruption
3 of Mt. Pinatubo in June 1991. Large amounts of aerosols, mainly sulfuric, were injected into
4 the stratosphere which led to decreases in TOC in the period 1991-1993, as can be seen from
5 Figure 1. In addition, measurements of spectral UV irradiance at Thessaloniki before June
6 1991 are sparse and available for less than two years.

7 Assuming that part of the long-term variability of the clear-sky UV irradiance is due to ozone,
8 one would expect that there should be two modes in this trend: one for the period of
9 decreasing total ozone and one for the period of stabilized or increasing total ozone. For some
10 northern hemisphere locations, it was found (Zerefos et al., 2012) that surface UV-B
11 irradiance increases faster in the period before 2006. As shown in the following such a turning
12 point exists also in the 20-year long UV data-series of Thessaloniki. Although quantitative
13 estimates of the trends for each of the two sub-periods have been derived they are discussed
14 briefly and in a more qualitative way, since the two time periods are too short to consider the
15 quantitative estimates of the trends reliable (Arola et al., 2003; Weatherhead et al., 1998).
16 Thus, in the following analysis long-term trends are comprehensively discussed only for the
17 entire period 1994-2014.

18 With respect to the presence of autocorrelation in the time series of TOC and UV and its
19 effects on trend analysis, Weatherhead et al. (1998) have reported that the deseasonalized
20 monthly mean UV irradiance data are autocorrelated in the first order. For the station of
21 Thessaloniki, the autocorrelation of the all-sky dataset is estimated to be small, less than 0.2.
22 In contrast, for the clear-sky dataset the autocorrelation is larger, ranging between 0.3 and 0.5
23 for different SZAs, and may affect the significance of the derived trends. In our analysis the
24 autocorrelation has not been removed either from the all-sky or the clear-sky datasets. As
25 discussed in Yang et al., (2006) removing the autocorrelation from time series with large
26 number of gaps, as in our case for the clear-sky UV data set, can induce artificial tendencies
27 and biases in the derived trends. In order to accurately detect the turning point in the trends of
28 the UV irradiance and to determine its statistical significance, the methodology of Yang et al.,
29 (2006) was applied on the monthly mean anomalies. This study aimed at detecting a turning
30 point in a time series of TOC with a large number of gaps using a non-autoregressive model.
31 The additional uncertainty due to the remaining autocorrelation was taken into account in the
32 estimation of the statistical significance associated with the detection of the turning point.

1 For the analysis of long-term changes we calculated daily anomalies for TOC, AOD and
2 spectral UV irradiance at different SZAs in order to remove the effect of SZA in the annual
3 variability of irradiance. These daily anomalies were calculated by subtracting from each data
4 point the climatological value for that day which was derived from the entire dataset. Monthly
5 mean anomalies were calculated by averaging the daily anomalies for months with at least 10
6 days of available data. As a next step, the effects of QBO and the 11-year solar cycle were
7 filtered from both the TOC and UV irradiance data sets, by applying a multilinear regression
8 analysis. The procedure described in, e.g., Zerefos et al., (2012) was followed with the only
9 difference being that in the present study we did not use an autoregressive model, due to gaps
10 in the clear-sky UV data set. It was found that the difference in the linear trends derived for
11 the period 1994 – 2014 with and without filtering the effects of these two natural cycles is
12 generally smaller than 0.5% per decade, thus smaller than the 1σ uncertainty of the trends
13 which ranges from about 1 to 3% per decade. However, it is not always negligible compared
14 to the magnitude of the derived trends which for all cases range between about -5 and 10%
15 per decade.

16 The number of gaps in the time-series of the UV irradiance is higher in the first half compared
17 to the second half of the period of study. In order to suppress the effect of the uneven
18 distribution of the measurements, the long-term changes in UV irradiance were calculated
19 from the yearly mean anomalies, instead of those derived from the multilinear model. After
20 removing the effects of the solar cycle and the QBO from the monthly mean anomalies, the
21 dataset was recomposed and the yearly mean anomalies were calculated. It was found again
22 that the difference in the linear trends derived from the yearly mean anomalies and directly
23 from the multilinear model is small, with the former being higher by only 0.1 – 0.4% per
24 decade. The statistical significance of the trends is derived from the Mann-Kendall test
25 (Burkey, 2006). In the following, a trend is considered significant when it is statistically
26 significant at the 95% confidence level.

27 **3.2 Comparison between the trends from the two Brewers for different SZAs**

28 Following the re-evaluation and quality control of the entire dataset, the trends of the UV
29 irradiance were calculated separately for the two Brewers operating at Thessaloniki. Mean
30 ratios between quasi-synchronous (within ± 1 min) spectral UV irradiance measurements from
31 B005 and B086 under all-sky conditions were calculated for different SZA's to evaluate the
32 applied corrections. Instrumental characteristics of Brewers (e.g., the slit function) may differ

1 for different instruments (e.g. Lakkala et al., 2008) leading to differences between, even
2 synchronous, single wavelength measurements. Comparing averages over small spectral
3 intervals suppress partly these effects. In the present study averages for 5 nm spectral
4 intervals were compared instead of irradiance measurements at single wavelengths (or
5 averages for narrower spectral intervals), because they are wide enough to suppress a great
6 part of the effect of these characteristics, while at the same time they are narrow enough to
7 assess if measurements are properly corrected for, e.g., the effects of temperature and SZA
8 with respect to wavelength. For the wavelength range 310 – 325 nm the ratios are very close
9 to 1 with a standard deviation of about 5%. For shorter wavelengths, the mean ratio gradually
10 decreases and for the 300 - 305nm range it is ~0.96 with a standard deviation of about 10%.
11 As already discussed, the uncertainties and the deviations from unity arise from the different
12 characteristics of the two instruments and from the imperfect synchronization of the
13 measurements (Garane et al., 2006). No dependency of the ratio from the temperature or the
14 solar zenith angle was found. The good agreement in the absolute levels of the measured
15 irradiance by the two instruments is an indication for the quality of the re-evaluated data. It
16 should be noted, however, that the synchronous measurements represent only ~50% of the
17 available data. The trends from both instruments for the entire period 1994-2014 were
18 compared for different solar zenith angles from 30° to 70° in steps of 10°. The results for
19 307.5 nm and 324 nm are presented in Figure 2 both for clear-sky and all-sky conditions.
20 For 307.5 nm, statistically significant trends were found for clear-skies for both instruments
21 and all SZAs, and for all-skies in B005 for 30° and 70° SZA. For 324 nm, only the clear-sky
22 trends from B005 and for 30° and 40° SZA are statistically significant. The trends for 324 nm
23 are generally smaller than those for 307.5 nm. The results are quite satisfactory and consistent
24 since for the same wavelengths and SZAs the irradiance trends from B005 and B086 do not
25 differ by more than 2% per decade and in most cases they agree within 1σ . The derived trends
26 both for clear- and all-sky data and for all SZAs are positive and range between 1% and 6%
27 per decade. Although the dependence of the trends on the SZA appears to be small and within
28 the uncertainty limits, at large SZAs the trends are greater. This dependence can be partially
29 attributed to the increasing optical path of radiation with SZA, which leads to stronger
30 absorption from ozone or aerosols. However, for different SZAs, the datasets comprise data
31 from different periods in the year (e.g. for SZA=30° data exist only from April to August,
32 while for SZA=70° data are available during the entire year). Since the long-term changes of

1 TOC, AOD and cloudiness are different for different seasons, the irradiance trends for
2 different SZAs should be affected differently by these factors.

3 **3.3 Seasonal trends**

4 Since the results from both instruments are generally similar, only the data from B086 are
5 used in the following, since this instrument has superior characteristics, at least with respect to
6 the rejection of stray light and angular response. Seasonal trends of the spectral UV irradiance
7 for 307.5, 324 and 350 nm (Figure 3) were calculated and compared to the corresponding
8 trends of the daily mean TOC and AOD (Figure 4). The trends in AOD are statistically
9 significant for all cases presented, in contrast to the trends in TOC which are not statistically
10 significant. For SZAs larger than 63° data are available during the whole year, thus, the
11 irradiance for 64° SZA (data ranging from 63° to 65°) is used in the analysis of trends. The
12 effect of the changes in cloudiness is assessed by comparing the trends of the clear-sky and
13 the all-sky irradiance.

14 As expected, the seasonal trends for 324 and the 350 nm are similar for both, clear-sky and
15 all-sky conditions. The changes of the solar irradiance at these wavelengths are practically
16 unaffected by the changes of TOC, while they are mainly affected by the changes in aerosols
17 and clouds. In general, the effects of changes in aerosol amount and/or properties on UV
18 irradiance are stronger for shorter wavelengths. Thus, the important negative trends of the
19 AOD at 320 nm that have been observed for Thessaloniki lead to slightly less positive trends
20 for the irradiance at 350 nm than at 324 nm. It must be clarified at this point that the
21 interaction of solar UV radiation with aerosols is very complex and the changes in the AOD
22 cannot explain the changes in UV irradiance without taking into account the absorption
23 efficiency of the aerosols (i.e., the single scattering albedo) for which no measurements are
24 available for this period. For example, decreases in the single scattering albedo (greater
25 absorption efficiency) counteract the effect of decreases in the AOD. As will be discussed
26 later, the fact that the changes in clear-sky UV-A irradiance (324 and 350 nm) cannot be fully
27 explained by the changes in the AOD is an indication that changes in other optical properties
28 of aerosols, such as the single scattering albedo may have occurred.

29 The greatest changes in irradiance at 324 and 350 nm were found in summer both for clear-
30 sky and all-sky conditions. The trend for clear skies at these wavelengths is about 3.5% per
31 decade, while for 307.5 nm it increases to about 5% per decade. The main driver for the

1 changes under clear skies appears to be the decreasing AOD, which for summer is more than
2 20% per decade. For all skies, the positive trends are almost double than those for clear skies
3 (about 7% for 324 and 350 nm and about 9% for 307.5 nm), suggesting that the attenuation of
4 irradiance by clouds is decreasing during the last two decades. All these trends are statistically
5 significant. For winter, the trends in irradiance for 324 and 350 nm are 3.5% and 3.0%
6 respectively both for clear skies and all-skies, suggesting that cloud effects during the last two
7 decades are very small in winter and changes in aerosols are the dominant factor. This
8 conclusion is confirmed by the negative trend of the AOD shown in Figure 4. For 307.5nm
9 the increases in TOC counteract the effects of changing aerosols, leading to a negative trend
10 of about -3% per decade for irradiance under clear skies. However, none of the trends in
11 winter is statistically significant.

12 For spring, the trends for clear skies are similar to those in summer for all three wavelengths,
13 while for all skies trends are smaller; by 0.5 - 1%. Thus, as for winter, the UV trends are due
14 mainly to decreasing AOD. Although for that season the trend in TOC is about 1% per
15 decade, this is not reflected in the trend of clear-sky irradiance at 307.5 nm which is slightly
16 larger than in the UV-A wavelengths, instead of being smaller. For this season only the trend
17 for 350 nm is statistically significant.

18 For autumn, the trends in clear-sky irradiance are approximately 7%, 3% and 1.5% for 307.5,
19 324 and 350 nm respectively, and statistically significant only for the first two wavelengths.
20 For all skies, the trends are 3-4% lower, suggesting an increasing attenuation by clouds during
21 this season. However, the differences between the clear sky and the all sky trends are within
22 the uncertainty limits of the later. The all-sky trends for autumn are not statistically
23 significant. One of the possible reasons for the stronger increase of the irradiance at 307.5 nm
24 compared to 324 and 350 nm is the small negative trend in TOC. Additionally, the relatively
25 large difference between the trends for 324 and 350 nm is explained by the decreasing
26 aerosols which have much stronger impact on shorter than on longer wavelengths.

27 Finally, the yearly averaged TOC is slightly increasing, by about 0.8% per decade, but this
28 change is not statistically significant. In contrast, the yearly mean AOD has been decreasing
29 by about 17% per decade, and therefore AOD is the dominant driver of the changes in the
30 yearly mean UV irradiance. The trends in UV irradiance range from 3 to 4% for clear skies,
31 while for all skies they are about 0.5% larger. At shorter wavelengths the trends are larger,

1 possibly due to the negative trend in TOC and the stronger effect of aerosols on the irradiance
2 at these wavelengths.

3 The results presented in Figure 3 lead to the conclusion that the enhanced attenuation of UV
4 radiation by clouds in summer is balanced by the decreased attenuation in autumn, leading to
5 a negligible effect on the yearly mean UV irradiance. However, as in all cases presented, the
6 differences between the trends in clear-sky and all-sky irradiance for summer and autumn are
7 similar to (or even lower than) their 1σ uncertainty, which is a strong indication that the
8 estimated changes in the attenuation of the UV irradiance by clouds are not significant.

9 **3.4 The role of ozone and aerosols on short- and long-term variability of
10 irradiance**

11 In the following we discuss in more detail the short- and long-term variability of clear-sky
12 UV irradiance at 307.5 and 350 nm in association with the evolution of factors causing this
13 variability. The analysis of the variability is performed on annual mean anomalies of UV
14 irradiance at 64° SZA, TOC and AOD, as well as for mean anomalies for the periods
15 December – May (winter - spring) and June – November (summer - autumn); the former
16 being affected mainly by changes in ozone, while the latter by changes in aerosols.
17 Furthermore we explore a potential turning point in the time series of irradiance at
18 Thessaloniki using monthly mean anomalies, in an attempt to confirm the findings of Zerefos
19 et al., (2012).

20 The results for 324 nm are not discussed since they are similar to those for 350 nm. For
21 monthly mean anomalies for the entire year, a turning point in the upward trend of irradiance
22 at both 307.5 and 350 nm has been detected in 2006, statistically significant at the 95%
23 confidence level. Since the same pattern occurs at both wavelengths and since no statistically
24 significant turning point has been detected for TOC, this piece-wise trend pattern in UV
25 irradiance has likely been caused by changes in aerosols. However, the small negative trend in
26 UV irradiance observed after 2006 (see Table 1 and Figure 5) does not comply with the
27 negative monotonic trend in AOD during the whole period of study which continues also after
28 2006. The behaviour of aerosols after 2006 has been verified by an independent dataset from
29 a collocated Cimel sun-photometer, which revealed a decreasing trend of about 0.1 per decade
30 in AOD at 440 nm from 2006 to 2014, similar to that of B005. Thus, the only factor that could
31 explain the small negative trend in UV irradiance during this period would be a negative trend

1 in the single scattering albedo (Bais et al., 2005; Nikitidou et al., 2013). This assumption
2 cannot be easily verified since the SSA data from the Cimel and from satellite overpasses
3 (e.g. <http://disc.sci.gsfc.nasa.gov/giovanni>) are sparse and inadequate to derive reliable trends.
4 However, simulations with UVSPEC revealed that for SZAs greater than 60° and for typical
5 aerosol properties and atmospheric conditions for Thessaloniki, the effect of a decrease in
6 AOD at 320 nm by 0.1 can be reversed by a simultaneous decrease in SSA by less than 0.1.
7 As shown in Table 1, the trends for 350 nm for winter-spring, summer-autumn and the entire
8 year are similar. For all the three cases the UV irradiance increases by about 10% from 1994
9 to 2006 and then it slightly decreases from 2006 to 2014 resulting to a mean rate of increase
10 of about 3.5% per decade for the entire period 1994 – 2014. For the three cases the mean rate
11 of decrease for the AOD is similar before and after 2006. Additionally, the year to year
12 variability of the mean anomalies for the AOD is not clearly anti-correlated with the year to
13 year variability of the mean anomalies for the UV irradiance at 350 nm, which can be only
14 attributed to changes in SSA. SSA may differ importantly for different types of aerosols
15 (Takemura et al., 2002). The aerosol mixture over Thessaloniki consists of several different
16 types of aerosols (e.g., urban, continental, marine, dust) and its composition varies (Amiridis
17 et al., 2005; Koukouli et al., 2006). This could lead to large variability of the SSA, even
18 within the same day (e.g., Ialongo et al., (2010)). An increase of the mean SSA in 1999
19 would, for example, explain why the very high annual mean levels of AOD in the specific
20 year are not depicted in the levels of UV irradiance.
21 The changes of the UV irradiance at 307.5 nm are highly affected by changes in TOC and
22 aerosols. For the winter-spring period no statistically significant turning point has been
23 detected in the trend for this wavelength. Additionally, the mean trend in irradiance for the
24 period 1994 – 2014 is weak (Table 1) compared to the corresponding trend for the period June
25 – November, and is likely caused by the combined, but opposing, effects of a statistically
26 significant positive trend in TOC and a negative trend in AOD. For the period June –
27 November no trend was detected in TOC, thus, as for 350 nm, the UV irradiance at 307.5 nm
28 increases steadily from 1994 to 2006 due to decreasing AOD and after 2006 remains
29 unchanged. A pattern which is similar to that for the period June – November appears also in
30 the annual means, with changes in irradiance dominated again by changes in aerosols of
31 opposite sign.

1 There are some interesting conclusions emerging from Figure 5: By comparing Figures 5(a) –
2 (c) with Figures 5(g) – (i), one can notice an anti-correlation between the year to year
3 variability of TOC and the year to year variability of UV irradiance at 307.5 nm, which
4 becomes stronger as the AOD decreases. For example, the low yearly mean TOC in 2000,
5 2008 and 2011, compared in each case with the yearly mean TOC for the nearest (e.g. 4 or 5)
6 years, coincides with high UV irradiance at 307.5 nm while correspondingly the high TOC in
7 1998, 2010 and 2013 coincides with low UV irradiance at 307.5 nm. Obviously, while the
8 year to year variability in irradiance at 307.5 nm is mainly driven by the changes in TOC, its
9 long-term changes are mainly driven by the changes in aerosols. For example, the yearly
10 mean TOC in 2010 is the highest that has been recorded during the entire period 1994 – 2014
11 (Steinbrecht et al., 2011) and has led to low yearly mean irradiance at 307.5 nm. However, the
12 yearly mean irradiance at 307.5 nm in 2010 is still higher than mean levels in the period 1994
13 – 1998, mainly due to the very high levels of aerosols in the atmosphere in the mid-1990s. As
14 the AOD decreases throughout the years, the anti-correlation between the short-term
15 variability of the TOC and the UV irradiance becomes clearer. Finally, it is noteworthy that
16 while the mean value of AOD for 2014 in the period summer – autumn is the lowest recorded
17 since 1994, the corresponding value for the period winter – spring is the highest of the last
18 seven years. These very high AOD values are probably due to the increased biomass-burning
19 aerosols arising from a shift in the type of fuel owing to the economic crisis in Greece after
20 2009 (Saffari et al., 2013). As a consequence of the increased aerosols, the levels of irradiance
21 at 350 nm for winter – spring 2014 are the lowest recorded during the last decade.

22 Since this paragraph aimed at attributing the short- and long-term variability of the UV
23 irradiance to the corresponding variability of TOC and AOD, the analysis was restricted to
24 clear-sky data. Although not shown here, a similar analysis has been performed for the
25 irradiance under all-sky conditions and a statistically significant turning point in 2006 has also
26 been detected in the trends of yearly mean irradiance for 307.5 and 350 nm. As already
27 discussed, changes in cloudiness do not have an important impact on the long-term changes of
28 the UV irradiance at Thessaloniki but are the main driver of the short-term variations in the
29 all-sky dataset.

30

4 Summary and conclusions

In the present study, spectral UV irradiance measurements from 1994 to 2014 at Thessaloniki, Greece have been used to investigate the short- and long-term variability of UV irradiance at specific wavelengths, affected differently by total ozone, aerosols and clouds. Although data are available since 1990 the analysis was restricted to the period 1994 – 2014 to avoid interferences in the trends from the volcanic aerosols injected into the stratosphere by the eruption of Mt. Pinatubo in 1991. Additionally, the parallel measurements from two co-located Brewer spectrophotometers after 1993 increase the confidence in the accuracy of the spectral measurements. Trends of clear-sky and all-sky UV irradiance at 307.5 and 324 nm were derived for the period 1994 – 2014 from both B005 and B086 for SZAs from 30° to 70°. The difference between the trends from the two instruments was found to be smaller than their 1σ uncertainty boundaries.

According to the results, the annual mean UV irradiance has increased during the last two decades. The increasing trends are similar for both clear-sky and all-sky data and are higher at shorter wavelengths and higher SZAs. The calculated trends range between 2% and 6% per decade, and for clear skies are statistically significant for most SZAs. For all skies, most of the irradiance trends are not statistically significant.

The impact of changes in TOC, aerosols and clouds on the changes in UV irradiance is different for different seasons. The negative trends in AOD, which are stronger in summer, lead to positive trends in UV irradiance at longer wavelengths (e.g., at 324 and 350 nm). For shorter wavelengths changes in TOC are also important. Thus, the effect of the small negative trend in AOD in winter is fully counteracted by the positive trend in TOC, resulting in a decreasing trend in clear-sky irradiance at 307.5 nm. Changes in clouds have a negligible effect on the trend of irradiance for winter and spring. The enhancement of the attenuation of irradiance by clouds in autumn is balanced by the reduced attenuation in summer, leading to similar changes in the annual means of clear-sky and all-sky irradiance. It is important to notice that the strongest changes in UV irradiance were found for summer when humans are more exposed to the Sun compared to the other seasons.

Moreover, it is shown that the period 1994 – 2014 can be divided in two sub-periods: during the first period (1994 - 2006) the annual mean UV irradiance is increasing fast while during the second period (2006 - 2014) the UV irradiance is relatively stable at 307.5 nm and is slightly decreasing at 350 nm. The long-term variability of UV irradiance for both short and

1 long wavelengths is mainly driven by the changes in aerosols. The short-term variability of
2 the clear-sky irradiance at 307.5 nm is mainly driven by the short-term variability of TOC.
3 The effect of the TOC changes on the year to year variability of UV irradiance becomes
4 clearer when AOD decreases. The short-term changes in irradiance at 350 nm cannot be fully
5 explained by the short-term changes in AOD, as the absorption efficiency of aerosols may
6 also change with time.

7

1 **References**

2 Amiridis, V., Balis, D. S., Kazadzis, S., Bais, A., Giannakaki, E., Papayannis, A., and
3 Zerefos, C.: Four-year aerosol observations with a Raman lidar at Thessaloniki, Greece, in the
4 framework of European Aerosol Research Lidar Network (EARLINET), *Journal of*
5 *Geophysical Research: Atmospheres*, 110, D21, doi:10.1029/2005jd006190, 2005.

6 Anderson, G. P., Clough, S., Kneizys, F., Chetwynd, J., and Shettle, E. P.: AFGL atmospheric
7 constituent profiles (0-120 km), *Tech. Rep. TR-86-0110*, AFGL, DTIC Document, 1986.

8 Antón, M., Serrano, A., Cancillo, M. L., GarcÍA, J. A., and Madronich, S.: Application of an
9 analytical formula for UV Index reconstructions for two locations in Southwestern Spain,
10 *Tellus B*, 63, 1052-1058, 2011.

11 Arola, A., Lakkala, K., Bais, A., Kaurola, J., Meleti, C., and Taalas, P.: Factors affecting
12 short- and long-term changes of spectral UV irradiance at two European stations, *J. Geophys.*
13 *Res.-Atmos.*, 108, 4549, doi:10.1029/2003jd003447, 2003.

14 Asta, J., Pål, B., Arne, D., Stefan, A.-E., Jörg, R., Kristin, M., Michael, F. H., William, B. G.,
15 and Johan, M.: Solar radiation and human health, *Reports on Progress in Physics*, 74, 066701,
16 doi:10.1088/0034-4885/74/6/066701, 2011.

17 Bais, A. F., Drosoglou, T., Meleti, C., Tourpali, K., and Kouremeti, N.: Changes in surface
18 shortwave solar irradiance from 1993 to 2011 at Thessaloniki (Greece), *International Journal*
19 *of Climatology*, 33, 2871-2876, 2013.

20 Bais, A. F., Gardiner, B. G., Slaper, H., Blumthaler, M., Bernhard, G., McKenzie, R., Webb,
21 A. R., Seckmeyer, G., Kjeldstad, B., Koskela, T., Kirsch, P. J., Gröbner, J., Kerr, J. B.,
22 Kazadzis, S., Leszczynski, K., Wardle, D., Josefsson, W., Brogniez, C., Gillotay, D., Reinen,
23 H., Weihs, P., Svenoe, T., Eriksen, P., Kuik, F., and Redondas, A.: SUSPEN intercomparison
24 of ultraviolet spectroradiometers, *Journal of Geophysical Research: Atmospheres*, 106,
25 12509-12525, 2001.

26 Bais, A. F., Kazantzidis, A., Kazadzis, S., Balis, D. S., Zerefos, C. S., and Meleti, C.:
27 Deriving an effective aerosol single scattering albedo from spectral surface UV irradiance
28 measurements, *Atmospheric Environment*, 39, 1093-1102, 2005.

1 Bais, A. F., McKenzie, R. L., Bernhard, G., Aucamp, P. J., Ilyas, M., Madronich, S., and
2 Tourpali, K.: Ozone depletion and climate change: impacts on UV radiation, Photochemical
3 & Photobiological Sciences, doi: 10.1039/c4pp90032d, 2015. 2015.

4 Bais, A. F., Tourpali, K., Kazantzidis, A., Akiyoshi, H., Bekki, S., Braesicke, P.,
5 Chipperfield, M. P., Dameris, M., Eyring, V., Garny, H., Iachetti, D., Jöckel, P., Kubin, A.,
6 Langematz, U., Mancini, E., Michou, M., Morgenstern, O., Nakamura, T., Newman, P. A.,
7 Pitari, G., Plummer, D. A., Rozanov, E., Shepherd, T. G., Shibata, K., Tian, W., and
8 Yamashita, Y.: Projections of UV radiation changes in the 21st century: impact of ozone
9 recovery and cloud effects, *Atmos. Chem. Phys.*, 11, 7533-7545, 2011.

10 Bais, A. F., Zerefos, C. S., and McElroy, C. T.: Solar UVB measurements with the double-
11 and single-monochromator Brewer ozone spectrophotometers, *Geophysical Research Letters*,
12 23, 833-836, 1996.

13 Bais, A. F., Zerefos, C. S., Meleti, C., Ziomas, I. C., and Tourpali, K.: Spectral measurements
14 of solar UVB radiation and its relations to total ozone, SO₂, and clouds, *Journal of*
15 *Geophysical Research: Atmospheres*, 98, 5199-5204, 1993.

16 Bais, A. F., Zerefos, C. S., Ziomas, I. C., Zoumakis, N., Mantis, H. T., Hofmann, D. J., and
17 Fiocco, G.: Decreases in the Ozone and the S02 Columns Following the Appearance of the El
18 Chichon Aerosol Cloud at Midlatitude. In: *Atmospheric Ozone*, Zerefos, C. S. and Ghazi, A.
19 (Eds.), Springer Netherlands, 1985.

20 Ballare, C. L., Caldwell, M. M., Flint, S. D., Robinson, S. A., and Bornman, J. F.: Effects of
21 solar ultraviolet radiation on terrestrial ecosystems. Patterns, mechanisms, and interactions
22 with climate change, *Photochemical & Photobiological Sciences*, 10, 226-241, 2011.

23 Bernhard, G.: Trends of solar ultraviolet irradiance at Barrow, Alaska, and the effect of
24 measurement uncertainties on trend detection, *Atmos. Chem. Phys.*, 11, 13029-13045, 2011.

25 Bernhard, G., Booth, C. R., Ehramjian, J. C., Stone, R., and Dutton, E. G.: Ultraviolet and
26 visible radiation at Barrow, Alaska: Climatology and influencing factors on the basis of
27 version 2 National Science Foundation network data, *J. Geophys. Res.-Atmos.*, 112, D09101,
28 doi:10.1029/2006jd007865, 2007.

29 Brewer, A. W.: A replacement for the Dobson spectrophotometer?, *PAGEOPH*, 106-108,
30 919-927, 1973.

1 Burkey, J.: A non-parametric monotonic trend test computing Mann-Kendall Tau, Tau-b, and
2 Sens Slope written in Mathworks-MATLAB implemented using matrix rotations., King
3 County, Department of Natural Resources and Parks, Science and Technical Services section.
4 Seattle, Washington. USA., 2006.

5 Cordero, R. R., Seckmeyer, G., Damiani, A., Labbe, F., and Laroze, D.: Monte Carlo-based
6 uncertainties of surface UV estimates from models and from spectroradiometers, *Metrologia*,
7 50, L1, 2013.

8 Damiani, A., Cordero, R. R., Cabrera, S., Laurenza, M., and Rafanelli, C.: Cloud cover and
9 UV index estimates in Chile from satellite-derived and ground-based data, *Atmospheric*
10 *Research*, 138, 139-151, 2014.

11 De Bock, V., De Backer, H., Van Malderen, R., Mangold, A., and Delcloo, A.: Relations
12 between erythema UV dose, global solar radiation, total ozone column and aerosol optical
13 depth at Uccle, Belgium, *Atmos. Chem. Phys.*, 14, 12251-12270, 2014.

14 Egorova, T., Rozanov, E., Gröbner, J., Hauser, M., and Schmutz, W.: Montreal Protocol
15 Benefits simulated with CCM SOCOL, *Atmos. Chem. Phys.*, 13, 3811-3823, 2013.

16 Eleftheratos, K., Kazadzis, S., Zerefos, C. S., Tourpali, K., Meleti, C., Balis, D., Zyrichidou,
17 I., Lakkala, K., Feister, U., Koskela, T., Heikkilä, A., and Karhu, J. M.: Ozone and
18 Spectroradiometric UV Changes in the Past 20 Years over High Latitudes, *Atmosphere-*
19 *Ocean*, doi: 10.1080/07055900.2014.919897, 2014. 1-9, 2014.

20 Farman, J. C., Gardiner, B. G., and Shanklin, J. D.: Large losses of total ozone in Antarctica
21 reveal seasonal ClO_x/NO_x interaction, *Nature*, 315, 207-210, 1985.

22 Fioletov, V. E., Kerr, J. B., Hare, E. W., Labow, G. J., and McPeters, R. D.: An assessment of
23 the world ground-based total ozone network performance from the comparison with satellite
24 data, *Journal of Geophysical Research: Atmospheres*, 104, 1737-1747, 1999.

25 Fioletov, V. E., Kimlin, M. G., Krotkov, N., McArthur, L. J. B., Kerr, J. B., Wardle, D. I.,
26 Herman, J. R., Meltzer, R., Mathews, T. W., and Kaurola, J.: UV index climatology over the
27 United States and Canada from ground-based and satellite estimates, *J. Geophys. Res.-*
28 *Atmos.*, 109, D22308, doi:10.1029/2004jd004820, 2004.

29 Fitzka, M., Simic, S., and Hadzimustafic, J.: Trends in spectral UV radiation from long-term
30 measurements at Hoher Sonnblick, Austria, *Theor. Appl. Climatol.*, 110, 585-593, 2012.

1 Garane, K., Bais, A. F., Kazadzis, S., Kazantzidis, A., and Meleti, C.: Monitoring of UV
2 spectral irradiance at Thessaloniki (1990–2005): data re-evaluation and quality control, Ann.
3 Geophys., 24, 3215–3228, 2006.

4 García, R. D., Cachorro, V. E., Cuevas, E., Toledano, C., Redondas, A., Blumthaler, M., and
5 Benounna, Y.: Comparison of measured and modelled spectral UV irradiance at Izaña high
6 mountain station: estimation of the underlying effective albedo, International Journal of
7 Climatology, doi: 10.1002/joc.4355, 2015. n/a-n/a, 2015.

8 Glandorf, M., Arola, A., Bais, A., and Seckmeyer, G.: Possibilities to detect trends in spectral
9 UV irradiance, Theor. Appl. Climatol., 81, 33-44, 2005.

10 Gröbner, J., Blumthaler, M., Kazadzis, S., Bais, A., Webb, A., Schreder, J., Seckmeyer, G.,
11 and Rembges, D.: Quality assurance of spectral solar UV measurements: results from 25 UV
12 monitoring sites in Europe, 2002 to 2004, Metrologia, 43, S66–S71, doi:10.1088/0026- 20
13 1394/43/2/S14, 2006.

14 Gröbner, J. and Meleti, C.: Aerosol optical depth in the UVB and visible wavelength range
15 from Brewer spectrophotometer direct irradiance measurements: 1991–2002, J. Geophys.
16 Res.- Atmos., 109, D09202, doi:10.1029/2003jd004409, 2004.

17 Häder, D.-P., Williamson, C. E., Wangberg, S.-A., Rautio, M., Rose, K. C., Gao, K.,
18 Helbling, E. W., Sinha, R. P., and Worrest, R.: Effects of UV radiation on aquatic ecosystems
19 and interactions with other environmental factors, Photochemical & Photobiological Sciences,
20 doi: 10.1039/c4pp90035a, 2015. 2015.

21 Häder, D. P., Helbling, E. W., Williamson, C. E., and Worrest, R. C.: Effects of UV radiation
22 on aquatic ecosystems and interactions with climate change, Photochemical &
23 Photobiological Sciences, 10, 242-260, 2011.

24 Hegglin, M. I. and Shepherd, T. G.: Large climate-induced changes in ultraviolet index and
25 stratosphere-to-troposphere ozone flux, Nature Geoscience, 2, 687-691, 2009.

26 Hofmann, D. J., Oltmans, S. J., Komhyr, W. D., Harris, J. M., Lathrop, J. A., Langford, A. O.,
27 Deshler, T., Johnson, B. J., Torres, A., and Matthews, W. A.: ozone loss in the lower
28 stratosphere over the United States in 1992–1993: Evidence for heterogeneous chemistry on
29 the Pinatubo aerosol, Geophysical Research Letters, 21, 65-68, 1994.

1 Ialongo, I., Buchard, V., Brogniez, C., Casale, G. R., and Siani, A. M.: Aerosol Single
2 Scattering Albedo retrieval in the UV range: an application to OMI satellite validation,
3 Atmos. Chem. Phys., 10, 331-340, 2010.

4 IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I
5 to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change Cambridge
6 University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp., 2013.

7 Karppinen, T., Redondas, A., García, R. D., Lakkala, K., McElroy, C. T., and Kyrö, E.:
8 Compensating for the Effects of Stray Light in Single-Monochromator Brewer
9 Spectrophotometer Ozone Retrieval, Atmosphere-Ocean, doi:
10 10.1080/07055900.2013.871499, 2014. 1-8, 2014.

11 Kazadzis, S., Bais, A., Amiridis, V., Balis, D., Meleti, C., Kouremeti, N., Zerefos, C. S.,
12 Rapsomanikis, S., Petrakakis, M., Kelesis, A., Tzoumaka, P., and Kelektsgolou, K.: Nine
13 years of UV aerosol optical depth measurements at Thessaloniki, Greece, Atmos. Chem.
14 Phys., 7, 2091-2101, 2007.

15 Kazadzis, S., Bais, A., Arola, A., Krotkov, N., Kouremeti, N., and Meleti, C.: Ozone
16 Monitoring Instrument spectral UV irradiance products: comparison with ground based
17 measurements at an urban environment, Atmos. Chem. Phys., 9, 585-594, 2009.

18 Kazantzidis, A., Bais, A., Garane, K., Kazadzis, S., and Meleti, C.: Estimation of UV
19 irradiance from ancillary data and comparison with measurements at Thessaloniki, Greece
20 (40.5° N, 23° E), 6362, doi:10.1111/12.689813, 2006.

21 Kazantzidis, A., Bais, A. F., Zempila, M. M., Kazadzis, S., den Outer, P. N., Koskela, T., and
22 Slaper, H.: Calculations of the human vitamin D exposure from UV spectral measurements at
23 three European stations, Photochemical & Photobiological Sciences, 8, 45-51, 2009.

24 Kazantzidis, A., Smedley, A., Kift, R., Rimmer, J., Berry, J., Rhodes, L. E., and Webb, A.: A
25 modeling approach to determine how much UV radiation is available across the UK and
26 Ireland for health risk and benefit studies, Photochemical & Photobiological Sciences, doi:
27 10.1039/c5pp00008d, 2015. 2015.

28 Kerr, J. B., Evans, W. F. J., and Asbridge, I. A.: Recalibration of Dobson Field
29 Spectrophotometers with a Travelling Brewer Spectrophotometer Standard. In: Atmospheric
30 Ozone, Zerefos, C. S. and Ghazi, A. (Eds.), Springer Netherlands, 1985.

1 Kerr, J. B. and Fioletov, V. E.: Surface ultraviolet radiation, *Atmos. Ocean*, 46, 159-184,
2 2008.

3 Kerr, J. B. and McElroy, C. T.: Evidence for Large Upward Trends of Ultraviolet-B Radiation
4 Linked to Ozone Depletion, *Science*, 262, 1032-1034, 1993.

5 Koukouli, M. E., Balis, D. S., Amiridis, V., Kazadzis, S., Bais, A., Nickovic, S., and Torres,
6 O.: Aerosol variability over Thessaloniki using ground based remote sensing observations and
7 the TOMS aerosol index, *Atmospheric Environment*, 40, 5367-5378, 2006.

8 Kreuter, A., Buras, R., Mayer, B., Webb, A., Kift, R., Bais, A., Kouremeti, N., and
9 Blumthaler, M.: Solar irradiance in the heterogeneous albedo environment of the Arctic coast:
10 measurements and a 3-D model study, *Atmos. Chem. Phys.*, 14, 5989-6002, 2014.

11 Kuttippurath, J., Lefèvre, F., Pommereau, J. P., Roscoe, H. K., Goutail, F., Pazmiño, A., and
12 Shanklin, J. D.: Antarctic ozone loss in 1979–2010: first sign of ozone recovery,
13 *Atmos. Chem. Phys.*, 13, 1625-1635, 2013.

14 Lakkala, K., Arola, A., Heikkilä, A., Kaurola, J., Koskela, T., Kyrö, E., Lindfors, A.,
15 Meinander, O., Tanskanen, A., Gröbner, J., and Hülsen, G.: Quality assurance of the Brewer
16 spectral UV measurements in Finland, *Atmos. Chem. Phys.*, 8, 3369-3383, 10.5194/acp-8-
17 3369-2008, 2008.

18 Li, Z., Wang, P., and Cihlar, J.: A simple and efficient method for retrieving surface UV
19 radiation dose rate from satellite, *Journal of Geophysical Research: Atmospheres*, 105, 5027-
20 5036, 2000.

21 Lucas, R. M., Norval, M., Neale, R. E., Young, A. R., de Gruijl, F. R., Takizawa, Y., and van
22 der Leun, J. C.: The consequences for human health of stratospheric ozone depletion in
23 association with other environmental factors, *Photochemical & Photobiological Sciences*, doi:
24 10.1039/c4pp90033b, 2015. 2015.

25 Mäder, J. A., Staehelin, J., Peter, T., Brunner, D., Rieder, H. E., and Stahel, W. A.: Evidence
26 for the effectiveness of the Montreal Protocol to protect the ozone layer, *Atmos. Chem. Phys.*,
27 10, 12161-12171, 2010.

28 Madronich, S., McKenzie, R. L., Björn, L. O., and Caldwell, M. M.: Changes in biologically
29 active ultraviolet radiation reaching the Earth's surface, *Journal of Photochemistry and*
30 *Photobiology B: Biology*, 46, 5-19, 1998.

1 Madronich, S., McKenzie, R. L., and Caldwell, M. M.: Changes in ultraviolet radiation
2 reaching the earth's surface, *Ambio*, 24, 143-152, 1995.

3 Madronich, S., Shao, M., Wilson, S. R., Solomon, K. R., Longstreth, J. D., and Tang, X. Y.:
4 Changes in air quality and tropospheric composition due to depletion of stratospheric ozone
5 and interactions with changing climate: implications for human and environmental health,
6 *Photochemical & Photobiological Sciences*, doi: 10.1039/c4pp90037e, 2015. 2015.

7 Mayer, B. and Kylling, A.: Technical note: The libRadtran software package for radiative
8 transfer calculations - description and examples of use, *Atmos. Chem. Phys.*, 5, 1855-1877,
9 2005.

10 McLinden, C. A. and Fioletov, V.: Quantifying stratospheric ozone trends: Complications due
11 to stratospheric cooling, *Geophysical Research Letters*, 38, L03808, 2011.

12 Meinander, O., Wuttke, S., Seckmeyer, G., Kazadzis, S., Lindfors, A., and Kyrö, E.: Solar
13 zenith angle asymmetry cases in polar snow UV albedo, *Geophysica*, 45, 183-198, 2009.

14 Meleti, C., Bais, A. F., Kazadzis, S., Kouremeti, N., Garane, K., and Zerefos, C.: Factors
15 affecting solar ultraviolet irradiance measured since 1990 at Thessaloniki, Greece, *Int. J. Rem.
16 Sens.*, 30, 4167-4179, 2009.

17 Meleti, C. and Cappellani, F.: Measurements of aerosol optical depth at Ispra: Analysis of the
18 correlation with UV-B, UV-A, and total solar irradiance, *Journal of Geophysical Research: Atmospheres*, 105, 4971-4978, 2000.

19 Meleti, C., Fragkos, K., Bais, A. F., Tourpali, K., Balis, D., and Zerefos, C. S.: Thirty years of
20 total ozone measurements at Thessaloniki with a MKII Brewer spectrophotometer,
21 Quadrennial Ozone Symposium 2012, Toronto, 2012.

22 Morgenstern, O., Braesicke, P., Hurwitz, M. M., O'Connor, F. M., Bushell, A. C., Johnson, C.
23 E., and Pyle, J. A.: The World Avoided by the Montreal Protocol, *Geophys. Res. Lett.*, 35,
24 L16811, doi:10.1029/2008gl034590, 2008.

25 Newchurch, M. J., Yang, E.-S., Cunnold, D. M., Reinsel, G. C., Zawodny, J. M., and Russell,
26 J. M.: Evidence for slowdown in stratospheric ozone loss: First stage of ozone recovery, *J.
27 Geophys. Res.-Atmos.*, 108, 4507, doi:10.1029/2003jd003471, 2003.

28 Newman, P. A. and McKenzie, R.: UV impacts avoided by the Montreal Protocol,
29 *Photochemical & Photobiological Sciences*, 10, 1152-1160, 2011.

1 Nikitidou, E., Kazantzidis, A., De Bock, V., and De Backer, H.: The aerosol forcing
2 efficiency in the UV region and the estimation of single scattering albedo at a typical West
3 European site, *Atmospheric Environment*, 69, 313-320, 2013.

4 Randel, W. J., Wu, F., Russell, J. M., Waters, J. W., and Froidevaux, L.: Ozone and
5 temperature changes in the stratosphere following the eruption of Mount Pinatubo, *Journal of*
6 *Geophysical Research: Atmospheres*, 100, 16753-16764, 1995.

7 Román, R., Bilbao, J., and de Miguel, A.: Erythemal ultraviolet irradiation trends in the
8 Iberian Peninsula from 1950 to 2011, *Atmos. Chem. Phys. Discuss.*, 14, 15545-15590, 2014.

9 Saffari, A., Daher, N., Samara, C., Voutsas, D., Kouras, A., Manoli, E., Karagkiozidou, O.,
10 Vlachokostas, C., Moussiopoulos, N., Shafer, M. M., Schauer, J. J., and Sioutas, C.: Increased
11 Biomass Burning Due to the Economic Crisis in Greece and Its Adverse Impact on
12 Wintertime Air Quality in Thessaloniki, *Environmental Science & Technology*, 47, 13313-
13 13320, 2013.

14 Schwander, H., Koepke, P., and Ruggaber, A.: Uncertainties in modeled UV irradiances due
15 to limited accuracy and availability of input data, *Journal of Geophysical Research: Atmospheres*, 102, 9419-9429, 1997.

17 Shettle, E. P.: Models of aerosols, clouds and precipitation for atmospheric propagation
18 studies,, In AGARD, Atmospheric Propagation in the UV, Visible, IR, and MM-Wave Region
19 and Related Systems Aspects 14 p, 1989. 1989.

20 Smedley, A. R. D., Rimmer, J. S., Moore, D., Toumi, R., and Webb, A. R.: Total ozone and
21 surface UV trends in the United Kingdom: 1979–2008, *International Journal of Climatology*,
22 32, 338-346, 2012.

23 Solomon, S., Garcia, R. R., Rowland, F. S., and Wuebbles, D. J.: On the depletion of
24 Antarctic ozone, *Nature*, 321, 755-758, 1986.

25 Steinbrecht, W., Köhler, U., Claude, H., Weber, M., Burrows, J. P., and van der A, R. J.: Very
26 high ozone columns at northern mid-latitudes in 2010, *Geophys. Res. Lett.*, 38, 1944-8007,
27 doi:10.1029/2010GL046634, 2011.

28 Takemura, T., Nakajima, T., Dubovik, O., Holben, B. N., and Kinne, S.: Single-Scattering
29 Albedo and Radiative Forcing of Various Aerosol Species with a Global Three-Dimensional
30 Model, *Journal of Climate*, 15, 333-352, 2002.

1 Tourpali, K., Bais, A. F., Kazantzidis, A., Zerefos, C. S., Akiyoshi, H., Austin, J., Brühl, C.,
2 Butchart, N., Chipperfield, M. P., Dameris, M., Deushi, M., Eyring, V., Giorgetta, M. A.,
3 Kinnison, D. E., Mancini, E., Marsh, D. R., Nagashima, T., Pitari, G., Plummer, D. A.,
4 Rozanov, E., Shibata, K., and Tian, W.: Clear sky UV simulations for the 21st century based
5 on ozone and temperature projections from Chemistry-Climate Models, *Atmos. Chem. Phys.*,
6 9, 1165-1172, 2009.

7 Turnock, S. T., Spracklen, D. V., Carslaw, K. S., Mann, G. W., Woodhouse, M. T., Forster, P.
8 M., Haywood, J., Johnson, C. E., Dalvi, M., Bellouin, N., and Sanchez-Lorenzo, A.: Modelled
9 and observed changes in aerosols and surface solar radiation over Europe between 1960 and
10 2009, *Atmos. Chem. Phys. Discuss.*, 15, 13457-13513, 2015.

11 UNEP: Environmental effects of ozone depletion and its interaction with climate change:
12 2010 assessment, report, 278 pp, Nairobi, Kenya., 2010.

13 van Dijk, A., Slaper, H., den Outer, P. N., Morgenstern, O., Braesicke, P., Pyle, J. A., Garny,
14 H., Stenke, A., Dameris, M., Kazantzidis, A., Tourpali, K., and Bais, A. F.: Skin Cancer Risks
15 Avoided by the Montreal Protocol—Worldwide Modeling Integrating Coupled Climate-
16 Chemistry Models with a Risk Model for UV, Photochemistry and Photobiology, 89, 234-
17 246, 2013.

18 Walker, D.: Cloud effects on erythemal UV radiation in a complex topography, doi:
19 <http://dx.doi.org/10.3929/ethz-a-005914035>, 2009. Diss., Eidgenössische Technische
20 Hochschule ETH Zürich, Nr. 18415, 2009, 2009.

21 Watanabe, S., Sudo, K., Nagashima, T., Takemura, T., Kawase, H., and Nozawa, T.: Future
22 projections of surface UV-B in a changing climate, *J. Geophys. Res.-Atmos.*, 116, D16118,
23 doi:10.1029/2011jd015749, 2011.

24 Weatherhead, E. C., Reinsel, G. C., Tiao, G. C., Meng, X.-L., Choi, D., Cheang, W.-K.,
25 Keller, T., DeLuisi, J., Wuebbles, D. J., Kerr, J. B., Miller, A. J., Oltmans, S. J., and
26 Frederick, J. E.: Factors affecting the detection of trends: Statistical considerations and
27 applications to environmental data, *Journal of Geophysical Research: Atmospheres*, 103,
28 17149-17161, 1998.

29 Webb, A. R., Kift, R., Durkin, M. T., O'Brien, S. J., Vail, A., Berry, J. L., and Rhodes, L. E.:
30 The role of sunlight exposure in determining the vitamin D status of the U.K. white adult
31 population, *British Journal of Dermatology*, 163, 1050-1055, 2010.

1 Weihs, P. and Webb, A. R.: Accuracy of spectral UV model calculations: 1. Consideration of
2 uncertainties in input parameters, *Journal of Geophysical Research: Atmospheres*, 102, 1541-
3 1550, 1997.

4 Williamson, C. E., Zepp, R. G., Lucas, R. M., Madronich, S., Austin, A. T., Ballare, C. L.,
5 Norval, M., Sulzberger, B., Bais, A. F., McKenzie, R. L., Robinson, S. A., Häder, D.-P., Paul,
6 N. D., and Bornman, J. F.: Solar ultraviolet radiation in a changing climate, *Nature Clim.
7 Change*, 4, 434-441, 2014.

8 WMO: Scientific assessment of ozone depletion: 2006, *Global Ozone Res. Monit. Proj. Rep.
9* 50, 572 pp., 2007.

10 Yang, E.-S., Cunnold, D. M., Salawitch, R. J., McCormick, M. P., Russell, J., Zawodny, J.
11 M., Oltmans, S., and Newchurch, M. J.: Attribution of recovery in lower-stratospheric ozone,
12 *J. Geophys. Res.-Atmos.*, 111, D17309, doi:10.1029/2005jd006371, 2006.

13 Zerefos, C.: Evidence of the El Chichón stratospheric volcanic cloud in Northern Greece,
14 *Geofísica Internacional*, 23, 299–304, 1984.

15 Zerefos, C. S.: Long-term ozone and UV variations at Thessaloniki, Greece, *Physics and
16 Chemistry of the Earth, Parts A/B/C*, 27, 455-460, 2002.


17 Zerefos, C. S., Tourpali, K., Eleftheratos, K., Kazadzis, S., Meleti, C., Feister, U., Koskela,
18 T., and Heikkilä, A.: Evidence of a possible turning point in solar UV-B over Canada, Europe
19 and Japan, *Atmos. Chem. Phys.*, 12, 2469-2477, 2012.

20

1 Table 1. Trends of TOC, AOD at 320 nm and spectral UV irradiance at 307.5 and 350 nm, for
 2 different periods. Asterisks denote the statistically significant trends

	period	WINTER-SPRING	SUMMER-AUTUMN	YEAR
307.5nm (change % per decade)	1994 – 2006	-	$11.0 \pm 3.3 *$	$7.1 \pm 2.1 *$
	2006 - 2014	-	-0.16 ± 7.7	-0.28 ± 5.0
	1994 – 2014	2.2 ± 1.9	$7.0 \pm 1.9 *$	$4.5 \pm 1.2 *$
350nm (change % per decade)	1994 – 2006	$6.9 \pm 1.8 *$	$6.7 \pm 1.6 *$	$7.0 \pm 1.4 *$
	2006 - 2014	-2.8 ± 4.1	-2.5 ± 3.7	-3.3 ± 3.2
	1994 – 2014	$3.8 \pm 1.0 *$	$3.4 \pm 1.0 *$	$3.3 \pm 0.9 *$
TOC (change % per decade)	1994 – 2014	$1.7 \pm 0.8 *$	0.0 ± 0.6	0.8 ± 0.6
320nm AOD (absolute change per decade)	1994 – 2014	$-0.06 \pm 0.02 *$	$-0.11 \pm 0.02 *$	$-0.09 \pm 0.01 *$

3




1

2

3 Figure 1. Time series of monthly mean all-sky and clear-sky irradiance at 63° ($\pm 1^\circ$) SZA for
 4 (a) 307.5 nm and (b) 324 nm. Monthly means derived from daily means are shown in (c) for
 5 TOC and (d) for AOD at 320 nm. Monthly means were calculated only for months with at
 6 least 10 days of data.

7



1
2
3 Figure 2. Linear trends (in % per decade) of spectral UV irradiance at 307.5 nm (left) and 324
4 nm (right) for clear-sky (upper) and all-sky (lower) conditions derived from Brewers #086
5 and #005, as a function of solar zenith angle. The shaded areas represent the $\pm 1\sigma$ uncertainty
6 of the derived trends.
7






Figure 3. Long-term changes (in % per decade) and associated 1σ uncertainty of the seasonal and the yearly mean spectral irradiance for 307.5, 324 and 350 nm at 64° SZA, for clear skies (a) and all skies (b) at Thessaloniki.



1
2
3 Figure 4. Long-term changes (in % per decade) and the associated 1σ uncertainty of the
4 seasonal and the yearly mean of TOC (blue rhombs) and the AOD at 320 nm (green rhombs).
5 The left (blue) axis corresponds to the changes in TOC while the right (green) axis to changes
6 in AOD.



1

2

3 Figure 5. Yearly mean anomalies and corresponding trends for clear-sky irradiance at 307.5
4 nm (a, b, c) and 350 nm (d, e, f), TOC (g, h, i) and AOD at 320 nm (j, k, l) for December –
5 May (left panels), June – November (middle panels) and for the entire year (right panels). A
6 piece-wise trend consisting of two linear trends has been drawn when a statistically
7 significant turning point has been detected; otherwise a linear trend for the entire period has
8 been drawn.

9