

Please note that the comments made by the reviewer are in black fonts while the responses are in blue fonts.

Review of manuscript version 3 of "Response of OH airglow emissions..." by Ghodpage, Hickey, Taori, Siingh, and Patil.

This version has been considerably revised and improved, and thereby nearly all of my previous objections removed. Now there are only some minor issues left which should be attended before the paper becomes fully acceptable. Therefore, I still qualify this as needing minor revision, but it should be easy to implement. Even a more reasonable ordering of figures 2 (present results, first, and comparisons, later), if the authors find that worthwhile, would not require complex text adjustments.

Response: We thank reviewer for helping us to enhance the quality of our manuscript and also for finding our revision improved.

As far as figure 2 is concerned, we want to give due importance to the earlier investigators, which is making the presentation complicated. Our aim is to reveal that there exist large deviations in gravity wave characteristics in so far large number of investigations and then present our results. To keep a proper balance among past, present studies and future directions, we hope that reviewer will appreciate that presentation in this figure is important.

Details, typos, wordy or unclear text, oversights, suggestions for easier reading:

L35: what does "and its usefulness" refer to? Certainly not to "the pioneering work of Meinel". Maybe, better separate Greet and Bittner references to a new sentence.

Response: Corrected as suggested.

L44: Meriweather -> Meriwether

Response: Modified as suggested.

L54: not clear what the Stubbs et al. reference stands for (it's not an update to Mies ?).

Response: Stubbs remained there by error. We have removed this reference.

L61: missing "d" in "defined as"

Response: Corrected.

L71: what's meant by "by measuring one under adiabatic conditions"?

Response: We have now explained it by adding text.

L73: delete "the" between "as" and "theoretical studies"

Response: Corrected.

L91: while "unique and first of its kind" claims have their role in reports for funding agencies; in a scientific paper, wording should be less provocative (like "such a model comparison over India has not been done before").

Response: Modified the sentence as suggested by the reviewer.

L105: "fillers" -> "filters"

Response: Corrected.

L110: better rearrange -> "with $\sim 10^\circ$ full field of view"

Response: Modified as suggested.

L112: "to to" -> "to"

Response: Done.

L113: better rearrange -> "amplify the very weak photomultiplier output current (in the range of nA) into..."

Response: Done.

L116: is there a need to repeat the 5 hours minimum obs time? At least, add "as mentioned".

Response: Text has been added with a reference.

L156: delete "at" between "atmosphere" and "in"

Response: Modified.

L157 - 161: not clear why ground-based results are discussed in this "space borne" section.

Response: This was written to specify the selection criteria of satellite data as suggested by the reviewer 1 in last review.

L162: delete "satellite" (SABER is an instrument). Main message (SABER data) should not be in parentheses.

Response: Done as suggested.

L171, 172: the previous temperature/intensity bug is still there: Fig. 1a is intensity, 1b is temperature (but figure caption is ok).

Response: Sorry for the error. Now this is corrected.

L176, 177, 180: cosine fit is mentioned three times (the one in L177 would be enough).

Response: Modified as suggested.

L182, 190: as in the previous version, the numerical values of eta in these examples must be based on relative amplitudes, also for temperature. I hope this is still only a typo (where "3.60 K", "1.019 K" should read "3.6%", "1.019%", because otherwise, the eta values derived are WRONG!

Response: Reviewer is correct, text has been modified.

L215-227: I doubt the usefulness of this long list of parameter ranges. If really necessary, it should maybe be put into a table and the discussion shortened (are there any conclusions drawn from this behaviour, except that it's similar for what other people have reported?).

Response: Our aim is to reveal that there exist large deviations in gravity wave characteristics in so far large number of investigations and then present our results.

L233, 234: model results have already been mentioned.

Response: Not really, the model results appear here first and are explained in a later section.

L235: what's new in figure 2a1 and 2b1 is that Kolhapur results have now been plotted separately (also, typo: "there" -> "their")

Response: This was as per the suggestion made by reviewer 1.

L253-255: see my comment on previous version (then, L225): it is not true that Reisin and Scheer 2004 had independent eta values "for the waves of 3000s". They only tried to understand the "relative scales of [their] intensity and temperature variances", in terms of mean eta values. So, citing that paper in the present context does not make sense.

Response: As suggested, we have removed the text.

L263: "2004" and the corresponding reference should therefore be deleted.

Response: As suggested, we have deleted this reference.

L266: The sentence should have ended at the end of the previous line, and "because of, it is also remains to be seen..." could be converted into a new statement (whatever the intended message; I am not sure).

Response: Text has been modified.

L315: there is an error in the Takahashi reference; 2011 is not in the list. It must be one of the earlier papers.

Response: It was supposed to be Takahashi et al. (1990).

L405: "chemical constituents composition" -> "chemical composition", or "minor constituents", or...?

Response: Corrected.

L537: Is there really an AGU report by Meinel with this title in 1950? I am only aware of the 2 part paper in the Astrophysical Journal "OH emission bands in the spectrum of the night sky. I. Astrophys. J. 111, 555-564, 1950", and "OH emission bands in the spectrum of the night sky. II. Astrophys. J. 112, 120-130, 1950".

Response: Yes this report exists.

L578: Tarasick and Hines 1990 paper must go in alphabetic order before Trasick and Shepherd, on next page.

Response: Corrected.

L630, 631: figure captions should explain the figure, but not interpret, like "A close resemblance... are notable"; such newspaper style should be avoided, in order not to distract from the essential info.

"The legends in the figure are as following" and other wordiness should be avoided. Caption should just briefly state that present results (1, 2) are compared to models (3 - 6) and previous observations (7..., 16:...).

It would be more logical to show the Kolhapur results (2a1, 2b1) first, not after 2a, 2b.

By the way: "verses" -> "versus".

Response: The captions have been modified as suggested.

L653, 654: shorten to read "As figure ..., but for..." ("similar to figure..." would be too weak, since it's more than merely "similar").

Response: Modified.

L655: Main message should be "Monthly mean OH emission rate profiles from SABER, for 2010 (solid lines) and 2011 (dashed lines)".

Response: Modified as suggested.

**Response of OH airglow emissions to the mesospheric gravity waves and its
comparisons with full wave model simulation at a low latitude Indian station**

R. N. Ghodpage¹, M. P. Hickey², A. Taori^{3,4}, Devendraa Siingh⁵ and P. T. Patil¹

[1] {Indian Institute of Geomagnetism, Shivaji University Campus, Kolhapur 416004, India}

[2] {Embry-Riddle Aeronautical University, FL - 32114, USA}

[3] {National Atmospheric Research Laboratory, Pakala Mandal, Gadanki (A. P.) 517112,
India}

[4] {now at- National Remote Sensing Center (NRSC), Hyderabad, 500037, India}

[5] {Indian Institute of Tropical Meteorology, Pune-411 008, Maharashtra, India}

Abstract:

The quasi-monochromatic gravity wave induced oscillations, monitored using the mesospheric OH airglow emission over Kolhapur (16.8°N and 74.2°E), India during January to April 2010 and January to December 2011, have been characterized using the Krassovsky method. The nocturnal variability reveals prominent wave signatures with periods ranging from 5.2-10.8 hr as the dominant nocturnal wave with embedded short period waves having wave periods 1.5-4.4 hr. The results show that the magnitude of the Krassovsky parameter, viz., $|\eta|$ ranged from 2.1 to 10.2 for principal or long nocturnal waves (5.2 to 10.8 hr observed periods), and, from 1.5 to 5.4 for the short waves (1.5 to 4.4 hr observed periods) during the years of 2010 and 2011, respectively. The phase, i.e., Φ values of the Krassovsky parameters exhibited larger variability and varied from -8.1° to -167° . The deduced mean vertical wavelengths are found to be approximately -60.2 ± 20 km and -42.8 ± 35 km for long and short wave periods for the year 2010. Similarly, for 2011 the mean vertical wavelengths are found to be approximately -77.6 ± 30 km and -59.2 ± 30 km for long and short wave periods, respectively, indicating that the observations over Kolhapur were dominated by upward propagating waves. We use a full wave model to simulate the response of OH emission to the wave motion and compare the results with observed values. We discuss the observed wave characteristics and cause of the noted differences.

Keywords: OH emissions, Mesospheric gravity wave, Full wave model

1. Introduction

The airglow Hydroxyl emissions (OH) have been oftenly used for studying atmospheric temperature variation in the mesopause region since the pioneering work of Meinel (1950) and its usefulness to derive ~~(Greet et al., 1998, Bittner et al., 2000)~~. ~~The the OH~~ rotational temperature ~~(Greet et al., 1998, Bittner et al., 2000)~~ ~~is one of useful parameters to monitor such variable atmospheric temperature in the mesopause region.~~ The collision frequency of OH with the neutral atmosphere in the neighborhood of 90 km of altitude should be in an order to 10^4 s^{-1} and the life time of the excited Hydroxyl emission is around 3 to 10 msec. (Mies, 1974). This ensures that the excited OH molecules in the rotational energy levels are in a thermal equilibrium with the atmospheric ambient gases (Sivjee & Hamwey, 1987, Takahashi et al., 1998). Thus, it is normally assumed that the rotational state of OH band is in Maxwell-Boltzmann distribution. The radiated light intensity provides a direct measure of OH quantum state distribution in the mesopause, if one knows the Einstein coefficients governing the emission. Meriweather (1975) arrived at an expression for the P1(2) and P1 (5) rotational lines of OH (8-3) band by making use of the vibration-rotation transition probabilities of Mies (1974). Therefore using two lines from a single band we can estimate the rotational temperature by the given quation (Mies, 1974):

$$T_{n,m} = \frac{E_{v'}(J'_m) - E_{v'}(J'_n)}{k \ln \left[\frac{I_n}{I_m} \frac{A(J'_m, v' \rightarrow J''_{m+1}, v'')}{A(J'_n, v' \rightarrow J''_{n+1}, v'')} \frac{2J'_m + 1}{2J'_n + 1} \right]}$$

Where $T_{n,m}$ is the rotational temperature calculated from two line intensities, I_n and I_m , from rotational levels J'_n, J'_m in the upper vibrational level v' , to J''_{n+1}, J''_{m+1} in the lower vibrational level v'' . $E_v(J)$ is the energy of the level (J, v) . $A(J'_n, v' \rightarrow J''_{n+1}, v'')$ is the Einstein coefficient,

for the transition from J'_n, v' to J''_m, v'' . The intensity ratio between P1 (2) and P1 (5) lines of the OH(8,3) band were used to obtain rotational temperature using the transition probabilities as given by Mies (1974) (~~Stubbs et al., 1983~~). Often the observed temporal variations in the mesospheric hydroxyl OH night airglow intensities and rotational temperatures are caused by propagating gravity waves from the lower to the upper atmosphere.

The interaction of these upward propagating waves with the ambient and other waves contribute to the dynamical variability, which in turn is reflected in observed airglow intensity and temperature perturbations (Hines, 1997). Krassovsky (1972) introduced a quantity ‘ η ’ to characterize the wave-induced perturbations. This parameter, termed as ‘Krassovsky’s parameter’, is now defined as $\eta = |\eta| e^{-i\Phi}$, where $|\eta|$ indicates the ratio of the amplitude variation between the emission intensity and temperature perturbations normalized to their time averages and Φ is the phase difference between the intensity wave and its temperature counterpart (e.g., Walterscheid et al., 1987; Taylor et al., 1991). It should also be mentioned here that apart from the pure dynamical processes η can also be affected by various other unknown parameters, such as the variation of local oxygen photochemistry (Hickey et al., 1993) and height variation of the emission layer which affects emission rates and temperature directly (Liu and Swenson 2003; Vargas et al., 2007). Although this can complicate studies of Krassovsky’s parameter, it offers an opportunity to study the above aspects at the same time. Overall, once the physics and chemistry of emissions are well understood, the η values would offer a good tool to study the perturbations caused in a parameter (temperature, brightness/intensity) by measuring one under the assumption that gravity wave induced perturbations are of adiabatic conditions nature.

Utilizing the above, many investigators have carried out observational as well as ~~the~~ theoretical studies on the identification and characterization of gravity wave and tidal signatures

with wave periodicities ranging from few minutes to several hours (e.g., Walterscheid et al., 1987; Hecht et al., 1987; Hickey 1988a, b; Taylor et al., 1991; Takahashi et al., 1992; Reisin and Scheer 1996; Taori and Taylor 2006; Guharay et al., 2008; Ghodpage et al., 2012, 2013). However, observational studies of the magnitude and phase of η over a range of wave periods for a given location and season are sparse. Some of the notable observations of η for the OH emission have been performed by Viereck and Deehr (1989) in the wave period range of $\sim 1 - 20$ hr and by Reisin and Scheer (1996) near to the semidiurnal tidal fluctuations.

In the present work, we utilize the mesospheric OH emission intensity and temperature data obtained during January - April 2010 and January - December 2011, when clear and moonless nights allowed observations to exceed 5 hours duration. We deduce the Krassovsky parameters as a function of observed wave period and also infer the vertical wavelengths for the observed mesospheric waves. Further, we compare our estimates with the earlier results reported by various investigators. We also employ a full-wave model to simulate the effects of wave motions on the OH airglow. This model has been used previously to compare observations and theory of airglow fluctuations (e.g., Hickey et al., 1998; Hickey and Yu 2005). Here, the model is used to estimate the values of the amplitudes and phases of Krassovsky's ratio which are compared to those derived from the observations, making the present study unique ~~and first of its kind over~~ as such model comparison over India has not been done before ~~Indian latitudes~~.

2. Instrumentation and Observations

The mesospheric OH observations are made using the multispectral photometer from Kolhapur (16.8°N, 74.2°E) (Ghodpage et al., 2013, 2014). We analyze the data from January - April 2010 and January-December 2011 when clear sky conditions prevailed for several nights. For the year

2010, 13 nights out of 45 nights of observation clearly showed wavelike features, while in 2011, 29 from 60 nights of data exhibited wavelike variations.

2.1 The multispectral photometer

Regular observations of the night airglow emissions, OI 630.0 nm, OI 557.7nm and OH Meinel (731 nm and 740 nm) band have been carried out at the low latitude station Kolhapur. We have operated multispectral photometer pointing to the zenith over Kolhapur. The filters have a band width of 1 nm and their temperature is controlled by a temperature controller at 24 °C. The temperature coefficient of filter is 0.011 nm/°C. At 24°C the transmission efficiency of filters is 40 - 70 %. We kept the integration time for each filter 15 seconds which results in repetition time of 90 seconds with an accuracy of approximately $\pm 0.5\%$ for line intensity. The photometer has F/2 optics with ~~full field of view~~ $\sim 10^\circ$ full field of view. The stepper motor rotation and sensing of the initial position is performed by computer controlled software. As the detector, the EMI9658B photomultiplier tube is used. An amplifier (high gain trans-impedance) is used to ~~to~~ convert and amplify the very weak photomultiplier's ~~very weak~~ output current (in the range of nA) ~~output current~~ in-to corresponding voltage form. In the absence of standard calibration source, we have used relative intensities (arbitrary units). In order to study the wave features present in the MLT region, we consider clear sky nights having more than 5 hours of continuous OH band data as mentioned in earlier reports (e.g., Taori et al., 2005).

2.2 Full Wave Model

The full-wave model is a linear, steady-state model that solves the linearized Navier-Stokes equations on a high resolution vertical grid to describe the vertical propagation of acoustic-

gravity waves in a windy background atmosphere including molecular viscosity and thermal conduction, ion drag, Coriolis force and the eddy diffusion of heat and momentum in the mesosphere. The model description, including equations, boundary conditions and method of solution has been described elsewhere (Hickey et al., 1997; Walterscheid and Hickey 2001; Schubert et al., 2003). The neutral perturbations are used as input to a linear, steady-state model describing OH airglow fluctuations (Hickey and Yu 2005).

The model solves the equations on a high resolution vertical grid subject to boundary conditions, and allows quite generally for the propagation in a height varying atmosphere (non-isothermal mean state temperature and height varying mean winds and diffusion). The linearized equations are numerically integrated from the lower to the upper boundary using the tri-diagonal algorithm described by Bruce et al. (1958) and Lindzen and Kuo (1969). The lower boundary is set well below the region of interest and a sponge layer is implemented to avoid effects of wave reflection in the airglow response. In this study the lower boundary (the bottom of the lower sponge layer) is placed at 250 km below $z = 0$ (i.e., -250 km). The wave forcing is through the addition of heat in the energy equation. The heating is defined by a Gaussian profile with a full-width-at-half-max of 0.125 km. It is centered at an altitude of 10 km. A Rayleigh-Newtonian sponge layer in addition to natural absorption by viscosity and heat conduction prevents spurious reflection from the upper boundary. At the upper boundary (here 300 km altitude) a radiation condition is imposed using a dispersion equation that includes viscous and thermal dissipation (Hickey and Cole 1987). The mean state is defined using the Mass Spectrometer Incoherent Scatter (MSIS) model (Hedin 1991).

A set of linear perturbation equations for the minor species involved in the OH emission chemistry is solved using the approach described in Hickey (1988). This assumes that these

minor species have the same velocity and temperature perturbations as the major gas (which are deduced from the full-wave model). A vertical integration of the volume emission rates through the vertical extent of the OH layer provides the brightness and brightness-weighted temperature perturbations, from which Krassovsky's ratio is determined. The OH chemistry we use is the same as that used previously (Hickey et al., 1997) and is for the OH (8-3) emission. We also determine the vertical wavelength at the peak of the OH emission layer evaluated from the phase variations of the temperature perturbations determined by the full-wave model.

2.3 Space borne measurements

The Sounding of the Atmosphere using Broadband Emission Radiometry (SABER), on-board the Thermosphere Ionosphere Mesosphere Energetic and Dynamics (TIMED) satellite, is a high-precision broadband radiometer which measures limb radiance (orbital inclination at 74°) of the terrestrial atmosphere ~~at~~ in 10 selected spectral bands ranging from 1.27 to 15 μm . In the present study, we note larger values of $|\eta|$ occur during 2011 compared to 2010 for long/principal waves, which indicates a larger intensity to temperature perturbation ratio over Kolhapur during the passage of the waves during 2011. This could be due to the differences in either the background atmosphere or the dynamical processes. To identify the differences in the OH emission layer in year 2010 and 2011, we scrutinize the OH volume emission rate profile for Kolhapur region (~~obtained from SABER~~)~~satellite~~. The selected latitude-longitude grids are 10°N to 20°N and 70°E to 90°E representing Kolhapur. The criteria for the selection of SABER data are such that: (i) the SABER pass should be during typical observation times (excluding twilight time).

3. Results and Discussion

To identify the wave structures in the data, we utilize the perturbation amplitudes normalized to their time averaged values (hereafter referred to as mean values) in the intensity and temperature data to calculate the Krassovsky ratio. To illustrate this, we show a typical example corresponding to the data obtained on 26-27 January 2011 in Figure 1. We plot the intensity deviations ~~temperature deviations~~ from their mean values in figure 1(a), while, the temperature deviations ~~intensity deviations~~ from their mean values are plotted in figure 1(b). We note that night airglow intensity variations show a long-period wave with embedded short-period oscillatory features. On this night, the mean airglow intensity is found to be ~ 1.83 arbitrary units and the mean temperature data is ~ 195.75 K. To identify the nocturnal variability plotted together with data as solid red lines are the of best-fit cosine model (e.g., Taori et al., 2005) as follows. ~~Also, best fit cosines are shown (solid red lines).~~

$$Y = A \cos \left[\pi \frac{(X - X_c)}{T} \right] \quad (1)$$

where, A is the amplitude of the fitted wave of half-period T with phase X_c , and X is the time. The solid red lines in figure 1 show the results of the best-fit cosine model. We observed the presence of $\sim 8.2 \pm 1.1$ hr and 8 ± 1.3 hr waves with relative amplitudes (normalized to their mean values and converted to corresponding % amplitude) $\sim 3.6\%$ ~~0 K~~ and 25.64%, in the nocturnal temperature and intensity variability, respectively. Given the uncertainties involved in the observations, we consider these to be the same waves. Further, we compute the $|\eta|$ value for this wave to be 7.12 ± 1.2 . To identify the shorter period features in the data we obtain residuals from the best-fit model values. The figure 1c and 1d panels show the nocturnal variability of the

residual intensity and temperature respectively. The best-fit model reveals the presence of $\sim 4.2 \pm 0.2$ and 3.0 ± 0.8 hr wave in the temperature and intensity residuals, respectively. Once again we treat these as the same wave for the reason explained above. The best-fit analysis shows the amplitudes of this wave to be ~ 1.019 ~~%K~~ and 3.75% arbitrary units in the temperature and intensity data, respectively. Hence, the $|\eta|$ value for short period waves is estimated to be 3.68 ± 0.9 . In general we note that in worst case, the maximum error in $|\eta|$ values are $<25\%$. The phase difference between the intensity and temperature waves is obtained with the help of best-fit parameters which were also verified with a cross correlation analysis. The phase of the principal waves (maxima) (period ~ 8.2 hr) was ~ 24.88 hr in the temperature data and 24.4 hr in the intensity data, which results in the phase difference of ~ 0.48 hr, i.e., Φ values of $-21.07 \pm 12^\circ$. Similarly, for the shorter period (period ~ 4.2 hr) the Φ values are estimated to be $-114.3 \pm 20^\circ$. We can also estimate the vertical wavelength with the help of Krassovsky's parameter following the approach elaborated by Tarasick & Hines (1990).

$$\lambda_z = \frac{2\pi\gamma H}{(\gamma - 1)|\eta| \sin(\phi)} \quad (2)$$

where $\gamma = C_p / C_v = 1.4$ is the ratio of specific heats, and $H = 6$ km is the scale height. This formula is valid for zenith observations and for plane waves. It is not valid for the evanescent waves. Equation (2), negative vertical wavelength corresponds to downward phase propagation (i.e., upward energy propagation), and that means that temperature oscillations precede the intensity oscillations in phase (e.g., Takahashi, ~~H.~~, et al. 1990). Using the above relation we find that vertical wavelength for the two cases discussed above are $\sim -51.5 \pm 15$ and -39.3 ± 40 km for the long period and the short period waves, respectively. Note that the long period wave estimates may be biased when the data length is comparable to that of the wave period and

therefore in our study we have considered only those waves whose periods are substantially less than the length of the available data.

The above analysis was carried out on nighttime events recorded during 2010 and 2011 when the prominent wave features were visible. During the 2010 period, the principal nocturnal waves in the data show the wave periods vary from 5.2 to 10.8 hr with corresponding temperature amplitudes ranging from 2 to 13.8 K. Similarly for 2011, wave periods vary between 5.2 and 8.4 hr with corresponding temperature amplitudes lying between 1.1 K and 15.7 K. However, the intensity amplitudes of the principal waves vary from 7.9% to 49.9 % and 5% to 90% for 2010 and 2011, respectively. We note that the estimated $|\eta|$ values were found to range from 2.1 - 10.5 for the principal wave. In the case of the short period waves, the periods ranged from 1.5 to 4.4 hr (for 2010) and 2.8 to 4.4 hr (for 2011) with corresponding temperature amplitudes ranging from 0.68 K to 12.2 K and 0.4 K to 14.2 K. The corresponding intensity amplitudes fall in range between $\sim 1.54\%$ to 46.8% and 1.32% to 46.8 for 2010 and 2011, respectively. The phase (Φ) values also exhibit large variability for long (short) period waves, range in between -27° and -167° (-27° and -150°) for 2010 and -8.1° and -65.2° (-39.1° and -122.6°) for 2011. For 2010 the deduced vertical wavelengths are found to vary from -32.2 km to -140 km and -24 km to -88 km for the long and short period waves, respectively. Similarly, for 2011 the deduced vertical wavelengths are found to vary from -40 km to -102 km, and -26 km to -92.4 km for the long and short period waves, respectively.

In Figure 2a we plot our results for $|\eta|$ (hereafter η) with pink half-filled squares indicating the estimates for the year 2010 and olive half-filled squares for the year 2011. We plot Φ in Figure 2b using the same symbols as used in Figure 2a. For a comparison, we also show the values of η and Φ of the results from other investigations are shown (Viereck and Deehr 1989;

Takahashi et al., 1992; Oznovich et al., 1995, 1997; Drob 1996; Reisin and Scheer 1996; Taylor et al., 2001; Lopez-Gonzalez et al., 2005). Also shown in the figure are the model estimates of Schubert et al. (1991), Tarasick and Shepherd (1992a, b), Walterscheid and Schubert (1995). We also plot observed η and Φ values against ~~there~~their observed period in figure (2a1 and 2b1). In general, we note that the parameter η increases with wave period. It is evident that the observed η and Φ values in our study show a large spread in their distribution as compared to the model values. A similar spread in the distribution of observed values of η (Figure 2a) from 1.03 to 7.85 has also been observed by other investigators (e.g., Takahashi et al., 1992). It may be noted that the values of η for the OH data in our study lie somewhere between the model estimates and the values observed by other investigators. Also noteworthy in this figure is that our η values are closer to the model values reported by Tarasick and Shepherd (1992a) for the waves with horizontal wavelength 500 km. The phase ' Φ ' values, on the other hand show significantly larger deviations from this model for 2010, while for 2011 the match between measured and modeled phases appear to be better. We note that our measurements of Φ matches somewhat with those reported by Viereck and Deehr (1989), while large differences with other investigators can be easily noted. The variation of Φ values with respect to the wave periodicity, obtained in the 2010 year clearly shows that most of the time we observe values to be higher than those obtained by different models.

Of the importance is that Reisin & Scheer (2001) found η values of 3.47 ± 0.07 corresponding to the wave periods between 1000s and 3h. Our observed values of η (arithmetic mean, 4.4 ± 1 for 2010 year and 5.7 ± 1.7 for 2011 year) for OH measurements agree well with this report. ~~In a further report, based on 5-year observations, Reisin and Scheer (2004), found the mean value of η to be ~ 5.6 for the nightly semidiurnal type waves and ~ 3.4 for the waves of 3000 s periodicity~~

~~which is in agreement with our values.~~ In another study based on long-term observations with a spectral airglow temperature imager (SATI) from a mid-latitude station, Lopez-Gonzalez et al. (2005) reported a mean value of η of approximately ~ 8.6 for the OH measurements with a larger variability than our observations show. In another report, Guharay et al. (2008), found that for wave periods ranging from 6 hr to 13 hr, values of η in between 1.7 to 5.4, while the phase varied from -13° to -90° . Similarly, Aushev et al. (2008) presented amplitudes of the Krassovsky parameter for wave periods of 2.2 to 4.7 hr which in range from 2.4 to 3.6 while the phase values in between -63° to -121° . It is noteworthy that our derived values broadly agree with Guharay et al. (2008, 2009), Reisin and Scheer (2001, 2004) and Viereck and Deehr (1989) while they are somewhat different from the values reported by Lopez-Gonzalez et al. (2005) which may be due to the fact that their observations corresponded to higher latitude than ours. ~~because of, it is~~ also remains to be seen that when mesopause altitude itself changes from low to high latitudes how would that reflects in the Krassovsky parameters.

The results of (η and Φ) shown in Figure 2 emphasize that there are significant differences in the Krassovsky parameters derived from one study to another. This we suspect to be caused by the variations in the altitudinal profile of oxygen and its effect on the η through the complex OH chemistry (Walterscheid et al., 1994). Another possibility over low latitudes was discussed by Makhlouf et al. (1995) who suggested the quenching caused by the perturbed molecules during their transitions from several vibrational levels. Winds also affect the OH response to gravity waves and therefore they will also contribute to the spread of values seen between the various observation studies (e.g., Sonnemann G. and M. Grygalashvyly, 2003).

Note that our observations as well as models show the phase Φ for OH to be a negative value indicating upward propagating waves (see Tarasick and Shepherd, 1992a, b). In general we

280 note that our Φ values, although on some occasions are closer to Viereck and Deehr (1989)
 281 observations, show deviations from other investigators and are larger than the model values on
 282 most occasions. Differences in theory and observation may be due to the horizontal wavelength
 283 assumed in the model and or the Prandtl number (ratio of kinematic viscosity to thermal
 284 diffusivity) assumed. The Prandtl number is important in theoretical calculations and modeling,
 285 especially when in terms of dissipating waves owing to molecular viscosity and thermal
 286 diffusivity while they propagate in the atmosphere (Hickey 1988). An error in the Prandtl
 287 number assumption will affect the derived wave parameters (λ_z , η etc.), which will successively
 288 mask the actual ones. In this regard, Makhlof et al. (1995) studied the variations in the η values
 289 by modifying the model proposed by Hines and using a photochemical dynamical model;
 290 however, they were still unable to explain the appearance of the negative phases appropriately.
 291 Hines and Tarasick (1987) found a wide range of η variability, a result supported by our
 292 measurements. Further, Hines and Tarasick (1997) subsequently discussed the necessary
 293 correction for thin and thick layer approximations for the calculation of η from airglow
 294 emissions due to gravity waves interaction. They also pointed out that OH emission intensity,
 295 which affects the derived η values, does not depend on the oxygen profile and other minor
 296 species, which contradicts the theory of Walterscheid et al. (1994), Schubert et al. (1991). The
 297 calculated vertical wavelengths (VW) for all the nights of the observation are shown in Figure 2c
 298 as pink half filled squares indicating the estimates for the year 2010 and olive half filled squares
 299 for the year 2011. Large differences exist from one night to another. The VW has a large
 300 variability ranging from -41 km to -102 km (2010) and -36.2 km to -140 km (2011) for long
 301 period waves, and, -26 to -92.4 and -24 to -88 km for short period waves period of 2010 and
 302 2011 years respectively. In 2010 (and 2011) years, the mean VW values for long and short

period waves are calculated to be -60.2 ± 20 km (-77.2 ± 40 km) and -42.8 ± 15 km (-59.2 ± 30) respectively. Further, unlike the clear dependency on the wave period noted in the Krassovsky parameters (η and Φ) no clear trend is noted in the calculated VW. We also plot the values reported by Reisin and Scheer (1996) and Lopez-Gonzalez et al. (2005) for a comparison. It is noteworthy that for all the days the VW for the long period wave are higher than the VW of short period waves. We also observed that VW values calculated for 2011 year are larger than 2010 year calculated values. We note that the values reported by Reisin and Scheer (1996) are approximately -30 km with about 40 km variability, which is a good agreement with our values. However, Lopez-Gonzalez et al. (2005) observed VW values to be approximately -10 km deduced from their OH observations, which do not agree with our values. Further, Ghodpage et al. (2012) analyzed the long-term nocturnal data of 2004-2007 and also observed that the VW lies between 28.6 and 163 km. Recently, Ghodpage et al. (2013) studied the simultaneous mesospheric gravity wave measurements in the OH emission from Gadanki and Kolhapur, inferring mean VWs varying from -26 to -60 km for the Kolhapur observations. Takahashi et al. (2011, 1990) reported vertical wavelengths varying from 20 to 80 km, which is in agreement with our values.

4. Comparison with the Full Wave Model Results

Wave simulations were performed using the Full Wave Model (FWM) for which the representative inputs were taken for the duration of observations reported in section 3. The observations were conducted over an approximate one month period spanning February 8th and March 13th, and accordingly we used the middle date of this observation period (February 25th) in the MSIS model to represent the undisturbed mean state. The latitude used was 16.8° N, and

the local time was midnight. Because the speed and direction of wave propagation were not determined from the observations, several simulations were performed for each wave period in which the direction of propagation (eastward, northward and westward propagation) and the phase speed (50 m/s, 100 m/s and 150 m/s) were varied. Note that the mean winds (not shown) in these simulations were derived from the Horizontal Wind Model (HWM) using the same input parameters as used for the MSIS model. The derived meridional winds (not shown) are far smaller than the zonal winds for the conditions considered here, and so while results for eastward and westward propagation differed quite markedly, those for northward and southward propagation did not. Hence we considered only a single direction (northward) for meridional propagation.

We also performed a tidal simulation using an equivalent gravity wave model (Lindzen 1970; Richmond 1975), as implemented in an earlier study (Walterscheid and Hickey 2001). The horizontal wavelength and Coriolis parameter are adjusted to give maximal correspondence with a given tidal mode. Here, we performed calculations for the terdiurnal (3,3), (3,4), (3,5) and (3,6) modes using parameters provided by Richmond (1975). The simplifications inherent in this approach are discussed by Walterscheid and Hickey (2001).

Comparisons between the full wave model results for η , Φ and λ_z and the values inferred from the observations are shown in figure 3a, 3b and 3c, respectively. In figure 3a we compare the observed values of η for 2010 and 2011. The observed values of η are represented as pink and olive lower half-filled squares for 2010 and 2011, respectively. In figure 3a we note that at few of the longer wave periods, the observed values of η are in good agreement with the full wave model results. For short period waves the values of η inferred from the observations appear

to be bounded by the model values for waves with horizontal phase velocities are 50 and 100 m/s, respectively. For example, for 3.6 hr wave periods, the average of the values of η inferred from the observations is 3.7, while the full wave model values lie between about 0.5 (for the 100 m/s wave) and 7 (for the 50 m/s, eastward propagating wave). For the 8 hr wave periods, the average of the values of η inferred from the observations is 5.7, which is bounded by the full wave model estimates for waves having a horizontal phase velocity of 50 m/s and different propagation directions.

Overall, we note that the comparison between the observed η values and the modeled values can be explained by gravity waves whose horizontal phase velocities range from 50 m/s to 100 m/s. In this regard, an earlier investigation by Pragati Sikha et al. (2010) reported observed gravity wave horizontal phase speeds (for periods 5 min to 17 min) varying between 10 m/s and 48 m/s. The propagation directions were reported to be preferentially towards the north. More recently, Taori et al. (2013) studied mesospheric gravity wave activity in the OH and OI 558 nm emissions from Gadanki. They observed that the gravity waves were moving in the north–west direction. The average phase velocity of the ripple-type waves was found to be 23.5 m/s. The other, band-type waves, with horizontal scales of about 40 km, were found to be propagating from south to north with an estimated phase speed of 90 m/s.

The vertical wavelengths (λ_z) calculated using the observed values of η and Φ differ significantly from the full wave model estimate for waves with phase velocities below 100 m/s. More typically, a comparison between those values inferred from the observations and those derived from the model tend to agree for phase velocities in the 100 - 150 m/s range. However, it should be noted that vertical wavelengths inferred from the observations are based on the use of the inferred Krassovsky's ratio, η , in Eq. (2). Please note that the errors in the determination of

371 the phase (Φ) of η may lead to significant errors (proportional to $\cot\Phi$) in the determination of
372 λ_z , especially as Φ approaches $\pm 180^\circ$.

373 The differences noted in the observed and modeled estimates of Krassovsky ratio
374 magnitudes η and phase (Φ) may be associated with the limitation arising due to dynamics as
375 well as the measurements. In terms of measurements limitation, the parameters achieved with the
376 best fit method may have leaked contribution from other wave components which may be
377 dynamically varying within a wave period. In terms of dynamics, that full wave model uses
378 climatological density (both major gas and minor airglow-related species) and wind profiles
379 which will introduce uncertainties. This point has been previously elaborated by Walterscheid et
380 al. (1994) with respect to the effect of a change in the [O] profile on the OH response to wave
381 motions.

382 It is interesting to note that the arithmetic mean values of $|\eta|$ for the years 2010 and 2011
383 were 4.4 and 5.7 respectively. When we look at each $|\eta|$ value from one wave period range to
384 other, the difference is found to be more than 30% which is well above the maximum errors in
385 the estimation. One may further argue that this difference may not be significant. For this, we
386 looked at the mode of the values for periods ranges 1-4 hr, 4-6 hr, 6-8 hr and 8-10 hr. We found
387 that in each case in the year 2011 mode values are larger than the year 2010. The differences
388 noted in the magnitude of the observed Krassovsky ratio η between 2010 and 2011 may be
389 associated with variations in the height and shape of the undisturbed OH emission profile. We
390 use the SABER data to investigate this aspect. To check whether there was a difference in the
391 OH emission layer structure, we selected the nighttime OH emission profile for a grid
392 encompassing 10°N to 20°N latitudes and 70°E to 90°E latitudes during February, March and
393 April months of the years 2010 and 2011. We have selected the February to March period

394 because the optical airglow data used in this study was acquired primarily during these months.
 395 The monthly mean values of OH emission rates are plotted in Figure 4. The solid curves
 396 correspond to 2010 data while the dashed curves correspond to 2011 data. We note that the peaks
 397 of OH emission layer during February, March and April of 2010 occurred at 84.2 km, 82.8 km
 398 and 85.1 km altitude, respectively, while the corresponding peaks for 2011 were found to occur
 399 at 85.8 km, 85.6 km and 85.2 km altitude. This suggests that the peak of the emission layer
 400 occurred at a somewhat lower altitude in 2010 compared to 2011. Also, the mission rates during
 401 February and March were found to be higher in 2010. It is important to note that in an earlier
 402 study, Ghodpage et al. (2013) compared the Krassovsky ratios at two different latitudes, Gadanki
 403 (13.5 N, 79.2E) and Kolhapur (16.8°N and 74.2°E) and noted a lower OH emission layer peak
 404 over Kolhapur and also larger estimated η values over Kolhapur. In the present case, instead of
 405 the location, it is the difference in the measurement year where the peak emission altitudes of the
 406 OH emission layer are somewhat different. As the peak emission layer arise due to the chemical
 407 reactions involving odd oxygen, it is proposed that chemical-~~constituents~~ composition was
 408 different from the year 2010 to the year 2011. Therefore, the noted emission rates may be
 409 responsible for the observed differences in the Krassovsky parameters. A further question that
 410 arise here is why the peaks should be different from one year to the other. As these months are
 411 pre-monsoon, when a large scale oscillation namely, El Niño/Southern Oscillation (ENSO)
 412 sweeps through the south Asian continent, we looked at the ENSO strength based on the
 413 Multivariate ENSO Index (MEI). This index is shown in table 1, where it is noteworthy that the
 414 MEI index for 2010 (January to May) is of opposite sign to that for the corresponding months in
 415 2011. We postulate that these large scale processes have a profound impact on the observed
 416 wave energetics and dynamics at mesospheric altitudes. Large scale processes induced the wave

oscillations associated with the ENSO. The ENSO generates a spectrum of waves which are of planetary scales. These are expected to generate a secular variation in temperature and density structure throughout the atmosphere. A difference in ENSO suggests that these forcing are different in the two years (2010, 2011). At present, we do not know through which process the ENSO may have implications in the observed wave characteristics. However, we believe that further investigation is required in order to confirm whether or not any such associations really do exist.

5. Conclusion:

We report the Krassovsky parameters for the observed gravity waves from Kolhapur (16.8°N and 74.2°E) and their comparison with the full wave model.

- 1) It is evident that the observed values of Krassovsky parameters in our study show a large spread in their distribution as compared to the model values (shown in Figure 2a). A similar spread in the distribution has also been reported by other investigators. We have also observed magnitude of η values is larger in the year 2011 than 2010.
- 2) It is also notable that the values of η for the OH data in our study lie between the model estimates and the values observed by other investigators. Whereas the phase values are more than the model values on most occasions. We note that our Φ measurements match with those reported by Viereck and Deehr (1989), while they show large differences with other investigators values.
- 3) Observed vertical wavelength (VW) values broadly agree with the range reported by other investigators and are found to vary from -26 to -140 km. We also noted that VW

values calculated for 2011 year are larger than 2010 year calculated values. Most of wave propagating upward in direction.

- 4) Comparison of observed η and Φ values agree fairly well with the full wave model results for waves with 50 and 100 m/s horizontal phase velocities. Vertical wavelengths tend to agree for waves with 100 and 150 m/s horizontal phase velocities, except for the longest period waves for which vertical wavelength cannot be reliably inferred from the observations.

The database used in the present study is limited in terms of the length and locations. Based on the above conclusions we emphasize that more rigorous study using coordinated observations and modeling are required to uncover the physics occurring at upper mesosphere.

Acknowledgements

This work is carried out under the research grant funded by Ministry of Science and Technology and Department of Space, Govt. of India. RNG thank the Director, Indian Institute of Geomagnetism (IIG), Navi Mumbai for encouragement to carry out this work. The night airglow observations at Kolhapur were carried out under the scientific collaboration program (MoU) between IIG, Navi Mumbai and Shivaji University, Kolhapur. MPH acknowledges the support of NSF grant AGS-1001074.

6. References

460 Aushev, V. M., Lyahov, V. V., Lopez-Gonzalez, M. J., Shepherd, M. G., and Dryna, E. A. :Solar
 461 eclipse of the 29 March 2006: results of the optical measurements by MORTI over Almaty
 462 (43.03°N, 76.58°E), J. Atmos. Sol. Terr. Phys., 70, 1088–1101, 2008.

463 Bruce, G. H., Peaceman, D. W., Rachford, Jr. H. H., and Rice, J. D.: Calculations of unsteady-
 464 state gas flow through porous media, Petrol. Trans. AIME, 198, 79-92, 1953.

465 Bittner, M., Offermann, D., Graeft.: Mesopause temperature variability above a midlatitude
 466 station in Europe. Journal of Geophysical Research, 105(D2): 2045–2058, 2000.

467 Drob, D. P.: Ground-based optical detection of atmospheric waves in the upper mesosphere and
 468 lower thermosphere, Ph. D. Thesis, University of Michigan, Ann Arbor, MI., 1996.

469 Ghodpage, R. N., Singh, D., Singh, R. P., Mukherjee, G. K., Vohat, P., and Singh, A. K.: Tidal
 470 and gravity waves study from the airglow measurements at Kolhapur (India), J. Earth Syst.
 471 Sci.121, 6, 1511–1525, 2012.

472 Ghodpage, R. N., Taori, A., Patil, P. T., and Gurubaran, S.: Simultaneous mesospheric gravity
 473 wave measurements in OH night airglow emission from Gadanki and Kolhapur – Indian low
 474 latitudes, Currents Science, 104, 1, 98-105, 2013.

475 Ghodpage, R.N., Taori, A., Patil, P. T., Gurubaran, S., Sharma, A. K., Nikte, S., and Nade, D.:
 476 Airglow Measurements of Gravity Wave Propagation and Damping over Kolhapur (16.8° N,
 477 74.2° E), International Journal of Geophysics (IJG), Volume 2014,1-9,
 478 <http://dx.doi.org/10.1155/2014/514937>, 2014.

479 Greet, P.A., French, WJR., Burns, G B., Williams, PFB., Lowe, R. P., & Finlayson, K. :OH (6-2)
 480 spectra and rotational temperature measurements at Davis, Antarctica, *Annales Geophysicae*,
 481 16(1), 77–89, 1998.

482 Guharay, A., Taori, A., Bhattacharjee, B., Pant, P., Pande, P., and Pandey, K.: First ground-
 483 based mesospheric measurements from central Himalayas, *Current Science*, 97, 664-669,
 484 2009.

485 Guharay, A., Taori, A., and Taylor, M.: Summer-time nocturnal wave characteristics in
 486 mesospheric OH and O₂ airglow emissions, *Earth Planets Space*, 60, 973–979, 2008.

487 Hecht, J. H., et al.: Observations of wave-driven fluctuations of OH nightglow emission bfrom
 488 Sondre Stromfjord, Greenland, *J. Geophys. Res.*, 92, 6091-6099, 1987.

489 Hedin, A. E. :Extension of the MSIS thermosphere model into the middle and lower atmosphere,
 490 *J. Geophys. Res.*, 96, 1159 – 1172, 1991.

491 Hickey, M. P., and Yu, Y.: A full-wave investigation of the use of a “cancellation factor” in
 492 gravity wave-OH airglow interaction studies, *J. Geophys. Res.*, 110, A01301,
 493 doi:10.1029/2003JA01372, 2005.

494 Hickey, M. P., Huang, T.-Y., and Walterscheid, R.: Gravity wave packet effects on chemical
 495 exothermic heating in the mesopause region, *J. Geophys. Res.*, 108(A12), 1448,
 496 doi:10.1029/2002JA009363, 2003.

497 Hickey, M. P., Walterscheid R. L., and Schubert, G.: Gravity wave heating and cooling in
 498 Jupiter’s thermosphere, *Icarus*, 148, 266-281, 2000.

499 Hickey, M. P., Walterscheid, R. L., Taylor, M. J., Ward, W., Schubert, G., Zhou, Q., Garcia, F.,
 500 Kelley, M. C., and Shepherd G. G.: Numerical simulations of gravity waves imaged over
 501 Arecibo during the 10-day January 1993 campaign, J. Geophys. Res., 102, 11,475-11,489,
 502 1997.

503 Hickey, M.P., Schubert, G., and Walterscheid, R. L.: Gravity wave-driven fluctuations in the O₂
 504 atmospheric (0-1) nightglow from an extended, dissipative emission region, J. Geophys.
 505 Res., 98(13),717-730, 1993.

506 Hickey, M. P.: Effects of eddy viscosity and thermal conduction and coriolis force in the
 507 dynamics of gravity wave driven fluctuations in the OH nightglow, J. Geophys. Res., 93,
 508 4077, 1988.

509 Hickey, M.P., Taylor, M.J., Gardner, C.S., and Gibbons, C.R. Full-wave modeling of small-
 510 scale gravity waves using Airborne Lidar and Observations of the Hawaiian Airglow
 511 (ALOHA-93) O(1S) images and coincident Na wind/ temperature lidar measurements, J.
 512 Geophys. Res. 103, 6439-6453,1998.

513 Hickey, M. P., and Cole, K. D.: A quartic dispersion equation for internal gravity waves in the
 514 thermosphere, J. Atmos. Terr. Phys., 49, 889-899, 1987.

515 Hines, C. O.: A fundamental theorem of airglow fluctuations induced by gravity waves, J.
 516 Atmos. Sol. Terr. Phys.,59, 319–326, 1997.

517 Hines, C. O., and Tarasick, D. W.: Layer truncation and the Eulerian/ Lagrangian duality in the
 518 theory of airglow fluctuations induced by gravity waves, J. Atmos. Sol. Terr. Phys., 59, 327–
 519 334, 1997.

520 Hines, C. O., and Tarasick, D. W. : On the detection and utilization of gravity waves in airglow
521 studies, Planet Space Sci., 35, 851–866, 1987.

522 Krassovsky, V. I.: Infrasonic variation of OH emission in the upper atmosphere, Ann. Geophys.,
523 28, 739–746, 1972.

524 Lindzen, R. S.: Internal gravity waves in atmospheres with realistic dissipation and temperature,
525 part I: Mathematical development and propagation of waves into the thermosphere, Geophys.
526 Fluid Dyn., 1, 303-355, 1970.

527 Lindzen, R. S., and Kuo, H. L.: A reliable method for the numerical integration of a large class
528 of ordinary and partial differential equations, Mon. Wea. Rev., 97, 732-734, 1969.

529 Liu, A. Z., and Swenson, G. R.: A modeling study of O₂ and OH airglow perturbations induced
530 by atmospheric gravity waves, J. Geophys. Res, 108, D4, 4151, doi:10.1029/2002JD002474,
531 2003.

532 Lopez-Gonzalez, M. J., et al.: Tidal variations of O₂ Atmospheric and OH(6-2) airglow and
533 temperature at mid-latitude from SATI observations, Ann. Geophys., 23, 3579–3590, 2005.

534 Makhlof, U. B., Picard, R. H., and Winick, J. R.: Photochemical-dynamical modeling of the
535 measured response of airglow to gravity waves, 1: basic model for OH airglow, J. Geophys.
536 Res., 100, 11,289–11,311, 1995.

537 | Mies, F. H.:Calculated vibrational transitions probabilities of OH ($X^2\pi$). Journal of —Molecular
538 | Spectroscopy, 53, 150–188, 1974.

539 | Meinel, A. B.:OH Emission bands in the spectrum of the night sky I. ~~American Geophysical~~
540 | ~~Union, 31 (21);Astrophys. J., 111, 555 – 564,~~ 1950.

541 Meriwether, J. W.: High latitude airglow observations of correlated short term fluctuations in the
 542 hydroxyl Meinel 8-3 band intensity and rotational temperature, Planet. Space Sci., 23, 1211–
 543 1221, 1975.

544 Oznovich, I., Walterscheid, R. L., Sivjee, G. G., and McEwen, D. J.: On Krassovsky's ratio for
 545 ter-diurnal hydroxyl oscillations in the winter polar mesopause, Planet Space Sci., 45(3), 385–
 546 394, 1997.

547 Oznovich, I., McEwen, D. J., and Sivjee, G. G.: Temperature and airglow brightness oscillations
 548 in the polar mesosphere and lower thermosphere, Planet Space Sci., 43, 1121–1130, 1995.

549 Pragati, R.S., Parihar, N., Ghodpage, R., Mukherjee, G.K.: Characteristics of gravity waves in
 550 the upper mesosphere region observed by OH airglow imaging, Current Science 98, 392–
 551 397, 2010.

552 Reisin, E.R., and Scheer, J.: Vertical propagation of gravity waves determined from zenith
 553 observations of airglow, Adv. Space Res. 27(10), 1743-1748, 2001.

554 ~~Reisin, E. R., and Scheer, J.: Gravity wave activity in the mesopause region from airglow~~
 555 ~~measurements at El Leoncito, J. Atmos. Sol. Terr. Phys., 66, 655–661, 2004.~~

556 Reisin, E. R., and Scheer, J.: Characteristics of atmospheric waves in the tidal period range
 557 derived from zenith observations of O₂(0-1) Atmospheric and OH (6-2) airglow at lower mid
 558 latitudes, J. Geophys. Res., 101, 21,223–21,232, 1996.

559 Richmond, A. D.: Energy relations of atmospheric tides and their significance to approximate
 560 methods of solution for tides with dissipative forces, J. Atmos. Sci., 32, 980-987, 1975.

561 Schubert, G., Hickey, M. P., and Walterscheid, R. L.: Heating of Jupiter's thermosphere by the
 562 dissipation of upward propagating acoustic waves, *Icarus*, 163, 398-413, 2003.

563 Schubert, G., Walterscheid, R. L., and. Hickey, M. P.: Gravity wave-driven fluctuations in OH
 564 nightglow from an extended, dissipative emission region, *J. Geophys. Res.*, 96 (A8), 13,869–
 565 13,880, 1991.

566 Sivjee, G. G., & Hamwey, R. M.: Temperature and chemistry of the polar mesopause OH.
 567 *Journal of Geophysical Research*, 92(A5): 4663–4672, 1987.

568 Stubbs, L. C., Boyd, J. S., and Bond, F. R.: Measurement of the OH rotational temperature at
 569 Mawson, East Antarctica, *Planet. Space Sci.*, 31 (8), 923–932, 1983.

570 Sonnemann G. and Grygalashvyly, M.: The zonal wind effect on the photochemistry within the
 571 mesosphere / menopause region, *Adv. Space Res. Vol. 32 (5)*, 719-724, 2003.

572 Takahashi, H., Buriti, R. A., Gobbi, D., and Batista, P. P.: Equatorial planetary wave signatures
 573 observed in mesospheric airglow emissions, *J. Atmos. Sol. Terr. Phys.*, 64, 1263–1272, 2002.

574 Takahashi, H., Sahai, Y., Batista, P. P., and Clemesha, B. R.: Atmospheric gravity wave effect
 575 on the airglow O₂(0-1) and OH (9-4) band intensity and temperature variations observed from
 576 a low latitude station, *Adv. Space Res.*, 12(10), 131–134, 1992.

577 Takahashi, H., Sahai, Y., and Teixeira, N.R.: Airglow intensity and temperature response to
 578 atmospheric wave propagation in the mesopause region, *Adv. Space Res.* 10, (10)77-(10)81,
 579 1990.

580 ~~Tarasick, D. W. and Hines, C. O.: The observable effects of gravity waves in airglow emission,~~
 581 ~~*Planet. Space Sci.*, 38, 1105–1119, 1990.~~

582 Takahashi, H, Batista, P. P, Buriti, R. A, Gobbi, D, Nakamura, T, Tsunda, T & Fukao S.
583 Simultaneous measurements of airglow OH emission and meteor wind by a scanning photometer
584 and the MU radar. Journal of Atmospheric and Solar-Terrestrial Physics, 60(17): 1649–1668,
585 1998.

586 [Tarasick, D. W. and Hines, C. O.: The observable effects of gravity waves in airglow emission,](#)
587 [Planet. Space Sci., 38, 1105–1119, 1990.](#)

588 Taylor, M. J., Gardner, L. C., and Pendleton, Jr. W. R.: Long-period wave signatures in
589 mesospheric OH Meinel (6,2) band intensity and rotational temperature at mid-latitudes, Adv.
590 Space Res., 27(6–7), 1171– 1179, 2001.

591 Taylor, M. J., Turnbull, D. N., and Lowe, R. P.: Coincident imaging and spectrometric
592 observations of zenith OH nightglow structure, Geophys. Res. Lett., 18, 1349–1352, 1991.

593 Taori, A., and Taylor, M. J.: Characteristics of wave induced oscillations in mesospheric O2
594 emission intensity and temperatures, Geophys. Res. Lett., 33, L01813,
595 doi:10.1029/2005GL024442, 2006.

596 Taori, A., Taylor, M. J., and Franke, S.: Terdiurnal wave signatures in the upper mesospheric
597 temperature and their association with the wind fields at low latitudes (20°N), J. Geophys.
598 Res., 110, D09S06, doi: 10.1029/2004JD004564, 2005.

599 Taori, A., Jayaraman, A., Kamalakar, V.: Imaging of mesosphere–thermosphere airglow
600 emissions over Gadanki (13.51N, 79.21E)-first results, J. Atmos. Sol. Terr. Phys. 93, 21–28,
601 <http://dx.doi.org/10.1016/j.jastp.2012.11.007>, 2013.

602 Tarasick, D. W., and Shepherd, G. G.: Effects of gravity waves on complex airglow chemistries:
603 1. $O_2(b^1\Sigma_g^+)$ emission, J. Geophys. Res., 97, 3185–3193, 1992a.

604 Tarasick, D. W., and Shepherd, G. G.: Effects of gravity waves on complex airglow chemistries:
605 2. OH emission, J. Geophys. Res., 97, 3195–3208, 1992b.

606 Vargas, F., Swenson, G., Liu, A., and Gobbi, D.: $O(^1S)$, OH, and $O_2(b)$ airglow layer
607 perturbations due to AGWs and their implied effects on the atmosphere, J. Geophys. Res., 112,
608 D14102, doi:10.1029/2006JD007642, 2007.

609 Viereck, R. A., and Deehr, C. S.: On the interaction between gravity waves and the OH Meinel
610 (6-2) and O_2 Atmospheric (0-1) bands in the polar night airglow, J. Geophys. Res., 94, 5397–
611 5404, 1989.

612 Walterscheid, R. L., and Schubert, G.: Dynamical-chemical model of fluctuations in the OH
613 airglow driven by migrating tides, stationary tides, and planetary waves, J. Geophys. Res.,
614 100, 17,443–17,449, 1995.

615 Walterscheid, R. L., and Hickey, M. P.: One-gas models with height-dependent mean molecular
616 weight: Effects on gravity wave propagation, J. Geophys. Res., 106, 28,831–28,839, 2001.

617 Walterscheid, R. L., Schubert, G., and Hickey, M. P.: A Comparison of Theories for Gravity
618 Wave Induced Fluctuations in Airglow Emissions, J. Geophys. Res., 99, 3935, 1994.

619 Walterscheid, R. L., Schubert, G., and Hickey, M. P.: Comparison of theories for gravity wave
620 fluctuations in airglow emissions, J. Geophys. Res., 99, 3935–3944, 1994.

621 Walterscheid, R. L., Schubert, G., and Straus, J. M.: A dynamical chemical model of wave-
622 driven fluctuations in the OH nightglow, J. Geophys. Res., 92, 1241 – 1254, 1987.

623

624

625

Figure captions:

Figure 1. Nocturnal variability in the mesospheric OH emissions on 26-27 January 2011. The upper panels represent the mean deviations in (a) intensity and (b) temperature data. Bottom panels represent (c) intensity and (d) temperature residuals. Solid line curves in each plot show the result of simple best-fit cosine model.

Figure 2 (a) ~~A comparison~~ Distribution of Krassovsky parameter ' η ', ~~s of data to their respective wave periods reported by investigators (list not exhaustive).~~ The x -axis shows the wave periodicity and the y-axis is for amplitude of Krassovsky parameters (η ~~and Φ~~) ~~in each plot. A close resemblance between the observational values and discrepancy between the observational and theoretical estimates are notable.~~ The legends in the figure are as following: (~~η ÷ 1 (for 2010 year) & 2 (for 2011 year);~~ present study ; 3, Schubert et al. 500 km; 4, Schubert et al. 1000 km; 5, Tarasick and Shepherd 500 km; 6, Tarasick and Shepherd 1000 km; 7, Takahashi et al. (1992); 8, Oznovich et al. (1995); 9, Drob et al. (1996); 10, Reisin and Scheer (1996); 11, Taylor et al. (2001); 12, Guharay et al (2008); 13, Walterscheid and Schubert (1995); 14, Lopez-Gonzalez et al. (2005); 15, Oznovich et al. (1997)); 16, Viereck and Deehr (1989).

Figure 2 (a1) Observed values of η ~~verses~~ versus wave period over Kolhapur alone.

Figure 2 (b) ~~A comparis~~ Distribution of ϕ phase values of Krassovsky parameter ' Φ ', reported by investigators (list not exhaustive) to their respective wave periods (~~Φ ÷ 1 (for year 2010); & 2 (for year 2011);~~ present study; 3, Schubert et al. 500 km; 4, Schubert et al. 1000 km; 5, Tarasick and Shepherd 500 km; 6, Tarasick and Shepherd 1000 km; 7, Viereck and Deehr (1989); 8, Oznovich et al. (1995); 9, Drob et al. (1996); 10, Reisin and Scheer (1996); 11, Taylor et al.

(2001); 12, Guharay et al.(2008); 13, Walterscheid and Schubert (1995); 14, Lopez-Gonzalez et al. (2005); 15, Oznovich et al. (1997); 16,Viereck and Deehr (1989);).

Figure 2 (b1) Observed values of Φ verses wave period over Kolhapur alone.

Figure 2(c) Deduced vertical wavelength (VW) for both the short and long period wave as function of wave periodicity. Also shown comparison with values obtained by other investigators.

Figure 3(a) Comparison with η calculated by observation of both year and Full wave model simulation with their respective wave period. Pink and olive lower half filled square shows the 2010 and 2011 year η observations (1 and 2 present study η ; 3,FWM simulation of η for 50 m/s horizontal phase velocity; 4, FWM simulation of η for 100 m/s horizontal phase velocity; 5, FWM simulation of η for 150 m/s horizontal phase velocity).

Figure 3(b) Simillar ~~as to~~ figure 3(a) but for phase values for both the short and long period wave. **Figure 3(c)** Simillar ~~to as~~ figure 3(a) but for deduced vertical wavelength (VW).

Figure 4. The monthly (February, March and April) mean ~~values of~~ OH emission rates ~~are profiles from SABER for the year 2010 (solid lines) and 2011 (dashed lines). shown in plot-(which are obtain from SABER data) . The solid lines plot the data for the year 2010 while the dashed lines represent the year 2011.~~

Table 1. Comparisons of deduced wave parameters in 2010,2011 years with MEI index and OH altitudes. The observed quantities are mean for their representative wave periods. (JFM-January, February and March months like this)

26 - 27 January 2011; Kolhapur








Figure 1.






678 Figure 2(b)




680

681 Figure 2(b1)

682



689 Figure 2(c)



691

692 Figure (3a)





Figure 3(b)



695

696 Figure 3 (c)




Figure 4.

704 Table 1.

Year	Mean η (\pm Errors)		Mean ($-\Phi$) (Deg.)		Mean ($-VW$) (km)		OH altitude (km)	MEI index				
	Long wave period	Short wave period	Long wave period	Short wave period	Long wave period	Short wave period		JFM	FMA	MAM	SON	OND
2010	4.4 \pm 1	2.3 ± 0.9	90.6 ± 40	70.4 \pm 45	60.2 ± 20	42.8 ± 15	82 km to 85.1 km during February – April	1.1	0.8	0.5	-1.4	-1.3
2011	5.7 \pm 1.7	2.7 ± 0.6	33.8 ± 40	64.4 \pm 40	77.6 ± 40	59.2 ± 30	85.1 km to 86 km during February – April	-1.1	-0.8	-0.6	-0.9	-0.9

705

706