Editor Initial Decision: Reconsider after minor revisions (Editor review) (12 Jun 2015) by Paul B. Shepson

Comments to the Author:

This paper is publishable, except that I would like to request one editorial change.

In your response to Reviewer 2's question about snowpack photochemistry on the original page 3647, lines 21-23, you stated "...HNO3 and HONO are also known to deposit to snow...".

It is also known that HONO is emitted from sunlit snowpacks, e.g. as discussed in Zhou et al. (GRL, 2001). This fact impacts the next part of your addition to the paper, in which you state "...a surface depletion of HO2 radicals would be expected". However, snowpack photochemistry is highly complex and while few emission fluxes have been measured, it is easy to hypothesize the reverse (HO2 being enhanced at the surface) as HOx precursors such as HONO and HCHO will be emitted from the snowpack. You might want to cite the appropriate studies from places like Summit, Alert, and Barrow about emission of HOx and NOx precursors from sunlit snowpacks, and be a bit more equivocal about the impact on HO2 near the surface. With this addition/change, the paper will be accepted.

Response:

The editor is of course correct in his assessment of the statements in question and we appreciate the criticism. The statement implying "a surface depletion of HO2 radicals..." was made considering only HO2NO2 as the contributing species. As the editor states this is incorrect as the system is significantly more complex than this. In order to improve this portion of the work we have reworked the manuscript to extend the content and description of these surface effects. The following changes have been made:

- The comments as to the direction of the HO2 gradient have been removed, indicating that the system is more complex and simultaneous flux measurements of reactive species (e.g. HONO, H2CO, H2O2) are necessary.
- The discussion on the impact of the surface fluxes on HOx and NOx has been moved up to the portion where figure 4 is discussed.
- Additional discussion has been added on the fluxes including references as the editor suggests which address previous studies focusing on snow surface emissions and chemistry.

Details of the changes made can be found in the attached annotated manuscript, where the majority of changes occur in section 4.

- 1 Peroxynitric acid (HO₂NO₂) measurements during the UBWOS 2013 and 2014 studies
- 2 using iodide ion chemical ionization mass spectrometry.
- 3 Patrick R. Veres^{a,b,*}, James M. Roberts^b, Robert J. Wild^{a,b}, Peter M. Edwards^{a,b,c}, Steven S.
- 4 Brown^b, Timothy S. Bates^{d,e}, Patricia K. Quinn^e, James E. Johnson^d, Robert J. Zamora^f,
- 5 Joost de Gouw^{a,b}
- 6
- 7 ^aCooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, Colorado, USA
- 8 ^bChemical Sciences Division, NOAA Earth System Research Laboratory, Boulder, CO, USA
- 9 °Now at Department of Chemistry, University of York, York, YO10 5DD, UK
- 10 ^dJoint Institute for the Study of the Oceans and Atmosphere, University of Washington, Seattle, WA, USA
- 11 °NOAA PMEL, Seattle, WA, USA
- 12 ^bPhysical Sciences Division, NOAA Earth System Research Laboratory, Boulder, CO, USA
- 13 *Correspondence to: patrick.veres@noaa.gov
- 14

15 Abstract

- 16 Laboratory work is reported here establishing iodide ion chemical ionization mass spectrometry
- 17 (I⁻ CIMS) as a sensitive method for the unambiguous detection of peroxynitric acid (HO₂NO₂,
- 18 PNA). A dynamic calibration source for HO₂NO₂, HO₂, and HONO was developed and
- 19 calibrated using a novel total NO_v detector (NO_v CaRDS). Photochemical sources of these
- 20 species were used for the calibration and validation of the Γ CIMS instrument for detection of
- 21 HO₂NO₂. Ambient observations of HO₂NO₂ using I CIMS were made during the 2013 and 2014
- 22 Uintah Basin Wintertime Ozone Study (UBWOS) are presented. Strong inversions leading to a
- 23 build-up of many primary and secondary pollutants as well as low temperatures drove daytime
- 24 HO₂NO₂ as high as 1.5 ppbv during the 2013 study. A comparison of HO₂NO₂ observations to
- 25 mixing ratios predicted using a chemical box model describing an ozone formation event

1	observed during the 2013 wintertime shows agreement in the daily maxima $\mathrm{HO}_2\mathrm{NO}_2$ mixing
2	ratio, but a differences of several hours in the timing of the observed maxima. Observations of
3	vertical gradients suggest that the ground snow surface potentially serves as both a net sink and
4	source of HO ₂ NO ₂ depending on time of day. Sensitivity tests using a chemical box model
5	indicate that the lifetime of HO ₂ NO ₂ with respect to deposition has a non-negligible impact on
6	ozone production rates on the order of 10%.
7	
8	
9	1 Introduction
10	Hydrogen oxides (HO _x = HO ₂ + OH) and nitrogen oxides (NO _x = NO ₂ + NO) play central roles
11	in atmospheric photochemistry. HO ₂ , a product of OH-initiated VOC oxidation, reacts with NO
12	to produce NO ₂ , a key step in the photochemical ozone formation cycle in the troposphere.
13	Peroxynitric acid (often referred to as PNA, HO ₂ NO ₂ , or HNO ₄) plays an important role in the
14	coupling of atmospheric HO _x and NO _x cycles (Niki et al., 1977), especially at low temperatures.
15	PNA serves as an important HO_x and NO_x reservoir species altering the oxidative capacity of the
16	atmosphere on regional and global scales (Kim et al., 2007;Chen et al., 2001;Davis et al.,
17	2001;Carpenter et al., 2000).
18	PNA is formed via the reaction of HO ₂ and NO ₂ (DeMore et al., 1997;Sander et al.,
19	2011).
• •	

 $HO_2 + NO_2 \xrightarrow{M} HO_2NO_2$

(1)

Formation via reaction (1) is favored at low temperatures and high pressures (Kim et al., 2007).
 Unimolecular decomposition is temperature dependent and occurs on a timescale of
 approximately 10 seconds at 1 atm, 298K with the lifetime rapidly increasing to hours at 258K
 (Gierczak et al., 2005);

5

$$6 \qquad HO_2NO_2 \xrightarrow{M} HO_2 + NO_2 \tag{2}$$

7

PNA can be lost *via* photolysis in the near ultraviolet (Jimenez et al., 2005) and near infrared *via*an overtone band (Roehl et al., 2002;Stark et al., 2008);

11
$$\frac{HO_2NO_2 \xrightarrow{\lambda} OH + NO_2}{}$$
(3)

12

13
$$\frac{HO_2NO_2 \xrightarrow{\lambda} HO_2 + NO_2}{}$$
(4)

14 Or by reaction with OH (Jimenez et al., 2004);

15

$$16 \qquad \qquad \underbrace{HO_2NO_2 + OH \longrightarrow H_2O + NO_2 + O_2}_{(5)}$$

17

In the lower troposphere, reactions (3), (4), and (5) typically occur on timescales of days to
months thereby implying that the dominant loss of PNA in the lower troposphere is typically
unimolecular dissociation or deposition.

21 Deposition of PNA on snow surfaces has been observed in various studies where the

22 atmospheric lifetime of PNA in Polar Regions is largely controlled by dry deposition (Huey et

1	al., 2004;Slusher et al., 2002;Jones et al., 2014). Additional laboratory studies have been
2	performed confirming the efficient uptake of PNA to ice (Li et al., 1996;Ulrich et al., 2012) and
3	sulfuric acid solutions (Zhang et al., 1997). Dependent on the fate of PNA after deposition, the
4	formation and subsequent deposition of PNA has been suggested to result in a net loss of HO_{x}
5	and an increase in NO (Grannas et al., 2007), with reactions (1) to (5) thereby having an impact
6	on tropospheric ozone formation (Salawitch et al., 2002).
7	Observations of PNA are generally limited in scope with most measurements focusing on
8	polar regions (Slusher et al., 2002;Huey et al., 2004;Slusher et al., 2010), the free troposphere
9	(Murphy et al., 2004;Singh et al., 2006;Singh et al., 2007;Keim et al., 2008;Kim et al., 2007),
10	and the stratosphere (Rinsland et al., 1996; Rinsland et al., 1986; Sen et al., 1998). Mean PNA
11	observations from these studies range from tens of pptv in polar surface regions to several
12	hundred pptv in the upper troposphere/lower stratosphere. The impacts of PNA on upper
13	tropospheric chemistry have been widely discussed (Brune et al., 1999;Wennberg et al.,
14	1999;Faloona et al., 2000), with one study in particular identifying a PNA contribution as high as
15	20% of the total NO _y budget (Murphy et al., 2004). Lower-tropospheric, mid-latitude
16	measurements, in contrast, remain largely unexplored with the exception of, to our knowledge, a
17	single airborne study conducted in Mexico (Spencer et al., 2009) where PNA concentrations up
18	to 600 pptv were observed, and were correlated with O ₃ formation.
19	The general lack of lower-tropospheric, mid-latitude observations is driven in part by
20	two factors (i) a diminished atmospheric impact of PNA due to higher rates of thermal
21	decomposition with respect to the generally colder polar and upper atmosphere, and (ii) a lack of
22	instrumentation capable of providing sensitive unambiguous measurements of PNA. Many of the
23	techniques available measure PNA as a component of NOy using O3/NO chemiluminescence

1 (Keim et al., 2008) or total peroxynitrates via thermal decomposition laser induced fluorescence 2 (Murphy et al., 2004). Currently available instrumentation capable of unambiguous measurement 3 of PNA is limited to remote sensing detection via IR absorption spectroscopy (Rinsland et al., 1986; Rinsland et al., 1996; Sen et al., 1998) or in-situ measurement via chemical ionization mass 4 5 spectrometry using the SiF₆⁻ ion (Slusher et al., 2002;Slusher et al., 2001;Huey, 2007), CF₃O⁻ 6 (Spencer et al., 2009; Huey et al., 1996), and I (Abida et al., 2011). Among these, CIMS 7 techniques have been shown to have sufficient sensitivity and time resolution for the in-situ 8 monitoring of PNA concentrations in the lower troposphere. 9 In this work, we present laboratory and ambient measurements illustrating the utility of 10 iodide ion CIMS for unambiguous measurement of PNA. Additionally, we show applicability of 11 this technique for the detection of both HO_2 and HONO, atmospheric species that are also 12 integral to HO_x and NO_x cycles. These results were necessary in order to rule out potential mass overlap or PNA interferences from the sampling of HO₂ and HONO. A photo-source has been 13 14 developed for dynamic production of PNA, HO₂, and HONO to assist with laboratory calibration 15 and elucidation of various ionization schemes. PNA observations made during the 2013 and 16 2014 Uintah Basin Wintertime Ozone Study (UBWOS) will be presented and compared to the 17 results of a chemically explicit box model developed to describe the air quality in the Uintah 18 Basin during a high ozone event observed during the 2013 study (Edwards et al., 2014). The 19 impact of PNA on HOx and NOx budgets, particularly as it relates to the photochemical 20 production of ozone, will also be discussed. 21

- 22
- 23 2 Experimental Setup

1 We present both laboratory and field data collected over a two-year period encompassing the 2 2013 and 2014 UBWOS field studies. The following describes the instrumentation used in this 3 work as well as a short description of the UBWOS study and field conditions. Information and links pertaining to the 2013 and 2014 Uintah Basin Winter Ozone Studies are available on the 4 5 web (http://esrl.noaa.gov/csd/groups/csd7/measurements/2013ubwos/ and 6 http://esrl.noaa.gov/csd/groups/csd7/measurements/2014ubwos/). 7 2.1 8 Instrumentation 9 2.1.1 Iodide Ion CIMS (I⁻ CIMS) 10 The I⁻ CIMS instrument consists of an ion flow tube coupled to a quadrupole mass spectrometer. Briefly, CH_3I in N_2 is passed through a ²¹⁰Po ionizer resulting in the production of iodide (Γ) 11 12 ions. Subsequently, I reacts in the flow tube with a sample gas resulting in the production of 13 anions that are mass filtered using a quadrupole mass spectrometer and detected. Additional 14 details about the instrument can be found elsewhere (Slusher et al., 2004). The instrumental 15 differences with respect to that described in Slusher et al. 2004 will be discussed as they relate to 16 the field deployments and laboratory measurements presented. 17 The sensitivity of the I CIMS to the species presented in this work (HO_2NO_2 , $HONO_3$), 18 and HO_2) is dependent on the degree of water clustering present in the flow tube. Some 19 ionization reactions occur faster through the I-H₂O⁻ cluster, likely as a result of ligand switching 20 via an intermediate cluster ion, while other reactions are independent of the water cluster 21 population. A detailed discussion of iodide-water cluster distributions and subsequent effect on 22 the iodide ion CIMS chemistry can be found in the supplementary material of Lee et al. 2014 23 (Lee et al., 2014). Even with a declustering region present, the ratio of the first water cluster to

the parent ion is a valuable diagnostic and can often be a more appropriate way to account for
 changes in the primary ion signal through normalization of measured signals to the sum of I⁻ and
 I-H₂O⁻, as was performed in this work.

The extent of the ion clustering in the I⁻ CIMS can be controlled using a collisional 4 5 dissociation chamber and through addition of water directly into the flow tube to modify the 6 expected ionization chemistry. This was achieved by flowing ~ 10 sccm N₂ bubbled through 7 water directly into the flow tube at a pressure of 30 torr. In this study, the ratio of the iodide water cluster ion (I-H₂O⁻, m/z 145) to the iodide ion (I⁻, m/z 127) ranged from approximately 5% 8 9 to 50% and was adjusted to maximize the instrument sensitivity towards individual species or 10 classes of compounds depending on the application. I CIMS was operated in the laboratory with 11 a switchable inlet to allow for sampling through a section of PFA tubing (0.025 m) either at room 12 temperature ('cold') or operated as an inlet dissociator (130°C, 'hot'). 13 The inlet configuration used during the 2014 UBWOS study was identical to that used in 14 the laboratory experiments, where inlet switching could be performed to sample from either a 15 'cold' or 'hot' inlet. During the 2013 UBWOS study, however, the I CIMS did not employ a 16 switchable inlet; rather a 'hot' inlet dissociator at 150°C was used throughout the entire 17 measurement period. Never-the-less post experiment tests with the inlet configuration used 18 during UBWOS 2013 permit us to quantify HO₂NO₂ during that study, albeit with increased 19 uncertainty. 20

21 2.1.2 Total NO_y Cavity Ring Down Instrument

22 A novel four channel laser diode based cavity ring-down (CRD) instrument was used to quantify

23 NO, NO₂ and NO_y, during the laboratory portion of this work. Details of the NO_y CRD

instrumentation can be found elsewhere (Wild et al., 2014). Briefly, NO_y is thermally dissociated
 to NO and NO₂ in a heated quartz inlet. Any NO in the sample air or formed *via* catalysis is
 converted to NO₂ by addition of O₃ after the quartz converter. Quantification of NO₂ in the
 sample is performed using cavity ring-down at 405 nm.

5 The quartz converter is typically operated with a gas temperature of 720°C, however, the 6 temperature can be adjusted downward to measure specific classes of compounds or individual 7 species that contribute to total NO_v while excluding unwanted interferences. A similar method 8 has been used previously for the speciation of NO_v in both ambient and field measurements 9 (Wooldridge et al., 2010;Day et al., 2002). In our experiments, the gas temperature in the 10 converter was optimized for detection of PNA and HONO, 130°C and 720°C respectively, a set 11 point determined by scanning the quartz heater through the entire temperature range as shown in 12 Wild et al. 2014 (Wild et al., 2014). A detailed discussion on the application of this technique 13 towards the quantification of HONO and PNA generated in a dynamic laboratory calibration 14 source is included below.

15

16 2.2 Field Site Description

The I⁻ CIMS instrument was deployed during the 2013 and 2014 UBWOS field studies
conducted in the Uintah Basin, UT. The goal of these studies was to improve our understanding
of the chemistry leading to wintertime observations of elevated ozone levels within the basin.
The field site configuration in 2014 is best described by the UBWOS 2012 field report (Lyman
and Shorthill, 2012) where a nearly identical site design was employed while a detailed
description of the 2013 measurements is available in the UBWOS 2013 field report (Stoeckenius
and McNally, 2014).

1	During both the 2013 and 2014 studies, observations of HO ₂ NO ₂ gradients over snow
2	were conducted by using a moveable (7m, 35°C) PFA inlet during the 2013 study and a
3	switchable dual inlet system (each inlet was 20m, 30°C) for the 2014 study. The total inlet flow
4	used during the 2014 study was approximately 20 slpm from which the I ⁻ CIMS subsampled 2
5	slpm. During the 2013 study, a total inlet flow of 2 slpm was sampled through a short stainless
6	steel section attached to the end of a PFA inlet and heated to a temperature of 65°C, to dissociate
7	N_2O_5 prior to sampling preventing CINO ₂ interferences due to reactions on the inlet surface
8	(Behnke et al., 1997). In this inlet, HO ₂ NO ₂ will dissociate as well as recombine, considering the
9	stability of HO_2 on PFA surfaces. The CIMS instrument in 2013 was not configured to monitor
10	the I-HO ₂ ⁻ or I-HO ₂ NO ₂ ⁻ ions, however, NO ₃ ⁻ , routinely monitored for diagnostic purposes,
11	provides a means to estimate ambient $\mathrm{HO}_2\mathrm{NO}_2$ as a result of the large sensitivity of the CIMS to
12	HO ₂ NO ₂ at this ion.
13	In addition to gas phase measurements conducted at the field site, snow samples were
14	collected daily throughout the UBWOS 2013 and 2014 study. Surface samples were collected
15	from the top 3 cm of snow from undisturbed locations within 1.3km of the field site. At each site,
16	2 to 6 1L glass jars were collected and kept frozen until analysis, typically within 48 hours of
17	sampling. Prior to analysis via an ion chromatograph (IC), samples were melted and filtered
18	using a 25mm diameter $0.4\mu m$ pore size nucleopore filter. Filtered samples were subsequently
19	injected directly into the IC and analyzed for anions (e.g. Cl ⁻ , NO ₂ ⁻ , NO ₃ ⁻) and cations (e.g. Na ⁺ ,
20	$NH_{4}^{+}, K^{+}).$
21	

23 3 Laboratory Results

Laboratory experiments using I CIMS were preformed both in preparation and upon the 1 2 conclusion of the UBWOS 2014 study. The goal of these experiments was to adapt the I⁻ CIMS technique for the sensitive detection of PNA and develop a method for calibration. The 3 4 development of an HO₂ based photolysis source for the production of HO₂NO₂ also led to the 5 recognition that the I⁻ CIMS can be applied to the direct measurement of HO₂ radicals. 6 Additionally, the HO₂ photolysis source is readily adaptable for the production of an on-line 7 HONO calibration standard. These laboratory developments in calibration standard production 8 and I CIMS detection of PNA, HONO, and HO2 will be discussed in detail in the following 9 sections.

10

11 3.1 Standard Generation and Detection

The following describes the ion molecule reactions with I⁻ ions resulting in the detection of
HO₂NO₂, HONO and HO₂:

15
$$I^- + HO_2 \longrightarrow I \cdot HO_2^-$$
 (5)

16

17
$$I^{-} + HO_2NO_2 \longrightarrow I \bullet HO_2NO_2^{-}$$
(6)

18

19
$$I^{-} + HO_2NO_2 \longrightarrow HOI + NO_3^{-}$$
(7)

20

21
$$\frac{HO_2NO_2 \xrightarrow{\Lambda} HO_2 + NO_2 \xrightarrow{I^-} I \cdot HO_2^-}{(8)}$$

22

1

3	where reaction (6) is only observed using a 'cold' inlet while reaction (8) occurs upon thermal
4	dissociation of PNA in the inlet. Experimentally, the above reactions, with the exception of
5	reaction (7), were observed in this work to occur predominantly via reaction of the hydrated
6	iodide cluster (I-H ₂ O ⁻ , m/z 145) based on the strong dependence of sensitivity on water vapor
7	observed during these experiments. This implies that the above reactions are ligand switching
8	reactions made faster by the ability of H ₂ O to accommodate excess energy of reaction through
9	extra degrees of freedom up to and including dissociation.
10	Normalized (10^6 cps I ^{\circ}), background corrected mass spectra are shown in Figure 1 for
11	each of the sources. The ratio of m/z 145 (I-H ₂ O ⁻) to m/z 127 (I ⁻) is displayed as a percentage on
12	each panel for reference. These spectra represent the result of the subtraction of a normalized
13	background mass spectrum from a sample spectrum. The background correction method that was
14	applied varies for each species and is dependent on the sample matrix, which will be described
15	separately in the following sections. In all cases, the mass ranges from 126 to 128 and 144 to 146
16	corresponding to the I $(m/z~127)$ and I-H_2O $(m/z/~145)$ ions were removed to simplify
17	interpretation of the mass spectra. These spectra allow for the identification of impurities in the
18	photolysis sources used as well as demonstrating the ions that were used for the unambiguous
19	detection of each analyte. Each of these sources will be discussed in detail in the following
20	sections.
21	

22 3.1.1 HO₂ Radical

1	HO_2 radicals are generated in the laboratory <i>via</i> photolysis of H_2O in the presence of O_2
2	(Dusanter et al., 2008). A mixture of approximately 100 sccm N_2 and 0.5 sccm O_2 was bubbled
3	through water and diluted into a 5 slpm flow of N_2 . The mixture is subsequently passed into a
4	PFA photolysis cell and irradiated with a 185 nm Pen-Ray $^{\ensuremath{\mathbb{R}}}$ lamp. The N_2 dilution flow is
5	produced using boil off from a high-pressure liquid nitrogen Dewar to limit the amount of $\mathrm{NO}_{\boldsymbol{x}}$
6	and VOC in the system. Detection of HO2 via I CIMS occurs through direct observation of the
7	parent ion cluster (I-HO ₂), therefore, it is not necessary to add CO in order to titrate OH as no
8	measurement interference is expected. Addition of trace amounts of CO was found to increase
9	the concentration of HO ₂ produced, though trace amounts of NO and NO ₂ from the steel cylinder
10	mixture resulted in an increase in PNA and HONO backgrounds with increasing CO.
11	Figure 1a shows the difference mass spectrum of the HO ₂ radical mixture less the
12	instrument background. In this case, the instrument background was the ion signal measured
13	prior to turning on the 185 nm lamp. It is clear from the mass spectrum that there is only a single
14	dominant peak observed at m/z 160 (I-HO2). A method for the quantitative calibration of the
15	HO_2 radical source and the I ⁻ CIMS instrument for the detection of HO_2 radicals will be
16	discussed in section 3.1.4. The sensitivity of Γ CIMS to the detection of HO ₂ was determined to
17	be a function of the mixing ratio of water in the flow tube as well as the extent of
18	clustering/declustering in the system, e.g. reduction of the I-H ₂ O ^{$:I^{-}$} ratio due to higher E/N in the
19	declustering region, results in a lower observed sensitivity. Additional work to characterize the
20	effect of humidity on the detection efficiency is necessary to refine the potential of this method
21	for ambient monitoring of HO _{2.}
22	The difficulty of quantitative sampling through an inlet is a significant limitation to the

23 measurement of radicals in the atmosphere. Laboratory experiments were performed to probe the

1	effect of inlet length on sampling of HO ₂ radicals produced in N ₂ . The results are shown in
2	Figure 2a, as the count rate at a given residence time normalized to the count rate at the shortest
3	residence time. This reaction is likely first order in HO ₂ therefore a log-linear fit is the most
4	appropriate representation of the data, however the data has been fit using a linear curve for
5	simplicity. HO_2 is lost at a rate of approximately 0.60 s ⁻¹ in a 6 mm o.d. PFA inlet. Results
6	indicate that the loss of HO_2 is driven not by surface losses, but loss of HO_2 via reaction with
7	residual NO ₂ in the system to produce HO ₂ NO ₂ , as can be observed in the nearly equivalent rate
8	of increase in observed HO ₂ NO ₂ (0.67 s ⁻¹ , Figure 2b). Qualitative observations of H ₂ O ₂ , at m/z
9	161 (I- $H_2O_2^-$), during the same experiment suggest that there is no loss of HO_2 via self-reaction
10	occurring on these timescales.

12 3.1.2 Peroxy Nitric Acid (HO₂NO₂, PNA)

13 Two methods were used in this work for the production of a PNA standard. In the first of these

14 methods, PNA was synthesized using the techniques described in Appelman and Gosztola 1995

15 (Appelman and Gosztola, 1995). Briefly, a nitrite-peroxide solution (NaNO₂ in 30% H₂O₂) is

16 mixed with a peroxide-acid solution (30% H₂O₂ in 70% HClO₄) at -20 °C to produce

17 approximately 1.7 M PNA in H₂O₂. The resulting solution is placed in a glass diffusion cell,

18 (Williams et al., 2000), at a temperature of -20 °C with zero air passed over the headspace to

19 produce a dynamic mixture of PNA. The 20 sccm diffusion source outflow was sampled directly

20 into the inlet flow of the I CIMS. This method of synthesis also results in the production of non-

13

21 negligible amounts of HNO₃ and H₂O₂. While nylon wool can be used to semi-selectively

22 remove HNO₃ from the calibration flow, no method for the selective removal of H₂O₂ was

1 identified. In any case, HNO₃ and H₂O₂ are observed at unique m/z ions and therefore do not

2 interfere with PNA measurement.

Alternatively, PNA can be dynamically generated using the output of the HO₂ source
described in section 2.2.1 (Ulrich et al., 2012). Addition of NO₂ to the output of the HO₂ radical
source results in the production of PNA. Due to the relative simplicity of this technique, photoproduction of PNA was used as the preferred I⁻ CIMS calibration method for the laboratory and
field measurements.

8 Figure 1 shows a difference mass spectrum of HO₂NO₂ detected using an iodide CIMS
9 instrument with a 'cold' inlet (Figure 1b) and an inlet dissociator temperature of 130°C (Figure
10 1c). In both cases, the instrument background was chosen as the ion signal prior to the addition
11 of NO₂. When using a 'cold' inlet, ~30°C, the dominant peak observed is m/z 62 (NO₃⁻),
12 equation 7. An ion signal at m/z 206 (I-HO₂NO₂⁻) is also observed, although to a much lesser

13 extent than m/z 62 (NO₃⁻), equation 6. When an inlet dissociator is used, HO₂NO₂ is observed at

14 m/z 62 (NO₃⁻) and m/z 160 (I-HO₂⁻), where the detection of PNA at m/z 62 (NO₃⁻) results from

15 incomplete thermal dissociation of HO_2NO_2 in the inlet.

- 16
- 17

18 **3.1.3 HONO**

19 Similarly to PNA, HONO can be formed by addition of NO to the output of the HO₂

source described in section 3.1.1. Addition of excess NO to the HO_2 calibration source results in

21 the production of HONO from titration of HO₂, as well as any OH produced in the source, via

22 the following reactions:

$$HO_2 + NO \longrightarrow OH + NO_2 \tag{10}$$

$$OH + NO \longrightarrow HONO$$
 (11)

5	Figure 1d shows the difference mass spectrum of the HONO calibration source, where the
6	instrument background here was chosen as the ion signal prior to the addition of NO. It is clear
7	from the figure that HONO is the only product formed and is detected by I ⁻ CIMS at m/z 174 (I-
8	HONO ⁻). This result is in contrast to the Abida et al. 2011 study which also reports m/z 46 as a
9	minor ion, a difference that can be attributed to the relatively stronger clustering used in our
10	work.
11	This method of HONO production is instantaneous and does not require the period of
12	stabilization that is necessary for acid-salt reaction based sources (Febo et al., 1995). HONO
13	standard production via the reaction of HO ₂ and NO provides a good alternative to previously
14	used I CIMS calibration methods (Roberts et al., 2010).
15	
16	3.2 Dynamic Source Calibration
16 17	3.2 Dynamic Source CalibrationQuantification of PNA and HONO produced using the above-described methods was performed
16 17 18	3.2 Dynamic Source Calibration Quantification of PNA and HONO produced using the above-described methods was performed using the quartz catalysis total NO _y instrument (Wild et al., 2014), described in section 2.2.2.
16 17 18 19	 3.2 Dynamic Source Calibration Quantification of PNA and HONO produced using the above-described methods was performed using the quartz catalysis total NO_y instrument (Wild et al., 2014), described in section 2.2.2. Laboratory experiments indicate that more than 99% of PNA is thermally dissociated above a
16 17 18 19 20	 3.2 Dynamic Source Calibration Quantification of PNA and HONO produced using the above-described methods was performed using the quartz catalysis total NO_y instrument (Wild et al., 2014), described in section 2.2.2. Laboratory experiments indicate that more than 99% of PNA is thermally dissociated above a temperature of 100 °C while HONO decomposition is negligible below 200 °C. The quartz inlet
16 17 18 19 20 21	 3.2 Dynamic Source Calibration Quantification of PNA and HONO produced using the above-described methods was performed using the quartz catalysis total NO_y instrument (Wild et al., 2014), described in section 2.2.2. Laboratory experiments indicate that more than 99% of PNA is thermally dissociated above a temperature of 100 °C while HONO decomposition is negligible below 200 °C. The quartz inlet was operated at gas temperatures of 160°C and 720°C for the measurement of PNA and HONO,

23 quantitative measurement of the PNA or HONO produced in the source.

1 During these experiments the calibration source flow was alternately sampled by the I 2 CIMS and CRD instruments. In order to eliminate any differences in radical reaction times, as a 3 result of inconsistencies in the inlet lengths between the two instruments, the gaseous mixture 4 was passed over glass wool subsequent to addition of NO or NO_2 to terminate the reaction by 5 removing any remaining HO_2 radicals. In this manner the I⁻ CIMS sensitivity is calculated as the 6 ratio of the I CIMS ion signal to the CaRDS measured concentrations. A summary of instrument 7 sensitivities and detections limits (3σ) is included in Table 1. For the calibration data reported in 8 Table 1, the m/z 145 to m/z 127 ratio was approximately 30%.

9 The I⁻ CIMS sensitivity towards HONO at the I-HONO⁻ cluster (m/z 174), reaction (9), was determined to be 1.7 Hz ppbv⁻¹ with a corresponding 3σ instrumental detection limit of 30 10 pptv. The detection limit for HONO is largely limited by the magnitude of the ever-present I-11 12 NO_2^{-} (m/z 173) ion and the low resolution of the quadrupole mass spectrometer used in this 13 work. The HONO sensitivity reported here represents detection with a 25°C inlet, however, the 14 instrument sensitivity was found to be nearly equivalent, within the stated uncertainties, using 15 the heated (130°C) dissociator. 16 The sensitivity towards PNA was determined for detection at the $I-HO_2NO_2^-$ (m/z 206, 17 25° C inlet), I-HO₂⁻ (m/z 160, 130°C inlet), and NO₃⁻ (m/z 62, 25°C inlet) ions, reactions (6) 18 though (8) respectively. The most sensitive method of detection was observed via NO_3^- in a 25°C inlet, 144 Hz pptv⁻¹ with a corresponding 3σ detection limit of 0.7 pptv. Detection at the I-HO₂⁻¹ 19 and I-HO₂NO₂⁻ ions are considerably less sensitive, 2.0 and 0.4 Hz pptv⁻¹ respectively. The 3σ 20 21 detection limits are approximately 40 and 20 pptv for I-HO₂⁻ and I-HO₂NO₂⁻ respectively. 22 While the most sensitive detection of PNA occurs via NO3, there exist several potential interferences that are also observed at that ion, see Table 1 in Wang et al. 2014 (Wang et al., 23

1 2014). Considering the sensitivity to PNA detection, relatively low daytime levels of PNA will 2 result in significant signals at m/z 62. In fact, a recent study has suggested that evidence exists 3 for a large daytime source of N₂O₅ detected via NO₃⁻ ion using I⁻ CIMS measurements (Wang et 4 al., 2014); however, low daytime levels of HO_2NO_2 would also be consistent with observations 5 presented in that work. Unfortunately, detection of PNA at $m/z \ 160 \ (I-HO_2)$ leads to overlap 6 with HO₂ radical detection at that mass. Therefore, we suggest that m/z 206 (I-HO₂NO₂) will 7 yield the most reliable, interference free method of PNA detection for ambient measurements. Increasing the I-H₂O cluster ratio beyond 30% (this study) should improve the instrument 8 9 sensitivity towards detection of PNA at the m/z 206 cluster ion. 10 An indirect calibration was performed for the quantification of the HO₂ source, as no 11 direct HO₂ measurement was readily available. An initial amount of HO₂ radicals are generated 12 and monitored via the I⁻ CIMS operated with a 25°C inlet. A small amount of NO₂ is then added 13 generating PNA with a corresponding reduction in observed Γ CIMS HO₂ signal. The 14 concentration of HO₂ lost by titration is assumed to be equivalent to the amount of PNA 15 produced in the reaction, as the formation of PNA involves the consumption of one HO₂ radical 16 per molecule. Using the previously described PNA calibration method, the sensitivity can be 17 calculated as the ratio of the reduction in the observed I^{-} CIMS I-HO₂⁻ signal to the NO_y 18 measured PNA concentration. The I CIMS detection sensitivity was determined using this method to be 2.6 Hz pptv⁻¹ with a corresponding 3σ detection limit of 20 pptv. As it was not the 19 20 focus of this work, the instrument inlet was not optimized for the sampling of radical species; 21 therefore changes in the inlet design and optimization of the iodide-water cluster distribution in 22 the flow tube could both serve to increase the instrument sensitivity and improve instrument 23 detection limits to levels more appropriate for ambient sampling of HO₂.

4 Uintah Basin Wintertime Ozone Study (UBWOS) Observations

3	Observations of HO_2NO_2 during the UBWOS 2014 study are shown in Figure 3 for the
4	entire duration of the measurement period. Measurements of the I-HO ₂ NO ₂ ⁻ ion using a cold
5	inlet were performed only during the initial and final portion of the study for thirty minutes every
6	hour. Ambient air was sampled through a heated inlet dissociator on alternating 30-minute
7	periods with 'cold' sampling performed on the opposite time periods during the following: Jan.
8	24 – 30 and Feb. 4 -14. For the remainder of the measurements, Jan. 30 - Feb. 4, ambient air was
9	continuously sampled through a heated dissociator. During sampling periods where the inlet
10	dissociator was used PNA was monitored on m/z 160 (I-HO ₂), and during periods of 'cold'
11	sampling via m/z 206 (I-HO ₂ NO ₂). It is clear from the correlation plot inset in Figure 3 that both
12	sampling methods ('cold' and hot) agree reasonably well, slope = 0.93, $R^2 = 0.785$. There does
13	appear to be a positive bias in the m/z 160 observations relative to detection at m/z 206, indicated
14	by a positive intercept and clear disagreement during certain periods shown in the Figure 3
15	timeseries. One possible explanation is the sensitivity at m/z 160 to ambient HO_2 or HO_2
16	generated in the inlet as a product of PNA decomposition. Given the length, temperature, and
17	residence time of the inlet used (20m, 30° C, ~ 4.8 sec), PNA is expected to decompose by
18	approximately 5% prior to sampling. The HO ₂ radicals produced as a result of PNA
19	decomposition would likely be detected via m/z 160 yet remain unobserved on the m/z 206 ion
20	leading to a positive bias in m/z 160 observations. Additionally from laboratory results presented
21	in Figure 2b, we expect there to be production of HO ₂ NO ₂ in the inlet; however, model
22	approximations of the HO ₂ mixing ratio during the 2013 study (15 pptv maximum) yields a

1 maximum formation of HO₂NO₂ in the inlet during peak ozone events of $\sim 1\%$ (Edwards et al.,

2 2014).

3 Aside from the sampling conditions already described, several days of PNA 4 measurements were performed during the 2014 study comparing a short unheated PFA inlet (3m, 5 ambient temperature) and a longer heated inlet (20m, 30°C). An average reduction of 5% is 6 observed in the PNA mixing ratio when sampling through the longer heated inlet as a result of 7 thermal and perhaps surface assisted decomposition. It is important to note that the data 8 presented here are not corrected for these inlet losses and are therefore considered a lower limit 9 on ambient PNA. 10 During UBWOS 2013, the I CIMS was equipped to monitor PAN compounds using a 11 thermal dissociation inlet, 150°C. This was done prior to the laboratory work described in this 12 manuscript and neither m/z 160 nor m/z 206 were monitored during the measurement period. 13 However, the NO_3^{-1} ion (m/z 62) was monitored throughout the entire campaign with observed 14 signals exceeding 2e5 counts per second at times. Using the calibration data obtained from the 15 laboratory portion of this work, an approximate PNA concentration was determined for the 2013 16 UBWOS study, assuming the same NO_3^- (m/z 62) sensitivity as measured during the laboratory 17 calibrations performed after the 2014 study (144 Hz pptv⁻¹), corrected for differences in 18 dissociator temperatures and transmission of HO₂NO₂ through the heated inlet tip used during 19 the 2013 study. The error associated with this method was calculated to be approximately 60%, 20 largely due to corrections applied to account for differences in instrument tuning and additional 21 HO₂NO₂ losses due to the different inlet conditions used during the two years. 22 Figure 4a shows the diurnal average of 2013 I⁻ CIMS observations of PNA for the entire 23 study. While the average diurnal mixing ratio peaks at 0.5 ppbv, mixing ratios up to 1.5 ppbv

1 were observed during the 2013 study and can be explained by the coincidence of high daytime 2 levels of NO₂ with the low temperatures in the Uintah basin. Similarly to the 2014 3 measurements, also included in Figure 4a, PNA reaches a peak after solar maximum ($\sim 15:00$ MST) with a minimum observed throughout the night. Concentrations of HO₂NO₂ observed 4 5 during the 2014 study were significantly lower relative to the 2013 study with a maximum 6 average mixing ratio of 0.1 ppby. During the 2013 and 2014 study, N₂O₅, a nighttime species and 7 potential interference on the NO_3^- ion, was not observed to contribute to the observed daytime 8 signal.

9 The PNA mixing ratios observed during the 2013 and 2014 studies are expected for cold 10 conditions with active photochemistry and sufficient NO_x pollution. Displayed in Figure 5 are 11 the conditions necessary to sustain an equilibrium concentration of 1 ppby PNA with respect to 12 temperature and the mixing ratios of HO₂ and NO₂. The dashed area superimposed on the figure 13 represents the approximate range of conditions encountered during the 2013 study, where HO_2 14 levels were approximated using model results that describe an ozone event observed during the 15 2013 study, described below. The Uintah basin provided a unique atmosphere that promotes the 16 formation of PNA for several reasons (1) a strong inversion during the wintertime allows 17 concentrations to build up in the boundary layer over several day periods (2) low ambient 18 temperatures favoring the formation of HO₂NO₂ over thermal decomposition and (3) radical 19 species propagation, e.g. HO₂ formation, is enhanced due to the active chemistry observed 20 during ozone formation events (Edwards et al., 2014). Lastly, as will be discussed later, the snow 21 surface acts as an important interface serving as both a source and a sink of HO₂NO₂. 22 The conditions encountered in the basin between the 2013 and 2014 season can be used 23 to explain the large difference in the observed PNA ambient mixing ratios, shown in Figure 4a.

1	Mainly during the 2013 study we observed strong inversions over multi-day periods allowing the	
2	buildup of primary and secondary pollutants, a phenomenon driven by snow surface cover and	
3	meteorological conditions encountered during the 2013 study (Ahmadov et al., 2015;Edwards et	
4	al., 2014). In contrast, relatively low snow cover during the 2014 season limited the formation of	
5	multi-day inversions yielding lower ambient mixing ratios of both primary and secondary	
6	pollutants. The resulting combination of lower NOx mixing ratios and higher ambient	
7	temperatures during the 2014 study thereby favored thermal dissociation of HO_2NO_2 and led to	
8	lower ambient mixing ratios than observed during the 2013 study. In addition, limited snow	
9	cover and reduced deposition of NO _y species to the snow surface, as a result of lower ambient	
10	mixing ratios, likely reduced the role of the snow surface as a source of HO_2NO_2 in 2014.	
11	As described previously, the vertical gradient of various species was probed through the	
12	use of a dual inlet system (<u>18.4m</u> and 1m heights) during the UBWOS 2014 study. A comparison	
13	of those measurements for PNA is shown in Figure 4a as diurnal averages of the measurements	
14	where dual inlet switching was applied. Figure 4b shows the result of the difference of the 1m	
15	minus the <u>18.4m</u> PNA measurements where the shaded region represents one standard deviation	
16	of the average for the entire study. Displayed in this fashion, a positive value is an indication that	
17	PNA is larger in the surface coupled layer, characteristic of daytime observations, than aloft with	
18	the opposite indicating a relative enhancement of PNA in the layer decoupled from the surface,	
19	which was typically observed at night. On average, HO_2NO_2 is depleted at the surface relative to	
20	air aloft by approximately 15 pptv with a reversal of nearly the same magnitude observed during	
21	midday. It is important to note that this data represents a 1-hour average over the entire <u>six-week</u>	
22	measurement period, and while the overall magnitude shown here is small observed ΔPNA	
23	values ranged from -150 pptv to 150 pptv.	
	1	

Patrick Veres 6/23/2015 3:39 PM Deleted: 30m

Patrick Veres 6/23/2015 3:39 PM Deleted: 30m

Patrick Veres 6/23/2015 11:59 AM Formatted: Subscript Patrick Veres 6/23/2015 11:59 AM Formatted: Subscript Patrick Veres 6/23/2015 11:47 AM Deleted: six week

1	One interpretation of these results is that deposition of PNA to the snow surface occurs	
2	throughout the night in the Uintah Basin with an emission from the snow surface observed in the	Deleted: there
3	early morning to midday. Through this mechanism snow surface photolysis of PNA could serve	Datrick Vorea 6
4	as an additional daytime surface source of NO_2 and HO_2 . A similar result has been reported by a	Deleted: Addit
5	recent study (Jones et al., 2014), where evidence for surface exchange was observed for HO ₂ NO ₂	Deleted: he
6	and HNO3 during the Antarctic winter. However, previous studied have highlighted the	Deleted: precur Patrick Veres 6
7	complexities of photochemistry chemistry occurring at or directly above snow surfaces (Chen et	Deleted: Jones
8	al., 2007; Chen et al., 2001), which complicates the interpretation of these observations.	Patrick Voros 6
9	Specifically, previous studies have shown enhancements of reactive species integral to	Deleted: Unfor not measured dire
10	HOx and NOx budgets at snow surfaces, e.g. HONO, CH2O, H2O2 (Hutterli et al., 1999;Hutterli	validation of this not possible.
11	et al., 2001;Honrath et al., 1999;Ridley et al., 2000;Zhou et al., 2001). Photolysis of these	Formatted: Ind
12	HO_x/NO_x precursors at or above the snow surface or a direct emission of NO_x could result in a	
13	net apparent surface source of HO_2NO_2 as it would shift the gas phase equilibrium, reactions (1)	
14	and (2), towards the formation of HO_2NO_2 . The net effect of this process on the HO_x budget at	
15	the snow surface is difficult to quantify without simultaneous observations of the other dominant	
16	HO _x precursors and an understanding of their chemical fates. Therefore further validation of our	Patrick Veres 6
17	interpretation, a bidirectional flux of HO ₂ NO ₂ from the snow surface, awaits direct flux	Formatted: Su
18	measurements of HO ₂ NO ₂ over these highly polluted snow surfaces.	
19	Regardless of the underlying mechanisms, the dynamics causing the observed vertical	Patrick Veres 6
20	distribution in PNA observations, whether mixing or deposition in origin, can have a	Deleted: In any
21	measureable impact on the ozone formation potential in the Uintah basin, To investigate the	Patrick Veres 6. Deleted: he dyn
22	effects, a chemical box model, based on the MCM v3.2 chemistry scheme, has been developed to	differences in PN mixing or deposit measureable imp
23	describe observed ozone production during a wintertime ozone pollution episode during	potential in the U discussed below

Patrick Veres 6/23/2015 12:38 PM Deleted: there is

Patrick Veres 6/23/2015 11:47 AM Deleted: Additionally t Patrick Veres 6/23/2015 11:48 AM Deleted: he Patrick Veres 6/23/2015 11:48 AM Deleted: precursors, such as Patrick Veres 6/23/2015 12:40 PM Deleted: Jones et al. 2014

Patrick Veres 6/23/2015 12:41 PM Deleted: Unfortunately, fluxes of PNA were not measured directly in this study so further validation of this interpretation of the data is not possible. Patrick Veres 6/23/2015 1:26 PM

Formatted: Indent: First line: 0.5"

Patrick Veres 6/23/2015 1:26 PM Formatted: Subscript

Patrick Veres 6/23/2015 11:49 AM Deleted: In any case, t

Patrick Veres 6/23/2015 1:31 PM

Deleted: he dynamics causing these differences in PNA observations, whether mixing or deposition in origin, have a measureable impact on the ozone formation potential in the Uintah basin as will be discussed below

Patrick Veres 6/23/2015 11:50 AM Deleted: A

UBWOS 2013 (Edwards et al., 2014). This 0-D model contains a near-explicit oxidation
 mechanism for 32 observed VOC and oxidized VOC, and is constrained using constant
 emissions of primary species (VOC and NO), with the concentrations of all other species
 calculated by the chemistry scheme. Physical loss processes, such as mixing and deposition, are
 represented via a bimodal 1st order loss process for all species, the rate of which has been
 optimized based on boundary layer height observations and the concentrations of long lived
 species, e.g. methane.

8 Figure 6a and 6b display the base model calculation of PNA and ozone reported in the 9 Edwards et al. 2014 study. While there is reasonable quantitative agreement between the model 10 and measurement daily maxima, there is a temporal shift in PNA measurements relative to model 11 predictions. While the reasons for this are as of yet unknown, possible explanations include 12 underestimation in modeled NO, additional daytime PNA loss mechanisms, or issues with the 13 simple parameterization of mixing used in the model. The latter seems unlikely as the model 14 reproduces the observed diurnal variation in ozone relatively well. As discussed above, there is 15 also evidence that PNA is lost to the snow surface, though the temporal trend in deposition 16 implied from the gradient measurements does not suggest a relatively higher rate in the morning 17 to early afternoon than in the evening, rather there is a possible indication of a mid-day snow 18 surface source. Nitrogen oxides within the model are parameterized using a constant source of 19 NO, with the partitioning of all nitrogen oxides calculated by the chemistry scheme. A 20 quantitative comparison of NO observations to modeled values suggests that the model does 21 typically underestimate the NO concentration throughout the morning. This underestimation 22 would lead to an underestimation of the loss of HO2 via reaction with NO, thereby slowing the 23 formation rate leading to an over-prediction of PNA. This process would have a particular

1 impact in the morning hours where observed NO is relatively large and the HO₂ source is

2 limited.

3 Deposition of PNA has the potential to result in a net loss of HO_x and thus an increase in NO (Grannas et al., 2007), due to reduced titration to NO₂, which would have an overall effect 4 5 on ozone formation potential. The sensitivity of ozone production in the UBWOS 2013 box 6 model to changes in the lifetime of PNA with respect to deposition (t_{PNA}) was investigated and the results of these tests are displayed in Figure 6. In addition to the 1st order physical loss term 7 applied to all species, predominantly representing losses due to mixing (Edwards et al., 2014), an 8 9 additional 1st order loss term for PNA was added to the model scheme to represent deposition. 10 Calculations with a lifetime for PNA with respect to deposition (tPNA) of 1-hour yield 11 approximately a 12% reduction in daily max ozone from the base case where no PNA deposition 12 was included. It is not possible to determine the observed lifetime of PNA with respect to deposition to the snow surface using data from the UBWOS 2013 or 2014 studies; however the 13 14 measurements and model suggest that surface deposition is occurring with a potentially 15 measureable effect on the ozone production in the Uintah basin. 16 The post-depositional fate of PNA is also important as this scenario assumes that the loss 17 of PNA to the snow surface is irreversible and thus a net HO_x and NO_x sink. Subsequent snow 18 chemistry resulting in storage and volatilization of HO_x (Jones et al., 2014) or NO_x, has the 19 potential to reduce the magnitude of the effects observed in Figure 6b. If PNA dissolves in an 20 aqueous solution, such as a quasi-liquid layer on the snow surface in the Uintah basin, it can 21 undergo the following dissociation and ionic reactions (Logager and Sehested, 1993; Zhu et al., 22 1993;Goldstein et al., 2005):

23

15 Measurements of snow surface nitrite made during the 2013 and 2014 study are shown in 16 Figure 7(a) and (b), respectively, along with observations of PNA. Two pollution events 17 observed in 2013 were separated by a cleanout event where additional precipitation (indicated by 18 the blue shaded region in figure 7a) accompanied by higher winds and unstable conditions 19 ventilated the basin. The nitrite content of the snow surface generally increases throughout the 20 first event at a rate proportional to the daily increase in PNA mixing ratio until fresh snow is 21 added and nitrite levels drop. Throughout the second event, nitrite levels in the snow surface 22 again build up as ambient PNA levels also increase. We note here that any HO₂NO₂ dissolved in

uptake, where chemical loss to NO_2^- is negligible (Zhang et al., 1997).

14

1 the snow can form NO_2^{-} with efficiencies as high as 56% depending on pH (Goldstein et al.,

2 2005).

3 A more detailed look at snow nitrite content shows a generally observed decrease in 4 surface layer concentrations throughout the day, Figure 7c, a result that is consistent with 5 reversible uptake of nitrite. This daytime depletion of snow nitrite could be evidence of the 6 reversible uptake of PNA, interpreted from data shown in Figure 4a, or an indication of the 7 formation and subsequent release of other nitrogen containing species, such as HONO or NO_x, 8 from the snow surface. It is interesting to note that the highest measurements of snow nitrite in a 9 given day typically occur prior to the build up of ambient PNA. This result is consistent with 10 nocturnal uptake of PNA or HONO to the snow surface, as shown in Figure 4b; however, no 11 evidence of this nighttime enrichment is available, as nighttime measurements of snow surface 12 nitrite were not made. Additionally, the extent to which reactions (13) and (15) were occurring, 13 and therefore PNA contributing to snow nitrite content, cannot be approximated as the pH of the 14 snow surface was not measured during this study. Furthermore, while the deposition and 15 potential volatilization of HO₂NO₂ can contribute to the net flux of nitrite at the snow surface, other species such as HNO₃ and HONO are also known to deposit to snow surfaces and should 16 also be considered as part of the net surface nitrogen budget. 17 18 Improved measurements of the deposition velocity on snow surfaces of varying acidities and the chemical fate of PNA after deposition is necessary to improve our understanding of the 19 20 impacts of PNA deposition to HO_x and NO_x budgets. As our model results indicate the post 21 depositional fate of PNA can have a non-negligible impact on ozone formation potential, 22 particularly in cold regions, such as the Uintah basin, where the lifetime of PNA is sufficiently 23 long such that deposition becomes a dominant sink. Considering the similar surface chemistry

Patrick Veres 6/24/2015 10:06 AM Deleted: shown in

Patrick Veres 6/23/2015 11:57 AM

Deleted: While the deposition and potential volatilization of HO2NO2 can contribute to the net flux of nitrite at the snow surface other species such as HNO3 and HONO are also known to deposit to snow surfaces and should also be considered as part of the net surface nitrogen budget (Reference). Furthermore, release of NOx from the snow surface in the form of NO2 could also result in a net apparent surface source as it would shift the gas phase equilibrium, reactions (1) and (2), towards the formation of HO₂NO₂. In this case, a surface depletion of HO2 radicals would be expected, however; measurements of HO2 were not made during the 2013 and 2014 studies. ... [1]

Patrick Veres 6/23/2015 11:42 AM Deleted: T

1 and influence on radical budgets for atmospheric HONO (VandenBoer et al., 2015),

2 simultaneous measurement must be performed to understand the tropospheric fate of PNA and

HONO species in cold regions and the extent of their involvement in tropospheric HO_x and NO_x
budgets.

- 5
- 6

7 4 Conclusions

8 Laboratory studies have been conducted to demonstrate the utility of I CIMS as a method 9 for the quantitative detection of HO₂, PNA, and HONO. Methods for the production and 10 calibration of on-line laboratory standards for HO₂, PNA, and HONO have been developed and 11 described above. The I CIMS technique offers an unambiguous method with sufficient 12 sensitivity for the detection of PNA in mid latitude tropospheric regions, an area with a general 13 lack of direct observations. Additionally, while it is not fully explored here, the application of I 14 CIMS for the direct detection of HO₂ has significant potential for use in future laboratory, 15 chamber, and ambient measurements. 16 The I CIMS instrument was deployed during the 2013 and 2014 UBWOS studies in the 17 Uintah basin, UT during several wintertime ozone events. Field-based experiments illustrate that 18 as a result of the thermal instability of PNA and the potential for formation within inlets, care 19 must be taken to both characterize sampling inlets and minimize residence times in order to 20 perform accurate quantitative measurements. PNA vertical gradient as well as correlation with 21 snow surface nitrite content suggests that deposition, dissociation and ionic reactions of PNA 22 could serve as a large surface nitrite source in the Uintah basin.

1	Comparison of the 2013 PNA observations to a chemically explicit box model yields
2	quantitative agreement of daily maximum mixing ratios, but with disagreement in the diurnal
3	cycle in late afternoon. Results of this comparison suggest that increased PNA deposition rates
4	lead to an overall decrease in the ozone production potential in the Uintah basin. This result is
5	dependent on the post-depositional fate of PNA, and more studies aimed at improving our
6	understanding of the deposited HO_x and NO_x and photochemistry of NO_y species on snow
7	surfaces are necessary to model this chemistry correctly. Results of this study demonstrate a
8	clear need for further studies of eddy covariance fluxes of PNA, and HONO, from snow surfaces
9	and investigation of subsequent chemistry to improve our understanding of these species as both
10	a source and sink of HO_x and NO_x .

12 Acknowledgements

- 13 The Uintah Basin Winter Ozone Studies were a joint project led and coordinated by the Utah
- 14 Department of Environmental Quality (UDEQ) and supported by the Uintah Impact Mitigation
- 15 Special Service District (UIMSSD), the Bureau of Land Management (BLM), the Environmental
- 16 Protection Agency (EPA) and Utah State University. This work was funded in part by the
- 17 Western Energy Alliance, and NOAA's Atmospheric Chemistry, Climate and Carbon Cycle
- 18 program. We thank Questar Energy Products for site preparation and support.

1 References

- 2 Abida, O., Mielke, L. H., and Osthoff, H. D.: Observation of gas-phase peroxynitrous and
- 3 peroxynitric acid during the photolysis of nitrate in acidified frozen solutions, Chem. Phys.
- 4 Lett., 511, 187-192, 10.1016/j.cplett.2011.06.055, 2011.
- 5 Ahmadov, R., McKeen, S., Trainer, M., Banta, R., Brewer, A., Brown, S., Edwards, P. M., de
- 6 Gouw, J. A., Frost, G. J., Gilman, J., Helmig, D., Johnson, B., Karion, A., Koss, A., Langford, A.,
- 7 Lerner, B., Olson, J., Oltmans, S., Peischl, J., Petron, G., Pichugina, Y., Roberts, J. M., Ryerson,
- 8 T., Schnell, R., Senff, C., Sweeney, C., Thompson, C., Veres, P. R., Warneke, C., Wild, R.,
- 9 Williams, E. J., Yuan, B., and Zamora, R.: Understanding high wintertime ozone pollution
- events in an oil- and natural gas-producing region of the western US, Atmos. Chem. Phys.,
- 11 15, 411-429, 10.5194/acp-15-411-2015, 2015.
- 12 Appelman, E. H., and Gosztola, D. J.: Aqueous Peroxynitric Acid (HOONO2): A Novel
- Synthesis and Some Chemical and Spectroscopic Properties, Inorganic Chemistry, 34, 4,1995.
- 15 IUPAC Task Group on Atmosperic Chemical Kinetic Data Evaluation: <u>http://iupac.pole-</u>
 <u>ether.fr</u>.
- 17 , access: December 2014, 2004.
- 18 Behnke, W., George, C., Scheer, V., and Zetzsch, C.: Production and decay of ClNO2, from the
- 19 reaction of gaseous N2O5 with NaCl solution: Bulk and aerosol experiments, J. Geophys.
- 20 Res.-Atmos., 102, 3795-3804, 10.1029/96jd03057, 1997.
- 21 Brune, W. H., Tan, D., Faloona, I. F., Jaegle, L., Jacob, D. J., Heikes, B. G., Snow, J., Kondo, Y.,
- 22 Shetter, R., Sachse, G. W., Anderson, B., Gregory, G. L., Vay, S., Singh, H. B., Davis, D. D.,
- 23 Crawford, J. H., and Blake, D. R.: OH and HO2 chemistry in the North Atlantic free
- 24 troposphere, Geophys. Res. Lett., 26, 3077-3080, 10.1029/1999gl900549, 1999.
- 25 Carpenter, L. J., Green, T. J., Mills, G. P., Bauguitte, S., Penkett, S. A., Zanis, P., Schuepbach, E.,
- 26 Schmidbauer, N., Monks, P. S., and Zellweger, C.: Oxidized nitrogen and ozone production
- efficiencies in the springtime free troposphere over the Alps, J. Geophys. Res.-Atmos., 105,
 14547-14559, 10.1029/2000jd900002, 2000.
- 29 Chen, G., Davis, D., Crawford, J., Nowak, J. B., Eisele, F., Mauldin, R. L., Tanner, D., Buhr, M.,
- 30 Shetter, R., Lefer, B., Arimoto, R., Hogan, A., and Blake, D.: An investigation of South Pole
- 31 HOx chemistry: Comparison of model results with ISCAT observations, Geophys. Res. Lett.,
- 32 28, 3633-3636, 10.1029/2001gl013158, 2001.
- 33 Chen, G., Huey, L. G., Crawford, J. H., Olson, J. R., Hutterli, M. A., Sjostedt, S., Tanner, D., Dibb,
- J., Lefer, B., Blake, N., Davis, D., and Stohl, A.: An assessment of the polar HOx photochemical
 budget based on 2003 Summit Greenland field observations, Atmos. Environ., 41, 7806-
- 36 7820, 10.1016/j.atmosenv.2007.06.014, 2007.
- 37 Davis, D., Nowak, J. B., Chen, G., Buhr, M., Arimoto, R., Hogan, A., Eisele, F., Mauldin, L.,
- 38 Tanner, D., Shetter, R., Lefer, B., and McMurry, P.: Unexpected high levels of NO observed at
- 39 South Pole, Geophys. Res. Lett., 28, 3625-3628, 10.1029/2000gl012584, 2001.
- 40 Day, D. A., Wooldridge, P. J., Dillon, M. B., Thornton, J. A., and Cohen, R. C.: A thermal
- 41 dissociation laser-induced fluorescence instrument for in situ detection of NO2, peroxy
- 42 nitrates, alkyl nitrates, and HNO3, J. Geophys. Res.-Atmos., 107, 10.1029/2001jd000779,

43 2002.



- 1 DeMore, W. B., Sander, S. P., Glolden, G. M., Hampson, R. F., Kurylo, M.J., Howeard, C. J.,
- 2 Ravishankara, A. R., Kolb, C. E., and Molina, M. J.: Chemical kinetics and photochemical data
- 3 for use in stratospheric modeling, Propulsion Laboratory, Pasadena, CA, 1997.
- 4 Dusanter, S., Vimal, D., and Stevens, P. S.: Technical note: Measuring tropospheric OH and
- 5 HO2 by laser-induced fluorescence at low pressure. A comparison of calibration
- 6 techniques, Atmos. Chem. Phys., 8, 321-340, 2008.
- 7 Edwards, P. M., Brown, S. S., Roberts, J. M., Ahmadov, R., Banta, R. M., deGouw, J. A., Dube, W.
- P., Field, R. A., Flynn, J. H., Gilman, J. B., Graus, M., Helmig, D., Koss, A., Langford, A. O., Lefer, 8
- 9 B. L., Lerner, B. M., Li, R., Li, S. M., McKeen, S. A., Murphy, S. M., Parrish, D. D., Senff, C. J.,
- 10 Soltis, J., Stutz, J., Sweeney, C., Thompson, C. R., Trainer, M. K., Tsai, C., Veres, P. R.,
- 11 Washenfelder, R. A., Warneke, C., Wild, R. J., Young, C. J., Yuan, B., and Zamora, R.: High
- 12 winter ozone pollution from carbonyl photolysis in an oil and gas basin, Nature, 514, 351-+, 13 10.1038/nature13767, 2014.
- 14 Faloona, I., Tan, D., Brune, W. H., Jaegle, L., Jacob, D. J., Kondo, Y., Koike, M., Chatfield, R.,
- 15 Pueschel, R., Ferry, G., Sachse, G., Vay, S., Anderson, B., Hannon, J., and Fuelberg, H.:
- 16 Observations of HOx and its relationship with NOx in the upper troposphere during SONEX,
- 17 J. Geophys. Res.-Atmos., 105, 3771-3783, 10.1029/1999jd900914, 2000.
- 18 Febo, A., Perrino, C., Gherardi, M., and Sparapani, R.: Evaluation of a high-purity and high-
- stability continuous generation system for nitrous acid, Environ. Sci. Technol., 29, 2390-19 20 2395.1995.
- 21 Gierczak, T., Jimenez, E., Riffault, V., Burkholder, J. B., and Ravishankara, A. R.: Thermal
- 22 decomposition of HO2NO2 (peroxynitric acid, PNA): Rate coefficient and determination of
- 23 the enthalpy of formation, Journal of Physical Chemistry A, 109, 586-596,
- 24 10.1021/jp046632f, 2005.
- 25 Goldstein, S., Lind, J., and Merenyi, G.: Chemistry of peroxynitrites as compared to
- 26 peroxynitrates, Chem. Rev., 105, 2457-2470, 10.1021/cr0307087, 2005.
- Grannas, A. M., Jones, A. E., Dibb, J., Ammann, M., Anastasio, C., Beine, H. J., Bergin, M., 27
- Bottenheim, J., Boxe, C. S., Carver, G., Chen, G., Crawford, J. H., Domine, F., Frey, M. M., 28
- 29 Guzman, M. I., Heard, D. E., Helmig, D., Hoffmann, M. R., Honrath, R. E., Huey, L. G., Hutterli,
- 30 M., Jacobi, H. W., Klan, P., Lefer, B., McConnell, J., Plane, J., Sander, R., Savarino, J., Shepson, P.
- 31 B., Simpson, W. R., Sodeau, J. R., von Glasow, R., Weller, R., Wolff, E. W., and Zhu, T.: An
- 32 overview of snow photochemistry: evidence, mechanisms and impacts, Atmos. Chem. Phys., 33 7,4329-4373,2007.
- 34 Honrath, R. E., Peterson, M. C., Guo, S., Dibb, J. E., Shepson, P. B., and Campbell, B.: Evidence
- 35 of NOx production within or upon ice particles in the Greenland snowpack, Geophys. Res. 36 Lett., 26, 695-698, 10.1029/1999gl900077, 1999.
- 37 Huev, L. G., Villalta, P. W., Dunlea, E. I., Hanson, D. R., and Howard, C. J.: Reactions of CF30-
- with atmospheric trace gases, Journal of Physical Chemistry, 100, 190-194, 38
- 39 10.1021/jp951928u, 1996.
- 40 Huey, L. G., Tanner, D. J., Slusher, D. L., Dibb, J. E., Arimoto, R., Chen, G., Davis, D., Buhr, M. P.,
- 41 Nowak, J. B., Mauldin, R. L., Eisele, F. L., and Kosciuch, E.: CIMS measurements of HNO3 and
- SO2 at the South Pole during ISCAT 2000, Atmos. Environ., 38, 5411-5421, 42
- 43 10.1016/j.atmosenv.2004.04.037, 2004.
- Huey, L. G.: Measurement of trace atmospheric species by chemical ionization mass 44
- 45 spectrometry: Speciation of reactive nitrogen and future directions, Mass Spec. Rev., 26,
- 46 166-184, 10.1002/mas.20118, 2007.

- 1 Hutterli, M. A., Rothlisberger, R., and Bales, R. C.: Atmosphere-to-snow-to-firn transfer
- 2 studies of HCHO at Summit, Greenland, Geophys. Res. Lett., 26, 1691-1694,
- 3 10.1029/1999gl900327, 1999.
- 4 Hutterli, M. A., McConnell, J. R., Stewart, R. W., Jacobi, H. W., and Bales, R. C.: Impact of
- 5 temperature-driven cycling of hydrogen peroxide (H2O2) between air and snow on the
- 6 planetary boundary layer, J. Geophys. Res.-Atmos., 106, 15395-15404,
- 7 10.1029/2001jd900102, 2001.
- 8 Jimenez, E., Gierczak, T., Stark, H., Burkholder, J. B., and Ravishankara, A. R.: Reaction of OH
- 9 with HO2NO2 (peroxynitric acid): Rate coefficients between 218 and 335 K and product
- 10 yields at 298 K, Journal of Physical Chemistry A, 108, 1139-1149, 10.1021/jp0363489,
- 11 2004.
- 12 Jimenez, E., Gierczak, T., Stark, H., Burkholder, J. B., and Ravishankara, A. R.: Quantum yields
- of OH, HO2 and NO3 in the UV photolysis of HO2NO2, Phys. Chem. Chem. Phys., 7, 342-348,
 10.1039/b413429i, 2005.
- 14 10.1039/0413429J, 2005.
- 15 Jones, A. E., Brough, N., Anderson, P. S., and Wolff, E. W.: H02N02 and HN03 in the coastal
- 16 Antarctic winter night: a "lab-in-the-field" experiment, Atmos. Chem. Phys., 14, 11843-
- 17 11851, 10.5194/acp-14-11843-2014, 2014.
- 18 Keim, C., Liu, G. Y., Blom, C. E., Fischer, H., Gulde, T., Hopfner, M., Piesch, C., Ravegnani, F.,
- 19 Roiger, A., Schlager, H., and Sitnikov, N.: Vertical profile of peroxyacetyl nitrate (PAN) from
- 20 MIPAS-STR measurements over Brazil in February 2005 and its contribution to tropical UT
- 21 NO(y) partitioning, Atmos. Chem. Phys., 8, 4891-4902, 2008.
- 22 Kim, S., Huey, L. G., Stickel, R. E., Tanner, D. J., Crawford, J. H., Olson, J. R., Chen, G., Brune, W.
- 23 H., Ren, X., Lesher, R., Wooldridge, P. J., Bertram, T. H., Perring, A., Cohen, R. C., Lefer, B. L.,
- 24 Shetter, R. E., Avery, M., Diskin, G., and Sokolik, I.: Measurement of H02N02 in the free
- 25 troposphere during the intercontinental chemical transport experiment North America
- 26 2004, J. Geophys. Res.-Atmos., 112, 10.1029/2006jd007676, 2007.
- 27 Lee, B. H., Lopez-Hilfiker, F. D., Mohr, C., Kurten, T., Worsnop, D. R., and Thornton, J. A.: An
- 28 Iodide-Adduct High-Resolution Time-of-Flight Chemical-Ionization Mass Spectrometer:
- Application to Atmospheric Inorganic and Organic Compounds, Environ. Sci. Technol., 48,
 6309-6317, 10.1021/es500362a, 2014.
- 31 Li, Z. J., Friedl, R. R., Moore, S. B., and Sander, S. P.: Interaction of peroxynitric acid with solid
- 32 H2O ice, J. Geophys. Res.-Atmos., 101, 6795-6802, 10.1029/96jd00065, 1996.
- Logager, T., and Sehested, K.: Formation and decay of peroxtnitric acid A pulse-radiolysis
 study Journal of Physical Chemistry, 97, 10047-10052, 1993.
- Lyman, S., and Shorthill, H.: Final Report: 2012 Uintah Basin Winter Ozone & Air Quality
- 36 Study, Utah State University, 281, 2012.
- 37 Murphy, J. G., Thornton, J. A., Wooldridge, P. J., Day, D. A., Rosen, R. S., Cantrell, C., Shetter, R.
- E., Lefer, B., and Cohen, R. C.: Measurements of the sum of HO2NO2 and CH3O2NO2 in the
- remote troposphere, Atmos. Chem. Phys., 4, 377-384, 2004.
- 40 Niki, H., Maker, P. D., Savage, C. M., and Breitenbach, L. P.: Fourier-Transform IR
- 41 spectroscopy observation of pernitric acid formed via HOO + NO2 -> HOONO2, Chem. Phys.
- 42 Lett., 45, 564-566, 10.1016/0009-2614(77)80090-0, 1977.
- 43 Ridley, B., Walega, J., Montzka, D., Grahek, F., Atlas, E., Flocke, F., Stroud, V., Deary, J., Gallant,
- 44 A., Boudries, H., Bottenheim, J., Anlauf, K., Worthy, D., Sumner, A. L., Splawn, B., and
- 45 Shepson, P.: Is the Arctic surface layer a source and sink of NOx in winter/spring?, J. Atmos.
- 46 Chem., 36, 1-22, 10.1023/a:1006301029874, 2000.

- 1 Rinsland, C. P., Zander, R., Farmer, C. B., Norton, R. H., Brown, L. R., Russell, J. M., and Park, J.
- 2 H.: Evidence for the presence of the 802.7 cm-1 band Q branch of H02N02 in high-
- 3 resolution solar absorption spectra of the stratosphere, Geophys. Res. Lett., 13, 761-764,
- 4 10.1029/GL013i008p00761, 1986.
- 5 Rinsland, C. P., Gunson, M. R., Salawitch, R. J., Michelsen, H. A., Zander, R., Newchurch, M. J.,
- 6 Abbas, M. M., Abrams, M. C., Manney, G. L., Chang, A. Y., Irion, F. W., Goldman, A., and
- 7 Mahieu, E.: ATMOS/ATLAS-3 measurements of stratospheric chlorine and reactive nitrogen
- 8 partitioning inside and outside the November 1994 Antarctic vortex, Geophys. Res. Lett.,
- 9 23, 2365-2368, 10.1029/96gl01474, 1996.
- 10 Roberts, J. M., Veres, P., Warneke, C., Neuman, J. A., Washenfelder, R. A., Brown, S. S.,
- 11 Baasandorj, M., Burkholder, J. B., Burling, I. R., Johnson, T. J., Yokelson, R. J., and de Gouw, J.:
- 12 Measurement of HONO, HNCO, and other inorganic acids by negative-ion proton-transfer
- 13 chemical-ionization mass spectrometry (NI-PT-CIMS): application to biomass burning
- 14 emissions, Atmos. Meas. Tech., 3, 981-990, 10.5194/amt-3-981-2010, 2010.
- 15 Roehl, C. M., Nizkorodov, S. A., Zhang, H., Blake, G. A., and Wennberg, P. O.:
- Photodissociation of peroxynitric acid in the near-IR, Journal of Physical Chemistry A, 106,
 3766-3772, 10.1021/jp013536v, 2002.
- 18 Salawitch, R. J., Wennberg, P. O., Toon, G. C., Sen, B., and Blavier, J. F.: Near IR photolysis of
- 19 HO2NO2: Implications for HOx, Geophys. Res. Lett., 29, 10.1029/2002gl015006, 2002.
- 20 Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies:
- 21 http://jpldataeval.jpl.nana.gov, access: 10-6, 2011.
- 22 Sen, B., Toon, G. C., Osterman, G. B., Blavier, J. F., Margitan, J. J., Salawitch, R. J., and Yue, G. K.:
- Measurements of reactive nitrogen in the stratosphere, J. Geophys. Res.-Atmos., 103, 3571 3585, 10.1029/97id02468, 1998.
- 25 Singh, H. B., Brune, W. H., Crawford, J. H., Jacob, D. J., and Russell, P. B.: Overview of the
- summer 2004 intercontinental chemical transport experiment North America (INTEX-A),
 J. Geophys. Res.-Atmos., 111, 10.1029/2006jd007905, 2006.
- 28 Singh, H. B., Salas, L., Herlth, D., Kolyer, R., Czech, E., Avery, M., Crawford, J. H., Pierce, R. B.,
- 29 Sachse, G. W., Blake, D. R., Cohen, R. C., Bertram, T. H., Perring, A., Wooldridge, P. J., Dibb, J.,
- 30 Huey, G., Hudman, R. C., Turquety, S., Emmons, L. K., Flocke, F., Tang, Y., Carmichael, G. R.,
- and Horowitz, L. W.: Reactive nitrogen distribution and partitioning in the North American
- 32 troposphere and lowermost stratosphere, J. Geophys. Res.-Atmos., 112,
- 33 10.1029/2006jd007664, 2007.
- 34 Slusher, D. L., Pitteri, S. J., Haman, B. J., Tanner, D. J., and Huey, L. G.: A chemical ionization
- 35 technique for measurement of pernitric acid in the upper troposphere and the polar
- 36 boundary layer, Geophys. Res. Lett., 28, 3875-3878, 10.1029/2001gl013443, 2001.
- 37 Slusher, D. L., Huey, L. G., Tanner, D. J., Chen, G., Davis, D. D., Buhr, M., Nowak, J. B., Eisele, F.
- 38 L., Kosciuch, E., Mauldin, R. L., Lefer, B. L., Shetter, R. E., and Dibb, J. E.: Measurements of
- 39 pernitric acid at the South Pole during ISCAT 2000, Geophys. Res. Lett., 29,
- 40 10.1029/2002gl015703, 2002.
- 41 Slusher, D. L., Huey, L. G., Tanner, D. J., Flocke, F. M., and Roberts, J. M.: A thermal
- 42 dissociation-chemical ionization mass spectrometry (TD-CIMS) technique for the
- 43 simultaneous measurement of peroxyacyl nitrates and dintrogen pentoxide, J. Geophys.
- 44 Res.-Atmos., 109, D19315, doi:10.1029/2004JD004670, 2004.
- 45 Slusher, D. L., Neff, W. D., Kim, S., Huey, L. G., Wang, Y., Zeng, T., Tanner, D. J., Blake, D. R.,
- 46 Beyersdorf, A., Lefer, B. L., Crawford, J. H., Eisele, F. L., Mauldin, R. L., Kosciuch, E., Buhr, M.
 - 32

- 1 P., Wallace, H. W., and Davis, D. D.: Atmospheric chemistry results from the ANTCI 2005
- 2 Antarctic plateau airborne study, J. Geophys. Res.-Atmos., 115, 10.1029/2009jd012605,
- 3 2010.
- 4 Spencer, K. M., McCabe, D. C., Crounse, J. D., Olson, J. R., Crawford, J. H., Weinheimer, A. J.,
- 5 Knapp, D. J., Montzka, D. D., Cantrell, C. A., Hornbrook, R. S., Mauldin, R. L., and Wennberg, P.
- 6 O.: Inferring ozone production in an urban atmosphere using measurements of
- 7 peroxynitric acid, Atmos. Chem. Phys., 9, 3697-3707, 2009.
- 8 Stark, H., Brown, S. S., Burkholder, J. B., Aldener, M., Riffault, V., Gierczak, T., and
- 9 Ravishankara, A. R.: Overtone dissociation of peroxynitric acid (HO2NO2): Absorption
- 10 cross sections and photolysis products, Journal of Physical Chemistry A, 112, 9296-9303,
- 11 10.1021/jp802259z, 2008.
- 12 Final Report 2013 Uinta Basin Winter Ozone Study:
- http://www.deq.utah.gov/locations/U/uintahbasin/studies/UBOS-2013.htm, access: July,
 2014.
- 15 Ulrich, T., Ammann, M., Leutwyler, S., and Bartels-Rausch, T.: The adsorption of
- 16 peroxynitric acid on ice between 230 K and 253 K, Atmos. Chem. Phys., 12, 1833-1845,
- 17 10.5194/acp-12-1833-2012, 2012.
- 18 VandenBoer, T. C., Young, C. J., Talukdar, R. K., Markovic, M. Z., Brown, S. S., Roberts, J. M.,
- 19 and Murphy, J. G.: Nocturnal loss and daytime source of nitrous acid through reactive
- 20 uptake and displacement, Nature Geosci, 8, 55-60, 10.1038/ngeo2298
- http://www.nature.com/ngeo/journal/v8/n1/abs/ngeo2298.html supplementary information, 2015.
- 23 Wang, X., Wang, T., Yan, C., Tham, Y. J., Xue, L., Xu, Z., and Zha, Q.: Large daytime signals of
- 24 N2O5 and NO3 inferred at 62 amu in a TD-CIMS: chemical interference or a real
- 25 atmospheric phenomenon?, Atmos. Meas. Tech., 7, 1-12, 10.5194/amt-7-1-2014, 2014.
- 26 Wennberg, P. O., Salawitch, R. J., Donaldson, D. J., Hanisco, T. F., Lanzendorf, E. J., Perkins, K.
- 27 K., Lloyd, S. A., Vaida, V., Gao, R. S., Hintsa, E. J., Cohen, R. C., Swartz, W. H., Kusterer, T. L.,
- and Anderson, D. E.: Twilight observations suggest unknown sources of HOx, Geophys. Res.
 Lett., 26, 1373-1376, 10.1029/1999gl900255, 1999.
- 30 Wild, R., Edwards, P., Dube, W. P., Baumann, K., Edgerton, E. S., Quinn, P. O., Roberts, J. M.,
- 31 Rollins, A. W., Veres, P. R., Warneke, C., WIlliams, E. J., Yuan, B., and Brown, S.: A
- 32 Measurement of Total Reactive Nitrogen, NOy, together with NO2, NO, and O3 via Cavity
- 33 Ring-down Spectroscopy., In Preparation, 2014.
- 34 Williams, J., Roberts, J. M., Bertman, S. B., Stroud, C. A., Fehsenfeld, F. C., Baumann, K., Buhr,
- 35 M. P., Knapp, K., Murphy, P. C., Nowick, M., and Williams, E. J.: A method for the airborne
- 36 measurement of PAN, PPN, and MPAN, J. Geophys. Res.-Atmos., 105, 28943-28960, 2000.
- 37 Wooldridge, P. J., Perring, A. E., Bertram, T. H., Flocke, F. M., Roberts, J. M., Singh, H. B., Huey,
- 38 L. G., Thornton, J. A., Wolfe, G. M., Murphy, J. G., Fry, J. L., Rollins, A. W., LaFranchi, B. W., and
- 39 Cohen, R. C.: Total Peroxy Nitrates (Sigma PNs) in the atmosphere: the Thermal
- 40 Dissociation-Laser Induced Fluorescence (TD-LIF) technique and comparisons to speciated
- 41 PAN measurements, Atmos. Meas. Tech., 3, 593-607, 10.5194/amt-3-593-2010, 2010.
- 42 Zhang, R. Y., Leu, M. T., and Keyser, L. F.: Heterogeneous chemistry of HO2NO2 in liquid
- 43 sulfuric acid, Journal of Physical Chemistry A, 101, 3324-3330, 10.1021/jp963321z, 1997.
- 44 Zhou, X. L., Beine, H. J., Honrath, R. E., Fuentes, J. D., Simpson, W., Shepson, P. B., and
- 45 Bottenheim, J. W.: Snowpack photochemical production of HONO: a major source of OH in

- the Arctic boundary layer in springtime, Geophys. Res. Lett., 28, 4087-4090,
- 10.1029/2001gl013531, 2001.
- Zhu, T., Yarwood, G., Chen, J., and Niki, H.: Evidence for the heterogeneous formation of nitrous acid from peroxynitric acid in environmental chambers, Environ. Sci. Technol., 27, 982-983, 10.1021/es00042a024, 1993.

Table 1. Summary of observed products, sensitivities, and detection limits (DL) for the reaction of I with HO₂NO₂, HONO and HO₂.

Analyte	Detected Ion (m/z)	Sensitivity ^a (Hz pptv ⁻¹)	DL (pptv, 3o)	Inlet ^a
HO ₂ NO ₂	I-HO ₂ NO ₂ ⁻ (206)	0.40 ± 0.06	20	Cold
	NO ₃ ⁻ (62)	144 ± 11^{b}	0.7	Cold, I
	I-HO ₂ ⁻ (160)	2.0 ± 0.04	40	Hot
HONO	I-HONO ⁻ (174)	1.7 ± 0.3	30	Cold, I
HO ₂	I-HO ₂ ⁻ (160)	2.6 ± 0.3	20	Cold, I

19 observed. The inlet dissociator was operated at a temperature of 130°C (Hot) or at ambient

20 temperature (Cold).

21 ^bSensitivity reported is for detection with an inlet dissociator at ambient temperature (Cold).





Figure 1. Shown are normalized (1e6 cps Γ), background corrected mass spectra acquired in the laboratory for calibration sources of HO₂ (a), HO₂NO₂ (b) and (c), and HONO (d). Spectra colored in blue were collected using a room temperature inlet (~25°C) while spectra in red indicate that an inlet dissociator at a temperature of 130°C was used. Background mass spectra have been subtracted from the displayed mass spectra to highlight the m/z ions that are produced *via* the Γ CIMS ion chemistry. All three species are detectable at a unique m/z, when an inlet dissociator is not used, allowing for simultaneous detection of HO₂, HO₂NO₂, and HONO.







3 sampled through PFA tubing at various flow rates (3-6 slpm) and lengths (0-3m). Concentrations

4 have been normalized to the initial concentration observed at the minimum reaction time

5 displayed. Reaction with NO₂ to form HO₂NO₂ appears to be the dominant loss for HO₂ on these

6 timescales.







- 3 dissociator (130°C) and the I-HO₂NO₂⁻ (m/z 206) ion with a cold dissociator (~25°C). Inset is a
- 4 correlation plot of the two measurements where comparison is possible.





Figure 4. Diurnal profiles of PNA are shown in panel (a) for the full measurement period during
the UBWOS 2013 and 2014 studies, with the 2014 study separated based on the sampling height
location. Panel (b) presents the difference in the 1m and <u>18.4m</u> PNA measurements for duration
of the 2014 study, where a positive value (brown) indicates larger concentrations at the ground
and a negative value (blue) suggests a relative PNA depletion in the 1m measurements. The

Patrick Veres 6/23/2015 3:39 PM Deleted: 30m

- 1 shaded region represents 1 standard deviation on the hourly average for the entire measurement
- 2 period.



Figure 5. Temperature and mixing ratios of HO₂ and NO₂ required to sustain an equilibrium
concentration of 1 ppbv HO₂NO₂. The region within the dashed circle superimposed on the
figure highlights the conditions encountered during the 2013 UBWOS study. Data shown were

⁷ calculated using the IUPAC database (Atkinson et al., 2004).









2 Figure 7. Displayed are gas phase HO₂NO₂ observations made during two wintertime ozone 3 events in the Uintah Basin, UT observed in 2013 (a), and through the entire 2014 study (b). 4 Measurements of the nitrite content of the snow surface taken during the same time periods are 5 also shown. In 2013, a precipitation event that occurred on the afternoon of February 8, shown as 6 the period in blue, added fresh surface snow and flushed pollutants out of the basin resulting in 7 lower ambient PNA and snow nitrite levels. Panel (c) shows in more detail periods from the 8 2013 measurements to better illustrate the daily reduction in snow surface nitrate that was 9 regularly observed.