

Author's response to the editor on "Partitioning and budget of inorganic and organic chlorine species observed by MIPAS-B and TELIS in the Arctic in March 2011" by G. Wetzel et al.

- 5 This file contains the response to the questions and comments of the editor together with the modified manuscript. All new changes (according to the suggestions of the editor) in the text of the manuscript are marked with blue colour.

10

Partitioning and budget of inorganic and organic chlorine species observed by MIPAS-B and TELIS in the Arctic in March 2011

- 15 **G. Wetzel¹, H. Oelhaf¹, M. Birk², A. de Lange³, A. Engel⁴, F. Friedl-Vallon¹, O. Kirner⁵, A. Kleinert¹, G. Maucher¹, H. Nordmeyer¹, J. Orphal¹, R. Ruhnke¹, B.-M. Sinnhuber¹, and P. Vogt²**

¹Karlsruhe Institute of Technology, Institute for Meteorology and Climate Research, Karlsruhe, Germany

- 20 ²Deutsches Zentrum für Luft und Raumfahrt, Institut für Methodik der Fernerkundung, Wessling, Germany

³SRON-Netherlands Institute for Space Research, Utrecht, the Netherlands

⁴Institut für Atmosphäre und Umwelt, J. W. Goethe Universität Frankfurt, Frankfurt, Germany

⁵Karlsruhe Institute of Technology, Steinbuch Centre for Computing, Karlsruhe, Germany

- 25 Correspondence to: G. Wetzel (gerald.wetzel@kit.edu)

Abstract

The Arctic winter 2010/2011 was characterized by a **persistent** vortex with extremely **low** temperatures in the lower stratosphere above northern Scandinavia leading to a strong activation of chlorine compounds (ClO_x) like Cl , Cl_2 , ClO , ClOOCl , OCIO , and HOCl which rapidly destroyed ozone when sunlight returned after winter solstice. The MIPAS (Michelson Interferometer for Passive Atmospheric Sounding) and TELIS (Terahertz and submillimeter Limb Sounder) balloon measurements obtained in northern Sweden on 31 March 2011 inside the polar vortex have provided vertical profiles of inorganic and organic chlorine species as well as diurnal variations of ClO around sunrise over the whole altitude range in which chlorine **has been** undergoing activation and deactivation. This flight was performed at the end of the winter during the last phase of ClO_x deactivation. The complete inorganic and organic chlorine partitioning and **budget for 31 March 2011, assumed to be representative for the late winter Arctic stratosphere**, has been derived by combining MIPAS-B and TELIS simultaneously observed molecules. A total chlorine amount of 3.41 ± 0.30 parts per billion by volume (ppbv) is inferred from the measurements (**above 24 km**). This value is in line with **previous** stratospheric observations carried out outside **the tropics** confirming the slightly decreasing chlorine **amount** in the stratosphere. Observations are compared and discussed with the output of a multi-year simulation performed with the Chemistry Climate Model EMAC (ECHAM5/MESSy Atmospheric Chemistry). The simulated stratospheric total chlorine amount is in accordance with the MIPAS-B/TELIS **observations** taking into account the fact that some chlorine source gases and very short-lived species are not included in the model.

1 Introduction

The discovery of the Antarctic stratospheric “ozone hole” in the **1970s** (Farman et al., 1985) strongly intensified research to unravel the reason for this ozone depletion. Chemically active chlorine (ClO_x) species like Cl , Cl_2 , ClO , ClOOCl , OCIO , and HOCl are part of total inorganic chlorine Cl_y ($\text{ClO}_x + \text{HCl} + \text{ClONO}_2$). They play a dominant role in the catalytic destruction of stratospheric ozone during polar winter when **low** temperatures and heterogeneous chemical reactions on polar stratospheric cloud (PSC) particles have previously enabled active chlorine compounds (mainly Cl_2) to be produced from its reservoir species ClONO_2 , HCl , and HOCl (e.g., Molina and Rowland, 1974; Solomon et al., 1986; Molina et al., 1987; **Solomon, 1999**; Crutzen and Oppenheimer, 2008). Due to the Montreal Protocol and successor agreements, emissions of dominant halocarbons were reduced such that total tropospheric (organic) chlorine

is decreasing since 1994 after reaching a peak value of nearly 3.7 parts per billion by volume (ppbv) (O'Doherty et al., 2004; WMO, 2011). The stratospheric total chlorine peak occurred
60 several years later, because **it takes this time for emitted air masses to propagate into the stratosphere** (Engel et al., 2002; WMO, 2011; Kohlhepp et al., 2012). The amount of equivalent effective stratospheric chlorine (chlorine and bromine halogens) is predicted to return to 1980 values around 2050 at mid-latitudes (Stolarski et al., 2010; WMO, 2011).

To assess and monitor the partitioning and budget of chlorine, a number of measurements of its
65 individual compounds have been carried out to calculate the amount of inorganic (Cl_y), organic (CCl_y), and finally total chlorine (Cl_{total}). An early observation based on data from the Atmospheric Trace Molecule Spectroscopy (ATMOS) instrument **was** published by Zander et al. (1992). A mean stratospheric total chlorine volume mixing ratio (VMR) of 2.58 ± 0.10 ppbv was observed at 30 °N in 1985. Significantly enhanced values between 3.4 and 3.5 (± 0.4) ppbv
70 in the 1992 Arctic lower stratosphere were estimated using retrieved data from the balloon-borne Michelson Interferometer for Passive Atmospheric Sounding (MIPAS-B) in combination with in-situ measurements (von Clarmann et al., 1995). A further slightly enhanced value of 3.53 ± 0.10 ppbv was detected during the ATMOS/ATLAS-3 November 1994 mission at northern mid-latitudes, also demonstrating the strong increase of stratospheric total chlorine
75 before regulating measures could alter this linear trend of 0.10 ppbv per year **in the time period from northern spring 1985 to fall 1994** (Zander et al., 1996). The same trend has been deduced between 1991 and 1995 by estimating total chlorine with the help of HCl observations from the Halogen Occultation Experiment (HALOE; Russell III et al., 1996). A further increased Cl_{total} value of 3.7 ± 0.2 ppbv was derived from MkIV balloon measurements carried out in the 1997
80 Arctic summer (Sen et al., 1999). This measurement **took place** close to the turn-over of the total stratospheric chlorine amount. Chlorine data obtained by the Atmospheric Chemistry Experiment Fourier Transform Spectrometer (ACE-FTS) in combination with in-situ measurements from the Stratospheric Aerosol and Gas Experiment (SAGE) III Ozone Loss and Validation Experiment (SOLVE) campaign (Schauffler et al., 2003; Nassar et al., 2006) were
85 used to estimate Cl_{total} between February 2004 and January 2005 in five latitude zones. A mean stratospheric Cl_{total} value of 3.65 ± 0.13 ppbv was determined for both the northern and southern mid-latitudes. This **start of a** temporal decrease of stratospheric chlorine was confirmed by observations **from** the Microwave Limb Sounder (MLS) from August 2004 until January 2006 (Froidevaux et al., 2006). A Cl_{total} value of 3.60 ± 0.13 ppbv at the end of this time period was

90 inferred from HCl measurements and a decrease of about 43 pptv in the stratospheric chlorine loading within this 18 month period was detected.

The long term trend of stratospheric inorganic chlorine was investigated by using data of multiple space-borne sensors like ACE-FTS, ATMOS, MLS, CLAES (Cryogenic Limb Array Etalon Spectrometer), CRISTA (Cryogenic Infrared Spectrometer and Telescope for the Atmosphere) and HALOE (Lary et al., 2007). This time series confirms that stratospheric Cl_y 95 peaked in the late 1990s and started to decrease as expected from the changing concentrations of tropospheric source gases and related transport times from the troposphere to the stratosphere. More recent published observations of Cl_{total} were performed by ACE-FTS covering the years 2004 until 2009. Nine chlorine containing species have been directly 100 measured by the satellite instrument (Brown et al., 2011, 2013). These data were supplemented by a number of further trace gases which were calculated using the SLIMCAT 3-dimensional Chemical Transport Model (Chipperfield, 2006). Global mean stratospheric chlorine was found to decrease by 0.46 % per year in the time period under investigation.

The purpose of this paper is to assess the partitioning and budget of inorganic and organic 105 stratospheric chlorine inside the late winter Arctic vortex. The winter 2010/2011 was characterized by a cold vortex defining a strong transport barrier until approximately mid-April (Manney et al., 2011; Sinnhuber et al., 2011). Temperatures were below the threshold associated for chlorine activation (~196 K) for more than 100 days between about 15 and 23 km. Consequently, an unprecedented Arctic ozone loss was observed which could be described 110 for the first time as an Arctic ozone hole since ozone profiles in late March resembled typical Antarctic late-winter profiles (Manney et al., 2011; Sinnhuber et al., 2011). Trace gas profiles of individual chlorine compounds were retrieved from limb emission spectra recorded during a balloon flight of MIPAS-B and the Terahertz and submillimeter Limb Sounder (TELIS) on 31 March 2011 inside the polar vortex. A description of the instruments, data analysis and chemical 115 modelling is given in Sect. 2. A discussion of the observed chlorine partitioning and budget follows in Sect. 3 and in the conclusions together with a comparison of the combined measured data to simulations of the Chemistry Climate Model (CCM) EMAC (ECHAM5/MESSy Atmospheric Chemistry model).

2 Instruments, data analysis and modelling

120 The MIPAS-B/TELIS flight took place on 31 March 2011 over northern Scandinavia inside the Arctic vortex at the end of the chlorine deactivation period that started slowly in early March

and accelerated towards the end of this month (Manney et al., 2011; Sinnhuber et al., 2011). The balloon gondola was launched from Esrange near Kiruna (Sweden, 67.9 °N, 21.1 °E) and reached its float level at about 35 km. Recorded limb sequences of MIPAS-B and TELIS are depicted in Fig. 1.

2.1 MIPAS-B instrument and data analysis

The balloon-borne limb-emission sounder MIPAS-B is a cryogenic Fourier Transform spectrometer which operates in the mid-infrared spectral range between about 4 and 14 μm . The maximum optical path difference of 14.5 cm of the beam in the interferometer allows a high unapodized spectral resolution of 0.0345 cm^{-1} (about 0.07 cm^{-1} after apodization with the Norton and Beer (1976) “strong” function) which allows the separation of individual spectral lines from continuum-like emissions in combination with a high radiometric accuracy of typically 1%. Values of the noise equivalent spectral radiance (NESR) are typically within 1×10^{-9} and $7 \times 10^{-9}\text{ W}(\text{cm}^2\text{ sr cm}^{-1})^{-1}$ for a single calibrated spectrum. Averaging over n spectra ($n \leq 5$) per single elevation scan reduces the spectral noise by a factor of $1/\sqrt{n}$. The instrument is characterized by a high performance and flexibility of the pointing system with a knowledge of the tangent altitude of better than 50 m at the $1\text{-}\sigma$ confidence limit. A comprehensive overview and description of the instrument together with processing of recorded interferograms to calibrated spectra including phase correction, Fourier Transformation to the spectral domain, and two-point calibration of the spectra from arbitrary to radiance units is given by Friedl-Vallon et al. (2004) and references therein. This includes instrument characterization in terms of the instrumental line shape, field of view, NESR, line of sight of the instrument, detector non-linearity (Kleinert, 2006) and the error budget of the calibrated spectra.

Forward radiance calculations were performed with the Karlsruhe Optimized and Precise Radiative transfer Algorithm (KOPRA; Stiller et al., 2002) which is a line-by-line and layer-by-layer model to simulate the infrared radiative transfer through the atmosphere. Molecular spectroscopic parameters for the calculation of limb emission spectra were taken from the high-resolution transmission molecular absorption database (HITRAN; Rothman et al., 2009) and a MIPAS dedicated spectroscopic data base (Raspollini et al., 2013). KOPRA also calculates derivatives of the radiance spectrum with respect to atmospheric state and instrument parameters and thus provides the Jacobians for the retrieval procedure KOPRAFIT (Höpfner et al., 2002). Since the vertical scan distance of adjacent tangent altitudes ranges between 1 and

1.5 km, the retrieval grid was set to 1 km up to the balloon float altitude. Above this level, the vertical spacing increases **gradually** to 10 km at the top altitude at 100 km. Considering the smoothing of the vertical part of the instrumental field of view, the retrieval grid is finer than the achievable vertical resolution of the measurement for a large part of the altitude region covered (especially above the observer altitude). To avoid retrieval instabilities due to this oversampling of the vertical retrieval grid, a Tikhonov-Phillips regularization approach (Phillips, 1962; Tikhonov, 1963) was applied **with a constraint** with respect to a first derivative a priori profile x_a of the target species:

$$x_{i+1} = x_i + [\mathbf{K}_i^T \mathbf{S}_y^{-1} \mathbf{K}_i + \mathbf{R}]^{-1} [\mathbf{K}_i^T \mathbf{S}_y^{-1} (y_{meas} - y(x_i)) - \mathbf{R}(x_i - x_a)] \quad (1)$$

where x_{i+1} is the vector of the desired state parameters for iteration $i+1$; y_{meas} is the measured radiance vector and $y(x_i)$ the calculation of the radiative transfer model using state parameters of iteration number i ; \mathbf{K} is the Jacobian matrix containing partial derivatives $\partial y(x_i) / \partial x_i$ while \mathbf{S}_y^{-1} is the inverse noise measurement covariance matrix and \mathbf{R} a regularization matrix with the first derivative operator and a regularization strength parameter.

In a first step, a temperature retrieval was performed using appropriate CO₂ lines of two separate bands around 810 cm⁻¹ and 950 cm⁻¹ and a priori pressure-temperature information from European Centre for Medium-Range Weather Forecasts (ECMWF) analyses together with a CO₂ VMR profile updated with data from NOAA ESRL GMD (National Oceanic and Atmospheric Administration, Earth System Research Laboratory, Global Monitoring Division; Montzka et al., 1999). The temperature retrieval 1- σ accuracy is estimated to be within about 0.7 K. Then, VMR profiles of the target species are individually retrieved in selected spectral regions (see Table 1). Profiles of species interfering with the target molecule were adjusted simultaneously during the retrieval procedure. An overview of **the analysis** of spectra with regard to chlorine- **and nitrogen**-containing molecules is given in von Clarmann et al. (1995) and Wetzel et al. (2002, 2010). The error estimation of the target parameter consists of random and systematic errors **that** were added in quadrature to yield the total error, which refers to the 1- σ confidence limit. Random errors include spectral noise as well as covariance effects of the simultaneously fitted parameters. Systematic errors mainly comprise spectroscopic data inaccuracies (band intensities), uncertainties in the line of sight, and gain calibration errors. The altitude resolution is calculated from the number of degrees of freedom of the retrieval, which

corresponds to the trace of the averaging kernel matrix. Typical values for the retrieved parameters are given in Table 1.

185 Many trace gases measured by the MIPAS-B instrument have been involved in a large number
of validation activities and cross-comparisons of satellite sensors like MIPAS, ILAS/ILAS-II
(Improved Limb Atmospheric Spectrometer) and SMILES (Superconducting Submillimeter-
Wave Limb-Emission Sounder). For species used in this work we explicitly mention for
evaluation the following studies: ClONO₂ (Höpfner et al., 2007; Wetzel et al., 2008, 2013),
190 CFC-11 and CFC-12 (Wetzel et al., 2008, 2013), ClO (Sagawa et al., 2013), and N₂O (Wetzel
et al., 2008; Payan et al., 2009).

2.2 TELIS instrument and data analysis

The cryogenic heterodyne balloon sounder TELIS was developed in a collaboration of three
partners: the German Aerospace Centre (DLR), Rutherford Appleton Laboratory (RAL),
195 United Kingdom, and the Netherlands Institute for Space Research (SRON). Each institute
generated one channel: 1.8 THz (DLR), 500 GHz (RAL), and 480-650 GHz (SRON). A
comprehensive description of the instrument is given by Birk et al. (2010) and de Lange et al.
(2012). The HCl and ClO results presented here were derived from spectra in the 480-650 GHz
channel with a tunable superconducting integrated receiver (SIR) developed and characterized
200 by de Lange et al. (2010) and de Lange et al. (2012). A local oscillator (LO) reference signal is
mixed with the atmospheric signal in a non-linear mixer. The measured spectrum is the
superposition of two spectra covering the frequency ranges $\nu_{LO} + \nu_{IF}$ and $\nu_{LO} - \nu_{IF}$, where ν_{IF} is
the measured difference (intermediate) frequency (IF).

The analysis of the TELIS spectra is carried out in a similar way as for the MIPAS-B retrieval
205 procedure. A forward line-by-line model is used to model the radiative transfer along the line-
of-sight of the instrument. Spectroscopic parameters are also taken from the HITRAN database
(Rothman et al., 2009). An instrument model to account for the specifics of the TELIS
instrument is included in the forward algorithm. Further details on the forward model are
described by de Lange et al. (2009) and references therein. The forward model is inverted with
210 a Gauss-Newton iteration scheme in combination with a Tikhonov-Phillips regularization
approach (Phillips, 1962; Tikhonov, 1963) as described in the previous section.

The HCl retrievals are performed for both chlorine isotopes H³⁵Cl and H³⁷Cl. The total amount
of HCl can be determined by taking into account the isotope abundance of H³⁵Cl (75.76 %) and

H³⁷Cl (24.23 %). While the random error of HCl is very small (~ 0.01 ppbv) the systematic error estimate yields between 0.05 and 0.4 ppbv resulting in a total error of about 10 to 15 % in the region of the VMR maximum. Systematic error sources are instrumental uncertainties such as instrumental line shape and side band ratio inaccuracies, detector non-linearity, and calibration and pointing errors. Furthermore, errors in the atmospheric pressure-temperature profile as well as spectroscopic data errors are taken into account. The largest uncertainty stems from the non-linear behaviour of the detector. This holds also for the ClO retrievals. The overall accuracy of the ClO measurement is almost entirely determined by systematic error sources. Similar to HCl, the total error for the species ClO typically remains within 10 and 15 % in the altitude region of its VMR maximum. An overview of the characteristics of the retrieved species is given in Table 2.

TELIS HCl and ClO observations have been evaluated using MLS measurements (de Lange et al., 2012). ClO was additionally compared to SMILES observations (Sagawa et al., 2013). The Sagawa study also includes a cross-comparison to MIPAS-B ClO observations.

2.3 Model calculations

Measured data are compared to simulations performed with the Chemistry Climate Model EMAC which is a numerical chemistry and climate simulation system that includes sub-models describing tropospheric and middle atmosphere processes (Jöckel et al., 2010). It uses the second version of the Modular Earth Submodel System (MESSy2) to link multi-institutional computer codes. The core atmospheric model is the 5th generation European Centre Hamburg general circulation model (ECHAM5, Roeckner et al., 2006). For the present study we applied EMAC (ECHAM5 version 5.3.02, MESSy version 2.50) in the T42L39MA-resolution, i.e., with a spherical truncation of T42 (corresponding to a quadratic Gaussian grid of approximately 2.8 by 2.8 degrees in latitude and longitude) with 39 vertical hybrid pressure levels from the ground up to 0.01 hPa. The applied model setup comprised, among others, the submodels MECCA (Sander et al., 2005) for the calculation of gas-phase chemistry and the submodel MSBM (Kirner et al., 2011) for the simulation of polar stratospheric clouds and the calculation of heterogeneous reaction rates. The PSC scheme was validated with the help of HNO₃, ClO, and O₃ data from the MLS instrument (Kirner et al., 2015).

A Newtonian relaxation technique of the prognostic variables temperature, vorticity, divergence and the surface pressure above the boundary layer and below 1 hPa towards the

245 ECMWF reanalysis ERA-Interim (Dee et al., 2011) has been applied to the model to simulate realistic synoptic conditions (van Aalst, 2005). Boundary conditions for greenhouse gases, chlorofluorocarbons (CFCs), and halons are adapted from observations (WMO, 2011; Meinshausen et al., 2011). Halogenated hydrocarbons are included according to the WMO-A1 scenario (WMO, 2011). Chlorine-containing tropospheric source gases considered in EMAC
250 are CFC-11, CFC-12, HCFC-22, CFC-113, CCl₄, CH₃Cl, and CH₃CCl₃. Photolysis rates of HCFC-22 and CFC-113 are the same as for CFC-12. The simulation includes a comprehensive chemistry setup from the troposphere to the lower mesosphere with 104 gas phase species, 234 gas phase reactions, 67 photolysis reactions, and 11 heterogeneous reactions on liquid aerosols, nitric acid trihydrate (NAT) and ice particles. Rate constants of gas-phase reactions are taken
255 from Atkinson et al. (2007) and Sander et al. (2011). The model output data were saved every 10 minutes. The temporally closest model output to the MIPAS-B measurements has been interpolated in space to the observed geolocations.

3 Chlorine partitioning and budget

The combination of two different sensors, MIPAS-B and TELIS, working in different spectral
260 regions (mid-infrared and microwave), enables the simultaneous measurement of virtually all relevant inorganic and organic chlorine molecules. The amount of inorganic chlorine [Cl_y] is defined as:

$$[\text{Cl}_y] = [\text{ClO}_x] + [\text{HCl}] + [\text{ClONO}_2] \quad (2)$$

where active chlorine [ClO_x] is calculated via:

$$265 \quad [\text{ClO}_x] = [\text{ClO}] + [\text{HOCl}] + 2 [\text{ClOOCl}] \quad (3)$$

The amount of organic chlorine [CCl_y] is composed of:

$$[\text{CCl}_y] = 2 [\text{CFC-12}] + 3 [\text{CFC-11}] + [\text{HCFC-22}] + 3 [\text{CFC-113}] + 4 [\text{CCl}_4] + [\text{CH}_3\text{Cl}] \quad (4)$$

Total chlorine [Cl_{total}] is given as the sum of both budgets:

$$[\text{Cl}_{\text{total}}] = [\text{Cl}_y] + [\text{CCl}_y] \quad (5)$$

270 Constituents, which are of minor importance for the Arctic stratospheric chlorine budget (like Cl₂, Cl, OClO, CH₃CCl₃, CFC-114, CFC-115, HCFC-141b, HCFC-142b, Halon-1211; see, e.g., Prinn et al., 2000) are neglected here. All the quantities defined in Eqs. (2) to (5) can be deduced from TELIS (measuring ClO and HCl) and MIPAS-B (measuring all gases except HCl) observations. However, the chlorine monoxide dimer ClOOCl is only measurable by MIPAS-

275 B under activated chlorine conditions ($[\text{ClOOCl}] > 0.5$ ppbv) without any PSC emissions in the recorded spectra (Wetzel et al., 2010). On 31 March 2011, no PSC signatures are visible in the MIPAS-B spectra but ClOOCl concentrations are below the detection limit. However, $[\text{ClOOCl}]$ can be estimated from $[\text{ClO}]$ with the following relation (Wetzel et al., 2012):

$$[\text{ClOOCl}] = ([\text{ClO}_{\text{noon}}] + 2 [\text{ClOOCl}_{\text{noon}}] - [\text{ClO}]) / 2 \quad (6)$$

280 while the amounts of $[\text{ClO}_{\text{noon}}]$ and $[\text{ClOOCl}_{\text{noon}}]$ which correspond to noon maximum and minimum values, respectively, can be both taken from EMAC simulations if the modelled ClO is constrained to the measured one.

MIPAS-B spectra have been recorded from night until day. The sunrise took place between 02:38 UTC at 36 km and 03:10 UTC at 9 km altitude. Fig. 2 shows the measured ClO cross section from 02:00 UTC to 04:38 UTC, corresponding to 64.0 °N, 30.1 °E and 63.5 °N, 28.9 °E. A temporal variation of ClO is visible. The concentration of this species is a measure of whether the air masses sounded are still chlorine-activated or not. After sunrise the mixing ratio of ClO increases in a layer between 16 and 22 km from nighttime values below 0.05 ppbv to daytime mixing ratios up to 0.4 ppbv. During periods of strong chlorine activation, significantly higher values around 2 ppbv are observed (see, e.g., Santee et al., 2003; Wetzel et al., 2012). The ClO increase is shown by both instruments, MIPAS-B and TELIS. The latter instrument measured not only with higher vertical resolution but also with higher temporal resolution compared to MIPAS-B, hence the TELIS data were transferred to the coarser temporal grid of MIPAS-B for better comparability. At altitudes above 26 km, MIPAS-B ClO temporal retrieval fluctuations are visible due to the large spectral noise error in this altitude region. As a consequence, the TELIS ClO data was used for calculating the chlorine partitioning and budget in the whole altitude range.

The decreasing ClO_x at the end of the Arctic winter in the lower stratosphere due to rising temperatures followed by shrinking ClO_x production from heterogeneous chemical reactions is in line with high amounts of ClONO_2 in this altitude region. The reaction of ClO with NO_2 produces the reservoir species ClONO_2 . The measured time evolution of this molecule is displayed in Fig. 3. Measured ClONO_2 data exhibit high values that are typical for observations in the late Arctic winter (see, e.g., Oelhaf et al., 1994; von Clarmann et al., 1997; Wetzel et al., 2002; von Clarmann et al., 2009). Only in an atmospheric layer around 19 km the vertical mixing ratio gradient is small since ClONO_2 values (around 19 km) are slightly lower than they would be if chlorine was completely deactivated. This observed signature is in line with the

enhanced ClO amounts around 19 km as seen in Fig. 2. A significant diurnal temporal variation is not visible in the ClONO₂ data.

The mean measured chlorine partitioning and budget for early morning is displayed in Fig. 4. A **spectral** noise error weighted averaging was applied to calculate the mean profiles, although statistical errors of the individual species profiles are similar. The molecules ClO and ClOOCl exhibit a temporal variation over the measured time period. However, since their mixing ratios are very low at this time in the year, vertical profiles of these species have also been averaged over the observed time period, with almost no consequence. To obtain a proxy of total inorganic chlorine, a N₂O-Cl_y correlation was derived from air samples collected with the balloon-borne cryogenic whole air sampler BONBON **in the Arctic** between 2009 and 2011 according to the method described in Engel et al. (2002) and Wetzel et al. (2010). **Cl_y from the cryosampler measurements is calculated as the difference between total chlorine and observed organic chlorine from the source gases CFC-11, CFC-12, CFC-113, CH₃CCl₃, CCl₄, HCFC-22, HCFC-141b, and HCFC-142b. In addition, an input of 50 pptv of chlorine from short lived source gases is taken into account which is assumed to be transformed immediately to inorganic chlorine. Total chlorine from the gases is propagated into the stratosphere in the same way as an inert tracer, as described in Engel et al. (2002), using global mean observation data from NOAA ESRL.** The proxy inorganic chlorine [Cl_y^{*}] is calculated with the following dependence on the amount of [N₂O], both given in ppbv:

$$[\text{Cl}_y^*] = 3.2008346 + 8.7786479 \times 10^{-6} [\text{N}_2\text{O}] - 2.9132361 \times 10^{-5} [\text{N}_2\text{O}]^2. \quad (7)$$

This correlation has been **applied** to MIPAS-B measured N₂O and yields up to 3.20 ppbv Cl_y^{*} in the stratosphere. The amount of inorganic chlorine is dominated by the chlorine reservoir species ClONO₂ and HCl, the latter one especially above 24 km. Above this altitude, where the Cl_y VMR is **(vertically) approximately** constant, the mean observed Cl_y amounts to 3.25 ± 0.30 ppbv which is in agreement with the deduced Cl_y^{*} within the error bars although there is a tendency **towards** a small positive **deviation** in the observations **compared to the Cl_y^{*} reference.** The **deviation** between Cl_y and Cl_y^{*} below 21 km **is caused** by different degrees of subsidence of the air masses in the case of the discussed balloon flight and the Cl_y^{*} reference which results in different N₂O mixing ratios in a specific altitude. Cl_y species play by far the largest part in the total chlorine budget from the lower to the upper Arctic winter stratosphere. **From about 17 km downwards**, the amount of organic chlorine gets increasingly dominant in the total chlorine budget. Source gases **that** contribute to **CCl_y are** visible in Fig. 4: CFC-12 (CCl₂F₂), CFC-11

(CCl₃F), HCFC-22 (CHClF₂), CFC-113 (C₂Cl₃F₃), CCl₄, and CH₃Cl. The mean amount of Cl_{total} is calculated as 3.41 ± 0.30 ppbv above 24 km. From the ratio Cl_y to Cl_{total} follows that about 95 % of total chlorine is inorganic in this altitude region.

The mean chlorine partitioning and budget as simulated by EMAC is shown in Fig. 5. The principal vertical profile shape of the measured chlorine species is well reproduced by the model. However, some differences in detail between simulated and observed data are visible. The modelled HCl VMR maximum appears slightly broader than the measured one. Below about 20 km, the simulation shows significantly lower values compared to the observation by TELIS. A difference is visible in the case of ClONO₂. The model underestimates this reservoir species and deviates by 0.8 ppbv (42 %) from the MIPAS-B data in the region of the VMR maximum at 22 km although simulated and measured NO_y and NO₂ (a necessary reactant in the production of ClONO₂ via NO₂ plus ClO) agree in this altitude region. Since simulated HCl and ClO_x (near 22 km) are in agreement with the observed data, the simulated Cl_y deviation from the measurement can be attributed to the ClONO₂ deficit in EMAC. Around 19 km, the difference in simulated and measured Cl_y is largest due to very low HCl values in EMAC compared to the HCl seen by TELIS. The amount of available Cl_y below about 24 km is dependent on the degree of downwelling of the air masses inside the polar vortex. In EMAC, the subsidence of the air masses in the course of the winter was underestimated such that we find higher values of tracers like N₂O and CFCs at a given altitude of the lower stratosphere compared to the measurements. These higher N₂O values are connected with lower Cl_y values according to the compact N₂O-Cl_y relationship, resulting in an underestimation of the chlorine reservoir species (especially ClONO₂). So, at least part of the ClONO₂ deficit in EMAC can be explained by the underestimation of the subsidence in the model.

The simulated Cl_y reaches its maximum VMR in the quasi altitude-constant region above 24 km with a mean value of 3.16 ppbv which is slightly lower than the measured one and close to the simulated value of Cl_y* (deduced from EMAC) which gives 3.19 ppbv. Below this altitude region, a similar bias between Cl_y and Cl_y* as in the case of the observations is visible.

The mean amount of Cl_{total} in the model run is calculated as 3.21 ppbv above 24 km, which is 0.20 ppbv lower than the observed one. About half of this simulated chlorine deficit can be explained by the fact that some minor CFCs (e.g. CFC-114 and CFC-115) and HCFCs (e.g. HCFC-141b and HCFC-142b) as well as halons are not included in the EMAC model. Their contribution to Cl_{total} is not more than 1% above 24 km (Brown et al., 2013). The remaining

deficit can be explained by very short-lived chlorine species which altogether amount to about 0.1 ppbv (Mébarki et al., 2010; WMO, 2011) and which are also not contained in the model simulation. However, the chlorine amount of these missing species is implicitly contained in the HCl measurement (since the short-lived chlorine species are converted to HCl after being photolyzed) and hence included in the observed chlorine budget. In the altitude region above 24 km, about 98 % of total chlorine in EMAC is inorganic. The shaded region of the budget profiles of ClO_x, Cl_y, CCl_y, and Cl_{total} shown in Fig. 5 takes into account all available chlorine species in EMAC that were not measured by MIPAS-B and TELIS. These molecules comprise Cl, Cl₂, OCIO (belonging to ClO_x and Cl_y) and CH₃CCl₃ (belonging to CCl_y) and add up to 0.1 ppbv at 16 km to the total chlorine budget (mainly due to Cl₂ and OCIO). However, at altitudes between 22 km and 36 km contributions of these gases to the chlorine budget are insignificant.

4 Conclusions

Observations from MIPAS-B/TELIS were performed at the end of the cold 2010/2011 stratospheric winter that was characterized by a persistent polar vortex enabling strong chlorine activation and ozone loss. The chlorine partitioning measured on 31 March 2011 reveals that in the outer part of the polar vortex (above Finland) the recovery of active chlorine (ClO_x) into the reservoir species (mainly ClONO₂) is almost completed by the end of March only a few days before the cold period had finished (Manney et al., 2011). This is verified by low amounts of daytime ClO of up to 0.4 ppbv around 19 km. The observed total stratospheric chlorine amounts to 3.41 ± 0.30 ppbv above 24 km (see Table 3). This is in accordance with the EMAC simulation (3.21 ppbv) taking into account the fact that some chlorine source gases and very short-lived species are not included in the model. The horizontal Cl_{total} distribution in EMAC (above 24 km) exhibits virtually no variation inside the polar vortex. The variation inside/outside vortex is no larger than 0.1 ppbv. That is smaller than the estimated Cl_{total} measurement accuracy of 0.3 ppbv such that the observations can be treated as representative at least for the geographical region of the Arctic vortex. Mean Cl_{total} values deduced from spectra recorded by the ACE-FTS instrument (Brown et al. 2013) give 3.44 ± 0.18 ppbv (morning occultations) and 3.50 ± 0.13 ppbv (evening occultations) for northern mid-latitudes and the Arctic in 2009. Extrapolating these data to 2011 with the chlorine trend (between 2004 and 2009) obtained from these ACE-FTS observations (about -0.4 % per year) yields Cl_{total} values of 3.41 ppbv (morning occultations) and 3.47 ppbv (evening occultations) comparable to the MIPAS-B/TELIS data. The accumulated amount of minor species (not measured by MIPAS-B/TELIS) like CFC-114,

CFC-115, HCFC-141b, HCFC-142b, and Halon-1211 was estimated to be about 0.7 % (~ 0.02 ppbv) of total chlorine at 30 km (Brown et al. 2013). Hence, the MIPAS-B/TELIS Cl_{total} value is in line with the data obtained from ACE-FTS solar occultations and is consistent with the decreasing amount of stratospheric chlorine. Considering the 2005 mean global tropospheric Cl_{total} from in-situ data of AGAGE (Advanced Global Atmospheric Gases Experiment) and NOAA ESRL databases, as compiled in WMO (2011), and transferring this value to 30 km taking into account a typical time lag of 6 years of stratospheric mean age of air (Engel et al., 2002, 2009; Stiller et al., 2008; WMO, 2011), we get an estimated Cl_{total} value of 3.40 ppbv for the year 2011, which is very close to the MIPAS-B/TELIS result (3.41 ppbv).

We finally conclude that the stratospheric total chlorine as deduced from Arctic MIPAS-B/TELIS observations on 31 March 2011 confirms previously published total chlorine assessments and their related trends. A recently published study by Mahieu et al. (2014) shows a HCl concentration increase between 2005/2006 and 2010/2011 in large parts of the northern hemispheric lower stratosphere in combination with an increase in the mean age of stratospheric air of up to 0.4 years. However, in the Arctic above 24 km, ascertained changes of mean age of stratospheric air are small (Mahieu et al., 2014) and, therefore, do not alter the findings above.

Acknowledgements

The balloon flight was funded by CNRS (Centre National de la Recherche Scientifique) and CNES (Centre National d'Etudes Spatiales) in the framework of the ENRICHED (European collaboration for Research on Stratospheric Chemistry and Dynamics) project coordinated by Nathalie Huret (Laboratoire de Physique et Chimie de l'Environnement et de l'Espace (LPC2E), Orléans, France) and Hermann Oelhaf (Karlsruhe Institute of Technology (KIT), Karlsruhe, Germany). We are grateful to the CNES balloon team for excellent balloon operations and the Swedish Space Cooperation for hosting the campaign and logistical assistance. We thank Nathalie Huret for project management and modeling support, and Katja Grunow from Free University of Berlin for meteorological support. The work presented here was funded in part by the European Space Agency (ESA) and the German Aerospace Center (DLR). We acknowledge support by Deutsche Forschungsgemeinschaft and Open Access Publishing Fund of Karlsruhe Institute of Technology.

References

- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin,
435 M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric
chemistry: Volume III – gas phase reactions of inorganic halogens, *Atmos. Chem. Phys.*, 7,
981-1191, doi:10.5194/acp-7-981-2007, 2007.
- Birk, M., Wagner, G., de Lange, G., de Lange, A., Ellison, B. N., Harman, M. R., Murk, A.,
Oelhaf, H., Maucher, G., and Sartorius, C.: TELIS: TeraHertz and subMMW Limb Sounder
440 – Project summary after first successful flight, in: Proceedings of the 21st International
Symposium on Space Terahertz Technology, University of Oxford and STFC Rutherford
Appleton Laboratory, Oxford, UK, 195–200, 2010.
- Brown, A. T., Chipperfield, M. P., Boone, C., Wilson, C., Walker, K. A., and Bernath, P. F.:
Trends in atmospheric halogen containing gases since 2004, *J. Quant. Spectrosc. Radiat.*
445 *Transfer*, 112, 2552-2566, 2011.
- Brown, A. T., Chipperfield, M. P., Dhomse, S., Boone, C., and Bernath, P. F.: Global
stratospheric chlorine inventories for 2004–2009 from Atmospheric Chemistry Experiment
Fourier Transform Spectrometer (ACE-FTS) measurements, *Atmos. Chem. Phys. Discuss.*,
13, 23491–23548, 2013.
- 450 Chipperfield, M. P.: New Version of the TOMCAT/SLIMCAT Off-Line Chemical Transport
Model: Intercomparison of Stratospheric Tracer Experiments, *Q. J. R. Meteorol. Soc.*, 132,
1179–1203, 2006.
- Crutzen, P. J., and M. Oppenheimer: Learning about ozone depletion, *Clim. Change*, 89, 143-
154, 2008.
- 455 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U.,
Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L.,
Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L.,
Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi,
M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., deRosnay,
460 P., Tavolato, C., Thépaut, J.-N., F. Vitart, F.: The ERA-Interim reanalysis: configuration and
performance of the data assimilation system, *Q. J. R. Meteorol. Soc.* 137, 553 – 597, 2011.

- de Lange, A., Landgraf, J., and Hoogeveen, R.: Stratospheric isotopic water profiles from a single submillimeter limb scan by TELIS, *Atmos. Meas. Tech.*, 2, 423–435, doi:10.5194/amt-2-423-2009, 2009.
- 465 de Lange, A., Birk, M., de Lange, G., Friedl-Vallon, F., Kiselev, O., Koshelets, V., Maucher, G., Oelhaf, H., Selig, A., Vogt, P., Wagner, G., and Landgraf, J.: HCl and ClO in activated Arctic air; first retrieved vertical profiles from TELIS submillimetre limb spectra, *Atmos. Meas. Tech.*, 5, 487-500, doi:10.5194/amt-5-487-2012, 2012.
- 470 de Lange, G., Birk, M., Boersma, D., Dercksen, J., Dmitriev, P., Ermakov, A., Filippenko, L., Golstein, H., Hoogeveen, R., de Jong, L., Khudchenko, A., Kinev, N., Kiselev, O., van Kuik, B., de Lange, A., van Rantwijk, J., Selig, A., Sobolev, A., Torgashin, M., de Vries, E., Wagner, G., Yagoubov, P., and Koshelets, V.: Development and characterization of the superconducting integrated receiver channel of the TELIS atmospheric sounder, *Supercond. Sci. Technol.*, 23, 045016, doi:10.1088/0953-2048/23/4/045016, 2010.
- 475 Engel, A., Strunk, M., Müller, M., Haase, H.-P., Poss, C., Levin, I., and Schmidt, U.: Temporal development of total chlorine in the high-latitude stratosphere based on reference distributions of mean age derived from CO₂ and SF₆, *J. Geophys. Res.*, 107, 4136, 10.1029/2001JD000584, 2002.
- 480 Engel, A., Möbius, T., Bönisch, H., Schmidt, U., Heinz, R., Levin, I., Atlas, E., Aoki, S., Nakazawa, T., Sugawara, S., Moore, F., Hurst, D., Elkins, J., Schauffler, S., Andrews, A., and Boering, K.: Age of stratospheric air unchanged within uncertainties over the past 30 years, *Nat. Geosci.*, 2, 28-31, 2009.
- Farman, J. C., B. G. Gardiner, and J. D. Shanklin: Large losses of total ozone in Antarctica reveal seasonal ClO_x/NO_x interaction, *Nature*, 315, 207-210, 1985.
- 485 Friedl-Vallon, F., Maucher, G., Kleinert, A., Lengel, A., Keim, C., Oelhaf, H., Fischer, H., Seefeldner, M., and Trieschmann, O.: Design and characterization of the balloon-borne Michelson Interferometer for Passive Atmospheric Sounding (MIPAS-B2), *Appl. Opt.*, 43, 3335-3355, 2004.
- 490 Froidevaux, L., Livesey, N. J., Read, W. G., Salawitch, R. J., Waters, J. W., Drouin, B., MacKenzie, I. A., Pumphrey, H. C., Bernath, P., Boone, C., Nassar, R., Montzka, S., Elkins, J., Cunnold, D., and Waugh, D.: Temporal decrease in upper atmospheric chlorine, *Geophys. Res. Lett.*, 33, L23812, doi:10.1029/2006GL027600, 2006.

- Höpfner, M., Oelhaf, H., Wetzell, G., Friedl-Vallon, F., Kleinert, A., Lengel, A., Maucher, G., Nordmeyer, H., Glatthor, N., Stiller, G., von Clarmann, T., Fischer, H., Kröger, C., and
495 Deshler, T.: Evidence of scattering of tropospheric radiation by PSCs in mid-IR limb emission spectra: MIPAS-B observations and KOPRA simulations, *Geophys. Res. Lett.*, 29(8), 1278, doi:10.1029/2001GL014443, 2002.
- Höpfner, M., von Clarmann, T., Fischer, H., Funke, B., Glatthor, N., Grabowski, U., Kellmann, S., Kiefer, M., Linden, A., Milz, M., Steck, T., Stiller, G. P., Bernath, P., Blom, C. E.,
500 Blumenstock, Th., Boone, C., Chance, K., Coffey, M. T., Friedl-Vallon, F., Griffith, D., Hannigan, J. W., Hase, F., Jones, N., Jucks, K. W., Keim, C., Kleinert, A., Kouker, W., Liu, G. Y., Mahieu, E., Mellqvist, J., Mikuteit, S., Notholt, J., Oelhaf, H., Piesch, C., Reddman, T., Ruhnke, R., Schneider, M., Strandberg, A., Toon, G., Walker, K. A., Warneke, T., Wetzell, G., Wood, S., and Zander, R.: Validation of MIPAS ClONO₂ measurements, *Atmos. Chem. Phys.*, 7, 257-281, doi:10.5194/acp-7-257-2007, 2007.
- 505
- Jöckel, P., Kerkweg, A., Pozzer, A., Sander, R., Tost, H., Riede, H., Baumgaertner, A., Gromov, S., and Kern, B.: Development cycle 2 of the Modular Earth Submodel System (MESSy2), *Geosci. Model Dev.*, 3, 717-752, doi:10.5194/gmd-3-717-2010, 2010.
- Kirner, O., Ruhnke, R., Buchholz-Dietsch, J., Jöckel, P., Brühl, C., and Steil, B.: Simulation
510 of polar stratospheric clouds in the chemistry-climate-model EMAC via the submodel PSC, *Geosci. Model Dev.*, 4, 169-182, 2011.
- Kirner, O., Müller, R., Ruhnke, R., and Fischer, H.: Contribution of liquid, NAT and ice particles to chlorine activation and ozone depletion in Antarctic winter and spring, *Atmos. Chem. Phys.*, 15, 2019-2030, doi:10.5194/acp-15-2019-2015, 2015.
- 515 Kleinert, A.: Correction of detector nonlinearity for the balloonborne Michelson Interferometer for Passive Atmospheric Sounding, *Appl. Opt.*, 45, 425-431, 2006.
- Kohlhepp, R., Ruhnke, R., Chipperfield, M. P., De Mazière, M., Notholt, J., Barthlott, S., Batchelor, R. L., Blatherwick, R. D., Blumenstock, Th., Coffey, M. T., Demoulin, P., Fast, H., Feng, W., Goldman, A., Griffith, D. W. T., Hamann, K., Hannigan, J. W., Hase, F.,
520 Jones, N. B., Kagawa, A., Kaiser, I., Kasai, Y., Kirner, O., Kouker, W., Lindenmaier, R., Mahieu, E., Mittermeier, R. L., Monge-Sanz, B., Morino, I., Murata, I., Nakajima, H., Palm, M., Paton-Walsh, C., Raffalski, U., Reddman, Th., Rettinger, M., Rinsland, C. P., Rozanov, E., Schneider, M., Senten, C., Servais, C., Sinnhuber, B.-M., Smale, D., Strong, K.,

- Sussmann, R., Taylor, J. R., Vanhaelewyn, G., Warneke, T., Whaley, C., Wiehle, M., and Wood, S. W.: Observed and simulated time evolution of HCl, ClONO₂, and HF total column abundances, *Atmos. Chem. Phys.*, 12, 3527-3556, doi:10.5194/acp-12-3527-2012, 2012.
- Lary, D. J., Waugh, D. W., Douglass, A. R., Stolarski, R. S., Newman, P. A., and Mussa, H.: Variations in stratospheric inorganic chlorine between 1991 and 2006, *Geophys. Res. Lett.*, 34, L21811, doi:10.1029/2007GL030053, 2007.
- 530 Mahieu, E., Chipperfield, M. P., Notholt, J., Reddman, T., Anderson, J., Bernath, P. F., Blumenstock, T., Coffey, M. T., Dhomse, S. S., Feng, W., Franco, B., Froidevaux, L., Griffith, D. W. T., Hannigan, J. W., Hase, F., Hossaini, R., Jones, N. B., Morino, I., Murata, I., Nakajima, H., Palm, M., Paton-Walsh, C., Russell III, J. M., Schneider, M., Servais, C., Smale, D. and Walker, K. A.: Recent northern hemisphere stratospheric HCl increase due to atmospheric circulation changes, *Nature*, 515, 104-107, doi:10.1038/nature13857, 2014.
- 535 Manney, G. L., Santee, M. L., Rex, M., Livesey, N. J., Pitts, M. C., Veefkind, P., Nash, E. R., Wohltmann, I., Lehmann, R., Froidevaux, L., Poole, L. R., Schoeberl, M. R., Haffner, D. P., Davies, J., Dorokhov, V., Gernandt, H., Johnson, B., Kivi, R., Kyro, E., Larsen, N., Levelt, P. F., Makshtas, A., McElroy, C. T., Nakajima, H., Parrondo, M. C., Tarasick, D. W., von der Gathen, P., Walker, K. A., and Zinoviev, N. S.: Unprecedented Arctic ozone loss in 2011, *Nature*, 478, 469-475, doi:10.1038/nature10556, 2011.
- 540 Mébarki, Y., Catoire, V., Huret, N., Berthet, G., Robert, C., and Poulet, G.: More evidence for very short-lived substance contribution to stratospheric chlorine inferred from HCl balloon-borne in situ measurements in the tropics, *Atmos. Chem. Phys.*, 10, 397-409, doi:10.5194/acp-10-397-2010, 2010.
- 545 Meinshausen, M., Smith, S. J., Calvin, K., Daniel, J. S., Kainuma, M. L. T., Lamarque, J.-F., Matsumoto, K., Montzka, S. A., Raper, S. C. B., Riahi, K., Thomson, A., Velders, G. J. M., van Vuuren, D. P. P.: The RCP greenhouse gas concentrations and their extensions from 1765 to 2300, *Clim. Change*, 109, 213-241, doi:10.1007/s10584-011-0156-z, 2011.
- 550 Molina, M. J., and F. S. Rowland: Stratospheric sink for chlorofluoromethanes: chlorine atom-catalysed destruction of ozone, *Nature*, 249, 810-812, 1974.
- Molina, M. J., Tso, T.-L., Molina, L. T., and Wang, F. C.-Y.: Antarctic stratospheric chemistry of chlorine nitrate, hydrogen chloride, and ice: Release of active chlorine, *Science*, 238, 1253-1257, 1987.

- 555 Montzka, S. A., Butler, J. H., Elkins, J. W., Thompson, T. M., Clarke, A. D. and Lock, L. T.:
Present and future trends in the atmospheric burden of ozone-depleting halogens, *Nature*,
398, 690-694, 1999.
- Nash, E. R., Newman, P. A., Rosenfield, J. E., and Schoeberl, M. R.: An objective
determination of the polar vortex using Ertel's potential vorticity, *J. Geophys. Res.*, 101,
560 9471-9478, 1996.
- Nassar, R., Bernath, P. F., Boone, C. D., Clerbaux, C., Coheur, P. F., Dufour, G., Froidevaux,
L., Mahieu, E., McConnell, J. C., McLeod, S. D., Murtagh, D. P., Rinsland, C. P., Semeniuk,
K., Skelton, R., Walker, K. A., and Zander, R.: A global inventory of stratospheric chlorine
in 2004, *J. Geophys. Res.*, 111, D22312, doi:10.1029/2006JD007073, 2006.
- 565 Norton, H., and Beer, R.: New apodization functions for Fourier spectroscopy, *J. Opt. Soc. Am.*,
66, 259-264 (Errata, *J. Opt. Soc. Am.*, 67, 419, 1977.) 1976.
- O'Doherty, S., Cunnold, D. M., Manning, A., Miller, B. R., Wang, R. H. J., Krummel, P. B.,
Fraser, P. J., Simmonds, P. G., McCulloch, A., Weiss, R. F., Salameh, P., Porter, L. W.,
Prinn, R. G., Huang, J., Sturrock, G., Ryall, D., Derwent, R. G., and Montzka, S. A.: Rapid
570 growth of hydrofluorocarbon 134a and hydrochlorofluorocarbons 141b, 142b, and 22 from
Advanced Global Atmospheric Gases Experiment (AGAGE) observations at Cape Grim,
Tasmania, and Mace Head, Ireland, *J. Geophys. Res.*, 109, D06310,
doi:10.1029/2003JD004277, 2004.
- Oelhaf, H., von Clarmann, T., Fischer, H., Friedl-Vallon, F., Fritsche, C., Linden, A., Piesch,
575 C., Seefeldner, M., Völker, W.: Stratospheric ClONO₂ and HNO₃ profiles inside the Arctic
vortex from MIPAS-B limb emission spectra obtained during EASOE, *Geophys. Res. Lett.*,
21, 1263-1266, 1994.
- Payan, S., Camy-Peyret, C., Oelhaf, H., Wetzell, G., Maucher, G., Keim, C., Pirre, M., Huret,
N., Engel, A., Volk, M. C., Kuellmann, H., Kuttippurath, J., Cortesi, U., Bianchini, G.,
580 Mencaraglia, F., Raspollini, P., Redaelli, G., Vigouroux, C., De Mazière, M., Mikuteit, S.,
Blumenstock, T., Velasco, V., Notholt, J., Mahieu, E., Duchatelet, P., Smale, D., Wood, S.,
Jones, N., Piccolo, C., Payne, V., Bracher, A., Glatthor, N., Stiller, G., Grunow, K., Jeseck,
P., Te, Y., and Butz, A.: Validation of version-4.61 methane and nitrous oxide observed by
MIPAS, *Atmos. Chem. Phys.*, 9, 413-442, doi:10.5194/acp-9-413-2009, 2009.

- 585 Phillips, D.: A technique for the numerical solution of certain integral equations of the first kind, *J. Assoc. Comput. Math.*, 9, 84–97, 1962.
- Prinn, R. G., Weiss, R. F., Fraser, P. J., Simmonds, P. G., Cunnold, D. M., Alyea, F. N., O'Doherty, S., Salameh, P., Miller, B. R., Huang, J., Wang, R. H. J., Hartley, D. E., Harth, C., Steele, L. P., Sturrock, G., Midgley, P. M., and McCulloch, A.: A history of chemically
590 and radiatively important gases in air deduced from ALE/GAGE/AGAGE, *J. Geophys. Res.*, 105, 17751-17792, 2000.
- Raspollini, P., Carli, B., Carlotti, M., Ceccherini, S., Dehn, A., Dinelli, B. M., Dudhia, A., Flaud, J.-M., López-Puertas, M., Niro, F., Remedios, J. J., Ridolfi, M., Sembhi, H., Sgheri, L., and von Clarmann, T.: Ten years of MIPAS measurements with ESA Level 2 processor
595 V6 – Part 1: Retrieval algorithm and diagnostics of the products, *Atmos. Meas. Tech.*, 6, 2419-2439, doi:10.5194/amt-6-2419-2013, 2013.
- Roeckner, E., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Koernblueh, L., Manzini, E., Schlese, U., and Schulzweida, U.: Sensitivity of simulated climate to horizontal and vertical resolution in the ECHAM5 atmosphere model, *J. Climate*, 19, 3771-3791, 2006.
- 600 Rothman, L. S., Gordon, I. E., Barbe, A., Benner, D. C., Bernath, P. F., Birk, M., Boudon, V., Brown, L. R., Campargue, A., Champion, J.-P., Chance, K., Coudert, L. H., Dana, V., Devi, V. M., Fally, S., Flaud, J.-M., Gamache, R. R., Goldman, A., Jacquemart, D., Kleiner, I., Lacome, N., Lafferty, W. J., Mandin, J.-Y., Massie, S. T., Mikhailenko, S. N., Miller, C. E., Moazzen-Ahmadi, N., Naumenko, O. V., Nikitin, A. V., Orphal, J., Perevalov, V. I., Perrin,
605 A., Predoi-Cross, A., Rinsland, C. P., Rotger, M., Šimečková, M., Smith, M. A. H., Sung, K., Tashkun, S. A., Tennyson, J., Toth, R. A., Vandaele, A. C., and Vander Auwera, J.: The HITRAN 2008 molecular spectroscopic database, *J. Quant. Spectrosc. Radiat. Transfer*, 110, 533–572, doi:10.1016/j.jqsrt.2009.02.013, 2009.
- Russell III, J. M., Luo, M., Cicerone, R. J., and Deaver, L. E.: Satellite confirmation of the
610 dominance of chlorofluorocarbons in the global stratospheric chlorine budget, *Nature*, 379, 526-529, 1996.
- Sagawa, H., Sato, T. O., Baron, P., Dupuy, E., Livesey, N., Urban, J., von Clarmann, T., de Lange, A., Wetzell, G., Connor, B. J., Kagawa, A., Murtagh, D., and Kasai, Y.: Comparison of SMILES ClO profiles with satellite, balloon-borne and ground-based measurements,
615 *Atmos. Meas. Tech.*, 6, 3325-3347, doi:10.5194/amt-6-3325-2013, 2013.

- Schauffler, S. M., Atlas, E. L., Donnelly, S. G., Andrews, A., Montzka, S. A., Elkins, J. W., Hurst, D. F., Romashkin, P. A., Dutton, G. S., and Stroud, V.: Chlorine budget and partitioning during the Stratospheric Aerosol and Gas Experiment (SAGE) III Ozone Loss and Validation Experiment (SOLVE), *J. Geophys. Res.*, 108, 4173, doi:10.1029/2001JD002040, 2003.
- 620
- Sander, R., Kerkweg, A., Jöckel, P., and Lelieveld, J.: Technical note: The new comprehensive atmospheric chemistry module MECCA, *Atmos. Chem. Phys.*, 5, 445-450, 2005.
- Sander, S. P., Friedl, R. R., Barker, J. R., Golden, D. M., Kurylo, M. J., Wine, P. H., Abbatt, J., Burkholder, J. B., Kolb, C. E., Moortgat, G. K., Huie, R. E., and Orkin, V. L.: Chemical kinetics and photochemical data for use in atmospheric studies Evaluation no. 17, JPL Publ. 10-6, Jet Propulsion Laboratory, Pasadena, CA, 2011.
- 625
- Santee, M. L., Manney, G. L., Waters, J. W., and Livesey, N. J.: Variations and climatology of ClO in the polar lower stratosphere from UARS Microwave Limb Sounder measurements, *J. Geophys. Res.*, 108, 4454, doi:10.1029/2002JD003335, 2003.
- 630
- Sen, B., Osterman, G. B., Salawitch, R. J., Toon, G. C., Margitan, J. J., Blavier, J.-F., Chang, A. Y., May, R. D., Webster, C. R., Stimpfle, R. M., Bonne, G. P., Voss, P. B., Perkins, K. K., Anderson, J. G., Cohen, R. C., Elkins, J. W., Dutton, G. S., Hurst, D. F., Romashkin, P. A., Atlas, E. L., Schauffler, S. M., and Loewenstein, M.: The budget and partitioning of stratospheric chlorine during the 1997 Arctic summer, *J. Geophys. Res.*, 104, 26653–26665, 1999.
- 635
- Solomon, S., Garcia, R. R., Rowland, F. S., and Wuebbles, D. J.: On the depletion of Antarctic ozone, *Nature*, 321, 755-758, 1986.
- Solomon, S.: Stratospheric ozone depletion: A review of concepts and history, *Rev. Geophys.*, 37, 275-316, 1999.**
- 640
- Sinnhuber, B.-M., Stiller, G., Ruhnke, R., von Clarmann, T., Kellmann, S., and Aschmann, J.: Arctic winter 2010/2011 at the brink of an ozone hole, *Geophys. Res. Lett.*, 38, L24814, doi:10.1029/2011GL049784, 2011.
- Stiller, G. P., von Clarmann, T., Funke, B., Glatthor, N., Hase, F., Höpfner, M., and Linden, A.: Sensitivity of trace gas abundances retrievals from infrared limb emission spectra to simplifying approximations in radiative transfer modeling, *J. Quant. Spectrosc. Radiat. Transfer*, 72(3), 249-280, 2002.
- 645

- 650 Stiller, G. P., von Clarmann, T., Höpfner, M., Glatthor, N., Grabowski, U., Kellmann, S., Kleinert, A., Linden, A., Milz, M., Reddmann, T., Steck, T., Fischer, H., Funke, B., López-Puertas, M., and Engel, A.: Global distribution of mean age of stratospheric air from MIPAS SF₆ measurements, *Atmos. Chem. Phys.*, 8, 677-695, doi:10.5194/acp-8-677-2008, 2008.
- Stolarski, R. S., Douglass, A. R., Newman, P. A., Pawson, P., and Schoeberl, M. R.: Relative contribution of greenhouse gases and ozone-depleting substances to temperature trends in the stratosphere: A Chemistry-Climate Model study, *J. Clim.*, 23, 28–42, 2010.
- 655 Tikhonov, A.: On the solution of incorrectly stated problems and a method of regularization, *Dokl. Acad. Nauk SSSR*, 151, 501–504, 1963.
- van Aalst, M. K.: *Dynamics and Transport in the Stratosphere - simulations with a general circulation model*, Ph.D. thesis, Institute for Marine and Atmospheric Research Utrecht, The Netherlands, 2005.
- 660 von Clarmann, T., Linden, A., Oelhaf, H., Fischer, H., Friedl-Vallon, F., Piesch, C., Seefeldner, M., Völker, W., Bauer, R., Engel, A., and Schmidt, U.: Determination of the stratospheric organic chlorine budget in the spring arctic vortex from MIPAS-B limb emission spectra and air sampling experiments, *J. Geophys. Res.*, 100, 13979-13997, 1995.
- 665 von Clarmann, T., Wetzel, G., Oelhaf, H., Friedl-Vallon, F., Linden, A., Maucher, G., Seefeldner, M., Trieschmann, O., and Lefèvre, F.: ClONO₂ vertical profile and estimated mixing ratios of ClO and HOCl in winter Arctic stratosphere from Michelson interferometer for passive atmospheric sounding limb emission spectra, *J. Geophys. Res.*, 102, 16157-16168, 1997.
- 670 von Clarmann, T., Höpfner, M., Kellmann, S., Linden, A., Chauhan, S., Funke, B., Grabowski, U., Glatthor, N., Kiefer, M., Schieferdecker, T., Stiller, G. P., and Versick, S.: Retrieval of temperature, H₂O, O₃, HNO₃, CH₄, N₂O, ClONO₂ and ClO from MIPAS reduced resolution nominal mode limb emission measurements, *Atmos. Meas. Tech.*, 2, 159-175, doi:10.5194/amt-2-159-2009, 2009.
- 675 Wetzel, G., Oelhaf, H., Ruhnke, R., Friedl-Vallon, F., Kleinert, A., Kouker, W., Maucher, G., Reddmann, T., Seefeldner, M., Stowasser, M., Trieschmann, O., von Clarmann, T., and Fischer, H.: NO_y partitioning and budget and its correlation with N₂O in the Arctic vortex and in summer mid-latitudes in 1997, *J. Geophys. Res.*, 107, 4280, doi:10.1029/2001JD000916, 2002.

- 680 Wetzel, G., Sugita, T., Nakajima, H., Tanaka, T., Yokota, T., Friedl-Vallon, F., Kleinert, A.,
Maucher, G., and Oelhaf, H.: Technical Note: Intercomparison of ILAS-II version 2 and 1.4
trace species with MIPAS-B measurements, *Atmos. Chem. Phys.*, 8, 1119-1126,
doi:10.5194/acp-8-1119-2008, 2008.
- 685 Wetzel, G., Oelhaf, H., Kirner, O., Ruhnke, R., Friedl-Vallon, F., Kleinert, A., Maucher, G.,
Fischer, H., Birk, M., Wagner, G., and Engel, A.: First remote sensing measurements of
ClOOCl along with ClO and ClONO₂ in activated and deactivated Arctic vortex conditions
using new ClOOCl IR absorption cross sections, *Atmos. Chem. Phys.*, 10, 931-945, 2010.
- Wetzel, G., Oelhaf, H., Kirner, O., Friedl-Vallon, F., Ruhnke, R., Ebersoldt, A., Kleinert, A.,
Maucher, G., Nordmeyer, H., and Orphal, J.: Diurnal variations of reactive chlorine and
nitrogen oxides observed by MIPAS-B inside the January 2010 Arctic vortex, *Atmos. Chem.*
Phys., 12, 6581-6592, doi:10.5194/acp-12-6581-2012, 2012.
- 690 Wetzel, G., Oelhaf, H., Friedl-Vallon, F., Kleinert, A., Maucher, G., Nordmeyer, H., and
Orphal, J.: Long-term intercomparison of MIPAS additional species ClONO₂, N₂O₅, CFC-
11, and CFC-12 with MIPAS-B measurements, *Annals of Geophysics*, 56, Fast Track-1,
10.4401/ag-6329, 2013.
- 695 WMO (World Meteorological Organization), Scientific Assessment of Ozone Depletion: 2010,
Global Ozone Research and Monitoring Project - Report No. 52, 516 pp., Geneva,
Switzerland, 2011.
- Zander, R., Gunson, M. R., Farmer, C. B., Rinsland, C. P., Irion, F. W., and Mahieu, E.: The
1985 chlorine and fluorine inventories in the stratosphere based on ATMOS observations at
30° north latitude, *J. Atmos. Chem.* 15, 171-186, 1992.
- 700 Zander, R., Mahieu, E., Gunson, M. R., Abrams, M. C., Chang, A. Y., Abbas, M., Aellig, C.,
Engel, A., Goldman, A., Irion, F. W., Kämpfer, N., Michelsen, H. A., Newchurch, M. J.,
Rinsland, C. P., Salawitch, R. J., Stiller, G. P., and Toon, G. C.: The 1994 northern
midlatitude budget of stratospheric chlorine derived from ATMOS/ATLAS-3 observations,
Geophys. Res. Lett., 23, 2357-2360, 1996.

705

Table 1. Setup for MIPAS-B trace species retrievals and typical ($1-\sigma$) errors. Results are given for different state parameters in corresponding spectral windows together with the retrieval altitude resolution (Alt. reso.).

Species	Spectral range (cm ⁻¹)	Noise error ^a (%)	Total error ^a (%)	Alt. reso. (km)
ClONO ₂	779.7 – 780.7	2 – 3	5 – 6	4 – 5
ClO	821.0 – 841.5	10 – 25 ^b	20 – 30 ^b	5 – 8
CFC-11	840.0 – 860.0	2 – 3	5 – 6	3 – 4
CFC-12	918.0 – 924.0	2 – 3	5 – 6	3 – 4
HCFC-22	828.0 – 830.0	3 – 10	8 – 15	3 – 6
CFC-113	813.0 – 830.0	3 – 10	20 – 25	3 – 6
CCl ₄	792.0 – 806.0	1 – 5	10 – 20	4 – 6
CH ₃ Cl	742.5 – 755.0	1 – 5	8 – 15	9 – 13
HOCl	1215.0 – 1265.0	10 – 15	35 – 50	6 – 8
N ₂ O	1227.8 – 1303.1	2 – 3	5 – 6	2 – 4

710 ^a in the altitude region around the VMR maximum;

^b daytime errors.

Table 2. Setup for TELIS HCl and ClO retrievals with typical ($1-\sigma$) errors and retrieval altitude resolution (Alt. reso.).

Species	Spectral line (GHz)	Noise error* (%)	Total error* (%)	Alt. reso. (km)
H ³⁵ Cl	625.9	< 1	10 – 15	2 – 5
H ³⁷ Cl	624.8	< 1	10 – 15	2 – 5
ClO	501.3	< 1	10 – 15	2 – 4

715 * in the altitude region around the VMR maximum.

720 **Table 3.** Mean stratospheric chlorine budgets (ppbv, including 1- σ total errors) as measured by MIPAS-B/TELIS and simulated by EMAC in comparison to ACE-FTS observations (Brown et al., 2013) and in-situ data from AGAGE and NOAA ESRL databases (WMO, 2011).

Budget	MIPAS-B/TELIS	EMAC	ACE-FTS	In-situ
Cl _{total}	3.41 ± 0.30 ^a	3.21 ^a	3.41 / 3.47 ^b	3.40 ^c
Cl _y	3.25 ± 0.30 ^a	3.16 ^a	-	-
Cl _y [*]	3.19 ± 0.002 ^a	3.19 ^a	-	-

^a mean value between 25 and 36 km;

^b mean value (for morning/evening occultations) between 30 and 70 °N for 2011, extrapolated from 2009 with trend between 2004 and 2009;

725 ^c mean global tropospheric value from 2005 corresponding to a stratospheric value of 2011 assuming a stratospheric mean age of 6 years.

730




Figure 1. Potential vorticity (PV) field (in $10^{-6} \text{ K m}^2 \text{ s}^{-1} \text{ kg}^{-1}$) from European Centre for Medium-Range Weather Forecasts (ECMWF) analysis on 31 March 2011, 00 UTC. MIPAS-B tangent points are plotted as black solid circles and TELIS tangent points as yellow solid circles (altitude range: 9.1 to 35.4 km). Both instruments look in the same direction. The vortex boundary which represents the strongest PV gradient (Nash et al., 1996) is shown as black dashed line. The insert (top left) shows the approximate measurement region (yellow marker) in relation to the position of the whole polar vortex.




Figure 2. Temporal evolution of chlorine monoxide (ClO) mixing ratios (ppbv) as observed by MIPAS-B (top) and TELIS (bottom) on 31 March 2011 above northern Scandinavia between 02:00 UTC and 04:38 UTC inside the Arctic vortex within the latitude/longitude sector shown in Fig. 1. The black solid line marks the sunrise terminator. A residual activation of chlorine is visible between 16 and 22 km with slightly enhanced ClO values up to 0.4 ppbv.




Figure 3. ClONO₂ mixing ratios (ppbv) as seen by MIPAS-B above northern Scandinavia on 31 March 2011 inside the late winter Arctic vortex. The black solid line marks the sunrise terminator.




Figure 4. Partitioning and budget of inorganic, organic, and total chlorine as measured by TELIS (HCl and ClO) and MIPAS-B (all other species) in the Arctic stratosphere on 31 March 2011 (see legend for line style and note non-linear abscissa). The reservoir species HCl and CIONO₂ dominate the stratospheric inorganic chlorine budget. Cl_y* deduced from observed N₂O data with the help of a N₂O-Cl_y correlation is shown for comparison. Note that for the calculation of the chlorine budgets the atomic content for each species has to be considered (some error bars have been omitted for clarity).




Figure 5. Partitioning and budget of inorganic, organic, and total chlorine as simulated with the chemistry climate model EMAC on 31 March 2011 (see legend for line style). Cl_y^* has been calculated from the simulated N_2O data according to Eq. (7). The budgets ClO_x , Cl_y , CCl_y , and Cl_{total} are calculated as listed in Eqs. (2) to (5), respectively. The shaded region of the budgets takes into account all minor chlorine species contained in EMAC (Cl_2 , Cl , OCIO , CH_3CCl_3) that were not measured by MIPAS-B and TELIS.