

We are thankful to the reviewer for her/his help in improving the quality of the manuscript. Our detailed responses to the comments are given below in italic

The authors have included an interesting update to the chemical transport model CHIMERE, involving an online coupling between aerosols and radiative effects on photolysis. This interaction has been shown before to be important in extremely polluted locations like Beijing but its importance in cleaner parts of the world is not clear. They focus on a wildfire event in Russia in order to address this importance for the model's performance against evaluation data in Moscow when the plume passes over. The difference between the model predictions with and without the new coupling are modest for ozone predictions. The major weakness of the paper is that the discussion sidesteps the comprehensive impacts of the wildfire, and instead only mentions the model's performance against itself (specifics described below). I would recommend the authors add some of this to the manuscript or perhaps submit to a more appropriate journal like Geoscientific Model Development.

Thanks to the reviewer's remarks, the aerosol radiative feedbacks on photochemistry have been now discussed in a more comprehensive way in the revised manuscript (see our detailed responses below). Moreover, as suggested by the reviewer 2, the evaluation of model results has been also improved by additional statistical comparisons for PM10 components (ammonium, nitrates and sulphates) in the section 3.2 of the revised manuscript:

| | NH_4^+ | | | | NO_3^- | | | | SO_4^{2-} | | | |
|---------|----------|------|-------|------|----------|------|-------|------|-------------|------|-------|------|
| | Mod. | Obs. | Corr. | RMSE | Mod. | Obs. | Corr. | RMSE | Mod. | Obs. | Corr. | RMSE |
| with | 1.09 | 0.86 | 0.48 | 1.03 | 0.14 | 0.12 | 0.20 | 0.15 | 2.24 | 0.48 | 0.23 | 1.80 |
| without | 1.16 | 0.86 | 0.42 | 1.04 | 0.19 | 0.12 | 0.26 | 0.23 | 2.33 | 0.48 | 0.45 | 1.88 |

Statistical comparisons between the near-surface concentrations of NH_4^+ , NO_3^- and SO_4^{2-} simulated with and without aerosol radiative feedbacks and measured at Moscow by an air quality station. Mod. and Obs. are the period-averaged modelled and observed concentration. Corr. and RMSE are the temporal correlation and the root mean square error.

As for NO_2 and O_3 , including the optical effect of aerosols in the photolysis calculations slightly improves the formation of secondary inorganic species in the CHIMERE model with a RMSE systematically reduced. The overestimation of sulphates levels is decreased and the simulated concentrations of ammonium and nitrates get close to the observed one. This result suggests that taking into account the aerosol solar extinction in the photolysis calculation gives an added value in the capacity of the model to reproduce the photochemistry under polluted environments.

Major Issues:

1) The introduction is brief, and focuses mostly on the problem from the perspective of model development. Yet the larger environmental impact perspective is somewhat lost. In other words, the authors present the fire episode as an example where the model might screw up if it doesn't have correct optical feedbacks. But what is the nature of this model problem in the context of the effects of the fire on the total environment, or model domain? Is it emitting so many BC particles that the effects of photolysis on secondary particles are kind of negligible? Are the uncertainties from photochemical reaction rates swamped by uncertainties in precursor emission rates? A reader could leave this paper thinking that fire plumes act to decrease O_3 concentrations, increase NO_2 and slightly decrease PM10, and that all of these effects are mild. Of course the whole story involves massive emissions of a host of pollutants. I think this piece of the story needs to be emphasized more

As suggested, questions relatives to fire events in the context of atmospheric composition and impacts have been better put into perspective in the Introduction of the revised manuscript by adding the following sentences :

« Also, the study of Chubarova et al. 2012 clearly shows that, during this specific wildfire episode, the aerosol optical thickness over the Moscow region was more than three times larger than the one observed during typical August conditions over the period 2001-2010. This suggests that, even if anthropogenic aerosols are present over the studied region, the contribution of smoke aerosols during this specific event is very large. Then, this case study represents an excellent opportunity to discuss how aerosol solar extinction, especially biomass burning particles, can affect photochemistry. Fires can affect atmospheric chemistry in several ways. They emit primary gaseous pollutants (such as CO, OH, NO, NO₂ and volatile organic compounds) that can react in the atmosphere to form ozone and other pollutants (Turquety, 2013). They also released aerosols that can directly affect air quality or indirectly by acting as a medium in complex heterogeneous reactions (Slade and Knopf 2013, Nie et al. 2015). Finally, they can affect the intensity of solar radiation, which in turn could affect photochemistry of the atmosphere. The latter impact is the subject of the present study. »

References quoted in the response and added in the revised manuscript :

- Chubarova, N., Nezval, Y., Sviridenkov, I., Smirnov, A., and Slutsker, I.: Smoke aerosol and its radiative effects during extreme fire event over Central Russia in summer 2010, *Atmos. Meas. Tech.*, 5, 557–568, 2012.

- Nie, W., Ding, A. J., Xie, Y. N., Xu, Z., Mao, H., Kerminen, V. M., Zheng, L. F., Qi, X. M., Huang, X., Yang, X. Q., Sun, J. N., Herrmann, E., Petäjä, T., Kulmala, M., and Fu, C. B.: Influence of biomass burning plumes on HONO chemistry in eastern China, *Atmospheric Chemistry and Physics*, 15, 1147–1159, 2015.

- Slade, J. H. and Knopf, D. A.: Heterogeneous OH oxidation of biomass burning organic aerosol surrogate compounds: assessment of volatilisation products and the role of OH concentration on the reactive uptake kinetics, *Phys. Chem. Chem. Phys.*, 15, 5898–5915, 2013.

- Turquety, S.: *The Atmospheric Impact of Wildfires. Fire Phenomena and the Earth System: An Interdisciplinary Guide to Fire Science*, Wiley Blackwell, 2013.


2) Figure 5 simplifies quite a lot of information and is a useful graphic. It also puzzles one at first glance. The JNO₂ values have changed by about half of the change in JO₃ as AOT increases. Yet the total change in O₃ barely shows up (at about -3% at most severe). Clearly, the concentrations of NO₂ and O₃ are playing important roles here. I think it would be interesting to probe the model a little further and look at the actual rates of formation and destruction of O₃ to get a more intuitive feeling of how the balance is affected by the aerosol direct effect.

Ozone formation is driven by two major precursors: nitrogen oxides (NO_x = NO + NO₂) and volatile organic compounds (VOC) in a complex photochemistry. However, it is possible to identify two regimes of ozone formation by looking at the ratio between the concentrations of VOC and NO_x: A NO_x-limited and a NO_x-saturated regime (Seinfeld and Pandis 1998). The Figure below indicates that the simulated photochemical regime was characterized by a NO_x-saturated situation over Moscow during the studied period. In this case, inclusion of the aerosol radiative impact on photochemistry leads to two antagonists responses : (1) Increase of NO_x concentration trough the reduction of their photolysis is unfavorable to ozone formation in a NO_x-saturated environment. In parallel, (2) reduction of the ozone photolysis is favorable to its accumulation. The overall impact

of aerosols on the ozone concentration is then small due to these two antagonist responses.

This point has been now clarified in the revised manuscript by adding the figure below and corresponding comments in the Section 3.2 of the revised manuscript :

« Response of the ozone concentration under the aerosol radiative influence is more complex. Indeed, ozone formation is driven by two major precursors: nitrogen oxides ($\text{NO}_x = \text{NO} + \text{NO}_2$) and volatile organic compounds (VOC) in a complex photochemistry. However, it is possible to identify two regimes of ozone formation by looking at the ratio between the concentrations of VOC and NO_x : A NO_x -limited and a NO_x -saturated regime (Seinfeld and Pandis 1998). Figure 6 indicates that the simulated photochemical regime was characterized by a NO_x -saturated situation over Moscow during the studied period. In this case, inclusion of the aerosol radiative impact on photochemistry leads to two antagonists responses : (1) Increase of NO_x concentration trough the reduction of their photolysis is unfavourable to ozone formation in a NO_x -saturated environment. In parallel, (2) reduction of the ozone photolysis is favourable to its accumulation. The overall impact of aerosols on the ozone concentration is then small due to these two antagonist responses, about 1 % per unit of AOT (see Figure 5). »



Mean VOC to NO_x ratio simulated over Moscow during the studied period

Reference quoted in the response and added in the revised manuscript :

- Seinfeld, J. H. and Pandis, S. N.: Atmospheric Chemistry and Physics. From air pollution to climate change, Wiley-Interscience publication, 1998.

Also, why were the values here plotted for midday? Was it because it makes it easier to calculate AOT through a single column. As the authors say in the text, it's the time of day with the least atmospheric path.


There is no specific reason for this choice. To keep coherence with previous discussions, Figure 5 has been rebuilt with daytime average values in the revised manuscript.

3) Page 7070-7071, lines 24-4: I'm uncomfortable with this discussion and the conclusion that taking the photolysis impacts into account could have a non-negligible effect on the air quality prediction. It is clear from figure 7 that the model is quite wrong on August 8. Yes, the model with aerosol feedbacks does very slightly better, but I think the authors are overstating the importance of the model improvement when they imply that it can make the difference between exceedance and attainment in air quality models. It seems like a relative coincidence after all, that the model without aerosol feedbacks falls above the threshold and the one with feedbacks falls below

We agree with the reviewer. This conclusion has been now removed.

4) The authors focus almost exclusively on O3 and include one plot at the end of the manuscript addressing PM mass. However, just by looking at the substantial reductions in sulfate and OA, I would expect there to be a significant effect on aerosol growth and the size distribution in general. Can the authors discuss this? How are there 8 aerosol bins affected?

This is a very interesting suggestion. Figure 8 has been modified to include some discussions on the aerosol size distribution in the Section 3.2 of the revised manuscript:



a) Daytime average percentage reduction of the near-surface concentration of sulphates, secondary organic aerosols and PM10 over Moscow due to the aerosol feedback. b) Repartition of this sulphate and SOA mass

« As illustrated in figure 9a, the maximum reduction in the near-surface concentrations of sulphates (oxidation product of SO₂) and SOA (oxidation product of COV) occurs on 8 August with daytime average values of 10 % and 4 %, respectively. For this day, figure 9b shows that these changes are mainly due to a reduced formation of very fine particles, i.e with a diameter comprised between 40 nm (bin 1) and 150 nm (bin 3). The overall impact is then a slightly reduction of the total aerosol mass concentration (PM10) comprised between 1 and 2 % over the entire period (Figure 9a) »

5) Since most of the discussion involves model development, I was surprised to not see a description of the computation time increase associated with incorporating the online coupling with the aerosols (including the core-shell calculations). This would help model developers decide whether or not they would like to implement the method in their own models (i.e. are the changes in O3 worth the time spent computing?)

The following sentence has been added in the Section 2.3 of the revised manuscript :

« It should be noted that adding the aerosol impact on solar extinction in simulation (2) induces a computation time increase of 50 % compared to the simulation (1).

Minor Issues:

1) The boundary conditions are not exactly consistent in time with the episode. Do the authors have an estimate for the influence of boundary conditions on their model domain? When predicting a very specific episode like this one, are they convinced that the lack of time dependence at the boundaries is acceptable?

The evaluation study of Péré et al. 2014 showed that the fire inventory of Kaiser et al. 2012 used in this work combined with the CHIMERE model have been shown to well capture the evolution of the Russian fire plume during the studied period, suggesting a low influence of these boundary conditions.

This point has been clarified by adding the following sentence in the section 2.1.1 of the revised manuscript :

« The evaluation study of Péré et al. 2014 showed that the fire inventory of Kaiser et al. 2012 used in this work combined with the CHIMERE model have been shown to well capture the evolution of the Russian fire plume during the studied period, suggesting a low influence of these boundary conditions. »

References quoted in the response and included in the revised manuscript :

- Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J. J., Razinger, M., Schultz, M. G., Suttie, M., and van der Werf, G. R.: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power, Biogeosciences, 9, 527–554, 2012.

- Péré, J. C., Bessagnet, B., Mallet, M., Waquet, F., Chiapello, I., Minvielle, F., Pont, V., and Menut, L.: Direct radiative effect of the Russian wildfires and its impact on air temperature and atmospheric dynamics during August 2010, Atmospheric Chemistry and Physics, 14, 1999–2013, 2014.

2) What version of WRF is being run offline? A little more detail on some of the weather selections for WRF would not be inappropriate. After all, the indirect effect calculations and the convective transport modules are relevant to this study.

The configuration of WRF used in this study has been now better described by adding the following sentences in the section 2.1.1 of the revised manuscript :

«The version 3.1 is used in this study with the same configuration as in Péré et al. (2011). It has 27 vertical levels from 40 m to about 20 km and includes the following parameterizations: the WRF single-moment five-class scheme of Hong et al. (2006) for the microphysics module, the Kain-Fritsch cumulus parametrization (Kain, 2004), the NOAH land surface module of Chen and Dudhia (2001) and the Yonsei University planetary boundary layer scheme (Hong et al., 2006; Hong, 2007).»

References quoted in the response and added in the revised manuscript :

- Chen, F. and Dudhia, J.: Coupling an advanced land-surface / hydrology model with the Penn State / NCAR MM5 modeling system. Part 1: Model description and implementation, Mon. Wea. Rev., 129, 569–585, 2001.
- Hong, S. Y., Noh, Y., and Dudhia, J.: A new vertical diffusion package with an explicit treatment of entrainment processes, Monthly Weather Review, 134, 2318–2341, 2006.
- Hong, S. Y.: Stable boundary layer mixing in a vertical diffusion scheme, The Korea Meteor. Soc., fall conference, Seoul, Korea, Oct. 25-26, 2007.
- Kain, J. S.: The Kain-Fritsch convective parameterization: An update, J. Appl. Meteor., 43, 170–181, 2004.
- Péré, J. C., Mallet, M., Pont, V., and Bessagnet, B.: Impact of the aerosol direct radiative forcing (ADRF) on the radiative budget, surface heat fluxes and atmospheric dynamics during the heatwave of summer 2003 over Western Europe. A modelling study., Journal of Geophysical Research, 116, d23119, doi:10.1029/2011JD016240, 2011.

This is quite relevant for figure 6, which might suggest that emissions from the fire are being injected too high.

As previously shown in details in the work of Péré et al. 2014, except for a lack of CALIOP measurements below 0.6 km, the model is within or close to the uncertainty range of observations except at 2 to 4 km where the model tends to largely overestimate CALIOP values (model biases from 30 to 350 % compared to observed median values). However, this part of the aerosol profile represents less than 30 % of the total aerosol extinction, suggesting that model uncertainties on the extinction vertical distribution remain limited.

3) What is TUV doing with sub-grid clouds? The grid cells (30 km) are somewhat coarse even for a regional simulation when the study is focusing on a plume

In TUV, clouds are assumed to be horizontally homogeneous layers. Hence, sub-grid cloud processes are not taken into account.

The following sentence has been added in the section 2.2 of the revised manuscript :

« In TUV, clouds are assumed to be horizontally homogeneous layers and are considered to be of three types : low, middle and high altitudes clouds. »

4) Page 7066, lines 10-23: While there is no denying that photolysis is critical for ozone formation in the atmosphere, the authors should also mention briefly the importance of VOCs in catalyzing ozone formation. The explanation given seems incomplete. This issue further relates back to the importance of emission inventory uncertainties compared to photolysis rate estimation uncertainties.

The reviewer is right. To clarify this point, the following sentence has been added in the Section 2.3 of the revised manuscript :

« The sequence of reactions (R1 to R4) is generally initiated by the reaction of various VOC with the OH radical. »

5) Figures 2 and 3: Please consider putting percent signs near the color axes for easy reference.

OK, This is now added.

Also, are these data for the lowest layer of the model or some column average?

Yes, the simulated values are for the lowest model layer. The term « near-surface » has been added in the captions.

May I suggest the authors replace “diurnal-average” with “daytime average” here and throughout the text?

Ok, this is now done.

6) Page 7069, lines 6-9: This statement seems true at first read, but certainly depends on the aerosol concentration, which will vary day to day and isn't necessarily a minimum at midday.

We agree with this remark. This statement has been removed and replaced by discussions on the daytime average values (please see response above about the major issue 2).

7) Figure 8: Are the SO₄ and SOA species for PM_{2.5} aerosols ?

Yes, they are present in the fine aerosol mode.

Presentation Issues and Typos:

1) Page 7065, line 28: Please consider putting the description of the simulations in their own section so that it is clearer to the reader that the computational experiment is being described. As it is now, the information is somewhat buried.

The subsection « On-line coupling between CHIMERE and TUV » has been renamed « Simulation set-up » in the revised manuscript.

2) Page 7058, line 15: In “terms” of air quality ...

3) Page 7058, line 26: “the exceedance of ...” and please change exceed to exceedance as appropriate throughout the text.

4) Page 7060, line 6: Please replace ‘emphasize’ with ‘focus’. There are many other English language typos throughout the paper. These don’t detract from the scientific quality of the paper, but they do distract. Please have someone look over the manuscript with the intention of correcting these mistakes

Thank you, the corresponding sentences have been now corrected in the revised manuscript. Also, we have now put a significant effort in improving its grammar quality.

We are thankful to the reviewer for her/his help in improving the quality of the manuscript. Our detailed responses to the comments are given below in italic

This manuscript describes a modeling study for aerosol radiative impact on photo-chemical pollutants near Moscow, Russia. This second version changed slightly from its first version, e.g. Figure 6's model extinction changed its wavelength from 400 nm to 500 nm. However, it still has very limited model-observation comparison. Other than the single comparison with CALIOP, all the manuscript describes the model-only result, which is hard to justify. Even that comparison, which is at night, has major issue since your discussion is about photochemical impact, which occurs during daytime, right?

In addition to the model comparison with NO₂ and O₃ observations, the evaluation of model results has been now improved by additional statistical comparisons for some PM10 components: ammonium, nitrates and sulphates (see last response below) . Moreover, thanks to both reviewers' remarks, the aerosol radiative feedbacks on photochemistry have been now discussed in a more comprehensive way in the revised manuscript.

Although this study is about the aerosol radiative impact from wildfire plume, it has very limited discussion the exact wildfire impact. How many aerosol impacts come from wildfire plume, and how many come from anthropogenic emission as Moscow is a megacity. The aerosol radiative impacts discuss throughout this paper could be partly due to anthropogenic influence. There is no discussion about the impact on wildfire event versus non-event.

We agree with the reviewer that the specific impact of wildfires on photochemistry has not been quantified. The aim of this study is to investigate the influence of aerosols on photochemistry during an extreme event (in terms of aerosol concentration), without differentiating particle's origin.

However, the study of Chubarova et al. 2012 clearly shows that, during this specific wildfire episode, the aerosol optical thickness over the Moscow region is more than three times larger than the one observed during typical August conditions over the period 2001-2010. This suggests that, even if anthropogenic aerosols are present over the studied region, the contribution of smoke aerosols during this specific event is very large.

This point has been clarified in the Introduction of the revised manuscript by adding the reference to the work of Chubarova et al 2012 :

« We focus on a major fire event that occurred in Russia during August 2010 as its episode was characterized by important concentrations of primary and secondary aerosols and large concentrations of ozone, especially over this specific region (Zvyagintsev et al. 2010, Konovalov et al. 2011, Popovicheva et al. 2014). Also, the study of Chubarova et al. 2012 clearly shows that, during this specific wildfire episode, the aerosol optical thickness over the Moscow region was more than three times larger than the one observed during typical August conditions over the period 2001-2010. This suggests that, even if anthropogenic aerosols are present over the studied region, the contribution of smoke aerosols during this specific event is very large. Then, this case study represents an excellent opportunity to discuss how aerosol solar extinction, especially biomass burning particles, can affect photochemistry. »

- Chubarova N., Y. Nezval, I. Sviridenkov, A. Smirnov and I. Slutsker: Smoke aerosol and its radiative effects during extreme fire event over Central Russia in summer 2010, Atmos. Meas. Tech., 5, 557-568, 2012.

Page 11, line 4: “the important concentrations of scattering aerosols”. In my knowledge, the major distinguished characteristic of biomass burning plume from anthropogenic aerosol is about its black carbon, or absorptive aerosol. Do you have any measurement near Moscow supporting your point ?

We agree with the reviewer that large values of SSA for biomass burning aerosols are not typical. However, as indicated in the Section 2.1.2, large values of SSA has been measured by AERONET over Moscow during the biomass burning event (0.95-0.96 in the visible/near infrared) and have been well simulated by CHIMERE (0.97 in the visible/near infrared). Such elevated SSA values are also supported by the study of Chubarova et al. 2012 highlighting the dominance of scattering species during this specific event.

This point has been clarified by adding the reference to Chubarova et al. 2012 in the section 2.1.2 of the revised manuscript :

« Such low aerosol absorption properties are supported by the study of Chubarova et al. 2012 highlighting elevated SSA values during this specific event. »

- N. Chubarova, Ye. Nezval, I. Sviridenkov, A. Smirnov, and I. Slutsker, Smoke aerosol and its radiative effects during extreme fire event over Central Russia in summer 2010, Atmos. Meas. Tech., 5, 557-568, 2012.

The major discussion is about the change of photolysis rate and its further impact on photochemical activities. However, there is no photolysis comparison with any observations, but just the model-only result? It is hard to verify whether the model gave the right result. The second version did not change it

We agree with the reviewer that it would be very interesting to make such comparisons. However, no photolysis frequencies measurements are available to us for this specific event. However, recently Palancar et al. 2013 realized a intercomparison exercise between the TUV model and UV actinic flux measurements over Mexico during the MILAGRO campaign. They highlighted the good performance of the model in reproducing observations both at the surface and in the lower troposphere over this highly polluted area. Hence, even though no comparison of photolysis simulations with measurements are made during this specific wildfire episode, the TUV validation study of Palancar et el. 2013 gives confidence in our estimation of photolysis rates perturbations by aerosols.

This point has been clarified in the section 2.2 of the revised manuscript by adding the reference to the work of Palancar et al. 2013 :

« Recently, Palancar et al. 2013 realized a intercomparison exercise between the TUV model and UV actinic flux measurements over Mexico during the MILAGRO campaign. They highlighted the good performance of the model in reproducing observations both at the surface and in the lower troposphere over this highly polluted area. This validation study gives confidence in our estimation of photolysis rates perturbations by aerosols during the 2010 Russian wildfires presented hereinafter. »

- Palancar G. G., B. L. Lefer, S. R. Hall, W. J. Shaw, C. A. Corr, S. C. Herndon, J. R. Slusser and S. Madronich, Effect of aerosols and NO₂ concentration on ultraviolet actinic flux near Mexico City during MILAGRO: measurements and model calculations, Atmos. Chem. Phys., 13, 1011–1022, 2013

In addition, to further investigate the performance of the model taking into account the aerosol radiative feedbacks, additional statistical comparisons have been included in the Section 3.2 of the revised manuscript for three inorganic aerosol species : ammonium, nitrates and sulphates :

| | NH_4^+ | | | | NO_3^- | | | | SO_4^{2-} | | | |
|---------|----------|------|-------|------|----------|------|-------|------|-------------|------|-------|------|
| | Mod. | Obs. | Corr. | RMSE | Mod. | Obs. | Corr. | RMSE | Mod. | Obs. | Corr. | RMSE |
| with | 1.09 | 0.86 | 0.48 | 1.03 | 0.14 | 0.12 | 0.20 | 0.15 | 2.24 | 0.48 | 0.23 | 1.80 |
| without | 1.16 | 0.86 | 0.42 | 1.04 | 0.19 | 0.12 | 0.26 | 0.23 | 2.33 | 0.48 | 0.45 | 1.88 |

Statistical comparisons between the near-surface concentrations of NH_4^+ , NO_3^- and SO_4^{2-} simulated with and without aerosol radiative feedbacks and measured at Moscow by an air quality station. Mod. and Obs. are the period-averaged modelled and observed concentration. Corr. and RMSE are the temporal correlation and the root mean square error.

As for NO_2 and O_3 , including the optical effect of aerosols in the photolysis calculations slightly improves the formation of secondary inorganic species in the CHIMERE model with a RMSE systematically reduced. The overestimation of sulphates levels is decreased and the simulated concentrations of ammonium and nitrates get close to the observed one. This result suggests that taking into account the aerosol solar extinction in the photolysis calculation gives an added value in the capacity of the model to reproduce the photochemistry under polluted environments.

Influence of the aerosol solar extinction on photochemistry during the 2010 Russian wildfires episode

J. C. Péré¹, B. Bessagnet², V. Pont³, M. Mallet³, and F. Minvielle¹

¹Laboratoire d'Optique Atmosphérique, Université Lille 1, 59655 Villeneuve d'Ascq, France.

²Institut National de l'Environnement Industriel et des Risques, Parc Technologique Alata, 60550 Verneuil en Halatte, France.

³Laboratoire d'Aérologie, Observatoire Midi-Pyrénées, 14 Avenue Edouard Belin, 31400 Toulouse, France.

Correspondence to: J. C. Péré
(jean-christophe.pere@univ-lille1.fr)

Abstract

In this work, impact of aerosol solar extinction on the photochemistry over eastern Europe during the 2010 wildfires episode is discussed for the period from 5 to 12 August 2010, which coincides to the peak of fire activity. The methodology is based on an on-line coupling between the chemistry-transport model CHIMERE (extended by an aerosol optical module) and the radiative transfer code TUV. Results of simulations indicate an important influence of the aerosol solar extinction, in terms of intensity and spatial extent, with a reduction of the photolysis rates of NO_2 and O_3 up to 50 % (in ~~diurnal-averaged~~ daytime average) along the aerosol plume transport. At a regional scale, these changes in photolysis rates lead to a 3–15 % increase in the NO_2 daytime concentration and to an ozone reduction near the surface of 1–12 %. The ozone reduction is shown to occur over the entire boundary layer, where aerosols are located. Also, ~~comparisons of simulations with air quality measurements over Moscow show that the inclusion of the aerosol feedback tends to slightly improve performance of the model in simulating NO_2 and O_3 ground concentrations. In term of air quality prediction, the O_3 peak reduction when including aerosol feedback results in a non-negligible difference in the predicted exceedance of alert threshold compared to the simulation without aerosol feedback, in coherence with measurements. Finally,~~ the total aerosol mass concentration (PM10) is shown to be decreased by 1–2 %, on average during the studied period, caused by a reduced formation of secondary aerosols such as sulphates and secondary organics (4–10 %) when aerosol impact on photolysis rates is included. In terms of model performance, comparisons of simulations with air quality measurements at Moscow indicate that an explicit representation of aerosols interaction with photolysis rates tend to improve the estimation of the near-surface concentration of ozone and nitrogen dioxide as well as the formation of inorganic aerosol species such as ammonium, nitrates and sulphates.

1 Introduction

For several years, it has been well recognized that air pollution of gaseous and particulate origin can have adverse health effects (Miller et al., 2012; Beelen et al., 2014). In consequence, efficient air pollution control strategies have now become a challenge for environmental policies. In the context of air quality monitoring, the ~~exceed~~ exceedance of certain thresholds of pollutant concentrations is a criterion often used by authorities of a country to prevent people from air pollution exposure. In general, the ~~exceed~~ exceedance of these thresholds is evaluated from air quality numerical forecast such as in France where the regional chemistry-transport model CHIMERE (Menut et al., 2013) is used in the French national air quality forecasting and monitoring system known as PREV'AIR (Honoré et al., 2008). Photochemical pollutants (ozone, secondary aerosols,...), which are formed from photo-dissociation of precursors such as nitrogen dioxide (NO₂) and volatile organic compounds (VOC) (Jenkin and Clemitshaw, 2000), are of particular interest for air quality monitoring (Honoré et al., 2008) due to their negative impacts on both environment and human health (Amin, 2014; Hunova et al., 2014).

The key parameter that governs the photo-dissociation of photochemical precursors in the atmosphere is the photolysis rate, which mainly depends on the available actinic flux (Seinfeld and Pandis, 2006). Aerosols are known to have large influence on the available actinic flux by interacting with solar radiation in the ultraviolet-visible wavelengths (Li et al., 2011a,b; Lou et al., 2014). For example, Wai and Tanner (2010) showed, by using a combination of remote sensing observations and chemical-transport model, that aerosol solar extinction could lead to a 7–32 % reduction in maximum ozone concentration over Hong-Kong during highly polluted days. Also, Li et al. (2011b) highlighted, with WRF-CHEM modelling experiments, that changes in photolysis rates due to the presence of particles led to a decrease of about, respectively, 2–17 % and 5–6 % in daytime ozone and secondary aerosols (nitrate, secondary organics) concentrations over Mexico City during the 2006 Megacity Initiative: Local and Global Re-

search Observations (MILAGRO) campaign.

To reduce computational time for operational purpose, one major characteristic of air quality modelling platforms is that impacts of aerosols and clouds on solar radiation are often taken into account as simplified attenuation factors when evaluating the photolysis rates (Honoré et al., 2008; Menut et al., 2013). However, Real and Sartelet (2011) highlighted that simplified parametrization of aerosol impact on photolysis rates could tend to worsen air quality model performance in simulating ozone and particulate concentration, especially under highly polluted environments.

The aim of the present study is to implement, in the chemistry-transport model CHIMERE,

an explicit representation of the alteration of photolysis rates by aerosols and discuss the impact in terms of modelled ozone budget and the formation of secondary aerosols at regional scale. We ~~emphasize~~ focus on a major fire event that occurred in Russia during August 2010 as its episode was characterized by important concentrations of primary and secondary aerosols and large concentrations of ozone, especially over this specific region (Zvyagintsev et al., 2010; Konovalov et al., 2011; Popovichcheva et al., 2014). ~~Then, it~~ Also, the study of Chubarova et al. 2012 clearly shows that, during this specific wildfire episode, the aerosol optical thickness over the Moscow region was more than three times larger than the one observed during typical August conditions over the period 2001–2010. This suggests that, even if anthropogenic aerosols are present over the studied region, the contribution of smoke aerosols during this specific event is very large. Then, this case study represents an excellent opportunity to discuss how aerosol solar extinction, especially biomass burning particles, can affect photochemistry. Fires can affect atmospheric chemistry in several ways. They emit primary gaseous pollutants (such as CO, OH, NO, NO₂ and volatile organic compounds) that can react in the atmosphere to form ozone and other pollutants (Turquety, 2013). They also released aerosols that can directly affect air quality or indirectly by acting as a medium in complex heterogeneous reactions (Slade and Knopf, 2013; Nie et al., 2015). Finally, they can affect the intensity of solar radiation, which in turn could affect photochemistry of the atmosphere. The latter impact is the subject of the present study.

The approach is based on an on-line coupling between the regional model CHIMERE, extended by an aerosol optical module (Péré et al., 2010), and the Tropospheric Ultraviolet and Visible (TUV) radiation model (Madronich and Flocke, 1998). In this methodology, the aerosol optical thickness, single scattering albedo and asymmetry parameter are first modelled by CHIMERE using an aerosol core-shell mixing hypothesis, as in Péré et al. (2009, 2010). This mixing approach has been previously used by Péré et al. (2014) to study the 2010 Russian wildfires direct radiative forcing and its feedback on the regional atmospheric dynamics. Results indicate that it can give a good representation of the absorption properties of particles during this specific period. In a second time, aerosol optical properties are used as inputs in the radiative transfer code TUV to evaluate the impact of aerosol short-wave solar extinction on photolysis rates and the formation of ozone and secondary particles. The advantage of such methodology is the use of two specific state-of-the-art models to explicitly simulate the interaction of physico-chemically resolved aerosols with the actinic flux and the associated impact on modelled photolysis rates and photochemistry.

Section 2 describes the configuration of each model as well as the development of their on-line coupling. In Section 3 are discussed modelled regional changes in the near surface concentrations of NO_2 , O_3 and secondary aerosols over Russia induced by modifications of photolysis rates by smoke aerosols during August 2010. Finally, conclusions and perspectives of future works are given in Section 4.

2 Methodology

2.1 Description of the CHIMERE model

2.1.1 Aerosol module

CHIMERE is a state-of-the-art 3D chemistry transport model that calculates the concentrations of numerous gaseous and particulate pollutants (Vautard et al., 2001; Menut

et al., 2013). The dynamics and gas phase parts of the model is regularly improved (Menut et al., 2013) and its documentation can be downloaded at <http://www.lmd.polytechnique.fr/chimere/>. In this work, the CHIMERE domain has a 30 km horizontal resolution and ranges from 43.40°N to 63.20°N in latitude and from 18.70°E to 57.30°E in longitude. The aerosol part is described by Bessagnet et al. (2004) and is composed of 10 chemical species: sulphates, nitrates, ammonium, primary organic and black carbon (OC and BC), secondary organic aerosols (SOA), sea salt, natural and anthropogenic dust and water. The evolution of aerosols is described with a 8-bins size distribution (from about 40 nm to 10 μ m in diameter) and includes the main physical processes such as nucleation, coagulation, condensation/evaporation, adsorption/desorption, wet and dry deposition and scavenging.

Anthropogenic emissions of gaseous and particulate origin come from the EMEP database. Concerning OC and BC emissions, the inventory of Junker and Liousse (2008) has been used. Natural soil dust are dynamically produced within the domain according to the methodology of Vautard et al. (2005). SOA formation is represented through oxidation processes of relevant precursors of biogenic and anthropogenic origin and gas particle partitioning schemes (Bessagnet et al., 2008). VOC and NO emissions from vegetation are calculated using the Model of Emissions of Gases and Aerosols from Nature (MEGAN) (Guenther et al., 2006). Aerosols and gases emitted by wildfires affecting Russia during 2010 are taken into account following the work described and validated by Kaiser et al. (2012). It consists in the assimilation of the fire radiative observations from the Moderate Resolution Imaging Spectroradiometer into the Global Fire assimilation System combined with the use of specific combustion rate and emission factors to estimate biomass burning emissions. More information on the methodology as well as a validation study for the 2010 Russian wildfires episode can be found in ~~(Kaiser et al., 2012)~~ [Kaiser et al. \(2012\)](#).

CHIMERE is forced at these boundaries by monthly climatologies, calculated over the 2000–2004 period, of the main gases and particles provided by the MOZART (Horowitz et al., 2003) and LMDzT-INCA global chemistry-transport models (Hauglus-

tainé et al., 2004), respectively. [The evaluation study of Péré et al. \(2014\) showed that the fire inventory of Kaiser et al. \(2012\) used in this work combined with the CHIMERE model have been shown to well capture the evolution of the Russian fire plume during the studied period, suggesting a low influence of these boundary conditions.](#) Also, CHIMERE is off-line driven by the Weather Research and Forecasting model (WRF) at a 30 km resolution. [The version 3.1 is used in this study with the same configuration as in Péré et al. \(2011\) . It has 27 vertical levels from 40 m to about 20 km and includes the following parametrizations: the WRF single-moment five-class scheme of Hong et al. \(2006\) for the microphysics module, the Kain–Fritsch cumulus parametrization \(Kain, 2004\) , the NOAH land surface module of Chen and Dudhia \(2001\) and the Yonsei University planetary boundary layer scheme \(Hong et al., 2006; Hong, 2007\) .](#)

2.1.2 Modelling aerosol optical properties

The calculation of optical properties of particles is the pre-requisite for the evaluation of their impacts on photolysis rates and photochemistry as they provide information on how aerosols will interact with the actinic flux. In that sense, we developed a numerical scheme dedicated to calculate aerosol optical properties from aerosol concentrations, size distribution and chemical composition modelled by CHIMERE. A complete description of this optical module is presented in the work of Péré et al. (2010). To compute the complex refractive index of a particle, the hypothesis of a core-shell mixing has been chosen with a core of primary species (BC, OC and dust) surrounded by a shell of secondary ones (sulphates, nitrates, ammonium, secondary organics) and sea salt and water. This mixing choice is supported by recent studies giving evidence of coatings of secondary particles on black carbon aerosols over Europe (Vester et al., 2007; McMeeking et al., 2011). Also, such mixing has been shown to correctly reproduce the absorbing properties of aerosols during the 2010 Russian wildfire episode (Péré et al., 2014). For each size bin, a volume average procedure is used to calculate the refractive index of the core and the shell (Lesins et al., 2002) which is then used as inputs in the Mie algorithm for n-layered spheres of Wu and Wang (1991) to calculate the scat-

tering and absorption coefficients. It should be noted that the volume of the core and the shell can vary during the simulation in function of the different physical processes influencing aerosol population. The optical properties of the total aerosol distribution needed in radiative transfer modelling, such as the Aerosol Optical Thickness (AOT), Single Scattering Albedo (SSA) and asymmetry parameter (g), are calculated as in Wu et al. (1996).

A detailed evaluation of the optical module for the 2010 Russian wildfire episode by using Aerosol Robotic Network (AERONET) sunphotometers measurements (AOT, SSA, aerosol size distribution) and Polarization and Directionality of the Earth's Reflectances (POLDER) and Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) aerosol extinction data is presented by Péré et al. (2014). Only a sum-up is given here. As displayed in Figure 1a-b, simulations for the period 5–12 August 2010 (peak of fire activity) show large AOT (at midday) over eastern Europe with modelled values above 2 (at 400 nm) in the aerosol plume. This aerosol plume was transported in an anticyclonic flow from the source region towards Moscow (5–8 August), to the north (9–10 August) and then back to the east (11–12 August). On average, simulations compare well with AOT POLDER data in terms of spatial distribution and intensity (correlation = 0.67–0.77, $-10\% < \text{biases} < 23\%$). The altitude of transport was shown to be below 5 km and comparisons between CHIMERE and CALIOP show good consistency. Figure 1a-b indicates that the simulated plume was advected over Moscow between 5–9 August, which is in accordance with AERONET measurements. During the study period, the aerosol composition was dominated by scattering organic species with modelled elevated SSA (0.97 between 300 nm and 1000 nm) close to AERONET values over Moscow (0.95–0.96 between 440 nm and 1020 nm). Such low aerosol absorption properties are supported by the study of Chubarova et al. (2012) highlighting elevated SSA values during this specific event. Globally, the comparisons between aerosol simulation and sunphotometers data highlighted the ability of the model to give an appropriate representation of the aerosol size distribution and scattering/absorption properties (Péré et al., 2014), which is the pre-requisite to evaluate its influence on photolysis

rates and the formation of photochemical pollutants.

2.2 Description of the TUV model

TUV is a widely-used state-of-the-art radiative transfer model developed at the National Centre for Atmospheric Research (Madronich and Flocke, 1998). In this study, we used the version 4.6 of the code (released in March 2009) freely available at the website: <http://cprm.acd.ucar.edu/Models/TUV/>. The model calculates the actinic flux and photolysis rates of a large number of photochemical species.

Photolysis is the process breaking the covalent bond of some reactive gaseous species by short-wave solar radiation. This process is very important in the atmosphere as it controls the abundance of numerous air pollutants such as ozone and nitrogen dioxide. The photolysis rate of a given specie J (s^{-1}) is calculated as follows:

$$J(s^{-1}) = \int_{\lambda_1}^{\lambda_2} \sigma(\lambda, T) \cdot \phi(\lambda, T) \cdot F(\lambda) d\lambda \quad (1)$$

where $\sigma(\lambda, T)$ and $\phi(\lambda, T)$ are, respectively, the absorption cross section (cm^2) and the quantum yield of a given molecule, T the air temperature (K) and $F(\lambda)$ the actinic flux between wavelengths λ_1 and λ_2 ($photons.cm^{-2}.s^{-1}.nm^{-1}$). The absorption cross section reflects the probability of collision between a photon and the molecule, while the quantum yield is the probability that the molecule is dissociated after collision with a photon. The dependence of both parameters on the air temperature is calculated by TUV by using the vertical profile of air temperature issued from the meteorological model WRF used to drive CHIMERE.

The actinic flux is calculated by integrating the solar flux over all sphere angles considering 5646 wavelengths between 120 nm and 1250 nm. When going through the

atmosphere, the actinic flux can be attenuated by molecular absorption and diffusion but also by the presence of clouds and aerosols. For clouds and aerosols, the attenuation is calculated by using their respective aerosol optical thickness, single scattering albedo and asymmetry parameter. In TUV, ~~three types of clouds are represented~~ clouds are assumed to be horizontally homogeneous layers and are considered to be of three types : low, middle and high altitude clouds. Altitudes of their bases and tops as well as their optical thicknesses are estimated by the meteorological model WRF. The single scattering albedo and asymmetry parameter are considered constant in the UV–visible wavelengths and are taken equal to, respectively, 0.99 and 0.85 for the three types of clouds (Madronich and Flocke, 1998). It should be noted that changes in the cloud optical properties due to the activation of aerosols into cloud condensation nuclei are not taken into account in our approach. However, the anticyclonic conditions that prevailed over eastern Europe during the studied period suggest a low impact of clouds on the modelled actinic flux and photolysis rates (Lau and Kim, 2012).

Concerning aerosols, the three optical properties (AOT, SSA and g) are calculated at 200, 300, 400, 500, 600 and 700 nm using the aerosol optical module and then interpolated to the TUV wavelength grid (120–1250 nm). To solve the radiative transfer equation and compute the actinic flux along the atmospheric column, the Eddington approximation has been chosen in TUV as it allows an accurate estimation of modelled radiative fluxes (Joseph et al., 1976).

Recently, Palancar et al. (2013) realized a intercomparison exercise between the TUV model and UV actinic flux measurements over Mexico during the MILAGRO campaign. They highlighted the good performance of the model in reproducing observations both at the surface and in the lower troposphere over this highly polluted area. This validation study gives confidence in our estimation of photolysis rates perturbations by aerosols during the 2010 Russian wildfires presented hereinafter.

2.3 On-line coupling between CHIMERE and TUV Simulation set-up

The methodology developed in this study consists of a one-way and on-line coupling between TUV and CHIMERE. In this approach, the radiative transfer code TUV has been implemented within CHIMERE so that each model runs simultaneously. During the simulation, the aerosol optical properties modelled by CHIMERE for a core-shell mixing (AOT, SSA, g) are used as inputs in TUV in order to take into account the influence of aerosol solar extinction on photolysis rates. Then, the photolysis rates estimated by TUV are in turn used by CHIMERE to calculate the concentrations of photochemical pollutants.

Two simulations are performed for the period of peak fire activity (5–12 August 2010):

(1) In the first one, the attenuation of actinic flux is only due to gases and clouds: CHIMERE–TUV(gases+clouds).

(2) In the second one, the impact of aerosols on solar extinction is added in the photolysis rates calculation: CHIMERE–TUV(gases+clouds+aerosols).

The impact of aerosols on photolysis rates and associated concentrations of photochemical pollutants are then estimated by differentiating the two simulations: (2) - (1).

It should be noted that adding the aerosol impact on solar extinction in simulation (2) induces a computation time increase of 50 % compared to the simulation (1).


We will focus on the aerosol impact on NO₂ and O₃ photolysis rates, which mainly drives the concentration of ozone, NO₂ and OH radicals in the troposphere. Indeed, the major source of ozone is the result of the NO₂ photolysis:



followed by the reaction of O(^3P) with a dioxygen molecule (M is a third body favouring the reaction):



Given that reaction (3) is rapid, the formation rate of ozone is mainly determined by the constant rate $J[\text{NO}_2]$. In parallel, the major sink of ozone during daytime is its photodissociation following the reaction:



$\text{O}(^1\text{D})$ will rapidly react with a water molecule to form OH radicals:



The latter reaction is a major source of OH radicals in the troposphere. They are involved in the formation of secondary particles as oxidants of their gaseous precursors. For example, they contribute to the oxidation of SO_2 , NO_2 and COV, which can result in the formation of, respectively, sulphate, nitrate and secondary organic aerosols. The sequence of reactions (R1 to R4) is generally initiated by the reaction of various VOC with the OH radical.

15 3 Results and discussion

3.1 Regional impact of the 2010 Russian wildfires on the formation of ozone and nitrogen dioxide

During the wildfire episode, the important concentrations of scattering aerosols have affected significantly the UV-visible solar radiation in terms of intensity and spatial extent. Figures 2a-b and 3a-b report the diurnal-averaged daytime average percentage changes in near surface photolysis rates of NO_2 and O_3 , respectively. Changes are shown to be negative over the entire area with mean diurnal-daytime values between -2 % and -50 % and closely follow the AOT spatial features (see Figures 1a-b). It is interesting to note that the impact of the aerosol solar extinction is more pronounced

25 for $J[\text{O}_3]$ than for $J[\text{NO}_2]$, for each day of the studied period. For both photolysis rate, the largest reduction is simulated along the transport of the aerosol plume (20–50 %). The photochemistry over the Moscow region has also been affected, especially during the arrival of the aerosol plume between 6 and 10 August 2010. This point will be discussed in further details hereafter. These modelled changes obtained here are comparable with the recent study of Real and Sartelet (2011) in which they simulate, by using the chemistry-transport model Polyphemus-Polair3D coupled with the radiative transfer code Fast-J, a reduction of $J[\text{NO}_2]$ and $J[\text{O}_3]$ reaching 30 % (in monthly mean)

5 during summer 2001 over European regions characterized by elevated AOT (0.6–0.7 at 550 nm). Also, Hodzic et al. (2007) simulated a 15–30 % NO_2 photolysis reduction during the 2003 European summer heatwave in case of absorbing biomass burning aerosols ($\text{AOT}(550 \text{ nm}) = 0.7\text{--}0.8$, $\text{SSA}(532 \text{ nm}) = 0.83\text{--}0.87$).

According to equations 2 to 4, the alteration of $J[\text{NO}_2]$ and $J[\text{O}_3]$ in the presence of

10 aerosols suggests, in turn, a modification of their concentrations near the surface. Figure 4 gives an example of these corresponding changes during the 8 August for NO_2 and O_3 . Some notable modification of the NO_2 concentration is simulated, as it is mainly driven during daytime by its photolytic destruction (see equation 2). The important diminution of $J[\text{NO}_2]$ due to aerosols (between 2 and 50 %) leads to an increase of

15 its near surface concentration reaching, in average, 3 to 15 %. Concerning ozone, its daytime concentration is influenced by both variations of $J[\text{NO}_2]$ (source of ozone, see equations 2 and 3) and variations of $J[\text{O}_3]$ (sink of ozone, see equation 4). We can deduce from figure 4 that the influence of $J[\text{NO}_2]$ reduction seems to slightly dominate the influence of $J[\text{O}_3]$ reduction, resulting in a decrease of the near surface concentration

20 of ozone between 1 % and 12 %. These results are comparable to those obtained by Real and Sartelet (2011) who simulated a 4–8 % reduction of near-surface ozone concentration (for July 2001) over areas where the decrease of $J[\text{NO}_2]$, due to the aerosol solar extinction, reached 30 %. Also, Mena-Carrasco et al. (2009) highlighted, by combining STEM-2K3 model experiments and aircraft observations from the MILAGRO

25 campaign during March 2006, a 40 % attenuation of $J[\text{NO}_2]$ associated to an intense

aerosol plume over Mexico City, resulting in a 5–10 % diminution of ozone production. Such impacts are however less pronounced than the ones obtained over some highly polluted Asian regions. For example, Bian et al. (2007) and Wai and Tanner (2010) showed, over China, a reduction of maximum ozone concentration reaching 30 % to 70 % associated to similar aerosol loadings as obtained in our study ($1 < \text{AOT}(550 \text{ nm}) < 2.5$). The low absorbing properties of the Russian smoke plume (P  r   et al., 2014) could be a reason for such a behaviour.

To further investigate aerosol feedback on the ozone cycle and the formation of secondary particles, we will now focus our study on the Moscow region where the aerosol solar extinction is pronounced, especially during the aerosol plume overpass (5–9 August 2010). Gaseous and particulate measurements from the Moscow air quality station will also be used in the analysis.

3.2 Impact of the 2010 Russian wildfires on the photochemistry over Moscow

On Figure 5, the impact of particles on the formation of ozone and nitrogen dioxide over Moscow is discussed in terms of ~~percentage changes (at midday) daytime~~ average percentage changes in their near-surface photolysis frequencies and concentrations as a function of modelled AOT (440 nm). As expected, changes appear to have a good linearity with AOT (440 nm) with a correlation of ~~0.960.90–0.98.95~~, i.e. modifications become more important when the aerosol loading increases. As shown previously over the entire area, $J[\text{O}_3]$ is more sensitive to the presence of particles (reduction of about ~~19–10~~ % per unit of AOT) than $J[\text{NO}_2]$ (reduction of about ~~10~~ 6 % per unit of AOT). These modifications of photolysis rates result in an increase of the ground NO_2 concentration of ~~4–3~~ % (per unit of AOT) ~~and a small reduction of ground O_3 of~~. Response of the ozone concentration under the aerosol radiative influence is more complex. Indeed, ozone formation is driven by two major precursors: nitrogen oxides ($\text{NO}_x = \text{NO} + \text{NO}_2$) and volatile organic compounds (VOC) in a complex photochemistry. However, it is possible to identify two regimes of ozone formation

by looking at the ratio between the concentrations of VOC and NO_x : A NO_x-limited and a NO_x-saturated regime (Seinfeld and Pandis, 1998) . Figure 6 indicates that the simulated photochemical regime was characterized by a NO_x-saturated situation over Moscow during the studied period. In this case, inclusion of the aerosol radiative impact on photochemistry leads to two antagonist responses : (1) Increase of NO_x concentration trough the reduction of their photolysis is unfavourable to ozone formation in a NO_x-saturated environment. In parallel, (2) reduction of the ozone photolysis is favourable to its accumulation. The overall impact of aerosols on the ozone concentration is then small due to these two antagonist responses, about 1 % (per unit of AOT). It should be noted that changes modelled here at midday are less important than regional changes previously obtained in diurnal-mean (see Figures 2a-b, 3a-b and 4) as the optical path (and then interaction of aerosols with solar radiation) is minimum at midday, compared to the rest of the day per unit of AOT (see Figure 5).

The influence of aerosols on photochemistry does not only occur at the surface but also in the low troposphere, as illustrated in Figure 6-7. This Figure presents the vertical profile of the ~~diurnal-averaged~~ daytime average percentage changes in the ozone concentration at the north of Moscow (59.9 °N, 37.6 °E) for the 9 August. The aerosol extinction coefficient (in km⁻¹) modelled by CHIMERE and measured by CALIOP at midnight is also indicated. The maximum ozone reduction (4–5 %) occurred below the two first km of the atmosphere within the aerosol plume (modelled extinction of 0.50-0.95 km⁻¹). Above, the influence of aerosols on the ozone formation gradually decrease to become negligible at an altitude of 5–6 km. It should be noted that below an altitude of 2 km (where more that 70 % of the aerosol solar extinction occurs), the modelled aerosol extinction is within the uncertainty range of CALIOP measurements, giving confidence in the estimated impact of aerosols on the ozone reduction.

In terms of model performance, it is interesting to see if an explicit representation of aerosol impact on photolysis rates tend to improve the simulation of the concentration of photochemical pollutants, compared to a simulation without aerosol feedback. For such analyse, statistical comparisons between the near-surface concentrations of NO₂

and O_3 simulated with and without aerosols and measured at Moscow by an air quality station has been made. Results are presented in Tables 1 and 2 for hourly values and daily maximum values, respectively. We can see that, for both configurations, scores for ozone and NO_2 are much lower than the ones usually obtained over western Europe with the CHIMERE model (Honoré et al., 2008), with RMSE of 48–88 $\mu g/m^3$ and correlation of 0.22–0.60. A possible reason could be that the EMEP emission database used in this study is less well documented for eastern Europe. However, Tables 1 and 2 indicates that the inclusion of aerosols in the simulation improves the correlation and slightly reduces biases with measurements for both species and for both hourly and maximum values.

To further investigate the aerosol feedback on the ozone ~~diurnal~~ daytime cycle, Figure 7–8 displays the temporal evolution (between 5 and 12 August 2010) of the near-surface ozone concentration (in $\mu g/m^3$) modelled with and without aerosol feedback along with corresponding observations at the Moscow monitoring station. ~~The European alert threshold of 240 $\mu g/m^3$ is also indicated by a straight line. First, figure 7–~~ This figure confirms the moderate overall impact of the aerosol solar extinction on the ozone production, with a maximum diminution reaching 7–10 % during the aerosol plume overpass (7, 8 and 9 August), which leads to slightly reduce the bias between model and observations. ~~For the other days, the alteration of photolysis rates by aerosols is also present but a similar impact on both the sink and source of ozone is simulated, leading to a negligible impact on the ozone production.~~ Depending on the day, the model simulates lower or higher hourly values compared to observations. As indicated previously, uncertainties on the EMEP emission database for this specific region could be a possible reason for these biases. Also, the model is shown to overestimate night-time concentrations over the period, which could be due to uncertainties in correctly estimating its dry deposition and titration by NO , as previously highlighted by Honoré et al. (2008) over Western Europe. It should be noted that a inadequate representation of the nocturnal air ventilation could also be a reason for such a model bias. Second, Figure

7 indicates that the European alert threshold of $240 \mu\text{g}/\text{m}^3$ is predicted to be exceeded during 8 August 2010 for the simulation without aerosol impacts, with a maximum modelled concentration reaching $251 \mu\text{g}/\text{m}^3$ in the middle of the afternoon. However, this limit is not exceeded when considering aerosol feedback effects with a reduction of the modelled ozone peak from 251 to $230 \mu\text{g}/\text{m}^3$, in coherence with observations ($160 \mu\text{g}/\text{m}^3$). Hence, taking into account the impact of aerosol solar extinction on photolysis rates in the CHIMERE numerical simulations could play a non-negligible role in term of air quality prediction, keeping in mind that better constrained emissions and nocturnal processes could also greatly improve model scores.

In parallel, the presence of aerosols tends to reduce the oxidising capacity of the atmosphere (through reduction of OH radicals, see equation 5), which leads to decrease the formation of secondary aerosols. As illustrated in figure 89a, the maximum reduction in the near-surface concentrations of sulphates (oxidation product of SO_2) and SOA (oxidation product of COV) occurs on 9 August with diurnal-averaged 8 August with daytime average values of 10 % and 4 %, respectively. For this day, figure 9b shows that these changes are mainly due to a reduced formation of very fine particles, i.e with a diameter comprised between 40 nm (bin 1) and 300 nm (bin 4). The overall impact is then a slightly reduction of the total aerosol mass concentration (PM10) comprised between 1 and 2 % over the entire period (Figure 9a). These results are comparable to the findings of Real and Sartelet (2011) and Li et al. (2011b) who showed a 5–10 % reduction of the formation of secondary aerosols due to the aerosol solar extinction, in case of intense particulate pollution over, respectively, Europe and Mexico City. As for NO_2 and O_3 , including the optical effect of aerosols in the photolysis calculations slightly improves the formation of secondary inorganic species in the CHIMERE model with a RMSE systematically reduced (see Table 3). The overestimation of sulphates levels is decreased and the simulated concentrations of ammonium and nitrates get close to the observed one. This result suggests that taking into account the aerosol solar extinction in the photolysis calculation gives an added value in the capacity of the

[model to reproduce the photochemistry under polluted environments.](#)

4 Conclusions

In the present study, we have developed an on-line coupling between the chemistry-transport model CHIMERE (complemented by an aerosol optical module) and the radiative transfer code TUV to study the impact of aerosol solar extinction on the photochemistry over eastern Europe during the 2010 wildfires episode. Simulations from 5 to 12 August 2010, corresponding to the peak of fire activity, have been performed with and without aerosol impact on photolysis rates and concentrations of gaseous and particulate pollutants. Large areas were affected by important concentrations of particles, with modeled AOT (440 nm) above 2 along the transport of the aerosol plume. A previous evaluation of the modeled optical properties during this specific episode (P  r   et al., 2014) showed good model performance in simulating both the magnitude and spectral dependence of the aerosol optical thickness and single scattering albedo.

The impact of aerosols on photolysis rates is shown to be regionally significant with a mean reduction of $J[\text{NO}_2]$ and $J[\text{O}_3]$ comprised between 2 % and 50 %, the maximum reduction being modelled in the aerosol plume. These modifications of photolysis frequencies result in an regional increase in the daytime concentration of NO_2 of 3–15 % and a decrease in the O_3 production near the surface comprised between 1 % and 12 % during 8 August.

The photochemistry over the Moscow region has also been affected, especially during the arrival of the aerosol plume between 6 and 10 August 2010. Over this area, results indicate that $J[\text{O}_3]$ is more sensitive to the presence of particles (reduction of about ~~19~~10 % per unit of AOT) than $J[\text{NO}_2]$ (reduction of about ~~10~~6 % per unit of AOT), resulting in an increase of the ground NO_2 concentration of ~~4~~3 % (per unit of AOT) and a small reduction of ozone of ~~2~~1 % (per unit of AOT). The photochemistry is shown to be impacted along the boundary layer where aerosols are located with, for example, a

25 4–5 % O₃ reduction modelled during 9 August within the first two km of the atmosphere. In terms of model performance, comparisons of simulations with air quality measurements at Moscow indicate that an explicit representation of aerosols interaction with photolysis rates tend to improve the estimation of the near-surface concentration of ozone and nitrogen dioxide, with some potential consequence for air quality prediction. For instance, the O₃ peak reduction when including aerosol feedback induces a non-negligible difference in the predicted exceedance of alert threshold during 8 August compared to the simulation without aerosol feedback, in coherence with measurements. In addition, the impact of aerosols on photolysis rates is shown to have an influence on the formation of secondary aerosols, through the modification of the OH concentration. Over Moscow, the aerosol plume tends to decrease the daytime concentrations of sulphates and secondary organics up to 4–10 %, which result in a small reduction of the total particulate concentration (PM₁₀) of 1–2 % on average over the period. The results presented in this work, issued from a modelling exercise, are consistent with those obtained in recent studies by combining model experiments and different sets of observations. In terms of model performance, comparisons of simulations with air quality measurements at Moscow indicate that an explicit representation of aerosols interaction with photolysis rates tend to improve the estimation of the near-surface concentration of ozone and nitrogen dioxide as well as the formation of inorganic aerosol species such as ammonium, nitrates and sulphates.

15 Recently, it has been suggested that some organic aerosols can absorb solar radiation, especially at the shorter visible and UV wavelengths (Zhong and Jang, 2011; Saleh et al., 2014). The methodology developed in this study provides a powerful tool to investigate the role of enhanced UV absorption by secondary organics on photo-chemistry at regional and urban scale.

20

Acknowledgements. Authors are grateful to the CRI at Lille 1, François Thieuleux at LOA and Anthony Ung at INERIS for their technical support. Olga Kislova at Mosecomonitoring (Moscow) is acknowledged for providing the surface air quality measurements used in this study. Authors

25 would also like to thank Isabelle Chiapello at LOA for her fruitful scientific discussion. The CNES french spatial agency is acknowledged for its financial support.

References

- Amin, N.: Effect of ozone on the relative yield of rice crop in Japan evaluated based on monitored concentrations, *Water, Air and Soil Pollution*, 225, 1–9, 2014.
- Beelen, R., Raaschou-Nielsen, O., Stafoggia, M., and co authors: Effects of long-term exposure to air pollution on natural-cause mortality: an analysis of 22 European cohorts within the
- 5 multicentre ESCAPE project, *The Lancet*, 383, 785–795, 2014.
- Bessagnet, B., Hodzic, A., Vautard, R., Beekmann, M., Cheinet, S., Honoré, C., Liousse, C., and Rouil, L.: Aerosol modeling with CHIMERE — Preliminary evaluation at the continental scale, *Atmospheric Environment*, 38, 2803–2817, 2004.
- Bessagnet, B., Menut, L., Curci, G., Hodzic, A., Guillaume, B., Liousse, C., Moukhtar, S.,
- 10 Pun, B., Seigneur, C., and Schulz, M.: Regional modeling of carbonaceous aerosols over Europe—focus on secondary organic aerosols, *Journal of Atmospheric Chemistry*, 61, doi:10.1007/s10874–009–9129–2, 2008.
- Bian, H., Han, S., Tie, X., Sun, M., and Liu, A.: Evidence of impact of aerosols on surface ozone concentration in Tianjin, China, *Atmospheric Environment*, 41, 4672–4681, 2007.
- 15 Chen, F. and Dudhia, J.: Coupling an advanced land-surface / hydrology model with the Penn State / NCAR MM5 modeling system. Part 1: Model description and implementation, *Mon. Wea. Rev.*, 129, 569–585, 2001.
- Chubarova, N., Nezval, Y., Sviridenkov, I., Smirnov, A., and Slutsker, I.: Smoke aerosol and its radiative effects during extreme fire event over Central Russia in summer 2010, *Atmos. Meas. Tech.*, 5, 557–568, 2012.
- 20 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmospheric Chemistry and Physics*, 6, 3181–3210, 2006.
- Hauglustaine, D. A., Hourdin, F., Jourdain, L., Filiberti, M. A., Walters, S., Lamarque, J. F., and Holland, E. A.: Interactive chemistry in the Laboratoire de Météorologie Dynamique general
- 25 circulation model: description and background tropospheric chemistry evaluation, *Journal of Geophysical Research*, 109, d04314.doi:10.1029/2003JD003957, 2004.

- Hodzic, A., Madronich, S., Bohn, B., Massie, S., Menut, L., and Wiedinmyer, C.: Wildfire particulate matter in Europe during summer 2003: meso-scale modeling of smoke emissions, transport and radiative effects, *Atmospheric Chemistry and Physics*, 7, 4043–4064, 2007.
- Hong, S. Y.: Stable boundary layer mixing in a vertical diffusion scheme, *The Korea Meteor. Soc.*, fall conference, Seoul, Korea, Oct. 25-26, 2007.
- Hong, S. Y., Noh, Y., and Dudhia, J.: A new vertical diffusion package with an explicit treatment of entrainment processes, *Monthly Weather Review*, 134, 2318–2341, 2006.
- Honoré, C., Rouil, L., Vautard, R., Beekmann, M., Bessagnet, B., Dufour, A., Elichegaray, C., Flaud, J. M., Malherbe, L., Meleux, F., Menut, L., Martin, D., Peuch, A., Peuch, V. H., and Poisson, N.: Predictability of European air quality: Assessment of 3 years of operational forecasts and analyses by the PREV'AIR system, *Journal of Geophysical Research*, 113, D04301, doi:10.1029/2007JD008761, 2008.
- Horowitz, L. W., Walters, S., Mauzeralles, D. Z., Emmonds, L. K., Rash, P. J., Granier, C., Tie, X., Lamarque, J. F., and Schultz, M. G.: A global simulation of tropospheric ozone and related tracers : Description and evaluation of MOZART, version 2, *Journal of Geophysical Research*, 108, 108(D24), 4784, 2003.
- Hunova, I., Maly, M., Rezacova, J., and Branis, M.: Association between ambient ozone and health outcomes in Prague, *International archives of occupational and environmental health*, 86, 89–97, 2014.
- Jenkin, M. E. and Clemitshaw, K. C.: Ozone and other secondary photochemical pollutants: chemical processes governing their formation in the planetary boundary layer, *Atmospheric Environment*, 34, 2499–2527, 2000.
- Joseph, J. H., Wiscombe, W. J., and Weinman, J. A.: The delta-Eddington approximation for radiative flux transfer, *Journal of the Atmospheric Sciences*, 33, 2452–2459, 1976.
- Junker, C. and Lioussé, C.: A global emission inventory of carbonaceous aerosol from historic record of fossil fuel and biofuel consumption for the period 1860-1997, *Atmospheric Chemistry and Physics*, 8, 1195–1207, 2008.
- Kain, J. S.: The Kain-Fritsch convective parameterization: An update, *J. Appl. Meteor.*, 43, 170–181, 2004.
- Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J. J., Razinger, M., Schultz, M. G., Suttie, M., and van der Werf, G. R.: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power, *Biogeosciences*, 9, 527–554, 2012.

- Konovalov, I. B., Beekmann, M., Kuznetsova, I. N., Yurova, A., and Zvyagintsev, A. M.: Atmospheric impacts of the 2010 Russian wildfires: integrating modelling and measurements of an extreme air pollution episode in the Moscow region, *Atmospheric Chemistry and Physics*, 11, 10 031–10 056, 2011.
- Lau, W. K. M. and Kim, K.-M.: The 2010 Pakistan flood and Russian heat wave: teleconnection of hydrometeorological extremes, *J. Hydrometeor.*, 13, 392–403, 2012.
- Lesins, G., Chylek, P., and Lohmann, U.: A study of internal and external mixing scenarios and its effect on aerosol optical properties and direct radiative forcing, *Journal of Geophysical Research*, 107, 4094, 2002.
- Li, G., Bei, N., Tie, X., and Molina, L. T.: Aerosol effects on the photochemistry in Mexico City during MCMA-2006/MILAGRO campaign, *Atmospheric Chemistry and Physics*, 11, 5169–5182, 2011b.
- Li, J., Wang, Z., Wang, X., Yamaji, K., Takigawa, M., Kanaya, Y., Pochanart, P., Liu, Y., Irie, H., Hu, B., Tanimoto, H., and Akimoto, H.: Impacts of aerosols on summertime tropospheric photolysis frequencies and photochemistry over Central Eastern China, *Atmospheric Environment*, 45, 1817–1829, 2011a.
- Lou, S., Liao, H., and Zhu, B.: Impacts of aerosols on surface-layer ozone concentrations in China through heterogeneous reactions and changes in photolysis rates, *Atmospheric Environment*, 85, 123–138, 2014.
- Madronich, S. and Flocke, S.: The role of solar radiation in atmospheric chemistry, *Handbook of Environmental Chemistry*, pp. 1–26, Springer-Verlag, Heidelberg, Germany, 1998.
- McMeeking, G. R., Morgan, W. T., Flynn, M., Highwood, E. J., Turnbull, K., Haywood, J., and Coe, H.: Black carbon aerosol mixing state, organic aerosols and aerosol optical properties over the United Kingdom, *Atmospheric Chemistry and Physics*, 11, 9037–9052, 2011.
- Mena-Carrasco, M., R. Carmichael, G., Campbell, J. E., Zimmerman, D., Tang, Y., Adhikary, B., D'allura, A., Molina, L. T., Zavala, M., Flocke, A. G. F., Campos, T., Weinheimer, A. J., Shetter, R., Apel, E., Montzka, D. D., Knapp, D. J., and Zheng, W.: Assessing the regional impacts of Mexico City emissions on air quality and chemistry, *Atmospheric Chemistry and Physics*, 9, 3731–3743, 2009.
- Menut, L., Bessagnet, B., Khvorostyanov, D., Beekmann, M., Blond, N., Colette, A., Coll, I., Curci, G., Foret, G., Hodzic, A., Mailler, S., Meleux, F., Monge, J.-L., Pison, I., Siour, G., Turquety, S., Valari, M., Vautard, R., and Vivanco, M. G.: CHIMERE 2013: a model for regional atmospheric composition modelling, *Geosci. Model Dev.*, 6, 981–1028, 2013.

- 30 Miller, M. R., Shaw, C. A., and Langrish, J. P.: From particles to patients: oxidative stress and the cardiovascular effects of air pollution, *Future Cardiology*, 8, 577–602, 2012.
- Nie, W., Ding, A. J., Xie, Y. N., Xu, Z., Mao, H., Kerminen, V. M., Zheng, L. F., Qi, X. M., Huang, X., Yang, X. Q., Sun, J. N., Herrmann, E., Petäjä, T., Kulmala, M., and Fu, C. B.: Influence of biomass burning plumes on HONO chemistry in eastern China, *Atmospheric Chemistry and Physics*, 15, 1147–1159, 2015.
- 5 Palancar, G. G., Lefer, B. L., Hall, S. R., Shaw, W. J., Corr, C. A., Herndon, S. C., Slusser, J. R., and Madronich, S.: Effect of aerosols and NO₂ concentration on ultraviolet actinic flux near Mexico City during MILAGRO: measurements and model calculations, *Atmospheric Chemistry and Physics*, 13, 1011–1022, 2013.
- Péré, J. C., Mallet, M., Bessagnet, B., and Pont, V.: Evidence of the aerosol core-shell mixing state over Europe during the heat wave of summer 2003 by using CHIMERE simulations and AERONET inversions, *Geophysical Research Letters*, 36, doi:10.1029/2009GL037334, 2009.
- 10 Péré, J. C., Mallet, M., Pont, V., and Bessagnet, B.: Evaluation of an aerosol optical scheme in the chemistry-transport model CHIMERE, *Atmospheric Environment*, 44, 3688–3699, 2010.
- Péré, J. C., Mallet, M., Pont, V., and Bessagnet, B.: Impact of the aerosol direct radiative forcing (ADRF) on the radiative budget, surface heat fluxes and atmospheric dynamics during the heatwave of summer 2003 over Western Europe. A modelling study., *Journal of Geophysical Research*, 116, d23119, doi:10.1029/2011JD016240, 2011.
- 15 Péré, J. C., Bessagnet, B., Mallet, M., Waquet, F., Chiapello, I., Minvielle, F., Pont, V., and Menut, L.: Direct radiative effect of the Russian wildfires and its impact on air temperature and atmospheric dynamics during August 2010, *Atmospheric Chemistry and Physics*, 14, 1999–2013, 2014.
- 20 Popovicheva, O., Kistler, M., Kireeva, E., Persiantseva, N., Timofeev, M., Kopeikin, V., and Kasper-Giebl, A.: Physicochemical characterization of smoke aerosol during large-scale wildfires: Extreme event of August 2010 in Moscow, *Atmospheric Environment*, 96, 405–414, 2014.
- 25 Real, E. and Sartelet, K.: Modeling of photolysis rates over Europe: impact on chemical gaseous species and aerosols, *Atmospheric Chemistry and Physics*, 11, 1711–1727, 2011.
- Saleh, R., Robinson, E. S., Tkacik, D. S., Ahern, A. T., Liu, S., Aiken, A. C., Sullivan, R. C., Presto, A. A., Dubey, M. K., Yokelson, R. J., Donahue, N. M., and Robinson, A. L.: Brownness

- 30 of organics in aerosols from biomass burning linked to their black carbon content, *Nature Geoscience*, 7, 647–650, 2014.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics. From air pollution to climate change*, Wiley-Interscience publication, 1998.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics. From air pollution to climate change (2nd Edition)*, Wiley-Interscience publication, 2006.
- 5 Slade, J. H. and Knopf, D. A.: Heterogeneous OH oxidation of biomass burning organic aerosol surrogate compounds: assessment of volatilisation products and the role of OH concentration on the reactive uptake kinetics, *Phys. Chem. Chem. Phys.*, 15, 5898–5915, 2013.
- Turquety, S.: *The Atmospheric Impact of Wildfires. Fire Phenomena and the Earth System: An Interdisciplinary Guide to Fire Science*, Wiley Blackwell, 2013.
- 10 Vautard, R., Beekmann, M., Roux, J., and Gombert, D.: Validation of a hybrid forecasting system for the ozone concentrations over the Paris area, *Atmospheric Environment*, 14, 2449–2461, 2001.
- Vautard, R., Bessagnet, B., Chin, M., and Menut, L.: On the contribution of natural aeolian sources to particulate matter concentrations in Europe: Testing hypotheses with a modeling approach, *Atmospheric Environment*, 39, 3291–3303, 2005.
- Vester, B. P., Ebert, M., Barnert, E. B., Schneider, J., Kandler, K., Schütz, L., and Weinbruch, L.: Composition and mixing state of the urban background aerosol in the Rhein-Main area (Germany), *Atmospheric Environment*, 41, 6102–6115, 2007.
- 645 Wai, K. M. and Tanner, P. A.: Variations of aerosol properties due to regional source contributions and impacts on ozone levels: a study in a south China city, *Environ. Chem.*, 7, 359–369, 2010.
- Wu, X., Seigneur, C., and Bergström, R.: Evaluation of a sectional representation of size distributions for calculating aerosol optical properties, *Journal of Geophysical Research*, 101, 19 277–19 283, 1996.
- 650 Wu, Z. P. and Wang, Y. P.: Electromagnetic scattering for multilayered spheres: Recursive algorithms, *Radio Science*, 26, 1393–1401, 1991.
- Zhong, M. and Jang, M.: Light absorption coefficient measurement of SOA using a UV–Visible spectrometer connected with an integrating sphere, *Atmospheric Environment*, 45, 4263–4271, 2011.
- 655

Zvyagintsev, A. M., Ivanova, N. S., Blyum, O. B., Kotel'nikov, S. N., Kruchenitskii, G. M., Kuznetsova, I. N., and Lapchenko, V. A.: Ozone content over the Russian Federation in the third quarter of 2010, *Russian Meteorology and Hydrology*, 35, 785–789, 2010.




Fig. 1a. Geographic distribution of the AOT (at midday) over Eastern Europe from 5 to 12 August 2010 modelled by CHIMERE (at 400 nm). Horizontal wind at 850 hPa simulated by WRF is also indicated.




Fig. 1b. Continuation of Figure 1




Fig. 2a. Geographic distribution of the modeled ~~diurnal-averaged~~ daytime average percentage changes in the near-surface $J[\text{NO}_2]$ due to the presence of aerosols.




Fig. 2b. Continuation of Figure 2.




Fig. 3a. Geographic distribution of the modelled ~~diurnal-averaged~~ daytime average percentage changes in the near-surface $J[O_3]$ due to the presence of aerosols.




Fig. 3b. Continuation of Figure 3.




Fig. 4. Geographic distribution of the diurnal-averaged daytime average percentage changes in the near-surface concentration of NO_2 (left) and O_3 (right) for the 8 August 2010, due to the presence of aerosols.




Fig. 5. Daytime average percentage changes (at midday) in the photolysis frequencies and concentrations of nitrogen dioxide and ozone over Moscow as a function of modelled AOT (440 nm).




Fig. 6. Mean VOC to NOx ratio simulated over Moscow during the studied period




Fig. 7. Vertical profile of the ~~diurnal-averaged~~ daytime average percentage changes in the ozone concentration at the north of Moscow (59.9 °N, 37.6 °E) for the 9 August. The aerosol extinction coefficient (in km⁻¹) modelled by CHIMERE and measured by CALIOP at midnight is also indicated.






Fig. 8. Temporal evolution (between 5 and 12 August 2010) of the near-surface ozone concentration (in $\mu\text{g}/\text{m}^3$) modelled with and without aerosol feedback along with corresponding observations at the Moscow monitoring station. The European alert threshold of $240 \mu\text{g}/\text{m}^3$ is also indicated by a straight line.



Diurnal-averaged percentage reduction of the near-surface concentration of sulphates, secondary organic aerosols and PM10 over Moscow due to the aerosol feedback.

Fig. 9. a) Daytime average percentage reduction of the near-surface concentration of sulphates, secondary organic aerosols and PM10 over Moscow due to the aerosol feedback. b) Repartition of this sulphate and SOA mass reduction between the 8 aerosol size bins for the 8 August

Table 1. statistical comparisons between the near-surface concentrations of NO₂ and O₃ simulated with and without aerosols and measured at Moscow by an air quality station. Mod. and Obs. are the period-averaged modelled and observed concentration. Corr. and RMSE are the temporal correlation and the root mean square error.

| | NO ₂ | | | | O ₃ | | | |
|---------|------------------------------|------------------------------|-------|------------------------------|------------------------------|------------------------------|-------|------------------------------|
| | Mod. (μg/m ³) | Obs. (μg/m ³) | Corr. | RMSE (μg/m ³) | Mod. (μg/m ³) | Obs. (μg/m ³) | Corr. | RMSE (μg/m ³) |
| with | 14 | 70 | 0.22 | 66 | 129 | 66 | 0.46 | 82 |
| without | 13 | 70 | 0.21 | 67 | 134 | 66 | 0.42 | 88 |

Table 2. Same as in Table 1 but for daily maximum values

| | NO ₂ | | | | O ₃ | | | |
|---------|------------------------------|------------------------------|-------|------------------------------|------------------------------|------------------------------|-------|------------------------------|
| | Mod. (µg/m ³) | Obs. (µg/m ³) | Corr. | RMSE (µg/m ³) | Mod. (µg/m ³) | Obs. (µg/m ³) | Corr. | RMSE (µg/m ³) |
| with | 32 | 90 | 0.60 | 66 | 180 | 161 | 0.19 | 48 |
| without | 31 | 90 | 0.60 | 67 | 185 | 161 | 0.16 | 54 |

Table 3. Same as in Table 1 but for the near-surface concentrations of ammonium, nitrates and sulphates

| | NH_4^+ | | | | NO_3^- | | | | SO_4^{2-} | | | |
|---------|--------------------------------------|--------------------------------------|-------|--------------------------------------|--------------------------------------|--------------------------------------|-------|--------------------------------------|--------------------------------------|--------------------------------------|-------|--------------------------------------|
| | Mod. ($\mu\text{g}/\text{m}^3$) | Obs. ($\mu\text{g}/\text{m}^3$) | Corr. | RMSE ($\mu\text{g}/\text{m}^3$) | Mod. ($\mu\text{g}/\text{m}^3$) | Obs. ($\mu\text{g}/\text{m}^3$) | Corr. | RMSE ($\mu\text{g}/\text{m}^3$) | Mod. ($\mu\text{g}/\text{m}^3$) | Obs. ($\mu\text{g}/\text{m}^3$) | Corr. | RMSE ($\mu\text{g}/\text{m}^3$) |
| with | 1.09 | 0.86 | 0.48 | 1.03 | 0.14 | 0.12 | 0.20 | 0.15 | 2.24 | 0.48 | 0.23 | 1.80 |
| without | 1.16 | 0.86 | 0.42 | 1.04 | 0.19 | 0.12 | 0.26 | 0.23 | 2.33 | 0.48 | 0.45 | 1.88 |

Influence of the aerosol solar extinction on photochemistry during the 2010 Russian wildfires episode

J. C. Péré¹, B. Bessagnet², V. Pont³, M. Mallet³, and F. Minvielle¹

¹Laboratoire d'Optique Atmosphérique, Université Lille 1, 59655 Villeneuve d'Ascq, France.

²Institut National de l'Environnement Industriel et des Risques, Parc Technologique Alata, 60550 Verneuil en Halatte, France.

³Laboratoire d'Aérologie, Observatoire Midi-Pyrénées, 14 Avenue Edouard Belin, 31400 Toulouse, France.

Correspondence to: J. C. Péré
(jean-christophe.pere@univ-lille1.fr)

Abstract

In this work, impact of aerosol solar extinction on the photochemistry over eastern Europe during the 2010 wildfires episode is discussed for the period from 5 to 12 August 2010, which coincides to the peak of fire activity. The methodology is based on an on-line coupling between the chemistry-transport model CHIMERE (extended by an aerosol optical module) and the radiative transfer code TUV. Results of simulations indicate an important influence of the aerosol solar extinction, in terms of intensity and spatial extent, with a reduction of the photolysis rates of NO_2 and O_3 up to 50 % (in ~~diurnal-averaged~~ daytime average) along the aerosol plume transport. At a regional scale, these changes in photolysis rates lead to a 3–15 % increase in the NO_2 daytime concentration and to an ozone reduction near the surface of 1–12 %. The ozone reduction is shown to occur over the entire boundary layer, where aerosols are located. Also, ~~comparisons of simulations with air quality measurements over Moscow show that the inclusion of the aerosol feedback tends to slightly improve performance of the model in simulating NO_2 and O_3 ground concentrations. In term of air quality prediction, the O_3 peak reduction when including aerosol feedback results in a non-negligible difference in the predicted exceedance of alert threshold compared to the simulation without aerosol feedback, in coherence with measurements. Finally,~~ the total aerosol mass concentration (PM10) is shown to be decreased by 1–2 %, on average during the studied period, caused by a reduced formation of secondary aerosols such as sulphates and secondary organics (4–10 %) when aerosol impact on photolysis rates is included. In terms of model performance, comparisons of simulations with air quality measurements at Moscow indicate that an explicit representation of aerosols interaction with photolysis rates tend to improve the estimation of the near-surface concentration of ozone and nitrogen dioxide as well as the formation of inorganic aerosol species such as ammonium, nitrates and sulphates.

1 Introduction

For several years, it has been well recognized that air pollution of gaseous and particulate origin can have adverse health effects (Miller et al., 2012; Beelen et al., 2014). In consequence, efficient air pollution control strategies have now become a challenge for environmental policies. In the context of air quality monitoring, the ~~exceed~~ exceedance of certain thresholds of pollutant concentrations is a criterion often used by authorities of a country to prevent people from air pollution exposure. In general, the ~~exceed~~ exceedance of these thresholds is evaluated from air quality numerical forecast such as in France where the regional chemistry-transport model CHIMERE (Menut et al., 2013) is used in the French national air quality forecasting and monitoring system known as PREV'AIR (Honoré et al., 2008). Photochemical pollutants (ozone, secondary aerosols,...), which are formed from photo-dissociation of precursors such as nitrogen dioxide (NO₂) and volatile organic compounds (VOC) (Jenkin and Clemitshaw, 2000), are of particular interest for air quality monitoring (Honoré et al., 2008) due to their negative impacts on both environment and human health (Amin, 2014; Hunova et al., 2014).

The key parameter that governs the photo-dissociation of photochemical precursors in the atmosphere is the photolysis rate, which mainly depends on the available actinic flux (Seinfeld and Pandis, 2006). Aerosols are known to have large influence on the available actinic flux by interacting with solar radiation in the ultraviolet-visible wavelengths (Li et al., 2011a,b; Lou et al., 2014). For example, Wai and Tanner (2010) showed, by using a combination of remote sensing observations and chemical-transport model, that aerosol solar extinction could lead to a 7–32 % reduction in maximum ozone concentration over Hong-Kong during highly polluted days. Also, Li et al. (2011b) highlighted, with WRF-CHEM modelling experiments, that changes in photolysis rates due to the presence of particles led to a decrease of about, respectively, 2–17 % and 5–6 % in daytime ozone and secondary aerosols (nitrate, secondary organics) concentrations over Mexico City during the 2006 Megacity Initiative: Local and Global Re-

search Observations (MILAGRO) campaign.

To reduce computational time for operational purpose, one major characteristic of air quality modelling platforms is that impacts of aerosols and clouds on solar radiation are often taken into account as simplified attenuation factors when evaluating the photolysis rates (Honoré et al., 2008; Menut et al., 2013). However, Real and Sartelet (2011) highlighted that simplified parametrization of aerosol impact on photolysis rates could tend to worsen air quality model performance in simulating ozone and particulate concentration, especially under highly polluted environments.

The aim of the present study is to implement, in the chemistry-transport model CHIMERE,

an explicit representation of the alteration of photolysis rates by aerosols and discuss the impact in terms of modelled ozone budget and the formation of secondary aerosols at regional scale. We ~~emphasize~~ focus on a major fire event that occurred in Russia during August 2010 as its episode was characterized by important concentrations of primary and secondary aerosols and large concentrations of ozone, especially over this specific region (Zvyagintsev et al., 2010; Konovalov et al., 2011; Popovichcheva et al., 2014). ~~Then, it~~ Also, the study of Chubarova et al. 2012 clearly shows that, during this

specific wildfire episode, the aerosol optical thickness over the Moscow region was more than three times larger than the one observed during typical August conditions over the period 2001–2010. This suggests that, even if anthropogenic aerosols are present over the studied region, the contribution of smoke aerosols during this specific event is very large. Then, this case study represents an excellent opportunity to discuss how aerosol solar extinction, especially biomass burning particles, can affect photochemistry. Fires can affect atmospheric chemistry in several ways. They emit primary gaseous pollutants (such as CO, OH, NO, NO₂ and volatile organic compounds) that can react in the atmosphere to form ozone and other pollutants (Turquety, 2013) . They also released aerosols that can directly affect air quality or indirectly by acting as a medium in complex heterogeneous reactions (Slade and Knopf, 2013; Nie et al., 2015) . Finally, they can affect the intensity of solar radiation, which in turn could affect photochemistry of the atmosphere. The latter impact is the subject of the present study.

The approach is based on an on-line coupling between the regional model CHIMERE, extended by an aerosol optical module (Péré et al., 2010), and the Tropospheric Ultra-violet and Visible (TUV) radiation model (Madronich and Flocke, 1998). In this methodology, the aerosol optical thickness, single scattering albedo and asymmetry parameter are first modelled by CHIMERE using an aerosol core-shell mixing hypothesis, as in Péré et al. (2009, 2010). This mixing approach has been previously used by Péré et al. (2014) to study the 2010 Russian wildfires direct radiative forcing and its feedback on the regional atmospheric dynamics. Results indicate that it can give a good representation of the absorption properties of particles during this specific period. In a second time, aerosol optical properties are used as inputs in the radiative transfer code TUV to evaluate the impact of aerosol short-wave solar extinction on photolysis rates and the formation of ozone and secondary particles. The advantage of such methodology is the use of two specific state-of-the-art models to explicitly simulate the interaction of physico-chemically resolved aerosols with the actinic flux and the associated impact on modelled photolysis rates and photochemistry.

Section 2 describes the configuration of each model as well as the development of their on-line coupling. In Section 3 are discussed modelled regional changes in the near surface concentrations of NO_2 , O_3 and secondary aerosols over Russia induced by modifications of photolysis rates by smoke aerosols during August 2010. Finally, conclusions and perspectives of future works are given in Section 4.

2 Methodology

2.1 Description of the CHIMERE model

2.1.1 Aerosol module

CHIMERE is a state-of-the-art 3D chemistry transport model that calculates the concentrations of numerous gaseous and particulate pollutants (Vautard et al., 2001; Menut

et al., 2013). The dynamics and gas phase parts of the model is regularly improved (Menut et al., 2013) and its documentation can be downloaded at <http://www.lmd.polytechnique.fr/chimere/>. In this work, the CHIMERE domain has a 30 km horizontal resolution and ranges from 43.40°N to 63.20°N in latitude and from 18.70°E to 57.30°E in longitude. The aerosol part is described by Bessagnet et al. (2004) and is composed of 10 chemical species: sulphates, nitrates, ammonium, primary organic and black carbon (OC and BC), secondary organic aerosols (SOA), sea salt, natural and anthropogenic dust and water. The evolution of aerosols is described with a 8-bins size distribution (from about 40 nm to 10 μ m in diameter) and includes the main physical processes such as nucleation, coagulation, condensation/evaporation, adsorption/desorption, wet and dry deposition and scavenging.

Anthropogenic emissions of gaseous and particulate origin come from the EMEP database. Concerning OC and BC emissions, the inventory of Junker and Liousse (2008) has been used. Natural soil dust are dynamically produced within the domain according to the methodology of Vautard et al. (2005). SOA formation is represented through oxidation processes of relevant precursors of biogenic and anthropogenic origin and gas particle partitioning schemes (Bessagnet et al., 2008). VOC and NO emissions from vegetation are calculated using the Model of Emissions of Gases and Aerosols from Nature (MEGAN) (Guenther et al., 2006). Aerosols and gases emitted by wildfires affecting Russia during 2010 are taken into account following the work described and validated by Kaiser et al. (2012). It consists in the assimilation of the fire radiative observations from the Moderate Resolution Imaging Spectroradiometer into the Global Fire assimilation System combined with the use of specific combustion rate and emission factors to estimate biomass burning emissions. More information on the methodology as well as a validation study for the 2010 Russian wildfires episode can be found in ~~(Kaiser et al., 2012)~~ [Kaiser et al. \(2012\)](#).

CHIMERE is forced at these boundaries by monthly climatologies, calculated over the 2000–2004 period, of the main gases and particles provided by the MOZART (Horowitz et al., 2003) and LMDzT-INCA global chemistry-transport models (Hauglus-

tainé et al., 2004), respectively. The evaluation study of Péré et al. (2014) showed that the fire inventory of Kaiser et al. (2012) used in this work combined with the CHIMERE model have been shown to well capture the evolution of the Russian fire plume during the studied period, suggesting a low influence of these boundary conditions. Also, CHIMERE is off-line driven by the Weather Research and Forecasting model (WRF) at a 30 km resolution. The version 3.1 is used in this study with the same configuration as in Péré et al. (2011) . It has 27 vertical levels from 40 m to about 20 km and includes the following parametrizations: the WRF single-moment five-class scheme of Hong et al. (2006) for the microphysics module, the Kain–Fritsch cumulus parametrization (Kain, 2004) , the NOAH land surface module of Chen and Dudhia (2001) and the Yonsei University planetary boundary layer scheme (Hong et al., 2006; Hong, 2007) .

2.1.2 Modelling aerosol optical properties

The calculation of optical properties of particles is the pre-requisite for the evaluation of their impacts on photolysis rates and photochemistry as they provide information on how aerosols will interact with the actinic flux. In that sense, we developed a numerical scheme dedicated to calculate aerosol optical properties from aerosol concentrations, size distribution and chemical composition modelled by CHIMERE. A complete description of this optical module is presented in the work of Péré et al. (2010). To compute the complex refractive index of a particle, the hypothesis of a core-shell mixing has been chosen with a core of primary species (BC, OC and dust) surrounded by a shell of secondary ones (sulphates, nitrates, ammonium, secondary organics) and sea salt and water. This mixing choice is supported by recent studies giving evidence of coatings of secondary particles on black carbon aerosols over Europe (Vester et al., 2007; McMeeking et al., 2011). Also, such mixing has been shown to correctly reproduce the absorbing properties of aerosols during the 2010 Russian wildfire episode (Péré et al., 2014). For each size bin, a volume average procedure is used to calculate the refractive index of the core and the shell (Lesins et al., 2002) which is then used as inputs in the Mie algorithm for n-layered spheres of Wu and Wang (1991) to calculate the scat-

tering and absorption coefficients. It should be noted that the volume of the core and the shell can vary during the simulation in function of the different physical processes influencing aerosol population. The optical properties of the total aerosol distribution needed in radiative transfer modelling, such as the Aerosol Optical Thickness (AOT), Single Scattering Albedo (SSA) and asymmetry parameter (g), are calculated as in Wu et al. (1996).

A detailed evaluation of the optical module for the 2010 Russian wildfire episode by using Aerosol Robotic Network (AERONET) sunphotometers measurements (AOT, SSA, aerosol size distribution) and Polarization and Directionality of the Earth's Reflectances (POLDER) and Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) aerosol extinction data is presented by P  r   et al. (2014). Only a sum-up is given here. As displayed in Figure 1a-b, simulations for the period 5–12 August 2010 (peak of fire activity) show large AOT (at midday) over eastern Europe with modelled values above 2 (at 400 nm) in the aerosol plume. This aerosol plume was transported in an anticyclonic flow from the source region towards Moscow (5-8 August), to the north (9-10 August) and then back to the east (11–12 August). On average, simulations compare well with AOT POLDER data in terms of spatial distribution and intensity (correlation = 0.67–0.77, $-10\% < \text{biases} < 23\%$). The altitude of transport was shown to be below 5 km and comparisons between CHIMERE and CALIOP show good consistency. Figure 1a-b indicates that the simulated plume was advected over Moscow between 5–9 August, which is in accordance with AERONET measurements. During the study period, the aerosol composition was dominated by scattering organic species with modelled elevated SSA (0.97 between 300 nm and 1000 nm) close to AERONET values over Moscow (0.95–0.96 between 440 nm and 1020 nm). [Such low aerosol absorption properties are supported by the study of Chubarova et al. \(2012\) highlighting elevated SSA values during this specific event.](#) Globally, the comparisons between aerosol simulation and sunphotometers data highlighted the ability of the model to give an appropriate representation of the aerosol size distribution and scattering/absorption properties (P  r   et al., 2014), which is the pre-requisite to evaluate its influence on photolysis

rates and the formation of photochemical pollutants.

2.2 Description of the TUV model

TUV is a widely-used state-of-the-art radiative transfer model developed at the National Centre for Atmospheric Research (Madronich and Flocke, 1998). In this study, we used the version 4.6 of the code (released in March 2009) freely available at the website: <http://cprm.acd.ucar.edu/Models/TUV/>. The model calculates the actinic flux and photolysis rates of a large number of photochemical species.

Photolysis is the process breaking the covalent bond of some reactive gaseous species by short-wave solar radiation. This process is very important in the atmosphere as it controls the abundance of numerous air pollutants such as ozone and nitrogen dioxide. The photolysis rate of a given specie J (s^{-1}) is calculated as follows:

$$J(s^{-1}) = \int_{\lambda_1}^{\lambda_2} \sigma(\lambda, T) \cdot \phi(\lambda, T) \cdot F(\lambda) d\lambda \quad (1)$$

where $\sigma(\lambda, T)$ and $\phi(\lambda, T)$ are, respectively, the absorption cross section (cm^2) and the quantum yield of a given molecule, T the air temperature (K) and $F(\lambda)$ the actinic flux between wavelengths λ_1 and λ_2 ($photons.cm^{-2}.s^{-1}.nm^{-1}$). The absorption cross section reflects the probability of collision between a photon and the molecule, while the quantum yield is the probability that the molecule is dissociated after collision with a photon. The dependence of both parameters on the air temperature is calculated by TUV by using the vertical profile of air temperature issued from the meteorological model WRF used to drive CHIMERE.

The actinic flux is calculated by integrating the solar flux over all sphere angles considering 5646 wavelengths between 120 nm and 1250 nm. When going through the

atmosphere, the actinic flux can be attenuated by molecular absorption and diffusion but also by the presence of clouds and aerosols. For clouds and aerosols, the attenuation is calculated by using their respective aerosol optical thickness, single scattering albedo and asymmetry parameter. In TUV, ~~three types of clouds are represented~~ clouds are assumed to be horizontally homogeneous layers and are considered to be of three types : low, middle and high altitude clouds. Altitudes of their bases and tops as well as their optical thicknesses are estimated by the meteorological model WRF. The single scattering albedo and asymmetry parameter are considered constant in the UV–visible wavelengths and are taken equal to, respectively, 0.99 and 0.85 for the three types of clouds (Madronich and Flocke, 1998). It should be noted that changes in the cloud optical properties due to the activation of aerosols into cloud condensation nuclei are not taken into account in our approach. However, the anticyclonic conditions that prevailed over eastern Europe during the studied period suggest a low impact of clouds on the modelled actinic flux and photolysis rates (Lau and Kim, 2012).

Concerning aerosols, the three optical properties (AOT, SSA and g) are calculated at 200, 300, 400, 500, 600 and 700 nm using the aerosol optical module and then interpolated to the TUV wavelength grid (120–1250 nm). To solve the radiative transfer equation and compute the actinic flux along the atmospheric column, the Eddington approximation has been chosen in TUV as it allows an accurate estimation of modelled radiative fluxes (Joseph et al., 1976).

Recently, Palancar et al. (2013) realized a intercomparison exercise between the TUV model and UV actinic flux measurements over Mexico during the MILAGRO campaign. They highlighted the good performance of the model in reproducing observations both at the surface and in the lower troposphere over this highly polluted area. This validation study gives confidence in our estimation of photolysis rates perturbations by aerosols during the 2010 Russian wildfires presented hereinafter.

2.3 On-line coupling between CHIMERE and TUV Simulation set-up

The methodology developed in this study consists of a one-way and on-line coupling between TUV and CHIMERE. In this approach, the radiative transfer code TUV has been implemented within CHIMERE so that each model runs simultaneously. During the simulation, the aerosol optical properties modelled by CHIMERE for a core-shell mixing (AOT, SSA, g) are used as inputs in TUV in order to take into account the influence of aerosol solar extinction on photolysis rates. Then, the photolysis rates estimated by TUV are in turn used by CHIMERE to calculate the concentrations of photochemical pollutants.

Two simulations are performed for the period of peak fire activity (5–12 August 2010):

(1) In the first one, the attenuation of actinic flux is only due to gases and clouds: CHIMERE–TUV(gases+clouds).

(2) In the second one, the impact of aerosols on solar extinction is added in the photolysis rates calculation: CHIMERE–TUV(gases+clouds+aerosols).

The impact of aerosols on photolysis rates and associated concentrations of photochemical pollutants are then estimated by differentiating the two simulations: (2) - (1).

It should be noted that adding the aerosol impact on solar extinction in simulation (2) induces a computation time increase of 50 % compared to the simulation (1).


We will focus on the aerosol impact on NO₂ and O₃ photolysis rates, which mainly drives the concentration of ozone, NO₂ and OH radicals in the troposphere. Indeed, the major source of ozone is the result of the NO₂ photolysis:



followed by the reaction of O(^3P) with a dioxygen molecule (M is a third body favouring the reaction):



Given that reaction (3) is rapid, the formation rate of ozone is mainly determined by the constant rate $J[\text{NO}_2]$. In parallel, the major sink of ozone during daytime is its photodissociation following the reaction:



$\text{O}(^1\text{D})$ will rapidly react with a water molecule to form OH radicals:



The latter reaction is a major source of OH radicals in the troposphere. They are involved in the formation of secondary particles as oxidants of their gaseous precursors. For example, they contribute to the oxidation of SO_2 , NO_2 and COV, which can result in the formation of, respectively, sulphate, nitrate and secondary organic aerosols. The sequence of reactions (R1 to R4) is generally initiated by the reaction of various VOC with the OH radical.

15 3 Results and discussion

3.1 Regional impact of the 2010 Russian wildfires on the formation of ozone and nitrogen dioxide

During the wildfire episode, the important concentrations of scattering aerosols have affected significantly the UV-visible solar radiation in terms of intensity and spatial extent. Figures 2a-b and 3a-b report the diurnal-averaged daytime average percentage changes in near surface photolysis rates of NO_2 and O_3 , respectively. Changes are shown to be negative over the entire area with mean diurnal-daytime values between -2 % and -50 % and closely follow the AOT spatial features (see Figures 1a-b). It is interesting to note that the impact of the aerosol solar extinction is more pronounced

25 for $J[\text{O}_3]$ than for $J[\text{NO}_2]$, for each day of the studied period. For both photolysis rate, the largest reduction is simulated along the transport of the aerosol plume (20–50 %). The photochemistry over the Moscow region has also been affected, especially during the arrival of the aerosol plume between 6 and 10 August 2010. This point will be discussed in further details hereafter. These modelled changes obtained here are comparable with the recent study of Real and Sartelet (2011) in which they simulate, by using the chemistry-transport model Polyphemus-Polair3D coupled with the radiative transfer code Fast-J, a reduction of $J[\text{NO}_2]$ and $J[\text{O}_3]$ reaching 30 % (in monthly mean)

5 during summer 2001 over European regions characterized by elevated AOT (0.6–0.7 at 550 nm). Also, Hodzic et al. (2007) simulated a 15–30 % NO_2 photolysis reduction during the 2003 European summer heatwave in case of absorbing biomass burning aerosols ($\text{AOT}(550 \text{ nm}) = 0.7\text{--}0.8$, $\text{SSA}(532 \text{ nm}) = 0.83\text{--}0.87$).

According to equations 2 to 4, the alteration of $J[\text{NO}_2]$ and $J[\text{O}_3]$ in the presence of

10 aerosols suggests, in turn, a modification of their concentrations near the surface. Figure 4 gives an example of these corresponding changes during the 8 August for NO_2 and O_3 . Some notable modification of the NO_2 concentration is simulated, as it is mainly driven during daytime by its photolytic destruction (see equation 2). The important diminution of $J[\text{NO}_2]$ due to aerosols (between 2 and 50 %) leads to an increase of

15 its near surface concentration reaching, in average, 3 to 15 %. Concerning ozone, its daytime concentration is influenced by both variations of $J[\text{NO}_2]$ (source of ozone, see equations 2 and 3) and variations of $J[\text{O}_3]$ (sink of ozone, see equation 4). We can deduce from figure 4 that the influence of $J[\text{NO}_2]$ reduction seems to slightly dominate the influence of $J[\text{O}_3]$ reduction, resulting in a decrease of the near surface concentration

20 of ozone between 1 % and 12 %. These results are comparable to those obtained by Real and Sartelet (2011) who simulated a 4–8 % reduction of near-surface ozone concentration (for July 2001) over areas where the decrease of $J[\text{NO}_2]$, due to the aerosol solar extinction, reached 30 %. Also, Mena-Carrasco et al. (2009) highlighted, by combining STEM-2K3 model experiments and aircraft observations from the MILAGRO

25 campaign during March 2006, a 40 % attenuation of $J[\text{NO}_2]$ associated to an intense

aerosol plume over Mexico City, resulting in a 5–10 % diminution of ozone production. Such impacts are however less pronounced than the ones obtained over some highly polluted Asian regions. For example, Bian et al. (2007) and Wai and Tanner (2010) showed, over China, a reduction of maximum ozone concentration reaching 30 % to 70 % associated to similar aerosol loadings as obtained in our study ($1 < \text{AOT}(550 \text{ nm}) < 2.5$). The low absorbing properties of the Russian smoke plume (P  r   et al., 2014) could be a reason for such a behaviour.

To further investigate aerosol feedback on the ozone cycle and the formation of secondary particles, we will now focus our study on the Moscow region where the aerosol solar extinction is pronounced, especially during the aerosol plume overpass (5–9 August 2010). Gaseous and particulate measurements from the Moscow air quality station will also be used in the analysis.

3.2 Impact of the 2010 Russian wildfires on the photochemistry over Moscow

On Figure 5, the impact of particles on the formation of ozone and nitrogen dioxide over Moscow is discussed in terms of ~~percentage changes (at midday) daytime~~ average percentage changes in their near-surface photolysis frequencies and concentrations as a function of modelled AOT (440 nm). As expected, changes appear to have a good linearity with AOT (440 nm) with a correlation of ~~0.960.90–0.98.95~~, i.e. modifications become more important when the aerosol loading increases. As shown previously over the entire area, $J[\text{O}_3]$ is more sensitive to the presence of particles (reduction of about ~~19–10~~ % per unit of AOT) than $J[\text{NO}_2]$ (reduction of about ~~10~~ 6 % per unit of AOT). These modifications of photolysis rates result in an increase of the ground NO_2 concentration of ~~4–3~~ % (per unit of AOT) ~~and a small reduction of ground O_3 of~~. Response of the ozone concentration under the aerosol radiative influence is more complex. Indeed, ozone formation is driven by two major precursors: nitrogen oxides ($\text{NO}_x = \text{NO} + \text{NO}_2$) and volatile organic compounds (VOC) in a complex photochemistry. However, it is possible to identify two regimes of ozone formation

by looking at the ratio between the concentrations of VOC and NO_x : A NO_x-limited and a NO_x-saturated regime (Seinfeld and Pandis, 1998) . Figure 6 indicates that the simulated photochemical regime was characterized by a NO_x-saturated situation over Moscow during the studied period. In this case, inclusion of the aerosol radiative impact on photochemistry leads to two antagonists responses : (1) Increase of NO_x concentration trough the reduction of their photolysis is unfavourable to ozone formation in a NO_x-saturated environment. In parallel, (2) reduction of the ozone photolysis is favourable to its accumulation. The overall impact of aerosols on the ozone concentration is then small due to these two antagonist responses, about 1 % (per unit of AOT). It should be noted that changes modelled here at midday are less important than regional changes previously obtained in diurnal-mean (see Figures 2a-b, 3a-b and 4) as the optical path (and then interaction of aerosols with solar radiation) is minimum at midday, compared to the rest of the day per unit of AOT (see Figure 5).

The influence of aerosols on photochemistry does not only occur at the surface but also in the low troposphere, as illustrated in Figure 6-7. This Figure presents the vertical profile of the diurnal-averaged daytime average percentage changes in the ozone concentration at the north of Moscow (59.9 °N, 37.6 °E) for the 9 August. The aerosol extinction coefficient (in km⁻¹) modelled by CHIMERE and measured by CALIOP at midnight is also indicated. The maximum ozone reduction (4–5 %) occurred below the two first km of the atmosphere within the aerosol plume (modelled extinction of 0.50-0.95 km⁻¹). Above, the influence of aerosols on the ozone formation gradually decrease to become negligible at an altitude of 5–6 km. It should be noted that below an altitude of 2 km (where more that 70 % of the aerosol solar extinction occurs), the modelled aerosol extinction is within the uncertainty range of CALIOP measurements, giving confidence in the estimated impact of aerosols on the ozone reduction.

In terms of model performance, it is interesting to see if an explicit representation of aerosol impact on photolysis rates tend to improve the simulation of the concentration of photochemical pollutants, compared to a simulation without aerosol feedback. For such analyse, statistical comparisons between the near-surface concentrations of NO₂

and O_3 simulated with and without aerosols and measured at Moscow by an air quality station has been made. Results are presented in Tables 1 and 2 for hourly values and daily maximum values, respectively. We can see that, for both configurations, scores for ozone and NO_2 are much lower than the ones usually obtained over western Europe with the CHIMERE model (Honoré et al., 2008), with RMSE of 48–88 $\mu g/m^3$ and correlation of 0.22–0.60. A possible reason could be that the EMEP emission database used in this study is less well documented for eastern Europe. However, Tables 1 and 2 indicates that the inclusion of aerosols in the simulation improves the correlation and slightly reduces biases with measurements for both species and for both hourly and maximum values.

To further investigate the aerosol feedback on the ozone ~~diurnal~~ daytime cycle, Figure 7–8 displays the temporal evolution (between 5 and 12 August 2010) of the near-surface ozone concentration (in $\mu g/m^3$) modelled with and without aerosol feedback along with corresponding observations at the Moscow monitoring station. ~~The European alert threshold of 240 $\mu g/m^3$ is also indicated by a straight line. First, figure 7–~~ This figure confirms the moderate overall impact of the aerosol solar extinction on the ozone production, with a maximum diminution reaching 7–10 % during the aerosol plume overpass (7, 8 and 9 August), which leads to slightly reduce the bias between model and observations. ~~For the other days, the alteration of photolysis rates by aerosols is also present but a similar impact on both the sink and source of ozone is simulated, leading to a negligible impact on the ozone production.~~ Depending on the day, the model simulates lower or higher hourly values compared to observations. As indicated previously, uncertainties on the EMEP emission database for this specific region could be a possible reason for these biases. Also, the model is shown to overestimate night-time concentrations over the period, which could be due to uncertainties in correctly estimating its dry deposition and titration by NO , as previously highlighted by Honoré et al. (2008) over Western Europe. It should be noted that a inadequate representation of the nocturnal air ventilation could also be a reason for such a model bias. Second, Figure

7 indicates that the European alert threshold of $240 \mu\text{g}/\text{m}^3$ is predicted to be exceeded during 8 August 2010 for the simulation without aerosol impacts, with a maximum modelled concentration reaching $251 \mu\text{g}/\text{m}^3$ in the middle of the afternoon. However, this limit is not exceeded when considering aerosol feedback effects with a reduction of the modelled ozone peak from 251 to $230 \mu\text{g}/\text{m}^3$, in coherence with observations ($160 \mu\text{g}/\text{m}^3$). Hence, taking into account the impact of aerosol solar extinction on photolysis rates in the CHIMERE numerical simulations could play a non-negligible role in term of air quality prediction, keeping in mind that better constrained emissions and nocturnal processes could also greatly improve model scores.

In parallel, the presence of aerosols tends to reduce the oxidising capacity of the atmosphere (through reduction of OH radicals, see equation 5), which leads to decrease the formation of secondary aerosols. As illustrated in figure 89a, the maximum reduction in the near-surface concentrations of sulphates (oxidation product of SO_2) and SOA (oxidation product of COV) occurs on 9 August with diurnal-averaged 8 August with daytime average values of 10 % and 4 %, respectively. For this day, figure 9b shows that these changes are mainly due to a reduced formation of very fine particles, i.e with a diameter comprised between 40 nm (bin 1) and 300 nm (bin 4). The overall impact is then a slightly reduction of the total aerosol mass concentration (PM10) comprised between 1 and 2 % over the entire period (Figure 9a). These results are comparable to the findings of Real and Sartelet (2011) and Li et al. (2011b) who showed a 5–10 % reduction of the formation of secondary aerosols due to the aerosol solar extinction, in case of intense particulate pollution over, respectively, Europe and Mexico City. As for NO_2 and O_3 , including the optical effect of aerosols in the photolysis calculations slightly improves the formation of secondary inorganic species in the CHIMERE model with a RMSE systematically reduced (see Table 3). The overestimation of sulphates levels is decreased and the simulated concentrations of ammonium and nitrates get close to the observed one. This result suggests that taking into account the aerosol solar extinction in the photolysis calculation gives an added value in the capacity of the

[model to reproduce the photochemistry under polluted environments.](#)

4 Conclusions

In the present study, we have developed an on-line coupling between the chemistry-transport model CHIMERE (complemented by an aerosol optical module) and the radiative transfer code TUV to study the impact of aerosol solar extinction on the photochemistry over eastern Europe during the 2010 wildfires episode. Simulations from 5 to 12 August 2010, corresponding to the peak of fire activity, have been performed with and without aerosol impact on photolysis rates and concentrations of gaseous and particulate pollutants. Large areas were affected by important concentrations of particles, with modeled AOT (440 nm) above 2 along the transport of the aerosol plume. A previous evaluation of the modeled optical properties during this specific episode (P  r   et al., 2014) showed good model performance in simulating both the magnitude and spectral dependence of the aerosol optical thickness and single scattering albedo.

The impact of aerosols on photolysis rates is shown to be regionally significant with a mean reduction of $J[\text{NO}_2]$ and $J[\text{O}_3]$ comprised between 2 % and 50 %, the maximum reduction being modelled in the aerosol plume. These modifications of photolysis frequencies result in an regional increase in the daytime concentration of NO_2 of 3–15 % and a decrease in the O_3 production near the surface comprised between 1 % and 12 % during 8 August.

The photochemistry over the Moscow region has also been affected, especially during the arrival of the aerosol plume between 6 and 10 August 2010. Over this area, results indicate that $J[\text{O}_3]$ is more sensitive to the presence of particles (reduction of about ~~19~~10 % per unit of AOT) than $J[\text{NO}_2]$ (reduction of about ~~10~~6 % per unit of AOT), resulting in an increase of the ground NO_2 concentration of ~~4~~3 % (per unit of AOT) and a small reduction of ozone of ~~2~~1 % (per unit of AOT). The photochemistry is shown to be impacted along the boundary layer where aerosols are located with, for example, a

25 4–5 % O₃ reduction modelled during 9 August within the first two km of the atmosphere. In terms of model performance, comparisons of simulations with air quality measurements at Moscow indicate that an explicit representation of aerosols interaction with photolysis rates tend to improve the estimation of the near-surface concentration of ozone and nitrogen dioxide, with some potential consequence for air quality prediction. For instance, the O₃ peak reduction when including aerosol feedback induces a non-negligible difference in the predicted exceedance of alert threshold during 8 August compared to the simulation without aerosol feedback, in coherence with measurements. In addition, the impact of aerosols on photolysis rates is shown to have an influence on the formation of secondary aerosols, through the modification of the OH concentration. Over Moscow, the aerosol plume tends to decrease the daytime concentrations of sulphates and secondary organics up to 4–10 %, which result in a small reduction of the total particulate concentration (PM₁₀) of 1–2 % on average over the period. The results presented in this work, issued from a modelling exercise, are consistent with those obtained in recent studies by combining model experiments and different sets of observations. In terms of model performance, comparisons of simulations with air quality measurements at Moscow indicate that an explicit representation of aerosols interaction with photolysis rates tend to improve the estimation of the near-surface concentration of ozone and nitrogen dioxide as well as the formation of inorganic aerosol species such as ammonium, nitrates and sulphates.

15 Recently, it has been suggested that some organic aerosols can absorb solar radiation, especially at the shorter visible and UV wavelengths (Zhong and Jang, 2011; Saleh et al., 2014). The methodology developed in this study provides a powerful tool to investigate the role of enhanced UV absorption by secondary organics on photo-chemistry at regional and urban scale.

20

Acknowledgements. Authors are grateful to the CRI at Lille 1, François Thieuleux at LOA and Anthony Ung at INERIS for their technical support. Olga Kislova at Mosecomonitoring (Moscow) is acknowledged for providing the surface air quality measurements used in this study. Authors

25 would also like to thank Isabelle Chiapello at LOA for her fruitful scientific discussion. The CNES french spatial agency is acknowledged for its financial support.

References

- Amin, N.: Effect of ozone on the relative yield of rice crop in Japan evaluated based on monitored concentrations, *Water, Air and Soil Pollution*, 225, 1–9, 2014.
- Beelen, R., Raaschou-Nielsen, O., Stafoggia, M., and co authors: Effects of long-term exposure to air pollution on natural-cause mortality: an analysis of 22 European cohorts within the
- 5 multicentre ESCAPE project, *The Lancet*, 383, 785–795, 2014.
- Bessagnet, B., Hodzic, A., Vautard, R., Beekmann, M., Cheinet, S., Honoré, C., Liousse, C., and Rouil, L.: Aerosol modeling with CHIMERE — Preliminary evaluation at the continental scale, *Atmospheric Environment*, 38, 2803–2817, 2004.
- Bessagnet, B., Menut, L., Curci, G., Hodzic, A., Guillaume, B., Liousse, C., Moukhtar, S.,
- 10 Pun, B., Seigneur, C., and Schulz, M.: Regional modeling of carbonaceous aerosols over Europe—focus on secondary organic aerosols, *Journal of Atmospheric Chemistry*, 61, doi:10.1007/s10874–009–9129–2, 2008.
- Bian, H., Han, S., Tie, X., Sun, M., and Liu, A.: Evidence of impact of aerosols on surface ozone concentration in Tianjin, China, *Atmospheric Environment*, 41, 4672–4681, 2007.
- 15 Chen, F. and Dudhia, J.: Coupling an advanced land-surface / hydrology model with the Penn State / NCAR MM5 modeling system. Part 1: Model description and implementation, *Mon. Wea. Rev.*, 129, 569–585, 2001.
- Chubarova, N., Nezval, Y., Sviridenkov, I., Smirnov, A., and Slutsker, I.: Smoke aerosol and its radiative effects during extreme fire event over Central Russia in summer 2010, *Atmos. Meas. Tech.*, 5, 557–568, 2012.
- 20 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmospheric Chemistry and Physics*, 6, 3181–3210, 2006.
- Hauglustaine, D. A., Hourdin, F., Jourdain, L., Filiberti, M. A., Walters, S., Lamarque, J. F., and Holland, E. A.: Interactive chemistry in the Laboratoire de Météorologie Dynamique general
- 25 circulation model: description and background tropospheric chemistry evaluation, *Journal of Geophysical Research*, 109, d04314.doi:10.1029/2003JD003957, 2004.

- Hodzic, A., Madronich, S., Bohn, B., Massie, S., Menut, L., and Wiedinmyer, C.: Wildfire particulate matter in Europe during summer 2003: meso-scale modeling of smoke emissions, transport and radiative effects, *Atmospheric Chemistry and Physics*, 7, 4043–4064, 2007.
- Hong, S. Y.: Stable boundary layer mixing in a vertical diffusion scheme, *The Korea Meteor. Soc.*, fall conference, Seoul, Korea, Oct. 25-26, 2007.
- Hong, S. Y., Noh, Y., and Dudhia, J.: A new vertical diffusion package with an explicit treatment of entrainment processes, *Monthly Weather Review*, 134, 2318–2341, 2006.
- Honoré, C., Rouil, L., Vautard, R., Beekmann, M., Bessagnet, B., Dufour, A., Elichegaray, C., Flaud, J. M., Malherbe, L., Meleux, F., Menut, L., Martin, D., Peuch, A., Peuch, V. H., and Poisson, N.: Predictability of European air quality: Assessment of 3 years of operational forecasts and analyses by the PREV'AIR system, *Journal of Geophysical Research*, 113, D04301, doi:10.1029/2007JD008761, 2008.
- Horowitz, L. W., Walters, S., Mauzeralles, D. Z., Emmonds, L. K., Rash, P. J., Granier, C., Tie, X., Lamarque, J. F., and Schultz, M. G.: A global simulation of tropospheric ozone and related tracers : Description and evaluation of MOZART, version 2, *Journal of Geophysical Research*, 108, 108(D24), 4784, 2003.
- Hunova, I., Maly, M., Rezacova, J., and Branis, M.: Association between ambient ozone and health outcomes in Prague, *International archives of occupational and environmental health*, 86, 89–97, 2014.
- Jenkin, M. E. and Clemitshaw, K. C.: Ozone and other secondary photochemical pollutants: chemical processes governing their formation in the planetary boundary layer, *Atmospheric Environment*, 34, 2499–2527, 2000.
- Joseph, J. H., Wiscombe, W. J., and Weinman, J. A.: The delta-Eddington approximation for radiative flux transfer, *Journal of the Atmospheric Sciences*, 33, 2452–2459, 1976.
- Junker, C. and Lioussse, C.: A global emission inventory of carbonaceous aerosol from historic record of fossil fuel and biofuel consumption for the period 1860-1997, *Atmospheric Chemistry and Physics*, 8, 1195–1207, 2008.
- Kain, J. S.: The Kain-Fritsch convective parameterization: An update, *J. Appl. Meteor.*, 43, 170–181, 2004.
- Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J. J., Razinger, M., Schultz, M. G., Suttie, M., and van der Werf, G. R.: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power, *Biogeosciences*, 9, 527–554, 2012.

- Konovalov, I. B., Beekmann, M., Kuznetsova, I. N., Yurova, A., and Zvyagintsev, A. M.: Atmospheric impacts of the 2010 Russian wildfires: integrating modelling and measurements of an extreme air pollution episode in the Moscow region, *Atmospheric Chemistry and Physics*, 11, 10 031–10 056, 2011.
- Lau, W. K. M. and Kim, K.-M.: The 2010 Pakistan flood and Russian heat wave: teleconnection of hydrometeorological extremes, *J. Hydrometeor.*, 13, 392–403, 2012.
- Lesins, G., Chylek, P., and Lohmann, U.: A study of internal and external mixing scenarios and its effect on aerosol optical properties and direct radiative forcing, *Journal of Geophysical Research*, 107, 4094, 2002.
- Li, G., Bei, N., Tie, X., and Molina, L. T.: Aerosol effects on the photochemistry in Mexico City during MCMA-2006/MILAGRO campaign, *Atmospheric Chemistry and Physics*, 11, 5169–5182, 2011b.
- Li, J., Wang, Z., Wang, X., Yamaji, K., Takigawa, M., Kanaya, Y., Pochanart, P., Liu, Y., Irie, H., Hu, B., Tanimoto, H., and Akimoto, H.: Impacts of aerosols on summertime tropospheric photolysis frequencies and photochemistry over Central Eastern China, *Atmospheric Environment*, 45, 1817–1829, 2011a.
- Lou, S., Liao, H., and Zhu, B.: Impacts of aerosols on surface-layer ozone concentrations in China through heterogeneous reactions and changes in photolysis rates, *Atmospheric Environment*, 85, 123–138, 2014.
- Madronich, S. and Flocke, S.: The role of solar radiation in atmospheric chemistry, *Handbook of Environmental Chemistry*, pp. 1–26, Springer-Verlag, Heidelberg, Germany, 1998.
- McMeeking, G. R., Morgan, W. T., Flynn, M., Highwood, E. J., Turnbull, K., Haywood, J., and Coe, H.: Black carbon aerosol mixing state, organic aerosols and aerosol optical properties over the United Kingdom, *Atmospheric Chemistry and Physics*, 11, 9037–9052, 2011.
- Mena-Carrasco, M., R.Carmichael, G., Campbell, J. E., Zimmerman, D., Tang, Y., Adhikary, B., D'allura, A., Molina, L. T., Zavala, M., Flocke, A. G. F., Campos, T., Weinheimer, A. J., Shetter, R., Apel, E., Montzka, D. D., Knapp, D. J., and Zheng, W.: Assessing the regional impacts of Mexico City emissions on air quality and chemistry, *Atmospheric Chemistry and Physics*, 9, 3731–3743, 2009.
- Menut, L., Bessagnet, B., Khvorostyanov, D., Beekmann, M., Blond, N., Colette, A., Coll, I., Curci, G., Foret, G., Hodzic, A., Mailler, S., Meleux, F., Monge, J.-L., Pison, I., Siour, G., Turquety, S., Valari, M., Vautard, R., and Vivanco, M. G.: CHIMERE 2013: a model for regional atmospheric composition modelling, *Geosci. Model Dev.*, 6, 981–1028, 2013.

- 30 Miller, M. R., Shaw, C. A., and Langrish, J. P.: From particles to patients: oxidative stress and the cardiovascular effects of air pollution, *Future Cardiology*, 8, 577–602, 2012.
- Nie, W., Ding, A. J., Xie, Y. N., Xu, Z., Mao, H., Kerminen, V. M., Zheng, L. F., Qi, X. M., Huang, X., Yang, X. Q., Sun, J. N., Herrmann, E., Petäjä, T., Kulmala, M., and Fu, C. B.: Influence of biomass burning plumes on HONO chemistry in eastern China, *Atmospheric Chemistry and Physics*, 15, 1147–1159, 2015.
- 5 Palancar, G. G., Lefer, B. L., Hall, S. R., Shaw, W. J., Corr, C. A., Herndon, S. C., Slusser, J. R., and Madronich, S.: Effect of aerosols and NO₂ concentration on ultraviolet actinic flux near Mexico City during MILAGRO: measurements and model calculations, *Atmospheric Chemistry and Physics*, 13, 1011–1022, 2013.
- Péré, J. C., Mallet, M., Bessagnet, B., and Pont, V.: Evidence of the aerosol core-shell mixing state over Europe during the heat wave of summer 2003 by using CHIMERE simulations and AERONET inversions, *Geophysical Research Letters*, 36, doi:10.1029/2009GL037334, 2009.
- 10 Péré, J. C., Mallet, M., Pont, V., and Bessagnet, B.: Evaluation of an aerosol optical scheme in the chemistry-transport model CHIMERE, *Atmospheric Environment*, 44, 3688–3699, 2010.
- Péré, J. C., Mallet, M., Pont, V., and Bessagnet, B.: Impact of the aerosol direct radiative forcing (ADRF) on the radiative budget, surface heat fluxes and atmospheric dynamics during the heatwave of summer 2003 over Western Europe. A modelling study., *Journal of Geophysical Research*, 116, d23119, doi:10.1029/2011JD016240, 2011.
- 15 Péré, J. C., Bessagnet, B., Mallet, M., Waquet, F., Chiapello, I., Minvielle, F., Pont, V., and Menut, L.: Direct radiative effect of the Russian wildfires and its impact on air temperature and atmospheric dynamics during August 2010, *Atmospheric Chemistry and Physics*, 14, 1999–2013, 2014.
- 20 Popovicheva, O., Kistler, M., Kireeva, E., Persiantseva, N., Timofeev, M., Kopeikin, V., and Kasper-Giebl, A.: Physicochemical characterization of smoke aerosol during large-scale wildfires: Extreme event of August 2010 in Moscow, *Atmospheric Environment*, 96, 405–414, 2014.
- 25 Real, E. and Sartelet, K.: Modeling of photolysis rates over Europe: impact on chemical gaseous species and aerosols, *Atmospheric Chemistry and Physics*, 11, 1711–1727, 2011.
- Saleh, R., Robinson, E. S., Tkacik, D. S., Ahern, A. T., Liu, S., Aiken, A. C., Sullivan, R. C., Presto, A. A., Dubey, M. K., Yokelson, R. J., Donahue, N. M., and Robinson, A. L.: Brownness

- 30 of organics in aerosols from biomass burning linked to their black carbon content, *Nature Geoscience*, 7, 647–650, 2014.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics. From air pollution to climate change*, Wiley-Interscience publication, 1998.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics. From air pollution to climate change (2nd Edition)*, Wiley-Interscience publication, 2006.
- 5 Slade, J. H. and Knopf, D. A.: Heterogeneous OH oxidation of biomass burning organic aerosol surrogate compounds: assessment of volatilisation products and the role of OH concentration on the reactive uptake kinetics, *Phys. Chem. Chem. Phys.*, 15, 5898–5915, 2013.
- Turquety, S.: *The Atmospheric Impact of Wildfires. Fire Phenomena and the Earth System: An Interdisciplinary Guide to Fire Science*, Wiley Blackwell, 2013.
- Vautard, R., Beekmann, M., Roux, J., and Gombert, D.: Validation of a hybrid forecasting system for the ozone concentrations over the Paris area, *Atmospheric Environment*, 14, 2449–2461, 2001.
- 10 Vautard, R., Bessagnet, B., Chin, M., and Menut, L.: On the contribution of natural aeolian sources to particulate matter concentrations in Europe: Testing hypotheses with a modeling approach, *Atmospheric Environment*, 39, 3291–3303, 2005.
- Vester, B. P., Ebert, M., Barnert, E. B., Schneider, J., Kandler, K., Schütz, L., and Weinbruch, L.: Composition and mixing state of the urban background aerosol in the Rhein-Main area (Germany), *Atmospheric Environment*, 41, 6102–6115, 2007.
- 645 Wai, K. M. and Tanner, P. A.: Variations of aerosol properties due to regional source contributions and impacts on ozone levels: a study in a south China city, *Environ. Chem.*, 7, 359–369, 2010.
- Wu, X., Seigneur, C., and Bergström, R.: Evaluation of a sectional representation of size distributions for calculating aerosol optical properties, *Journal of Geophysical Research*, 101, 19 277–19 283, 1996.
- 650 Wu, Z. P. and Wang, Y. P.: Electromagnetic scattering for multilayered spheres: Recursive algorithms, *Radio Science*, 26, 1393–1401, 1991.
- Zhong, M. and Jang, M.: Light absorption coefficient measurement of SOA using a UV–Visible spectrometer connected with an integrating sphere, *Atmospheric Environment*, 45, 4263–4271, 2011.
- 655

Zvyagintsev, A. M., Ivanova, N. S., Blyum, O. B., Kotel'nikov, S. N., Kruchenitskii, G. M., Kuznetsova, I. N., and Lapchenko, V. A.: Ozone content over the Russian Federation in the third quarter of 2010, *Russian Meteorology and Hydrology*, 35, 785–789, 2010.




Fig. 1a. Geographic distribution of the AOT (at midday) over Eastern Europe from 5 to 12 August 2010 modelled by CHIMERE (at 400 nm). Horizontal wind at 850 hPa simulated by WRF is also indicated.




Fig. 1b. Continuation of Figure 1




Fig. 2a. Geographic distribution of the modeled ~~diurnal-averaged~~ daytime average percentage changes in the near-surface $J[\text{NO}_2]$ due to the presence of aerosols.




Fig. 2b. Continuation of Figure 2.




Fig. 3a. Geographic distribution of the modelled ~~diurnal-averaged~~ daytime average percentage changes in the near-surface $J[O_3]$ due to the presence of aerosols.




Fig. 3b. Continuation of Figure 3.




Fig. 4. Geographic distribution of the ~~diurnal-averaged~~ daytime average percentage changes in the near-surface concentration of NO_2 (left) and O_3 (right) for the 8 August 2010, due to the presence of aerosols.




Fig. 5. Daytime average percentage changes (at midday) in the photolysis frequencies and concentrations of nitrogen dioxide and ozone over Moscow as a function of modelled AOT (440 nm).




Fig. 6. Mean VOC to NOx ratio simulated over Moscow during the studied period




Fig. 7. Vertical profile of the ~~diurnal-averaged~~ daytime average percentage changes in the ozone concentration at the north of Moscow (59.9 °N, 37.6 °E) for the 9 August. The aerosol extinction coefficient (in km⁻¹) modelled by CHIMERE and measured by CALIOP at midnight is also indicated.






Fig. 8. Temporal evolution (between 5 and 12 August 2010) of the near-surface ozone concentration (in $\mu\text{g}/\text{m}^3$) modelled with and without aerosol feedback along with corresponding observations at the Moscow monitoring station. The European alert threshold of $240 \mu\text{g}/\text{m}^3$ is also indicated by a straight line.



Diurnal-averaged percentage reduction of the near-surface concentration of sulphates, secondary organic aerosols and PM10 over Moscow due to the aerosol feedback.

Fig. 9. a) Daytime average percentage reduction of the near-surface concentration of sulphates, secondary organic aerosols and PM10 over Moscow due to the aerosol feedback. b) Repartition of this sulphate and SOA mass reduction between the 8 aerosol size bins for the 8 August

Table 1. statistical comparisons between the near-surface concentrations of NO₂ and O₃ simulated with and without aerosols and measured at Moscow by an air quality station. Mod. and Obs. are the period-averaged modelled and observed concentration. Corr. and RMSE are the temporal correlation and the root mean square error.

| | NO ₂ | | | | O ₃ | | | |
|---------|--------------------------------------|--------------------------------------|-------|--------------------------------------|--------------------------------------|--------------------------------------|-------|--------------------------------------|
| | Mod. ($\mu\text{g}/\text{m}^3$) | Obs. ($\mu\text{g}/\text{m}^3$) | Corr. | RMSE ($\mu\text{g}/\text{m}^3$) | Mod. ($\mu\text{g}/\text{m}^3$) | Obs. ($\mu\text{g}/\text{m}^3$) | Corr. | RMSE ($\mu\text{g}/\text{m}^3$) |
| with | 14 | 70 | 0.22 | 66 | 129 | 66 | 0.46 | 82 |
| without | 13 | 70 | 0.21 | 67 | 134 | 66 | 0.42 | 88 |

Table 2. Same as in Table 1 but for daily maximum values

| | NO ₂ | | | | O ₃ | | | |
|---------|--------------------------------------|--------------------------------------|-------|--------------------------------------|--------------------------------------|--------------------------------------|-------|--------------------------------------|
| | Mod. ($\mu\text{g}/\text{m}^3$) | Obs. ($\mu\text{g}/\text{m}^3$) | Corr. | RMSE ($\mu\text{g}/\text{m}^3$) | Mod. ($\mu\text{g}/\text{m}^3$) | Obs. ($\mu\text{g}/\text{m}^3$) | Corr. | RMSE ($\mu\text{g}/\text{m}^3$) |
| with | 32 | 90 | 0.60 | 66 | 180 | 161 | 0.19 | 48 |
| without | 31 | 90 | 0.60 | 67 | 185 | 161 | 0.16 | 54 |

Table 3. Same as in Table 1 but for the near-surface concentrations of ammonium, nitrates and sulphates

| | NH_4^+ | | | | NO_3^- | | | | SO_4^{2-} | | | |
|---------|--------------------------------------|--------------------------------------|-------|--------------------------------------|--------------------------------------|--------------------------------------|-------|--------------------------------------|--------------------------------------|--------------------------------------|-------|--------------------------------------|
| | Mod. ($\mu\text{g}/\text{m}^3$) | Obs. ($\mu\text{g}/\text{m}^3$) | Corr. | RMSE ($\mu\text{g}/\text{m}^3$) | Mod. ($\mu\text{g}/\text{m}^3$) | Obs. ($\mu\text{g}/\text{m}^3$) | Corr. | RMSE ($\mu\text{g}/\text{m}^3$) | Mod. ($\mu\text{g}/\text{m}^3$) | Obs. ($\mu\text{g}/\text{m}^3$) | Corr. | RMSE ($\mu\text{g}/\text{m}^3$) |
| with | 1.09 | 0.86 | 0.48 | 1.03 | 0.14 | 0.12 | 0.20 | 0.15 | 2.24 | 0.48 | 0.23 | 1.80 |
| without | 1.16 | 0.86 | 0.42 | 1.04 | 0.19 | 0.12 | 0.26 | 0.23 | 2.33 | 0.48 | 0.45 | 1.88 |