

ACPD Detailed response to referees

We thank the reviewers for their comments. These comments helped us to relate our work to previous works in the field and to consider an additional learning algorithm that was designed for non-stationarity sequences. Please find below our detailed response to the referees' comments along with a summary of the changes made in the revised version.

Anonymous Referee #1

We thank reviewer #1 for a very detailed and informative review.

Strong similarities to past work

It appears the authors were unaware of prior work, but unfortunately this submission cannot be published without major revision, because both the proposed problem (using learning algorithms to compute weightings to improve the predictions of the multi-model ensemble of AOGCMs) and the type of machine learning approach ("sequential learning algorithms") were already presented in previous work.

- C. Monteleoni, G. Schmidt, and S. Saroha. Tracking Climate Models. In NASA Conference on Intelligent Data Understanding (CIDU), pages 1–15, 2010.
- C. Monteleoni, G. Schmidt, S. Saroha, and E. Asplund. Tracking Climate Models. In Journal of Statistical Analysis and Data Mining: Special Issue: Best of CIDU 2010. Volume 4, Issue 4, pages 72–392, August 2011.
- S. McQuade and C. Monteleoni. MRF-Based Spatial Expert Tracking of the Multi-Model Ensemble. In New Approaches for Pattern Recognition and Change Detection, session at American Geophysical Union (AGU) Fall Meeting, 2013.

The first 2 are the most similar to the present submission; the 3rd is also highly related and was presented at AGU.

We thank the referee for pointing out these important works.

In the revised version, we cite and discuss these works (the published works that we could read and refer to) and the differences between them and our work.

Moreover, while the stated task is identical to the first two works listed above, i.e. to use "sequential learning algorithms" (also known as "online learning with expert advice"), to combine the temperature projections of a CMIP ensemble (CMIP3 was used in the 2010 paper), the machine learning methods from the 2010 paper are more recent than the ones in the present submission and essentially subsume the methods used here.

To see this, note that while the authors cite a textbook to refer to the algorithms (Cesa-Bianchi and Lugosi 2006, which we will refer to as CB06), the actual algorithms used are from the 1990's and earlier.

The algorithm denoted as EWA in the submission is originally due to:

- N. Littlestone and M. K. Warmuth. The weighted majority algorithm. In IEEE Symposium on Foundations of Computer Science, pages 256–261, 1989.
- V. Vovk. Aggregating strategies. In Proc. 3rd Annu. Workshop on Comput. Learning Theory, pages 371–383. Morgan Kaufmann, 1990.

The algorithm denoted as EGA in the submission is originally due to:

- Jyrki Kivinen and Manfred K. Warmuth: Exponentiated gradient versus gradient descent for linear predictors. Information and Computation 132(1):1-63, 1997.

In contrast, the algorithm in the 2010 paper (which we will hereby refer to as M10) is from the early 2000's:

- C. Monteleoni and T. Jaakkola. Online Learning of Non-stationary Sequences. In Advances in Neural Information Processing Systems 16. pages 1093–1100, 2003.

In the intervening time, the field of learning algorithms had advanced significantly (and subsequently it has advanced even further). In particular, relevant to the study of climate change and the problem in question, online learning algorithms were designed expressly for non-stationary data. The algorithms applied in the present submission were not designed to handle non-stationary data. Their performance guarantees (regret bounds) are with respect to the best fixed expert. However many climate scientists have shown that in any CMIP ensemble, some climate models perform better at some times and others at other times. An influential paper on online learning algorithms (extending algorithms with exponentiated weight updates for a variety of prediction loss functions) provided modified weight updates that allow for “switching” among “best” experts:

- M. Herbster and M. K. Warmuth, Tracking the best expert, Mach Learn 32 (1998), 151 – 178.

The algorithm used in M10 was a further extension from that work which actually learns the switching rate online (sequentially), simultaneous to updating the weights over the experts. It can still track the best fixed-expert as a special case, and in that sense it is significantly more flexible than the algorithms used in the present submission.

In our field, it is common to cite a book (or a review article) that summarizes previous results.

Note that the application of the EWA and EGA in our work involves optimization of the learning rate.

In order to compare the results of these algorithms to the more recent LAA, we considered the LAA in the revised version, and we present the results of all the learning schemes for an easy comparison of their performances.

In the revised version, we added citations of the original works related to the EWA, EGA, fixed-share, and LAA . In addition, we added the results of the LAA to Figures 3 and 5 and changed the Results and Discussion sections to reflect these changes and additions.

More recently, there have been several more machine learning approaches to the multi-model ensemble prediction problem proposed in the present submission. Note that the present submission treats geospatial locations separately. This was also the case in M10, but subsequently, they did an extension to explicitly model geospatial neighborhood influence in the online learning weight updates. This is another highly related work to the present submission.

- S. McQuade and C. Monteleoni. Global Climate Model Tracking using Geospatial Neighborhoods. In Proceedings of the Twenty-Sixth AAAI Conference on Artificial Intelligence, Special Track on Computational Sustainability and AI, pp. 335–341, 2012.

The above was on a regular grid over latitude and longitude. However there has also been some recent work on learning the geospatial structure (need not be a grid) from the data such that regional features such as mountain ranges may cause some geospatially neighboring regions to have lesser spatial influence on one another.

- A. R. Goncalves, P. Das, S. Chatterjee, V. Sivakumar, and A. Banerjee. Global climate models combination using Multitask Sparse Structure Learning. Fourth Workshop on Understanding Climate Change Through Data, At National Center for Atmospheric Research (NCAR), Boulder, Colorado, USA, 2014.

The above is not a “sequential learning” technique. Additionally, there have been other kinds of machine learning algorithms applied to the problem, e.g.

- M. Ghafarianzadeh and C. Monteleoni. Climate Prediction via Matrix Completion. In Proceedings of the Twenty-Seventh AAAI Conference on Artificial Intelligence, Late-Breaking Papers Track, 2013.

Geospatial correlations do exist in the climate system and accounting for these correlations have the potential to improve future climate predictions as discussed by the referee. However, this is beyond the scope of the current work. Moreover, the results show a relatively smooth spatial distribution of the RMSE and uncertainty, which suggests that the geospatial correlations are well captured by the models in the ensemble.

Note, when the authors discuss using methods that predict “according to the best model,” they should see the related literature on learning algorithms. (In CB06, see Follow the Leader and Follow the Perturbed Leader).

The meaning of the “best” model is explicitly explained in the text. It is important to note that we only used this type of prediction to show that the predictions of the SLAs outperform those of the models that obtain the highest weights.

In summary, the main problem with the present submission is its very strong similarity to previous work, along with a significant gap in the discussion of related work on using learning algorithms in climate predictions, in particular for the problem in question.

The works mentioned by the referee analyzed long-term climate predictions, most of them based on the CMIP3. In our work, we focused only on decadal climate predictions. These are different simulations initialized with real observed conditions unlike the long-term climate predictions. These experiments were first introduced in the CMIP5. The introduction of the paper elaborates on the differences between the two types of predictions.

Our results show that for decadal climate predictions, adding the climatology as an additional expert in the ensemble significantly improves the predictions. For obvious reasons, adding the climatology to long-term climate projections is not expected to have the same effect.

The revised versions of Figures 3 and 5 (and the globally averaged results provided in the text) show that the LAA performs similarly to the EWA. However, the EGA outperforms both the EWA and the LAA. The similar performance of the EWA and the LAA suggests that for decadal climate predictions, the non-stationarity nature of the dynamics is not significant. None of the previous works presented such a comparison between different SLAs. All these details are discussed in the revised version.

Previous works focused on improving the predictions using ensembles of models and learning algorithms. Here, we show that SLAs not only improve the predictions but they also reduce the associated uncertainties. The uncertainties in future climate predictions are crucial for evaluating the predictions and, clearly, for any practical use of the results.

In the revised manuscript, we attempt to better emphasize these differences from previous works.

Technical corrections

There’s a mistake in the equations. While it is more standard to state the weights in terms of the losses, L , than the regret, R , the authors chose to use R , and in the process a mistake was introduced into the

equation for weights. When using L , there's a negative sign before the learning rate η in the exponent. However since R is defined as the loss of the expert subtracted from the loss of the learning algorithm, when using R in the exponent, there should not be negative sign. Intuitively, experts' weights should be decreased not increased, proportionally to their prediction losses. So the authors should either switch to L or else delete the negative in the exponent of Equation (3). (See pages 14, 72, and 304 of CB06).

We thank the referee for pointing out this typo.

In the revised version, we use the loss rather than the regret in order to maintain consistency with previous works.

Differences from past work, and positive aspects of the submission

On the positive side, the paper is very thoughtful and well-written. Moreover, there are also some novel experiments and ways of looking at the results, along with interesting insights from climate science. These should be emphasized in a revision.

- The input ensemble is from CMIP5 and has other differences from the ensembles used in several of the past works.
- Learning is done at a much higher spatial granularity than in M10 and some of the past works.
- The visualizations are different than M10 (although not all are different than all the past work on the problem, cf. Goncalves).
- There are some interesting findings, e.g. they point out that their results show that there is more model disagreement at polar regions.
- The authors use a variety of different metrics to evaluate their results instead of just prediction loss, e.g. RMSE (which however is closely related to the chosen prediction loss, also used in M10), as well as global, area-weighted uncertainty, and some interesting significance tests.
- The algorithm used in M10 has a per-time-iteration complexity that's on the order of a factor of T times that of the algorithm used in the present submission, where T is the desired number of temporal epochs (e.g. number of months for the whole experiment). This is for the optimized version of the algorithm in M10; one can also use lower complexity for slightly degraded results. This increased complexity is used in handling non-stationary data.

This is a difference, although not necessarily an advantage, from M10: The evaluations are done using a training and a test period. However this might not be the best technique, given that the observations and climate models exhibit non-stationarity. In such a setting, online learning algorithms are typically evaluated using progressive validation error and/or cumulative prediction loss (see M10).

We thank the referee for the positive comments.

Recommendations and suggested resources

It is very encouraging to see climate scientists using machine learning in their research. However this reviewer recommends resubmission, taking these major concerns into account. Given that there is already significant prior work in this area, the authors are strongly advised to re-think the framing of their submission, discussing past work, emphasizing the differences, and focusing much more on what is novel in their submission. Perhaps given the strong overlaps with past work, the authors will decide to run additional experiments with other learning algorithms with different properties, and compare them.

We followed the referee's advice and revised the paper to include a better review of previous works in the field, and in particular, we added the most recent learning algorithm and compared its performance to the older algorithms used in the original submission. We also now emphasize the difference between

decadal climate predictions and long-term predictions, as well as the finding that on decadal time scales, the non-stationarity of the climate predictions do not seem to be crucial.

There is a field of interdisciplinary research using learning algorithms in climate science called “climate informatics” (see <http://climateinformatics.org> for a variety of links and a recent video tutorial surveying work in the field). There is an annual international climate informatics workshop (since 2011; the next workshop will be held at NCAR in Colorado, in September 2015). These authors are doing work in this area and might benefit from this growing research community.

We thank the referee for informing us of this important workshop.

Anonymous Referee #2

We thank the reviewer for the helpful report.

This paper presents a machine learning-based approach for improving model-generated projections of past and future climate. The work is well motivated citing appropriate literature, and the technical approach is presented in sufficient detail.

We thank the referee for the positive comments.

Unfortunately, the authors appear to have missed some important literature in this area, most notably work by C. Monteleoni and colleagues over the past several years, see e.g.,:
C. Monteleoni, G. Schmidt, and S. Saroha. Tracking Climate Models. In NASA CIDU, 2010. S. McQuade and C. Monteleoni. Global Climate Model Tracking using Geospatial Neighborhoods. In The 2nd Int’l Workshop on Climate Informatics, 2012. C. Monteleoni, G. Schmidt, and S. McQuade, Climate Informatics: Accelerating Discovery in Climate Science with Machine Learning. IEEE Computing in Science and Engineering (CISE) Magazine, Special Issue on Machine Learning. 15(5), 32–40, 2013.

Indeed, in the previous version of the manuscript, we missed these important works. Please see our detailed response to reviewer #1 for a thorough discussion of the differences between our work and previous ones.

In the revised manuscript, the learn- α algorithm was considered and compared with the other learning algorithms.

After much deliberation, this omission demands rejection of the manuscript. In order to be reconsidered, the authors should thoroughly familiarize themselves with this and any other prior work (in both climate / Earth science and machine learning as it is an interdisciplinary contribution), re-frame the presentation of their approach in the context of the literature, and qualitatively and/or quantitatively compare their approach to existing methods where appropriate.

In the revised manuscript, we discuss previous works and the differences between them and our work. In addition, we consider the algorithm previously used and compare its performance to the other sequential learning algorithms.

List of changes:

1. We added a brief discussion of the differences between decadal climate predictions and long-term climate projection to the Introduction.
2. We added paragraphs to the Introduction discussing previous works in which SLAs were applied to an ensemble of climate models.
3. In the description of the SLAs, we now use the loss rather than the regret, and we fixed the typo pointed out by reviewer #1.
4. In the SLA section, we introduced the learn- α algorithm (LAA).
5. In sections 3-5, we added the results of the LAA for the CMIP5 ensemble that we considered. The results are compared with the results of the other SLAs.
6. Figures 3 and 5 were modified to include the results of the LAA. Their captions were changed accordingly.
7. The Discussion and Summary section was modified to include a discussion of the differences between the LAA and the other SLAs considered. In addition, we now discuss the implications of the results found for decadal climate predictions.
8. We added references to:
Herbster, M. and Warmuth, M. K.: Tracking the best expert, *Mach Learn*, 32, 151–178, 1998.
Kivinen, J. and Warmuth, M.: Exponentiated gradient versus gradient descent for linear predictors. *Information and Computation*, 132(1):1–63, 1997.
Littlestone, N. and Warmuth, M.: The weighted majority algorithm. *Information and Computation*, 108:212–261, 1994.
Monteleoni, C. and Jaakkola, T.: Online Learning of Non-stationary Sequences, in *Advances in Neural Information Processing Systems*, 16, 1093–1100, 2003.
Monteleoni, C., Saroha, S., and Schmidt, G.: Tracking Climate Models, in *NASA Conference on Intelligent Data Understanding (CIDU)*, pages 1–15, 2010.
Monteleoni, C., Saroha, S., Schmidt, G., and Asplund, E.: Tracking Climate Models, in *Journal of Statistical Analysis and Data Mining: Special Issue: Best of CIDU 2010, Volume 4, Issue 4*, pages 72–392, 2011.

Manuscript prepared for Atmos. Chem. Phys. Discuss.
with version 2014/09/16 7.15 Copernicus papers of the \LaTeX class copernicus.cls.
Date: 26 May 2015

Improvement of climate predictions and reduction of their uncertainties using learning algorithms

E. Strobach and G. Bel

Department of Solar Energy and Environmental Physics, Blaustein Institutes for Desert Research,
Ben-Gurion University of the Negev, Sede Boqer Campus, 84990 Israel

Correspondence to: G. Bel (bel@bgu.ac.il)

Abstract

Simulated climate dynamics, initialized with observed conditions, is expected to be synchronized, for several years, with the actual dynamics. However, the predictions of climate models are not sufficiently accurate. Moreover, there is a large variance between simulations initialized at different times and between different models. One way to improve climate predictions and to reduce the associated uncertainties is to use an ensemble of climate model predictions, weighted according to their past ~~performance~~performances. Here, we show that skillful predictions, for a decadal time scale, of the 2 m-temperature can be achieved by applying a sequential learning algorithm to an ensemble of decadal climate model simulations. The predictions generated by the learning algorithm are shown to be better than those of each of the models in the ensemble, the better performing simple average and a reference climatology. In addition, the uncertainties associated with the predictions are shown to be reduced relative to those derived from an equally weighted ensemble of ~~bias corrected~~bias-corrected predictions. The results show that learning algorithms can help to better assess future climate dynamics.

1 Introduction

A new group of global climate simulations, referred to as the decadal experiments, was introduced in the Coupled Model Intercomparison Project (CMIP5) multi-model ensemble (Taylor et al., 2012; Meehl et al., 2009). The ~~idea behind these~~decadal climate predictions differ from the long-term climate projections in their duration, aims and meaningful output. The idea behind the decadal experiments was to investigate the predictability of the climate by atmosphere ocean general circulation models (AOGCMs) in time scales of up to 30 years whereas long-term climate projections use the same type of models to predict the forced response of the climate system to different future atmospheric compositions over the next century (Meehl et al., 2009; Taylor et al., 2012).

The AOGCMs [in the decadal experiments](#) were initialized with interpolated observations of the ocean, sea ice and atmospheric conditions, together with the atmospheric composition (Taylor and Meehl, 2011) [\(note that long-term projections are initialized with a quasi-equilibrium pre-industrial state \(Taylor et al., 2012\)\)](#). Therefore, they were expected to reproduce the monthly and annual averages of the climate variables and the response of the climate system to changes in the atmospheric composition (Warner, 2011; Collins, 2007; Kim et al., 2012). Indeed, it was shown (Kim et al., 2012) that in some regions, the CMIP5 simulations have some prediction skill. It was also confirmed (Kim et al., 2012) that the multi-model average provides better predictions than each of the models, similar to what was found for other climate simulations (Doblas-Reyes et al., 2000; Palmer et al., 2004; Hagedorn et al., 2005; Feng et al., 2011). However, the simple multi-model average does not take into account the quality differences between the models; therefore, it is expected that a weighted average, with weights based on the past performances of the models, will provide better predictions than the simple average. As expected, it was shown that the weighted average of climate models can improve predictions when using ensembles of AGCMs (Rajagopalan et al., 2002; Robertson et al., 2004; Yun et al., 2003), AOGCMs (YUN et al., 2005; Pavan and Doblas-Reyes, 2000; Chakraborty and Krishnamurti, 2009) and regional climate models (Feng et al., 2011; Samuels et al., 2013).

The uncertainties in climate predictions can be attributed to three main sources—: [the](#) internal variability of the model, inter-model variability and future forcing scenario uncertainties. The internal variability of the model stems from the sensitivity of the model to the initial conditions, sensitivity to the values of the parameters and the discretization method used. The inter-model variability is the result of different parameterization schemes and modeling approaches adopted in different models. The uncertainties due to different forcing scenarios are mostly related to different scenarios assumed regarding future greenhouse gas emissions. On a decadal time scale, forcing scenario uncertainties and uncertainties due to the internal variability of each model are considerably smaller than the inter-model uncertainties (Meehl et al., 2009; Hawkins and Sutton, 2009) (we also verified that the internal variability of each of the models we used is much smaller than the inter-model vari-

ability). Therefore, estimation of the uncertainties from an ensemble of climate models is expected to give a meaningful estimation of the total climate prediction uncertainties.

Different methods were used to improve climate predictions using an ensemble of models. A common approach is the simple regression (Krishnamurti et al., 2000; Krishnamurti, 1999). The regression does not assign a weight to each member of the ensemble but rather attempts to find the set of coefficients yielding the minimal square error for a linear combination of the ensemble model predictions. Bayesian methods have also been used for weighting ensembles of climate model projections (Rajagopalan et al., 2002; Robertson et al., 2004; Tebaldi and Knutti, 2007; Smith et al., 2009; Buser et al., 2009, 2010). The weighting scheme of these methods relies on a certain distribution of the errors and other prior assumptions regarding the models; these assumptions are not necessarily valid for climate dynamics and predictions. Many variations of the Bayesian methods were applied to weather forecasting in order to establish the ensemble of models (Kalnay et al., 2006); these methods are less useful for climate predictions in which the variability between different models is larger than the internal variability of each model (Meehl et al., 2009; Hawkins and Sutton, 2009).

Recently, sequential learning algorithms (SLAs) (Cesa-Bianchi and Lugosi, 2006) were applied to ensembles of climate models in order to improve the predictions (Mallet et al., 2009; Mallet, 2010; Monteleoni et al., 2010, 2011). Mallet et al. (2009); Mallet (2010) combined data assimilation and SLAs in order to improve seasonal to annual ozone concentration forecasts. Monteleoni et al. (2010, 2011) applied an improved version (Monteleoni and Jaakkola, 2003) of a method for learning non-stationary sequences (Herbster and Warmuth, 1998) to long-term climate predictions.

Here, we use a sequential learning algorithm (SLA) method several SLAs to weight climate models in the CMIP5 decadal experiments (Taylor and Meehl, 2011) and thereby to improve both global and regional predictions. In addition, we show that the uncertainties associated with these improved predictions are smaller than those of the unweighted ensemble. The first algorithm is the Exponentiated Weighted Average (EWA) (Littlestone, 1994) and the second is the Exponentiated Gradient Average (EGA) (Kivinen, 1997). The two

~~original algorithms were modified and adjusted to improve decadal climate predictions. A more recent algorithm, the learn- α algorithm (LAA), which is adopted from the field of game theory (Gesa-Bianchi and Lugosi, 2006; Mallet et al., 2009; Mallet, 2010), to weight ensemble members of global climate models. more suitable for the study of nonstationary sequences, was also used (Monteleoni and Jaakkola, 2003). The decadal climate predictions allow us to have a learning period and a validation period for testing the SLAs' performances. In addition, the use of methods for nonstationary sequences helps to assess the stationarity of the climate predictions in decadal time scales.~~

It is important to note that the SLA method assigns real weights (taking values between zero and one) to the ensemble models rather than to future climate paths (it is straightforward to use the weights of the models to get the probabilities of future climate paths ~~which are common in~~, which are the common products of the Bayesian approaches); this characteristic makes the SLA method appropriate for model evaluation. The SLA method has several advantages compared with other weighting schemes: (i) it makes no assumptions regarding the distribution of the climate variables and the model parameters. Therefore, it can be used for all climate variables and all types of predictions; (ii) there is an upper bound for the deviation of the weighted ensemble average from the best model. For a sufficiently lengthy learning period (the duration of this period depends on the variable, the learning rate (which is described later) and the number of models in the ensemble), the SLA prediction is at least as good as the prediction of the best model in the ensemble; (iii) the weights can be dynamically updated, when new measurements are introduced, with no significant computational cost. ~~The original method (Gesa-Bianchi and Lugosi, 2006) was modified and adjusted to improve decadal climate predictions.~~

2 The sequential learning ~~algorithm~~algorithms

~~The~~A sequential learning algorithm (SLA) assigns weights to the climate models (*the experts*) in the ensemble based on their past performance. In this work, the output of the models was divided into two periods—: a learning period during which the weights were

updated and a prediction period during which the weights remained fixed and equal to the weights assigned by the SLA in the last step of the learning process. In order to capture the spatial variability in model performance, the weights were spatially distributed and the weight of each model in each grid cell was determined by the local past performance of the model. For the sake of clarity, the algorithm is described below without spatial indexes although the calculations were done for each grid cell separately. The prediction of the SLA *forecasters* is the weighted average of the ensemble (Cesa-Bianchi and Lugosi, 2006). The weights are assigned to minimize the cumulative regret with respect to each one of the climate models. The cumulative regret of *expert E* is defined as:

$$R_{E,n} \equiv \sum_{t=1}^n (l(p_t, y_t) - l(f_{E,t}, y_t)) \equiv L_n - L_{E,n}. \quad (1)$$

t is a discrete time, l denotes some loss function that is a measure of the difference between the predicted (p_t by the *forecaster* and $f_{E,t}$ by *expert E*) and the true (y_t) values. In this work, we defined the loss function to be the square of the difference between the *forecaster* prediction and the “real” value, namely, $l(p_t, y_t) \equiv (p_t - y_t)^2$. $L_n \equiv \sum_{t=1}^n l(p_t, y_t)$, $L_{E,n} \equiv \sum_{t=1}^n l(f_{E,t}, y_t)$ are the cumulative loss functions of the *forecaster* and *expert E*, respectively. The outcome of the *forecaster*, after $n - 1$ steps of learning, is weights assigned to the climate models in the ensemble to be used for forecasting the value at $t = n$. The forecast for $t = n$ is the weighted average of the climate models, that is:

$$p_n \equiv \sum_{E=1}^N w_{E,n-1} \underline{(R_{E,n-1})} \cdot f_{E,n}. \quad (2)$$

Here, N is the number of models (*experts*) and $w_{E,n-1}$ is the weight of *expert E*, which is determined by the regret up to time $n - 1$. We used two *forecasters* (weighting schemes): the Exponentiated Weighted Average (EWA) and the Exponentiated Gradient Average (EGA).

The EWA weight is defined as:

$$w_{E,n} \equiv \frac{e^{-\eta \cdot R_{E,n}}}{\sum_{E=1}^N e^{-\eta \cdot R_{E,n}}} \frac{e^{-\eta \cdot L_{E,n}}}{\sum_{E=1}^N e^{-\eta \cdot L_{E,n}}} \quad (3)$$

and its prediction at time n is:

$$p_n = \frac{\sum_{E=1}^N e^{-\eta L_{E,n-1}} f_{E,n}}{\sum_{E=1}^N e^{-\eta L_{E,n-1}}}. \quad (4)$$

The EGA is similar to the EWA but with the cumulative regret loss calculated from the summation of the loss gradients. The cumulative regret loss for the EGA *forecaster* is defined as:

$$\underline{RL}_{E,n}^G \equiv \sum_{t=1}^n l_{E'}'(p_t, y_t) - \underline{\sum_{t=1}^n l'(f_{E,t}, y_t)} \equiv L_n^G - L_{E,n}^G \quad (5)$$

where,

$$l_{E'}'(p_t, y_t) \equiv \frac{\partial l(p_t, y_t)}{\partial w_{E,t-1}} \frac{\partial l_E(p_t, y_t)}{\partial w_{E,t-1}} = 2 \cdot (p_t - y_t) \cdot f_{E,t}. \quad (6)$$

For both *forecasters*, $\eta > 0$ is a parameter representing the learning rate.

The deviation between the forecast and the “real” trajectory was quantified using the root mean square error (RMSE). The RMSE of a grid cell with coordinates (i, j) , over a period of n time steps (months in our case), is defined as:

$$\text{RMSE}(i, j) \equiv \sqrt{(1/n) \sum_{t=1}^n (p_t(i, j) - y_t(i, j))^2}, \quad (7)$$

where $p_t(i, j)$ is the value predicted by the *forecaster* and $y_t(i, j)$ is the “real” value. The global, area-weighted RMSE is defined as:

$$G_{\text{RMSE}} \equiv (1/A_{\text{Earth}}) \sum_{i,j} A_{i,j} \text{RMSE}(i, j), \quad (8)$$

where A_{Earth} is the earth’s surface area and $A_{i,j}$ is the area of the (i, j) grid cell.

The learning rate, η , was chosen to minimize the metric $M \equiv \text{RMSE} \cdot (1 + \text{floor}(\max(\Delta w / \Delta t) / (1/N)))$ during the learning period. This metric provides a minimal deviation of the forecast climate trajectory from the observed one and also ensures stable weights of the models (a significant change in the weight of a model was considered the weight a model would be assigned in the absence of learning). We also tested [the](#) optimization of η using only a fraction of the learning period and found that as long as the optimization period was of the same order of the prediction period, there was no significant change in the outcome. An important difference between the EWA and EGA methods is that after a long enough learning period under ideal conditions (stationary time series), the former converges to the best model while the latter converges to the “real” value assuming that the real value is known. Figure 1 illustrates this difference using a simple case.

This difference between the *forecasters* implies that for [a](#) long enough learning period, using an ensemble that includes one model that [perform](#) [performs](#) better throughout the learning period, the weights will be distributed such that the prediction of the EWA will be determined by this best model and the uncertainty will be very small (due to the small weights of the other models). Under the same conditions, the EGA would still assign more significant weights to the other models in order to extract the information they contain regarding the dynamics of the “real” value ~~and this will lead~~, [leading](#) to larger uncertainty (and often better predictions).

[The learn- \$\alpha\$ algorithm \(Monteleoni and Jaakkola, 2003\) is based on the fixed-share algorithm developed by \(Herbster and Warmuth, 1998\). The fixed-share algorithm is a generalization of the EWA algorithm that increases the ability to switch between *experts* \(or between climate models in our case\) in response to changes in their performance. It is done](#)

by adding a switching probability parameter, α , that ensures that all *experts* are considered at all times. Monteleoni and Jaakkola (2003) improved this algorithm by learning the optimal switching rate between *experts*. This algorithm was already tested for long-term climate projections using the CMIP3 long-term experiments (Monteleoni et al., 2010, 2011), and here we also test its performance in decadal climate predictions for comparison with the EWA and EGA methods.

The learn- α algorithm assigns weights for each expert and for each value of the switching rate $\alpha_j \in [0, 1]$; the discrete index, $j \in 1, \dots, m$ represents the optimal discretization of α (Monteleoni and Jaakkola, 2003). The weight of each expert for a given value of α , $w_{E,t=1}(\alpha_j)$, is set initially to $1/N_e$ (N_e is the number of experts in the ensemble), and the weight of each α_j , $w_{t=1}(\alpha_j)$ is set initially to $1/N_\alpha$ (N_α is the number of discrete values of $\alpha \in [0, 1]$ that are considered). The weights are updated as follows. (i) At each time step, the loss of each model, E , is calculated in a similar manner to the EWA, $l_{E,t} \equiv (f_{E,t} - y_t)^2$. (ii) For each α_j , the loss per α is calculated, $l_t(\alpha_j) \equiv -\log\left(\sum_{E=1}^{N_e} w_{E,t}(\alpha_j) e^{-l_{E,t}}\right)$, and the weight of α_j is updated according to

$$w_{t+1}(\alpha_j) = \frac{1}{Z_t} w_t(\alpha_j) e^{-l_t(\alpha_j)}, \quad (9)$$

where Z_t normalizes the weights. (iii) For each model, E , and switching rate, α_j , the weight $w_{E,t}(\alpha_j)$ is updated according to

$$w_{E,t+1}(\alpha_j) = \frac{1}{Z_t(\alpha_j)} \sum_{E^*=1}^{N_e} w_{E^*,t}(\alpha_j) e^{-l_{E^*,t}} S(E, E^*; \alpha_j), \quad (10)$$

where,

$$S(E, E^*; \alpha_j) \equiv (1 - \alpha_j) \delta(E, E^*) + \frac{\alpha_j}{N_e - 1} (1 - \delta(E, E^*)). \quad (11)$$

$\delta(\cdot, \cdot)$ is the Kronecker delta and $Z_t(\alpha_j)$ normalizes the weights per alpha.

The prediction at $t = n$ is a weighted average of the experts and the different values of α :

$$p_n = \sum_{E=1}^{N_e} \sum_{j=1}^{N_\alpha} w_{n-1}(\alpha_j) \cdot w_{E,n-1}(\alpha_j) \cdot f_{E,n}. \quad (12)$$

One can see that in the LAA, the learning rate, $\eta = 1$, and the switching rate, α , is sequentially optimized, while for the EWA and EGA, the learning rate, η , was set to achieve the best performance during the learning period. As a successor of the EWA, the LAA also tends to converge to the best model. However, as mentioned, it can shift between models faster than the EWA, which is important when the sequences learned are nonstationary.

3 Improved predictions

We consider an ensemble of eight global climate models for the period of 1981–2011, whose results are part of the CMIP5 decadal experiments (Taylor and Meehl, 2011). Table 1 describes the eight models that we used in this study. These models were first linearly interpolated to the spatial resolution of the NCEP/NCAR reanalysis data using the NCAR command language (NCL) (NCL, 2011). We focus on the model predictions of the 2 m-temperature. The decadal experiments of the CMIP5 project include a set of runs for each of the models, representing different initial conditions. In agreement with the common knowledge (Meehl et al., 2009), we found that on decadal time scales, the internal variability of each model is smaller than the variability between the models. Therefore, we chose, arbitrarily, the first run for each of the ensemble models. The results presented here are based on a learning period of 20 years (1981–2001), followed by predictions for a 10 year (2001–2011) validation period.

The learning period served for both learning (i.e., weight assignment) and ~~also to correct~~ correcting the bias of the models. This was simply done by subtracting the average of

each of the models during the learning period and adding the average of the NCEP/NCAR reanalysis data (Kalnay et al., 1996) (considered here as reality). This bias correction was applied to each grid cell separately. ~~The bias correction and~~ was done to ~~assure ensure~~ that the improvement achieved by the forecasters ~~is was~~ beyond the impact of a simple bias correction. In addition, we chose a long enough learning period to ensure that our results ~~are were~~ not affected by the drift of the models from the ~~initial initial~~ condition toward their climate dynamics (Meehl et al., 2009).

The performance of the models was determined by comparing the model predictions to the NCEP/NCAR reanalysis data (Kalnay et al., 1996). We are aware of the spurious variability and trends in the NCEP data and of other reanalysis projects (Uppala et al., 2005; Onogi et al., 2007); however, in order to demonstrate the capability of the SLA to improve global and regional climate predictions, the reanalysis data is the best dataset ~~to use~~.

Using the predictions of the climate models only 20 years after they were initialized can ~~debate cast doubt on~~ their ability to generate skillful predictions since it is believed that climate models' skill tends to vanish after that long a period. However, we found that, for most of the models we used, this is not the case. This fact is illustrated in Fig. 2, which shows that the globally averaged RMSE of most of the climate models did not increase considerably during the 30 year-long simulations. Another noticeable and important feature of the ~~CMIP5's~~ climate models of the ~~GMIP5~~ is the fact that, globally, climatology performs much better than each of the models. In Sect. 5, we show that, despite this fact, the SLA can use the models and the climatology to provide a forecast ~~which that~~ is better than the climatology.

~~Three Four~~ forecasting methods (*forecasters*) were tested: the ~~Exponentiated Weighted Average (EWA), the Exponentiated Gradient Average (EGA)~~ ~~EWA, the EGA, the LAA~~ and a simple average. The simple average represents no learning and is presented to illustrate the superior performance of the ~~SLASLAs~~. The performance of the *forecasters* is measured by the root mean square error (RMSE), during the validation period, which quantifies the deviation of the predicted climate trajectory from the observed one.

Figure 3 shows the RMSE in the 2m-temperature monthly average prediction, during the 10 year validation period, for each grid cell. Panels a, ~~band c~~, ~~c and d~~ correspond to the RMSE of the EWA, EGA, LAA and simple average weighting schemes, respectively. ~~Both the EWA and the EGA~~ The EWA, EGA and LAA forecasters give better predictions than the simple average. The improvement achieved by the ~~two three~~ forecasters, compared with the simple average, is more apparent close to the poles and in ~~western~~ South America. In these regions, the models deviate more from each other, and the weighting schemes favor those that perform better. Over the oceans and low to ~~mid-latitudes~~ mid-latitudes, the models showed better agreement ~~and therefore~~, and therefore, the weighting schemes did not yield a large improvement.

The global, area-weighted RMSE can be used to quantify the improvement achieved by the SLA *forecasters*, that is, 1.316°C for the EWA, 1.297°C for the EGA, 1.372°C for the LAA and 1.390°C for the simple average. Since the EWA has the tendency to converge to the best model (if the ensemble includes a model that is always better than the others in certain regions), we also ~~compare~~ compared the performance of the EWA and EGA *forecasters* with two forecasting methods that predict according to the best model (defined as the model that was assigned the highest weight according to either the EWA or the EGA) in each grid cell. The global, area-weighted RMSE was found to be 1.568°C for the best model based on the EWA and 1.633°C for the best model based on the EGA. These results show that the SLA *forecasters* outperform the best models in the ensemble. In general, we found that a longer learning period improves the predictions of the *forecasters*. Figure 4 shows that the area-weighted RMSE of the *forecasters* (during the validation period) is reduced when the learning period is extended. By increasing the learning rate, we found that shorter learning periods can be selected with no significant increase in error; however, we chose a learning period ~~which that~~ is of the order of the prediction period in order to capture the climate dynamics in all the time scales that are relevant to the prediction period.

4 Reduced uncertainties

The weights obtained from the SLA method can be used to better estimate the uncertainties of the predictions. The uncertainties are quantified by the square root of the time average of the weighted variance of the ensemble. This quantity (for a period of n time steps) in the (i, j) grid cell is defined as:

$$\text{STD}(i, j) \equiv \sqrt{(1/n) \sum_{t=1}^n \sum_{E=1}^N w_E(i, j) (f_{E,t}(i, j) - p_t(i, j))^2}. \quad (13)$$

Here, $f_{E,t}(i, j)$ is the prediction of model E for grid cell (i, j) , at time t ; $p_t(i, j)$ is the prediction of the *forecaster* for grid cell (i, j) , at time t (i.e., the weighted average of all the models); and $w_E(i, j)$ is the weight assigned to model E at grid cell (i, j) (the weights remain constant during the validation period for which the STD is calculated). The global, area-weighted uncertainty is defined as:

$$G_{\text{STD}} \equiv (1/A_{\text{Earth}}) \sum_{i,j} A_{i,j} \text{STD}(i, j). \quad (14)$$

Figure 5 shows the uncertainty of the 2m-temperature during the validation period for the three forecasting methods; panels a, ~~b and c~~, c and d correspond to the EWA, EGA, LAA and simple average *forecasters*, respectively. It is important to note that this uncertainty is only due to the different predictions of the ensemble models; other sources of uncertainty are not affected by our forecasting schemes. ~~Both the EWA and EGA~~ The three learning algorithms, EWA, EGA and LAA forecasters, yield smaller uncertainties than does the simple average. The improvement is significant in regions where the uncertainties are larger, such as toward the poles and over South America and Africa. The global, area-weighted ~~uncertainties~~ uncertainties are: ~~1.242, 1.381, 1.242°C, 1.381°C, 1.078°C~~ and 1.593°C for the EWA, EGA, LAA and simple average *forecasters*, respectively. These values show that in addition to improving the predictions, the SLA *forecasters* also reduce the uncertainties of these predictions. Note that the smaller uncertainty of the EWA and the LAA forecaster forecasters is

simply due to the fact that ~~this~~ ~~these~~ ~~forecaster~~ ~~forecasters~~ ~~converges~~ ~~converge~~ to the best model in each grid cell (if the ensemble includes a model that is always the best). The uncertainty of the EGA provides a better estimate of the ~~predictions~~ ~~predictions1000~~ uncertainty because its predictions ~~converges~~ ~~converge~~ to the observations.

5 Skillful forecast

The skill of a *forecaster* may be defined as its ability to provide better predictions than the reference climatology. In our study, the natural choice is the climatology of the learning period, that is:

$$C_m \equiv \frac{1}{L} \sum_{i=1}^L y_{i,m}, \quad (15)$$

where, $y_{i,m}$ is the value of the variable (in this study, it is the 2 m-temperature as reported in the reanalysis data) in the calendar month m of the year i ; the learning period duration is L years; and the climatology, C_m , is just the average of that variable during the L years. ~~Prediction~~ ~~A prediction~~ that is based on climatology assumes that for each month of the prediction period, the value of the variable will be equal to the climatology of the corresponding calendar month. Therefore, it is reasonable to expect that a skillful model should provide more information on the variability of the climate than the average of previous years (the climatology).

Figure 6a shows the differences between the 10 year RMSE of the 2 m-temperature monthly mean, of the climatology and of the EGA *forecaster*. Positive values represent locations where the EGA *forecaster* has a smaller RMSE and is, therefore, considered as a skillful *forecaster*. In most regions, the climatology performs better than the EGA *forecaster* (and, obviously, better than the best model!); however, some regions indicate the EGA's advantage, such as eastern North America up to Greenland. We found that the regions in which the EGA *forecaster* performs better are characterized by larger variability

(which increases the deviations from the climatology). The global, area-weighted RMSE is 1.188°C for the climatology and 1.373°C for the EGA. One could conclude that the EGA *forecaster* is not skillful.

To circumvent this problem, we decided to add the climatology of the learning period as an additional model to our ensemble. In Fig. 6b, we show the difference between the RMSE of the EGA *forecaster*, for the model ensemble that includes the climatology, and the RMSE of the climatology itself. In this figure, one can see that the EGA *forecaster*, for the model ensemble that includes the climatology, provides predictions that are at least as good as the climatology over most of the globe. Adding the climatology to the ensemble reduced the global, area-weighted RMSE of the EGA *forecaster* to 1.156°C —a small improvement (a reduction of about 2.7%) over the climatology. The global, area-weighted RMSE of the EWA, LAA and simple average with climatology are 1.187°C , 1.180°C and 1.337°C , respectively. The global, area-weighted uncertainties of the 10 year validation period, in this case are: 0.118°C , 0.953°C , 0.118°C , 0.953°C , 0.836°C , and 1.552°C for the EWA, EGA, LAA and simple average *forecasters*, respectively. Note that as we mentioned earlier, the small uncertainty associated with the EWA *forecaster* is not representative of the climate prediction uncertainty. In what follows, we focus on the significance of the results of the EGA *forecaster*.

6 Significance tests

There is more than one test that can be done to demonstrate the significance of the results. We focus on testing whether the EGA *forecaster* improves the predictions beyond climatology (as shown earlier, each of the models performs poorer—worse than the climatology) and whether it reduces the uncertainties below those of an equally weighted ensemble. Both tests were done globally and regionally. We start by defining two properties. The first is the difference between the absolute error of the climatology and the absolute error of the EGA *forecaster* at a given grid cell and time point, that is, $|(C_t(i, j) - y_t(i, j))| - |(p_t(i, j) - y_t(i, j))|$. The second is the difference between the uncer-

tainties of the equally weighted ensemble and the ensemble weighted according to [the EGA forecaster](#) at a given grid cell and time point, that is
$$\sqrt{\frac{1}{N} \sum_{E=1}^N (f_{E,t}(i,j) - f_{,t}(i,j))^2} - \sqrt{\sum_{E=1}^N w_E(i,j) \cdot (f_{E,t}(i,j) - p_t(i,j))^2}$$
 (the dot replacing the E index $f_{,t}$ represents averaging over that index). For both quantities, positive values represent [a better performance of the EGA forecaster](#). The 10 year validation period yields [for each of these quantities, a time series with 120 points in each grid cell](#). The fraction of the time series ([the number of points out of the total 120](#)) showing positive values can be used to test the significance of the improvement. We define a significant improvement by the EGA forecaster to be when the number of successes are above 66 (i.e., when the null hypothesis that the quantities defined above are symmetrically distributed around zero is rejected with $\sim 90\%$ confidence).

Figure 7 shows the spatial distributions of the number of positive values (out of the total 120 time points) for the two quantities. The upper panel corresponds to the difference between the absolute error of the climatology and the EGA forecaster, and the lower panel corresponds to the difference between the uncertainties of the equally weighted and EGA weighted ensembles.

The upper panel in Fig. 7 shows that there are large regions of improvement, which is more apparent over land, close to the poles and to the equator. The lower panel shows that in regions in which the EGA reduces the uncertainty, it does so for almost all time points and vice versa. No correlation between significant improvement of the predictions and significant reduction of the uncertainties was identified.

The global test we performed was done by calculating the area weighted average of the two quantities defined above and ~~to plot~~ [plotting](#) the histograms of their time series. These are shown in Fig. 8. The upper panel shows the ~~global average~~ [globally averaged](#) absolute error difference between the climatology and the EGA forecaster, and the lower panel shows the ~~global average~~ [globally averaged](#) difference between the uncertainties of the equally weighted and EGA weighted ensembles. The x axis is in units of $^{\circ}\text{C}$ and is zero centered to emphasize the ~~non-symmetric~~ [nonsymmetrical](#) distribution of the data. The upper panel shows that there are only 11 negative values out of 120 and a positive

peak at around 0.03°C . The probability of more than 108 positive values out of 120 in a ~~symmetric distribution with~~ symmetrical distribution with a zero mean is practically zero; therefore, we conclude that, globally, the EGA forecaster predicts better than climatology. The ~~uncertainties difference~~ difference in uncertainties shows that the EGA *forecaster* has lower uncertainty than the equally weighted ensemble for all the time points ~~and therefore,~~ and therefore, we can also conclude that the reduction of the globally averaged uncertainties is significant.

7 Summary and discussion

The SLA method does not rely on any assumptions regarding the distributions of the climate variables; therefore, it is robust and can be used for any climate variable. The updating scheme of the weights does not require a considerable computational cost and allows for a fast and easy update of the weights when new measurements become available. In the results presented here, we used the deviation from the trajectory of the climate variable as the metric for the weighting, but other weighting methods can also be applied. For example, one can use a measure of the statistical distance, such as the Kullback–Leibler divergence (Kullback and Leibler, 1951) or the Jensen–Shannon divergence (Manning and Schütze, 1999); a model that yields a probability density function (PDF) ~~which that~~ is closer to the measured PDF of a variable will get a higher weight.

One disadvantage of the SLA method (which may also be considered as an advantage for some applications) is the fact that the weights are between zero and one. This means that if the measurements are not spanned by the predictions of the models, the SLA algorithm will not be able to track the measurements but would converge to the best model since ~~by definition,~~ by definition, the SLA predictions are bounded by the predictions of the models of the ensemble. In this case, other methods, such as the regression that can yield any linear combination of the model predictions, may achieve better predictions than the SLA *forecasters* but will not be able to reduce the ensemble uncertainties.

We showed that climate predictions (on a decadal time scale) of the 2m-temperature monthly average can be improved and that the associated uncertainties can be reduced using the [SLA algorithms](#). [The largest improvement was found using the EGA forecaster](#). [The small improvement achieved by the EWA and LAA, when the climatology was added as an expert to the ensemble, is only associated with the fact that they tracked the climatology in regions where it performed the best. Therefore, the EWA and LAA did not show significant improvement compared to climatology.](#)

[The improvement of the LAA over the EWA is too small to be considered significant. Moreover, both the EWA and the LAA performed worse than the EGA. These results suggest that in decadal climate prediction, the nonstationary nature of the climate system does not play a major role. The more significant improvement is achieved when focusing on tracking the best prediction rather than the best model \(Cesa-Bianchi and Lugosi, 2006\).](#)

[The improved predictions and reduced uncertainties considered here are only those arising from the variability between different models. This is because the ensemble used in this study consists of only one run \(corresponding to one initial condition\) of each of the models. The uncertainties due to the internal variability of each of the models remained unaffected. In principle, the SLA method can be used to quantify the quality of different initialization methods. However, there is no justification for weighting initial conditions generated by the same method at times that are of the same order of magnitude before the prediction period. Therefore, the SLA method cannot reduce uncertainties associated with the internal variability of the models.](#)

The SLA method provided better predictions than each one of the models and their simple average. All the models, including the simple average, considered in this study showed no global skill; namely, in averaging over the globe, the climatology provided a better prediction than each of the models. The SLA *forecasters* do not resolve this issue unless the climatology is added as an additional model to the ensemble. When the model ensemble includes the climatology, the SLA *forecasters* can yield better predictions than the climatology itself by assigning high weight to the climatology in the regions where the models fail and

high weight to the best models in regions where they perform better than the climatology (namely, regions where the best models are skillful).

The method and the results presented here provide performance-based, ~~spatially-distributed~~ spatially distributed weights of climate models, which lead to improved climate predictions and reduced uncertainties. These can be relevant for many applications in agriculture and ecology, and for decision makers and other stakeholders. The ~~spatially-distributed~~ spatially distributed weights may also be used for testing new parameterization and physics schemes in global circulation models.

Acknowledgements. The research leading to these results has received funding from the European Union Seventh Framework Programme (FP7/2007-2013) under grant number [293825].

References

- Buser, C. M., Künsch, H. R., Lüthi, D., Wild, M., and Schär, C.: Bayesian multi-model projection of climate: bias assumptions and interannual variability, *Clim. Dynam.*, 33, 849–868, 2009.
- Buser, C., Künsch, H., and Schär, C.: Bayesian multi-model projections of climate: generalization and application to ENSEMBLES results, *Clim. Res.*, 44, 227–241, 2010.
- Cesa-Bianchi, N. and Lugosi, G.: *Prediction, Learning, and Games*, Cambridge University Press, Cambridge, UK, 2006.
- Chakraborty, A. and Krishnamurti, T. N.: Improving global model precipitation forecasts over India using downscaling and the FSU superensemble. Part II: Seasonal climate, *Mon. Weather Rev.*, 137, 2736–2757, 2009.
- Collins, M.: Ensembles and probabilities: a new era in the prediction of climate change, *Philos. T. R. Soc. A*, 365, 1957–1970, 2007.
- Doblas-Reyes, F. J., Déqué, M., and Piedelievre, J.-P.: Multi-model spread and probabilistic seasonal forecasts in PROVOST, *Q. J. Roy. Meteor. Soc.*, 126, 2069–2087, 2000.
- Feng, J., Lee, D.-K., Fu, C., Tang, J., Sato, Y., Kato, H., McGregor, J., and Mabuchi, K.: Comparison of four ensemble methods combining regional climate simulations over Asia, *Meteorol. Atmos. Phys.*, 111, 41–53, 2011.
- Hagedorn, R., Doblas-Reyes, F. J., and Palmer, T. N.: The rationale behind the success of multi-model ensembles in seasonal forecasting – I. Basic concept, *Tellus A*, 57, 219–233, 2005.

- Hawkins, E. and Sutton, R.: The potential to narrow uncertainty in regional climate predictions, *B. Am. Meteorol. Soc.*, 90, 1095–1107, 2009.
- [Herbster, M. and Warmuth, M. K.: Tracking the best expert, *Mach Learn.* 32, 151–178, 1998.](#)
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *B. Am. Meteorol. Soc.*, 77, 437–471, 1996.
- Kalnay, E., Hunt, B., Ott, E., and Szunyogh, I.: Ensemble forecasting and data assimilation: two problems with the same solution?, in: *Predictability of Weather and Climate*, edited by: Palmer, T. N. and Hagedorn, R., Cambridge University Press, Cambridge, 157–180, 2006.
- Kim, H.-M., Webster, P. J., and Curry, J. A.: Evaluation of short-term climate change prediction in multi-model CMIP5 decadal hindcasts, *Geophys. Res. Lett.*, 39, L10701, doi:10.1029/2012GL051644, 2012.
- [Kivinen, J. and Warmuth, M.: Exponentiated gradient versus gradient descent for linear predictors. *Information and Computation*, 132\(1\):1–63, 1997.](#)
- Krishnamurti, T. N.: Improved weather and seasonal climate forecasts from multimodel superensemble, *Science*, 285, 1548–1550, 1999.
- Krishnamurti, T. N., Kishtawal, C. M., Zhang, Z., LaRow, T., Bachiochi, D., Williford, E., Gadgil, S., and Surendran, S.: Multimodel ensemble forecasts for weather and seasonal climate, *J. Climate*, 13, 4196–4216, 2000.
- Kullback, S. and Leibler, R. A.: On information and sufficiency, *Ann. Math. Stat.*, 22, 79–86, doi:10.1214/aoms/1177729694, 1951.
- [Littlestone, N. and Warmuth, M.: The weighted majority algorithm. *Information and Computation*, 108:212–261, 1994.](#)
- Mallet, V.: Ensemble forecast of analyses: coupling data assimilation and sequential aggregation, *J. Geophys. Res.-Atmos.*, 115, D24303, doi:10.1029/2010JD014259, 2010.
- Mallet, V., Stoltz, G., and Mauricette, B.: Ozone ensemble forecast with machine learning algorithms, *J. Geophys. Res.-Atmos.*, 114, D05307, doi:10.1029/2008JD009978, 2009.
- Manning, C. D. and Schütze, H.: *Foundations of Statistical Natural Language Processing*, MIT Press, Cambridge, MA, USA, 1999.
- Meehl, G. A., Goddard, L., Murphy, J., Stouffer, R. J., Boer, G., Danabasoglu, G., Dixon, K., Giorgetta, M. A., Greene, A. M., Hawkins, E., Hegerl, G., Karoly, D., Keenlyside, N., Kimoto, M., Kirt-

- man, B., Navarra, A., Pulwarty, R., Smith, D., Stammer, D., and Stockdale, T.: Decadal prediction, *B. Am. Meteorol. Soc.*, 90, 1467–1485, 2009.
- [Monteleoni, C. and Jaakkola, T.: Online Learning of Non-stationary Sequences, in Advances in Neural Information Processing Systems, 16, 1093–1100, 2003.](#)
- [Monteleoni, C., Saroha, S., and Schmidt, G.: Tracking Climate Models, In NASA Conference on Intelligent Data Understanding \(CIDU\), pages 1–15, 2010.](#)
- [Monteleoni, C., Saroha, S., Schmidt, G., and Asplund, E.: Tracking Climate Models, In Journal of Statistical Analysis and Data Mining: Special Issue: Best of CIDU 2010, Volume 4, Issue 4, pages 72–392, 2011.](#)
- Onogi, K., TsuTsu, J., Koide, H., Sakamoto, M., Kobayashi, S., Hatsushika, H., Matsumoto, T., Yamazaki, N., Kamahori, H., Takahashi, K., Kadokura, S., Wada, K., Kato, K., Oyama, R., Ose, T., Mannoji, N., and Taira, R.: The JRA-25 reanalysis, *J. Meteorol. Soc. Jpn.*, 85, 369–432, 2007.
- Palmer, T. N., Doblas-Reyes, F. J., Hagedorn, R., Alessandri, A., Gualdi, S., Andersen, U., Feddersen, H., Cantelaube, P., Terres, J.-M., Davey, M., Graham, R., Délecluse, P., Lazar, A., Déqué, M., Guérémy, J.-F., Díez, E., Orfila, B., Hoshen, M., Morse, A. P., Keenlyside, N., Latif, M., Maisonave, E., Rogel, P., Marletto, V., and Thomson, M. C.: Development of a European multimodel ensemble system for seasonal-to-interannual prediction (DEMETER), *B. Am. Meteorol. Soc.*, 85, 853–872, 2004.
- Pavan, V. and Doblas-Reyes, F. J.: Multi-model seasonal hindcasts over the Euro-Atlantic: skill scores and dynamic features, *Clim. Dynam.*, 16, 611–625, 2000.
- Rajagopalan, B., Lall, U., and Zebiak, S. E.: Categorical climate forecasts through regularization and optimal combination of multiple GCM ensembles, *Mon. Weather Rev.*, 130, 1792–1811, 2002.
- Robertson, A. W., Lall, U., Zebiak, S. E., and Goddard, L.: Improved combination of multiple atmospheric GCM ensembles for seasonal prediction, *Mon. Weather Rev.*, 132, 2732–2744, 2004.
- Samuels, R., Harel, M., and Alpert, P.: A new methodology for weighting high-resolution model simulations to project future rainfall in the Middle East, *Clim. Res.*, 57, 51–60, 2013.
- Smith, R. L., Tebaldi, C., Nychka, D., and Mearns, L. O.: Bayesian modeling of uncertainty in ensembles of climate models, *J. Am. Stat. Assoc.*, 104, 97–116, 2009.
- Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: A Summary of the CMIP5 Experiment Design, available at: http://cmip-pcmdi.llnl.gov/cmip5/experiment_design.html (last access: 7 March 2015), 2011.
- Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An overview of CMIP5 and the experiment design, *B. Am. Meteorol. Soc.*, 93, 485–498, 2012.

- The NCAR Command Language (Version 6.0.0) [Software], doi:10.5065/D6WD3XH5, 2011.
- Tebaldi, C. and Knutti, R.: The use of the multi-model ensemble in probabilistic climate projections, *Philos. T. R. Soc. A*, 365, 2053–2075, 2007.
- Uppala, S. M., Kållberg, P. W., Simmons, A. J., Andrae, U., Bechtold, V. D. C., Fiorino, M., Gibson, J. K., Haseler, J., Hernandez, A., Kelly, G. A., Li, X., Onogi, K., Saarinen, S., Sokka, N., Allan, R. P., Andersson, E., Arpe, K., Balmaseda, M. A., Beljaars, A. C. M., Berg, L. V. D., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher, M., Fuentes, M., Hagemann, S., Hólm, E., Hoskins, B. J., Isaksen, L., Janssen, P. A. E. M., Jenne, R., McNally, A. P., Mahfouf, J.-F., Morcrette, J.-J., Rayner, N. A., Saunders, R. W., Simon, P., Sterl, A., Trenberth, K. E., Untch, A., Vasiljevic, D., Viterbo, P., and Woollen, J.: The ERA-40 re-analysis, *Q. J. Roy. Meteor. Soc.*, 131, 2961–3012, 2005.
- Warner, T. T.: *Numerical Weather and Climate Prediction*, Cambridge University Press, Cambridge, UK, 2011.
- Yun, W. T., Stefanova, L., and Krishnamurti, T. N.: Improvement of the multimodel superensemble technique for seasonal forecasts, *J. Climate*, 16, 3834–3840, 2003.
- Yun, W. T., Stefanova, L., Mitra, A. K., Kumar, T. S. V. V., Dewar, W., and Krishnamurti, T. N.: A multi-model superensemble algorithm for seasonal climate prediction using DEMETER forecasts, *Tellus A*, 57, 280–289, 2005.

Table 1. ~~Models Availability~~Model Availabilities.

Institute ID	Model Name	Modeling Center (or Group)	Grid (lat × lon)
BCC	BCC-CSM1.1	Beijing Climate Center, China Meteorological Administration	64 × 128
CCCma	CanCM4	Canadian Centre for Climate Modelling and Analysis	64 × 128
CNRM-CERFACS	CNRM-CM5	Centre National de Recherches Meteorologiques/Centre Europeen de Recherche et Formation Avancees en Calcul Scientifique	128 × 256
LASG-IAP	FGOALS-s2	LASG, Institute of Atmospheric Physics, Chinese Academy of Sciences	108 × 128
IPSL	IPSL-CM5A-LR	Institut Pierre-Simon Laplace	96 × 96
MIROC	MIROC5 MIROC4h	Atmosphere and Ocean Research Institute (The University of Tokyo), National Institute for Environmental Studies, and Japan Agency for Marine-Earth Science and Technology	128 × 256 320 × 640
MRI	MRI-CGCM3	Meteorological Research Institute	160 × 320




Figure 1. An ideal experiment with two experts. The first always predicts zero and the second always predicts one. The true value is always 0.7. The EWA forecaster converges to the best model (predicting one) while the EGA forecaster converges to the true value.




Figure 2. Temporal evolution of the global and annual average of the 2-m-temperature RMSE for the eight climate models (after bias correction) and the climatology. During the 30 years of the simulations, the skill of most of the models did not decline. In fact, a simple linear fit to the models indicates that some of them increased their skill with time.




Figure 3. 10 year RMSE of the 2-m-temperature for three forecasting methods: (a) EWA, (b) EGA, and (c) LAA, and (d) the simple average. The colors represent the RMSE of each grid cell. Both All the SLA forecasters yield a smaller global RMSE than the simple average. The improvements achieved by the EWA and EGA forecasters, compared with the simple average, are more apparent close to the poles and in southwestern America.




Figure 4. Global, area-weighted RMSE of the 2 m-temperature, during the 10 year validation period, as a function of the learning time. The presented RMSE was calculated for the EGA forecaster; however, a similar trend was obtained for the EWA [and LAA](#). In general, a longer learning period improves the *forecaster* predictions.




Figure 5. The 2 m-temperature uncertainty during the 10 year validation period for three forecasting methods: (a) EWA, (b) EGA, and (c) LAA, and (d) the simple average. The uncertainties of the EWA and LAA are smaller than those of the EGA; however, the predictions of the EGA are better (see the text for a more detailed explanation). Both the EGA and All the EWA forecasters yield smaller uncertainties than the simple average. The uncertainties, corresponding to the SLA forecasting schemes, are significantly reduced in regions where the uncertainties are larger, such as toward the poles and over South America and Africa.




Figure 6. The difference between the 10 year validation period average 2m-temperature RMSE of the climatology and the EGA forecaster, **(a)** EGA with an ensemble that includes eight models, **(b)** EGA with an ensemble that includes the same eight models and also the climatology of the learning period as an additional model. The results demonstrate that when the ensemble includes the climatology, the EGA forecaster is skillful.




Figure 7. The number of time points in which the EGA *forecaster* performs better. The upper panel shows the spatial distribution of the number of time points in which the absolute error of the EGA *forecaster* is smaller than that of the climatology. The lower panel shows the spatial distribution of the number of time points in which the uncertainty of the EGA weighted ensemble is smaller than that of the equally weighted ensemble. White circles represent significant improvement by the EGA *forecaster* and black circles represent its significantly poorer performance. Both quantities show better performance of the EGA *forecaster* over most of the [globglobe](#).




Figure 8. The histograms of the globally averaged differences of absolute error and uncertainty. The upper panel shows the histogram of the globally averaged difference between the absolute error of the climatology and that of the EGA *forecaster*. The lower panel shows the histogram of the difference between the uncertainties of equally weighted and EGA weighted ensembles. Both quantities show significantly improved performance of the EGA *forecaster*.