

Potential source regions and processes of the aerosol in the summer Arctic

Jost Heintzenberg¹, Caroline Leck², and Peter Tunved³

1: Leibniz-Institute for Tropospheric Research, Permoser Str. 15, 04318 Leipzig, Germany

2: Department of Meteorology, Arrhenius Laboratory, Stockholm University, 10691
Stockholm, Sweden

3: Department of Applied Environmental Science, Stockholm University, 10691 Stockholm,
Sweden

Abstract

Sub-micrometer particle size distributions measured during four summer cruises of the Swedish icebreaker *Oden* 1991, 1996, 2001, and 2008 were combined with dimethyl sulfide gas data, back trajectories and daily maps of pack ice cover in order to investigate source areas and aerosol formation processes of the boundary layer aerosol in the central Arctic. With a clustering algorithm potential aerosol source areas were explored. Clustering of particle size distributions together with back-trajectories delineated five potential source regions and three different aerosol types that covered most of the Arctic basin: Marine, Newly formed and aged particles over the pack ice. Most of the pack ice area with < 15% percent of open water under the trajectories exhibited the aged aerosol type with only one major mode around 40 nm. For newly formed particles to occur two conditions had to be fulfilled over the pack ice: The air had spent ten days while traveling over ever more contiguous ice with less than 30% open water during the last five days. Additionally, the air had experienced more open water (at least twice as much as in the cases of aged aerosol) during the last four days before arrival in heavy ice conditions at *Oden*. Thus we hypothesize that these two conditions were essential factors for the formation of ultrafine particles over the central Arctic pack ice. A comparison the *Oden* data with summer size distribution data from Alert, Nunavut and Mt. Zeppelin, Spitsbergen confirmed the *Oden* findings with respect to particle sources over the central Arctic. Future more frequent broken-ice or open water patches in summer will spur biological activity in surface water promoting the formation of biological particles. Thereby low clouds and fogs and subsequently the surface energy balance and ice melt may be affected.

1. Introduction

The investigation of the summer aerosol over the central Arctic Ocean began with the first Swedish Arctic icebreaker expedition (*Ymer*-80) in 1980 (Lannefors et al., 1983) followed up later in a series of four international ice-breaker expeditions to the summer central Arctic Ocean on the Swedish icebreaker *Oden* in the years 1991 (Leck et al., 1996), 1996 (Leck et al., 2001), 2001 (Leck et al., 2004), and 2008 (Tjernström et al., 2014).

As illustrated in Fig. 1, several hypothesized sources may contribute to the aerosol over the central Arctic Ocean, and thus to the formation of low-level stratiform clouds and their effects on the surface energy balance. Long-range transported biomass burning or pollution plumes has been observed in helicopter profiles. These plumes always occurred in the free troposphere well above the top of the boundary layer and were rarely mixed down to the surface (Kupiszewski et al., 2013). This finding is consistent with light absorbing surface aerosol measurements over the summer pack ice indicating extremely low concentrations on the order of a few nanograms of black carbon per cubic meter (Heintzenberg, 1982; Maenhaut et al., 1996).

Transport of precursor gases and marine biogenic particles (specifically polymer gels¹) from the marginal ice zone (MIZ) or locally from open leads² over the pack ice has been found to result in raised concentrations of accumulation mode particles within the high Arctic boundary layer (Chang et al., 2011; Heintzenberg and Leck, 2012; Heintzenberg et al., 2006;

¹ Marine gels or polymer gels are produced by phytoplankton and biological secretions of sea ice algae at the sea-air interface. The polymer gels are made up of water-insoluble, heat resistant, highly surface-active and highly hydrated (99% water) polysaccharide molecules spontaneously forming 3-dimensional networks inter-bridged with divalent ions ($\text{Ca}^{2+}/\text{Mg}^{2+}$), to which other organic compounds, such as proteins and lipids, are readily bound E.K. Bigg, C. Leck and L. Tranvik, Particulates of the surface microlayer of open water in the central Arctic Ocean in summer, *Mar. Chem.* **91**(2004), 131-141, Q. Gao, P. Matrai and C. Leck, On the chemical dynamics of extracellular polymeric secretions (polysaccharides) in the high Arctic surface microlayer, *ibid.* **8**(2011), 401-418, C. Leck, Q. Gao, F. Mashayekhy Rad and U. Nilsson, Size-resolved atmospheric particulate polysaccharides in the high summer Arctic, *Atmos. Chem. Phys.* **13**(2013), 12573-12588, M.V. Orellana *et al.*, Marine microgels as a source of cloud condensation nuclei in the high Arctic, *PNAS* **108**(2011), 13612–13617.

² The high Arctic open leads can be described as ever-changing open water channels comprising 10-30% of the ice pack ice area, ranging from a few meters up to a few kilometers in width.

1 Hellén et al., 2012; Kupiszewski et al., 2013; Leck et al., 2013; Nilsson and Leck, 2002).
2 This may involve both direct emissions of primary larger accumulation mode marine
3 particles, as well as growth of smaller particles via two processes, namely heterogeneous
4 condensation and aerosol cloud processing.

5 Particles advected into the central Arctic within the boundary layer frequently experience
6 efficient scavenging processes associated with low clouds and fog near the MIZ
7 (Heintzenberg and Leck, 2012; Nilsson and Leck, 2002) which explains their later very low
8 near-surface aerosol concentrations.

9 Heterogeneous condensation and aerosol cloud processing occurs when the oxidation
10 products of dimethyl sulfide (DMS) released by phytoplankton advected from open waters
11 south of and along the marginal ice edge, (Leck and Persson, 1996a), condense on non-
12 activated particles which then are incorporated into cloud droplets. In the latter droplets
13 liquid-phase oxidation of absorbed gases can add further material to the droplet constituents.
14 Evaporated cloud droplets leave behind raised concentrations of accumulation mode particles,
15 grown via the two processes described. This process creates the bimodal particle size
16 distribution characteristic of cloud-processed air (Hoppel et al., 1994).

17 New particle formation (nucleation) occurred about 15% of the observed time period (Karl
18 et al., 2013). However, these events often manifested themselves as a simultaneous increase
19 of particle number concentrations in the < 10 nm and 20–50 nm size ranges, and not as the
20 prototypical “banana growth” (e.g., c.f. Kulmala et al., 2001). Conventional nucleation
21 paradigms (Karl et al., 2012) fail to explain this phenomenon. An alternate hypothesis
22 explaining this could be fragmentation and/or dispersion of primary marine polymer gels,
23 200–500 nm diameter in size, into the nanogel size fractions down to a few nanometer
24 polymers (Karl et al., 2013; Leck and Bigg, 2010).

25 While the four expeditions provided a wealth of new observations and understanding of the
26 system of low-level clouds, their formation, and their effects on the boundary-layer and

1 surface energy balance over the Arctic pack ice area, the ultimate partitioning of aerosol
2 particles among potential source regions and processes remains elusive. The present paper
3 continues the analysis of the aerosol data from the four *Oden* cruises with a focus on the
4 above discussed potential source regions and related aerosol formation processes. The ship
5 positions during the cruises shown in Fig. 2 indicate that the measured data only cover a small
6 part of the European Arctic sector. However, with back trajectories the data coverage can be
7 extended over the whole Arctic basin. This approach was first followed with aerosol data
8 measuring during the *Ymer*-80 expedition by Jaenicke and Schütz (1982) and with Norwegian
9 Arctic aerosol data by Heintzenberg and Larsen (1983). For the present study back trajectory
10 information was complemented with daily maps of ice concentrations. Sections 2.3 and 2.4
11 give more details. For the combination of aerosol data and information of air origin and ice
12 data a dedicated cluster algorithm was developed. For a test of the clustering algorithm the
13 aerosol database was complemented with the data on atmospheric dimethyl sulfide (DMS(g))
14 concentrations taken during all four cruises (Kettle et al., 1999; Leck and Persson, 1996b).

15 To date, 23 years after the first *Oden* expedition, there are still no other surface aerosol
16 data from the central Arctic to compare with. The nearest land stations are Mt. Zeppelin,
17 Spitsbergen and Alert, Nunavut. The present paper therefore also makes an attempt to
18 connect the size distributions taken on *Oden* and the clusters derived with them with size
19 resolved aerosol number data and trajectories from these two land stations.

20 With the combined data set and the clustering algorithm the main goal of the present study
21 is to identify potential source regions of aerosol particles observed over the central summer
22 Arctic. Specifically, we would like to differentiate between local sources within the pack ice
23 region and distant sources. Extending our previous analyses discussed above with the locally
24 measured parameters to different source regions we aim at identifying factors controlling the
25 aerosol life cycle over the inner Arctic.

2. Experimental data

2.1 Sampling conditions on icebreaker *Oden*

All four icebreaker expeditions utilized an identical sampling manifold upstream of all gas phase and aerosol instrumentation. This manifold extended at an angle of 45° to about three meters above the container roof of the laboratory container on *Odens'* 4th deck to optimize the distance both from the sea and from the ship's superstructure. The height of the sampling manifold was ~ 25 m above sea level and consisted of two masts (PM_1 : Diameter $< 1\mu m$ and PM_{10} : Diameter $< 10\mu m$), with one additional sampling line for volatile organic compounds including DMS. Direct contamination from the ship was minimized with a pollution controller. Provided that the wind was within $\pm 70^\circ$ of the direction of the bow and stronger than 2 ms^{-1} , no pollution reached the sample inlets. Further details of the instrumentation and precautions to exclude contaminated periods can be found in Leck et al., (2001) and in Tjernström et al., (2014).

2.2 Data collected onboard *Oden*

2.2.1 Gas data

As compared to the 1271 hourly DMS values, which were concurrent with contamination-free aerosol data a total of 2035 hours of DMS data were available in the four cruises for clustering.

During the expedition in 1991, integrated samples of DMS were analyzed by a Gas Chromatograph (GC)-Flame Photometric Detection (FPD) system where a glass-fiber-wool cold-trap was used in the pre-concentration step. The sampling duration was 20 min (Persson and Leck, 1994). During the three subsequent cruises, DMS was automatically collected with a time resolution of 15 min and pre-concentrated in the following two steps: first, a gold trap (gold wire in a Pyrex glass tube) for collection, and second, a (TENAX®) medium to achieve a sharp injection of the analyte into the GC-FPD. To remove atmospheric oxidants prior to collection, a high-capacity scrubber based on 100% cotton wadding was used (Persson and Leck, 1994) in all four cruises. The overall accuracy, valid for both GC-FPD methods described above, was within $\pm 12\%$ with a detection limit of $0.045 \text{ nmol m}^{-3}$.

To further improve on time resolution, we added a Proton Transfer Reaction Mass Spectrometer system (PTR-MS) (Lindinger and Hansel, 1998) during the 2001 cruise with a sampling frequency of 2 min and in the 2008 experiment DMS was measured a PTR-TOFMS (Aerosol Time of Flight Mass Spectrometer) built at Innsbruck University. The PTR-TOFMS was calibrated by applying a dynamically diluted DMS gas standard (Apel & Riemer Environmental Inc.). Zero-calibrations were performed every 2–6 h using catalytically scrubbed air. The sampling frequency of the PTR- TOFMS system was 1 min. The instrument is described in detail in (2010). For the benefit of time resolution of the PTR systems, the detection limit was increased by a factor of ten to 0.45 nmol m^{-3} . $1 \text{ nmol m}^{-3} = 22.4 \text{ ppt(v)}$ at 0° C and 1013.25 mbar .

2.2.2 Aerosol data

The *Oden* aerosol database is essentially the same as in Heintzenberg and Leck (2012) with 2645 hours of sub-micrometer particle number size distributions between 5 and 560 nm diameter. Tandem Differential Mobility Particle sizers (TDMPS) were used to measure the number size distributions of dry sub-micrometer particles with pairs of very similar

differential mobility analyzers (DMAs). The TSI 3010 counters used in the DMAs were size and concentration calibrated against an electrometer and the TSI 3025 counters for particle sizes below 20 nm diameter in the standard way after Stolzenburg (1988). In 1996 a second, modified TSI 3010 was utilized to extend the data from 20 to 5 nm instead of a TSI 3025. The harmonized size range for all cruises comprised 36 channels, which were spaced in equidistant fashion on a logarithmic scale. Before taking hourly averages the data had been cleaned thoroughly for possible pollution from the ship (cf. Heintzenberg and Leck (2012) for details). In 1991 the Arctic part of the cruise covered the time from August 18 through September 26. In 1996 the icebreaker stayed in the pack ice region from July 26 to September 4. The corresponding period in 2001 was July 10 through August 25, and in 2008 August 4 through September 5. A total of 2645 hours of aerosol data after the data processing (cf. Table 1).

2.3 Aerosol data from Arctic land stations

During the most recent two cruises in 2001 and 2008 sub-micrometer size distribution measurements were taken at the observatory on Mt. Zeppelin 78.9° N, 11.86° E; elevation 474 m asl) (Tunved et al., 2013). For comparison with the *Oden* data 1968 hourly average number size distributions in 20 diameter channels from 20 to 600 nm were available with concurrent five-day back trajectories.

The Dr. Neil Trivett Global Atmosphere Watch (GAW) Observatory at Alert, Nunavut (82.5° N, 75° W; elevation 210 m asl) is the only other site close to the central Arctic with comparable aerosol measurements, i.e. regular sub-micrometer particle size distribution measurements since 2011 (Leaith et al., 2013). Thus, no Alert size distributions are available during any *Oden* cruise. Instead, Alert data during the core month August of the

Oden cruises will be utilized for comparison. Specifically, we have 1517 hourly average size distributions in 54 channels between 10 and 500 nm during the Augusts of 2011, 2012, and 2013 with concurrent five-day back trajectories arriving at 250 m over Alert.

2.4 Back trajectories

Three dimensional back trajectories have been calculated for the three different receptor sites used in this study: to the icebreaker *Oden* arriving at 100 m, above sea level (a.s.l.), to the Zeppelin observatory located at the Zeppelin mountain near Ny Ålesund, Svalbard, at 474 m and to Alert at 250 m. The trajectories have been calculated backward for 10 days using the HYSPLIT2 model (Draxler and Rolph, 2003) with meteorological data provided by NCEP/NCAR project for years 1991-1996. (for more information consult <http://www.esrl.noaa.gov/psd/data/gridded/data.nmc.reanalysis.html>). For 2008 we applied the HYSPLIT4 model with GDAS data (Global Data Assimilation System). More information about the GDAS dataset can be found at Air Resources Laboratory (ARL), NOAA (<http://ready.arl.noaa.gov/>), where meteorological data also can be downloaded).

We are aware of the limitations in trajectory accuracy. On one hand the data sparse Arctic region limits the validity of the meteorological fields on which the trajectory calculations are based. On the other hand, out to the nearest continental borders the meteorological setting, surface conditions and the resulting atmospheric fields in the central Arctic are relatively simple. Figure 9 in Leck and Persson (1996b) shows an example where the trajectories were able to resolve an influence of the settlements Barentsburg and Longyearbyen on Spitsbergen in the measurements onboard *Oden* which was located near the North Pole. If we assume some 30% position uncertainty relative to the trajectory length yielding on average 3000 km for a ten-day back trajectory (cf. Stohl, 1998) this will in general not allow us to differentiate

1 between distant regions such as Beaufort Sea, Chukchi Sea, and Laptev Sea outside the pack
2 ice. A distinction between these seas and Kara Seas is however possible. The meteorological
3 information calculated along the trajectories was utilized in the analysis.

4 Instead of discussing paths of uncertain individual trajectories we plotted geographic
5 results on maps of stereographic projection centered on the North Pole. These maps were
6 covered with a coarse grid of 35 x 39 geocells, in which the passage of trajectories or the
7 occurrence of other results of this study was quantified. Fig. 2 shows that the geographical
8 region covered by the back trajectories extends to and partly beyond the pack ice limits of the
9 studied summers.

11 **2.5 Ice data**

13 Daily ice concentrations were taken from the NSIDC database (<https://nsidc.org/data>). The
14 orbits of the ice-sensing satellites excluded the area north of about 86°N. Here we assumed
15 100% ice cover. The ice data were interpolated for each hour along all back trajectories
16 because the maps of ice concentrations for the four cruises given in Fig. 3 clearly show that
17 not only did the extent of the sea vary considerably over the 17 years time of the whole data
18 set but also strongly within the study area. As integral parameters the average sum of open
19 water in percentage of each back trajectory were calculated and will be referred to: a) OS5
20 (shorter than five days before arrival at *Oden*), b) OG5 (greater than five days). From the
21 cruise-average gridded ice concentrations rough average ice limits were calculated for each
22 cruise. For that purpose contiguous lines of 10% ice concentrations north of 76° N were
23 formed and added to Fig. 2 and to maps of individual cruise years.

26 **3. Clustering approach of aerosol and trajectory data**

Many clustering approaches have been developed in exploratory data analysis (Jain et al., 1999). In atmospheric aerosol research they are used to find groups of similar aerosol data, particle origin or formation. The basic clustering algorithm of the present study has been introduced in Heintzenberg et al. (2013). Input aerosol data were pre-processed with the common Standard Normal Variate (SNV) transformation by subtracting their respective grand average and dividing them by their respective standard deviations. The same SNV transformation was applied to the trajectories after projecting them onto a stereographic map centered at the North Pole. The clustering algorithm collects the clustered data in up to nine clusters based on different input information or coordinates:

- X, y, and/or height information of the projected trajectories,
- Percentage open water along the projected trajectories, and
- Particle number size distributions.

The algorithm can utilize any combination of these three sets of clustering coordinates, i.e. the projected horizontal coordinates of the trajectories or their combination with their height coordinates and/or open water information can be clustered but also their height coordinates alone. In each case the resulting clusters of aerosol properties are calculated if available. Vice versa, aerosol properties could be clustered and for each of such clusters the resulting trajectory clusters are calculated. Finally, clusters can be sought based on aerosol, trajectory, and ice information.

The search algorithm is constrained by the four parameters N_{init} , X_{av} , P , and C_{fin} . N_{init} sets the initial minimum number of members, i.e. hourly data points that any cluster is required to have before further processing. The parameter X_{av} is defined according to

$$X_{av} = \frac{\sum_{j=1, N_j} \sum_{k=1, m} (x_{j,k} - \bar{x}_{j,k})^2}{m \cdot N}. \quad (1)$$

m is the number of coordinates to be clustered. If particle size distributions are clustered m corresponds to the number of diameters. N is the number of members in the respective cluster. $x_{j,k}$ is the coordinate k of cluster member j and $\bar{x}_{j,k}$ is the corresponding average cluster coordinate. The average distance of cluster members from a cluster average coordinate stays below a set upper limit of $X = X_{av}$. The similarity of cluster members can be improved by eliminating outliers in order of their distance from the cluster average.

Initially, the algorithm allows the input data to be segregated into a maximum of nine clusters. The algorithm will then eliminate the cluster with the maximum value of $(x_{j,k} - \bar{x}_{j,k})^2$ for any j and k until the number of cluster members is reduced to P ($P \leq 1$) times the initial number of members. Finally, the clusters will be compared to each other in order to eliminate cluster i with the minimum difference X' of average coordinates from any other cluster j

$$X' = \frac{\sum_{k=1,m} (\bar{x}_{i,k} - \bar{x}_{j,k})^2}{m} \quad (2)$$

until a given final number of clusters C_{fin} is reached. The non-sequential cluster numbering in the results discussed below reflects this elimination process, i.e. any cluster number missing in the results was eliminated in this process.

Tests of the cluster algorithm with Arctic 10-day trajectories only yielded clusters with very few members. Meteorologically this finding is easily understood: After a short time very little similarity in air pathways extending over ten days can be expected. Consequently, we limited all clustering experiments involving trajectories to five days. In Fig. 2 we see that the *Oden* cruises mainly covered the European plus western Russian sector of the inner Arctic. The trajectory coverage in Fig. 2 also shows that air from the longitudinal sector opposite to the *Oden* tracks, i.e. longitudes from about 150 to about 230 degrees partly took more than five days to reach *Oden*. Due to meteorological variability, transport pathways

from this sector to the measuring point were less similar than in other Arctic sectors and within five days the clustering algorithm could not often find many similar trajectories. Thus, in order not to miss potential source regions in this sector a conventional longitudinal sector cluster named “LC” was added to the algorithm that combined all unclustered data, the back trajectories of which had spent at least three days in this sector.

Any aerosol clustering experiment lies in between two extreme approaches. In the first one as many members as possible with somewhat similar properties are combined in each cluster, trying to cover the total data set as completely as possible with as few clusters as possible. Considering aerosol dynamics and the multitude of atmospheric processes much information will be lost in this approach. The other extreme clustering approach would attempt to be as specific as possible considering either air history and properties or aerosol properties in order reveal as much information as possible about potential aerosol source regions and formation processes. For the present study the clustering was directed towards the second extreme while trying to maintain sufficient coverage and statistical relevance in order to allow general conclusions.

For the geographic spread of the trajectories of any derived Cluster i the metric X_i is defined as

$$X_i = \sum_{j=1, N_i} \frac{1}{n_j}, \quad (3)$$

with n_j being the number of trajectory hits in any of the N_i geocells that are being crossed by trajectories of the respective cluster. The wider (and less regionally specific) the trajectory distribution of a cluster is the larger becomes X_i , and the more trajectories pass through any one cell, the narrower the spread becomes. Taken over all cruise years the 5-day back trajectories cover a total of 554 geocells. The corresponding number for 10-day trajectories is 870. Thus, with X_i the fraction of possible geocells covered by the trajectories of any cluster i can be visualized.

In subsequent maps potential source regions are identified by different colors. However, each geocell can only have only one color per map. Thus, as a measure of overlapping regions the parameter $P_{unique,i}$ is calculated as a parameter quantifying the uniqueness of the geographic area of Cluster i . $P_{unique,i}$ is the sum of the N_i geocells that are being crossed by trajectories of Cluster i but of no trajectories of any other cluster; the sum is divided by N_i and reported in percent. To sharpen this parameter only geocells that have been passed by a minimum number of trajectories (usually 25) are being counted. Assuming independent trajectory distributions 25 hits per geocells would correspond to a 25% uncertainty.

The quality of the particle size distributions in the derived clusters is described in two ways. With $x_{j,k} = dn(dlogD_p)/dlog(D_p)$ being the differential number concentration of cluster member j at diameter k and $\overline{dn}_{j,k}$ being the arithmetic cluster average of $dn_{j,k}$ the similarity of particle size distributions can be quantified for each cluster i . Additionally, in the graphical display of cluster-average size distributions the size-dependent standard deviations of the cluster averages are shown.

4. Test of the trajectory clustering with DMS

Leck and Persson (1996a, b) reported evidence for a substantial DMS source at the fringe of the central Arctic Ocean just along the MIZ, the Barents and Kara Seas being particularly strong source areas, releasing the gas to the atmosphere from the uppermost ocean. This is a result of the melting ice, which is favorable for the production of the DMS precursor dimethyl-sulfoniopropionate, released by the marine microbial food web. By using a three-dimensional numerical model Lundén et al. (2007) clearly showed that DMS(g) is advected with a photochemical turnover time to ca. 2.4 days, (Nilsson and Leck, 2002), over the pack ice in plumes originating from the source at the ice edge or in the adjacent sea just south

thereof. The above findings show that over the pack ice area, local contributions to the atmospheric DMS concentrations are negligible. At the same time DMS(g) advected from the marine source is reduced by more than an order of magnitude (Leck and Persson, 1996b).

The 5-day back trajectories (vertical dimension excluded) of all 2035 hours in all four cruises with DMS(g) data were clustered in experiment “DMS”. Run parameters of this and all other clustering experiments are listed in Table 2. Four well-separated trajectory clusters were found. An additional fifth cluster “LC” comprised the unclustered data in the longitudinal sector as defined in section 3. On average over all years the five clusters cover 22% of the DMS data. Key data of the five clusters are collected in Table 3. The regional distribution of the trajectories in the clusters is plotted in Fig. 4. Clusters 2 (red) and 3 (yellow) have the highest median (DMS)g values: 11 and 2.5 nmol m⁻³, respectively. The trajectories of these two clusters clearly point towards highly source rich ice-free areas of Greenland Sea, and Barents Sea also identified by Leck and Persson (1996a, b), and Lundén et al. (2007). The high average percentage of open water under the related trajectories, (parameter OS5 in Table 3), corroborates these results. The remaining Clusters 7, 8, and “LC” exhibited the low median DMS(g) values 0.5, 0.6, and 0.8 nmol m⁻³, respectively, together with low percentages of open water. Consequently, the test of the clustering algorithm with all available DMS(g) data has the following outcome: The potential source regions identified by the algorithm in the MIZ and adjacent open waters agree with previous DMS studies. Thus, we expect the clustering algorithm to be able to identify other potential source regions of the surface aerosol over the Arctic summer pack ice.

5. Regional distribution of potential aerosol source areas

Encouraged by the results of the test of the clustering algorithm introduced in the previous section we sought clusters of similar parameters in our total data set covering 2645 hours in four Arctic summers. The combination of horizontal trajectory information and particle size distribution was segregated into five clusters covering 25% of all hourly data in experiment “All aerosol”. The regional trajectory distributions of these clusters are plotted in Fig. 5 together with the average size distributions of the clusters. The trajectories of the five clusters cover different areas of the central Arctic and the open waters of the adjacent Arctic seas. All clusters except the longitudinal sector Cluster “LC” have geographic coverage values X_i of ≈ 20 or less. The uniqueness parameter $P_{unique,i}$ of the five clusters are 93%, 66%, 35%, 44%, and 40%, respectively, i.e. only 7% of the geocells of Cluster 1 are hit by trajectories of other clusters whereas 65% of the geocells of Cluster 4 are passed by trajectories of other clusters as well. Low-level advection of air from the open waters of the Barents Sea (OS5 = 65%) yields the typical bimodal marine size distributions, (cf. Heintzenberg et al., 2004), found in Cluster 1. Its median total number of 110 cm^{-3} is lower than the typical 250 cm^{-3} found for remote marine regions in lower latitudes (Heintzenberg et al., 2004). Cluster 3 with its potential source region over the Kara Sea has a similar bimodal shape, albeit with a much lower median total number of 37 cm^{-3} . The small tail of the average number size distribution of Cluster 3 towards the lower size limit indicates the occurrence of new particle formation in its potential source region, which is largely ice covered (OS5 = 26%). Cluster 4 stems from a potential source region north of Greenland and around the North Pole with extremely low values of open water (OS5 = 7%, with the caveat of limited satellite coverage). In its median total number of 60 cm^{-3} the accumulation mode comprises but a small shoulder. The more distant Cluster 5 is located in the pack ice covered region of the Beaufort Sea and the Canadian archipelago (OS5 = 16%). Whereas the average particle number size distribution associated with Cluster 5 is similar in shape to that in Cluster 4 the median total number in this cluster is 80 cm^{-3} with somewhat higher median concentrations of particles below 10 nm

referred to as ultrafine particles. The average particle size distribution of Cluster “LC” differs strongly from that of the other clusters: Not only is the total number about twice as high as in any of the other clusters but these high numbers also are found at smaller diameters than in any of the other clusters, i.e. largely below 30 nm. Median open water percentages below 30% in Cluster “LC” clearly demonstrate that the air masses with such high concentrations of ultrafine particles have spent long times of the pack ice. In general, median open water percentages in cluster experiment “All aerosol” differed strongly in between the clusters indicating that the amount of open water may be a controlling factor on the particle size distributions measured on *Oden*.

Besides the trajectories the ice maps yield the only system parameters that cover the whole Arctic basin. Figures 2, and 3 clearly show that both, the limits and internal variability of the Arctic pack ice varied strongly during the present study. In order to explore this potentially controlling factor we clustered the open water information along the trajectories for the total data set in the experiment “Open water” and found two groups of clusters, each with systematic differences of open water percentages in between the groups. Fig. 6a collects all trajectories of the group “Marginal ice” whereas Fig. 6b comprises the trajectory distribution of the group “Pack ice”. With some overlap in the marginal ice zone reaching from Greenland to the Laptev Sea the geographic regions of the two subpopulations are largely complementary.

In group one, named “Marginal ice”, comprising Clusters 4, 5, 7, and 8 (cf. Table 3), the median values of OG5 and OS5 were 95%, and 53%, respectively whereas the corresponding values of OG5 and OS5 were 27%, and 25%, respectively in the group two, named “Pack ice”, comprising Clusters 1, 2, 3, and 6. The clusters of the open water experiment will not be considered in detail further down. Instead a reclustering within the two groups will be discussed next.

As in experiment “All aerosol” we clustered horizontal trajectory information and particle size distributions in the subpopulation “Marginal ice”, comprising 787 hours after constraining the data input by requiring both, OG5 and OS5 being greater than 50%. 26% of the data were collected in the four clusters displayed in Figure 7 in terms of their average number size distributions together with the geographic distributions of the respective cluster trajectories. Clusters 1, and 2 exhibit typical bimodal marine size distributions as already found in clusters 1, and 3 in experiment “All aerosol” (cf. Fig. 5), albeit with more distinct potential source regions (cf. parameter P_{unique} in Table 3) over the open waters of Kara Sea and Laptev sea (Cluster 1), and North Atlantic and Barents Sea (Cluster 2). The high median DMS concentration of 2.8 nmol m^{-3} in the latter cluster reflects the highly productive open waters of the respective source region.

The trajectories of Cluster 3 come from the northern part of Greenland and the average trajectory height of 1600 m during the last five days before trajectory arrival clearly point towards a free tropospheric origin of this cluster. This character is also reflected by its average number size distribution in Fig. 7c, which is essentially monomodal with its peak around 40 nm. This monomodal distribution may be the result of very long aging of polluted air in the free troposphere (e.g., Leaitch and Isaac, 1991; Parungo et al., 1990) or may indicate new particle formation with modest growth over the Greenland ice cap.

The average number size distribution of Cluster 4 is shown in Fig. 7b and reflects another special case of input of polluted air into the pack ice region. The small diameter of 22 nm of its main peak indicates a rather fresh aerosol generated in the air mass that passed over Spitsbergen. Due to the low average trajectory travel height of some 500 m the air seemingly picked up a small accumulation mode around 150 nm. A more detailed analyses of a similar case is discussed in Bigg et al., (1996) and Leck and Persson (1996b).

Next, we discuss in more detail the subpopulation “pack ice” (cf. Fig. 6b). Clustering the open water information along the trajectories yielded the four Clusters 1, 2, 3, and 6 which

1 according to their OS5 values clearly were associated with the inner pack ice region (cf. Table
2 3). Their cluster-average size distributions and respective geographic trajectory distributions
3 are displayed in Fig. 8. The monomodal size distribution with low total numbers of Cluster 1
4 in Fig. 8a strongly reminds us of the aged aerosol in Clusters 4, and 5 in experiment “All
5 aerosol”. Cluster 1 practically covers the whole pack ice region in Fig. 6b, i.e. this type of
6 aged aerosol may appear all over the inner Arctic. While the geographic distributions of
7 Clusters 2, 3, and 6 largely are located in the same inner pack ice region their size
8 distributions in Fig. 8b look very different. Several peaks below 50 nm appear with high
9 number concentrations (up to 900 cm^{-3} at sizes down to the lower diameter limit of the
10 instruments). This type of aerosol strongly likens that of Cluster LC in the experiment “All
11 aerosol” (cf. Fig. 5).

12 Finally we explored in greater detail the large geographic region of Cluster 1 in experiment
13 “Open water” by reclustering its 636 hours of aerosol data with the information of horizontal
14 trajectories and size distributions in experiment “Pack ice low”. The results are plotted in
15 terms of average size distributions and geographic distributions of trajectories in Fig. 9.
16 Similar to Clusters 4, and 5 all clusters of the experiment “Pack ice low” have one main
17 number peak around 40 nm and a varying second mode around 100 nm which may indicate
18 some cloud processing. The similarity in size distribution while being associated with
19 different potential source regions is due to the fact that the prescribed tight ice conditions
20 occurred in different areas of the pack ice in different years (cf. Fig. 3).

23 6. Comparison with the nearest land stations

24
25 As pointed out in Section 1, 23 years after the first *Oden* expedition there are still no other
26 surface aerosol data from the central Arctic to compare with. The nearest land stations are

Mt. Zeppelin, Spitsbergen, (78.9 °N, 11.86 °E), and Alert, Nunavut, (82.5°N, 75°W). In this section the size distributions taken on *Oden* and the clusters derived with them and with back trajectories will be connected with aerosol data and trajectories from these two land stations.

6.1 Comparison *Oden* / Mt. Zeppelin

For a first comparison of particle size distributions observed at the location of the icebreaker *Oden* and at Mt. Zeppelin during the summers of 2001 and 2008, and the back trajectories to Mt. Zeppelin were employed. The closest points with distances less than 360 km between a trajectory point to the concurrent position of the icebreaker were sought along each trajectory. A total of 296 hours fulfilled this condition with an average travel time between *Oden* and Mt. Zeppelin of 36 hours and an average minimal distance between back trajectory and *Oden* of 177 km. Size distributions measured on *Oden* at the time of minimal distance were compared to size distributions measured on Mt. Zeppelin at the time of trajectory arrival. Fig. 10 gives the statistics of this comparison in terms of 25%, 50%, and 75% percentiles.

Absolute concentration levels, and the shapes of the size distributions with their main peaks roughly compare at the two points, encouraging further investigations. In all three percentiles a similar systematic change is apparent in Fig. 10. During the travel from the more central pack ice covered *Oden* area to Mt. Zeppelin concentrations decreased at all diameters larger than some 30 nm, which could be due to cloud scavenging in the marginal ice zone.

Encouraged by this statistical comparison of trajectory-connected data at the two stations cluster experiment “Oden-Zeppelin” was set up, clustering the combined particle size distribution data from the two stations. For this experiment the size distributions on *Oden* and at Mt. Zeppelin had to be harmonized. The Zeppelin data of the two largest channels (501 nm,

1 and 631 nm) were interpolated at the largest *Oden* diameter of 570 nm. All *Oden* data were
2 interpolated at the more coarsely spaced Mt. Zeppelin channels between 20 nm and 570 nm.
3 This harmonization yielded size distributions with 15 common diameter channels plus 11
4 channels from 5.1 nm to 20 nm that were only measured on *Oden*. These channels were set to
5 “missing data” at Mt. Zeppelin and were not utilized in the clustering algorithm. Averages of
6 three very different clusters of combined size distributions are shown in Fig. 11.

7 Despite our disregarding in this cluster analysis any direct trajectory connection we
8 derived quite similar cluster-average size distributions in terms of shape and absolute
9 concentrations. Because of its lower size limit the Mt. Zeppelin instrument could not detect
10 freshly formed ultrafine particles. However, the steep rise towards 20 nm in Cluster 2 of the
11 Mt. Zeppelin data in Fig. 11 is in good agreement with the right flank of the main peak about
12 15 nm that only shows up in the *Oden* data.

13 For each cluster the geographical distribution of five-day back trajectories were calculated.
14 For the *Oden* data in the clusters we utilized trajectories at 500 m arrival height in order to be
15 more compatible with the M. Zeppelin trajectories arriving at 474 m. Common potential
16 source areas were explored by plotting the average relative occurrence of trajectory points
17 only in geocells that were hit by back trajectories to both stations. Fig. 12 presents the
18 geographical distribution of jointly hit geocells for the three clusters in Fig. 11. The high
19 standard deviations of the *Oden* data in cluster 1 below 20 nm indicate the rather episodic
20 occurrence of ultrafine particles. Thus, we separated two cases of potential source areas for
21 cluster 1 in the *Oden* data, one for all cases with number concentrations below 10 nm
22 ($N_{10}=0\text{ cm}^{-3}$ measured on *Oden*, and one for all cases with $N_{10} > 1\text{ cm}^{-3}$.

23 The cases of newly formed ultrafine particles were only connected with air masses from
24 the central Arctic. Except for one geocell north of Nordaustlandet, Svalbard Cluster 2 with its
25 main peak around 15 nm also was connected with air from the central Arctic. Only cluster 4
26 had back trajectories leading out of the pack ice limit into the North Atlantic.

6.2 Comparison with Alert, Nunavut

With the cluster experiment “Oden-Alert” commonalities were sought in the shape of the size distributions measured on *Oden* and at Alert. For this exercise the data from both sites had to be harmonized in a fashion similar to the corresponding exercise with Mt. Zeppelin data. The higher resolution Alert data were interpolated at all possible diameters of the *Oden* data (11 to 435 nm). The interpolation yielded aerosol data at 34 common diameters, which could be clustered. Disregarding the fact that they were not synchronized we pooled the harmonized data from both sites into a set of 4877 hours of size distributions for the clustering. With the run parameters listed in Table 2 31% of the set were sorted into three clusters of similar shapes of size distribution. Fig. 13 presents average size distributions at both sites for these three clusters. In these cluster averages the *Oden* data extend the distributions to diameters between five and 11 nm. Clusters 1 and two have bimodal shapes albeit with the Aitken mode diameter of Cluster 1 being about 10% smaller than that of Cluster 2. Cluster 3 with highest number concentration has only one mode in the Alert size range with its peak between 40 and 50 nm.

The geographic distribution of back trajectories for the three clusters in experiment “Oden-Alert” is collected in Fig 14. Only geocells that are hit by back trajectories from both sites are marked. Additionally, two subpopulations of trajectories were formed. For Fig. 14left only data without any particles less than 10 nm measured on *Oden* were utilized. No joint geocells occurred for Cluster 2 in this subpopulation. The joint geocells for Clusters 1 and 3 cover most of the central Arctic with branches into the open water areas of the Eurasian Arctic

sectors from the Fram Strait to the Laptev Sea. In Fig 14right only cases with $N_{10} > 1\text{cm}^{-3}$ are collected. For Cluster 1, into which typical bimodal marine size distributions were sorted the geographic distribution of potential source areas did not change much in Fig. 14right. Fig. 14 indicates, however, that even in this type of air new particle formation was recorded on *Oden*. Cluster 3 with the strongest cases of new particle formation was focused onto the central Arctic when N_{10} on *Oden* was greater than 1cm^{-3} . Also, joint cells of Cluster 2 with its main mode around 30 nm appeared over the ice covered area between Greenland and the North Pole in Fig. 14right.

7. Synopsis and conclusions

The present paper continues the analysis of the aerosol data from the four summer cruises of the Swedish icebreaker *Oden* in 1991, 1996, 2001, and 2008 with a focus on potential source regions and related aerosol formation processes as illustrated in Fig. 1. While the four cruises provided a wealth of new observations there appears to be an inconsistency when comparing direct observations of a local particle flux from an open lead (Held et al., 2011) suggesting the pack ice area to be a net sink of aerosols, to statistical interpretations of aerosol concentrations (Heintzenberg and Leck, 2012), which suggests the inner most Arctic to be a source of sub-micrometer particles. Further support of the latter findings relates to the fact that near-surface airborne aerosol, as well as low-level cloud and fog droplets, contained the same type of polymer gel material as found in the open-lead surface microlayer (Bigg et al., 2004; Gao et al., 2012; Leck and Bigg, 2005; Leck et al., 2013; Orellana et al., 2011).

When comparing the course of open water under the trajectories in this study for the two aerosol types, i.e. Cluster 1, 2, 3, and 6 in the experiment “Open water” (Fig. 8) with Clusters 2, 3, and 8 of the experiment “Pack ice high” (cf. Fig. 8b) significant differences between

1 newly formed and aged aerosol over the pack ice become clear which lends further support to
2 the findings of particle sources over the inner Arctic.

3 In both subpopulations the air had spent ten days over pack ice with less than 50% open
4 water while traveling over ever more contiguous ice. Trajectories connected with high
5 concentrations of newly formed small particles, however, experienced more open water
6 during the last four days before arrival in heavy ice conditions at *Oden*. Thus we hypothesize
7 that both, long travel times over the more contiguous ice, combined with more open water
8 conditions during the last days before air mass arrival were an essential factor controlling the
9 simultaneous occurrence of high number concentrations at several discrete particle sizes in the
10 < 10 nm and 20–50 nm size ranges over the Arctic pack ice. An hypothesis fitting with this
11 chain of events could be fragmentation and/or dispersion of primary marine polymer gels,
12 200–500 nm diameter in size, into the nanogel size fractions down to a few nanometer
13 polymers (Karl et al., 2013; Leck and Bigg, 2010). Fragmentation was suggested previously
14 to be favored by evaporation of cloud or haze drops and promoted by long travel times over
15 the pack ice (e.g., Heintzenberg et al., 2006). The fragmentation hypotheses appears to be
16 consistent with the findings of a polymer gel source at the air-sea interface (Bigg et al., 2004;
17 Gao et al., 2012; Leck and Bigg, 2005; Leck et al., 2013; Orellana et al., 2011) and may also
18 explain why only a few percent of the observed total particle number variability at the ship
19 was explained by the direct measurements of particle number fluxes (Held et al., 2011).
20 Based on past and present results we conclude the inner most Arctic to be a source of sub-
21 micrometer particles.

22 Even though the Alert data had been taken in later years they still confirm the findings
23 from the other sites with respect to particle sources over the central Arctic (cf. Figs. 13, 14).
24 Also, our comparison with Spitsbergen data clearly identified similarities in the structure of
25 the size distributions and, again, pointed towards particle sources in the inner Arctic (Figs. 11,
26 12). Conventional nucleation paradigms (Karl et al., 2012) fail to explain observations of

1 small particle formation over the inner Arctic and those south of the pack ice. Previously
2 reported results from Alert in summer, (Leaith et al., 2013), and on Mt. Zeppelin,
3 Spitsbergen in spring, (Engvall et al., 2008), showed nucleation events. On Spitsbergen they
4 were followed by prototypical “banana growth” (e.g., c.f. Kulmala et al., 2001). The
5 nucleation events at both Alert and Zeppelin are explained by a conventional nucleation
6 mechanism involving solar radiation in concert with the presences of precursor gases and
7 attendant low condensational sinks. A major difference between the two land stations and the
8 inner Arctic lies in the different DMS levels. To our best knowledge (Karl et al., 2013) the
9 extremely low DMS concentrations, (Leck and Persson, 1996a, b) in the inner Arctic are not
10 sufficient for the conventional nucleation mechanism. Given that, perhaps the main
11 difference between the studies concerns how efficiently nucleation and growth of particles
12 resulting from DMS oxidation are predicted by the choice of model and lack of observations
13 to constrain the model assumptions.

14 With a clustering the open water information along the trajectories a clear separation of
15 marine versus pack ice aerosol was achieved. Then the total data set was divided into two
16 subpopulations above and below the 50% value of average open water during the course of
17 the trajectories. The two constrained data sets were investigated further for potential source
18 regions of pack ice and marine aerosols by clustering their horizontal trajectory components.
19 In the marine aerosol this clustering yielded two main source regions over Laptev and Kara
20 Seas, the aerosol showing bimodal features (cf. Fig. 7a). Beyond that two special cases
21 emerged in the marine aerosol: The first case covers polluted North Atlantic air that had
22 passed over Svalbard (cf. Fig. 7b). The second case covers free tropospheric air that had
23 crossed Greenland before arriving at *Oden* (cf. Fig. 7c).

24 The subpopulations below the 50% value of average open water during the course of the
25 trajectories indicated two different aerosol types in addition to the case of small particle
26 formation discussed above: Bimodal marine aerosol from the marginal ice zone and open seas

1 around the pack ice (cf. Figs. 5a, and 9c) and an aged aerosol that also occurred frequently
2 over the pack ice (Fig. 5c, 8a, and 9). For the former case this may involve both direct
3 emissions of larger polymer gel accumulation mode particles, as well as growth of smaller
4 particles via two processes, namely heterogeneous condensation and aerosol cloud processing
5 in which the bimodal particle size distribution characteristic of cloud-processed air is created
6 (Hoppel et al., 1994). Previous studies in the same area and season (Chang et al., 2011;
7 Heintzenberg and Leck, 2012; Heintzenberg et al., 2006; Hellén et al., 2012; Kupiszewski et
8 al., 2013; Leck and Bigg, 2005; Leck et al., 2013; Leck and Persson, 1996b; Nilsson and
9 Leck, 2002) have shown raised concentrations of accumulation mode particles within the high
10 Arctic boundary layer which the authors attribute to sources upwind *Oden*: transport of
11 precursor gases and marine biogenic particles from the MIZ or locally from open leads over
12 the pack ice. Previous reported result of individual particles by Bigg and Leck (2001, 2008),
13 Leck et al. (2002), and Leck and Bigg (2005a, b, 2010) collected over the pack ice however
14 have failed to find evidence of sea salt particles of less than 200 nm in diameter. Larger,
15 Super-micrometer particles contained a varied and appreciable organic component shown to
16 be polymer gels but also significant amounts of sodium chloride (Leck et al., 2002; Leck et
17 al., 2013).

18 The frequent occurrence of the aged aerosol (Figs. 8a, and 9) belonged to the
19 subpopulation in which the air had spent ten days over pack ice with less than 50% open
20 water while traveling over ever more contiguous ice (cf. Fig. 15) but had experienced less
21 open water during the last four days before arrival at *Oden* relative to the subpopulation
22 newly formed particles (cf. Fig. 8b). The noted relative losses of the accumulation mode can
23 be explained by an efficient scavenging processes associated with low clouds and fog near the
24 MIZ and during the first days of advection over the pack ice (Heintzenberg and Leck, 2012;
25 Nilsson and Leck, 2002). The loss in the sub-Aitken mode particle sizes would have resulted

1 from coagulation processes most efficient and thus most realistic when involving clod/fog
2 droplets (Karl et al., 2012).

3 What are the possible implications of our findings for the Arctic climate system? In the
4 course of the ongoing reduction of the summer pack ice favorable biological conditions for
5 new particle formation might increase over the Central Arctic with more frequent broken-ice
6 or open water patches. More open water increases biological activity in surface water
7 promoting the formation of biological particles. Consequently, number concentrations of
8 small particles might increase over the inner Arctic. Provided that enough condensates are
9 available, e.g., DMS oxidation products or emissions from increasing Arctic shipping, more
10 cloud condensation nuclei might result, which would affect the prevalent low clouds and fogs
11 in the summer Arctic. Changing clouds would affect the surface energy balance, which in
12 turn would have effects on ice melt.

13 14 Acknowledgements

15
16 We are most grateful to NSIDC for their providing Arctic sea ice data. In particular we are
17 indebted to Sara, Lisa, and Terry from the NSIDC user service, who helped JH with endless
18 patience to understand the formalities of NSIDC's ice data. Richard Leitch very kindly
19 processed and provided the Alert aerosol data for this study.

Literature

- E.K. Bigg, C. Leck and E.D. Nilsson, Sudden changes in Arctic atmospheric aerosol concentrations during summer and autumn, *Tellus* **48B**(1996), 254-271.
- E.K. Bigg, C. Leck and L. Tranvik, Particulates of the surface microlayer of open water in the central Arctic Ocean in summer, *Mar. Chem.* **91**(2004), 131-141.
- R.Y.W. Chang *et al.*, Aerosol composition and sources in the Central Arctic Ocean during ASCOS, *Atmos. Chem. Phys.* **11**(2011), 10619-10636.
- R. Draxler and G. Rolph, HYSPLIT (HYbrid Single-Particle Lagrangian Integrated Trajectory) Model access via NOAA ARL READY. NOAA Air Resources Laboratory, Silver Spring, MD (2003).
- A.-C. Engvall *et al.*, Changes in aerosol properties during spring-summer period in the Arctic troposphere, *Atmos. Chem. Phys.* **8**(2008), 445-462.
- Q. Gao, C. Leck, C. Rauschenberg and P.A. Matrai, On the chemical dynamics of extracellular polysaccharides in the high Arctic surface microlayer, *Ocean Sci. Discuss.* **9**(2012), 215–259.
- Q. Gao, P. Matrai and C. Leck, On the chemical dynamics of extracellular polymeric secretions (polysaccharides) in the high Arctic surface microlayer, *Mar. Chem.* **8**(2011), 401-418.
- M. Graus, M. Müller and A. Hansel, High Resolution PTR-TOF: Quantification and Formula Confirmation of VOC in Real Time, *Journal of the American Society for Mass Spectrometry* **21**(2010), 1037-1044.
- J. Heintzenberg, Size-segregated measurements of particulate elemental carbon and aerosol light absorption at remote Arctic locations., *Atmos. Environ.* **16**(1982), 2461-2469.
- J. Heintzenberg *et al.*, Mapping the aerosol over Eurasia from the Zotino Tall Tower (ZOTTO), *Tellus B* **65**(2013), doi:<http://dx.doi.org/10.3402/tellusb.v3465i3400.20062>.

- 1 J. Heintzenberg, W. Birmili, A. Wiedensohler, A. Nowak and T. Tuch, Structure, variability
2 and persistence of the submicrometer marine aerosol, *Tellus* **56B**(2004), 357-367.
- 3 J. Heintzenberg and S. Larssen, SO₂ and SO₄ in the Arctic: Interpretation of observations at
4 three Norwegian Arctic-subArctic stations, *Tellus* **35B**(1983), 255-265.
- 5 J. Heintzenberg and C. Leck, The summer aerosol in the central Arctic 1991 - 2008: did it
6 change or not?, *Atmos. Chem. Phys.* **12**(2012), 3969-3983.
- 7 J. Heintzenberg *et al.*, Aerosol number-size distributions during clear and fog periods in the
8 summer high Arctic: 1991, 1996, and 2001, *Tellus* **58B**(2006), 41-50.
- 9 A. Held, I.M. Brooks, C. Leck and M. Tjernström, On the potential contribution of open lead
10 particle emissions to the central Arctic aerosol concentration, *Atmos. Chem. Phys.*
11 **11**(2011), 3093-3105.
- 12 H. Hellén, C. Leck, J. Paatero, A. Virkkula and H. Hakola, Summer concentrations of
13 NMHCs in ambient air of the Arctic and Antarctic, *Bor. Env. Res.* **17**(2012), 385–397.
- 14 W.A. Hoppel, G.M. Frick, J.W. Fitzgerald and R.E. Larson, Marine boundary layer
15 measurements of new particle formation and the effects nonprecipitating clouds have
16 on aerosol size distribution, *J. Geophys Res.* **99**(1994), 14443-14459.
- 17 R. Jaenicke and L. Schütz, Arctic aerosols in surface air, *Idöjaras* **86**(1982), 235-241.
- 18 A.K. Jain, M.N. Murty and P.J. Flynn, Data Clustering: A Review, *ACM Comp. Surv.*
19 **31**(1999), 264–323.
- 20 M. Karl, C. Leck, E. Coz and J. Heintzenberg, Marine nanogels as a source of atmospheric
21 nanoparticles in the high Arctic, *Geophys. Res. Lett.* **40**(2013), 3738–3743.
- 22 M. Karl, C. Leck, A. Gross and L. Pirjola, A Study of New Particle Formation in the Marine
23 Boundary Layer Over the Central Arctic Ocean using a Flexible Multicomponent
24 Aerosol Dynamic Model, *Tellus* **64B**(2012),
25 doi:<http://dx.doi.org/10.3402/tellusb.v3464i3400.17158>.

- 1 A.J. Kettle *et al.*, A global database of sea surface dimethylsulfide (DMS) measurements and
2 a simple model to predict sea surface DMS as a function of latitude, longitude and
3 month, *Global Biogeochem. Cycles* **13**(1999), 399-444.
- 4 M. Kulmala *et al.*, On the formation, growth and composition of nucleation mode particles,
5 *Tellus* **53B**(2001), 479-490.
- 6 P. Kupiszewski *et al.*, Vertical profiling of aerosol particles and trace gases over the central
7 Arctic Ocean during summer, *Atmos. Chem. Phys.* **13**(2013), 12405-12431.
- 8 H. Lannefors, J. Heintzenberg and H.-C. Hansson, A comprehensive study of physical and
9 chemical parameters of the Arctic summer aerosol; results from the Swedish
10 expedition Ymer-80, *Tellus* **35B**(1983), 40-54.
- 11 W.R. Leaitch and G.A. Isaac, Tropospheric aerosol size distributions from 1982 to 1988 over
12 Eastern North America, *Atmos. Environ.* **25A**(1991), 601-619.
- 13 W.R. Leaitch *et al.*, Dimethyl sulfide control of the clean summertime Arctic aerosol and
14 cloud, *Elem. Sci. Anth.* **1**(2013), 000017.
- 15 C. Leck and E.K. Bigg, Biogenic particles in the surface microlayer and overlaying
16 atmosphere in the central Arctic Ocean during summer, *Tellus* **57B**(2005), 305–316.
- 17 C. Leck and E.K. Bigg, New particle formation of marine biological origin, *Aerosol Sci.*
18 *Technol.* **44**(2010), 570-577.
- 19 C. Leck *et al.*, Overview of the atmospheric research program during the International Arctic
20 Ocean Expedition of 1991 (IAOE-91) and its scientific results, *Tellus* **48B**(1996), 136-
21 155.
- 22 C. Leck, Q. Gao, F. Mashayekhy Rad and U. Nilsson, Size-resolved atmospheric particulate
23 polysaccharides in the high summer Arctic, *Atmos. Chem. Phys.* **13**(2013), 12573-
24 12588.

- 1 C. Leck, E.D. Nilsson, E.K. Bigg and L. Bäcklin, The atmospheric program on the Arctic
2 Ocean Expedition 1996 (AOE-96): An overview of scientific goals, experimental
3 approach, and instruments, *J. Geophys. Res.* **106**(2001), 32051-32067.
- 4 C. Leck and C. Persson, The central Arctic Ocean as a source of dimethyl sulfide: Seasonal
5 variability in relation to biological activity, *Tellus* **48B**(1996a), 156-177.
- 6 C. Leck and C. Persson, Seasonal and short-term variability in dimethyl sulfide, sulfur
7 dioxide and biogenic sulfur and sea salt aerosol particles in the arctic marine boundary
8 layer, during summer and autumn, *Tellus* **48B**(1996b), 272-299.
- 9 C. Leck, M. Tjernström, P. Matrai, E. Swietlicki and K. Bigg, Can marine micro-organisms
10 influence melting of the Arctic pack ice?, *EOS* **85**(2004), 25-36.
- 11 W. Lindinger and A. Hansel, Proton-transfer-reaction mass spectrometry (PTR-MS): On-line
12 monitoring of volatile organic compounds at ppt levels, *Chem. Soc. Rev.* **27**(1998),
13 347-354.
- 14 J. Lundén, G. Svensson and C. Leck, Influence of meteorological processes on the spatial and
15 temporal variability of atmospheric dimethyl sulfide in the high Arctic summer, *J.*
16 *Geophys. Res.* **112**(2007), D13308, doi:13310.11029/12006JD008183.
- 17 W. Maenhaut, G. Ducastel, C. Leck, E.D. Nilsson and J. Heintzenberg, Multi-elemental
18 composition and sources of the high Arctic atmospheric aerosol during summer and
19 autumn, *Tellus* **48B**(1996), 300-321.
- 20 E.D. Nilsson and C. Leck, A pseudo-Lagrangian study of the sulfur budget in the remote
21 Arctic marine boundary layer, *Tellus B* **54**(2002), 213-230.
- 22 M.V. Orellana *et al.*, Marine microgels as a source of cloud condensation nuclei in the high
23 Arctic, *PNAS* **108**(2011), 13612–13617.
- 24 F.P. Parungo, C.T. Nagamoto, P.J. Sheridan and R.C. Schnell, Aerosol characteristics of
25 Arctic haze sampled during AGASP-II, *Atmos. Environ.* **21A**(1990), 937-949.

- 1 C. Persson and C. Leck, Determination of reduced sulfur compounds in the atmosphere using
2 a cotton scrubber for oxidant removal and GC with flame photometric detection, *Anal.*
3 *Chem.* **66**(1994), 983-987.
- 4 A. Stohl, Computations, accuracy and applications of trajectories - A review and
5 bibliography, *Atmos. Environ.* **32**(1998), 947-966.
- 6 M.R. Stolzenburg, An ultrafine aerosol size distribution measuring system. *Department of*
7 *Mechanical Engineering*, University of Minnesota, Minneapolis (1988).
- 8 M. Tjernström *et al.*, The Arctic Summer Cloud-Ocean Study (ASCOS): Overview and
9 experimental design, *Atmos. Chem. Phys.* **14**(2014), 2823-2869.
- 10 P. Tunved, J. Ström and R. Krejci, Arctic aerosol life cycle: linking aerosol size distributions
11 observed between 2000 and 2010 with air mass transport and precipitation at Zeppelin
12 station, Ny-Ålesund, Svalbard, *Atmos. Chem. Phys.* **13**(2013), 3643–3660.

1

Year	Start date and time	End date and time	Hours
1991	1991-08-18 16:00	1991-09-26 23:00	768
1996	1996-07-24 19:00	1996-09-04 23:00	581
2001	2001-07-10 00:00	2001-08-25 23:00	676
2008	2008-08-04 06:00	2008-09-07 17:00	620

2

3 Table 1 Start and end date and time (UTC) of the hourly *Oden* aerosol data utilized in the
4 present paper in 1991, 1996, 2001, and 2008, and the number of hourly averages
5 after screening for possible pollution from the ship.

6

7

Experiment	Constraint	X	Y	OW	PSD	N_{init}	X_{av}	P	C_{fin}
DMS	None	x	x			48	0.2	0.75	4
All aerosol	None	x	x		x	72	0.2	0.75	4
Open water				x		24	0.2	0.9	8
Marginal ice	OG5 and OS5 > 50%	x	x		x	24	0.2	0.9	4
Pack ice low	Data from cluster 1 of “Open water”	x	x		x	48	0.7	0.5	3
Oden- Zeppelin	Concurrent time periods in 2001, 2008				x	48	0.05	0.90	3
Oden-Alert					x	48	0.05	0.90	3

Table 2 Run parameters of the cluster experiments. Constraint = Constraints on data input to clustering algorithm. OW = Percentage of open water along the trajectories. PSD = Particle size distribution. N_{init} = Initial number of hours required in each cluster. X_{av} = Average distance of the cluster members from a cluster average of normalized coordinates (cf. Eq. 1). P ($P \leq 1$) = Outlier reduction factor to be applied to each cluster (cf. Section 3). C_{fin} = Number of clusters after eliminating clusters with smallest average distance from any other cluster (cf. Section 3).

Experiment	Cluster	X_i	P_{unique}	OS5	ZAVT	NT0	N10	N26	P24	P48	P5D
DMS	2	4.7	96	69	580	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
	3	8.4	87	91	260	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
	6	14	65	50	50	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
	7	12	27	20	260	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
	LC	27	28	29	480	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
All aerosol	1	16	93	65	200	110	0	16	0	0	0.1
	3	19	66	26	440	37	0.5	6	0.3	0.4	2
	4	15	35	7	480	61	0.3	8	0	0	3
	5	21	44	16	400	81	0.1	30	0.6	1.2	5
	LC	33	40	27	690	208	30	72	0	0	5
Open water	1	21	23	10	380	100	0.4	11	0	0.1	4
	2	24	4	25	670	180	13	61	0	0	3.9
	3	33	0	29	300	150	8	63	0	0	4.3
	4	31	2	96	460	160	0.6	13	0	0	3

[Geben
Sie
Text
ein]

Experiment	Cluster	X _i	P _{unique}	OS5	ZAVT	NTO	N10	N26	P24	P48	P5D
	5	69	13	93	540	210	1	18	0	0	5
	6	20	6	25	390	130	60	64	0	0.9	1
	7	21	18	88	110	170	0	23	1	1	1
	8	52	50	96	620	90	1.9	22	0	0.1	5
Marginal ice	1	18	100	57	490	50	0	4	0	0.2	12
	2	15	97	65	240	120	0	18	0	0	0.3
	3	24	95	64	1600	130	0	22	0	0	4
	4	21	92	86	550	80	2	19	0	0	5
Pack ice low	1	8	44	6	510	50	0	7	0	0	2.7
	2	8	62	10	240	60	0	10	2.3	3.6	5
	3	10	60	12	190	40	0	5	1.3	1.9	6.6
Oden- Zeppelin	1	104	56 ¹	n.a.	890 ²	10	0	5	0	0	0
	2	38	0 ¹	n.a.	900 ²	60	2	50	0	0	0
	4	51	6 ¹	n.a.	740 ²	50	0	13	0	0	0
	LC	61	58 ¹	n.a.	1080 ²	130	1.4	30	0	0	0

Experiment	Cluster	X_i	P_{unique}	OS5	ZAVT	NTO	N10	N26	P24	P48	P5D
Oden-Alert	1	190	162 ¹	23	710	30	0	7	0	0	0
	2	70	0 ¹	29	640	110	5	70	0	0	0
	3	170	41 ¹	25	910	90	0	10	0	0	0

Table 3 Key data of the clusters of the cluster experiments. LC = Longitudinal cluster (cf. Section 3). X_i = Width of geographic coverage (vis. Eq. 4); P_{unique} = Parameter of the uniqueness of geographic coverage (%; vis. Section 3); ZAVT = Average height of trajectories during the last five days before arrival at *Oden*; OS5 = cluster-median open water (%) under the back trajectories during the last five days before arrival at *Oden*; NTO = cluster-median total particle number concentration (cm^{-3}); N10 = cluster-median particle number concentration below 10 nm diameter (cm^{-3}); N26= cluster-median particle number concentration below 26 nm diameter (cm^{-3}); P24 = Median sum of precipitation along the last 24 hours along the trajectories; P48 = Median sum of precipitation along the last 48 hours along the trajectories; P5D = Median sum of precipitation along the last five days along the trajectories; n.a. = non applicable; n.d. = no data. Aerosol and gas values for experiment Oden-Zeppelin hold for *Oden* data only. ¹ Number of cells hit jointly by trajectories to *Oden* and to Mt. Zeppelin. ² At *Oden* trajectories with 500 m arrival heights were employed.




Fig. 1 Schematic view of the sources and transport mechanisms of aerosol particles over the summer Arctic pack ice, adapted from Kupiszewski et al. (2013).




Fig. 2 Map of the working area of the present study: White: Cruise tracks during the four *Oden* expeditions in 1991, 1996, 2001, and 2008. Red symbol: North Pole. Dark grey geocells: Area covered with at least 100 trajectory hits per geocells by 5-day back trajectories in all four cruises. Additional geocells in light grey are covered likewise by 10-day back trajectories. Colored lines: Ten percent limit of sea ice cover north of 76° N estimated from average sea concentrations (<https://nsidc.org/data>) during each of the four *Oden* cruises.




Fig. 3 Gridded average Arctic sea ice cover in % during the four *Oden* cruises in 1991, 1996, 2001, and 2008. Only cells with at least 100 ice pixels per cell are plotted. The “blind spot” of satellite data north of 86 N is assumed to have 100% ice cover.




Fig. 4 Regional distribution of five clusters of back trajectories with hourly DMS values of all *Oden* cruises. Cluster 2= Red; Cluster 3= Yellow; Cluster 7= Mocha; Cluster 8= Cyan; Longitudinal cluster “LC”= Grey. The color saturation indicates the number of trajectory hits per geocells in percent. Only geocells with at least 25 trajectory hits are shown. The symbol indicates the North Pole.




Fig. 5 Left: Average particle number size distributions of the five clusters of horizontal trajectory coordinates, combined with particle size distributions. The clusters are separated into a) marine, b) pack ice high, and c) pack ice low. Cluster 1= Blue; Cluster 3= Yellow; Cluster 4= Green; Cluster 5= Magenta; Cluster “LC”= Copper. Error bars give one standard deviation about the cluster-average. Right: Corresponding regional distributions of median total number concentrations, (NTO, cm⁻³). The color saturation indicates the total number associated with the respective trajectory. Only geocells with at least 25 trajectory hits are shown. The symbol indicates the North Pole.




Fig. 6 a) Geographic distribution of trajectories of the subpopulations “Marginal ice”, and
b) geographic distribution of trajectories of all data in the subpopulation “Pack ice”.




Fig. 7 Left: Average particle number size distributions of the four clusters of horizontal trajectory coordinates, combined with particle size distributions in the subpopulation “Marginal ice”. The clusters are separated into a) marine, b) Spitsbergen and c) Greenland. Error bars give one standard deviation about the group average. Blue= Cluster 1; Yellow= Cluster 3; Green = Cluster 4. Right: Corresponding regional distributions of median total number concentrations, (NTO, cm^{-3}). The color saturation indicates the total number associated with the respective trajectory. Only geocells with at least 25 trajectory hits are shown. The symbol indicates the North Pole.




Fig. 8 Left: Average particle number size distributions of the four clusters of horizontal trajectory coordinates, combined with particle size distributions in the subpopulation “Open water”. The clusters are separated into a) Pack ice low, b) Pack ice high. Error bars give one standard deviation about the group average. Blue= Cluster 1; Yellow= Cluster 3.

Right: Corresponding regional distributions of median total number concentrations, (NTO, cm^{-3}). The color saturation indicates the total number associated with the respective trajectory. Only geocells with at least 25 trajectory hits are shown. The symbol indicates the North Pole.




Fig. 9 Left: Average particle number size distributions of the four clusters of horizontal trajectory coordinates, combined with particle size distributions in the subpopulation “Pack ice low”. Error bars give one standard deviation about the group average. Blue= Cluster 1; Red = Cluster 2; Yellow= Cluster 3.

Right: Corresponding regional distributions of median total number concentrations, (NTO, cm^{-3}). The color saturation indicates the total number associated with the respective trajectory. Only geocells with at least 25 trajectory hits are shown. The symbol indicates the North Pole.




Fig. 10 25%, 50%, and 75% percentiles of trajectory-connected number size distributions taken during the *Oden* cruises in 2001 and 2008 on the icebreaker *Oden* and on Mt. Zeppelin, Spitsbergen.




Fig. 11 Average number size distributions in three clusters of harmonized size distribution data taken on *Oden* and on Mt. Zeppelin during the *Oden* cruises in the summers of 2001 and 2008. There are no Mt. Zeppelin data below 20 nm diameter. For Cluster 1 standard deviations about the average *Oden* data are shown.




Fig. 12 Geographic distribution of back trajectories for the three clusters in Fig. 11 with joint occurrences of at least 25 trajectory hits per geocells. Left: For Cluster 1 only cases without particles less than 10 nm measured on *Oden* were considered. Right: For Cluster 1 only cases with particle concentrations less than 10 nm $>1 \text{ cm}^{-3}$ measured on *Oden* were considered. The cluster coloring corresponds to that in Fig. 11. Colored lines: Ten percent limits of sea ice cover north of 76° N estimated from average sea concentrations (<https://nsidc.org/data>) during the *Oden* cruises of 2001 and 2008.




Fig. 13 Average number size distributions in three clusters of harmonized particle size distributions measured on *Oden* during all cruises and at Alert, Nunavut during the Augusts of 2011, 2012, and 2013. There are no Alert data below 11 nm diameter.




Fig. 14 Geographic distribution of back trajectories for the three clusters in Fig. 13 with joint occurrences of at least 25 trajectory hits per geocells. Left: Only cases without particles less than 10 nm measured on *Oden* were considered, (no joint geocells for Cluster 2). Right: Only cases with particle concentrations less than $10 \text{ nm} > 1 \text{ cm}^{-3}$ measured on *Oden* were considered. The colors correspond to those in Fig. 13.




Fig. 15 Median open water percentages along the trajectories of Clusters 1-3 in experiment “Pack ice low” (cf. Fig. 9) and those of Clusters 1, 2, 3, and 6 in experiment “Open water” (cf. Fig. 8).