

Manuscript prepared for Atmos. Chem. Phys. Discuss.  
with version 2014/09/16 7.15 Copernicus papers of the  $\LaTeX$  class copernicus.cls.  
Date: 10 March 2015

# **Comment on “Observation and modelling of $\text{HO}_x$ radicals in a boreal forest” by Hens et al. (2014)**

**D. Mogensen and M. Boy**

Department of Physics, P.O. Box 48, University of Helsinki, 00014, Finland

Correspondence to: D. Mogensen (ditte.mogensen@helsinki.fi)

## 1 Correction on reference

Though as a researcher one is always happy when peers find ones work useful and accordingly cite it, it is unfortunate when the respective paper is cited incorrectly. Hens et al. (2014) write “Studies on oxidation processes in monoterpene–dominated environments are rare. Direct OH reactivity measurements in a boreal forest, conducted by Sinha et al. (2008), and a box model study investigating the OH reactivity budget (Mogensen et al., 2011) revealed a significant fraction of “unknown OH reactivity.” Firstly, the reference of Sinha et al. (2008) must be an error, since that refers to a publication about the method development of the Comparative Reactivity Method (CRM) together with first field tests conducted in the tropical rainforest of Suriname and the urban atmosphere of Mainz, but not in a boreal forest. We assume that Hens et al. (2014) instead are thinking of the paper by Sinha et al. (2010). Further, it is correct that Mogensen et al. (2011) found a large fraction of unaccounted OH reactivity, however they did not reach that conclusion using a box model. Instead they presented the first model study of the total OH reactivity in a boreal forest using a one dimensional column model with near-explicit chemistry (Boy et al., 2011; Mogensen et al., 2011). There exist earlier publications where the OH reactivity has been addressed using a 1-D model; e.g. the OH loss has been simulated with respect to certain primary emitted organic compounds (e.g., Stroud et al., 2005).

By including the dimension of the entire column of the boundary layer, we are capable of providing valuable information. E.g., as shown in Mogensen et al. (2011), the reactivity of OH has a distinct vertical profile, which is obviously not captured by a box model. Due to practical limitations, one can only measure the reactivity of OH (and other parameters of any other compound) at one or a few points in space. To our knowledge, at the time of publication of Mogensen et al. (2011), there existed no publications describing that the OH reactivity had been measured at more than one height simultaneously at one site. Mao et al. (2012) had conducted such experiments in 2009, however, the manuscript appeared later than Mogensen et al. (2011). Another exception is the aircraft measurements conducted by Mao et al. (2009), however those measurements did not capture the lowest part of the

boundary layer. Mogensen et al. (2011) therefore gave the first hints about how the space dependent canopy emission and boundary layer meteorology could affect the reactivity in a boreal forest. Hereby we got the first glimpse of how the reactivity of OH could look outside of that one measured height. Upon publication of Mogensen et al. (2011) has followed several measurement and modelling papers investigating the vertical dimension of the OH reactivity (e.g., Mao et al., 2012; Pratt et al., 2012; Hansen et al., 2014).

## References

- Boy, M., Sogachev, A., Lauros, J., Zhou, L., Guenther, A., and Smolander, S.: SOSA – a new model to simulate the concentrations of organic vapours and sulphuric acid inside the ABL – Part 1: Model description and initial evaluation, *Atmos. Chem. Phys.*, 11, 43–51, doi:10.5194/acp-11-43-2011, 2011.
- Hansen, R. F., Griffith, S. M., Dusanter, S., Rickly, P. S., Stevens, P. S., Bertman, S. B., Carroll, M. A., Erickson, M. H., Flynn, J. H., Grossberg, N., Jobson, B. T., Lefer, B. L., and Wallace, H. W.: Measurements of total hydroxyl radical reactivity during CABINEX 2009 – Part 1: field measurements, *Atmos. Chem. Phys.*, 14, 2923–2937, 10.5194/acp-14-2923-2014, 2014.
- Hens, K., Novelli, A., Martinez, M., Auld, J., Axinte, R., Bohn, B., Fischer, H., Keronen, P., Kubistin, D., Nölscher, A. C., Oswald, R., Paasonen, P., Petäjä, T., Regelin, E., Sander, R., Sinha, V., Sipilä, M., Taraborrelli, D., Tatum Ernest, C., Williams, J., Lelieveld, J., and Harder, H.: Observation and modelling of HO<sub>x</sub> radicals in a boreal forest, *Atmos. Chem. Phys.*, 14, 8723–8747, doi:10.5194/acp-14-8723-2014, 2014.
- Mao, J., Ren, X., Brune, W. H., Olson, J. R., Crawford, J. H., Fried, A., Huey, L. G., Cohen, R. C., Heikes, B., Singh, H. B., Blake, D. R., Sachse, G. W., Diskin, G. S., Hall, S. R., and Shetter, R. E.: Airborne measurement of OH reactivity during INTEX-B, *Atmos. Chem. Phys.*, 9, 163–173, doi:10.5194/acp-9-163-2009, 2009.
- Mao, J., Ren, X., Zhang, L., Van Duin, D. M., Cohen, R. C., Park, J. -H., Goldstein, A. H., Paulot, F., Beaver, M. R., Crounse, J. D., Wennberg, P. O., DiGangi, J. P., Henry, S. B., Keutsch, F. N., Park, C., Schade, G. W., Wolfe, G. M., Thornton, J. A., and Brune, W. H.: Insights into hydroxyl measurements and atmospheric oxidation in a California forest, *Atmos. Chem. Phys.*, 12, 8009–8020, doi:10.5194/acp-12-8009-2012, 2012.

- Mogensen, D., Smolander, S., Sogachev, A., Zhou, L., Sinha, V., Guenther, A., Williams, J., Nieminen, T., Kajos, M. K., Rinne, J., Kulmala, M., and Boy, M.: Modelling atmospheric OH-reactivity in a boreal forest ecosystem, *Atmos. Chem. Phys.*, 11, 9709–9719, doi:10.5194/acp-11-9709-2011, 2011.
- Pratt, K. A., Mielke, L. H., Shepson, P. B., Bryan, A. M., Steiner, A. L., Ortega, J., Daly, R., Helmig, D., Vogel, C. S., Griffith, S., Dusanter, S., Stevens, P. S., and Alaghmand, M.: Contributions of individual reactive biogenic volatile organic compounds to organic nitrates above a mixed forest, *Atmos. Chem. Phys.*, 12, 10125-10143, 10.5194/acp-12-10125-2012, 2012.
- Sinha, V., Williams, J., Crowley, J. N., and Lelieveld, J.: The Comparative Reactivity Method – a new tool to measure total OH Reactivity in ambient air, *Atmos. Chem. Phys.*, 8, 2213–2227, doi:10.5194/acp-8-2213-2008, 2008.
- Sinha, V., Williams, J., Lelieveld, J., Ruuskanen, T. M., Kajos, M. K., Patokoski, J., Hellén, H., Hakola, H., Mogensen, D., Boy, M., Rinne, J., and Kulmala, M.: OH reactivity measurements within a boreal forest: evidence for unknown reactive emissions, *Environ. Sci. Technol.*, 44, 6614–6620, 2010.
- Stroud, C., Makar, P., Karl, T., Guenther, A., Geron, C., Turnipseed, A., Nemitz, E., Baker, B., Potosnak, M., and Fuentes, J. D.: Role of canopy-scale photochemistry in modifying biogenic-atmosphere exchange of reactive terpene species: results from the CELTIC field study, *J. Geophys. Res.*, 110, D17303, doi:10.1029/2005JD005775, 2005.