

Interactive comment on “The global tropospheric ammonia distribution as seen in the 13 year AIRS measurement record” by J. X. Warner et al.

L. Yurganov

yurganov@umbc.edu

Received and published: 22 January 2016

The paper definitely should be published.

However, one point should be clarified, namely the mode of calculation of the jacobian matrix. SARTA is set on the 100 atmospheric layers with different widths of the layers in hPa (or in molecules/cm² of air partial columns). Calculation of the jacobian for a given gas requires taking these widths into account. Otherwise jacobians must depend on the mode for dividing the atmosphere on layers. If perturbations are in ppb (or ppt) and the same for all layers, then corresponding perturbations in mol/cm² would be different for different atmospheric layers: the less width in hPa, the less number of molecules for a given layer. In the lowest limit of zero width in hPa, the perturbed number of

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



molecules will be zero for any perturbation in ppb. Please, add some words, describing this effect and how do you take it into account.

Interactive comment on Atmos. Chem. Phys. Discuss., 15, 35823, 2015.

ACPD

15, C11924–C11925,
2016

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C11925

