

Short- and long-term variability of spectral solar UV irradiance at Thessaloniki, Greece: effects of changes in aerosols, total ozone and clouds

I. Fountoulakis¹, A.F. Bais¹, K. Fragkos¹, C. Meleti¹, K. Tourpali¹, M.M. Zempila^{1,2}

[1]{Aristotle University of Thessaloniki, Laboratory of Atmospheric Physics, Thessaloniki, Greece}

[2]{[Now to: Natural Resource Ecology Laboratory, Colorado State University, Fort Collins, USA](#)}

Correspondence to: I. Fountoulakis (iliasnf@auth.gr)

Abstract

In this study, we discuss the short- and the long-term variability of spectral UV irradiance at Thessaloniki, Greece using a long, quality-controlled data set from two Brewer spectrophotometers. Long-term changes in spectral UV irradiance at 307.5, 324 and 350 nm for the period 1994 – 2014 are presented for different solar zenith angles and discussed in association to changes in total ozone column (TOC), aerosol optical depth (AOD) and cloudiness observed in the same period. Positive changes in annual mean anomalies of UV irradiance, ranging from 2% to 6% per decade, have been detected both for clear- and all-sky conditions. The changes are generally greater for larger solar zenith angles and for shorter wavelengths. For clear skies, these changes are, in most cases, statistically significant at the 95% confidence limit. Decreases in the aerosol load and weakening of the attenuation by clouds lead to increases in UV irradiance in the summer, of 7 – 9% per decade for 64° solar zenith angle. The increasing TOC in winter counteracts the effect of decreasing AOD for this particular season, leading to small, statistically insignificant, negative long-term changes in irradiance at 307.5 nm. Annual mean UV irradiance levels are increasing from 1994 to 2006 and remain relatively stable thereafter, possibly due to the combined changes in the amount and optical properties of aerosols. However, no statistically significant corresponding turning

1 point has been detected in the long-term changes of AOD. ~~Trends in irradiance during the two~~
2 ~~sub-periods are not discussed, because the length of the two datasets is too short for deriving~~
3 ~~statistically significant estimates.~~ The absence of signatures of changes in AOD in the short-
4 term variability of irradiance in the UV-A may have been caused by changes in the single
5 scattering albedo of aerosols, which may counteract the effects of changes in AOD on
6 irradiance. The anti-correlation between the year-to-year variability of the irradiance at
7 307.5nm and TOC is clear and becomes clearer as the AOD decreases.

9 **1 Introduction**

10 Although ultraviolet (UV) radiation is only a small fraction (<10%) of the total solar radiation
11 that reaches the earth's surface, it is vital for the life on earth (Asta et al., 2011; ~~Hader~~Häder et
12 al., 2015; Lucas et al., 2015; Madronich et al., 2015; UNEP, 2010; Williamson et al., 2014).
13 The amount of solar UV radiation reaching the atmosphere is changing periodically due to
14 changes in the earth-sun distance and the solar activity. Solar radiation with wavelengths
15 shorter than 290nm is entirely blocked by the atmosphere, while for longer wavelengths the
16 fraction that penetrates to the surface depends mainly on the solar zenith angle, the
17 composition of the atmosphere and the characteristics of the surface (Kerr and Fioletov,
18 2008). The interaction between the UV radiation, the atmospheric constituents and the
19 characteristics of the surface is complicated and not yet fully understood (Bernhard et al.,
20 2007; Kerr and Fioletov, 2008; Meinander et al., 2009). The geophysical parameters that
21 mainly affect the levels of the surface UV irradiance are: ozone, clouds, surface reflectivity
22 and aerosols (Arola et al., 2003; Bais et al., 1993; Bernhard et al., 2007; WMO, 2007).

23 From the early 1980s until the mid-1990s, the sharp decline of the stratospheric ozone over
24 the mid and high latitudes led to important increases in the surface UV irradiance (Kerr and
25 McElroy, 1993; Madronich et al., 1998). The successful implementation of the Montreal
26 protocol decelerated the weakening of the ozone layer (Egorova et al., 2013; Mäder et al.,
27 2010; Newman and McKenzie, 2011) and many recent studies indicate the first signs of
28 recovery over the northern hemisphere (Kuttippurath et al., 2013; McLinden and Fioletov,
29 2011; Newchurch et al., 2003; Smedley et al., 2012). Signs from the onset of ozone recovery
30 since the late 1990s on surface UV-B irradiance have been mainly detected over the northern
31 high latitudes; recent studies indicate that UV-B has been declining during the last two
32 decades (Bernhard, 2011; Eleftheratos et al., 2014). However, during the same period, both

1 the UV-B and the UV-A irradiance are increasing over many locations in the northern
2 hemisphere mid-latitudes, mainly due to the negative trends in the amount of clouds and
3 aerosols (De Bock et al., 2014; Fitzka et al., 2012; Román et al., 2014; Smedley et al., 2012;
4 Zerefos et al., 2012). In a recent study, Fragkos et al. (2015) showed that in Thessaloniki,
5 even under extreme high (low) TOC conditions, the erythemal irradiance can be lower
6 (higher) than its climatological values due to the dominant effect of aerosols. Zerefos et al.
7 (2012) ~~show suggest~~ that over Canada, Europe and Japan, ~~the UV irradiance at 305 and 325~~
8 ~~nm increases by -0.9% per year during the period 1995—2006, while after 2006—~~ there is a
9 statistically significant evidence of a slowdown or even a turning point in the upward UV-B
10 trends after 2006, ~~which is an indication that since then the negative trends of aerosols are no~~
11 ~~more the main driver of the long-term changes in UV-B irradiance. , mainly due to a~~
12 ~~corresponding turning point in the negative trends of aerosols. They attribute the strong~~
13 ~~positive trends of UV B irradiance to the effect of decreasing aerosol that dominates over the~~
14 ~~effect of increasing TOC.~~

15 In the next decades, ozone, aerosols, clouds and surface reflectivity are projected to undergo
16 important changes (IPCC, 2013). Changes in these factors may alter the levels of the surface
17 UV irradiance (Bais et al., 2015; Bais et al., 2011; Hegglin and Shepherd, 2009; Tourpali et
18 al., 2009; Watanabe et al., 2011) with important impacts on the human health and the balance
19 of the ecosystems (UNEP, 2010; Williamson et al., 2014). However, the uncertainties in the
20 spatial and temporal variability, the magnitude, and the direction of the projected changes of
21 surface UV irradiance are still high (Bais et al., 2015). Thus, good quality measurements of
22 the spectral UV irradiance and the main factors controlling its levels at the earth's surface are
23 of great importance for achieving better understanding and more accurate modelling of the
24 interactions among UV radiation, ozone, aerosols, clouds and surface reflectivity (García et
25 al., 2015; Kreuter et al., 2014; Mayer and Kylling, 2005; Schwander et al., 1997).

26 Accurate knowledge of the levels of spectral surface UV irradiance is necessary in order to
27 quantify effects on the health of humans (Kazantzidis et al., 2015; Webb et al., 2010) and
28 ecosystems (Ballare et al., 2011; ~~HaderHäder~~ et al., 2011), and prevent potential impacts from
29 over- or under-exposure to UV radiation (Lucas et al., 2015). Additionally, reliable
30 estimations of the trends of spectral surface UV irradiance provide useful information for
31 assessing these impacts and for adopting proper measures (Morgenstern et al., 2008; Newman
32 and McKenzie, 2011; van Dijk et al., 2013). Climatologies and trends of surface UV

1 irradiance (spectral or broadband) can be derived either directly from ground based
2 measurements (Fitzka et al., 2012; Glandorf et al., 2005; Zerefos, 2002), or indirectly from
3 measurements of surface reflectivity, ozone, aerosols and cloudiness derived either from
4 satellites (Damiani et al., 2014; Fioletov et al., 2004; Li et al., 2000), or from ground based
5 instruments (Antón et al., 2011; Román et al., 2014; Walker, 2009). The uncertainties of these
6 parameters and the applied methodologies increase the uncertainty of the indirectly derived
7 UV irradiance, when compared to measurements (Cordero et al., 2013; Weihs and Webb,
8 1997). Thus, long records of good quality measurements of UV irradiance lead to more
9 reliable estimations of its short-term and long-term changes (Arola et al., 2003; Weatherhead
10 et al., 1998).

11 The present study aims at the quantification of the long-term changes in surface UV
12 irradiance using spectral measurements which are recorded since 1990 at Thessaloniki (Bais
13 et al., 2001; Garane et al., 2006; Gröbner et al., 2006), one of the longest time series globally
14 (Glandorf et al., 2005). An important aspect is also the attribution of the trends and variability
15 of UV irradiance to changes in the total ozone column (TOC), the aerosol optical depth
16 (AOD) and cloudiness during the same period. Special emphasis is given to the reported
17 slowdown of the [positive trends of UV irradiance \(Zerefos et al., 2012\) and their correlation](#)
18 [with the reported](#) aerosol decline over the northern hemisphere (Turnock et al., 2015; ~~Zerefos~~
19 ~~et al., 2012~~).

20

21 **2 Instrumentation and data**

22 In the 1980s, the increased concern for the stratospheric ozone depletion (Farman et al., 1985;
23 Solomon et al., 1986) and its effect on the levels of UV radiation at the Earth's surface (Kerr
24 and McElroy, 1993; Madronich et al., 1995; Zerefos, 2002), led to increased deployment of
25 ground-based instruments worldwide (Fioletov et al., 1999), to monitor the TOC and the
26 surface UV irradiance. Among these instruments, several Brewer spectrophotometers
27 (Brewer, 1973; Kerr et al., 1985) were deployed at different locations including Thessaloniki,
28 Greece, where the first commercially available single-monochromator Brewer with serial
29 number 005 (B005) was installed in 1982. Since then B005 performs continuous
30 measurements of the TOC and the columnar SO₂ (Bais et al., 1993; Bais et al., 1985; Meleti et
31 al., 2012; Zerefos, 1984). These measurements are also used to derive the aerosol optical
32 depth at specific UV-B wavelengths (Kazadzis et al., 2007). Monitoring of spectral UV

1 irradiance with B005 started in 1990 (Bais et al., 1996; Bais et al., 1993; Garane et al., 2006).
2 Since 1993, a second, double-monochromator, Brewer spectrophotometer with serial number
3 086 (B086) is also operating at Thessaloniki for continuous monitoring of the spectral UV
4 irradiance (Bais et al., 1996). Both instruments are located at the facilities of the Laboratory
5 of Atmospheric Physics (latitude 40.634° N, longitude 22.956° N, altitude 60 m above sea
6 level).

7 The spectral measurements of B005 cover the wavelength range 290-325nm in steps of 0.5
8 nm and spectral resolution of about 0.55 nm (FWHM). The corresponding spectral range for
9 B086 is 290-363nm, with the same wavelength step and very similar spectral resolution. The
10 UV dataset of both instruments was quality checked and re-evaluated up to the end of 2005
11 (Garane et al., 2006) and has been used in different studies (Kazadzis et al., 2009; Kazantzidis
12 et al., 2006; Kazantzidis et al., 2009; Meleti et al., 2009). Garane et al., (2006) presented a
13 comprehensive analysis of the uncertaintys from all known sources that may affect the
14 calibration procedures and the spectral measurements of B005 and B086. In this study the
15 overall 1σ uncertainty of the ~~measurements-measured irradiance was estimated is to about~~ 5%
16 for B086 and ~~ranges~~ from 6.5% near 305nm to 5% near 320nm for B005 ~~(Garane et al.,~~
17 ~~2006)~~. Recently, the quality control and the re-evaluation of the post-2005 dataset have been
18 completed and the time series is now extended to the end of 2014, comprising about 170.000
19 spectra for B005 and 140.000 spectra for B086.

20 Direct spectral irradiance measurements performed with B005 at 306.3, 310.0, 313.5, 316.8
21 and 320.1 nm are used to derive the TOC (Kerr et al. 1981) and the AOD (Gröbner and
22 Meleti, 2004; Meleti and Cappellani, 2000). The uncertainty of the TOC measurements is
23 estimated to about 1% or less (Kerr et al., 1985), while for the AOD the uncertainty is of the
24 order of 0.04 at 320 nm for air mass 1.4 (Kazadzis et al., 2007). Comparisons with AOD data
25 for the period 2005 – 2014 provided from a collocated Cimel sun-photometer which is part of
26 AERONET (<http://aeronet.gsfc.nasa.gov/>) revealed an overall agreement to within 0.1 for air
27 mass values up to 3.2.

28 For the trend analysis, which will be discussed later, data for the 11-year solar cycle and the
29 Quasi-Biennial Oscillation (QBO) of the winds in the equatorial stratosphere have been used.
30 Monthly means for the solar flux at 10.7 cm were downloaded from the NOAA national
31 geophysical data centre ([http://www.ngdc.noaa.gov/stp/space-weather/solar-data/solar-](http://www.ngdc.noaa.gov/stp/space-weather/solar-data/solar-features/solar-radio/noontime-flux/penticton/)
32 [features/solar-radio/noontime-flux/penticton/](http://www.ngdc.noaa.gov/stp/space-weather/solar-data/solar-features/solar-radio/noontime-flux/penticton/)), while for the QBO wind data were

1 downloaded from the Freie Universität Berlin ([http://www.geo.fu-](http://www.geo.fu-berlin.de/en/met/ag/strat/produkte/qbo/index.html)
2 [berlin.de/en/met/ag/strat/produkte/qbo/index.html](http://www.geo.fu-berlin.de/en/met/ag/strat/produkte/qbo/index.html)).

3 The spectral irradiances used in this study are averages of five measurements (within ± 1 nm
4 about the nominal wavelength) and the analysis is performed for 307.5, 324 and 350 nm. The
5 irradiance for the former two wavelengths is derived from both instruments while for 350 nm
6 it is derived only from B086. The solar irradiance at 307.5nm is strongly absorbed by ozone
7 in contrast to the other two wavelengths where the ozone absorption is either weak (324 nm)
8 or negligible (350 nm). In many studies, the irradiance at 305nm has been used to estimate the
9 effect of TOC on UVB irradiance. However, for large solar zenith angles and/or under cloudy
10 conditions the effects from the dark signal and the stray light are very important for such low
11 wavelength, especially for the single monochromator Brewers (Karppinen et al., 2014).
12 Therefore, in the present study the irradiance at 307.5 nm has been chosen because it is
13 stronger than at 305 nm, with significantly higher signal-to-noise ratio required for more
14 accurate determination of trends, while the effect of ozone absorption remains very strong.
15 Additionally, the changes in irradiance for 307.5 nm are more representative of the changes in
16 the erythemal irradiance, which is an important, human health-related metric. Finally, the
17 effect of SO₂ absorption in relation to the effect of ozone absorption at 307.5 nm is the
18 weakest in the range 306 - 309 nm. The changes in irradiance at 324 nm and 350 nm for clear
19 skies are mainly determined by the changes in the amount and optical properties of the
20 aerosols. Thus the effect of TOC on the long-term trends can be estimated by comparing the
21 trend in irradiance at 307.5 nm with the trend at these longer wavelengths. In order to detect
22 the presence of clouds during measurements and separate the data under clear skies, data from
23 a collocated pyranometer (type Kipp & Zonen CM-21) recorded at high temporal resolution
24 (~~60 Hz~~ once per minute) have been used (Bais et al., 2013).

25 The long-term variability of the spectral UV irradiance was investigated for specific solar
26 zenith angles, in order to minimize interferences from the different paths of radiation in the
27 atmosphere. Specifically, averages of measurements corresponding to SZAs within $\pm 1^\circ$ about
28 the nominal SZA were used. In order to eliminate remaining biases induced by these slightly
29 different SZAs, correction factors were derived with the radiative transfer model UVSPEC,
30 which is included in version 1.7 of the libRadtran package (Mayer and Kylling, 2005). The
31 simulations were made for a range of TOC and AOD values within the expected range of
32 variability over Thessaloniki: 250 – 550 DU for TOC and 0 – 1.4 for the AOD at 320 nm,

1 using the US standard atmospheric profile (Anderson et al., 1986), the aerosol profile
2 suggested by Shettle (1989), and typical values of the surface reflectivity, the single scattering
3 albedo and the asymmetry factor of 0.05, 0.85 and 0.7 respectively (Bais et al., 2005). The
4 simulations revealed that while for small SZAs and long wavelengths the differences in clear-
5 sky UV irradiance are small, at 305 nm the differences escalate to 60% for a change in SZA
6 from 69° to 71°. The following empirical relationship has been derived to correct the
7 measured irradiance at SZAs different than the nominal:

$$8 \quad \frac{I_0}{I_\theta} = 1 + \alpha(\lambda, \theta_0) \cdot \left(\frac{1}{\cos \theta} - \frac{1}{\cos \theta_0} \right) \quad (1)$$

9 Where θ_0 is the nominal SZA, θ is the actual SZA, I_0 is the irradiance for θ_0 , I_θ is the
10 irradiance for θ and $\alpha(\lambda, \theta_0)$ is the correction factor which depends on wavelength λ and θ_0 .
11 After applying the correction, the differences between irradiances for SZAs which differ by
12 up to 2° do not exceed 10% ($\pm 5\%$ about the mean) for wavelengths ranging from 290 to 400
13 nm and SZAs from 15° to 80°. For the wavelengths above 310 nm and SZAs smaller than 70°
14 the remaining discrepancies are generally below 2%, while for the same SZAs and
15 wavelengths between 305 and 310 nm the remaining discrepancies range between 1% and
16 5%. Thus, for 307.5 nm, the remaining uncertainties due to differences from the nominal SZA
17 range from 1 to 5% for SZAs between 30° and 70°. For the same range of SZAs the
18 corresponding uncertainties are lower than 2% and 1% for 324 and 350 nm, respectively.

19 The monthly mean values of the irradiance at 307.5 nm and at 324 nm, the TOC, and the
20 AOD derived from B005 since 1990 are presented in Figure 1. The period from June 1991
21 until December 1993 has been shaded to highlight the low TOC values due to the Mt.
22 Pinatubo volcanic eruption in June of 1991 (Hofmann et al., 1994; Randel et al., 1995). The
23 annual cycle of the irradiance at 307.5 nm is clearly anti-correlated with the annual cycle of
24 TOC while for the irradiance at 324 nm the annual variability is mainly caused by changes in
25 cloudiness. There is an indication of increasing tendency in irradiance, for both wavelengths
26 since the beginning of the record, which will be further discussed in the following sections.
27 The monthly mean TOC generally ranges between about 280 and 400 DU with no obvious
28 long-term trend. The high aerosol load over Thessaloniki is depicted in the monthly mean
29 values of AOD that range between about 0.3 and 0.9. However, during the last two decades,
30 the mean levels of the AOD are decreasing and its inter-annual variability becomes weaker.

1 As will be shown in the following, changes in aerosols play a key role in both the short- and
2 the long-term variability of the UV irradiance over Thessaloniki.

3

4 **3 Data Analysis and Results**

5 **3.1 Methodology**

6 The present study has two main objectives: First, to quantify and discuss the long- and short-
7 term changes in UV irradiance, and second, to investigate whether a turning point exists in the
8 long-term variability of the UV time-series, during the period 1994-2014. Data before 1994
9 are not used in the analysis, to eliminate the effect on total ozone from the volcanic eruption
10 of Mt. Pinatubo in June 1991. Large amounts of aerosols, mainly sulfuric, were injected into
11 the stratosphere which led to decreases in TOC in the period 1991-1993, as can be seen from
12 Figure 1. In addition, measurements of spectral UV irradiance at Thessaloniki before June
13 1991 are sparse and available for less than two years.

14 Assuming that part of the long-term variability of the clear-sky UV irradiance is due to ozone,
15 one would expect that there should be two modes in this trend: one for the period of
16 decreasing total ozone and one for the period of stabilized or increasing total ozone. For some
17 northern hemisphere locations, it was found (Zerefos et al., 2012) that surface UV-B
18 irradiance increases faster in the period before 2006. ~~If-As shown in the following~~ such a
19 turning point exists also in the 20-year long UV data-series of Thessaloniki~~-. Although~~
20 ~~quantitative estimates of the trends for each of~~ the two sub-periods ~~have been derived they are~~
21 ~~discussed briefly and in a more qualitative way, since the two time periods are would be too~~
22 short to ~~derive reliable~~ ~~consider the quantitative~~ estimates of the trends ~~reliable in each sub-~~
23 ~~period~~ (Arola et al., 2003; Weatherhead et al., 1998). Thus, in the following ~~the analysis~~ long-
24 term trends are comprehensively discussed only for the entire period 1994-2014.

25 With respect to the presence of autocorrelation in the time series of TOC and UV and its
26 effects on trend analysis, Weatherhead et al. (1998) have reported that the deseasonalized
27 monthly mean UV irradiance data are autocorrelated in the first order. For the station of
28 Thessaloniki, the autocorrelation of the all-sky dataset is estimated to be small, less than 0.2
29 ~~and should not affect significantly the magnitude of the trends~~. In contrast, for the clear-sky
30 dataset the autocorrelation is larger, ranging between 0.3 and 0.5 for different SZAs, ~~and may~~
31 ~~affect the significance of the derived trends~~. ~~In our analysis the autocorrelation has not been~~

1 | ~~removed either from the all-sky or the clear-sky datasets.~~ As discussed in Yang et al., (2006)
2 | removing the autocorrelation from time series with large number of gaps, as in our case for
3 | the clear-sky UV data set, can induce artificial tendencies and biases in the derived trends.
4 | ~~Therefore, the autocorrelation was treated differently in the calculation of the long term~~
5 | ~~changes in UV irradiance than in the detection of the turning point in its trends.~~

6 | In order to accurately detect the turning point in the trends of the UV irradiance and to
7 | determine its statistical significance, the methodology of Yang et al., (2006) was applied on
8 | the monthly mean anomalies. This study aimed at detecting a turning point in a time series of
9 | TOC with a large number of gaps using a non-autoregressive model. The additional
10 | uncertainty due to the remaining autocorrelation was taken into account in the estimation of
11 | the statistical significance associated with the detection of the turning point.

12 | For the analysis of long-term changes we calculated daily anomalies for TOC, AOD and
13 | spectral UV irradiance at different SZAs in order to remove the effect of SZA in the annual
14 | variability of irradiance. These daily anomalies were calculated by subtracting from each data
15 | point the climatological value for that day which was derived from the entire dataset. ~~Then~~
16 | ~~using the daily anomalies we derived monthly mean anomalies~~ Monthly mean anomalies were
17 | calculated by averaging the daily anomalies for months with at least 10 days of available data.

18 | As a next step, the effects of QBO and the 11-year solar cycle were filtered from both the
19 | TOC and UV irradiance data sets, by applying a multilinear regression analysis. The
20 | procedure described in, e.g., Zerefos et al., (2012) was followed with the only difference
21 | being that in the present study we did not use an autoregressive model, due to gaps in the
22 | clear-sky UV data set. It was found that the difference in the linear trends derived for the
23 | period 1994 – 2014 with and without filtering the effects of these two natural cycles is
24 | generally smaller than 0.5% per decade, thus smaller than the 1 σ uncertainty of the trends
25 | which ranges from about 1 to 3% per decade. However, it is not always negligible
26 | compared to the magnitude of the derived trends which for all cases range between about -5
27 | and 10% per decade.

28 | -The number of gaps in the time-series of the UV irradiance is higher in the first half
29 | compared to the second half of the period of study. In order to suppress the effect of the
30 | uneven distribution of the measurements, the long-term changes in UV irradiance were
31 | calculated from the yearly mean anomalies, instead of those derived from the multilinear
32 | model. After removing the effects of the solar cycle and the QBO from the monthly mean

1 anomalies, the dataset was recomposed and the yearly mean anomalies were calculated. ~~This~~
2 ~~way the problem of the autocorrelation was also solved since the yearly mean anomalies are~~
3 ~~not autocorrelated. However, i~~It was found again that the difference in the linear trends
4 derived from the yearly mean anomalies and directly from the multilinear model is small,
5 with the former being higher by only 0.1 – 0.4% per decade. The statistical significance of the
6 trends is derived from the Mann-Kendall test (Burkey, 2006). In the following, a trend is
7 considered significant when it is statistically significant at the 95% confidence level.

8 **3.2 Comparison between the trends from the two Brewers for different SZAs**

9 Following the re-evaluation and quality control of the entire dataset, the trends of the UV
10 irradiance were calculated separately for the two Brewers operating at Thessaloniki. Mean
11 ratios between quasi-synchronous (within ± 1 min) spectral UV irradiance measurements from
12 B005 and B086 under all-sky conditions ~~averaged over 5 nm spectral intervals~~ were
13 calculated for different SZA's to evaluate the applied corrections. Instrumental characteristics
14 of Brewers (e.g., the slit function) may differ for different instruments (e.g. Lakkala et al.,
15 2008) leading to differences between, even synchronous, single wavelength measurements.
16 Comparing averages over small spectral intervals suppress partly these effects. In the present
17 study averages for 5 nm spectral intervals were compared instead of irradiance measurements
18 at single wavelengths (or averages for narrower spectral intervals), because they are wide
19 enough to suppress a great part of the effect of these characteristics, while at the same time
20 they are narrow enough to assess if measurements are properly corrected for, e.g., the effects
21 of temperature and SZA with respect to wavelength. For the wavelength range 310 – 325 nm
22 the ratios are very close to 1 with a standard deviation of about 5%. For shorter wavelengths,
23 the mean ratio gradually decreases and for the 300 - 305nm range it is ~ 0.96 with a standard
24 deviation of about 10%. As already discussed, ~~T~~the uncertainties and the deviations from
25 unity arise from the different characteristics of the two instruments ~~(e.g., angular response,~~
26 ~~stray light rejection, spectral resolution and temperature)~~ and from the imperfect
27 synchronization of the measurements (Garane et al., 2006). No dependency of the ratio from
28 the temperature or the solar zenith angle was found. The good agreement in the absolute
29 levels of the measured irradiance by the two instruments is an indication for the quality of the
30 re-evaluated data. It should be noted, however, that the synchronous measurements represent
31 only $\sim 50\%$ of the available data. The trends from both instruments for the entire period 1994-
32 2014 were compared for different solar zenith angles from 30° to 70° in steps of 10° . The

1 results for 307.5 nm and 324 nm are presented in Figure 2 both for clear-sky and all-sky
2 conditions.

3 For 307.5 nm, statistically significant trends were found for clear-skies for both instruments
4 and all SZAs, and for all-skies in B005 for 30° and 70° SZA. For 324 nm, only the clear-sky
5 trends from B005 and for 30° and 40° SZA are statistically significant. ~~¶The trends for 324 nm~~
6 ~~are generally smaller than those for 307.5 nm. For this wavelength and all SZAs the all-sky~~
7 ~~trends are not statistically significant for both instruments, as they are for clear skies and for~~
8 ~~SZAs between 50° and 70°. For smaller SZAs only the trends derived from B005 data are~~
9 ~~statistically significant.~~

10 The results are quite satisfactory and consistent since for the same wavelengths and SZAs the
11 irradiance trends from B005 and B086 do not differ by more than 2% per decade and in most
12 cases they agree within 1σ . The derived trends both for clear- and all-sky data and for all
13 SZAs are positive and range between 1% and 6% per decade. Although the dependence of the
14 trends on the SZA appears to be small and within the uncertainty limits, at large SZAs the
15 trends are greater. This dependence can be partially attributed to the increasing optical path of
16 radiation with SZA, which leads to stronger absorption from ozone or aerosols. However, for
17 different SZAs, the datasets comprise data from different periods in the year (e.g. for
18 SZA=30° data exist only from April to August, while for SZA=70° data are available during
19 the entire year). Since the long-term changes of TOC, AOD and cloudiness are different for
20 different seasons, the irradiance trends for different SZAs should be affected differently by
21 these factors.

22 **3.3 Seasonal trends**

23 Since the results from both instruments are generally similar, only the data from B086 are
24 used in the following, since this instrument has superior characteristics, at least with respect to
25 the rejection of stray light and angular response. Seasonal trends of the spectral UV irradiance
26 for 307.5, 324 and 350 nm (Figure 3) were calculated and compared to the corresponding
27 trends of the daily mean TOC and AOD (Figure 4). The trends in AOD are statistically
28 significant for all cases presented-, in contrast to the trends in TOC which are not statistically
29 significant. For SZAs larger than 63° data are available during the whole year, thus, the
30 irradiance for 64° SZA (data ranging from 63° to 65°) is used in the analysis of trends. The

1 effect of the changes in cloudiness is assessed by comparing the trends of the clear-sky and
2 the all-sky irradiance.

3 As expected, the seasonal trends for 324 and the 350 nm are similar for both, clear-sky and
4 all-sky conditions. The changes of the solar irradiance at these wavelengths are practically
5 unaffected by the changes of TOC, while they are mainly affected by the changes in aerosols
6 and clouds. In general, the effects of changes in aerosol amount and/or properties on UV
7 irradiance are stronger for shorter wavelengths. Thus, the important negative trends of the
8 AOD at 320 nm that have been observed for Thessaloniki lead to slightly less positive trends
9 for the irradiance at 350 nm than at 324 nm. It must be clarified at this point that the
10 interaction of solar UV radiation with aerosols is very complex and the changes in the AOD
11 cannot explain the changes in UV irradiance without taking into account the absorption
12 efficiency of the aerosols (i.e., the single scattering albedo) for which no measurements are
13 available for this period. For example, decreases in the single scattering albedo (greater
14 absorption efficiency) counteract the effect of decreases in the AOD. As will be discussed
15 later, the fact that the changes in clear-sky UV-A irradiance (324 and 350 nm) cannot be fully
16 explained by the changes in the AOD is an indication that changes in other optical properties
17 of aerosols, such as the single scattering albedo may have occurred.

18 The greatest changes in irradiance at 324 and 350 nm were found in summer both for clear-
19 sky and all-sky conditions. The trend for clear skies at these wavelengths is about 3.5% per
20 decade, while for 307.5 nm it increases to about 5% per decade. The main driver for the
21 changes under clear skies appears to be the decreasing AOD, which for summer is more than
22 20% per decade. For all skies, the positive trends are almost double than those for clear skies
23 (about 7% for 324 and 350 nm and about 9% for 307.5 nm), suggesting that the attenuation of
24 irradiance by clouds is decreasing during the last two decades. All these trends are statistically
25 significant. For winter, the trends in irradiance for 324 and 350 nm are 3.5% and 3.0%
26 respectively both for clear skies and all-skies, suggesting that cloud effects during the last two
27 decades are very small in winter and changes in aerosols are the dominant factor. This
28 conclusion is confirmed by the negative trend of the AOD shown in Figure 4. For 307.5nm
29 the increases in TOC counteract the effects of changing aerosols, leading to a negative trend
30 of about -3% per decade for irradiance under clear skies. However, none of the trends in
31 winter is statistically significant.

1 For spring, the trends for clear skies are similar to those in summer for all three wavelengths,
2 while for all skies trends are smaller; by 0.5 - 1%. Thus, as for winter, the UV trends are due
3 mainly to decreasing AOD. Although for that season the trend in TOC is about 1% per
4 decade, this is not reflected in the trend of clear-sky irradiance at 307.5 nm which is slightly
5 larger than in the UV-A wavelengths, instead of being smaller. For this season only the trend
6 for 350 nm is statistically significant.

7 For autumn, the trends in clear-sky irradiance are approximately 7%, 3% and 1.5% for 307.5,
8 324 and 350 nm respectively, and statistically significant only for the first two wavelengths.
9 For all skies, the trends are 3-4% lower, suggesting an increasing attenuation by clouds during
10 this season. However, the differences between the clear sky and the all sky trends are within
11 the uncertainty limits of the later. The all-sky trends for autumn are not statistically
12 significant. One of the possible reasons for the stronger increase of the irradiance at 307.5 nm
13 compared to 324 and 350 nm is the small negative trend in TOC. Additionally, the relatively
14 large difference between the trends for 324 and 350 nm is explained by the decreasing
15 aerosols which have much stronger impact on shorter than on longer wavelengths.

16 Finally, the yearly averaged TOC is slightly increasing, by about 0.8% per decade, but this
17 change is not statistically significant. In contrast, the yearly mean AOD has been decreasing
18 by about 17% per decade, and therefore AOD is the dominant driver of the changes in the
19 yearly mean UV irradiance. The trends in UV irradiance range from 3 to 4% for clear skies,
20 while for all skies they are about 0.5% larger. At shorter wavelengths the trends are larger,
21 possibly due to the negative trend in TOC and the stronger effect of aerosols on the irradiance
22 at these wavelengths.

23 ~~A general conclusion is~~ The results presented in Figure 3 lead to the conclusion that the
24 enhanced attenuation of UV radiation by clouds in summer is balanced by the decreased
25 attenuation in autumn, leading to a negligible effect on the yearly mean UV irradiance.
26 However, as in all cases presented-, the differences between the trends in clear-sky and all-sky
27 irradiance for summer and autumn are similar to (or even lower than) their 1σ uncertainty,
28 which is a strong indication that the estimated changes in the attenuation of the UV irradiance
29 by clouds are not significant.

3.4 The role of ozone and aerosols on short- and long-term variability of irradiance

In the following we discuss in more detail the short- and long-term variability of clear-sky UV irradiance at 307.5 and 350 nm in association with the evolution of factors causing this variability. ~~Furthermore we explore a potential turning point in the time series of irradiance at Thessaloniki, in an attempt to confirm the findings of Zerefos et al., (2012).~~ The analysis of the variability is performed on annual mean anomalies of UV irradiance at 64° SZA, TOC and AOD, as well as for mean anomalies for the periods December – May (winter - spring) and June – November (summer - autumn); ~~since the former is being is~~ affected mainly by changes in ozone, ~~changes~~ while the ~~latter second is affected mainly~~ by changes in aerosols. Furthermore we explore a potential turning point in the time series of irradiance at Thessaloniki using monthly mean anomalies, in an attempt to confirm the findings of Zerefos et al., (2012).

The results for 324 nm are not discussed since they are similar to those for 350 nm. For ~~the monthly mean yearly mean~~ anomalies for the entire year, a turning point in the upward trend of irradiance at both 307.5 and 350 nm has been detected in 2006, statistically significant at the 95% confidence level. Since the same pattern occurs at both wavelengths and since no statistically significant turning point has been detected for TOC, this piece-wise trend pattern in UV irradiance has likely been caused by changes in aerosols. However, the small negative trend in UV irradiance observed after 2006 (see Table 1 and Figure 5) does not comply with the negative monotonic trend in AOD during the whole period of study which continues also after 2006. The behaviour of aerosols after 2006 has been verified by an independent dataset from a collocated Cimel sun-photometer, which revealed a decreasing trend of about 0.1 per decade in AOD at 440 nm from 2006 to 2014, similar to that of B005. Thus, the only factor that could explain the small negative trend in UV irradiance during this period would be a negative trend in the single scattering albedo (Bais et al., 2005; Nikitidou et al., 2013). This assumption cannot be easily verified since the SSA data from the Cimel and from satellite overpasses (e.g. <http://disc.sci.gsfc.nasa.gov/giovanni>) are sparse and inadequate to derive reliable trends. However, simulations with UVSPEC revealed that for SZAs greater than 60° and for typical aerosol properties and atmospheric conditions for Thessaloniki, the effect of a decrease in AOD at 320 nm by 0.1 can be reversed by a simultaneous decrease in SSA by less than 0.1.

1 As shown in Table 1, the trends for 350 nm for winter-spring, summer-autumn and the entire
2 year are similar. For all the three cases the UV irradiance increases by about 10% from 1994
3 to 2006 and then it slightly decreases from 2006 to 2014 resulting to a mean rate of ~~decrease~~
4 increase of about 3.5% per decade for the entire period 1994 – 2014. For ~~all~~ the three cases
5 the mean rate of decrease for the AOD is similar before and after 2006. Additionally, the year
6 to year variability of the mean anomalies for the AOD is not clearly anti-correlated with the
7 year to year variability of the mean anomalies for the UV irradiance at 350 nm, which can be
8 only attributed to changes in SSA. SSA may differ importantly for different types of aerosols
9 (Takemura et al., 2002). The aerosol mixture over Thessaloniki consists of several different
10 types of aerosols (e.g., urban, continental, marine, dust) and its composition varies (Amiridis
11 et al., 2005; Koukouli et al., 2006). This could lead to large variability of the SSA, even
12 within the same day (e.g., Ialongo et al., (2010)). An increase of the mean SSA in 1999
13 would, for example, explain why the very high annual mean levels of AOD in the specific
14 year are not depicted in the levels of UV irradiance.

15 The changes of the UV irradiance at 307.5 nm are highly affected by changes in TOC and
16 aerosols. For the winter-spring period no statistically significant turning point has been
17 detected in the trend for this wavelength. Additionally, the mean trend in irradiance for the
18 period 1994 – 2014 is weak (Table 1) compared to the corresponding trend for the period June
19 – November, and is likely caused by the combined, but opposing, effects of a statistically
20 significant positive trend in TOC and a negative trend in AOD. For the period June –
21 November no trend was detected in TOC, thus, as for 350 nm, the UV irradiance at 307.5 nm
22 increases steadily from 1994 to 2006 due to decreasing AOD and after 2006 remains
23 unchanged. A ~~similar-pattern which is similar with to this period that-pattern~~ for the period
24 June – November appears also in the annual means, with changes in irradiance dominated
25 again by changes in aerosols of opposite sign.

26 There are some interesting conclusions emerging from Figure 5: By comparing Figures 5(a) –
27 (c) with Figures 5(g) – (i), one can notice an ~~obvious~~ anti-correlation between the year to year
28 variability of TOC and the year to year variability of UV irradiance at 307.5 nm, which
29 becomes stronger as the AOD decreases. For example, the low yearly mean TOC in 2000,
30 2008 and 2011, compared in each case with the yearly mean TOC for the nearest (e.g. 4 or 5)
31 years, coincides with high UV irradiance at 307.5 nm while correspondingly the high TOC in
32 1998, 2010 and 2013 coincides with low UV irradiance at 307.5 nm. Obviously, while the

1 | ~~year to year variability long term changes~~ in irradiance at 307.5 nm ~~are is~~ mainly driven by
2 | the changes in ~~aerosol~~TOC, its ~~long-term changes year to year variability is are~~ mainly driven
3 | by the changes in ~~TOC~~aerosols. For example, the yearly mean TOC in 2010 is the highest that
4 | has been recorded during the entire period 1994 – 2014 (Steinbrecht et al., 2011) and has led
5 | to ~~extremely~~ low yearly mean irradiance at 307.5 nm. However, the yearly mean irradiance at
6 | 307.5 nm in 2010 is still higher than mean levels in the period 1994 – 1998, mainly due to the
7 | very high levels of aerosols in the atmosphere in the mid-1990s. As the AOD decreases
8 | throughout the years, the anti-correlation between the short-term variability of the TOC and
9 | the UV irradiance becomes clearer. Finally, it is noteworthy that while the mean value of
10 | AOD for 2014 in the period summer – autumn is the lowest recorded since 1994, the
11 | corresponding value for the period winter – spring is the highest of the last seven years. These
12 | very high AOD values are probably due to the increased biomass-burning aerosols arising
13 | from a shift in the type of fuel owing to the economic crisis in Greece after 2009 (Saffari et
14 | al., 2013). As a consequence of the increased aerosols, the levels of irradiance at 350 nm for
15 | winter – spring 2014 are the lowest recorded during the last decade.

16 | Since this paragraph aimed at attributing the short- and long-term variability of the UV
17 | irradiance to the corresponding variability of TOC and AOD, the analysis was restricted to
18 | clear-sky data. Although not shown here, a similar analysis has been performed for the
19 | irradiance under all-sky conditions and a statistically significant turning point in 2006 has also
20 | been detected in or the trends of yearly mean irradiance for 307.5 and 350 nm. As already
21 | discussed, changes in cloudiness do not have an important impact on the long-term changes of
22 | the UV irradiance at Thessaloniki but are the main driver of the short-term variations in the
23 | all-sky dataset.

24 |

25 | **4 Summary and conclusions**

26 | In the present study, spectral UV irradiance measurements from ~~1990–1994~~ to 2014 at
27 | Thessaloniki, Greece have been used to investigate the short- and long-term variability of UV
28 | irradiance at specific wavelengths, affected differently by total ozone, aerosols and clouds.
29 | ~~Although data are available since 1990~~ ~~the analysis was restricted to the period 1994 – 2014,~~
30 | to avoid interferences in the trends from the volcanic aerosols injected into the stratosphere by
31 | the eruption of Mt. Pinatubo in 1991. Additionally, the parallel measurements from two co-
32 | located Brewer spectrophotometers after 1993 ~~ensure data of superior quality~~ increase the

1 | confidence in the accuracy of the spectral measurements. Trends of clear-sky and all-sky UV
2 irradiance at 307.5 and 324 nm were derived for the period 1994 – 2014 from both B005 and
3 B086 for SZAs from 30° to 70°. The difference between the trends from the two instruments
4 was found to be smaller than their 1σ uncertainty boundaries.

5 According to the results, the annual mean UV irradiance has increased during the last two
6 decades. The increasing trends are similar for both clear-sky and all-sky data and are larger
7 higher at shorter wavelengths and higher SZAs. The calculated trends range between 2% and
8 6% per decade, and for clear skies are statistically significant for most SZAs. For all skies,
9 most of the irradiance trends are not statistically significant.

10 The impact of changes in TOC, aerosols and clouds on the changes in UV irradiance is
11 different for different seasons. The negative trends in AOD, which are stronger in summer,
12 lead to positive trends in UV irradiance at longer wavelengths (e.g., at 324 and 350 nm). For
13 shorter wavelengths changes in TOC are also important. Thus, the effect of the small negative
14 trend in AOD in winter is fully counteracted by the positive trend in TOC, resulting in a
15 decreasing trend in clear-sky irradiance at 307.5 nm. Changes in clouds have a negligible
16 effect on the trend of irradiance for winter and spring. The enhancement of the attenuation of
17 irradiance by clouds in autumn is balanced by the reduced attenuation in summer, leading to
18 similar changes in the annual means of clear-sky and all-sky irradiance. It is important to
19 notice that the strongest changes in UV irradiance were found for summer when humans are
20 more exposed to the Sun compared to the other seasons.

21 Moreover, it is shown that the period 1994 – 2014 can be divided in two sub-periods: during
22 the first period (1994 - 2006) the annual mean UV irradiance is increasing fast while during
23 the second period (2006 - 2014) the UV irradiance is relatively stable at 307.5 nm and is
24 slightly decreasing at 350 nm. The long-term variability of UV irradiance for both short and
25 long wavelengths is mainly driven by the changes in aerosols. The short-term variability of
26 the clear-sky irradiance at 307.5 nm is mainly driven by the short-term variability of TOC.
27 The effect of the TOC changes on the year to year variability of UV irradiance becomes
28 clearer when AOD decreases. The short-term changes in irradiance at 350 nm cannot be fully
29 explained by the short-term changes in AOD, as the absorption efficiency of aerosols may
30 also change with time.

31

References

[Amiridis, V., Balis, D. S., Kazadzis, S., Bais, A., Giannakaki, E., Papayannis, A., and Zerefos, C.: Four-year aerosol observations with a Raman lidar at Thessaloniki, Greece, in the framework of European Aerosol Research Lidar Network \(EARLINET\), Journal of Geophysical Research: Atmospheres, 110, D21, doi:10.1029/2005jd006190, 2005.](#)

Anderson, G. P., Clough, S., Kneizys, F., Chetwynd, J., and Shettle, E. P.: AFGL atmospheric constituent profiles (0-120 km), Tech. Rep. TR-86-0110, AFGL, DTIC Document, 1986.

Antón, M., Serrano, A., Cancillo, M. L., GarcÍA, J. A., and Madronich, S.: Application of an analytical formula for UV Index reconstructions for two locations in Southwestern Spain, *Tellus B*, 63, 1052-1058, 2011.

Arola, A., Lakkala, K., Bais, A., Kaurola, J., Meleti, C., and Taalas, P.: Factors affecting short- and long-term changes of spectral UV irradiance at two European stations, *J. Geophys. Res.-Atmos.*, 108, 4549, doi:10.1029/2003jd003447, 2003.

Asta, J., Pål, B., Arne, D., Stefan, A.-E., Jörg, R., Kristin, M., Michael, F. H., William, B. G., and Johan, M.: Solar radiation and human health, *Reports on Progress in Physics*, 74, 066701, doi:10.1088/0034-4885/74/6/066701, 2011.

Bais, A. F., Drosoglou, T., Meleti, C., Tourpali, K., and Kouremeti, N.: Changes in surface shortwave solar irradiance from 1993 to 2011 at Thessaloniki (Greece), *International Journal of Climatology*, 33, 2871-2876, 2013.

Bais, A. F., Gardiner, B. G., Slaper, H., Blumthaler, M., Bernhard, G., McKenzie, R., Webb, A. R., Seckmeyer, G., Kjeldstad, B., Koskela, T., Kirsch, P. J., Gröbner, J., Kerr, J. B., Kazadzis, S., Leszczynski, K., Wardle, D., Josefsson, W., Brogniez, C., Gillotay, D., Reinen, H., Weihs, P., Svenoe, T., Eriksen, P., Kuik, F., and Redondas, A.: SUSPEN intercomparison of ultraviolet spectroradiometers, *Journal of Geophysical Research: Atmospheres*, 106, 12509-12525, 2001.

Bais, A. F., Kazantzidis, A., Kazadzis, S., Balis, D. S., Zerefos, C. S., and Meleti, C.: Deriving an effective aerosol single scattering albedo from spectral surface UV irradiance measurements, *Atmospheric Environment*, 39, 1093-1102, 2005.

1 Bais, A. F., McKenzie, R. L., Bernhard, G., Aucamp, P. J., Ilyas, M., Madronich, S., and
2 Tourpali, K.: Ozone depletion and climate change: impacts on UV radiation, *Photochemical*
3 *& Photobiological Sciences*, doi: 10.1039/c4pp90032d, 2015. 2015.

4 Bais, A. F., Tourpali, K., Kazantzidis, A., Akiyoshi, H., Bekki, S., Braesicke, P.,
5 Chipperfield, M. P., Dameris, M., Eyring, V., Garny, H., Iachetti, D., Jöckel, P., Kubin, A.,
6 Langematz, U., Mancini, E., Michou, M., Morgenstern, O., Nakamura, T., Newman, P. A.,
7 Pitari, G., Plummer, D. A., Rozanov, E., Shepherd, T. G., Shibata, K., Tian, W., and
8 Yamashita, Y.: Projections of UV radiation changes in the 21st century: impact of ozone
9 recovery and cloud effects, *Atmos. Chem. Phys.*, 11, 7533-7545, 2011.

10 Bais, A. F., Zerefos, C. S., and McElroy, C. T.: Solar UVB measurements with the double-
11 and single-monochromator Brewer ozone spectrophotometers, *Geophysical Research Letters*,
12 23, 833-836, 1996.

13 Bais, A. F., Zerefos, C. S., Meleti, C., Ziomas, I. C., and Tourpali, K.: Spectral measurements
14 of solar UVB radiation and its relations to total ozone, SO₂, and clouds, *Journal of*
15 *Geophysical Research: Atmospheres*, 98, 5199-5204, 1993.

16 Bais, A. F., Zerefos, C. S., Ziomas, I. C., Zoumakis, N., Mantis, H. T., Hofmann, D. J., and
17 Fiocco, G.: Decreases in the Ozone and the SO₂ Columns Following the Appearance of the El
18 Chichon Aerosol Cloud at Midlatitude. In: *Atmospheric Ozone*, Zerefos, C. S. and Ghazi, A.
19 (Eds.), Springer Netherlands, 1985.

20 Ballare, C. L., Caldwell, M. M., Flint, S. D., Robinson, S. A., and Bornman, J. F.: Effects of
21 solar ultraviolet radiation on terrestrial ecosystems. Patterns, mechanisms, and interactions
22 with climate change, *Photochemical & Photobiological Sciences*, 10, 226-241, 2011.

23 Bernhard, G.: Trends of solar ultraviolet irradiance at Barrow, Alaska, and the effect of
24 measurement uncertainties on trend detection, *Atmos. Chem. Phys.*, 11, 13029-13045, 2011.

25 Bernhard, G., Booth, C. R., Ehramjian, J. C., Stone, R., and Dutton, E. G.: Ultraviolet and
26 visible radiation at Barrow, Alaska: Climatology and influencing factors on the basis of
27 version 2 National Science Foundation network data, *J. Geophys. Res.-Atmos.*, 112, D09101,
28 doi:10.1029/2006jd007865, 2007.

29 Brewer, A. W.: A replacement for the Dobson spectrophotometer?, *PAGEOPH*, 106-108,
30 919-927, 1973.

1 Burkey, J.: A non-parametric monotonic trend test computing Mann-Kendall Tau, Tau-b, and
2 Sens Slope written in Mathworks-MATLAB implemented using matrix rotations., King
3 County, Department of Natural Resources and Parks, Science and Technical Services section.
4 Seattle, Washington. USA., 2006.

5 Cordero, R. R., Seckmeyer, G., Damiani, A., Labbe, F., and Laroze, D.: Monte Carlo-based
6 uncertainties of surface UV estimates from models and from spectroradiometers, *Metrologia*,
7 50, L1, 2013.

8 Damiani, A., Cordero, R. R., Cabrera, S., Laurenza, M., and Rafanelli, C.: Cloud cover and
9 UV index estimates in Chile from satellite-derived and ground-based data, *Atmospheric*
10 *Research*, 138, 139-151, 2014.

11 De Bock, V., De Backer, H., Van Malderen, R., Mangold, A., and Delcloo, A.: Relations
12 between erythemal UV dose, global solar radiation, total ozone column and aerosol optical
13 depth at Uccle, Belgium, *Atmos. Chem. Phys.*, 14, 12251-12270, 2014.

14 Egorova, T., Rozanov, E., Gröbner, J., Hauser, M., and Schmutz, W.: Montreal Protocol
15 Benefits simulated with CCM SOCOL, *Atmos. Chem. Phys.*, 13, 3811-3823, 2013.

16 Eleftheratos, K., Kazadzis, S., Zerefos, C. S., Tourpali, K., Meleti, C., Balis, D., Zyrichidou,
17 I., Lakkala, K., Feister, U., Koskela, T., Heikkilä, A., and Karhu, J. M.: Ozone and
18 Spectroradiometric UV Changes in the Past 20 Years over High Latitudes, *Atmosphere-*
19 *Ocean*, doi: 10.1080/07055900.2014.919897, 2014. 1-9, 2014.

20 Farman, J. C., Gardiner, B. G., and Shanklin, J. D.: Large losses of total ozone in Antarctica
21 reveal seasonal ClO_x/NO_x interaction, *Nature*, 315, 207-210, 1985.

22 Fioletov, V. E., Kerr, J. B., Hare, E. W., Labow, G. J., and McPeters, R. D.: An assessment of
23 the world ground-based total ozone network performance from the comparison with satellite
24 data, *Journal of Geophysical Research: Atmospheres*, 104, 1737-1747, 1999.

25 Fioletov, V. E., Kimlin, M. G., Krotkov, N., McArthur, L. J. B., Kerr, J. B., Wardle, D. I.,
26 Herman, J. R., Meltzer, R., Mathews, T. W., and Kaurola, J.: UV index climatology over the
27 United States and Canada from ground-based and satellite estimates, *J. Geophys. Res.-*
28 *Atmos.*, 109, D22308, doi:10.1029/2004jd004820, 2004.

29 Fitzka, M., Simic, S., and Hadzimustafic, J.: Trends in spectral UV radiation from long-term
30 measurements at Hoher Sonnblick, Austria, *Theor. Appl. Climatol.*, 110, 585-593, 2012.

1 Garane, K., Bais, A. F., Kazadzis, S., Kazantzidis, A., and Meleti, C.: Monitoring of UV
2 spectral irradiance at Thessaloniki (1990–2005): data re-evaluation and quality control, *Ann.*
3 *Geophys.*, 24, 3215–3228, 2006.

4 García, R. D., Cachorro, V. E., Cuevas, E., Toledano, C., Redondas, A., Blumthaler, M., and
5 Benounna, Y.: Comparison of measured and modelled spectral UV irradiance at Izaña high
6 mountain station: estimation of the underlying effective albedo, *International Journal of*
7 *Climatology*, doi: 10.1002/joc.4355, 2015. n/a-n/a, 2015.

8 Glandorf, M., Arola, A., Bais, A., and Seckmeyer, G.: Possibilities to detect trends in spectral
9 UV irradiance, *Theor. Appl. Climatol.*, 81, 33-44, 2005.

10 Gröbner, J., Blumthaler, M., Kazadzis, S., Bais, A., Webb, A., Schreder, J., Seckmeyer, G.,
11 and Rembges, D.: Quality assurance of spectral solar UV measurements: results from 25 UV
12 monitoring sites in Europe, 2002 to 2004, *Metrologia*, 43, S66–S71, doi:10.1088/0026- 20
13 1394/43/2/S14, 2006.

14 Gröbner, J. and Meleti, C.: Aerosol optical depth in the UVB and visible wavelength range
15 from Brewer spectrophotometer direct irradiance measurements: 1991–2002, *J. Geophys.*
16 *Res.- Atmos.*, 109, D09202, doi:10.1029/2003jd004409, 2004.

17 ~~Hader~~Häder, D.-P., Williamson, C. E., Wangberg, S.-A., Rautio, M., Rose, K. C., Gao, K.,
18 Helbling, E. W., Sinha, R. P., and Worrest, R.: Effects of UV radiation on aquatic ecosystems
19 and interactions with other environmental factors, *Photochemical & Photobiological Sciences*,
20 doi: 10.1039/c4pp90035a, 2015. 2015.

21 ~~Hader~~Häder, D. P., Helbling, E. W., Williamson, C. E., and Worrest, R. C.: Effects of UV
22 radiation on aquatic ecosystems and interactions with climate change, *Photochemical &*
23 *Photobiological Sciences*, 10, 242-260, 2011.

24 Hegglin, M. I. and Shepherd, T. G.: Large climate-induced changes in ultraviolet index and
25 stratosphere-to-troposphere ozone flux, *Nature Geoscience*, 2, 687-691, 2009.

26 Hofmann, D. J., Oltmans, S. J., Komhyr, W. D., Harris, J. M., Lathrop, J. A., Langford, A. O.,
27 Deshler, T., Johnson, B. J., Torres, A., and Matthews, W. A.: ozone loss in the lower
28 stratosphere over the United States in 1992–1993: Evidence for heterogeneous chemistry on
29 the Pinatubo aerosol, *Geophysical Research Letters*, 21, 65-68, 1994.

1 [Ialongo, I., Buchard, V., Brogniez, C., Casale, G. R., and Siani, A. M.: Aerosol Single](#)
2 [Scattering Albedo retrieval in the UV range: an application to OMI satellite validation,](#)
3 [Atmos. Chem. Phys., 10, 331-340, 2010.](#)

4 IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I
5 to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change Cambridge
6 University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp., 2013.

7 Karppinen, T., Redondas, A., García, R. D., Lakkala, K., McElroy, C. T., and Kyrö, E.:
8 Compensating for the Effects of Stray Light in Single-Monochromator Brewer
9 Spectrophotometer Ozone Retrieval, Atmosphere-Ocean, doi:
10 10.1080/07055900.2013.871499, 2014. 1-8, 2014.

11 Kazadzis, S., Bais, A., Amiridis, V., Balis, D., Meleti, C., Kouremeti, N., Zerefos, C. S.,
12 Rapsomanikis, S., Petrakakis, M., Kelesis, A., Tzoumaka, P., and Kelektsoğlu, K.: Nine
13 years of UV aerosol optical depth measurements at Thessaloniki, Greece, Atmos. Chem.
14 Phys., 7, 2091-2101, 2007.

15 Kazadzis, S., Bais, A., Arola, A., Krotkov, N., Kouremeti, N., and Meleti, C.: Ozone
16 Monitoring Instrument spectral UV irradiance products: comparison with ground based
17 measurements at an urban environment, Atmos. Chem. Phys., 9, 585-594, 2009.

18 Kazantzidis, A., Bais, A., Garane, K., Kazadzis, S., and Meleti, C.: Estimation of UV
19 irradiance from ancillary data and comparison with measurements at Thessaloniki, Greece
20 (40.5° N, 23° E), 6362, doi:10.1117/12.689813, 2006.

21 Kazantzidis, A., Bais, A. F., Zempila, M. M., Kazadzis, S., den Outer, P. N., Koskela, T., and
22 Slaper, H.: Calculations of the human vitamin D exposure from UV spectral measurements at
23 three European stations, Photochemical & Photobiological Sciences, 8, 45-51, 2009.

24 Kazantzidis, A., Smedley, A., Kift, R., Rimmer, J., Berry, J., Rhodes, L. E., and Webb, A.: A
25 modeling approach to determine how much UV radiation is available across the UK and
26 Ireland for health risk and benefit studies, Photochemical & Photobiological Sciences, doi:
27 10.1039/c5pp00008d, 2015. 2015.

28 Kerr, J. B., Evans, W. F. J., and Asbridge, I. A.: Recalibration of Dobson Field
29 Spectrophotometers with a Travelling Brewer Spectrophotometer Standard. In: Atmospheric
30 Ozone, Zerefos, C. S. and Ghazi, A. (Eds.), Springer Netherlands, 1985.

1 Kerr, J. B. and Fioletov, V. E.: Surface ultraviolet radiation, *Atmos. Ocean*, 46, 159-184,
2 2008.

3 Kerr, J. B. and McElroy, C. T.: Evidence for Large Upward Trends of Ultraviolet-B Radiation
4 Linked to Ozone Depletion, *Science*, 262, 1032-1034, 1993.

5 [Koukouli, M. E., Balis, D. S., Amiridis, V., Kazadzis, S., Bais, A., Nickovic, S., and Torres,](#)
6 [O.: Aerosol variability over Thessaloniki using ground based remote sensing observations and](#)
7 [the TOMS aerosol index, *Atmospheric Environment*, 40, 5367-5378, 2006.](#)

8 Kreuter, A., Buras, R., Mayer, B., Webb, A., Kift, R., Bais, A., Kouremeti, N., and
9 Blumthaler, M.: Solar irradiance in the heterogeneous albedo environment of the Arctic coast:
10 measurements and a 3-D model study, *Atmos. Chem. Phys.*, 14, 5989-6002, 2014.

11 Kuttippurath, J., Lefèvre, F., Pommereau, J. P., Roscoe, H. K., Goutail, F., Pazmiño, A., and
12 Shanklin, J. D.: Antarctic ozone loss in 1979–2010: first sign of ozone recovery,
13 *Atmos. Chem. Phys.*, 13, 1625-1635, 2013.

14 [Lakkala, K., Arola, A., Heikkilä, A., Kaurola, J., Koskela, T., Kyrö, E., Lindfors, A.,](#)
15 [Meinander, O., Tanskanen, A., Gröbner, J., and Hülsen, G.: Quality assurance of the Brewer](#)
16 [spectral UV measurements in Finland, *Atmos. Chem. Phys.*, 8, 3369-3383, 10.5194/acp-8-](#)
17 [3369-2008, 2008.](#)

18 Li, Z., Wang, P., and Cihlar, J.: A simple and efficient method for retrieving surface UV
19 radiation dose rate from satellite, *Journal of Geophysical Research: Atmospheres*, 105, 5027-
20 5036, 2000.

21 Lucas, R. M., Norval, M., Neale, R. E., Young, A. R., de Gruijl, F. R., Takizawa, Y., and van
22 der Leun, J. C.: The consequences for human health of stratospheric ozone depletion in
23 association with other environmental factors, *Photochemical & Photobiological Sciences*, doi:
24 10.1039/c4pp90033b, 2015. 2015.

25 Mäder, J. A., Staehelin, J., Peter, T., Brunner, D., Rieder, H. E., and Stahel, W. A.: Evidence
26 for the effectiveness of the Montreal Protocol to protect the ozone layer, *Atmos. Chem. Phys.*,
27 10, 12161-12171, 2010.

28 Madronich, S., McKenzie, R. L., Björn, L. O., and Caldwell, M. M.: Changes in biologically
29 active ultraviolet radiation reaching the Earth's surface, *Journal of Photochemistry and*
30 *Photobiology B: Biology*, 46, 5-19, 1998.

1 Madronich, S., McKenzie, R. L., and Caldwell, M. M.: Changes in ultraviolet radiation
2 reaching the earth's surface, *Ambio*, 24, 143-152, 1995.

3 Madronich, S., Shao, M., Wilson, S. R., Solomon, K. R., Longstreth, J. D., and Tang, X. Y.:
4 Changes in air quality and tropospheric composition due to depletion of stratospheric ozone
5 and interactions with changing climate: implications for human and environmental health,
6 *Photochemical & Photobiological Sciences*, doi: 10.1039/c4pp90037e, 2015. 2015.

7 Mayer, B. and Kylling, A.: Technical note: The libRadtran software package for radiative
8 transfer calculations - description and examples of use, *Atmos. Chem. Phys.*, 5, 1855-1877,
9 2005.

10 McLinden, C. A. and Fioletov, V.: Quantifying stratospheric ozone trends: Complications due
11 to stratospheric cooling, *Geophysical Research Letters*, 38, L03808, 2011.

12 Meinander, O., Wuttke, S., Seckmeyer, G., Kazadzis, S., Lindfors, A., and Kyrö, E.: Solar
13 zenith angle asymmetry cases in polar snow UV albedo, *Geophysica*, 45, 183-198, 2009.

14 Meleti, C., Bais, A. F., Kazadzis, S., Kouremeti, N., Garane, K., and Zerefos, C.: Factors
15 affecting solar ultraviolet irradiance measured since 1990 at Thessaloniki, Greece, *Int. J. Rem.*
16 *Sens.*, 30, 4167-4179, 2009.

17 Meleti, C. and Cappellani, F.: Measurements of aerosol optical depth at Ispra: Analysis of the
18 correlation with UV-B, UV-A, and total solar irradiance, *Journal of Geophysical Research:*
19 *Atmospheres*, 105, 4971-4978, 2000.

20 Meleti, C., Fragkos, K., Bais, A. F., Tourpali, K., Balis, D., and Zerefos, C. S.: Thirty years of
21 total ozone measurements at Thessaloniki with a MKII Brewer spectrophotometer,
22 *Quadrennial Ozone Symposium 2012*, Toronto, 2012.

23 Morgenstern, O., Braesicke, P., Hurwitz, M. M., O'Connor, F. M., Bushell, A. C., Johnson, C.
24 E., and Pyle, J. A.: The World Avoided by the Montreal Protocol, *Geophys. Res. Lett.*, 35,
25 L16811, doi:10.1029/2008gl034590, 2008.

26 Newchurch, M. J., Yang, E.-S., Cunnold, D. M., Reinsel, G. C., Zawodny, J. M., and Russell,
27 J. M.: Evidence for slowdown in stratospheric ozone loss: First stage of ozone recovery, *J.*
28 *Geophys. Res.-Atmos.*, 108, 4507, doi:10.1029/2003jd003471, 2003.

29 Newman, P. A. and McKenzie, R.: UV impacts avoided by the Montreal Protocol,
30 *Photochemical & Photobiological Sciences*, 10, 1152-1160, 2011.

1 Nikitidou, E., Kazantzidis, A., De Bock, V., and De Backer, H.: The aerosol forcing
2 efficiency in the UV region and the estimation of single scattering albedo at a typical West
3 European site, *Atmospheric Environment*, 69, 313-320, 2013.

4 Randel, W. J., Wu, F., Russell, J. M., Waters, J. W., and Froidevaux, L.: Ozone and
5 temperature changes in the stratosphere following the eruption of Mount Pinatubo, *Journal of*
6 *Geophysical Research: Atmospheres*, 100, 16753-16764, 1995.

7 Román, R., Bilbao, J., and de Miguel, A.: Erythemat ultraviolet irradiation trends in the
8 Iberian Peninsula from 1950 to 2011, *Atmos. Chem. Phys. Discuss.*, 14, 15545-15590, 2014.

9 Saffari, A., Daher, N., Samara, C., Voutsas, D., Kouras, A., Manoli, E., Karagkiozidou, O.,
10 Vlachokostas, C., Moussiopoulos, N., Shafer, M. M., Schauer, J. J., and Sioutas, C.: Increased
11 Biomass Burning Due to the Economic Crisis in Greece and Its Adverse Impact on
12 Wintertime Air Quality in Thessaloniki, *Environmental Science & Technology*, 47, 13313-
13 13320, 2013.

14 Schwander, H., Koepke, P., and Ruggaber, A.: Uncertainties in modeled UV irradiances due
15 to limited accuracy and availability of input data, *Journal of Geophysical Research:*
16 *Atmospheres*, 102, 9419-9429, 1997.

17 Shettle, E. P.: Models of aerosols, clouds and precipitation for atmospheric propagation
18 studies., In *AGARD, Atmospheric Propagation in the UV, Visible, IR, and MM-Wave Region*
19 *and Related Systems Aspects* 14 p, 1989. 1989.

20 Smedley, A. R. D., Rimmer, J. S., Moore, D., Toumi, R., and Webb, A. R.: Total ozone and
21 surface UV trends in the United Kingdom: 1979–2008, *International Journal of Climatology*,
22 32, 338-346, 2012.

23 Solomon, S., Garcia, R. R., Rowland, F. S., and Wuebbles, D. J.: On the depletion of
24 Antarctic ozone, *Nature*, 321, 755-758, 1986.

25 Steinbrecht, W., Köhler, U., Claude, H., Weber, M., Burrows, J. P., and van der A, R. J.: Very
26 high ozone columns at northern mid-latitudes in 2010, *Geophys. Res. Lett.*, 38, 1944-8007,
27 doi:10.1029/2010GL046634, 2011.

28 [Takemura, T., Nakajima, T., Dubovik, O., Holben, B. N., and Kinne, S.: Single-Scattering](#)
29 [Albedo and Radiative Forcing of Various Aerosol Species with a Global Three-Dimensional](#)
30 [Model, *Journal of Climate*, 15, 333-352, 2002.](#)

1 Tourpali, K., Bais, A. F., Kazantzidis, A., Zerefos, C. S., Akiyoshi, H., Austin, J., Brühl, C.,
2 Butchart, N., Chipperfield, M. P., Dameris, M., Deushi, M., Eyring, V., Giorgetta, M. A.,
3 Kinnison, D. E., Mancini, E., Marsh, D. R., Nagashima, T., Pitari, G., Plummer, D. A.,
4 Rozanov, E., Shibata, K., and Tian, W.: Clear sky UV simulations for the 21st century based
5 on ozone and temperature projections from Chemistry-Climate Models, *Atmos. Chem. Phys.*,
6 9, 1165-1172, 2009.

7 Turnock, S. T., Spracklen, D. V., Carslaw, K. S., Mann, G. W., Woodhouse, M. T., Forster, P.
8 M., Haywood, J., Johnson, C. E., Dalvi, M., Bellouin, N., and Sanchez-Lorenzo, A.: Modelled
9 and observed changes in aerosols and surface solar radiation over Europe between 1960 and
10 2009, *Atmos. Chem. Phys. Discuss.*, 15, 13457-13513, 2015.

11 UNEP: Environmental effects of ozone depletion and its interaction with climate change:
12 2010 assessment, report, 278 pp, Nairobi, Kenya., 2010.

13 van Dijk, A., Slaper, H., den Outer, P. N., Morgenstern, O., Braesicke, P., Pyle, J. A., Garny,
14 H., Stenke, A., Dameris, M., Kazantzidis, A., Tourpali, K., and Bais, A. F.: Skin Cancer Risks
15 Avoided by the Montreal Protocol—Worldwide Modeling Integrating Coupled Climate-
16 Chemistry Models with a Risk Model for UV, Photochemistry and Photobiology, 89, 234-
17 246, 2013.

18 Walker, D.: Cloud effects on erythemal UV radiation in a complex topography, doi:
19 <http://dx.doi.org/10.3929/ethz-a-005914035>, 2009. Diss., Eidgenössische Technische
20 Hochschule ETH Zürich, Nr. 18415, 2009, 2009.

21 Watanabe, S., Sudo, K., Nagashima, T., Takemura, T., Kawase, H., and Nozawa, T.: Future
22 projections of surface UV-B in a changing climate, *J. Geophys. Res.-Atmos.*, 116, D16118,
23 doi:10.1029/2011jd015749, 2011.

24 Weatherhead, E. C., Reinsel, G. C., Tiao, G. C., Meng, X.-L., Choi, D., Cheang, W.-K.,
25 Keller, T., DeLuisi, J., Wuebbles, D. J., Kerr, J. B., Miller, A. J., Oltmans, S. J., and
26 Frederick, J. E.: Factors affecting the detection of trends: Statistical considerations and
27 applications to environmental data, *Journal of Geophysical Research: Atmospheres*, 103,
28 17149-17161, 1998.

29 Webb, A. R., Kift, R., Durkin, M. T., O'Brien, S. J., Vail, A., Berry, J. L., and Rhodes, L. E.:
30 The role of sunlight exposure in determining the vitamin D status of the U.K. white adult
31 population, *British Journal of Dermatology*, 163, 1050-1055, 2010.

1 Weihs, P. and Webb, A. R.: Accuracy of spectral UV model calculations: 1. Consideration of
2 uncertainties in input parameters, *Journal of Geophysical Research: Atmospheres*, 102, 1541-
3 1550, 1997.

4 Williamson, C. E., Zepp, R. G., Lucas, R. M., Madronich, S., Austin, A. T., Ballare, C. L.,
5 Norval, M., Sulzberger, B., Bais, A. F., McKenzie, R. L., Robinson, S. A., ~~Hader~~Häder, D.-P.,
6 Paul, N. D., and Bornman, J. F.: Solar ultraviolet radiation in a changing climate, *Nature*
7 *Clim. Change*, 4, 434-441, 2014.

8 WMO: Scientific assessment of ozone depletion: 2006, *Global Ozone Res. Monit. Proj. Rep.*
9 50, 572 pp., 2007.

10 Yang, E.-S., Cunnold, D. M., Salawitch, R. J., McCormick, M. P., Russell, J., Zawodny, J.
11 M., Oltmans, S., and Newchurch, M. J.: Attribution of recovery in lower-stratospheric ozone,
12 *J. Geophys. Res.-Atmos.*, 111, D17309, doi:10.1029/2005jd006371, 2006.

13 Zerefos, C.: Evidence of the El Chichón stratospheric volcanic cloud in Northern Greece,
14 *Geofísica Internacional*, 23, 299–304, 1984.

15 Zerefos, C. S.: Long-term ozone and UV variations at Thessaloniki, Greece, *Physics and*
16 *Chemistry of the Earth, Parts A/B/C*, 27, 455-460, 2002.


17 Zerefos, C. S., Tourpali, K., Eleftheratos, K., Kazadzis, S., Meleti, C., Feister, U., Koskela,
18 T., and Heikkilä, A.: Evidence of a possible turning point in solar UV-B over Canada, Europe
19 and Japan, *Atmos. Chem. Phys.*, 12, 2469-2477, 2012.

20

1 Table 1. Trends of TOC, AOD at 320 nm and spectral UV irradiance at 307.5 and 350 nm, for
 2 different periods. Asterisks denote the statistically significant trends


	period	WINTER-SPRING	SUMMER-AUTUMN	YEAR
307.5nm (change % per decade)	1994 – 2006	-	11.0 ± 3.3 *	7.1 ± 2.1 *
	2006 - 2014	-	-0.16 ± 7.7	-0.28 ± 5.0
	1994 – 2014	2.2 ± 1.9	7.0 ± 1.9 *	4.5 ± 1.2 *
350nm (change % per decade)	1994 – 2006	6.9 ± 1.8 *	6.7 ± 1.6 *	7.0 ± 1.4 *
	2006 - 2014	-2.8 ± 4.1	-2.5 ± 3.7	-3.3 ± 3.2
	1994 – 2014	3.8 ± 1.0 *	3.4 ± 1.0 *	3.3 ± 0.9 *
TOC (change % per decade)	1994 – 2014	1.7 ± 0.8*	0.0 ± 0.6	0.8 ± 0.6
320nm AOD (absolute change per decade)	1994 – 2014	-0.06 ± 0.02*	-0.11 ± 0.02 *	-0.09 ± 0.01 *

3




1
2
3
4
5
6
7

Figure 1. Time series of monthly means of all-sky and clear-sky irradiances at 63° (±1°) SZA for (a) 307.5 nm irradiance, and (b) 324 nm irradiance. Monthly means derived from daily means are shown in (c) daily mean for TOC, and (d) daily mean for AOD at 320 nm. Monthly means were calculated only for months with at least 10 days of data.



1
2
3
4
5
6
7

Figure 2. Linear trends (in % per decade) of spectral UV irradiance at 307.5 nm (left) and 324 nm (right) for clear-sky (upper) and all-sky (lower) conditions derived from Brewers #086 and #005, as a function of solar zenith angle. The shaded areas represent the $\pm 1\sigma$ uncertainty of the derived trends.




1

2

3 Figure 3. Long-term changes (in % per decade) and associated 1σ uncertainty of the seasonal
 4 and the yearly mean spectral irradiance for 307.5, 324 and 350 nm at 64° SZA, for clear skies
 5 (a) and all skies (b) at Thessaloniki.

6




1

2

3 Figure 4. Long-term changes (in % per decade) and the associated 1σ uncertainty of the
 4 seasonal and the yearly mean of TOC (blue rhombs) and the AOD at 320 nm (green rhombs).
 5 The left (blue) axis corresponds to the changes in TOC while the right (green) axis to changes
 6 in AOD. ~~The changes in AOD are statistically significant (95%) while the changes in TOC~~
 7 ~~are not.~~

8



1
2
3
4
5
6
7
8
9

Figure 5. Yearly mean anomalies and corresponding trends for clear-sky irradiance at 307.5 nm (a, b, c) and 350 nm (d, e, f), TOC (g, h, i) and AOD at 320 nm (j, k, l) for December – May (left panels), June – November (middle panels) and for the entire year (right panels). A piece-wise trend consisting of two linear trends has been drawn when a statistically significant turning point has been detected; otherwise a linear trend for the entire period has been drawn.