

Ice phase in Altopcumulus Clouds over Leipzig: Remote sensing observations and detailed modelling

Martin Simmel¹, Johannes Bühl¹, Albert Ansmann¹, and Ina Tegen¹

¹TROPOS, Leibniz Institute for Tropospheric Research, Permoser Str. 15, 04318 Leipzig, Germany

Correspondence to: Martin Simmel
(simmel@tropos.de)

Abstract. The present work combines remote sensing observations and detailed cloud modeling to investigate two altopcumulus cloud cases observed over Leipzig, Germany. A suite of remote sensing instruments was able to detect primary ice at rather warm-high temperatures of -6°C . For comparison, a second mixed phase case at about -25°C is introduced. To further look into the details of cloud microphysical processes a simple dynamics model of the Asai-Kasahara type is combined with detailed spectral microphysics forming the model system AK-SPECS. Vertical velocities are prescribed to force the dynamics as well as main cloud features to be close to the observations. Subsequently, sensitivity studies with respect to ice microphysical parameters are carried out with the aim to quantify the most important sensitivities for the cases investigated. For the cases selected, the liquid phase is mainly determined by the model dynamics (location and strength of vertical velocity) whereas the ice phase is much more sensitive to the microphysical parameters (ice nuclei (IN) nucleating particle (INP) number, ice particle shape). The choice of ice particle shape may induce large uncertainties which are in the same order as those for the temperature-dependent IN-INP number distribution.

1 Introduction

Altopcumulus clouds. According to Warren et al. (1998a,b) altopcumulus and altostratus clouds together cover 22 % of the earth's surface. For single-layered altopcumulus clouds, observations by Buehl et al. (2013) show the typical feature with a maximum of liquid water in the upper part of the cloud (increasing with height) and an ice maximum in the lower part of the cloud, mostly below liquid cloud base down in the virgae. This was previously reported from Fleishauer et al. (2002) and Carey et al. (2008).

Fleishauer et al. (2002) also emphasized a lack of significant temperature inversions or wind shears as a major feature of these clouds. Kanitz et al. (2011) show that the ratio of ice-containing clouds increases with decreasing temperature. However, the numbers are different for different locations with similar dynamics but with different aerosol burden, e.g., at northern and southern midlatitudes, underlining the question for the influence of ice-nucleating particles (INP). The observations with the highest temperatures are close to the limit at which the best atmospheric ice nuclei are known to nucleate ice in the immersion mode. This can only be attributed to the aerosol particles which are formed out of or at least contain biological material such as bacteria (Hartmann et al., 2013), fungi, or pollen. This is corroborated by the review of Murray et al. (2012) stating that only biological particles are known to form ice above -15°C . However, these observations are from laboratory studies and it is still unclear whether or to which-what extent these extremely efficient ice nuclei are abundant in atmosphere, especially above the boundary layer. One idea is that freezing is caused by soil dust with biological particles dominating the freezing behaviour (O'Sullivan et al., 2014) which could explain on the one hand the atmospheric abundance of biological material and on the other hand the relatively high freezing temperatures above -15°C of ambient measurements. Seeding from ice clouds above can be excluded for the cases presented ; however, preconditioning of ice nuclei (IN) could be a reason for these high ice activation temperatures - which means that ice has formed at the cloud temperatures observed.

Ice nucleation still is a large source of uncertainty in cloud modeling. Recently, several studies use combinations of vertically fine resolved models with rather detailed representation of the ice nucleation processes. Often, wave clouds are used as comparison since they represent rather ideal condi-

tions when they are not influenced by ice seeding from layers above. Field et al. (2012) apply a 1D kinematic model with bulk microphysics but prognostic ~~IN-INP~~. Eidhammer et al. (2010) use a Lagrangian parcel model for the comparison of the ice nucleation schemes of Phillips et al. (2008) and DeMott et al. (2010) under certain constraints. A 1D column model with a very detailed 2D spectral description of liquid and ice phase is employed by Dearden et al. (2012). Sun et al. (2012) use a 1.5D model with spectral microphysics for shallow convective clouds for a sensitivity study of immersion freezing due to bacteria and its influence on precipitation formation.

Most ice microphysics descriptions in models lack from the fact that ice nuclei are not represented as a prognostic variable. These models diagnose the number of ice particles based on thermodynamical parameters such as temperature and humidity (MEYERS et al., 1992) and are, therefore, not able to consider whether ~~IN-INP~~ were already activated at previous time steps in the model.

However, despite its important contribution, ice nucleation does not determine the entire microphysics of mixed-phase clouds alone. It is rather the complex transfer between the three phases of water: water vapor, liquid water and ice described by the Wegener-Bergeron-Findeisen (WBF) mechanism (Wegener, 1911; Bergeron, 1935; Findeisen, 1938).

It is well-known that due to the different saturation pressures of water vapor with respect to liquid water and ice, a mixed-phase is in a non-equilibrium state which, nevertheless, may lead to a quasi-steady existence (Korolev and Field, 2008). The main drivers for this phase transfer are vertical velocity (leading to supersaturation and subsequent droplet formation) and ice particle formation and growth (WBF starts) leading to sedimentation of the typically fast growing ice particles (WBF ends due to removal of ice). The motivation of this work is to shed more light on the relative contributions of the different processes involved in these complex interactions.

The paper is structured as follows. Section 2 describes the remote sensing observations of two mixed-phase altocumulus cloud cases above Leipzig. The dynamical model as well as the process descriptions and initial data used for this study are specified in section 3. Section 4 refers to changes in the dynamic parameters of the model to identify base cases ~~–The results for the which describe the observations sufficiently well to perform~~ sensitivity studies with respect to ~~the microphysical parameters~~ microphysical parameters. The results for those sensitivity studies are presented in section 5 and section 6 closes with a discussion of the results.

2 Remote sensing observations

Altocumulus and altostratus clouds are regularly observed with the Leipzig Aerosol and Cloud Remote Observations

System (LACROS) at [the Leibniz Institute for Tropospheric Research TROPOS](#). LACROS combines the capabilities of Raman/depolarization lidar (Althausen et al., 2009), a MIRA-35 cloud radar (Bauer-Pfundstein and G6rsdorf, 2007), a Doppler lidar (B6uhl et al., 2012), a microwave radiometer, a sun-photometer and a disdrometer to measure height-resolved properties of aerosols and clouds. The Cloudnet framework (Illingworth et al., 2007) is used to derive microphysical parameters like liquid-water content (Pospichal et al., 2012) or ice-water content (Hogan et al., 2006). The following two cases have been selected to illustrate this variety and to serve as examples to be compared to model results.

2.1 Case 1: Warm mixed-phase cloud

One of the warmest mixed-phase clouds within the data set was observed on 17 September 2011 between 00:00 and 00:22 UTC (see Fig. ?? 1). The liquid part of the cloud extends from about 4250 m to 4450 m height at temperatures of about -6°C according to the GDAS ([Global Data Assimilation System](#)) reanalysis data for Leipzig. ~~Liquid water~~ Liquid water content (LWC) is between 0.1 g/m^{-3} to 1 g/m^{-3} whereas ice water content (IWC) is about 3–4 orders of magnitude smaller and reaches its maximum value within the virgae (see Fig. 2). Liquid water path (LWP) measured by a microwave radiometer varies between 20 and 50 g/m^2 (mostly about 25 g/m^2) whereas ice water path (IWP) is only slightly above the detection limit of about 0.01 g/m^2 implying a rather large uncertainty with correspondingly large error bars. Virgae (falling ice) are observed down to about 3000 m, which is close to the 0°C level. This is supported by Fig. 3 where the cloud radar (right panel) mainly shows particles falling from the top layer. Therefore, particles are mainly moving downwards (green color) and can be identified as ice particles by their size. Only at the very top at about 4300 m particles are small enough to still be lifted upwards (yellow colors). The Doppler lidar (left panel), however, shows the motion of small cloud droplets at the predominantly liquid cloud top. Hence, in this plot the cloud-top turbulence becomes visible. Vertical windspeeds range from about -1.5 m/s to 1.0 m/s with pdf maxima at -0.5 m/s and 0.5 m/s , respectively (Fig. ?? 3).

2.2 Case 2: Colder mixed-phase cloud

A much colder case ~~could be~~ was observed on 2 August 2012 between 21:00 UTC and 21:40 UTC (see Fig. ?? 4). Liquid water was measured around 7500 m at about -25°C with ~~an~~ a LWP between 10 and 30 g/m^2 ~~–and a LWC of up to~~ 0.1 g/m^{-3} which is much smaller than for case 1. As can be expected due to the lower temperature, the ice phase was much more massive than in case 1 and reached down to about 5500 m with an IWP of about $1\text{--}10\text{ g/m}^2$ ~~–and an IWC of up to~~ 0.01 g/m^{-3} which means that in some parts of the cloud,

ice and liquid water reach the same order of magnitude (see Fig. 5). Vertical wind speeds were in the same range as for the warmer case described above (Fig. 6).

Accuracy of the IWC is $\pm 50\%$. For the LWC calculated by the scaled adiabatic approach the same order of magnitude applies. Vertical wind speeds are measured directly by evaluation from the recorded cloud radar and Doppler lidar spectra. Errors are ± 0.15 m/s for the cloud radar and ± 0.05 m/s for the Doppler lidar. These errors are mainly due to the pointing accuracy of the two systems.

3 Model Description and initialization

For the model studies an Asai-Kasahara type model is used (Asai and Kasahara, 1967). The model geometry is axisymmetric and consists of an inner and an outer cylinder with radii of 100 m and 1000 m, respectively, resulting in a radius ratio of 1:10 which is typical for this setup. However, the geometric configuration of the model is not intended to match the geometry of the clouds (and the cloud-free spaces between the clouds) but is rather meant to provide the possibility of horizontal exchange between clouds and a cloud-free background.

The vertical resolution is constant with height and is chosen to be $\Delta z = 25$ m to give a sufficient resolution of the cloud layer and to roughly match the vertical resolution of the observations. In contrast to a parcel model, the vertically resolved model grid allows for a description of hydrometeor sedimentation. This is important especially for the fast growing ice crystals to realistically describe their interaction with the vapor and liquid phase (Bergeron-Findeisen Wegener-Bergeron-Findeisen process). A time step of 1 s was used for the dynamics as well as for the microphysics.

However, in contrast to other Asai-Kasahara model studies, updrafts are not initialized by a heat and/or humidity pulse in certain layers for a given period of time. Instead, vertical velocity (updrafts and downdrafts) in the inner cylinder is prescribed, which is more similar to a kinematic model like the Kinematic Driver Model KiD (Shipway and Hill, 2012). In that way dynamics can be controlled to make sure that it is close to the observations.

The cloud microphysics is described by the mixed-phase spectral microphysics module SPECS (Simmel and Wurzler, 2006; Diehl et al., 2006). SPECS provides a joint spectrum for the liquid phase (soluble wetted aerosol particles as well as cloud and rain drops) and one spectrum for the ice phase.

For this case study, collision processes between ice particles and drops (riming) and between ice particles and ice particles (accretion) are not taken into account. One the one hand, this is to exclude further uncertainties which would be introduced by the collision/collection kernel for those interactions, one the other hand, only small or neglectable effects are expected. Clouds are shallow which means that

there is not much time for the ice particles to interact with droplets (especially when the ice is preferentially formed near cloud base and sediments out soon). In addition, for case 1 ice particle concentrations are low which highly limits the probability of collisions. At the low temperatures of case 2 sticking efficiency is expected to be low. This assumption is corroborated by the findings of Smith et al. (2009) stating that water vapor deposition (and sublimation), balanced by sedimentation are more important than accretional growth.

3.1 Description of ice microphysics

In the following, the differences in the description of the microphysics compared to Diehl et al. (2006) are described.

3.1.1 Immersion freezing

For this study, immersion freezing is assumed to be the only primary ice formation process. Since during the above mentioned observations no *in situ* measurements of the IN-INP are available, the parameterization of DeMott et al. (2010) is used assuming that all IN-INP are active in the immersion freezing mode. The parameterization of DeMott et al. (2010) is based on an empirical relation of IN-INP and the number of aerosol particles with radii > 250 nm ($N_{AP,r>250nm}$). To cover case 1, the parameterization is extrapolated to -5°C despite the fact that the underlying measurements were only taken at -9°C and below. As base case $N_{AP,r>250nm} = 10^5 \text{ kg}^{-1}$ air is used as input data for the parameterization resulting in about 0.01 active IN-INP per liter for -6°C (case 1) and about 0.5 IN-INP per liter for -25°C (case 2), respectively, at standard conditions. This corresponds to a relatively low number of larger aerosol particles but is well within the range observed by DeMott et al. (2010).

For the potential IN-INP a prognostic temperature resolved field with 20 temperature bins with a resolution of 1 K is introduced into SPECS. It ranges from -5°C to -25°C to cover the temperature range for the selected cases and can easily be changed for other case studies. This is a simplified version of the method used by Fridlind et al. (2007). The potential IN-INP field is initially defined in every grid cell (layer) and is transported vertically with the given up/downdrafts and horizontally exchanged between inner and outer cylinder in the same way as the other hydrometeor fields (drops and ice crystals). Immersion freezing occurs as soon as liquid drops above a certain size limit are present and the temperature of a certain potential IN-INP bin is reached. Then the respective amount of drops freezes (if available) instantaneously and is transferred from the liquid to the frozen spectrum. If more drops larger than the size threshold of $10 \mu\text{m}$ than active IN-INP are present, the IN-INP are distributed evenly over all drop size bins above the threshold value. The drop size threshold was chosen to restrict freezing to droplets and to prevent (large) non-activated aerosol particles at high relative humidity (but subsaturated

275 wrt water) outside the cloud from freezing. If ice crystals 325
melt below the freezing level, they contribute to the poten-
tial ~~IN-INP~~ field at that level.

3.1.2 Ice particle shape

280 It is well known that ice particle shape highly influences wa-
ter vapor deposition (described by changing the capacitance
of the particle) as well as terminal fall velocity of the ice
particle. Therefore, instead of the previously chosen spher-
ical ice particle shape, ice particles now can be prescribed
285 as hexagonal columns or plates. The aspect ratio can be ei-
ther constant for all size bins or be changed with size fol-
lowing the approach of Mitchell (1996). Typically, with in-
creasing particle size, the deviation from an uniform aspect 335
ratio increases. In our simulations, a constant uniform aspect
ratio ($ar=1$) is used as base case. From Mitchell (1996) the
size-varying aspect ratios for plates ~~and columns are (ranging~~
290 ~~from 15 to 3000 μm with a single description) and columns~~
~~(for size ranges of 30 to 100 μm , 100 to 300 μm , and above~~
~~300 μm in diameter) are calculated from the mass-dimension 340~~
~~power laws and used for sensitivity studies.~~

295 The (relative) capacitance needed for the calculation of
deposition growth of the ice crystals is modeled using the
method of Westbrook et al. (2008) for the aspect ratios given
above. Ice crystal terminal fall velocities are calculated ac- 345
cording to Heymsfield and Westbrook (2010) using the same
aspect ratios. 300

3.2 Model initialization

3.2.1 Thermodynamics

The Asai-Kasahara model has to be initialized with vertical
profiles of temperature and dewpoint temperature either from
305 reanalysis data (here GDAS) or radiosonde profiles from
nearby stations (here Meiningen, Thuringia). Fig. 7 shows
profiles of temperature and relative humidities with respect
to liquid water and to ice, respectively, for both cases. For
case 1, profiles from both methods show a similar general
behaviour but the radiosonde profile of Meiningen measured
310 at 00 UTC is used since it provides a finer vertical resolution
than the GDAS reanalysis data ~~(cp. Fig. 7).~~ However, for 360
case 2 the Meiningen RS profile misses the humidity layer
at the level where the clouds were observed ~~and, therefore,~~
315 ~~This means that the profile is not representative for the given~~
~~meteorological situation. Therefore,~~ GDAS reanalysis data
for Leipzig at 21 UTC were chosen. Finally, both profiles 365
used show a sufficiently humid layer where the clouds were
observed, so that lifting of these layers lead to supersatura-
tion and subsequent cloud formation. 320

~~In contrast to other Asai-Kasahara model studies, updrafts~~
~~are not initialized by a heat and/or humidity pulse in certain~~
~~layers for a given period of time. Instead, As mentioned 370~~
~~above,~~ vertical velocity (updrafts and downdrafts) in the in-

ner cylinder is prescribed at cloud level ranging from h_{bot} to
 h_{top} . The center of this interval is given by $h_{mid} = (h_{top} +$
 $h_{bot})/2$ and its half-depth by $h_{depth} = (h_{top} - h_{bot})/2$. h_{bot}
ranges from 3800 m to 4100 m for case 1 and from 7000 m to
7300 m for case 2. The respective values for h_{top} are 4500 m
and 7700 m. The vertical dependency (compare Fig. 8, left)
is given by

$$f_h(h) = \frac{h_{depth}^2 - (h - h_{mid})^2}{h_{depth}^2} \quad \text{for } h_{bot} \leq h \leq h_{top} \quad (1)$$

resulting in the time- and height-dependent function

$$w(h, t) = w_{mid}(t) f_h(h) \quad \text{for } h_{bot} \leq h \leq h_{top} \quad (2)$$

and $w(h, t) = 0$ otherwise, defining $w_{mid}(t)$ as the updraft
velocity at h_{mid} . In order to match the observed wind field
distributions rather closely, $w_{mid}(t)$ is chosen as a stochastic
function

$$w_{mid}(t) = w_{ave} + f_{scal} \frac{\delta(t)^3}{|\delta(t)|} \quad (3)$$

where w_{ave} is the average ('large-scale') updraft velocity at
 h_{mid} varying between 0.1 m/s and 0.4 m/s, f_{scal} is the scal-
ing factor determining the range of updraft velocities (chosen
as 4 m/s to obtain a difference of minimum and maximum ve-
locity of 2 m/s), and $\delta(t)$ is a random number ranging from
-0.5 to +0.5 obtained from a linear stochastic process pro-
vided by FORTRAN. After 30 s model time a new $\delta(t)$ is
created. Different realizations of the stochastic process are
tested (see below). E.g., $w_{mid}(t)$ ranges from -0.7 m/s to
1.3 m/s if $w_{ave} = 0.3$ m/s and $f_{scal} = 4$ m/s as it is shown in
350 the temporal evolution and the histogram in ~~Figure Fig.~~ 8.

Due to the height dependent vertical velocity w , a horizon-
tal transport velocity u_k (exchange between inner and outer
cylinder) is induced in the Asai-Kasahara formulation for a
given model layer k .

$$u_k = - \frac{w_{k+\frac{1}{2}} \rho_{k+\frac{1}{2}} - w_{k-\frac{1}{2}} \rho_{k-\frac{1}{2}}}{f_r \Delta z \rho_k} \quad (4)$$

Full indices k indicate values at level centers whereas
half indices ($k + \frac{1}{2}$, $k - \frac{1}{2}$) describe values at level inter-
faces. $f_r = 2/r_i$ is a geometry parameter with the radius
 ~~$r_i = 1000$~~ $r_i = 100$ m of the inner cylinder.

The prescribed velocity field leads to the following effects
(all descriptions are related to the inner cylinder if not stated
otherwise explicitly):

- In the updraft phase: In the upper part (between h_{mid}
and h_{top}) of the updraft, mixing occurs from the inner
to the outer cylinder whereas in the lower part (between
 h_{low} and h_{mid}), horizontal transport is from the outer
cylinder into the inner one
- For downdrafts it is the other way: This means that be-
low h_{mid} drops and ice particles are transported from
the inner cylinder to the outer one and are therefore re-
moved from the inner cylinder

- below h_{low} or above h_{top} , no horizontal exchange takes place.

The question arises to which extent this dynamical behaviour reflects the real features of the observed clouds and whether this is critical for the topics aimed at in this study.

Prescribing vertical velocity in any way also means that a feedback of microphysics on dynamics due to phase changes (e.g., release of latent heat for condensing water vapor or freezing/melting processes) is not considered by the model.

3.2.2 Aerosol distribution

Since no in situ aerosol measurements are available, literature data is used. The Raman lidar observations do not show any polluted layers for both cases; therefore data from LACE98 (Petzold et al., 2002) are used which should be representative for the free troposphere over Leipzig. For case 1 values for the lower free troposphere (M6), for case 2 those from the upper free troposphere (M1) are used (see Petzold et al., 2002, Tab. 6) (see Petzold et al., 2002, Tab. 6).

4 Model results: Dynamics

In a first step, the aim is to achieve a sufficient agreement concerning macroscopic cloud features as well as (liquid phase) microphysics as far as they were observed. The following parameters describing model dynamics (updraft velocity) are varied to identify a ‘best case’ which in the second step can be used to perform sensitivity studies with respect to (ice) microphysics (see also Tables 1 and 2).

- h_{low} : ranging from 3800 m to 4100 m for the warmer and from 7000 m to 7300 m for the colder case. This parameter influences the vertical cloud extent and, therefore, liquid water content and liquid water path.
- w_{ave} : ranging from 0.1 m/s to 0.4 m/s. Higher average updraft also leads to higher LWC. Due to the lateral mixing processes the model setup requires a positive updraft velocity in average to form and maintain clouds.
- δ : Four different realizations of the stochastic process are used. This influences the timing of the cloud occurrence as well as LWC and LWP but not systematically.

All model results shown refer to the inner cylinder.

4.1 Case 1: Warm mixed-phase cloud

Figs. 9 and 11 show time-height plots of the liquid (contours, linear scale) and ice (colours, logarithmic scale) water mixing-ratio content for case 1 illustrating the cloud sensitivity with respect to variation of cloud base (h_{bot}), average vertical updraft (w_{ave}), and the realization of the stochastic

process, respectively. Liquid clouds form in the updraft regions (cp. Fig. 8) whereas in the downdrafts the liquid phase vanishes at least partly. ~~Ice forms mainly at cloudbase and immediately starts.~~ If active INP are available ice formation can take place within the liquid part of the cloud. The INP are partly already active near liquid cloud base which means that they trigger freezing as soon as the droplets are formed. Less efficient INP become active after further cooling above cloud base. After ice formation rapid depositional growth takes place and the ice particles almost immediately start to sediment. Due to the supersaturation with respect to ice even below liquid cloud base, ice particles still grow while sedimenting, reaching their maximum size before, finally, subsaturated regions are reached and ~~evaporation~~ sublimation sets in. Figs. 10 and 12 show the time evolution of liquid (lower panel) and ice water path (upper panel) for the same parameters varied, reflecting the same temporal patterns. ~~Table Tab. 1~~ summarizes the maximum values for liquid and ice water ~~mixing-ratio~~ (LWMCcontent (LWC/IWMC)), liquid and ice water path (LWP/IWP) as well as cloud droplet and ice particle number concentration (CDN/IPN) for all dynamics sensitivity runs for case 1.

One can clearly observe, that a lower h_{bot} (Fig. 9) results in a lower cloud base, larger vertical cloud extent as well as more liquid water. ~~The same trend with similar intensity~~ LWC maxima are within a factor of 2 for varying h_{bot} . A similar trend is observed for the ice phase (see also Fig. 10) ~~10~~, but IWC maxima differ only by about 25 %. However, the values of the two maxima of the condensed phase after about 15–20 min and about 40 min model time are quite different. The first maximum is more pronounced for the ice phase whereas the second one is larger for the liquid phase. While the liquid phase is dominated by the updraft velocity (see Fig. 8) the ice phase additionally depends on ~~IN~~ INP supply. In the first ice formation event at 15 min, all ~~IN~~ INP active at the current temperature actually form ice leading to an ~~IN-IPN~~ INP depletion. Due to the horizontal exchange with the outer cylinder the ~~IN-IPN~~ INP reservoir is refilled, but only to a certain extent when the second cloud event after 40 min sets in. Due to the limited ~~IN-IPN~~ INP supply the second ice maximum is weaker than the first one. The stochastic velocity fluctuations cause fluctuations in relative humidity, which are directly reflected by the liquid phase parameters whereas the ice phase generally reacts much slower. Sensitivity of CDN and IPN with respect to change of h_{bot} does not seem to be systematic.

Increasing the average updraft velocity w_{ave} leads to a similar increase of liquid water and ice as lowering h_{bot} (see Figs. 11, upper panel and 12, left). This can be expected since more water vapor flows through the cloud and is able to condense. However, a certain limit seems to be reached for W_w04, since the increase of LWP slows down (see maximum value at 40 min in Fig. 12, left). This is due to the enhanced horizontal exchanged following eq. (4). Additionally, the stronger updrafts allow the ice particles a longer presence

time in the vicinity of the cloud and, therefore, an enhanced growth at comparably high supersaturation with respect to ice before sedimentation sets in at larger sizes. This also leads to an accumulation of ice particles and, therefore, to a higher IPN. Surprisingly, CDN ~~only depends~~ depends only weakly and not systematically on w_{ave} which is in contrast to the typical enhancement of CDN with increasing updraft velocities.

Figs. 11 (lower panel) and 12 (right) show that different realizations of the stochastic process (as explained above in section 3.2.1) lead to different temporal cloud evolutions. However, differences in maximum LWP and ~~LWLR-LWC~~ LWLR-LWC are much smaller than those discussed above. Variations in maximum IWP and ~~IWLR-IWC~~ IWLR-IWC as well as CDN and IPN are in the range of about 30%. This is also true for average LWP ranging from 18 g/m² for W_r1 to 26 g/m² for W_r3. However, despite the different maxima and temporal evolutions of IWP, average IWP is almost identical for the different stochastic realizations (0.023 g/m²). This shows that changing the stochastic realization influences cloud evolution in detail (timing) but does not change the overall picture.

With maximum values between 17 and 57 g/m² the modeled liquid water path is in the same range as the observed values (20–50 g/m²), especially for the 'wetter' runs (smaller h_{bot} , larger w_{ave}). Average LWP typically is about half (40–60%) of the maximum value for most of the runs which also fits well into the observations. Ice forms within the liquid layer and sediments to about 3800 m for most runs which is less than for the observations. The (maximum) modeled ice mixing ratio is in the same order of magnitude as the observed one (about 10⁻⁷ kg/m³). The same holds for the ice water path with values of about 0.01 g/m² for both, model and observation. For the other values, no observational data is available for comparison.

4.2 Case 2: Colder mixed-phase cloud

Due to the ~~colder-lower~~ colder-lower temperatures of case 2 much more ~~INP are active and much more~~ ice is produced than in case 1 (see Figs. 13–16 as well as Tab. 2). ~~Therefore This~~ also means that near cloud base much more active INP are available and that a further cooling within the clouds only slightly increases the number of active INP leading again to a preferential ice nucleation near liquid cloud base. Due to the lower temperatures and the more massive ice formation, the virgae reach down to more than 1500 m below liquid cloud base which is in concordance with the observations. The principal behaviour with respect to the sensitivity parameters is similar as in case 1: The liquid phase is enhanced by either decreasing h_{bot} or increasing w_{ave} , showing the 'saturation' effect slightly more pronounced as in case 1. Different stochastic realizations only weakly influence the maximum and average values of the liquid phase but change the timing of occurrence. Generally, the variability of the ice phase is weaker than in case 1. The different stochastic realizations

show the highest variability in ~~IWLR-IWC~~ IWLR-IWC and IWP. Different variations of h_{bot} show almost identical IWPs, whereas changing w_{ave} at least slightly influences maximum ~~IWLR-IWC~~ IWLR-IWC and IWP, which again can be attributed to the ice particle accumulation in the updraft. Liquid water path is smaller than in case 1 and reaches maximum values between 10 and 43 g/m² which well covers the observed maximum value of about 20 g/m². Cloudnet observations show an IWC of 10⁻⁷ – 10⁻⁵ kg/m³ which is an increase by a factor of 10–100 compared to case 1. Similar values are obtained by the model results underlining the strong temperature dependency of the ice nucleation process.

5 Sensitivity studies

In the previous section it could be shown that dynamical parameters can be chosen in a way that the model results (in terms of LWP, IWP as well as cloud geometry) are in good agreement with the observations. This allows to perform sensitivity studies with respect to cloud microphysics. To cover the proper sensitivities we have to answer the question which microphysical parameters are expected to have a large influence on mixed phase microphysics and are rather uncertain to be estimated. This leads to (temperature-dependent) ~~IN~~ number (N_{INP}) ~~number (N_{INP})~~ number (N_{INP}) which directly influences ice particle number but mostly is poorly known. To be consistent with the freezing parameterization of the model, ~~N_{INP}~~ N_{INP} is varied by changing $N_{AP,r>250nm}$ which additionally is easier to observe in most cases. A second parameter is the shape of the ice particles which does not influence the primary freezing process but the subsequent growth by water vapor deposition onto existing ice particles and, therefore, the total ice mass produced. Their relative importance shall be quantified and also be compared to the influence of dynamics discussed above.

5.1 ~~IN-INP~~ number

Changing $N_{AP,r>250nm}$ leads to a temperature-dependent change of ~~IN-INP~~ number which is relatively small for warmer conditions. However, the effect increases with decreasing temperature. This is illustrated by the following numbers. The parameterization of DeMott et al. (2010) gives about 0.009 active ~~IN-INP~~ number per liter at standard conditions (~~N_{INP}~~ N_{INP}) when $N_{AP,r>250nm} = 10^5 \text{ kg}^{-1}$ at $T = -5^\circ\text{C}$. A tenfold increase to $N_{AP,r>250nm} = 10^6 \text{ kg}^{-1}$ results in about 0.012 active ~~IN-INP~~ number per liter which is a rise of only about 35%. For $T = -7^\circ\text{C}$, ~~IN-INP~~ number rises by about 65% for a tenfold increase of $N_{AP,r>250nm}$. This shows that for those rather ~~warm-high~~ temperatures considered for case 1, a massive change in $N_{AP,r>250nm}$ leads to relatively small changes in ~~N_{INP}~~ N_{INP} and only a small effect on the ice phase can be expected. This is confirmed by Fig. 17 (left) showing liquid and ice water ~~mixing ratios~~ mixing ratios

575 [contents](#) for W_in6. [Ice-mass-IWC](#) is enhanced by less than 60 % for W_in6 and by about 160 % for W_in7 which is consistent for the given temperature range (see Tab. 3). Similar values are obtained for the change in IPN. This directly leads to the conclusion that the individual ice particles grow independently from each other. Their individual growth history is (in contrast to drop growth) only influenced by thermodynamics as long as their number is low enough which seems to be the case here.

580 This is confirmed by Fig. 18 showing drop and ice particle size distributions at the time when the maximum IWP is reached (16 min for case 1, 17 min for case 2). For case 1 (upper panel), the liquid phase (contours) is unaffected by the [IN-INP](#) enhancement. Despite the increase of ice particle number and mass the shape of the ice particle size distribution (colors) is not changed. The smallest ice particles can be observed at three discrete height (and temperature) levels caused by the temperature resolved parameterization of the potential [IN-INP](#) described in section 3.1.1. In reality this part of the spectrum showing rather freshly nucleated and fast growing ice particles should be continuous over the height range from about 4100 m to 4400 m. Nevertheless, the total number of ice particles formed is described correctly.

585 One can conclude that increasing [IN-INP](#) number therefore increases ice particle number as well as ice mass proportionally. Generally, the ice mass remains small and the liquid phase is not affected by the ice mass increase. Those results are supported by Fig. 19 (left) showing an unchanged LWP and a proportionally growing IWP for increased [IN-INP](#) numbers.

590 For the colder case 2 the parameters are varied in the same way. However, one big difference is that a tenfold increase of $N_{AP,r>250nm}$ at $T = -25^\circ\text{C}$ results in a much larger change in active [IN-INP](#). Their number rises by 300 % from about 0.5 per liter to about 2 per liter following the parameterization. This is reflected by the IPN values in [Table Tab. 4](#). Fig. 17 (right) and [Table Tab. 4](#) show that ice mass increases in such a way that liquid water is depleted partially (C_in6 by about 50 %) or almost totally (C_in7) due to the [Bergeron-Findeisen-Wegener-Bergeron-Findeisen](#) process. Compared to C_base, ice is enhanced by a factor of 3–4 for C_in6 and about 10 for C_in7 whereas IPN increases by a factor of 12. This can also be seen in the IWP (Fig. 19, right, red lines) showing a limited increase for C_in7, especially for the first maximum after 17 min. This means that the results for C_in6 are still consistent with an independent growth of the individual ice particles (as described above) despite the relatively high ice occurrence.

595 This is verified by the size distributions in Fig. 18 (lower panel). As in case 1 the ice particle size distributions only differ by the number/mass, but not by shape. Additionally, the decrease in the liquid phase is reflected also in the drop spectrum showing a more shallow liquid part of cloud as well as droplet distribution shifted to smaller sizes.

However, for C_in7 the ice particles compete for water vapor which becomes clear from (i) the depletion of liquid water (resulting in a lower supersaturation with respect to ice) and (ii) the ice mass enhancement factor being below the value expected from the ice nucleation parameterization and below that of IPN. This means that despite the higher number of [IN-INP](#) and, therefore, ice particles, the amount of ice is limited by the thermodynamic conditions which results in the production of more but smaller ice particles, similar to the Twomey effect for drop activation.

[As mentioned earlier, ice particle growth is not restricted to the liquid part of the cloud but also occurs in the layer below liquid cloud base, which is still supersaturated with respect to ice. This leads to a decrease in relative humidity in this part of the cloud, which in turn weakens or suppresses droplet formation by shifting liquid cloud base to higher altitudes. The lower LWC for the runs with higher IWC therefore cannot only be attributed to the WBF processes but also to this indirect effect.](#)

5.2 Ice particle shape

As discussed previously, for most of the cases (except for C_in7) changing the parameters in the section above does neither influence the ice particles themselves nor their individual growth. Additionally, due to their low number, there is almost no competition of the ice particles for water vapor, and, therefore, ice water content scales linearly with ice particle number. In contrast to this, changing the ice particle shape from quasi-spherical ($ar=1$) to columns or plates with size-dependent axis ratios deviating from unity results in an increase of water vapor deposition on the individual ice particles leading to enhanced ice water content due to larger individual particles when ice particle numbers remain unchanged. This is due to (i) enhanced relative capacitance resulting in faster water vapor deposition and (ii) lower terminal velocities of the ice particles leading to longer residence times in vicinity of conditions with supersaturation with respect to ice.

Fig. 20 (left) shows the results for the runs using hexagonal columns (W_col) as prescribed ice particle type. Compared to the previous results (W_base, W_in6, W_in7) more ice mass is produced (see [Table Tab. 3](#)) but still the liquid part of the cloud remains unaffected (compare also LWP and IWP in Fig. 19, left). Similar results are obtained for the assumption of plate-like ice particles (W_pla). The mass increase results from the larger ice particle size due to the reasons discussed above which can be seen from Fig. 21 showing the size distributions for W_col at different times. On the upper left panel W_col is shown after 16 min corresponding to Fig. 18. Compared to W_base, larger ice particles are produced leading to more ice mass (equivalent radius up to $300\ \mu\text{m}$ compared to $189\text{--}238\ \mu\text{m}$ for the base case). Additionally, due to the lower fall speed of the columns ($1.03\ \text{m/s}$ vs. $1.75\text{--}2.24\ \text{m/s}$), the maximum of the ice is at about 4200 m compared to 4100 m

for the base case. On the upper right panel, size distributions after 21 min are shown corresponding to the IWP maximum of W_col. Ice particles have grown larger (equivalent radius up to $378 \mu\text{m}$, length of the columns increases from about 3 mm to 4.5 mm) and sedimentation has developed further with increasing terminal velocity (1.13 m/s). Similar results are obtained for plates (W_pla) with terminal velocities of 0.89–1.21 m/s, equivalent radii of 300–476 μm and maximum dimension of 1.8–3.2 mm.

The lower terminal velocity of columns and plates despite their larger size is leading to the stronger tilting of the virgae. Additionally, ice particle number IPN is enhanced by about 30% although ice nucleation is identical to W_base. This can be attributed to the lower fall velocities, too, leading to an accumulation of ice particles. The differences between W_col and W_pla are caused by both, the higher relative capacitances of and lower terminal fall velocities of plates compared to columns (at least when their axis ratios are chosen following Mitchell (1996)).

For case 2 (C_col and C_pla), the liquid water reduction due to the Bergeron-Findeisen process is similar to C_in6 (see Fig. 20, right, and Table Tab. 4). In contrast to the respective case 1 runs, less ice is produced than for C_in7. The tilting of the virgae is not as strong as in W_col which is due to the larger ice particle sizes leading to higher terminal fall velocities (1.43–1.60 m/s). Additionally, the lower air density leads to an increase of terminal velocity of more than 10% independently from shape. Fig. 21 (lower panels) show panel shows the size distributions for C_col at different times. Due to the longer growth time larger individual ice particles than in case 1 are produced (equivalent radius up to 600 μm compared to 300 μm for the base case).

To decide whether independent ice particle growth or competition occurs, further runs with less IN-INP (C_col_in4 and C_pla_in4) are discussed (see Fig. 19, right). IWMR-IWC and IWP of these runs (in4) are about one third of the values of the respective runs with more IN-INP (in5). For ice particle number, a factor of slightly more than three occurs which means that a weak competition for water vapor occurs for C_col and C_pla resulting in slightly smaller individual ice particles compared to C_col_in4 and C_pla_in4.

6 Conclusions

The model system AK-SPECS was applied to simulate dynamical and microphysical processes within altocumulus clouds. Sensitivity studies on relative contributions on cloud evolution as well as comparisons to observations were made.

Variation of the dynamic parameters as it was done in section 4 leads to systematic differences mainly in the liquid phase (LWMRLWC, LWP) which can easily be explained. More liquid water is produced when either cloud base is lowered (corresponding to a larger vertical cloud extent) or vertical wind velocity is increased. However, the effects of

the dynamics on the ice phase are surprisingly small, at least smaller than those on the liquid phase. Increasing vertical velocity leads to an accumulation of the smaller ice particles in the enhanced updraft.

On the other hand, much larger differences in terms of IWMR-IWC and IWP were found when microphysical parameters like IN-INP number or ice particle shape were varied under identical dynamic conditions. This is valid for both cases studied. However, at least for the ice nucleation parameterization used, sensitivity of IN-INP number strongly increased with decreasing temperature.

This means that relatively large differences concerning the ice phase can only be reached when either IN-INP number differs considerably or ice particle shape is different (which should not be the case for relatively similar thermodynamical conditions). After Fukuta and Takahashi (1999) for case 1 with temperatures of about -6°C column-like ice particles with $ar = 0.1$ could be expected (corresponding to W_col) whereas for case 2 ($T < -24^\circ\text{C}$) hexagonal particles with $ar = 1$ are most likely (e.g., C_base). Those ice shapes were observed in laboratory studies at water saturation which was also valid for the observed cases when ice formed by immersion freezing within the liquid layer of the cloud. However, below liquid cloud base supersaturation with respect to ice decreases. These ice shapes can also explain why a depletion of the liquid phase was not observed in case 2 as it was predicted by the sensitivity studies using either columns or plates as prescribed shape. Generally, the liquid phase is affected considerably only when enough ice particles are present which typically is the case for cold conditions with a sufficient amount of IN-INP and fast growing ice particle shapes (most effective for large deviations from spherical shapes).

Acknowledgements. Funding—This study was supported by the Deutsche Forschungsgemeinschaft DFG-project UDINE-

urlstyle

Althausen, D., Engelmann, R., Baars, H., Heese, B., Ansmann, A., Müller, D., and Komppula, M.: Portable Raman Lidar PollyXT for Automated Profiling of Aerosol Backscatter, Extinction, and Depolarization, *J. Atmos. Oceanic Technol.*, 26, 2366–2378, 2009.

Asai, T. and Kasahara, A.: A theoretical study of compensating downward motions associated with cumulus clouds, *JOURNAL OF THE ATMOSPHERIC SCIENCES*, 24, 487–496, 1967.

Bauer-Pfundstein, M. R. and Grsdorf, U.: Target Separation and Classification Using Cloud Radar Doppler Spectra, in: *Proceedings of the 33rd Conference on Radar Meteorology, 2007.*

Buehl, J., Ansmann, A., Seifert, P., Baars, H., and Engelmann, R.: Toward a quantitative characterization of heterogeneous ice formation with lidar (DFG) under Grant AN 258/radar: Comparison of CALIPSO/CloudSat with ground-based observations,

- GEOPHYSICAL RESEARCH LETTERS, 40, 4404–4408, 2013.
- Bühl, J., Engelmann, R., and Ansmann, A.: Removing the Laser-Chirp Influence from Coherent Doppler Lidar Datasets by Two-Dimensional Deconvolution, *J. Atmos. Oceanic Technol.*, 29, 1042–1051, 2012.
- Dearden, C., Connolly, P. J., Choulaton, T., Field, P. R., and Heymsfield, A. J.: Factors influencing ice formation and growth in simulations of a mixed-phase wave cloud, *JOURNAL OF ADVANCES IN MODELING EARTH SYSTEMS*, 4, 2012.
- DeMott, P. J., Prenni, A. J., Liu, X., Kreidenweis, S. M., Petters, M. D., Twohy, C. H., Richardson, M. S., Eidhammer, T., and Rogers, D. C.: Predicting global atmospheric ice nuclei distributions and their impacts on climate, *PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA*, 107, 11217–11222, 2010.
- Diehl, K., Simmel, M., and Würzler, S.: Numerical sensitivity studies on the impact of aerosol properties and drop freezing modes on the glaciation, microphysics, and dynamics of clouds, *JOURNAL OF GEOPHYSICAL RESEARCH-ATMOSPHERES*, 111, 2006.
- Eidhammer, T., DeMott, P. J., Prenni, A. J., Petters, M. D., Twohy, C. H., Rogers, D. C., Stith, J., Heymsfield, A., Wang, Z., Pratt, K. A., Prather, K. A., Murphy, S. M., Seinfeld, J. H., Subramanian, R., and Kreidenweis, S. M.: Ice Initiation by Aerosol Particles: Measured and Predicted Ice Nuclei Concentrations versus Measured Ice Crystal Concentrations in an Orographic Wave Cloud, *JOURNAL OF THE ATMOSPHERIC SCIENCES*, 67, 2417–2436, 2010.
- Field, P. R., Heymsfield, A. J., Shipway, B. J., DeMott, P. J., Pratt, K. A., Rogers, D. C., Stith, J., and Prather, K. A.: Ice in Clouds Experiment Layer Clouds. Part II: Testing Characteristics of Heterogeneous Ice Formation in Lee-Wave Clouds, *JOURNAL OF THE ATMOSPHERIC SCIENCES*, 69, 1066–1079, 2012.
- Fridlind, A. M., Aekerman, A. S., McFarquhar, G., Zhang, G., Poellot, M. R., DeMott, P. J., Prenni, A. J., and Heymsfield, A. J.: Ice properties of single-layer stratocumulus during the Mixed-Phase Arctic Cloud Experiment: 2. Model results, *JOURNAL OF GEOPHYSICAL RESEARCH-ATMOSPHERES*, 112, 2007.
- Fukuta, N. and Takahashi, T.: The Growth of Atmospheric Ice Crystals: A Summary of Findings in Vertical Supercooled Cloud Tunnel Studies, *JOURNAL OF THE ATMOSPHERIC SCIENCES*, 56, 1963–1979, 1999.
- Hartmann, S., Augustin, S., Clauss, T., Wex, H., Santl-Temkiv, T., Voigtlander, J., Niedermeier, D., and Stratmann, F.: Immersion freezing of ice nucleation active protein complexes, *Atmospheric Chemistry and Physics*, 13, 5751–5766, 2013.
- Heymsfield, A. J. and Westbrook, C. D.: Advances in the Estimation of Ice Particle Fall Speeds Using Laboratory and Field Measurements, *JOURNAL OF THE ATMOSPHERIC SCIENCES*, 67, 2469–2482, 2010.
- Hogan, R. J., Mittermaier, M. P., and Illingworth, A. J.: The Retrieval of Ice Water Content from Radar Reflectivity Factor and Temperature and Its Use in Evaluating a Mesoscale Model, *J. Appl. Meteor. Climatol.*, 45, 301–317, 2006.
- Illingworth, A. J., Hogan, R. J., O'Connor, E. J., Bouniol, D., Delanoë, J., Pelon, J., Protat, A., Brooks, M. E., Gaussiat, N., Wilson, D. R., Donovan, D. P., Baltink, H. K., van Zadelhoff, G. J., Eastment, J. D., Goddard, J. W. F., Wrench, C. L., Haeffelin, M., Krasnov, O. A., Russechenberg, H. W. J., Piriou, J.-M., Vinit, F., Seifert, A., Tompkins, A. M., and Willén, U.: Cloudnet, *Bull. Amer. Meteor. Soc.*, 88, 883–898, 2007.
- Meyers, M., DeMott, P., and Cotton, W.: NEW PRIMARY ICE-NUCLEATION PARAMETERIZATIONS IN AN EXPLICIT CLOUD MODEL, *JOURNAL OF APPLIED METEOROLOGY*, 31, 708–721, 1992.
- Mitchell, D.: Use of mass-15. We also acknowledge funding from the EU FP7-ENV-2013 programme “impact of Biogenic vs. Anthropogenic emissions on Clouds and area-dimensional power laws for determining precipitation particle terminal velocities Climate: towards a Holistic Understanding” (BACCHUS), *JOURNAL OF THE ATMOSPHERIC SCIENCES*, 53, 1710–1723, 1996. project.no.603445.
- Murray, B. J., O'Sullivan, D., Atkinson, J. D., and Webb, M. E.: Ice nucleation by particles immersed in supercooled cloud droplets, *Chem. Soc. Rev.*, 41, 6519–6554, 2012.
- O'Sullivan, D., Murray, B. J., Malkin, T. L., Whale, T. F., Umo, N. S., Atkinson, J. D., Price, H. C., Baustian, K. J., Browse, J., and Webb, M. E.: Ice nucleation by fertile soil dusts: relative importance of mineral and biogenic components, *Atmospheric Chemistry and Physics*, 14, 1853–1867, 2014.

References

- Althausen, D., Engelmann, R., Baars, H., Heese, B., Ansmann, A., Müller, D., and Komppula, M.: Portable Raman Lidar PollyXT for Automated Profiling of Aerosol Backscatter, Extinction, and Depolarization, *J. Atmos. Oceanic Technol.*, 26, 2366–2378, doi:10.1175/2009JTECHA1304.1, 2009.
- Asai, T. and Kasahara, A.: A theoretical study of compensating downward motions associated with cumulus clouds, *JOURNAL OF THE ATMOSPHERIC SCIENCES*, 24, 487–496, doi:10.1175/1520-0469(1967)024<0487:ATSOTC>2.0.CO;2, 1967.
- Bauer-Pfundstein, M. R. and Görsdorf, U.: Target Separation and Classification Using Cloud Radar Doppler-Spectra, in: Proceedings of the 33rd Conference on Radar Meteorology, 2007.
- Bergeron, T.: On the physics of clouds and precipitation, in: *Process verbaux de l'association de Meteorologie*, pp. 156–178, International Union of Geodesy and Geophysics, Lisboa, Portugal, 1935.
- Buehl, J., Ansmann, A., Seifert, P., Baars, H., and Engelmann, R.: Toward a quantitative characterization of heterogeneous ice formation with lidar/radar: Comparison of CALIPSO/CloudSat with ground-based observations, *GEOPHYSICAL RESEARCH LETTERS*, 40, 4404–4408, doi:10.1002/grl.50792, 2013.
- Bühl, J., Engelmann, R., and Ansmann, A.: Removing the Laser-Chirp Influence from Coherent Doppler Lidar Datasets by Two-Dimensional Deconvolution, *J. Atmos. Oceanic Technol.*, 29, 1042–1051, doi:10.1175/JTECH-D-11-00144.1, 2012.
- Carey, L. D., Niu, J., Yang, P., Kankiewicz, J. A., Larson, V. E., and Vonder Haar, T. H.: The Vertical Profile of Liquid and Ice

- Water Content in Midlatitude Mixed-Phase Altocumulus Clouds, *J. Appl. Meteorology and Climate*, 47, 2487–2495, 2008. ⁹⁵⁵
- Dearden, C., Connolly, P. J., Choullarton, T., Field, P. R., and Heymsfield, A. J.: Factors influencing ice formation and growth in simulations of a mixed-phase wave cloud, *JOURNAL OF ADVANCES IN MODELING EARTH SYSTEMS*, 4, doi:10.1029/2012MS000163, 2012. ⁹⁶⁰
- DeMott, P. J., Prenni, A. J., Liu, X., Kreidenweis, S. M., Petters, M. D., Twohy, C. H., Richardson, M. S., Eidhammer, T., and Rogers, D. C.: Predicting global atmospheric ice nuclei distributions and their impacts on climate, *PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA*, 107, 11 217–11 222, doi:10.1073/pnas.0910818107, 2010. ⁹⁶⁵
- Diehl, K., Simmel, M., and Wurzler, S.: Numerical sensitivity studies on the impact of aerosol properties and drop freezing modes on the glaciation, microphysics, and dynamics of clouds, *JOURNAL OF GEOPHYSICAL RESEARCH-ATMOSPHERES*, 111, doi:10.1029/2005JD005884, 2006. ⁹⁷⁰
- Eidhammer, T., DeMott, P. J., Prenni, A. J., Petters, M. D., Twohy, C. H., Rogers, D. C., Stith, J., Heymsfield, A., Wang, Z., Pratt, K. A., Prather, K. A., Murphy, S. M., Seinfeld, J. H., Subramanian, R., and Kreidenweis, S. M.: Ice Initiation by Aerosol Particles: Measured and Predicted Ice Nuclei Concentrations versus Measured Ice Crystal Concentrations in an Orographic Wave Cloud, *JOURNAL OF THE ATMOSPHERIC SCIENCES*, 67, 2417–2436, doi:10.1175/2010JAS3266.1, 2010. ⁹⁸⁰
- Field, P. R., Heymsfield, A. J., Shipway, B. J., DeMott, P. J., Pratt, K. A., Rogers, D. C., Stith, J., and Prather, K. A.: Ice in Clouds Experiment-Layer Clouds. Part II: Testing Characteristics of Heterogeneous Ice Formation in Lee Wave Clouds, *JOURNAL OF THE ATMOSPHERIC SCIENCES*, 69, 1066–1079, doi:10.1175/JAS-D-11-026.1, 2012. ⁹⁸⁵
- Findeisen, W.: Kolloid-meteorologische Vorgänge bei Niederschlagsbildung, *Meteorolog. Z.*, 55, 121–133, 1938. ⁹⁹⁰
- Fleishauer, R. P., Larson, V. E., and Vonder Haar, T. H.: Observed Microphysical Structure of Midlevel, Mixed-Phase Clouds, *J. Atmos. Sci.*, 59, 1779–1804, 2002. ⁹⁹⁵
- Fridlind, A. M., Ackerman, A. S., McFarquhar, G., Zhang, G., Poellot, M. R., DeMott, P. J., Prenni, A. J., and Heymsfield, A. J.: Ice properties of single-layer stratocumulus during the Mixed-Phase Arctic Cloud Experiment: 2. Model results, *JOURNAL OF GEOPHYSICAL RESEARCH-ATMOSPHERES*, 112, doi:10.1029/2007JD008646, 2007. ¹⁰⁰⁰
- Fukuta, N. and Takahashi, T.: The Growth of Atmospheric Ice Crystals: A Summary of Findings in Vertical Supercooled Cloud Tunnel Studies, *JOURNAL OF THE ATMOSPHERIC SCIENCES*, 56, 1963–1979, doi:10.1175/1520-0469(1999)056(1963:TGOAIC)2.0.CO;2, 1999. ¹⁰⁰⁵
- Hartmann, S., Augustin, S., Clauss, T., Wex, H., Santl-Temkiv, T., Voigtlander, J., Niedermeier, D., and Stratmann, F.: Immersion freezing of ice nucleation active protein complexes, *Atmospheric Chemistry and Physics*, 13, 5751–5766, doi:10.5194/acp-13-5751-2013, 2013. ¹⁰¹⁰
- Heymsfield, A. J. and Westbrook, C. D.: Advances in the Estimation of Ice Particle Fall Speeds Using Laboratory and Field Measurements, *JOURNAL OF THE ATMOSPHERIC SCIENCES*, 67, 2469–2482, doi:10.1175/2010JAS3379.1, 2010. ¹⁰¹⁵
- Hogan, R. J., Mittermaier, M. P., and Illingworth, A. J.: The Retrieval of Ice Water Content from Radar Reflectivity Factor and Temperature and Its Use in Evaluating a Mesoscale Model, *J. Appl. Meteor. Climatol.*, 45, 301–317, doi:10.1175/JAM2340.1, 2006. ¹⁰²⁰
- Illingworth, A. J., Hogan, R. J., O'Connor, E. J., Bouniol, D., Delanoë, J., Pelon, J., Protat, A., Brooks, M. E., Gaussiat, N., Wilson, D. R., Donovan, D. P., Baltink, H. K., van Zadelhoff, G.-J., Eastment, J. D., Goddard, J. W. F., Wrench, C. L., Haeffelin, M., Krasnov, O. A., Russchenberg, H. W. J., Piriou, J.-M., Vinit, F., Seifert, A., Tompkins, A. M., and Willén, U.: Cloudnet, *Bull. Amer. Meteor. Soc.*, 88, 883–898, doi:10.1175/JAM2340.1, 2006. ¹⁰²⁵
- Kanitz, T., Seifert, P., Ansmann, A., Engelmann, R., Althausen, D., Casiccia, C., and Rohwer, E. G.: Contrasting the impact of aerosols at northern and southern midlatitudes on heterogeneous ice formation, *Geophysical Res. Letters*, 38, L17 802, {doi:10.1029/2011GL048532}, 2011. ¹⁰³⁰
- Korolev, A. and Field, P. R.: The Effect of Dynamics on Mixed-Phase Clouds: Theoretical Considerations, *J. Atmos. Sci.*, 65, 66–86, 2008. ¹⁰³⁵
- MEYERS, M., DEMOTT, P., and COTTON, W.: NEW PRIMARY ICE-NUCLEATION PARAMETERIZATIONS IN AN EXPLICIT CLOUD MODEL, *JOURNAL OF APPLIED METEOROLOGY*, 31, 708–721, doi:10.1175/1520-0450(1992)031(0708:NPINPI)2.0.CO;2, 1992. ¹⁰⁴⁰
- Mitchell, D.: Use of mass- and area-dimensional power laws for determining precipitation particle terminal velocities, *JOURNAL OF THE ATMOSPHERIC SCIENCES*, 53, 1710–1723, doi:10.1175/1520-0469(1996)053(1710:UOMAAD)2.0.CO;2, 1996. ¹⁰⁴⁵
- Murray, B. J., O'Sullivan, D., Atkinson, J. D., and Webb, M. E.: Ice nucleation by particles immersed in supercooled cloud droplets, *Chem. Soc. Rev.*, 41, 6519–6554, 2012. ¹⁰⁵⁰
- O'Sullivan, D., Murray, B. J., Malkin, T. L., Whale, T. F., Umo, N. S., Atkinson, J. D., Price, H. C., Baustian, K. J., Browse, J., and Webb, M. E.: Ice nucleation by fertile soil dusts: relative importance of mineral and biogenic components, *Atmospheric Chemistry and Physics*, 14, 1853–1867, doi:10.5194/acp-14-1853-2014, 2014. ¹⁰⁵⁵
- Petzold, A., Fiebig, M., Flentje, H., Keil, A., Leiterer, U., Schroder, F., Stifter, A., Wendisch, M., and Wendling, P.: Vertical variability of aerosol properties observed at a continental site during the Lindenberg Aerosol Characterization Experiment (LACE 98), *JOURNAL OF GEOPHYSICAL RESEARCH-ATMOSPHERES*, 107, doi:10.1029/2001JD001043, 2002. ¹⁰⁶⁰
- Phillips, V. T. J., DeMott, P. J., and Andronache, C.: An empirical parameterization of heterogeneous ice nucleation for multiple chemical species of aerosol, *J. Atmos. Sci.*, 65, 2757–2783, 2008. ¹⁰⁶⁵
- Pospichal, B., Kilian, P., and Seifert, P.: Performance of cloud liquid water retrievals from ground-based remote sensing observations over Leipzig, in: *Proceedings of the 9th International Symposium on Tropospheric Profiling (ISTP)*, 2012. ¹⁰⁷⁰
- Shipway, B. J. and Hill, A. A.: Diagnosis of systematic differences between multiple parametrizations of warm rain microphysics using a kinematic framework, *Q. J. R. Meteorol. Soc.*, 138, 2196–2211, doi:10.1002/qj.1913, 2012. ¹⁰⁷⁵
- Simmel, M. and Wurzler, S.: Condensation and activation in sectional cloud microphysical models, *ATMOSPHERIC RE-* ¹⁰⁸⁰

- SEARCH, 80, 218–236, doi:10.1016/j.atmosres.2005.08.002, 2006.
- 1015 Smith, A. J., Larson, V. E., Niu, J., Kankiewicz, J. A., and Carey, L. D.: Processes that generate and deplete liquid water and snow in thin midlevel mixed-phase clouds, *J. Geophys. Res.*, 114, D12 203, doi:10.1029/2008JD011531, 2009.
- 1020 Sun, J., Ariya, P. A., Leighton, H. G., and Yau, M. K.: Modeling Study of Ice Formation in Warm-Based Precipitating Shallow Cumulus Clouds, *JOURNAL OF THE ATMOSPHERIC SCIENCES*, 69, 3315–3335, doi:10.1175/JAS-D-11-0344.1, 2012.
- 1025 Warren, S. G., Hahn, C. J., London, J., Chervin, R. M., and Jenne, R.: Global distribution of total cloud cover and cloud type amount over land, Tech. Rep. Tech. Note TN-317 STR, NCAR, 1998a.
- Warren, S. G., Hahn, C. J., London, J., Chervin, R. M., and Jenne, R.: Global distribution of total cloud cover and cloud type amount over the ocean, Tech. Rep. Tech. Note TN-317 STR, NCAR, 1998b.
- 1030 Wegener, A.: *Thermodynamik der Atmosphäre*, J. A. Barth Verlag, 1911.
- Westbrook, C. D., Hogan, R. J., and Illingworth, A. J.: The capacitance of pristine ice crystals and aggregate snowflakes, *JOURNAL OF THE ATMOSPHERIC SCIENCES*, 65, 206–219, doi:10.1175/2007JAS2315.1, 2008.
- ~~Petzold, A., Fiebig, M., Flentje, H., Keil, A., Leiterer, U., Schroder, F., Stifter, A., Wendisch, M., and Wendling, P.: Vertical variability of aerosol properties observed at a continental site during the Lindenberg Aerosol Characterization Experiment (LACE-98), *JOURNAL OF GEOPHYSICAL RESEARCH-ATMOSPHERES*, 107, , 2002.~~
- ~~Phillips, V. T. J., DeMott, P. J., and Andronache, C.: An empirical parameterization of heterogeneous ice nucleation for multiple chemical species of aerosol, *J. Atmos. Sci.*, 65, 2757–2783, 2008.~~
- ~~Pospichal, B., Kilian, P., and Seifert, P.: Performance of cloud liquid water retrievals from ground-based remote sensing observations over Leipzig, in: *Proceedings of the 9th International Symposium on Tropospheric Profiling (ISTP)*, 2012.~~
- ~~Simmel, M. and Wurzler, S.: Condensation and activation in sectional cloud microphysical models, *ATMOSPHERIC RESEARCH*, 80, 218–236, , 2006.~~
- ~~Sun, J., Ariya, P. A., Leighton, H. G., and Yau, M. K.: Modeling Study of Ice Formation in Warm-Based Precipitating Shallow Cumulus Clouds, *JOURNAL OF THE ATMOSPHERIC SCIENCES*, 69, 3315–3335, , 2012.~~
- 1060 ~~Westbrook, C. D., Hogan, R. J., and Illingworth, A. J.: The capacitance of pristine ice crystals and aggregate snowflakes, *JOURNAL OF THE ATMOSPHERIC SCIENCES*, 65, 206–219, , 2008.~~

Table 1. Overview of the model results for the dynamic sensitivity runs for the warmer case 1 (maximum values of L/FWMR/IWC: liquid/ice water mixing-ratio/content, L/IWP: liquid/ice water path, CDN: cloud drop number, IPN: Ice particle number).

run	parameter value differing from base case	<u>LW</u> <u>MR</u> - <u>LWC</u> g/kg·m ⁻³	<u>IW</u> <u>MR</u> - <u>IWC</u> 10 ⁻³ g/kg·m ⁻³	LWP g/m ²	IWP 10 ⁻³ g/m ²	CDN cm ⁻³	IPN l ⁻¹
W_base	—	<u>0.277</u> - <u>0.355</u>	<u>0.304</u> - <u>0.379</u>	41.33	62.27	46.89	0.0197
W_h38	$h_{bot} = 3800$ m	<u>0.332</u> - <u>0.426</u>	<u>0.329</u> - <u>0.408</u>	57.05	73.11	48.63	0.0235
W_h40	$h_{bot} = 4000$ m	<u>0.225</u> - <u>0.289</u>	<u>0.286</u> - <u>0.357</u>	28.58	58.12	61.48	0.0240
W_h41	$h_{bot} = 4100$ m	<u>0.170</u> - <u>0.219</u>	<u>0.259</u> - <u>0.324</u>	18.23	45.81	59.53	0.0208
W_w01	$w_{ave} = 0.1$ m/s	<u>0.147</u> - <u>0.187</u>	<u>0.160</u> - <u>0.200</u>	17.41	31.73	43.36	0.0138
W_w02	$w_{ave} = 0.2$ m/s	<u>0.232</u> - <u>0.297</u>	<u>0.241</u> - <u>0.300</u>	32.86	47.18	54.57	0.0175
W_w04	$w_{ave} = 0.4$ m/s	<u>0.297</u> - <u>0.382</u>	<u>0.359</u> - <u>0.448</u>	44.48	78.25	52.66	0.0219
W_r1	stoch. realiz. r1	<u>0.261</u> - <u>0.336</u>	<u>0.254</u> - <u>0.316</u>	40.32	54.85	64.26	0.0163
W_r3	stoch. realiz. r3	<u>0.296</u> - <u>0.381</u>	<u>0.252</u> - <u>0.314</u>	42.88	54.48	43.03	0.0167
W_r4	stoch. realiz. r4	<u>0.269</u> - <u>0.346</u>	<u>0.197</u> - <u>0.245</u>	40.91	46.93	47.42	0.0151

Table 2. Overview of the model results for the dynamic sensitivity runs for the colder case 2 (maximum values of L/FWMR/IWC: liquid/ice water mixing-ratio/content, L/IWP: liquid/ice water path, CDN: cloud drop number, IPN: Ice particle number).

run	parameter value differing from base case	<u>LW</u> <u>MR</u> - <u>LWC</u> g/kg·m ⁻³	<u>IW</u> <u>MR</u> - <u>IWC</u> g/kg·m ⁻³	LWP g/m ²	IWP g/m ²	CDN cm ⁻³	IPN l ⁻¹
C_base	—	<u>0.215</u> - <u>0.377</u>	<u>0.026</u> - <u>0.041</u>	29.35	10.71	70.56	0.462
C_h70	$h_{bot} = 7000$ m	<u>0.258</u> - <u>0.452</u>	<u>0.030</u> - <u>0.048</u>	43.06	11.34	71.33	0.432
C_h72	$h_{bot} = 7200$ m	<u>0.169</u> - <u>0.296</u>	<u>0.022</u> - <u>0.035</u>	18.71	10.11	90.51	0.396
C_h73	$h_{bot} = 7300$ m	<u>0.122</u> - <u>0.215</u>	<u>0.018</u> - <u>0.028</u>	10.54	9.27	77.61	0.337
C_w01	$w_{ave} = 0.1$ m/s	<u>0.126</u> - <u>0.219</u>	<u>0.025</u> - <u>0.040</u>	17.19	8.01	76.98	0.292
C_w02	$w_{ave} = 0.2$ m/s	<u>0.181</u> - <u>0.316</u>	<u>0.027</u> - <u>0.044</u>	25.89	9.42	74.40	0.415
C_w04	$w_{ave} = 0.4$ m/s	<u>0.229</u> - <u>0.402</u>	<u>0.028</u> - <u>0.045</u>	30.58	11.85	98.37	0.439
C_r1	stoch. realiz. r1	<u>0.209</u> - <u>0.366</u>	<u>0.014</u> - <u>0.023</u>	29.37	6.57	86.64	0.257
C_r3	stoch. realiz. r3	<u>0.228</u> - <u>0.399</u>	<u>0.029</u> - <u>0.046</u>	30.22	9.95	79.65	0.341
C_r4	stoch. realiz. r4	<u>0.213</u> - <u>0.373</u>	<u>0.031</u> - <u>0.049</u>	29.53	8.33	95.89	0.419

Table 3. Overview of the model results for the microphysical sensitivity runs for the warmer case 1 (maximum values of L/FWMR/IWC: liquid/ice water mixing-ratio/content, L/IWP: liquid/ice water path, CDN: cloud drop number, IPN: Ice particle number).

run	parameter value differing from base case	<u>LW</u> <u>MR</u> - <u>LWC</u> g/kg·m ⁻³	<u>IW</u> <u>MR</u> - <u>IWC</u> 10 ⁻³ g/kg·m ⁻³	LWP g/m ²	IWP g/m ²	CDN cm ⁻³	IPN l ⁻¹
W_in6	$N_{AP,r>250nm} = 10^6$ kg ⁻¹	<u>0.276</u> - <u>0.354</u>	<u>0.496</u> - <u>0.619</u>	41.31	0.10	46.69	0.0296
W_in7	$N_{AP,r>250nm} = 10^7$ kg ⁻¹	<u>0.276</u> - <u>0.354</u>	<u>0.801</u> - <u>1.000</u>	41.24	0.17	41.61	0.0450
W_col	ice shape: columns	<u>0.275</u> - <u>0.353</u>	<u>1.467</u> - <u>1.830</u>	41.20	0.27	42.90	0.0257
W_pla	ice shape: plates	<u>0.275</u> - <u>0.353</u>	<u>2.285</u> - <u>2.850</u>	41.13	0.45	43.41	0.0267

Fig. 1. Lidar and radar observations on 17 September 2011 (case 1). Left: Ice-water-content-IWC, right: Lidar range-corrected 1064 nm signal (in logarithmic scale, arbitrary units a. u.), right: radar reflectivity. The dashed box denotes the region for which case 1 observations are shown in the following figures.

Table 4. Overview of the model results for the microphysical sensitivity runs for the colder case 2 (maximum values of L/IWMP/IWC: liquid/ice water ~~mixing-ratio~~content, L/IWP: liquid/ice water path, CDN: cloud drop number, IPN: Ice particle number).

run	parameter value differing from base case	LWMP-LWC g/kg-m ³	IWMP-IWC g/kg-m ³	LWP g/m ²	IWP g/m ²	CDN cm ⁻³	IPN l ⁻¹
C.in6	$N_{AP,r>250nm} = 10^6 \text{ kg}^{-1}$	0.128 <u>0.224</u>	0.089 <u>0.140</u>	13.09	34.75	80.29	1.380
C.in7	$N_{AP,r>250nm} = 10^7 \text{ kg}^{-1}$	0.021 <u>0.036</u>	0.265 <u>0.446</u>	2.58	57.98	46.67	5.208
C.col	ice shape: columns	0.135 <u>0.237</u>	0.139 <u>0.223</u>	14.33	46.78	78.40	0.462
C.col.in4	ice shape: columns, $N_{AP,r>250nm} = 10^4 \text{ kg}^{-1}$	0.216 <u>0.378</u>	0.048 <u>0.076</u>	30.01	14.93	74.75	0.139
C.pla	ice shape: plates	0.104 <u>0.182</u>	0.183 <u>0.294</u>	9.94	57.11	39.41	0.472
C.pla.in4	ice shape: plates, $N_{AP,r>250nm} = 10^4 \text{ kg}^{-1}$	0.207 <u>0.362</u>	0.064 <u>0.102</u>	27.80	19.21	74.44	0.129

Fig. 2. ~~Vertical velocity-Cloudnet derived water contents~~ for case 1. Left: ~~derived from radar observations (valid for large particles; virgae)~~ Liquid water content, right: ice water content (both in logarithmic scale).

Fig. 3. Vertical velocity for case 1. Left: derived from lidar (valid for more numerous smaller droplets at cloud base), right: derived from radar observations (valid for large particles; virgae).

Fig. 4. Lidar and radar observations on 2 August 2012 (case 2). Left: Ice-water content IWC, right: 1064-532 nm attenuated backscatter coefficient, right: radar reflectivity.

~~Vertical velocity for case 2, derived from radar observations.~~

Fig. 5. Cloudnet derived water contents for case 2. Left: Liquid water content, right: ice water content (both in logarithmic scale).

Fig. 6. Vertical velocity for case 2. Left: derived from lidar (valid for more numerous smaller droplets at cloud base), right: derived from radar observations (valid for large particles; virgae).

Fig. 7. Vertical profiles of temperature (left) and relative humidity (right) with respect to liquid water (full lines) and ice (dashed lines) based on a radiosonde observation (Meiningen) for case 1 (black) and from GDAS (grid point Leipzig) for case 2 (red).

Fig. 8. Vertical velocity field of the inner cylinder for case 1. Left: Height dependence (red line) and temporal evolution of one realization of the stochastic vertical velocity field (black line) for $w_{ave} = 0.3 \text{ m/s}$ at h_{mid} . Right: Histogram of velocity field. Vertical velocity for case 2 is identical but for heights between 7100 m and 7700 m.

Fig. 9. Liquid LWC (contours) and ice-water-mixing-ratio-LWC (colours, logarithmic scale) for case 1. Comparison of different values for h_{bot} (Upper left: W_base, $h_{bot} = 3900$ m, upper right: W_h38, $h_{bot} = 3800$ m, lower left: W_h40, $h_{bot} = 4000$ m, lower right: W_h41, $h_{bot} = 4100$ m.)

Fig. 10. Liquid (lower panel) and ice water paths (upper panel) for case 1. Comparison of the different values for h_{bot} .

Fig. 11. Liquid-LWC (contours) and ice-water-mixing-ratio-IWC (colours, logarithmic scale) for case 1. Comparison of different average updraft velocities w_{ave} (Upper panel: Left: W_w01, $w_{ave} = 0.1$ m/s, right: W_w04, $w_{ave} = 0.4$ m/s.) and different stochastic realizations (Lower: Left: W_r1, r1, right: W_r4, r4).

Fig. 12. Liquid (lower panels) and ice water paths (upper panels) for case 1. Comparison of the different values for w_{ave} (left) and the different stochastic realizations (right).

Fig. 13. Liquid-LWC (contours) and ice-water-mixing-ratio-IWC (colours, logarithmic scale) for case 2. Comparison of different values for h_{bot} (Upper left: C_base, $h_{bot} = 7100$ m, upper right: C_h70, $h_{bot} = 7000$ m, lower left: C_h72, $h_{bot} = 7200$ m, lower right: C_h73, $h_{bot} = 7300$ m.)

Fig. 14. Liquid (lower panel) and ice water paths (upper panel) for case 2. Comparison of the different values for h_{bot} .

Fig. 15. Liquid-LWC (contours) and ice-water-mixing-ratio-IWC (colours, logarithmic scale) for case 2. Comparison of different average updraft velocities w_{ave} (Upper: Left: C_w01, $w_{ave} = 0.1$ m/s, right: C_w04, $w_{ave} = 0.4$ m/s) and the different stochastic realizations (Lower: Left: C_r1, r1, right: C_r4, r4).

Fig. 16. Liquid (lower panels) and ice water paths (upper panels) for case 2. Comparison of the different values for w_{ave} (left) and the different stochastic realizations (right).

Fig. 17. Liquid-LWC (colours) and ice-water-mixing-ratio-IWC (contours, logarithmic scale) for case 1 (W_in6, left) and case 2 (C_in6, right). Enhancing IN by increasing $N_{AP,r>250nm}$ by a factor of 10.

Fig. 18. Liquid-LWC (colourscontours) and ice-water-mass-IWC per bin (contourscolours, both logarithmic scale) for case 1 (upper panel) and case 2 (lower panel) for the respective base case (left) and the case with enhanced IN number (right; in6) after 16 and 17 minutes model time, respectively, corresponding to the IWP maximum of the base case runs.

Fig. 19. Liquid (lower panel) and ice water paths (upper panel) for case 1 (left) and case 2 (right). Comparison of the sensitivities with respect to IN number and ice particle shape.

Fig. 20. Liquid-LWC (colourscontours) and ice-water-mixing-ratio-IWC (contourscolours, logarithmic scale). Results for changing ice particle shape to hexagonal columns for case 1 (W_col, left) and case 2 (C_col, right).

Fig. 21. Liquid-LWC (colourscontours) and ice water mass per bin (contourscolours, both logarithmic scale) for case 1 (upper panel) and case 2 (lower panel) assuming columns as ice particle shape at IWP maximum of the respective base case (left) and at IWP of the run (right).