
Atmos. Chem. Phys. Discuss., 15, C270–C273, 2015 www.atmos-chem-phys-discuss.net/15/C270/2015/ © Author(s) 2015. This work is distributed under the Creative Commons Attribute 3.0 License.



ACPD 15, C270–C273, 2015

> Interactive Comment

Interactive comment on "Influence of along-valley terrain heterogeneity on exchange processes over idealized valleys" *by* J. S. Wagner et al.

Anonymous Referee #1

Received and published: 18 February 2015

General evaluation

The authors present idealized simulations dealing with the dependence of thermally driven valley winds on various aspects of the valley geometry. The study is interesting and provides new scientific results, is technically well done (with some exceptions mentioned below) and is presented in a far-above-average writing style. I therefore recommend acceptance for publication subject to minor revisions.



Interactive Discussion



Major comments

1. The literature review is rather incomplete, giving the wrong impression that there has been no scientific work on valley winds before 2000. In fact, there have been a lot of important studies on valley winds in the 1980's and 1990's, e.g. by Egger and Whiteman. At least a few seminal papers should be mentioned and put into context with more recent work.

2. At the end of the introduction, it does not become sufficiently clear which aspects of the present work are entirely new and/or an extension of previous work. This is certainly not the first idealized study addressing the impact of a sloping valley floor or a varying valley cross-section on the thermal valley wind circulation.

3. This work is one of many idealized process studies being motivated by the need of improving the physics parameterizations of our numerical models without sufficiently addressing this issue later in the discussion and/or the conclusions. In their concluding sentence, the authors just state "Future boundary layer parameterization schemes ... should consider these valley geometry parameters besides other effects ...".

Well ... how is this supposed to work in practice, in particular at model resolutions nowadays used for regional weather forecasting, which marginally resolve large valleys, like e.g. the Inn Valley, and heavily under-resolve smaller tributaries? What actually would need to be parameterized is the difference between the thermal circulation the model produces at its operational resolution and the circulation it produces at a much higher resolution representing the orography reasonably accurate. No one knows if this is possible in a generic way. Some more substantiated thoughts on this issue would be very welcome.

ACPD 15, C270–C273, 2015

> Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Minor comments

p. 424, ln 27: The way by which the contribution of various parameters of the model orography to the valley wind intensity is computed is a bit confusing. Intuitively, I would say that 2.62*REF / 1.62*REF yields a factor of about 1.6, rather than taking 262% - 162% (or 162% - 62% after subtracting REF) to obtain 100% or a factor of 2.

p. 425, top: The authors report that a wide valley forces substantially weaker valley winds than a narrower valley (comparison REF-W30) and argue that this because of the smaller fractional volume reduction. While the volume effect is likely the most important reason, I think that the depth/width ratio of the valley also plays a role. What happens when widening the valley and the mountain ridge in between by the same factor (by enlarging the model domain)?

p. 425, In. 7: "overestimation" \rightarrow "discrepancy"

p. 432, In. 14: Bad wording. Suggestion: "Simulations with inclined valley floors reveal a significant increase of ..."

Fig. 2: Please indicate the contour interval in the caption.

Fig. 4: Please use the same θ contour interval in all panels and indicate it in the caption. Moreover, the thick contour lines for along-valley wind speed are barely visible. My suggestion would be to make them green, and to use violet or purple for the present green dashed line.

Fig. 5: Please use the same θ contour interval in all panels and indicate it in the caption.

Fig. 6, 7, 8, 9, 13, 14: The lines are difficult to distinguish. At least, SL should be dash-dotted in order to be clearly distinguishable from Ixxx.

In addition, Figs. 4, 5, 9 and 10 should be enlarged. In particular Fig. 9.

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Interactive comment on Atmos. Chem. Phys. Discuss., 15, 415, 2015.

ACPD

15, C270–C273, 2015

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

