

[Interactive
Comment](#)

Interactive comment on “A comparison of chemical mechanisms using Tagged Ozone Production Potential (TOPP) analysis” by J. Coates and T. M. Butler

J. Coates and T. M. Butler

jane.coates@iass-potsdam.de

Received and published: 23 July 2015

We would like to thank Dr Jenkin for the helpful review that will enable us to improve our manuscript; our responses to the review points and comments are found below.

Review Point 1: Given that some inventory speciations contain several hundred VOCs for typical urban sources, it could be argued that this is already a substantially lumped representation – or maybe more correctly an incomplete speciation, as it is based on reported measurements of a subset of species (although probably the generally more important ones). As a result, the reference MCM simulations are themselves already

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

[Discussion Paper](#)



a reduction, using only a subset of the mechanism. The numbers of species and reactions listed for MCM v3.2 and MCM v3.1 (and CRI v2) in Table 1 should therefore probably more correctly correspond to the VOC speciation, as many species in the complete mechanisms are not participating in the chemistry. These can be obtained using the subset mechanism assembling facilities at the relevant MCM and CRI websites (see end of review). If the authors also wish to keep the existing full mechanism numbers, the subset numbers could be included in a footnote.

Author Response: We agree with Dr Jenkin that it would be more complete stating the number of organic reactions used in our study for each mechanism listed in Table 1. We have updated Section 2.1 (page 12395, line 10) of the manuscript stating : *We used a subset of each chemical mechanism containing all the reactions needed to fully describe the degradation of the VOC in Table 2.*

Accordingly, we have updated Table 1 in the manuscript with the number of organic species and reactions need to fully describe the degradation of the VOC used in the study for each mechanism. We have retained the information on the total number of organic reactions in each mechanism by updating the mechanism description in Section 2.1 to include this information.

Review Point 2: On page 12400 (line 12) the high Ox formation from propane degradation in RACM2 is attributed to the mechanism species HC3 producing about 17 times the amount of acetaldehyde that is produced from propane in MCM v3.2. This high ratio arises because acetaldehyde is not a significant first generation product of propane degradation, and therefore not formed in MCM v3.2 until the second-generation chemistry (specifically the further oxidation of the relatively minor product, propanal). Acetaldehyde is therefore formed in RACM2 instead of other products formed in MCM v3.2 (mainly acetone, and some propanal). Acetone, has a low OH reactivity (and photolysis rate) and is not significantly oxidised on the several day timescale of these calculations, thereby largely halting Ox formation after the first-generation chemistry. I suggest it is therefore the failure of RACM2 to represent

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

the high yield of an unreactive product that results in its overestimate in Ox. Although the high relative formation of acetaldehyde on day 1 with RACM2 and MCM v3.2 is one consequence of this, it is not itself the source of the Ox overestimate with RACM2.

Author Response: We would first like to thank Dr Jenkin for further insight into the differences in Ox production between RADM2 and MCM v3.2. As Dr Jenkin suggests, the yield of the less reactive ketone products during propane degradation is lower than in the MCM v3.2. We have updated Section 3.1 of the manuscript (page 12400, line 11) as follows: *Propane is represented as HC3 in RADM2 (Stockwell et al., 1990) and the degradation of HC3 has a lower yield of the less reactive ketones compared to the MCM. The further degradation of ketones hinders Ox production due to the low OH reactivity and photolysis rate of ketones. Secondary degradation of HC3 proceeds through the degradation of acetaldehyde (CH₃CHO) propagating Ox production through the reactions of CH₃CO₃ and CH₃O₂ with NO. Thus the lower ketone yields leads to increased Ox production from propane degradation in RADM2 compared to the MCM v3.2.*

Section 3.2.2 (page 12403, line 26) was also updated as follows: *The secondary chemistry of HC3 is tailored to produce O_x from these different VOC and differs from alkane degradation in the MCM v3.2 by producing less ketones in RADM2.*

Review Point 2-Cont: On page 12402 (line 28), where the second day maximum in Ox from toluene degradation with RACM2 and CRI v2 is attributed to "...increased C₂H₅O₂ production from degradation of unsaturated dicarbonyls; C₂H₅O₂ is not produced during degradation of unsaturated dicarbonyls in the MCM v3.2." I am particularly familiar with CRI v2, and I agree that a likely contributor to the discrepancy is that the formation of the (relatively reactive) unsaturated dicarbonyls (UDCARB8 and UDCARB11) is too efficient. C₂H₅O₂ is indeed used as a representative peroxy radical, formed from one channel of the oxidation of UDCARB8 - this being the surrogate for butenedial (MALDIAL) in MCM v3.2. However, MALDIAL is also oxidised to peroxy radicals (MALDIALO₂ and MALDIALCO₃) in MCM v3.2, with the fraction not leading to anhy-

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

drone formation being represented by $C_2H_5O_2$ in CRI v2. I therefore do not think this is an unreasonable assignment (note that contributions of MALDIALO₂ + NO and MALDIALCO₃ + NO are probably hidden within the large “production others” category for MCM v3.2). It is more that the formation of the unsaturated dicarbonyls is too efficient in CRI v2, and that their degradation produces Ox, regardless of which peroxy radicals are used as representatives. Although the fluxes through the reactions of NO with the specific peroxy radicals ($C_2H_5O_2$ from UDCARB8 and RN10O₂ from UDCARB11) are how this is quantified in the present study, I think that highlighting increased production of $C_2H_5O_2$ as the sole specific cause is not particularly instructive, as it is once again a consequence of the real cause.

Author Response: Again, we would like to thank Dr Jenkin for his insight into the approach of toluene degradation in CRI v2. Based on this insight, we have revised Section 3.2.1 (page 12402, line 29) of the manuscript (third paragraph from the end of the Section) to: *The second day maximum of O_x production in CRI v2 and RACM2 from toluene degradation results from more efficient production of unsaturated dicarbonyls than the MCM v3.2. The degradation of unsaturated dicarbonyls produces peroxy radicals such as $C_2H_5O_2$ which promote O_x production via reactions with NO.*

Minor Comments 1: General “Volatile organic compounds” seems to be abbreviated as either “VOC” or “VOCs” at different points in the manuscript. Given that the original definition on line 3 of the Introduction is “VOCs”, I would suggest using this consistently throughout, unless talking about an individual VOC.

Author Response: We have corrected the manuscript to defined volatile organic compounds as VOC (page 12390, line 22), and updated the manuscript to use this acronym consistently. Changes were made to page 12391, line 22; page 12397 lines 3 and 10; and to the caption of Table 3.

Minor Comments 2: Page 12391, line 22: perhaps it should be stated that VOCs are oxidised mainly by reaction with the OH radical, to acknowledge the existence of other

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

initiation pathways.

Minor Comments 3: Page 12392: perhaps it should be clarified that Reaction (R4) specifically illustrates the abstraction of H from a VOC by reaction with OH, as occurs exclusively for alkanes. The main routes for the reactions of OH with alkenes and aromatics proceed by OH addition.

Author Response to Minor Comments 2 and 3: We have updated page 12391, line 22 to: *VOC (RH) are mainly oxidised in the troposphere by the hydroxyl radical (OH) forming peroxy radicals (RO₂) in the presence of O₂. For example, (Reaction R4) describes the OH-oxidation of alkanes proceeding through abstraction of an H from the alkane. In high-NO_x conditions,*

Minor Comments 4: Page 12394, line 13: As stated, the full CRI v2 does lump degradation products into common representatives. Although not used in the present study, its further reduced variants (e.g. CRI v2-R5) also lump emitted VOCs using POCP as a criterion (Watson et al., 2008), so that they are subsets of the full mechanism. As indicated in comment 1 above, the present work also uses a subset of the full mechanism, so I'm not sure that use of the "full CRI" can be claimed on page 12395, line 19.

Author Response: Our subset of the CRI v2 was taken from the full CRI not any of the reduced variants of the CRI that use further reduction techniques as described in Watson et al. (2008). We have updated page 12395, line 19 as follows: *The CRI v2 is available in more than one reduced variant, described in Watson et al. (2008). We used a subset of the full version of the CRI v2 (<http://mcm.leeds.ac.uk/CRI>).*

Minor Comments 5: Page 12398, line 9: The use of a family of Ox species is a sensible approach. However, it might be worth giving a formal definition of "other species involved in fast cycling with NO₂", as those shown have a wide range of cycling lifetimes. Are PANs sufficiently short-lived?

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

Author Response: In our study, we simulate the conditions within the planetary boundary layer thus PAN chemistry is dominated by its production and thermal decomposition. Ox budgets when not including PAN as part of the Ox family are thus dominated by these cycles of PAN formation and thermal decomposition. For this reason we include PANs as part of the Ox family.

Other Comments 1: Page 12395, lines 13-22: MCM v3.2 is used as the reference mechanism in this study, with MCM v3.1 also considered for completeness, because it was the reference for the original development of CRI v2 (Jenkin et al., 2008). Because an overview description of MCM v3.2 has never been published in the open literature, I provide here a short summary of the updates. This is mainly for information, and not necessarily for reproduction in the paper, unless deemed helpful by the authors.

Because there is no overview publication, the authors have used the citation "Rickard et al. (2015)" for the MCM v3.2 website. I suggest the author list of this citation is expanded to include those listed as "current contributors" on the citation tab of the MCM v3.2 website (<http://mcm.leeds.ac.uk/MCMv3.2/contributors.htm>).

Author Response: We would like to thank Dr Jenkin for providing the updates from the MCM v3.1 to MCM v3.2, and we shall update the reference to the MCM v3.2 to (Rickard et al., 2015) including those listed as current contributors to the MCM v3.2.

Other Comments 2: I note from the response to another reviewer (Coates and Butler, Atmos. Chem. Phys. Discuss., 15, C3816, 2015) that the authors are proposing to include references to the studies of Bloss et al. (2005) and Pinho et al. (2005) to illustrate that the MCM has had difficulties in reproducing the results of chamber studies for aromatic VOCs and isoprene. Whilst I agree with this for aromatics, the main conclusion of Pinho et al. (2005) was that the MCM v3 isoprene scheme (written in 2001) performed very well. The major factor responsible for deviations in performance of the MCM v3 scheme from the SAPRC chamber data was the absence of the reaction of O(3P) with isoprene in MCM v3, this reaction being insignificant under atmospheric conditions. A

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

number of other less important refinements were also identified by Pinho et al. (2005), and these were all implemented long before release of MCM v3.2 in 2011. I therefore think it is misleading to report that the MCM has had difficulties in reproducing the results of traditional chamber studies for isoprene.

Author Response: Based upon the comments of Dr Jenkin, we shall not include the Pinho et al. (2005) study as an example of the MCM having difficulties reproducing chamber study results. We have updated page 12395, line 15 of the manuscript accordingly: *The MCM v3.2 is the reference mechanism in this study due to its level of detail (16 349 organic reactions). Despite this level of detail, the MCM had difficulties in reproducing the results of chamber study experiments involving aromatic VOC (Bloss et al., 2005).*

References

- Bloss, C., Wagner, V., Jenkin, M. E., Volkamer, R., Bloss, W. J., Lee, J. D., Heard, D. E., Wirtz, K., Martin-Reviejo, M., Rea, G., Wenger, J. C., and Pilling, M. J.: Development of a detailed chemical mechanism (MCMv3.1) for the atmospheric oxidation of aromatic hydrocarbons, *Atmos. Chem. Phys.*, 5, 641–664, 2005.
- Jenkin, M. E., Watson, L. A., Utembe, S. R., and Shallcross, D. E.: A Common Representative Intermediates (CRI) mechanism for VOC degradation. Part 1: Gas phase mechanism development, *Atmos. Environ.*, 42, 7185–7195, 2008.
- Pinho, P. G., Pio C. A., and Jenkin M. E.: Evaluation of isoprene degradation in the detailed tropospheric chemical mechanism, MCM v3, using environmental chamber data, *Atmospheric Environment*, 39, 7, 1303–1322, 2005.
- Rickard, A., Young, J., Pilling, M., Jenkin. M., Pascoe, S., and Saunders S.: The Master Chemical Mechanism Version MCM v3.2, available at: <http://mcm.leeds.ac.uk/MCMv3.2/> (last access: 09 July 2015), 2015.
- Watson, L. A., Shallcross, D. E., Utembe, S. R., and Jenkin, M. E.: A Common Representative Intermediates (CRI) mechanism for VOC degradation. Part 2: Gas phase mechanism reduction, *Atmospheric Environment*, 42, 7196–7204, 2008.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Interactive comment on Atmos. Chem. Phys. Discuss., 15, 12389, 2015.

ACPD

15, C5144–C5151, 2015

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C5151

