

Supplementary Material

Summited to: Atmospheric Chemistry and Physics (ACP)

Title: Quantitative Assessment of Atmospheric Emissions of Toxic Heavy Metals from Anthropogenic Sources of China: Historical trend, spatial variation distribution, uncertainties and control policies

Authors: H. Z. Tian¹, C. Y. Zhu¹, J. J. Gao¹, K. Cheng^{1,2}, J. M. Hao³, K. Wang¹, S. B. Hua¹, Y. Wang¹, J. R. Zhou¹

¹State Key Joint Laboratory of Environmental Simulation & Pollution Control, School of Environment, Beijing Normal University, Beijing 100875, China

²School of Environment, Henan Normal University, Xinxiang, Henan Key Laboratory for Environmental Pollution Control, Key Laboratory for Yellow River and Huai River Water Environment and Pollution Control, Ministry of Education, Xinxiang 453007, China

³State Environmental Protection Key Laboratory of Sources and Control of Air Pollution Complex, School of Environment, Tsinghua University, Beijing 10084, China

Correspondence to: H.Z. Tian (hztian@bnu.edu.cn)

Number of pages: 51

Number of section: 3

Number of tables: 22

Number of figures: 8

Section List

Section S1. Mathematical description of bootstrap simulation Table List

Section S2. Removal efficiencies of 12 HMs through coal cleaning and coke process

Section S3. Temporal variation trends of HM emissions from other primary anthropogenic sources

31	Table List
32	Table S1. Summary of heavy metal species and the associated emission sources categories
33	Table S2. The emission source classification by coal combustion sector
34	Table S3. Statistical parameters of bootstrap mean contents of Hg, As, Se, Pb, Cd, Cr, Ni and Sb
35	in produced coal by provinces
36	Table S4. Mn Content of Raw Coal as Mined in China, by Province
37	Table S5. Co Content of Raw Coal as Mined in China, by Province
38	Table S6. Cu Content of Raw Coal as Mined in China, by Province
39	Table S7. Zn Content of Raw Coal as Mined in China, by Province
40	Table S8. Averaged concentrations of Hg, As, Se, Pb, Cd, Cr, Ni, Sb, Mn, Co, Cu and Zn in coals
41	as consumed by province (unit: $\mu\text{g/g}$).
42	Table S9. Removal efficiencies of 12 HMs through coal cleaning and coke process
43	Table S10. Release rates of Mn, Co, Cu and Zn from coal-fired facilities.
44	Table S11. Removal efficiencies of Mn, Co, Cu and Zn by different control devices
45	Table S12. TSP emission factors for vehicle brake and tyre wear
46	Table S13. Composition of tyre and brake wear in term of heavy metals, ppm
47	Table S14. Parameter values used in the transformed normal distribution function computation of
48	the variation of heavy metals emission factors over time
49	Table S15. Abatement efficiencies for nonferrous metals smelting
50	Table S16. Heavy metals emission factors for non-coal combustion sources: temporal, and process
51	variations
52	Table S17. Abatement efficiencies for iron and steel production

53 **Table S18.** The emission limits of air pollutants of relative industrial process in China and
54 developed regions, mg/m³

55 **Table S19.** Data source of activity data for the main heavy metals emitting sectors in China

56 **Table S20.** Selected parameters showing method and assumption for uncertainty analysis

57 **Table S21.** The atmospheric concentrations of As, Pb, Cr and Cu in PM_{2.5} in Beijing during 2000
58 to 2012

59 **Table S22.** Uncertainties in the sectoral emissions of heavy metals in China in 2010

60

61 **Figure List**

62 **Fig. S1.** Coal consumption by different sectors in China, 2000-2012

63 **Fig. S2.** The output of nonferrous metals in China, 2000-2012

64 **Fig. S3.** The output of pig iron and steel products in China, 2000-2012

65 **Fig. S4.** The output of non-metallic minerals manufacturing in China, 2000-2012

66 **Fig. S5.** The number of civil vehicles in China, 2000-2012

67 **Fig. S6.** Comparison between historical HM (As, Cr, Pb, and Cu) emissions and temporal
68 variation of atmospheric concentrations in PM_{2.5} in Beijing during 2000-2012

69 **Fig. S7.** The distribution of point sources in China

70 **Fig. S8.** The uncertainty bounds for China's anthropogenic atmospheric emissions of twelve HMs
71 during 1949 to 2012

Section

Section S1. Mathematical description of bootstrap simulation

Bootstrap is a numerical technique originally developed for the purpose of estimating confidence intervals for statistics. This method can provide solutions of confidence intervals in situations where exact analytical solutions may be unavailable and in which approximate analytical solutions are inadequate. Confidence intervals for a statistic are inferred from its sampling distribution. For example, the 2.5th and 97.5th percentiles of sampling distribution enclose a 95% confidence interval. The brief mathematical description of bootstrap simulation is as follows:

A random sample $X=(x_1, x_2, \dots, x_n)$ of size n is observed from a completely unspecified probability distribution F . The sampling distribution $R(X, F)$ is the function of X and F . Assume $\theta=\theta(F)$ is a parameter of F , F_n is the empirical distribution function of X , $\hat{\theta} = \hat{\theta}(F_n)$ is the estimator of θ , and the estimation error can be expressed as:

$$R(X, F) = \hat{\theta}(F_n) - \theta(F) \triangleq T_n \quad (1)$$

The basic steps of computing the distribution $R(X, F)$ by bootstrap simulation are summarized as follows:

- (1) The value of observed samples $X=(x_1, x_2, \dots, x_n)$ are finite overall samples (called original samples), $x_i \sim F(x)$, $i=1, 2, \dots, n$. The empirical distribution function of original samples is shown as:

$$F_n = \begin{cases} 0 & x < x_{(1)} \\ k/n & x_{(k)} \leq x < x_{(k+1)} \\ 1 & x \geq x_{(n)} \end{cases} \quad (2)$$

where, $x_{(1)} \leq x_{(2)} \leq \dots \leq x_{(n)}$ is the statistics of x_1, x_2, \dots, x_n sorted in ascending order.

- (2) Monte Carlo simulation is used to randomly simulate N groups of samples $x_{(j)}^* = (x_1^*, x_2^*, \dots,$

$x_n^*), j=1, 2, \dots, N$ (a very large number) from F_n , and these regeneration samples called

bootstrap samples. The generation method of empirical distribution function by Monte Carlo simulation can be expressed as: (a) generate a random integer η with independence and uniformity between 0 and M ($M \gg n$) by computer; (b) let $i = \eta \% n$, and i is the remainder of n divide η ; (c) find the sample x_i as the regeneration sample x^* in observed samples, and x^* is the needed random sample.

- (3) Calculate the statistics of bootstrap samples:

$$R^*(X^*, F_n) = \hat{\theta}(F_n^*) - \hat{\theta}(F_n) \rightarrow R_n \quad (3)$$

where, F_n^* is the empirical distribution function of bootstrap samples. As small samples

can't derive $\theta(F)$, $\hat{\theta}(F_n)$ is used to approximate it.

- (4) Use the distribution of R_n (under given situation) to simulate the distribution of T_n , say:

$\theta(F) \approx \hat{\theta} - R_n$, which can receive N numbers of $\theta(F)$. Then, the distribution and eigenvalue of unknown parameter θ can be obtained.

Section S2. Removal efficiencies of 12 HMs through coal cleaning and coke process

Some studies have reported that coal cleaning is an effective and feasible way to reduce atmospheric emissions of heavy metals before coal burning (Luttrell et al., 2000; Wang et al., 2006). By the year of 2012, only about 20.9% of total raw coal production is washed before burning, and is primarily used for coke making in iron and steel industry (NBS, 2013a). In view of the operation characteristics and the application situation of coal cleaning processes in China, we assume the average removal efficiency of Hg, As, Se, Pb, Cd, Cr, Ni, Sb, Mn, Co, Cu and Zn to be 50.0%, 54.0%, 30.0%, 36.3%, 32.2%, 58.0%, 58.5%, 35.7%, 68.2%, 39.3%, 31.8% and 48.6% (Quick and Irons, 2002; Bai, 2003; Wang et al., 2003a; Yao et al., 2012), respectively.

Due to excessive temperature range (400 °-1000 °) in coke process (Zajusz-Zubek and Koniecznyński, 2003), high emission will be found, especially for volatile substance. According to analyze the data of heavy metals discharge in coke process as described in other studies (Helble et al., 1996; Guo et al., 2002; Guo et al., 2003; Yi et al., 2007; Koniecznyński et al., 2012), we presume that 10.0% of Hg, 70.0% of As, 60.0% of Se, 68.5% of Pb, 80.0% of Cd, 76.0% of Cr, 90.3% of Ni, 30.0% of Sb, 92.4% of Mn, 92.9% of Co, 80.0% of Cu and 73.9% of Zn remains after the coking process.

Section S3. Temporal variation trends of HM emissions from other primary anthropogenic

sources

1 HM emissions from liquid fuels combustion

Although liquid fuels only take up about 8.9% of the total primary energy production and account for nearly 18.8% of total energy consumption in 2012, the liquid fuels consumption is also one of major contributors for atmospheric Ni emissions due to the relatively high content of Ni in fuel oil (Tian et al., 2012b). Furthermore, with the rapid growth of vehicle/plane populations and transport turnover (including passenger and cargo turnover), the consumptions of gasoline, diesel oil and kerosene of China have reached 116.0, 184.1 and 22.0 Mt in 2012, respectively.

Because of the large usage of leaded gasoline in China before 2001, none can afford to neglect the accumulated emissions of Pb from gasoline consumption by vehicles during 1949 to 2012, although the leaded gasoline has been forbidden to produce and use since 2001.

In this study, we estimate that the discharge of Ni from liquid fuels combustion have increased from 12.8 t in 1949 to 604.5 t in 2012. Therein, fuel oil combustion contributes over 82.1% of the total liquid fuels consumption category in 2012. Notably, the total Ni emission from liquid fuels consumption category has increased slightly (less than 2% annually) since 1980 despite of the rapid growth of distillate oils (gasoline, diesel oil, and kerosene), which is mainly because of the lower Ni content in distillate oils and relatively constant supply of fuel oil in China in the past three decades (NBS, 2013b; Wang et al., 2003b; Tian et al., 2012b).

In term of Pb emission from gasoline combustion category, the reduced lead content of gasoline is the primary reason for the sharp decrease in total Pb emissions in 1991 and 2001 (Li et al., 2012), as with national total Pb emission. For the first sharp emission decline, the total emission has decreased by 36.8% from 12 832.2 t in 1990 to 8107.5 t in 1991. For the other sharp decline, the total emission has decreased by 98.1% from 12 866.7 t in 2000 to 248.3 t in 2001. However, the Pb emissions from this category have continued to increase in the following years due to the gradually increase of gasoline consumption with the rapid growth of urban vehicle populations (please see Fig. S5).

2 HM emissions from brake and tyre wear

During the period of 1949 to 2012, the amount of civilian vehicles has increased from 0.1 million units to 109.3 million units. Furthermore, the passenger turnover of highways and freight turnover of highways have increased continuously to 1846.8 billion passenger-kilometer and 5953.5 billion ton-kilometer, respectively (NBS, 2013b). As a result, the total Pb, Cr, Sb, Mn, Cu and Zn emissions from brake and tyre wear have increased remarkably to 333.5, 124.0, 530.1, 133.8, 2720.1 and 954.7 t in 2012, respectively. Especially during 2000 to 2012, the annual growth rate of these HM emissions from brake and tyre wear is up to about 17.5%, which is closely related to the rapid growth of civilian vehicle population (see Fig. S5). For other HMs (As, Se, Cd, Ni and Co), the extraordinarily low emissions from brake and tyre wear category are estimated due to trace level of these elements in brake linings.

163 **Table**

164

165

Table S1 Summary of heavy metal species and the associated emission sources categories

Sector	Category	Hg	As	Se	Pb	Cd	Cr	Ni	Sb	Mn	Co	Cu	Zn
Coal combustion	Power plant	Raw coal	•	•	•	•	•	•	•	•	•	•	•
		Cleaned coal	•	•	•	•	•	•	•	•	•	•	•
		Briquette	•	•	•	•	•	•	•	•	•	•	•
		Coke	•	•	•	•	•	•	•	•	•	•	•
	Industrial sector	Raw coal	•	•	•	•	•	•	•	•	•	•	•
		Cleaned coal	•	•	•	•	•	•	•	•	•	•	•
		Briquette	•	•	•	•	•	•	•	•	•	•	•
		Coke	•	•	•	•	•	•	•	•	•	•	•
	Residential sector	Raw coal	•	•	•	•	•	•	•	•	•	•	•
		Cleaned coal	•	•	•	•	•	•	•	•	•	•	•
		Briquette	•	•	•	•	•	•	•	•	•	•	•
		Coke	•	•	•	•	•	•	•	•	•	•	•
	Other sector	Raw coal	•	•	•	•	•	•	•	•	•	•	•
		Cleaned coal	•	•	•	•	•	•	•	•	•	•	•
		Briquette	•	•	•	•	•	•	•	•	•	•	•
		Coke	•	•	•	•	•	•	•	•	•	•	•
Non-coal combustion	Biomass burning	Straw	•	•	•	•	•	•	•	•	•	•	•
		Wood	•	•	•	•	•	•	•	•	•	•	•
	Liquid fuel combustion	Crude oil	•	•	•	•	•	•		•	•	•	•
		Gasoline	•	•	•	•	•	•		•	•	•	•
		Diesel for stationary sources	•	•	•	•	•	•		•	•	•	•

Nonferrous smelting	Diesel for transportation	•	•	•	•	•	•	•		•	•	•	•
	Fuel oil	•	•	•	•	•	•	•		•	•	•	•
	Kerosene for stationary sources	•	•	•	•	•	•	•		•	•	•	•
	Kerosene for transportation			•	•	•	•	•		•	•	•	•
	Primary copper	•	•	•	•	•	•	•	•	•		•	•
	Secondary copper	•	•	•	•	•	•	•	•			•	•
	Primary lead	•	•	•	•	•	•	•	•			•	•
	Secondary lead	•	•		•	•	•					•	•
	Primary zinc	•	•	•	•	•	•	•	•			•	•
	Secondary zinc	•	•		•	•	•						•
	Primary aluminum					•	•	•					•
	Secondary aluminum	•	•		•	•	•	•		•		•	•
	Nickel							•					
	Antimony								•				
Non-metallic minerals manufacturing	Gold (large scale)	•											
	Mercury mining	•											
	Cement	•	•	•	•	•	•	•		•		•	•
	Glass	•	•	•	•	•	•	•				•	•
Ferrous smelting	Brick	•	•	•	•	•	•	•	•	•	•		
	Pig iron	•	•	•	•	•	•	•	•	•		•	•
	Steel	•	•	•	•	•	•	•	•	•		•	•
Municipal solid waste (MSW) incineration	Municipal solid waste	•	•	•	•	•	•	•	•	•		•	•
Brake and Tyre wear (B&TW)	Brake pad		•	•	•	•	•	•	•	•	•	•	•
	Tyre		•	•	•	•	•	•	•	•	•	•	•

Table S2 The emission source classification by coal combustion sector

Economic sector	Fuel type	Boiler type	PM control device	SO ₂ control device	NO _x control device
Coal-fired power plant	raw coal	pulverized-coal boiler	ESP	WFGD	SCR
	raw coal	pulverized-coal boiler	ESP	WFGD	
	raw coal	pulverized-coal boiler	ESP		
	raw coal	pulverized-coal boiler	FF	WFGD	SCR
	raw coal	pulverized-coal boiler	FF	WFGD	
	raw coal	pulverized-coal boiler	FF		
	raw coal	pulverized-coal boiler	wet scrubber	WFGD	
	raw coal	pulverized-coal boiler	wet scrubber		
	raw coal	pulverized-coal boiler	cyclone	WFGD	
	raw coal	pulverized-coal boiler	cyclone		
	raw coal	fluidized-bed furnace	ESP	WFGD	SCR
	raw coal	fluidized-bed furnace	ESP	WFGD	
	raw coal	fluidized-bed furnace	ESP		
	raw coal	fluidized-bed furnace	FF	WFGD	SCR
	raw coal	fluidized-bed furnace	FF	WFGD	
	raw coal	fluidized-bed furnace	FF		
	raw coal	fluidized-bed furnace	wet scrubber	WFGD	
	raw coal	fluidized-bed furnace	wet scrubber		
	raw coal	fluidized-bed furnace	cyclone		
	raw coal	stoker fired boiler	ESP	WFGD	
	raw coal	stoker fired boiler	ESP		
	raw coal	stoker fired boiler	FF		
	raw coal	stoker fired boiler	wet scrubber		

	raw coal	stoker fired boiler	cyclone		
	cleaned coal	pulverized-coal boiler	ESP	WFGD	SCR
	cleaned coal	pulverized-coal boiler	ESP	WFGD	
	cleaned coal	pulverized-coal boiler	ESP		
	cleaned coal	pulverized-coal boiler	FF	WFGD	SCR
	cleaned coal	pulverized-coal boiler	FF	WFGD	
	cleaned coal	pulverized-coal boiler	FF		
	cleaned coal	pulverized-coal boiler	wet scrubber	WFGD	
	cleaned coal	pulverized-coal boiler	wet scrubber		
	cleaned coal	pulverized-coal boiler	cyclone	WFGD	
	cleaned coal	pulverized-coal boiler	cyclone		
	cleaned coal	fluidized-bed furnace	ESP	WFGD	SCR
	cleaned coal	fluidized-bed furnace	ESP	WFGD	
	cleaned coal	fluidized-bed furnace	ESP		
	cleaned coal	fluidized-bed furnace	FF	WFGD	SCR
	cleaned coal	fluidized-bed furnace	FF	WFGD	
	cleaned coal	fluidized-bed furnace	FF		
	cleaned coal	fluidized-bed furnace	wet scrubber		
	cleaned coal	fluidized-bed furnace	cyclone		
	cleaned coal	stoker fired boiler	ESP	WFGD	
	cleaned coal	stoker fired boiler	ESP		
	cleaned coal	stoker fired boiler	wet scrubber		
	cleaned coal	stoker fired boiler	cyclone		
	briquette	pulverized-coal boiler	ESP	WFGD	
	coke	pulverized-coal boiler	ESP	WFGD	

Coal-fired industrial boiler	raw coal	stoker fired boiler	ESP
	raw coal	stoker fired boiler	FF
	raw coal	stoker fired boiler	wet scrubber
	raw coal	stoker fired boiler	cyclone
	raw coal	stoker fired boiler	
	raw coal	fluidized-bed furnace	wet scrubber
	raw coal	coke furnace	FF
	raw coal	coke furnace	wet scrubber
	raw coal	coke furnace	
	cleaned coal	stoker fired boiler	ESP
	cleaned coal	stoker fired boiler	FF
	cleaned coal	stoker fired boiler	wet scrubber
	cleaned coal	stoker fired boiler	cyclone
	cleaned coal	stoker fired boiler	
	cleaned coal	fluidized-bed furnace	wet scrubber
	cleaned coal	coke furnace	FF
	cleaned coal	coke furnace	wet scrubber
	cleaned coal	coke furnace	
	briquette	stoker fired boiler	wet scrubber
	briquette	stoker fired boiler	cyclone
	briquette	stoker fired boiler	
	coke	stoker fired boiler	wet scrubber
	coke	stoker fired boiler	cyclone
	coke	stoker fired boiler	
Coal-fired residential	raw coal	stove	

sector	cleaned coal briquette coke	stove stove stove	
Coal-fired other sectors	raw coal	stoker fired boiler	wet scrubber
	raw coal	stoker fired boiler	cyclone
	raw coal	stoker fired boiler	
	cleaned coal	stoker fired boiler	
	briquette	stoker fired boiler	
	coke	stoker fired boiler	

168 Table S3 Statistical parameters of bootstrap mean contents of Hg, As, Se, Pb, Cd, Cr, Ni and Sb in
169 produced coal by provinces (Tian et al., 2013)

Provinces	Hg	As	Se	Pb	Cd	Cr	Ni	Sb
Anhui	0.43	2.89	7.54	13.24	0.11	31.25	19.57	0.25
Beijing								
Chongqing	0.31	5.66	3.69	30.44	1.22	28.44	20.9	1.71
Fujian	0.07	9.93	1.22	25.53	0.31	30.48	16.42	0.38
Gansu	0.27	4.14	0.51	8.35	0.08	23.7	19.3	0.7
Guangdong	0.07	8.3	0.6	24.4	0.25	74	24.9	
Guangxi	0.33	16.94	5.03	29.94	0.41	116.41	22.48	5.55
Guizhou	0.39	6.68	3.82	23.81	0.79	28.47	22.87	6.01
Hainan								
Hebei	0.15	4.88	2.31	29.3	0.23	32.52	14.61	0.41
Heilongjiang	0.12	3.42	0.9	22.15	0.13	15.48	10.49	0.79
Henan	0.2	2.2	4.86	16.78	0.54	24.94	11.84	0.37
Hubei	0.2	5.3	8.76	47.39	0.36	40.52	18.61	1.17
Hunan	0.12	10.59	3.72	26.29	0.64	37.03	13.25	1.54
Inner Mongolia	0.22	5.77	1.1	26.67	0.1	13.02	6.35	0.7
Jiangsu	0.69	2.74	6.11	20.98	0.06	19.82	15.48	0.55
Jiangxi	0.16	7.41	8.39	19.33	0.56	39.75	22.66	1.83
Jilin	0.4	11.57	4.06	29	0.15	23.09	15.34	1.02
Liaoning	0.17	5.51	0.85	19.68	0.16	26.24	24.13	0.81
Ningxia	0.22	3.65	4.27	14.05	1.1	10.63	10.95	0.27
Qinghai	0.25	2.68	0.3	10.72	0.03	30.82	12.2	0.91
Shaanxi	0.21	3.87	3.43	35.17	0.75	32.73	18.86	2.95
Shandong	0.18	5.23	3.66	16.64	0.39	20.62	23.77	0.47
Shanghai								
Shanxi	0.17	3.84	3.85	26.23	0.75	21.57	15.41	1.13
Sichuan	0.29	5.38	3.31	28.29	1.95	33	19.28	1.7
Tianjin								
Xinjiang	0.06	2.97	0.24	2.68	0.12	7.83	8.26	0.67
Yunnan	0.36	8.82	1.48	42.54	0.8	73.62	24.32	0.97
Zhejiang	0.65	12.04	12.02	17.25	0.47	24.2	9.95	0.73

170

171 Table S4 Mn Content of Raw Coal as Mined in China, by Province

Provinces ^a	Number of samples	Minimum (µg/g)	Maximum (µg/g)	Arithmetic mean (µg/g)	Literature cited
Anhui	47	0.80	76.30	27.69	(Tang et al., 2002; Wu, 2006; Li et al., 2011)
Beijing ^b	/	/	/	45.80	/
Chongqing	20	5.23	291.00	66.65	(Zhao et al., 2002; Zhuang et al., 2003)

Fujian	7	30.00	459.00	134.28	(Yan and Lu, 1995)
Gansu	13	31.00	1820.00	671.32	(Ren et al., 2006)
Guangxi	15	4.00	128.70	52.49	(Yan and Lu, 1995; Tang et al., 2002)
Guizhou	101	7.00	937.00	152.62	(Zhuang et al., 2000; Tang et al., 2002; Wu et al., 2008)
Hebei	5	20.00	111.00	45.80	(Zhao et al., 2002)
Heilongjiang	1	219.80	219.80	219.80	(Tang et al., 2002; Ren et al., 2006)
Henan	10	22.53	367.46	101.39	(Yan and Lu, 1995; Guo et al., 2005)
Hubei	9	4.00	100.00	49.53	(Yan and Lu, 1995)
Hunan	7	4.00	690.00	266.01	(Wang and Mo, 1999; Tang et al., 2002)
Inner Mongolia	10	12.70	510.00	149.43	(Tang et al., 2002; Guo et al., 2005; Li et al., 2008)
Jiangsu	3	3.90	188.00	95.95	(Tang et al., 2002; Xiu and Wen, 2004; Ren et al., 2006)
Jiangxi	21	8.00	224.00	79.59	(Tang et al., 2002)
Jilin	10	3.30	270.90	84.39	(Ma et al., 2000; Tang et al., 2002; Ren et al., 2006)
Liaoning	6	7.00	200.34	120.56	(Tang et al., 2002; Guo et al., 2005; Ren et al., 2006)
Ningxia	16	7.75	209.50	48.49	(Zhao et al., 2002)
Qinghai	4	22.08	212.00	82.54	(Ren et al., 2006)
Shaanxi	31	6.39	3950.00	398.87	(Dou et al., 1998; Yang et al., 2008a; Yang et al., 2008b)
Shandong	19	9.00	239.50	87.06	(Yan and Lu, 1995; Tang et al., 2002; Guo et al., 2005)
Shanxi	64	0.20	1624.00	80.90	(Tang et al., 2002; Guo et al., 2005)
Sichuan	14	7.20	412.00	121.37	(Tang et al., 2002; Zhuang et al., 2003)
Xinjiang	99	2.00	501.00	52.18	(Cui et al., 2004; Zhou et al., 2010)
Yunnan	9	31.00	125.30	51.41	(Tang et al., 2002; Guo et al., 2005; Dai et al., 2009)
Zhejiang	3	28.00	41.24	32.71	(Li et al., 1993; Tang et al., 2002)

^a Hong Kong, Macao, Taiwan are not included in this table, Guangdong, Hainan, Shanghai, Tianjin and Tibet do not produce raw coal.

^b Beijing lack of corresponding date, in this study, we choose the Mn content of Hebei instead.

Table S5 Co Content of Raw Coal as Mined in China, by Province

Provinces ^c	Number of samples	Minimum (µg/g)	Maximum (µg/g)	Arithmetic mean (µg/g)	Literature cited
Anhui	97	1.32	65.70	12.12	(Tang et al., 2002; Wu, 2006; Chen et al., 2009)
Beijing ^d	/	/	/	8.91	/
Chongqing	38	1.38	90.30	13.38	(Zhao et al., 2002; Bai et al., 2007)

Fujian	4	1.24	15.50	7.55	(Yan and Lu, 1995; Wang et al., 1997; Xu et al., 2001)
Gansu	3	1.54	15.90	7.05	(Ren et al., 2006)
Guangxi	35	2.24	19.90	7.05	(Yan and Lu, 1995; Wang et al., 1997; Xu et al., 2001; Zeng et al., 2005)
Guizhou	148	0.40	119.00	11.91	(Zhuang et al., 1999; Zhuang et al., 2000; Tang et al., 2002; Yang, 2006; Wu et al., 2008)
Henan	9	3.25	12.77	5.93	(Yan and Lu, 1995; Xu et al., 2001; Tang et al., 2002)
Hubei	13	3.00	45.00	8.91	(Yan and Lu, 1995; Xu et al., 2001)
Hebei	38	1.00	24.40	6.80	(Wang et al., 1997; Zhuang et al., 1999; Tang et al., 2002; Dai et al., 2003; Tang et al., 2005; Tang et al., 2009)
Heilongjiang	7	5.60	25.50	12.42	(Tang et al., 2002; Ren et al., 2006)
Hunan	12	0.80	18.50	6.15	(Wang and Mo, 1999; Tang et al., 2002)
Inner Mongolia	99	0.20	28.20	4.08	(Wang et al., 1997; Tang et al., 2002; Dai et al., 2003; Li et al., 2008)
Jilin	13	4.98	38.50	10.91	(Ma et al., 2000; Ren et al., 2006)
Jiangsu	3	1.30	20.10	11.20	(Tang et al., 2002; Xiu and Wen, 2004)
Jiangxi	20	1.00	13.00	5.48	(Xu et al., 2001; Tang et al., 2002)
Liaoning	24	3.60	53.66	13.59	(Kong et al., 2001; Tang et al., 2002; Ren et al., 2004)
Ningxia	18	0.88	22.60	7.29	(Zhao et al., 2002; Bai, 2003)
Qinghai	4	2.19	4.03	2.85	(Zhao et al., 2002; Bai, 2003)
Sichuan	21	0.80	47.60	9.39	(Tang et al., 2002; Zhuang et al., 2003)
Shandong	73	0.34	46.30	5.89	(Yan and Lu, 1995; Huang et al., 2000; Xu et al., 2001; Tang et al., 2002)
Shaanxi	34	0.94	32.90	8.65	(Dou et al., 1998; Tang et al., 2002; Yang et al., 2008a; Yang et al., 2008b)
Shanxi	69	0.40	28.30	4.82	(Wang et al., 1997; Zhuang et al., 1999; Tang et al., 2002; Dai et al., 2003)
Xinjiang	62	0.45	25.80	6.63	(Tang et al., 2002; Zhou et al., 2010)
Yunnan	40	1.79	37.86	11.84	(Tang et al., 2002; Dai et al., 2009; Hu et al., 2009)
Zhejiang	3	2.65	7.39	4.64	(Li et al., 1993; Tang et al., 2002)

^c Hong Kong, Macao, Taiwan are not included in this table, Guangdong, Hainan, Shanghai, Tianjin and Tibet do not produce raw coal.

^d Beijing lack of corresponding date, in this study, we choose the Co content of Hebei instead.

Table S6 Cu Content of Raw Coal as Mined in China, by Province

Province ^e	Number of	Minimum (µg/g)	Maximum (µg/g)	Arithmetic mean	Literature cited
-----------------------	-----------	----------------	----------------	-----------------	------------------

samples					($\mu\text{g/g}$)
Anhui	85	5.03	140.00	36.21	(Tang et al., 2002; Wu, 2006; Chen et al., 2009)
Beijing ^f	/	/	/	27.37	/
Chongqing	28	14.50	156.00	42.57	(Zhuang et al., 2003; Bai et al., 2007; Zhu and Li, 2009)
Fujian	4	21.60	59.00	38.48	(Yan and Lu, 1995; Xu et al., 2001)
Gansu	1	7.25	7.25	7.25	(Bai, 2003; Ren et al., 2006)
Guangxi	45	3.00	69.00	25.79	(Yan and Lu, 1995; Xu et al., 2001; Zeng et al., 2005)
Guizhou	131	0.90	370.00	55.04	(Zhuang et al., 2000; Yang, 2006; Wu et al., 2008; Cheng et al., 2013)
Henan	8	23.30	60.37	40.86	(Yan and Lu, 1995; Xu et al., 2001; Tang et al., 2002)
Hubei	9	19.00	81.00	33.89	(Yan and Lu, 1995; Xu et al., 2001)
Hebei	31	6.90	78.40	27.37	(Zhuang et al., 1999; Tang et al., 2002; Dai et al., 2003; Tang et al., 2005; Tang et al., 2009)
Heilongjiang	12	4.10	69.00	15.62	(Tang et al., 2002; Ren et al., 2006)
Hunan	5	4.28	51.50	25.79	(Wang and Mo, 1999; Tang et al., 2002)
Inner Mongolia	93	1.60	92.20	18.63	(Tang et al., 2002; Dai et al., 2003; Li et al., 2008)
Jilin	10	5.00	98.70	28.17	(Ma et al., 2000; Tang et al., 2002)
Jiangsu	2	21.60	76.30	48.95	(Tang et al., 2002; Xiu and Wen, 2004)
Jiangxi	20	7.00	60.70	21.13	(Xu et al., 2001; Bai, 2003)
Liaoning	19	7.90	85.00	30.38	(Kong et al., 2001; Ren et al., 2004)
Ningxia	4	1.49	8.07	4.52	(Zhao et al., 2002; Bai, 2003)
Qinghai ^g	/	/	/	15.71	/
Sichuan	12	11.20	65.90	33.52	(Tang et al., 2002; Bai, 2003; Zhuang et al., 2003)
Shandong	37	2.64	238.00	34.78	(Yan and Lu, 1995; Liu et al., 2001; Xu et al., 2001; Tang et al., 2002)
Shaanxi	31	5.60	164.00	31.93	(Dou et al., 1998; Tang et al., 2002; Yang et al., 2008a; Yang et al., 2008b)
Shanxi	57	0.00	264.00	27.89	(Zhuang et al., 1999; Tang et al., 2002; Dai et al., 2003)
Xinjiang	96	0.80	36.00	6.58	(Zhao et al., 2002; Bai, 2003; Zhou et al., 2010)
Yunnan	24	0.00	169.00	59.38	(Tang et al., 2002; Dai et al., 2009; Hu et al., 2009)
Zhejiang	1	93.28	93.28	93.28	(Zhao et al., 2002; Bai, 2003)

^e Hong Kong, Macao, Taiwan are not included in this table, Guangdong, Hainan, Shanghai, Tianjin and Tibet do not produce raw coal.

^{f,g} Beijing and Qinghai lack of corresponding date, in this study, we choose the Cu content of

Hebei and the average Cu content of surrounding province (Gansu, Sichuan and Xinjiang) instead, respectively.

Table S7 Zn Content of Raw Coal as Mined in China, by Province

Provinces ^h	Number of samples	Minimum (μg/g)	Maximum (μg/g)	Arithmetic mean (μg/g)	Literature cited
Anhui	100	1.00	112.00	26.17	(Tang et al., 2002; Wu, 2006; Chen et al., 2009; Li et al., 2011)
Beijing ⁱ	/	/	/	49.54	/
Chongqing	26	1.00	39.00	23.41	(Zhao et al., 2002; Zhuang et al., 2003)
Fujian	4	90.00	299.00	174.75	(Yan and Lu, 1995; Wang et al., 1997)
Gansu	2	6.40	54.30	30.35	(Ren et al., 2006)
Guangxi	38	1.41	212.00	56.88	(Yan and Lu, 1995; Wang et al., 1997; Zeng et al., 2005)
Guizhou	157	0.79	561.00	56.97	(Zhuang et al., 1999; Zhuang et al., 2000; Yang, 2006; Li et al., 2011; Wei et al., 2012; Cheng et al., 2013)
Henan	8	10.41	60.00	31.93	(Yan and Lu, 1995)
Hubei	11	5.00	384.00	63.46	(Yan and Lu, 1995)
Hebei	40	5.13	131.00	49.54	(Zhuang et al., 1999; Tang et al., 2002; Dai et al., 2003; Tang et al., 2005; Tang et al., 2009)
Heilongjiang	2	21.40	33.00	27.20	(Ren et al., 2006)
Hunan	4	19.80	158.00	60.35	(Zhao et al., 2002)
Inner Mongolia	97	23.90	257.00	43.18	(Wang et al., 1997; Dai et al., 2003; Li et al., 2008)
Jilin	14	5.40	360.00	79.71	(Ma et al., 2000; Tang et al., 2002)
Jiangsu	2	17.20	18.94	18.07	(Tang et al., 2002; Xiu and Wen, 2004)
Jiangxi	17	3.40	173.00	92.12	(Zhao et al., 2002)
Liaoning	20	22.00	310.00	70.71	(Kong et al., 2001; Tang et al., 2002; Ren et al., 2004)
Ningxia	9	7.30	73.96	21.60	(Song et al., 2011)
Qinghai ^j	/	/	/	30.89	/
Sichuan	13	22.30	99.50	45.65	(Zhao et al., 2002; Zhuang et al., 2003)
Shandong	62	2.67	68.70	16.38	(Yan and Lu, 1995; Huang et al., 2000; Liu et al., 2001; Tang et al., 2002)
Shaanxi	33	8.75	1511.00	114.64	(Dou et al., 1998; Yang et al., 2008a; Yang et al., 2008b)
Shanxi	62	0.56	864.85	65.05	(Wang et al., 1997; Zhuang et al., 1999; Tang et al., 2002; Dai et al., 2003)
Xinjiang	65	4.00	112.00	16.55	(Tang et al., 2002; Zhou et al., 2010)
Yunnan	40	0.00	204.00	59.11	(Dai et al., 2009; Hu et al., 2009)

Zhejiang	1	14.81	14.81	14.81	(Zhao et al., 2002)
----------	---	-------	-------	-------	---------------------

189 ^h Hong Kong, Macao, Taiwan are not included in this table, Guangdong, Hainan, Shanghai, Tianjin
190 and Tibet do not produce raw coal.

191 ^{i,j} Beijing and Qinghai lack of corresponding data, in this study, we choose the Zn content of Hebei
192 and the average Zn content of surrounding province (Gansu, Sichuan and Xinjiang) instead,
193 respectively.

Table S8 Averaged concentrations of Hg, As, Se, Pb, Cd, Cr, Ni, Sb, Mn, Co, Cu and Zn in coals as consumed by province (unit: $\mu\text{g/g}$).

Provinces	Hg	As	Se	Pb	Cd	Cr	Ni	Sb	Mn	Co	Cu	Zn
Anhui	0.40	2.94	7.19	13.82	0.16	30.37	19.16	0.28	34.85	11.37	36.36	27.00
Beijing	0.17	4.02	3.59	26.74	0.66	23.39	15.27	1.01	75.49	5.16	27.82	63.24
Chongqing	0.32	5.74	3.70	29.91	1.19	28.44	21.06	2.05	73.65	13.29	43.67	26.11
Fujian	0.12	4.48	2.32	15.23	0.29	18.53	12.93	0.37	86.32	6.55	29.83	68.99
Gansu	0.23	3.74	0.43	7.48	0.08	21.31	16.30	0.72	166.54	6.50	8.05	27.64
Guangdong	0.14	4.04	2.78	19.20	0.53	18.67	12.29	1.41	99.33	6.09	27.99	59.28
Guangxi	0.14	6.31	1.75	14.04	0.22	44.78	9.73	1.86	52.55	6.78	23.12	38.71
Guizhou	0.39	6.69	3.81	23.86	0.79	28.60	22.87	6.00	153.52	11.88	55.20	57.30
Hainan	0.09	2.07	2.08	14.16	0.40	11.64	8.32	0.61	68.33	5.23	21.91	46.98
Hebei	0.19	5.17	1.94	27.45	0.25	20.71	10.50	0.68	104.34	5.08	22.95	49.06
Heilongjiang	0.16	4.39	0.97	23.76	0.12	14.97	9.46	0.76	187.49	9.30	17.24	35.33
Henan	0.19	2.64	4.58	19.47	0.59	24.61	12.87	0.61	96.01	5.77	37.72	41.57
Hubei	0.21	3.85	4.22	33.40	0.70	31.65	17.70	2.26	98.98	8.00	32.77	92.46
Hunan	0.14	8.10	3.80	25.46	0.66	31.66	13.72	1.34	200.45	5.75	27.19	59.80
Inner Mongolia	0.21	5.57	1.13	25.54	0.11	12.97	6.63	0.69	142.62	4.18	18.60	42.68
Jiangsu	0.25	4.13	3.47	26.28	0.44	24.72	15.73	1.50	103.15	7.38	30.50	64.74
Jiangxi	0.18	5.25	6.19	22.91	0.61	33.52	19.09	1.71	88.56	6.28	27.96	81.93
Jilin	0.28	7.77	2.38	26.86	0.15	18.10	11.33	0.86	128.46	8.23	22.62	57.68
Liaoning	0.18	5.42	1.18	21.94	0.16	20.25	16.05	0.77	130.45	9.77	24.51	56.64
Ningxia	0.21	3.67	4.22	15.44	1.06	11.88	11.46	0.37	51.87	7.00	7.19	26.70
Qinghai	0.24	2.89	1.16	11.44	0.26	26.46	11.93	0.77	75.08	3.81	13.29	28.88
Shaanxi	0.21	3.85	3.51	33.16	0.78	30.63	18.11	2.70	102.92	8.48	29.21	104.21
Shandong	0.19	5.05	2.76	22.77	0.37	18.26	15.05	0.71	108.20	4.97	27.19	38.62

Shanghai	0.28	4.92	2.73	24.01	0.20	17.07	10.94	0.68	118.85	5.76	25.58	39.93
Shanxi	0.16	3.76	3.64	24.81	0.70	20.74	14.96	1.08	79.91	4.88	27.06	63.26
Sichuan	0.29	5.45	3.27	29.64	1.74	34.39	19.63	1.76	110.71	9.90	35.28	49.16
Tianjin	0.17	3.98	3.65	26.52	0.69	22.39	15.14	1.05	76.65	5.09	27.84	63.76
Xinjiang	0.06	3.00	0.25	2.93	0.12	7.88	8.24	0.67	53.35	6.62	6.71	16.83
Yunnan	0.36	8.73	1.58	41.73	0.80	71.66	24.26	1.19	55.82	11.90	59.63	59.34
Zhejiang	0.20	4.05	3.41	27.76	0.60	24.73	15.81	1.63	95.01	6.55	28.43	74.94

195

Table S9 Removal efficiencies of 12 HMs through coal cleaning and coke process

Categories	Hg	As	Se	Pb	Cd	Cr
Coal cleaning	50.0%	54.0%	30.0%	36.3%	32.2%	58.0%
Coking process	90%	30%	40%	31.5%	20%	24%
Category	Ni	Sb	Mn	Co	Cu	Zn
Coal cleaning	58.5%	35.7%	68.2%	39.3%	31.8%	48.6%
Coking process	9.7%	70%	7.6%	7.1%	20%	26.1%

196

197

Table S10 Release rates of Mn, Co, Cu and Zn from coal-fired facilities.

Categories	Release Rates (%) ^k				Literature cited
	Mn	Co	Cu	Zn	
Pulverized-coal boiler	91.70			94.70	(Jin et al., 2003)
	67.00	88.00			(Nodelman et al., 2000)
	57.00	90.00			
	58.00	62.00			(Llorens et al., 2001)
	94.40	92.86	92.31		(Benson et al., 1995)
	86.00	94.00	93.00	96.00	(Xu et al., 2004)
Stoker fired boiler				84.00	(Álvarez-Ayuso et al., 2006)
	16.24	42.44	25.69	33.34	(Wang et al., 1996)
	6.30	11.40			(Zhang et al., 2003)
	26.00	21.88			(Song et al., 2006a)
				15.00	(He et al., 2005)
				5.00	
Fluidized-bed furnace				12.00	(Reddy et al., 2005)
	47.70	57.50	50.20	51.40	
	50.20	55.30	43.40	49.70	(Benson et al., 1995)
	42.70	56.20	45.75	44.80	
	64.29	66.70	83.30		(Klika et al., 2001)
		76.45	78.90	82.01	
		82.29	87.26	77.94	(Bartoňová and Klika, 2009)
		47.00	30.00		
		61.00	68.00		(Zajusz-Zubek and Koniecznyński, 2003)
	28.00				
Coke furnace	23.00	22.00	11.00		(Chen et al., 2008)
	38.00	36.00	22.00		
	40.00	37.00	33.00		(Guo et al., 2004)
	12.00				
				58.00	(Helble et al., 1996)
				30.00	(Wei et al., 2012)

198

199

200

^k the release rate of Hg, As, Se, Pb, Cd, Cr, Ni and Sb from different combustion boilers can be referred in our previously studies (Tian et al., 2010; Tian et al., 2011; Tian et al., 2012a; Tian et al., 2012b).

201
202

Table S11. Removal efficiencies of Mn, Co, Cu and Zn by different control devices

Categories	Release Rates (%) ¹				Literature cited
	Mn	Co	Cu	Zn	
ESPs	93.00	97.00	97.00		(Benson et al., 1995)
	95.00				
	97.90	97.20		92.90	(Ondov et al., 1979)
	99.90	99.80			
	99.10	99.80			
	97.20	99.40			(Nyberg et al., 2009)
		97.70			
	98.50	98.20			(Helble, 2000)
	86.00	94.00	93.00	96.00	(Xu et al., 2004)
		90.02			(Han et al., 2002)
FFs	87.00	93.00	97.75	97.50	(Nodelman et al., 2000)
	98.00	99.00			
	99.70	99.90			
	99.80	99.90			(Nyberg et al., 2009)
Cyclone	67.00	72.00	60.00	64.00	(Gogebakan and Selçuk, 2009)
Wet scrubber	98.97	99.82	98.97	99.03	(Ondov et al., 1979)
	65.79	76.19	55.56	29.09	
	37.50	66.67	86.49	52.50	(Córdoba et al., 2012a)
WFGD	72.08	78.95	24.56	71.38	(Córdoba et al., 2012b)
		68.93	35.26	80.00	
		32.30	27.27		
		41.95	22.17		(Tang et al., 2013)
		32.88	31.29		

203 ¹ the removal efficiencies of Hg, As, Se, Pb, Cd, Cr, Ni and Sb by different control device can be
204 referred in our previously studies (Tian et al., 2010;Tian et al., 2012a;Tian et al., 2012b;Tian et al.,
205 2011).

206
207

Table S12 TSP emission factors for vehicle brake and tyre wear

Vehicle categories	TSP emission factor (g/km)	Uncertainty range(g/km)
TSP emission factors for vehicle tyre wear		
Two-wheel vehicles	0.0046	0.0042–0.0053
Passenger cars	0.0107	0.0067–0.0162
Light-duty trucks	0.0169	0.0088–0.0217
Heavy-duty vehicles	0.0412	0.0227–0.0898
TSP emission factors for vehicle brake wear		
Two-wheel vehicles	0.0037	0.0022 –0.0050
Passenger cars	0.0075	0.0044 –0.0100
Light-duty trucks	0.0117	0.0088 –0.0145
Heavy-duty vehicles	0.0365	0.0235 –0.0420

208

Table S13 Composition of tyre and brake wear in term of heavy metals, ppm

Element	tyre			brake		
	mean	min	max	mean	min	max
As	3.8	1.6	6	67.5	10	130
Se	20	/	/	20	/	/
Pb	176	6.3	670	6072	120	20000
Cd	4.7	1.4	9	22.4	1.5	57
Cr	23.8	2	61	2311	115	8050
Ni	29.9	2.4	63	327	80	600
Sb	2	/	/	10000	/	/
Mn	51	2	100	2460	1700	3220
Co	12.8	0.9	24.8	6.4	/	/
Cu	174	1.8	490	51112	370	142000
Zn	7434	430	13494	8676	270	21800

Table S14 Parameter values used in the transformed normal distribution function computation of the variation of heavy metals emission factors over time

Elements	Parameters	Hg	As	Se	Pb	Cd	Cr	Ni	Sb	Mn	Cu	Zn
Copper	ef _a	27.50	3333.33	300.00	4000.00	1250.00	25.00	5000.00	336.67	100.00	8333.33	6000.00
	ef _b	8.50	100.00	15.00	200.00	50.00	1.00	50.00	10.10	4.50	250.00	300.00
	s	40	30	30	30	30	30	30	30	30	30	30
Lead	ef _a	43.60	400.00	330.00	8000.00	500.00	57.50	166.67	506.67	/	83.33	680.00
	ef _b	6.00	1.00	16.50	200.00	5.00	2.30	5.00	15.20	/	5.00	20.00
	s	40	30	30	30	30	30	30	30	/	30	30
Zinc	ef _a	75.00	600.00	66.67	2900.00	500.00	39.00	68.00	200.00	/	420.00	16000.00
	ef _b	17.00	5.00	10.00	50.00	5.00	1.17	1.36	6.00	/	25.00	500.00
	s	40	30	30	30	30	30	30	30	30	30	30
Gold (large scale)	ef _a	520	/	/	/	/	/	/	/	/	/	/
	ef _b	25	/	/	/	/	/	/	/	/	/	/
	s	36	/	/	/	/	/	/	/	/	/	/
Mercury mining	ef _a	182	/	/	/	/	/	/	/	/	/	/
	ef _b	45	/	/	/	/	/	/	/	/	/	/
	s	36	/	/	/	/	/	/	/	/	/	/
Iron	ef _a	0.06	3.50	0.26	3.50	1.60	2.67	12.00	0.40	0.83	20.00	57.14
	ef _b	0.04	0.08	0.01	0.07	0.02	0.08	0.12	0.00	0.08	0.40	4.00
	s	40	30	30	30	30	30	30	30	30	30	30
Steel	ef _a	0.05	0.56	0.12	74.32	2.47	4.11	3.15	0.20	129.27	5.79	190.00
	ef _b	0.01	0.01	0.00	1.49	0.02	0.12	0.03	0.00	2.02	0.12	6.05
	s	40	30	30	30	30	30	30	30	30	30	30

Cement	ef _a	0.10	13.94	2.54	54.43	2.28	11.79	8.12	/	55.34	24.20	57.20
	ef _b	0.02	0.07	0.01	0.38	0.01	0.05	0.04	/	0.28	0.12	0.29
	s	35	25	25	25	25	25	25	/	25	25	25
MSW	ef _a	2.80	2.14	0.50	107.00	5.45	4.49	3.93	6.00	9.00	14.00	60.00
	ef _b	0.06	0.05	0.01	0.12	0.01	0.04	0.09	3.00	0.21	0.13	0.11
	s	32	28	28	28	28	28	28	28	28	28	28

Table S15 Abatement efficiencies for nonferrous metals smelting (Pacyna et al., 2002)

Elements	Efficiency, %	95% confidence interval	
		lower, %	upper, %
Hg	0	0	67
As	97	91	99
Se	85	55	95
Pb	95	85	98
Cd	99	96	100
Cr	90	70	97
Ni	97	90	99
Cu	94	81	98
Zn	80	40	93

Table S16 Heavy metals emission factors for non-coal combustion sources: temporal, and process variations

Categories		Durable year	Hg	As	Se	Pb	Cd	Cr	Literature cited
Liquid fuel combustion (g t ⁻¹ fuel	Crude oil	1949-2012	0.01	0.17	0.09	0.19	0.05	0.11	(de Souza et al., 2006; UK, 2013)
	Fuel oil for stationary	1949-2012	0.01	0.17	0.09	0.19	0.05	0.11	(de Souza et al., 2006; UK, 2013)

combustion)	sources								
	Kerosene for stationary sources	1949-2012	0.0019	0.02	0.0115	0.000009	0.0000504	0.0000504	(US EPA, 1996a)
	Diesel oil for stationary sources	1949-2012	0.0019	0.02	0.0115	0.000009	0.0000504	0.0000504	(US EPA, 1996c)
	Gasoline	1949-2012	0.06	0.02	0.3	/	0.01	0.01	(UK, 2013)
	Diesel oil for transportation	1949-2012	0.06	0.02	0.3	0.0325	0.04	0.04	(Wang et al., 2003b; UK, 2013)
	Kerosene for transportation	1949-2012	/	/	0.14	0.06	0.01	0.01	(UK, 2013)
	Primary Al	1949-1996	/	/	/	/	0.197	/	
		2012	/	/	/	/	0.1	/	
	Secondary Al	1949-1996	0.032	0.319	/	3.734	0.175	0.841	(UK, 2000, 2013)
		2012	0.0161	0.162	/	1.896	0.0887	0.427	
Nonferrous metal smelting (g t ⁻¹ nonferrous metal production)	Primary Cu	Pre-1900	27.5	3333	300	4000	1250	25	
		2012	8.5	100	15	200	50	1	(Habashi, 1978; Nriagu, 1979;
	Secondary Cu	1949-1996	1	2	5	90	3	1	Pacyna, 1984; Nriagu and
		2012	0.4	0.8	2	36	1.2	0.4	Pacyna, 1988; Skeaff and
	Primary Pb	Pre-1900	43.6	400	330	8000	500	57.5	Dubreuil, 1997; EC, 2001b;
		2012	6	1	16.5	200	5	2.3	Pacyna and Pacyna, 2001; Pacyna
	Secondary Pb	1949-1996	1	0.5	/	100	2.5	1.773	et al., 2002; Theloke et al., 2008;
		2012	0.4	0.2	/	40	1	0.709	Streets et al., 2011; EEA, 2009,
	Primary Zn	Pre-1900	75	600	67	2900	500	39.0	2013)
		2012	17	5	10	50	5	1.170	

Ferroous metals smelting (g t ⁻¹ ferrous metal production)	Secondary Zn	1949-1996	0.013	0.945	/	10.439	5.515	1.799	(Hylander and Meili, 2005; Pacyna, 2006; Pacyna, 2010; Pirrone, 2010; Streets, 2011)
		2012	0.0065	0.48	/	5.3	2.8	0.913	
	Ni smelting	1949-1996	/	/	/	/	/	/	
		2012	/	/	/	/	/	/	
	Sb smelting	1949-1996	/	/	/	/	/	/	
		2012	/	/	/	/	/	/	
	Gold (large scale)	Pre-1900	520	/	/	/	/	/	
		2012	25	/	/	/	/	/	
	Mercury mining	Pre-1900	182	/	/	/	/	/	
		2012	45	/	/	/	/	/	
	Pig iron	Pre-1900	0.06	3.5	0.26	3.5	1.6	2.7	
		2015	0.04	0.08	0.013	0.0699	0.016	0.08	
	Steel produced	Pre-1900	0.05	0.5584	0.12	74.3	2.5	4.1	
		2012	0.008	0.011168	0.003	1.5	0.025	0.123	
	Glass	1949-1996	0.124	0.248	49.556	24.8	0.372	6.194	
		2012	0.050	0.101	20.153	10.1	0.151	2.519	
	Cement	Pre-1900	0.1	13.94	2.54	54.4	2.28	11.8	
		2012	0.0202	0.0697	0.0127	0.38	0.0114	0.0511	
The output of Non-metallic minerals manufacturing in China, 2000-2012 (g t ⁻¹ material production)	Brick	1949-1996	0.044	0.059	0.104	0.068	0.007	0.023	(US EPA, 1996b; NPI, 1998)
		2012	0.015	0.020	0.036	0.023	0.002	0.008	
Municipal solid waste	MSWI	Pre-1900	2.8	2.14	0.5	107	5.45	4.49	(Nriagu, 1979; Pacyna, 1984; Nriagu and Pacyna, 1988; US

incineration (g t ⁻¹ waste)		2016	0.060	0.053	0.0117	0.118	0.012	0.037	EPA, 1996a; UK, 2000)
Biomass burning (g t ⁻¹ residue)	Crop straw	1949-2012	0.008	0.058	0.036	0.865	0.049	0.22	(US EPA, 1996; Li et al., 2007;
	Firewood	1949-2012	0.03	0.03	0.09	0.91	0.08	0.9	EEA, 2013; UK, 2013)
Category		Durable year	Ni	Sb	Mn	Co	Cu	Zn	Literature cited
	Crude oil	1949-2012	10.6	/	0.223	0.151	0.460	1.035	(de Souza et al., 2006; UK, 2013)
	Fuel oil for stationary sources	1949-2012	10.6	/	0.223	0.151	0.460	1.035	(de Souza et al., 2006; UK, 2013)
	Kerosene for stationary sources	1949-2012	0.06	/	0.0504	0.101	0.030	0.489	(US EPA, 1996a)
Liquid fuel combustion (g t ⁻¹ fuel combustion)	Diesel oil for stationary sources	1949-2012	0.06	/	0.0504	0.101	0.030	0.489	(US EPA, 1996c)
	Gasoline	1949-2012	0.04	/	0.004	0.002	0.02	0.0275	(UK, 2013)
	Diesel oil for transportation	1949-2012	0.04	/	0.040	0.0151	0.221	0.234	(Wang et al., 2003b; UK, 2013)
	Kerosene for transportation	1949-2012	0.03	/	0.004	0.002	0.034	0.01	(UK, 2013)
Nonferrous metal smelting (g t ⁻¹ nonferrous metal	Primary Al	1949-1996 2012	19.7 10	/ /	/ /	/ /	/ /	19.7 10	(UK, 2000; 2013)
	Secondary Al	1949-1996 2010	0.802 0.407	/ /	1.16 0.588	/ /	3.2 1.621	15.2 7.734	
	Primary Cu	Pre-1900	5000	337	100	/	8333	6000	(Nriagu, 1979; Pacyna, 1984;

production)	Secondary Cu	2012	50	10.1	4.5	/	250	300	Nriagu and Pacyna, 1988; Skeaff and Dubreuil, 1997; EC, 2001a; Pacyna and Pacyna, 2001; Pacyna et al., 2002; Theloke et al., 2008; EEA, 2009, 2013)	
		1949-1996	1	3	/	/	100	200		
	Primary Pb	2012	0.4	1.2	/	/	40	80		
		Pre-1900	167	507	/	/	83	680		
	Secondary Pb	2012	5	15.2	/	/	5	20		
		1949-1996	/	/	/	/	1	20		
	Primary Zn	2012	/	/	/	/	0.4	8		
		Pre-1900	68	200	/	/	420	16000		
	Secondary Zn	2012	1.36	6	/	/	25	500		
		1949-1996	0	0	/	/	/	270		
	Ni smelting	2012	0	0	/	/	/	137.1		
		1949-1996	900	/	/	/	/	/		
	Sb smelting	2012	360	/	/	/	/	/		
		1949-1996	/	173	/	/	/	/		
	Ferrous metals smelting (g t ⁻¹ ferrous metal production)	Pig iron	2012	/	70	/	/	/		/
Pre-1900			12	0.4	0.830	/	20	57.143	(Nriagu, 1979; Pacyna, 1984; Nriagu and Pacyna, 1988; Kakareka et al., 1998; UK, 2000; EC, 2001a; Theloke et al., 2008; EEA, 2009; Pirrone et al., 2010; UK, 2013)	
Steel produced		2015	0.12	0.004	0.082	/	0.4	4	(Tian et al., 2012c)	
		Pre-1900	3.2	0.2	129	/	5.790	190	(Nriagu and Pacyna, 1988; Nriagu and Pacyna, 1988; Theloke et al., 2008; Passant et al., 2002; NPI, 2008; US EPA, 2012)	
Glass		2012	0.032	0.004	2.016	/	0.116	6.0	(EEA, 2000; EC, 2001c)	
		1949-1996	5.0	/	/	/	1.239	24.8	(Nriagu and Pacyna, 1988; Passant et al., 2002; NPI, 2008; US EPA, 2012)	
Cement		2012	2.015	/	/	/	0.504	10.1	(Nriagu and Pacyna, 1988; Passant et al., 2002; NPI, 2008; US EPA, 2012)	
		Pre-1900	8.1	/	55.3	/	24.2	57.2	(Nriagu and Pacyna, 1988; Passant et al., 2002; NPI, 2008; US EPA, 2012)	
Non-metallic minerals manufacturing (g t ⁻¹ material production)		Cement	2012	0.0406	/	0.277	/	0.121	0.286	(Nriagu and Pacyna, 1988; Passant et al., 2002; NPI, 2008; US EPA, 2012)
			Pre-1900	8.1	/	55.3	/	24.2	57.2	(Nriagu and Pacyna, 1988; Passant et al., 2002; NPI, 2008; US EPA, 2012)
Non-metallic minerals manufacturing (g t ⁻¹ material production)		Cement	2012	0.0406	/	0.277	/	0.121	0.286	(Nriagu and Pacyna, 1988; Passant et al., 2002; NPI, 2008; US EPA, 2012)
			Pre-1900	8.1	/	55.3	/	24.2	57.2	(Nriagu and Pacyna, 1988; Passant et al., 2002; NPI, 2008; US EPA, 2012)

Municipal solid waste incineration (g t ⁻¹ waste)	Brick	1949-1996	0.033	0.012	0.132	0.001	/	/	(US EPA, 1996c; NPI, 1998)
		2012	0.011	0.004	0.045	0.0003	/	/	
	MSWI	Pre-1900	3.93	6	9	/	14	60	(Nriagu, 1979; Pacyna, 1984; Nriagu and Pacyna, 1988; US EPA, 1996b; UK, 2000)
		2016	0.086	3	0.208	/	0.127	0.109	
	Crop straw	1949-2012	0.177	0.019	0.0955	0.0045	0.1	0.028	(US EPA, 1996a; Li et al., 2007; EEA, 2013; UK, 2013)
	Firewood	1949-2012	0.98	0.0728	0.652	0.0045	0.1	1.25	

Table S17 Abatement efficiencies for iron and steel production (Kakareka et al., 1998)

Elements	Efficiency, %	95% confidence interval	
		lower, %	upper, %
Pb	96	93	98
Cd	96	91	98
Ni	94	88	97
Zn	95	90	98

Table S18 The emission limits of air pollutants of relative industrial process in China and developed regions, mg/m³

Pollutants	GB 25467-2 010	GB 25466-2 010	GB 30770-2 014	GB 25465-2 010	GB 28663-2 012	GB 28664-2 012	GB 4915-2 013	GB 29620-2 013	GB 26453-2 011	GB 18485-2 014	EU 2000/76/ EC	EU 2001/80/ EC
PM	80	80	30	20-100	50	50-100	30	100	50	20	10~30	50~100
SO ₂	400	400	400	400	100	/	200	400	400	80	50	200~850
NO _x	/	/	200	/	300	/	400	/	700	250	200	200~400
As & compounds	0.4	/	0.5	/	/	/	/	/	/	/	/	/

Pb & compounds	0.7	/	0.5	/	/	/	/	/	/	/	/	/
Hg & compounds	0.012	0.05	0.01	/	/	/	0.05	/	/	0.05	0.05	/
Cd & compounds	/	/	0.05	/	/	/	/	/	/	/	0.05	/
(Cd+Tl) & their compounds	/	/	/	/	/	/	/	/	/	0.1	0.05~0.1	/
(Sb+As+Pb+Cr+Co+Cu+Mn +Ni) & their compounds	/	/	/	/	/	/	/	/	/	1.0	0.5~1.0	/

219 Table S19 Data source of activity data for the main heavy metals emitting sectors in China

Emission sectors	Data Sources
Coal consumption by power plants	China Electric Power Yearbook China Editorial Power Industry Statistics China Mechanical Industry Yearbook
Coal consumption by industrial boilers	China Coal Industry Yearbook China Energy Statistical Yearbook
Coal consumption by residential sectors	China Energy Statistical Yearbook
Coal consumption by other sectors	China Energy Statistical Yearbook
Biomass burning	China Statistical Yearbook
Liquid fuels combustion	China Energy Statistical Yearbook
Nonferrous metals smelting	The Yearbook of Nonferrous Metals Industry of China
Non-metallic minerals manufacturing	China Cement Almanac
Ferrous metal smelting	China Steel Yearbook
Municipal solid waste incineration	China Energy Statistical Yearbook
Brake and tyre wear	China Automotive Industry Yearbook China's Auto Market Almanac

220

221 Table S20 Selected parameters showing method and assumption for uncertainty analysis

Categories	Parameter description	Distribution	Sources or methods
Coal combustion sources			
Coal consumption	power plant	Normal (CV: 5%)	(Zhao et al., 2008)
	Industrial sectors	Normal (CV: 5%)	(Tian et al., 2012a)
	Residential sectors	Normal (CV: 14%)	(Tian et al., 2012a)
	Other sectors	Normal (CV: 16%)	(Tian et al., 2012a)
Release rate	Pulverized-coal boiler	Triangular	Data fitting
	Stoker fired boiler	Triangular	Data fitting
	Fluidized-bed furnace	Triangular	Data fitting
	Coke furnace	Triangular	Data fitting
Removal efficiency	ESPs	Normal (CV: 5%)	Subject judgment
	EFs	uniform	Data fitting
	Wet scrubber	Triangular	Subject judgment
	Cyclone	Normal (CV: 20%)	Subject judgment
	Wet-FGD	Triangular	Subject judgment
	Coal washing	Uniform	Data fitting
Non-coal combustion sources			
Biomass burning	Biofuel consumption	Normal (CV: 20%)	(Zhao et al., 2011)
	Emission factors	Triangular	(Zhao et al., 2011)
	Ratio of biomass burning	Normal (province dependent)	(Zhao et al., 2011)
	straw-to-crop ratio	Uniform (product dependent)	(Zhao et al., 2011)
Liquid fuel	Liquid fuel consumption	Normal (CV: 5%)	(Zhao et al., 2011)

combustion	Emission factors	Normal (CV: 25%)	Subject judgment
Nonferrous metal smelting	Nonferrous metal production	Normal (CV: 5%)	(Zhao et al., 2011)
Non-metallic minerals	Emission factors	Triangular	Data fitting
manufacturing	Output of Cement/ glass / brick	Normal (CV: 20%)	Subject judgment
	emission factors (cement, glass)	Normal (CV: 25%)	Subject judgment
	emission factors (brick)	Normal (CV: 30%)	Subject judgment
Ferrous metal smelting	Pig iron and steel yield	Normal (CV: 15%)	(Zhao et al., 2011)
	Emission factors	Triangular	Subject judgment
Municipal solid waste incineration	MSW consumption	Normal (CV: 20%)	Subject judgment
	Emission factors	Normal (CV: 20%)	Subject judgment
Brake and tyre wear	Vehicle number	Normal (CV: 5%)	(Zhao et al., 2011)
	Average vehicle mileage	Normal (CV: 5%)	(Zhao et al., 2011)
	TSP emission factors	Uniform	Data fitting
	Heavy metal content	Triangular	Data fitting

222

223 Table S21 The atmospheric concentrations of As, Pb, Cr and Cu in PM_{2.5} in Beijing during 2000 to
224 2012

Year	Atmospheric concentration (ng m ⁻³)				Literature cited
	As	Pb	Cr	Cu	
2000	30.0	300.0	20.0	30.0	(Song et al., 2006b)
2001		170.0		50.0	(Duan et al., 2006)
2002	38.3	218.3	26.7	58.3	(Sun et al., 2004)
2003	38.3	218.3	26.7	58.3	(Sun et al., 2004)
2005	16.0	189.5	50.0	53.0	(Yu et al., 2012; Zhang et al., 2012)
2006	20.2	173.3	73.6	43.1	(Cui et al., 2008; Yang et al., 2008c, d; Yu et al., 2012)
2007	19.0	189.7	31.7	51.3	(Gao, 2012; Wang et al., 2010; Yu et al., 2012; Zhang et al., 2010)
2008	7.6	67.5	5.8	25.8	(Mu et al., 2010; Yu et al., 2012; Zhang, 2012; Zhang et al., 2010)
2009	17.2	135.5	13.6	40.0	(Tao et al., 2014; Zhao et al., 2013)
2010	22.8	142.7	16.4	36.8	(Tao et al., 2014; Yu et al., 2013)
2011	15.6		13.4	47.7	(Wang et al., 2014)
2012	23.4	158.0	24.6	54.7	(Guo, 2014; Yang et al., 2015; Zhang et al., 2014)

225

Table S22 Uncertainties in the sectoral emissions of heavy metals in China in 2010

Categories	Hg	As	Se	Pb
Coal-fired power plant	139.4 (-21.5%, 25.7%)	406.4 (-22.6%, 25.2%)	538.6 (-25.5%, 22.5%)	833.0 (-25.6%, 25.9%)
Coal-fired industrial boiler	171.0 (-44.6%, 60.7%)	821.3 (-49.4%, 53.4%)	965.1 (-49.3%, 56.6%)	5449.5 (-57.5%, 58.7%)
Coal-fired residential sector	7.4 (-57.4%, 55.7%)	10.9 (-54.2%, 64.7%)	74.3 (-57.3%, 65.7%)	417.2 (-53.2%, 65.3%)
Coal-fired other sector	22.4 (-66.1%, 71.4%)	485.2 (-70.2%, 78.6%)	282.9 (-69.4%, 71.9%)	1224.1 (-57.0%, 80.9%)
Biomass burning	9.1 (-47.5%, 48.4%)	34.0 (-48.3%, 66.6%)	33.3 (-50.2%, 42.3%)	585.6 (-57.8%, 56.4%)
Liquid fuel combustion	15.9 (-62.9%, 63.6%)	11.8 (-48.2%, 41.3%)	83.5 (-56.0%, 51.3%)	678.4 (-89.3%, 68.2%)
Nonferrous metal smelting	222.5 (-45.2%, 54.8%)	338.9 (-57.1%, 67.6%)	146.7 (-39.5%, 37.4%)	1602.9 (39.6%, 37.8%)
Ferrous metal smelting	29.9 (-52.0%, 52.1%)	57.8 (-55.1%, 61.7%)	9.9 (-47.2%, 58.4%)	1047.3 (-53.5%, 72.3%)
Non-metallic minerals manufacturing	52.9 (-81.6%, 98.5%)	152.4 (-90.2%, 107.5%)	744.0 (-95.1%, 103.2%)	1085.0 (-82.3%, 98.7%)
Municipal solid waste incineration	1.6 (-69.4%, 67.9%)	1.2 (-65.3%, 75.1%)	0.3 (-55.6%, 64.9%)	3.8 (-65.3%, 69.2%)
Brake and tyre wear	/	3.1 (-80.9%, 97.3%)	1.9 (-81.0%, 159.6%)	267.7 (-89.0%, 157.6%)
Total emission	672.1 (-34.2%, 46.7%)	2322.9 (-36.4%, 48.8%)	2880.5 (-39.1%, 50.6%)	13194.5 (-32.7%, 46.9%)
Categories	Cd	Cr	Ni	Sb
Coal-fired power plant	15.5 (-19.0%, 27.0%)	598.3 (-27.2%, 26.4%)	537.8 (-19.1%, 23.7%)	97.9 (-29.4%, 23.9%)
Coal-fired industrial boiler	104.0 (-50.1%, 55.8%)	5317.6 (-61.9%, 53.8%)	1005.8 (-46.8%, 52.3%)	214.8 (-51.6%, 63.6%)
Coal-fired residential sector	3.7 (-58.6%, 60.4%)	58.9 (-56.6%, 67.1%)	33.7 (-58.2%, 68.8%)	1.0 (-58.9%, 77.9%)
Coal-fired other sector	25.7 (-62.5%, 66.5%)	773.7 (-57.5%, 73.0%)	185.7 (-69.8%, 80.0%)	105.9 (-61.1%, 78.7%)
Biomass burning	38.0 (-41.5%, 52.5%)	263.1 (-45.3%, 59.6%)	255.4 (-46.5%, 54.6%)	21.9 (-60.7%, 65.1%)
Liquid fuel combustion	9.6 (-61.0%, 57.9%)	11.9 (-47.7%, 50.3%)	422.1 (-58.2%, 67.9%)	/
Nonferrous metal smelting	200.6 (-54.5%, 56.7%)	19.7 (-35.9%, 37.3%)	422.5 (-39.9%, 38.9%)	122.6 (-38.1%, 33.6%)
Ferrous metal smelting	28.3 (-48.7%, 56.7%)	131.3 (-40.7%, 47.5%)	102.7 (-55.0%, 45.8%)	5.4 (-49.0%, 53.0%)
Non-metallic minerals	28.8 (-80.6%, 90.0%)	190.2 (-85.3%, 93.6%)	155.4 (-90.4%, 86.9%)	3.4 (-96.9%, 106.2%)

manufacturing				
Municipal solid waste incineration	0.3 (-66.5%, 76.2%)	0.9 (-58.8%, 68.3%)	2.0 (-59.3%, 76.0%)	69.5 (-75.7%, 71.0%)
Brake and tyre wear	1.2 (-77.1%, 120.7%)	99.6 (-97.0%, 171.5%)	15.5 (-70.9%, 85.4%)	425.7 (-91.1%, 170.0%)
Total emission	455.8 (-35.7%, 49.3%)	7465.2 (-33.8%, 47.8%)	3138.6 (-38.2%, 49.4%)	1068.1 (-35.6%, 48.3%)
Categories	Mn	Co	Cu	Zn
Coal-fired power plant	3072.1 (-23.5, 29.1%)	164.9 (-33.5%, 25.2%)	1477.5 (-21.6%, 28.7%)	2367.1 (-23.0%, 31.5%)
Coal-fired industrial boiler	4472.2 (-43.6%, 45.1%)	517.4 (-39.8%, 43.5%)	2004.4 (-55.0%, 56.9%)	4449.7 (-51.6%, 49.0%)
Coal-fired residential sector	43.4 (-57.8%, 56.7%)	5.3 (-62.6%, 58.4%)	24.0 (-61.3%, 57.0%)	284.6 (-59.7%, 57.0%)
Coal-fired other sector	2130.7 (-60.6%, 74.7%)	218.8 (-59.7%, 63.2%)	933.0 (-65.2%, 73.4%)	1058.8 (-65.6%, 65.2%)
Biomass burning	158.8 (-52.3%, 53.4%)	3.0 (-45.7%, 56.3%)	66.8 (-45.7%, 46.3%)	227.4 (-54.5%, 63.6%)
Liquid fuel combustion	15.6 (-51.4%, 53.8%)	8.6 (-46.7%, 55.1%)	56.4 (-60.9%, 52.2%)	81.5 (-62.6%, 44.4%)
Nonferrous metal smelting	15.8 (-59.8%, 56.5%)	/	990.1 (-66.8%, 53.5%)	3960.2 (-56.3%, 41.6%)
Ferrous metal smelting	1431.4 (-56.9, 68.2%)	/	331.2 (-43.7%, 48.5%)	6421.2 (-56.4%, 43.9%)
Non-metallic minerals manufacturing	563.5 (-89.9%, 85.7%)	0.3 (-93.3%, 94.1%)	247.6 (-88.7%, 103.3%)	889.8 (-86.4%, 94.2%)
Municipal solid waste incineration	4.9 (-87.9%, 60.9%)	/	3.1 (-61.4%, 66.5%)	3.1 (-79.2%, 61.9%)
Brake and tyre wear	107.4 (-45.1%, 61.8%)	0.9 (-79.7%, 159.8%)	2184.8 (-93.4%, 139.4%)	760.2 (-84.3%, 113.9%)
Total emission	12015.9 (-32.2%, 42.0%)	919.2 (-38.1%, 41.4%)	8318.8 (-37.5%, 50.8%)	20503.7 (-32.2%, 45.5%)

227 **Figures**




Fig. S1. Coal consumption by different sectors in China, 2000-2012




Fig. S2. The output of nonferrous metals in China, 2000-2012




Fig. S3. The output of pig iron and steel products in China, 2000-2012




Fig. S4. The output of non-metallic minerals manufacturing in China, 2000-2012




Fig. S5. The number of civil vehicles in China and Beijing, 2000-2012




Fig. S6. Comparison between historical HM (As, Cr, Pb, and Cu) emissions and temporal variation of atmospheric concentrations in $PM_{2.5}$ in Beijing during 2000-2012




Fig. S7 The distribution of point sources in China




Fig. S8. The uncertainty bounds for China's anthropogenic atmospheric emissions of twelve HMs during 1949 to 2012

Reference

- Álvarez-Ayuso, E., Querol, X., and Tomás, A.: Environmental impact of a coal combustion-desulphurisation plant: Abatement capacity of desulphurisation process and environmental characterisation of combustion by-products, *Chemosphere*, 65, 2009-2017, doi:10.1016/j.chemosphere.2006.06.070, 2006.
- Bai, X. F.: The distributions, modes of occurrence and volatility of trace elements in coals of China, China Coal Science Research Institute, 2003 (in Chinese).
- Bai, X. F., Li, W. H., Chen, Y. F., and Jiang, Y.: The general distributions of trace elements in Chinese coals, *Coal Qual. Technol.*, 1, 1-4, 2007 (in Chinese with English abstract).
- Bartoňová, L., and Klika, Z.: Volatility of Cu, Ni, Cr, Co, Pb, and As in fluidized-bed combustion chamber in relation to their modes of occurrence in coal, *World Acad. Sci. Eng. Technol.*, 3, 29-32, 2009.
- Benson, S. A., Erickson, T. A., and Brekke, D. W.: Comparison of HAPs from advanced and conventional power systems: Tidd versus Cardinal, North Dakota Univ., Grand Forks, ND (United States). Energy and Environmental Research Center, 1995.
- Córdoba, P., Font, O., Izquierdo, M., Querol, X., Leiva, C., López-Antón, M. A., D íz-Somoano, M., Ochoa-González, R., Rosa Martínez-Tarazona, M., and Gómez, P.: The retention capacity for trace elements by the flue gas desulphurisation system under operational conditions of a co-combustion power plant, *Fuel*, 102, 773-788, doi:10.1016/j.fuel.2012.06.059, 2012a.
- Córdoba, P., Ochoa-Gonzalez, R., Font, O., Izquierdo, M., Querol, X., Leiva, C., López-Antón, M. A., D íz-Somoano, M., Rosa Martínez-Tarazona, M., and Fernandez, C.: Partitioning of trace inorganic elements in a coal-fired power plant equipped with a wet Flue Gas Desulphurisation system, *Fuel*, 92, 145-157, doi:10.1016/j.fuel.2011.07.025, 2012b.
- Chen, J., Liu, W. Z., and Chen, P.: The application of SPSS for trace elements in Huainan Mining coal, *Coal Prep. Technol.*, 46-50, 2009 (in Chinese).
- Chen, Y. W., Liu, G. J., Wang, L., Kang, Y., and Yang, J. L.: Occurrence and fate of some trace elements during pyrolysis of Yima coal, China, *Energy Fuels*, 22, 3877-3882, doi:10.1021/ef800485w, 2008.
- Cheng, W., Yang, R. D., Zhang, Q., Cui, Y. C., and Gao, J. B: Distribution characteristics, occurrence modes and controlling factors of trace elements in Late Permian coal from Bijie City, Guizhou Province, *J. China Coal Soc.*, 38, 103-113, 2013 (in Chinese with English abstract).
- Cui, G. L., Quan, S. J., and Wu, C. D.: Jurassic coal in Yanqi Basin (Xinjiang, China): geochemical characteristics of trace elements and their implications, *Acta Scientiarum Naturalium Universitatis Peknensis*, 40, 594-600, 2004 (in Chinese with English abstract).
- Cui, Y., Guo, X. B., Deng, F. R., and Liu, H.: Analysis of water-soluble ions and elements in PM₁₀ and PM_{2.5}, *J. Environ. Health*, 25, 291-294, 2008 (in Chinese with English abstract).
- Dai, S. F., Ren, D. Y., Li, S. S., Song, J. F., and Wu, C. H.: Concentrations of minor elements and regional distribution of arsenic in Late Paleozoic Coals from North China Platform, *J. China Univ. Min. Technol.*, 32, 111-114, 2003 (in Chinese).
- Dai, S. F., Ren, D. Y., Zhou, Y., Wang, X., Zhao, L., and Zhu, X.: Composite formation of trace elements in coals, synsedimentary volcanic ashes from mineral enrichment and underwater jet, *Science in China Press*, 53, 3120-3126, 2009 (in Chinese).

de Souza, R. M., Meliande, A. L., da Silveira, C. L., and Auciú, R. Q.: Determination of Mo, Zn, Cd, Ti, Ni, V, Fe, Mn, Cr and Co in crude oil using inductively coupled plasma optical emission spectrometry and sample introduction as detergentless microemulsions, *Microchem. J.*, 82, 137-141, doi:10.1016/j.microc.2006.01.005, 2006.

Dou, T. H., Xiao, D. X., Dong, Y. Q., Zhang, S. N., and Zhang, Q. L.: A preliminary study on trace elements in Dongsheng coal mining area in Shenfu, *Coal Geol. Explor.*, 26, 11-15, 1998 (in Chinese with English abstract).

Duan, F. K., He, K. B., Ma, Y. L., Yang, F. M., Yu, X. C., Cadle, S. H., Chan, T., and Mulawa, P. A.: Concentration and chemical characteristics of PM_{2.5} in Beijing, China: 2001–2002, *Sci. Total Environ.*, 355, 264–275, doi:10.1016/j.scitotenv.2005.1003.1001, 2006.

European Commission (EC): Integrated Pollution Prevention and Control (IPPC), Best Available Techniques Reference Document on the Production of Iron and Steel, 2001a.

European Commission (EC): Integrated Pollution Prevention and Control (IPPC), Reference Document on Best Available Techniques (BREF) in the Non-Ferrous Metal Industries, 2001b.

European Commission (EC): Integrated Pollution Prevention and Control (IPPC), reference document on Best Available Techniques (BREF) in the glass manufacturing industries, December 2001 2001c.

European Environment Agency (EEA): EMEP/EEA air pollutant emission inventory guidebook 2000, available at: <http://www.eea.europa.eu/publications/emep-eea-guidebook-2000> (last access: 18 August 2013), 2000.

European Environment Agency (EEA): EMEP/EEA air pollutant emission inventory guidebook 2009, available at: <http://www.eea.europa.eu/publications/emep-eea-emission-inventory-guidebook-2009> (last access: 24 December 2013), 2009.

European Environment Agency (EEA): EMEP/EEA air pollutant emission inventory guidebook 2013, available at: <http://www.eea.europa.eu/publications/emep-eea-guidebook-2013> (last access: 12 November 2013), 2013.

Gao, S.: Source apportionment of PM₁₀ and PM_{2.5} in five cities in China, Master Thesis, Tianjin Medical University, 2012 (in Chinese with English abstract).

Gogebakan, Z., and Selçuk, N.: Trace elements partitioning during co-firing biomass with lignite in a pilot-scale fluidized bed combustor, *J. Hazard. Mater.*, 162, 1129-1134, doi:10.1016/j.jhazmat.2008.05.149, 2009.

Guo, R. X., Yang, J. L., Liu, D. Y., and Liu, Z. Y.: Transformation behavior of trace elements during coal pyrolysis, *Fuel Process. Technol.*, 77, 137-143, doi:10.1016/S0378-3820(02)00041-3, 2002.

Guo, R. X., Yang, J. L., Liu, D. Y., and Liu, Z. Y.: The fate of As, Pb, Cd, Cr and Mn in a coal during pyrolysis, *J. Anal. Appl. Pyrolysis*, 70, 555-562, doi:10.1016/S0165-2370(03)00025-1, 2003.

Guo, R. X., Yang, J. L., and Liu, Z. Y.: Behavior of trace elements during pyrolysis of coal in a simulated drop-tube reactor, *Fuel*, 83, 639-643, doi:10.1016/j.fuel.2003.08.021, 2004.

Guo, R. X., Yang, J. L., and Liu, Z. Y.: Volatility of trace harmful elements in coal, *China Environ. Sci.*, 24, 641-645, 2005 (in Chinese with English abstract).

Guo, X. S.: Digestion method and pollution characteristics of heavy metals of the PM_{2.5} in typical

341 northeast Beijing, Master Thesis, Jiangxi Agricultural University, 2014 (in Chinese with
 342 English abstract).

343 Habashi, F.: Metallurgical plants: how mercury pollution is abated, *Environ. Sci. Technol.*, 12,
 344 1372-1376, doi:10.1021/es60148a011, 1978.

345 Han, J., Xu, M. H., Cheng, J. F., Qiao, Y., and Ceng, H. C.: Study of trace element emission factor
 346 in coal-fired boilers, *J. Eng. Thermophys.*, 23, 770-772, 2002 (in Chinese with English
 347 abstract).

348 He, Z. L., Yang, X. E., and Stoffella, P. J.: Trace elements in agroecosystems and impacts on the
 349 environment, *Journal of Trace Elements in Medicine and Biology*, 19, 125-140,
 350 doi:10.1016/j.jtemb.2005.02.010, 2005.

351 Helble, J.: A model for the air emissions of trace metallic elements from coal combustors
 352 equipped with electrostatic precipitators, *Fuel Process. Technol.*, 63, 125-147,
 353 doi:10.1016/S0378-3820(99)00093-4, 2000.

354 Helble, J. J., Mojtahedi, W., Lyyränen, J., Jokiniemi, J., and Kauppinen, E.: Trace element
 355 partitioning during coal gasification, *Fuel*, 75, 931-939, doi:10.1016/0016-2361(96)00056-7,
 356 1996.

357 Hu, R. Z., Qi, H. W., Zhou, M. F., Su, W. C., Bi, X. W., Peng, J. T., and Zhong, H.: Geological
 358 and geochemical constraints on the origin of the giant Lincang coal seam-hosted germanium
 359 deposit, Yunnan, SW China: a review, *Ore Geol. Rev.*, 36, 221-234,
 360 doi:10.1016/j.oregeorev.2009.02.007, 2009.

361 Huang, W. H., Yang, Q., Tang, D. Z., Kang, X. D., and Liu, D. M.: Trace elements geochemistry
 362 of the coals in the Taiyuan formation from Zaozhuang coal field, *Geoscience*, 14, 61-68,
 363 2000 (in Chinese with English abstract).

364 Hylander, L. D.; Meili, M. The rise and fall of mercury: converting a resource to refuse after 500
 365 years of mining and pollution. *Crit. Rev. Env. Sci. Technol.*, 35, 1-36,
 366 doi:10.1080/10643380490492485, 2005.

367 Jin, B. P., Huang, Y. J., Zhong, Z. P., Xiao, R., Dong, C. Q., and Zhou, H. C.: Occurrence of
 368 several trace elements in pulverized coal boiler, *J. Combust. Sci. Technol.*, 9, 323-328, 2003
 369 (in Chinese with English abstract).

370 Kakareka, S., Khomich, V., Kukharchyk, T., and Loginov, V.: Heavy Metal emission factors
 371 assessment for the CIS countries, Institute for problems of Natural Resources use and
 372 Ecology of the National Academy of Sciences of Belarus, Minsk, 1998.

373 Klika, Z., Bartoňová, L., and Spears, D.: Effect of boiler output on trace element partitioning
 374 during coal combustion in two fluidised-bed power stations, *Fuel*, 80, 907-917,
 375 doi:10.1016/S0016-2361(00)00164-2, 2001.

376 Kong, H., Zeng, R., Zhuang, X., and Xu, W.: Research on trace elements in Beipiao coal field in
 377 Liaoning, *Geoscience*, 15, 415-420, 2001 (in Chinese).

378 Koniecznyński, J., Zajusz-Zubek, E., and Jabłońska, M.: The release of trace elements in the
 379 process of coal coking, *The Sci. World J.*, doi.org/10.1100/2012/294927, 2012.

380 Li, H. Y., Hao, Y., Yang, L., and Liu, Y. J.: Characteristics of bitumen trace elements and rare
 381 earth elements of coal in southeastern margin of Ordos basin, *Xinjiang Petrol. Geol.*, 29,
 382 159-162, 2008.

383 Li, H., Zheng, L. G., and Liu, G. J.: The concentration characteristics of trace elements in coal
 384 from the Zhangji mining area, Huainan coalfield, *Acta Petrol. ET Mineral.*, 30, 696-700,

2011 (in Chinese with English abstract).

Li, Q., Cheng, H. G., Zhou, T., Lin, C. Y., and Guo, S.: The estimated atmospheric lead emissions in China, 1990–2009, *Atmos. Environ.*, 60, 1–8, doi:10.1016/j.atmosenv.2012.06.025, 2012.

Li, W. H., Xiong, F., and Jiang, N.: Trace elements in three high sulfur coals, *Coal Anal. Utilization*, 1, 7-9, 1993 (in Chinese).

Li, X. H., Wang, S. X., Duan, L., Hao, J. M., Li, C., Chen, Y. S., and Yang, L.: Particulate and trace gas emissions from open burning of wheat straw and corn stover in China, *Environ. Sci. Technol.*, 41, 6052-6058, doi:10.1021/es0705137, 2007.

Liu, G. J., Peng, Z. C., Yang, P. Y., Wang, G. L., and Song, C.: Changes of trace elements in coal during combustion, *J. Fuel Chem. Technol.*, 29, 119-123, 2001 (in Chinese with English abstract).

Llorens, J. F., Fernandez-Turiel, J. L., and Querol, X.: The fate of trace elements in a large coal-fired power plant, *Environ. Geol.*, 40, 409-416, doi:10.1007/s002540000191, 2001.

Luttrell, G. H., Kohmuench, J. N., and Yoon, R. H.: An evaluation of coal preparation technologies for controlling trace element emissions, *Fuel Process. Technol.*, 65, 407-422, doi:10.1016/S0378-3820(99)00107-1, 2000.

Ma, Z. W., Wang, Q. C., and Fang, F. M.: The contents and constituent characteristic of trace elements in coal in Tonghua and Baishan region of Jilin province, *Heilongjiang Environ. J.*, 24, 38-41, 2000 (in Chinese with English abstract).

Mu, L. T., Wang, J. J., Li, N., Tong, J. Y., and Pan, X. C.: Feature analysis of metal components of PM_{2.5} and PM₁₀ during sand dust weather, *J. Environ. Health*, 27, 755–758, 2010 (in Chinese with English abstract).

National Pollutant Inventroy (NPI): Emission estimation technique manual for bricks, cermmics, & clay product manufacturing, availalbe at: <http://www.npi.gov.au/reporting/industry-reporting-materials/emission-estimation-technique-manuals> (last access: 12 June 2013), 1998.

National Pollutant Inventroy (NPI): Emission estimation technique manual for cement manufacturing, availalbe at: <http://www.npi.gov.au/reporting/industry-reporting-materials/emission-estimation-technique-manuals> (last access: 20 June 2013), 2008.

National Bureau of Statistics (NBS), P. R. China: China Energy Statistical Yearbook, China Statistics Press, Beijing, 2013a.

National Bureau of Statistics (NBS), P. R. China: China Statistical Yearbook, China Statistics Press, Beijing, 2013b.

Nodelman, I. G., Pisupati, S. V., Miller, S. F., and Scaroni, A. W.: Partitioning behavior of trace elements during pilot-scale combustion of pulverized coal and coal-water slurry fuel, *J. Hazard. Mater.*, 74, 47-59, doi:10.1016/S0304-3894(99)00198-3, 2000.

Nriagu, J. O.: Global inventory of natural and anthropogenic emissions of trace metals to the atmosphere, *Nature*, 279, 409-411, doi:10.1038/279409a0, 1979.

Nriagu, J. O., and Pacyna, J. M.: Quantitative assessment of worldwide contamination of air, water and soils by trace metals, *Nature*, 333, 134–139, doi:10.1038/333134a0, 1988.

Nyberg, C. M., Thompson, J. S., Zhuang, Y., Pavlish, J. H., Brickett, L., and Pletcher, S.: Fate of trace element haps when applying mercury control technologies, *Fuel Process. Technol.*, 90, 1348-1353, doi:10.1016/j.fuproc.2009.06.025, 2009.

- Ondov, J. M., Ragaini, R. C., and Biermann, A. H.: Elemental emissions from a coal-fired power plant. Comparison of a venturi wet scrubber system with a cold-side electrostatic precipitator, *Environ. Sci. Technol.*, 13, 598-607, doi:10.1021/es60153a009, 1979.
- Pacyna, J., Van der Most, P., Hobson, M., Wieser, M., Müller, B., Duval, L., Spezzano, P., Lotz, T., and Kakareka, S.: Combustion and Industry Expert Panel workshop, European Joint Research Centre (JRC), Ispra, 2002.
- Pacyna, J. M.: Estimation of the atmospheric emissions of trace elements from anthropogenic sources in Europe, *Atmos. Environ.*, 18, 41-50, doi:10.1016/0004-6981(84)90227-0, 1984.
- Pacyna, J. M., and Pacyna, E. G.: An assessment of global and regional emissions of trace metals to the atmosphere from anthropogenic sources worldwide, *Environ. Rev.*, 9, 269-298, doi:10.1139/a01-012, 2001.
- Pacyna, E. G., Pacyna, J. M., Steenhuisen, F., and Wilson, S.: Global anthropogenic mercury emission inventory for 2000. *Atmos. Environ.*, 40, 4048-4063, doi:10.1016/j.atmosenv.2006.03.041, 2006.
- Pacyna, E. G., Pacyna, J. M., Sundseth, K., Munthe, J., Kindbom, K., Wilson, S., Steenhuisen, F., Maxson, P.: Global emission of mercury to the atmosphere from anthropogenic sources in 2005 and projections to 2020, *Atmos. Environ.*, 44, 2487-2499, doi:10.1016/j.atmosenv.2009.06.009, 2010.
- Passant, N., Peirce, M., Rudd, H. J., Scott, D. W., Marlowe, I., and Watterson, J. D.: UK Particulate and Heavy Metal Emissions from Industrial Processes, Netcen, AEA Technology, Harwell, Oxfordshire, Report No AEAT-6270, 2002.
- Pirrone, N., Cinnirella, S., Feng, X., Finkelman, R. B., Friedli, H. R., Leaner, J., Mason, R., Mukherjee, A. B., Stracher, G. B., Streets, D. G., Telmer, K.: Global mercury emissions to the atmosphere from anthropogenic and natural sources. *Atmos. Chem. Phys.*, 10, 5951-5964, doi:10.5194/acp-10-5951-2010, 2010.
- Quick, W. J., and Irons, R. M. A.: Trace element partitioning during the firing of washed and untreated power station coals, *Fuel*, 81, 665-672, doi:10.1016/S0016-2361(01)00197-1, 2002.
- Reddy, M. S., Basha, S., Joshi, H. V., and Jha, B.: Evaluation of the emission characteristics of trace metals from coal and fuel oil fired power plants and their fate during combustion, *J. Hazard. Mater.*, 123, 242-249, doi:10.1016/j.jhazmat.2005.04.008, 2005.
- Ren, D. Y., Xu, D. W., and Zhao, F. H.: A preliminary study on the enrichment mechanism and occurrence of hazardous trace elements in the Tertiary lignite from the Shenbei coalfield, China, *Int. J. Coal Geol.*, 57, 187-196, doi:10.1016/j.coal.2003.10.001, 2004.
- Ren, D. Y., Zhao, F. H., Dai, S., and Zhang, J.: Geochemistry of trace elements in coal, Science Press, Beijing, 2006 (in Chinese).
- Skeaff, J. M., and Dubreuil, A. A.: Calculated 1993 emission factors of trace metals for Canadian non-ferrous smelters, *Atmos. Environ.*, 31, 1449-1457, doi:10.1016/S1352-2310(96)00319-6, 1997.
- Song, D. Y., Qin, Y., Zhang, J. Y., and Zheng, C. G.: Volatility of environmentally-sensitive trace elements during coal combustion, *J. Huazhong Univ. of Sci. & Tech. (Nature Science Edition)* 33, 36-38, 2006 (in Chinese with English abstract).
- Song, D. Y., Ma, Y. J., Qin, Y., Wang, W. F., and Zheng, C. G.: Volatility and mobility of some trace elements in coal from Shizuishan Power Plant, *J. Fuel Chem. Technol.*, 39, 328-332, doi:10.1016/S1872-5813(11)60024-8, 2011.

- Song, Y., Xie, S. D., Zhang, Y. H., Zeng, L. M., Salmon, L. G., and Zheng, M.: Source apportionment of PM_{2.5} in Beijing using principal component analysis/absolute principal component scores and UNMIX, *Sci. Total Environ.*, 372, 278–286, doi:10.1016/j.scitotenv.2006.1008.1041, 2006b.
- Streets, D. G., Devane, M. K., Lu, Z., Bond, T. C., Sunderland, E. M., and Jacob, D. J.: All-time releases of mercury to the atmosphere from human activities, *Environ. Sci. Technol.*, 45, 10485–10491, doi:10.1021/es202765m, 2011.
- Sun, Y. L., Zhuang, G. S., Wang, Y., Han, L. H., Guo, J. H., Dan, M., Zhang, W. J., Wang, Z. F., and Hao, Z. P.: The air-borne particulate pollution in Beijing—concentration, composition, distribution and sources, *Atmos. Environ.*, 38, 5991–6004, doi:10.1016/j.atmosenv.2004.5907.5009, 2004.
- Tang, Q., Liu, G. J., Zhou, C. C., and Sun, R. Y.: Distribution of trace elements in feed coal and combustion residues from two coal-fired power plants at Huainan, Anhui, China, *Fuel*, 107, 315–322, doi:10.1016/j.fuel.2013.01.009, 2013.
- Tang, X. Y., Zhao, J. Y., and Huang, W. H.: Nine metal elements in coal of China, *Coal Geol. China*, 14, 43–54, 2002 (in Chinese).
- Tang, Y. G., Yin, Z. R., Chang, C. X., Zhang, Y. Z., Song, H. B., Wang, S., and Hao, L.: Distribution of trace elements in the Kailuan coalfield, *J. China Coal Soc.*, 30, 80–84, 2005 (in Chinese).
- Tang, Y. G., Chang, C. X., Zhang, Y. Z., and Li, W. W.: Migration and distribution of fifteen toxic trace elements during the coal washing of the Kailuan Coalfield, Hebei Province, China, *Energy Explor. Exploit.*, 27, 143–152, doi:10.1260/0144-5987.27.2.143, 2009.
- Tao, J., Zhang, R. J., Duan, J. C., Xing, J. S., Zhu, L. H., Chen, Z. M., Zhao, Y., and Cao, J. J.: Seasonal variation of carcinogenic heavy metals in PM_{2.5} and source analysis in Beijing, *Environ. Sci.*, 35, 411–417, doi:10.13227/j.hjhx.12014.13202.13012, 2014 (in Chinese with English abstract).
- Theloke, J., Kummer, U., Nitter, S., Geftler, T., and Friedrich, R.: Überarbeitung der Schwermetallkapitel im CORINAIR Guidebook zur Verbesserung der Emissionsinventare und der Berichterstattung im Rahmen der Genfer Luftreinhaltekonvention, Report for Umweltbundesamt, 2008.
- Tian, H. Z., Wang, Y., Xue, Z. G., Cheng, K., Qu, Y. P., Chai, F. H., and Hao, J. M.: Trend and characteristics of atmospheric emissions of Hg, As, and Se from coal combustion in China, 1980–2007, *Atmos. Chem. Phys.*, 10, 11905–11919, doi:10.5194/acp-10-11905-2010, 2010.
- Tian, H., Zhao, D., He, M. C., Wang, Y., and Cheng, K.: Temporal and spatial distribution of atmospheric antimony emission inventories from coal combustion in China, *Environ. Pollut.*, 159, 1613–1619, doi:10.1016/j.envpol.2011.02.048, 2011.
- Tian, H. Z., Cheng, K., Wang, Y., Zhao, D., Lu, L., Jia, W. X., and Hao, J. M.: Temporal and spatial variation characteristics of atmospheric emissions of Cd, Cr, and Pb from coal in China, *Atmos. Environ.*, 50, 157–163, doi:10.1016/j.atmosenv.2011.12.045, 2012a.
- Tian, H. Z., Lu, L., Cheng, K., Hao, J. M., Zhao, D., Wang, Y., Jia, W. X., and Qiu, P. P.: Anthropogenic atmospheric nickel emissions and its distribution characteristics in China, *Sci. Total Environ.*, 417, 148–157, doi:10.1016/j.scitotenv.2011.11.069, 2012b.
- Tian, H. Z., Zhao, D., Cheng, K., Lu, L., He, M. C., and Hao, J. M.: Anthropogenic atmospheric emissions of antimony and its spatial distribution characteristics in China, *Environ. Sci.*

Technol., 46, 3973–3980, doi:10.1021/es2041465, 2012c.

Tian, H., Lu, L., Hao, J. M., Gao, J. J., Cheng, K., Liu, K. Y., Qiu, P. P., and Zhu, C. Y.: A review of key hazardous trace elements in Chinese coals: Abundance, occurrence, behavior during coal combustion and their environmental impacts, *Energy Fuels*, 27, 601–614, doi:10.1021/ef3017305, 2013.

United Kingdom (UK): emission factor databases of NAEI, available at: <http://naei.defra.gov.uk/data/> (last access: 8 August 2014), 2000.

United Kingdom (UK): emission factor databases of NAEI, available at: <http://naei.defra.gov.uk/data/> (last access: 21 September 2014), 2013.

US Environmental Protection Agency (US EPA): AP-42, fifth edition, volume 1, Chapter 1.3: available at: <http://www.epa.gov/ttn/chief/ap42/> (last access: 12 October 2014), 1996a.

US Environmental Protection Agency (US EPA): AP 42, fifth Edition, volume 1, chapter 2: Solid Waste Disposal, available at: <http://www.epa.gov/ttn/chief/ap42/ch02/index.html> (last access: 23 July 2014), 1996b.

US Environmental Protection Agency (US EPA): AP 42, fifth edition, volume 1, chapter 11: Mineral Products Industry, available at: <http://www.epa.gov/ttn/chief/ap42/ch11/index.html> (last access: 26 October 2014), 1996c.

US Environmental Protection Agency (US EPA): Web Factor Information Retrieval System (WebFIRE), available at: <http://cfpub.epa.gov/webfire/index.cfm?action=fire.FactorsBasedOnDetailedSearch> (last access: 21 September 2014), 2012.

Wang, Q. C., Shao, Q. C., Kang, S. L., Wang, Z. G., and Zou, S. T.: Distribution of 15 trace elements in the combustion products of coal, *J. Fuel Chem. Technol.*, 24, 137–142, 1996 (in Chinese with English abstract).

Wang, Q. Q., Ma, Y. L., Tan, J. H., Yang, F. M., Wei, L. F., Duan, J. C., and He, K. B.: Characterization of water-soluble heavy metals of PM_{2.5} during winter in Beijing, China, *Environ. Sci.*, 34, 2204–2210, 2014 (in Chinese with English abstract).

Wang, W., Tao, H., Kim, D., and Pan, X. C.: Changes of Elements in PM_{2.5} and PM₁₀ during sand-dust weather in Beijing and Alashan League, *J. Environ. Health* 27, 763–766, 2010 (in Chinese with English abstract).

Wang, W. F., Qin, Y., and Song, D. Y.: Cleaning potential of hazardous elements during coal washing, *J. Fuel Chem. Technol.*, 31, 295–299, 2003a (in Chinese with English abstract).

Wang, W. F., Qin, Y., Sang, S. X., Jiang, B., Guo, Y. H., Zhu, Y. M., and Fu, X. H.: Partitioning of minerals and elements during preparation of Taixi coal, China, *Fuel*, 85, 57–67, doi:10.1016/j.fuel.2005.05.017, 2006.

Wang, Y. F., Huang, K. L., Li, C. T., Mi, H. H., Luo, J. H., and Tsai, P. J.: Emissions of fuel metals content from a diesel vehicle engine, *Atmos. Environ.*, 37, 4637–4643, doi:10.1016/j.atmosenv.2003.07.007, 2003b.

Wang, Y. Q., Ren, D. Y., Lei, J., Tang, Y., Yang, S., and Yang, Y.: Preliminary study on the distribution and characteristic of trace elements in coals, *Scientia Geologica Sinica*, 32, 65–73, 1997 (in Chinese).

Wang, Y. Q., and Mo, J. Y., Ren, D. Y.: Distribution of minor and trace elements in magmatic hydrothermal metamorphic coal of Meitian Coal Mined, Hunan province, *Geochimica*, 28, 289–296, 1999 (in Chinese with English abstract).

- Wei, X. F., Zhang, G. P., Cai, Y. B., Li, L., and Li, H. X.: The volatilization of trace elements during oxidative pyrolysis of a coal from an endemic arsenosis area in southwest Guizhou, China, *J. Anal. Appl. Pyrolysis*, 98, 184-193, doi:10.1016/j.jaap.2012.08.015, 2012.
- Wu, Y., Wang, S. X., Streets, D. G., Hao, J. M., Chan, M., and Jiang, J. K.: Trends in anthropogenic mercury emissions in China from 1995 to 2003, *Environ. Sci. Technol.*, 40, 5312–5318, doi:10.1021/es060406x, 2006.
- Wu, Y. Y., Qin, Y., Yi, T. S., and Xia, X. H.: Enrichment and geochemical origin of some trace elements in high-sulfur coal from Kaili, eastern Guizhou Province, *Geochimica*, 37, 615-622, 2008 (in Chinese with English abstract).
- Xiu, Y., and Wen, H.: Trace elements in Chinese coal, Commercial Press, 2004 (in Chinese).
- Xu, M. H., Zheng, C. G., Feng, R., Qiao, Y., and Yan, R.: Overview of trace elements research in coal combustion process, *Proceedings of the CSEE*, 21, 33-38, 2001 (in Chinese with English abstract).
- Xu, M. H., Yan, R., Zheng, C. G., Qiao, Y., Han, J., and Sheng, C. D.: Status of trace element emission in a coal combustion process: a review, *Fuel Process. Technol.*, 85, 215-237, doi:10.1016/S0378-3820(03)00174-7, 2004.
- Yan, R., and Lu, X.: The distribution of trace elements in several typical coals in China, *Analysis Laboratory*, 14, 43-47, 1995 (in Chinese).
- Yang, J.: Contents and occurrence modes of trace elements in the Late Permian coals from Puan Coalfield, Guizhou Province, *J. Fuel Chem. Technol.*, 34, 129-135, 2006 (in Chinese with English abstract).
- Yang, J., Deng, J., Li, L., Wang, X., and Liu, Z.: Organic residual hydrocarbon-inorganic mineral affinity of trace elements and their influence on coal-formed hydrocarbon, *Acta Geoscientica Sinica*, 29, 235-240, 2008a (in Chinese with English abstract).
- Yang, J., Fu, Q., Guo, X. S., Chu, B. L., Yao, Y. W., Teng, Y. G., and Wang, Y. Y.: Concentrations and seasonal variation of ambient PM_{2.5} and associated metals at a typical residential area in Beijing, China, *Bull. Environ. Contam. Toxicol.*, 94, 232–239, doi:10.1007/s00128-00014-01443-y, 2015.
- Yang, L., Liu, C. Y., and Li, H. Y.: Geochemistry of trace elements and rare earth elements of coal in Chenjiashan coal mine, *Coal Geol. Explor.*, 36, 10-14, 2008b (in Chinese with English abstract).
- Yang, Y. J., Wang, Y. S., Wen, T. X., and Li, L.: Characteristics and sources of elements of atmospheric particles before and in heating period in Beijing, *Environ. Sci.*, 29, 3275–3279, 2008c (in Chinese with English abstract).
- Yang, Y. J., Wang, Y. S., Wen, T. X., and Xu, H. H.: The mass concentration of PM₁₀ and PM_{2.5} in Beijing and their chemical composition characteristics analysis, *Environ. Chem.*, 27, 117–118, 2008d (in Chinese).
- Yao, Q. X., Du, M. L., Wang, S. L., Liu, J., Yang, J. L., and Shang, H. T.: Modes of occurrence and cleaning potential of hazardous trace elements in Huanglong coal, *J. Xian Univ. Sci. Technol.*, 32, 214-220, 2012 (in Chinese with English abstract).
- Yu, L. D., Wang, G. F., Zhang, R. J., Zhang, L. M., Song, Y., Wu, B. B., Li, X. F., An, K., and Chu, J. H.: Characterization and Source Apportionment of PM_{2.5} in an Urban Environment in Beijing, *Aerosol and Air Quality Res.*, 13, 574–583, doi: 10.4209/aaqr.2012.07.0192, 2013.
- Yu, Y., Cen, K., Stefan, N., Nina, S., and Chen, Y.: Concentration characteristics and seasonal

- trend of main heavy metal elements of PM_{2.5} in Beijing, *Geoscience*, 5, 975–982, 2012 (in Chinese with English abstract).
- Zajusz-Zubek, E., and Koniecznyński, J.: Dynamics of trace elements release in a coal pyrolysis process, *Fuel*, 82, 1281–1290, doi:10.1016/S0016-2361(03)00031-0, 2003.
- Zeng, R. S., Zhuang, X. G., Koukouzas, N., and Xu, W. D.: Characterization of trace elements in sulphur-rich Late Permian coals in the Heshan coal field, Guangxi, South China, *Int. J. Coal Geol.*, 61, 87–95, doi:10.1016/j.coal.2004.06.005, 2005.
- Zhang, G. W.: Pollution characteristics and source analysis of elements of the PM_{2.5} in Northeast Beijing, Master Thesis, Shandong Normal University, 2012 (in Chinese with English abstract).
- Zhang, J., Han, C. L., and Xu, Y. Q.: The release of the hazardous elements from coal in the initial stage of combustion process, *Fuel Process. Technol.*, 84, 121–133, doi:10.1016/S0378-3820(03)00049-3, 2003.
- Zhang, L. L., Gao, Y. X., Dao, X., Wang, C., Teng, E. J.: Composition and distribution of elements in air particulate matters during heating season of Beijing-Tianjin-Hebei megacities, China, *Environ. Monit. in China*, 30, 53–61, 2014 (in Chinese with English abstract).
- Zhang, X. L., Zhao, X. J., Pu, W. W., Xu, J.: Comparison of elemental characteristics of suspended particles PM_{2.5} in urban and rural area of Beijing, China *Powder Sci. Technol.*, 16, 28–34, 2010 (in Chinese with English abstract).
- Zhao, J. Y., Tang, X. Y., and Huang, W. H.: Abundance of trace elements in coal of China, *Coal Geol. China*, 14, 5–13, 2002 (in Chinese with English abstract).
- Zhao, P. S., Dong, F., He, D., Zhao, X. J., Zhang, X. L., Zhang, W. Z., Yao, Q., and Liu, H. Y.: Characteristics of concentrations and chemical compositions for PM_{2.5} in the region of Beijing, Tianjin, and Hebei, China, *Atmos. Chem. Phys.*, 13, 4631–4644, doi:10.5194/acp-13-4631-2013, 2013.
- Zhao, Y., Wang, S. X., Duan, L., Lei, Y., Cao, P. F., and Hao, J. M.: Primary air pollutant emissions of coal-fired power plants in China: Current status and future prediction, *Atmos. Environ.*, 42, 8442–8452, doi:10.1016/j.atmosenv.2008.08.021, 2008.
- Zhao, Y., Nielsen, C. P., Lei, Y., McElroy, M. B., and Hao, J. M.: Quantifying the uncertainties of a bottom-up emission inventory of anthropogenic atmospheric pollutants in China, *Atmos. Chem. Phys.*, 11, 2295–2308, doi:10.5194/acp-11-2295-2011, 2011.
- Zhou, J. B., Zhuang, X. G., Alastuey, A., Querol, X., and Li, J. H.: Geochemistry and mineralogy of coal in the recently explored Zhundong large coal field in the Junggar basin, Xinjiang province, China, *Int. J. Coal Geol.*, 82, 51–67, doi:10.1016/j.coal.2009.12.015, 2010.
- Zhu, C. S., and Li, D. H.: Occurrences of trace elements in the No.2 coal of the Changhebian Coal Mine, Chongqing, China, *Acta Metallurgica Sinica*, 28, 259–263, 2009.
- Zhuang, X. G., Yang, S. K., Zeng, R. S., and Xu, W. D.: Characteristics of trace elements in coals from several main coal districts in China, *Geol. Sci. Technol. Inf.*, 18, 63–66, 1999 (in Chinese with English abstract).
- Zhuang, X. G., Querol, X., Zeng, R. S., Xu, W. D., Alastuey, A., Lopez-Soler, A., and Plana, F.: Mineralogy and geochemistry of coal from the Liupanshui mining district, Guizhou, south China, *Int. J. Coal Geol.*, 45, 21–37, doi:10.1016/S0166-5162(00)00019-7, 2000.
- Zhuang, X. G., Querol, X., Plana, F., Alastuey, A., Lopez-Soler, A., and Wang, H.: Determination of elemental affinities by density fractionation of bulk coal samples from the Chongqing coal

649 district, Southwestern China, Int. J. Coal Geol., 55, 103-115,
650 doi:10.1016/S0166-5162(03)00081-8, 2003.