Atmos. Chem. Phys. Discuss., 3, 3411–3449, 2003 www.atmos-chem-phys.org/acpd/3/3411/ © European Geosciences Union 2003



ACPD

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.

Global distribution of total ozone and lower stratospheric temperature variations

W. Steinbrecht¹, B. Hassler¹, H. Claude¹, P. Winkler¹, and R. S. Stolarski²

¹German Weather Service, Hohenpeissenberg, Germany ²NASA, Goddard Space Flight Center, Greenbelt, Maryland, USA

Received: 23 May 2003 - Accepted: 23 June 2003 - Published: 3 July 2003

Correspondence to: W. Steinbrecht (wolfgang.steinbrecht@dwd.de)

| Title Page | | |
|-------------------|--------------|--|
| Abstract | Introduction | |
| Conclusions | References | |
| Tables | Figures | |
| | | |
| | | |
| • | • | |
| Back | Close | |
| Full Screen / Esc | | |
| | | |
| Print Version | | |
| | | |
| Interactive | Discussion | |

Abstract

This study gives an overview of interannual variations of total ozone and 50 hPa temperature. It is based on newer and longer records from the 1979 to 2001 Total Ozone Monitoring Spectrometer (TOMS) and Solar Backscatter Ultraviolet (SBUV) instruments, and on US National Center for Environmental Prediction (NCEP) reanaly-5 ses. Multiple linear least squares regression is used to quantify various natural and anthropogenic influences. For most influences the total ozone and 50 hPa temperature responses look very similar, reflecting a very close coupling. As a rule of thumb, a 10 Dobson Unit (DU) change in total ozone corresponds to a 1 K change of 50 hPa temperature. Large influences come from the linear trend term, up to -30 DU or 10 -1.5 K/decade, from terms related to polar vortex strength, up to 50 DU or 5 K (typical, minimum to maximum), from tropospheric meteorology, up to 30 DU or 3 K, or from the Quasi-Biennial Oscillation (QBO), up to 25 DU or 2.5 K. The 11-year solar cycle, up to 25 DU or 2.5 K, El Niño/Southern Oscillation (ENSO), up to 10 DU or 1 K, are somewhat smaller influences. Stratospheric aerosol after the 1991 Pinatubo eruption 15 lead to warming up to 3 K at low latitudes and to ozone depletion up to 40 DU at high latitudes. Response to QBO, polar vortex strength, and to a lesser degree to ENSO, exhibit an inverse correlation between low latitudes and higher latitudes. Responses to the solar cycle or 400 hPa temperature, however, have the same sign over most of the globe. Responses are usually zonally symmetric at low and mid-latitudes, but 20 asymmetric at high latitudes. There, solar cycle, QBO or ENSO influence position and

1. Introduction

The discovery of the "ozone-hole" over Antarctica in 1984 has triggered enormous interest in stratospheric ozone, both from the general public and from the scientific community. Continuing improvement of our understanding of the various processes

strength of the stratospheric anti-cyclones over the Aleutians and south of Australia.

ACPD

3, 3411–3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.

| Title Page | | |
|------------------------|--------------|--|
| Abstract | Introduction | |
| Conclusions | References | |
| Tables | Figures | |
| | | |
| | | |
| • | • | |
| Back | Close | |
| Full Screen / Esc | | |
| | | |
| Print Version | | |
| Interactive Discussion | | |
| | | |

controlling the stratospheric ozone layer has since been documented by a wealth of publications (Solomon, 1999; Staehelin et al., 2001) and by a series of ozone assessments (SPARC, 1998; WMO, 2003). Production of harmful chlorofluorocarbons was successfully curbed by international agreements, i.e. the Vienna Convention for the

- ⁵ Protection of the Ozone Layer (1985), the Montreal Protocol (1987) and its amendments. A reduction of chlorine loading and a recovery of the ozone layer are expected at some time in this century (Engel et al., 1998; WMO, 2003). However, stratospheric chlorine will remain high for the next years. Bromine, which also destroys ozone, is still increasing. The ozone layer will remain vulnerable. Many details of the expected
- ¹⁰ ozone recovery are unclear, e.g. possible interactions with climate change, or the consequences of increasing bromine (Salby and Callaghan, 2002; Ramaswamy et al., 2001; WMO, 2003). In order to actually "see" a recovery, high quality measurements are needed for many years to come. For the early detection of signs of a recovery, or of a possible worsening, a good quantitative understanding is necessary for the various
- natural processes, which contribute to the variation of ozone levels from year to year. Many studies have investigated variations of ozone and temperature on interannual time scales. This includes trends (Bojkov et al., 1990; Staehelin et al., 2001; Ramaswamy et al., 2001), influences from the QBO (Yang and Tung, 1995; Baldwin et al., 2001), the 11-year solar cycle (Zerefos et al., 1997; Hood, 1997; Labitzke and van Loon, 2000; Lee and Smith, 2003), from El Niño/Southern Oscillation (*ENSO*) (Shiotani, 1992; Zerefos et al., 1992; Reid, 1994), or from meteorological factors (Hood et al., 1999; Appenzeller et al., 2000; Steinbrecht et al., 2001).

The purpose of this study is to give an updated and more global overview of the most relevant influences. Newer and longer data records are used. Rather than focus

on global means or zonal averages, the full geographical distribution is presented. A large fraction of the ozone column resides in the lower stratosphere where ozone and temperature are closely linked (Randel and Cobb, 1994). Therefore, analysis of the ozone column data is complemented by analysis of lower stratospheric temperature data, at 50 hPa. Where possible, a simple conceptual view is presented, linking total

ACPD

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



ozone and lower stratospheric temperature variations to a given influence.

2. Data sources and method

series are the basis of our investigation.

25

The investigation is based on total column ozone data from the series of Total Ozone Mapping Spectrometers (TOMS) and Solar Backscatter Ultraviolet (SBUV and SBUV/2) instruments on various satellite platforms. All data are calibrated to be consistent with the 1996 to 1999 Earth-Probe TOMS instrument. See Stolarski and Hollandsworth (2002) for details. The combined data-set currently provides nearly continuous global coverage from late 1978 to December 2001, on a 5° latitude, 10° longitude grid. For each grid cell and each month of the year climatological mean values were defined by the average value over the entire 1979 to 2001 time period. Monthly mean total ozone anomalies were then calculated for each grid cell by subtracting the climatological mean value for the appropriate month of the year. The resulting anomaly time

The same procedure was applied to lower stratospheric temperature fields at 50 hPa from meteorological reanalyses by the US National Center for Environmental Prediction (NCEP, Kistler et al., 2001). The NCEP data were regridded to the same latitudelongitude grid as the TOMS/SBUV data. Here we generally report on NCEP data from the same 1979 to 2001 time period where TOMS/SBUV data are available. However, most of the results for 50 hPa temperature change very little, when the longer time period 1958 to 2001 is considered.

In order to quantify different influences on total ozone or 50 hPa temperature, we use a multiple regression approach (Steinbrecht et al., 2001). This has become standard practice (Bojkov et al., 1990; SPARC, 1998; WMO, 2003), but here more predictors are used, in particular for meteorological influences. Anomaly time series Y of total ozone or 50 hPa temperature are described as a linear combination of predictor time series (=explanatory variables), each of which accounts for a different influence on

ACPD

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



total ozone or temperature:

$$\begin{split} Y &= c_{TR} TR + c_{FS} FS + c_A A + c_{T(400)} T(400) \\ &+ c_{QBO(10)} QBO(10) + c_{QBO(30)} QBO(30) \\ &+ c_{u(60N)} u(60N) + c_{u(60S)} u(60S) \\ &+ c_{ENSO} ENSO + residual \end{split}$$

Given time series for the explanatory variables trend TR, solar cycle FS, stratospheric aerosol A, 400 hPa temperature anomaly T(400), equatorial wind at 10 and ₅ 30 hPa QBO(10/30), zonal wind anomaly at 60° North and South, u(60N/S), and El Niño/Southern Oscillation, ENSO, as well as observed time series of ozone or temperature at each grid cell, Y, the coefficients c can be determined by linear least squares regression. Global fields for each coefficient are obtained as a result. They describe the geographical distribution of the influence for each explanatory variable. Regression was done separately for seasonal mean anomalies, yielding separate sets 10 of coefficients c_{TB}, c_{FS}, c_{OBO(10)}, and so on, for winter (December, January, February = DJF), spring (March, April, May = MAM), summer (June, July, August = JJA), and fall (September, October, November = SON). Thus a seasonal variation is allowed for. From the "full" set of possible explanatory variables, only those contributing significantly, at the 90% confidence level of partial F-tests, were left in the regression, 15 following the stepwise regression procedure outlined in Draper and Smith (1998).

Obviously Eq. (1) is only a very simple approximation. Non-linear effects, as well as possible couplings between explanatory variables, are neglected. Generally, only an "average" response to each predictor can be obtained. In some years a much stronger or much weaker response must be expected. Nevertheless, our results indicate that

- even this simple approach can still give quite a useful quantification. A more complete picture can be obtained, e.g. with 3-dimensional, fully coupled chemistry globalcirculation model simulations. However, such simulations are much more complex and more difficult to interpret.
- ²⁵ The choice of explanatory variables in Eq. (1) is based on the criteria that all should

ACPD

3, 3411–3449, 2003

(1)

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



be relevant for total ozone from past experience, and that the data should be easy to obtain and should be updated regularly. The linear trend *TR* describes long-term ozone changes related to the more or less linear increase of stratospheric chlorine since the 1970s. Most recent data show slowing chlorine increase, or very recently, even the beginnings of a decrease (WMO, 2003). However, for compatibility with many other studies, the linear increase was used as a good enough simple description. Improvements would be marginal, if a more detailed curve were used (Steinbrecht et al., 2001). For 50 hPa temperature, the linear trend simply describes long-term changes.

10.7 cm solar radio flux *FS* is a common proxy for UV irradiance changes related to the 11-year solar cycle (Zerefos et al., 1997). Monthly mean data observed in Ottawa and Penticton, Canada, were obtained from ftp://ftp.ngdc.noaa.gov/STP/ SOLAR_DATA/SOLAR_RADIO/FLUX.

For influences from the Quasi-Biennial Oscillation (QBO), equatorial zonal winds at 10 and 30 hPa *QBO*(10) and *QBO*(30) from FU Berlin are included (B. Naujokat, priv. comm. 2003). Correct phasing of the QBO related signal is allowed for by using winds at two levels, 10 and 30 hPa, which are out of phase by nearly $\pi/2$ (see Bojkov and Fioletov, 1995).

Large enhancements of stratospheric aerosol after major volcanic eruptions in the tropics affect ozone and temperature by changing heating rates and atmospheric trans-

- ²⁰ ports. Ozone is additionally affected by heterogeneous reactions on the much increased aerosol surface area (Solomon, 1999; Robock, 2000). To account for these effects, the time dependent stratospheric aerosol optical depth *A* is included as one explanatory variable. We use zonal mean data compiled by Sato et al. (2003), which allow for latitudinal variation. For the period of interest, the data are largely based on
- ²⁵ measurements by the Stratospheric Aerosol and Gas Experiments (SAGE I and II). Note that using stratospheric aerosol optical depth measured by lidar at a single location (Garmisch-Partenkirchen in Southern Germany) (Jäger et al., 1995) gives very similar results.

The meteorological situation in the troposphere can have a large influence on ozone

ACPD

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



column and on lower stratospheric temperature (Steinbrecht et al., 1998; Appenzeller et al., 2000). Therefore temperature anomalies at 400 hPa, T(400), were used as one explanatory variable. Temperature in the free troposphere and lower stratosphere is highly correlated with other parameters such as tropopause height, geopotential ⁵ heights, or tropospheric circulation indices. The T(400) anomaly field was derived from NCEP reanalyses.

Influences from the El Niño–Southern Oscillation phenomenon are accounted for by the *ENSO* explanatory variable, obtained from the US National Oceanic and Atmospheric Administration, Climate Prediction Center (NOAA-CPC) (http://www.cpc.ncep. 10 noaa.gov/data/indices/).

The strength of the polar winter vortex is another important explanatory variable. A strong vortex is a prerequisite for the formation of an "ozone hole" (Solomon, 1999). Vortex strength is also related to meridional transports in the Brewer-Dobson circulation (Perlwitz and Graf, 1995; Salby and Callaghan, 2002). Zonal wind anomalies at 50 hPa, 60° North and 60° South from NCEP reanalyses, u(60N/S), are a good measure of

vortex strength, and were included as explanatory variables. Other possibilities are indices for the large scale stratospheric circulation, such as the Arctic Oscillation Index (Baldwin and Dunkerton, 1999), or the divergence of the Eliassen-Palm flux (Salby and Callaghan, 2002). However, we preferred the zonal-winds, because they are easy to obtain from the NCEP reanalyses and seem to work guite well.

Linear least squares regression requires that the explanatory variables are sufficiently independent (uncorrelated). While we found that this is generally fulfilled in our case, there are two important exceptions: 1.) The strength of the polar vortex is related to the phase of the QBO and the solar cycle (Holton and Tan, 1980; Labitzke and van Loon, 2000). 2.) Tropospheric temperatures *T*(400) are highly correlated with El Niño/La Niña over large parts of the globe. When two explanatory variables are highly correlated, the stepwise approach will include only the more significant explanatory.

tory variable, and leave the less significant explanatory variables out of the regression (Draper and Smith, 1998). This avoids technical problems with highly correlated ex-

ACPD

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.

| Title Page | | | |
|------------------------|-------------------|--|--|
| Abstract | Introduction | | |
| Conclusions | References | | |
| Tables | Figures | | |
| | | | |
| | | | |
| | ► | | |
| Back | Close | | |
| Full Scre | Full Screen / Esc | | |
| | | | |
| Print Version | | | |
| Interactive Discussion | | | |
| | | | |

planatory variables. For the critical cases, results obtained with the full set of predictors are compared to results with a reduced set of predictors. Generally, interdependencies between the explanatory variables are small and hardly influence the results.

- The residual in Eq. (1) is typically a first-order autocorrelated noise series (Bojkov et al., 1990; SPARC, 1998). However, since seasonal mean data are used, and since the regression usually describes a very large part of the observed variance, autocorrelation was found to be negligible. One years total ozone or temperature residual has little to do with the residual of the previous year. This simplifies the statistical tests and the stepwise linear regression model. Nevertheless, particularly for the solar cycle, it must be realized that the 23 years of 1979 to 2001 data only cover 3 solar maxima and 2 solar minima. Thus we have less than 23 completely independent samples of the solar cycle effect, but likely more than 3. The statistical significance, i.e. the confidence level, of our results may therefore be overestimated, in particular for the longer timescales of solar cycle or *ENSO*. Since for 50 hPa temperature very similar results are
- ¹⁵ obtained when using 42 years of data (1958 to 2001), there does not seem to be a major problem.

3. Results

20

An important measure for the performance of a regression is R^2 , the ratio of variance described by the regression, i.e. the variance of all terms on the right side of Eq. (1) except for the residual, to observed variance, i.e. to the variance of left side of Eq. (1).

- A perfect regression has $R^2 = 1$ and fully explains the observed variance. Small $R^2 \approx 0$ indicates that the regression explains only a small part of the observed variance. This can happen when random noise, e.g. from the measurement itself, accounts for a large fraction of the observed variance, or when important explanatory variables are missing.
- ²⁵ Generally for our study, R^2 values are higher in winter and spring, and lower in fall, in the respective hemispheres.

As examples for good and relatively poor regression, Fig. 1 shows maps of R^2 for

ACPD

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



Northern Hemisphere winter (DJF) and Northern Hemisphere fall (SON). In DJF, R^2 for total ozone exceeds 0.7 over most of the observed globe, indicating that more than 70 percent of the observed total ozone variance are accounted for by the regression. In some regions the regression works less well, e.g. around 10° North and 10° South, over parts of Russia, or over the Southern Pacific. Regression also works less well for SON, where R^2 falls below 0.4 for a substantial fraction of the Northern Hemisphere, and for some regions in the Southern Hemisphere. An important contributor to the low R^2 regions around 10° latitude, on either side of the equator, is the low natural variance of total ozone there. This makes atmospheric and observational noise more relevant.

- ¹⁰ To a lesser degree the same applies to the lower R^2 over the fall hemispheres. The low R^2 regions over the Southern Pacific in Southern Hemisphere spring (SON), or over Russia in northern winter (DJF) are more pronounced in the R^2 maps for 50 hPa temperature (right panels of Fig. 1).
- For 50 hPa temperature, R^2 is similar for all seasons. R^2 is high (> 0.7) and the regression works well in the tropics and subtropics and in the polar region in winter. However, there are broad bands around 50° to 60° latitude of each hemisphere where R^2 is low (< 0.4). There, the regression does not give a good description. To a lesser degree, these bands also appear in the R^2 for total ozone. These bands might come from variations in position and size of jet-streams or polar vortices, not well accounted to by the present set of explanatory variables. Additional explanatory variables might
- for by the present set of explanatory variables. Additional explanatory variables might improve the regression in this latitude band. However, within the scope of the present study, simple additional explanatory variables could not be found. As a consequence, results for the 50° to 60° latitude bands have to be interpreted with care.

3.1. Linear trend

Figure 2 shows maps of the linear trend term from Eq. (1) for northern and southern spring. This term describes long-term changes in total ozone, which can largely be attributed to chemical ozone depletion by anthropogenic chlorine and bromine (Stae-

3419

Interannual variations of total ozone and temperature

W. Steinbrecht et al.

| Title | Title Page | | |
|------------------------|-------------------|--|--|
| Abstract | Introduction | | |
| Conclusions | References | | |
| Tables | Figures | | |
| 14 | ÞI | | |
| | | | |
| • | ► | | |
| Back | Close | | |
| Full Scr | Full Screen / Esc | | |
| | | | |
| Print Version | | | |
| Interactive Discussion | | | |

helin et al., 2001; WMO, 2003). Total ozone has been decreasing over the last 20 years outside of the tropics. The decrease is larger at higher latitudes. In both hemispheres, the decrease is larger in spring than in fall. By far the largest ozone trends, up to -50 Dobson Units (DU) per decade, are found in the "ozone hole" south of 50° S in south⁵ ern spring. Fairly high ozone trends, up to -20 DU/decade are also found in northern

spring over Siberia. Trends for DJF and JJA (both not shown) are similar to those for MAM and SON, respectively.

Temperature at 50 hPa also shows a significant negative trend over most of the globe, ranging between -0.5 and -2 K per decade. The trend appears to be larger in the Southern Hemisphere, and is generally larger in the tropics and subtropics and

In the Southern Hemisphere, and is generally larger in the tropics and subtropics and in the summer and fall hemispheres. Note that the temperature trends exhibit less seasonal variation than the total ozone trends. Smaller negative temperature trends by 0.5 to 1 K, with a similar geographical distribution are found when the regression is applied to NCEP reanalyses for the longer 1958 to 2001 period, not only the 1979 to 2001 period corresponding to the total ozone record.

The ozone trends presented here are fully consistent with other recent investigations, e.g. Fioletov et al. (2002), or WMO (2003). Apart from the region south of 60° S, where the specific time period considered and questions regarding consistency of the NCEP reanalysis before 1979 appear to be important, our temperature trends are similar
to Ramaswamy et al. (2001). As shown there, the 50 hPa temperature trends are consistent with reduced heating by reduced lower stratospheric ozone, and increased cooling from increasing carbon dioxide and water vapor.

3.2. Meteorological influence

Temperature anomalies at 400 hPa were included in the regression to account for links between tropospheric meteorological parameters and total ozone or lower stratospheric temperature (Steinbrecht et al., 1998; Appenzeller et al., 2000; WMO, 2003). Figure 3 shows the size of ozone and 50 hPa temperature fluctuations attributed to T(400). 3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.

| Title Page | | | |
|------------------------|-------------------|--|--|
| Abstract | Introduction | | |
| Conclusions | References | | |
| Tables | Figures | | |
| | | | |
| | | | |
| • | • | | |
| Back | Close | | |
| Full Scre | Full Screen / Esc | | |
| Print Version | | | |
| Interactive Discussion | | | |

As size of an influence, here and in most following plots and discussions, we define 2 standard deviations of the corresponding time series term $c_X X$ on the right side of Eq. (1). This size is close to the average minimum-to-maximum variation attributed to a specific influence. The color plots in Fig. 3, and the following, show this size as a function of latitude and longitude over the globe. Red and yellow colors indicate positive correlation between ozone, or temperature, and explanatory variable X. Blue colors (negative values) indicate inverse correlation. Note that the regression coefficients c_X can be estimated by dividing the 2 standard deviations of $c_X X$, given in Fig. 3 and the following, by 2 standard deviations of the corresponding explanatory variable X, given

10 in Table 1.

5

As Fig. 3 shows, both ozone and 50 hPa temperature fluctuations are inversely correlated to tropospheric temperature. The size of this influence generally increases with latitude. In the tropics little or no ozone fluctuations are associated with tropospheric temperature changes. Fluctuations associated with 400 hPa temperature are largest

¹⁵ in spring at northern high latitudes, where they reach up to 30 DU or 3 K. The overall picture is, however, similar for all seasons and both hemispheres. Total ozone is low and the lower stratosphere is cold when the troposphere is warm, whereas total ozone and lower stratospheric temperature are high when the troposphere is cold.

This behavior can be explained by up- and downward air motions in the lower strato-

- ²⁰ sphere, which compensate opposite vertical motions in the troposphere (Reed, 1950; Steinbrecht et al., 1998; Labitzke and van Loon, 1999). In a tropospheric high pressure system, sinking air in the troposphere leads to adiabatic warming there. The tropopause and air in the lower stratosphere rise. This results in adiabatic cooling in the lower stratosphere and moves ozone poor air up. Total ozone decreases. A
- tropospheric low pressure system has vertical motions in the opposite direction, and consequently a cold troposphere and a warm, ozone rich lower stratosphere with high total ozone. Note that increased/decreased radiative heating by more/less ozone also tends to give a positive correlation between total ozone and lower stratospheric ozone, especially on longer time-scales (Ramaswamy et al., 2001).

ACPD

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



The QBO with its downwards propagating zonal wind anomalies is an intriguing example of wave mean-flow interaction in the atmosphere. It has far-ranging effects that can be observed in many atmospheric parameters (Baldwin et al., 2001). Size and

- ⁵ geographic distribution of ozone and 50 hPa temperature fluctuations attributed to the QBO by the regression in this study are shown in Fig. 4. When equatorial zonal wind is westerly at 30 hPa, total ozone and 50 hPa temperature are high in the tropics (yellow and red colors) and are low (green and blue colors) in a subtropical belt from about 20° to 40° or 50°. In the winter hemisphere, subtropical ozone and temperature anomalies
- are stronger, up to -25 DU or -2.5 K, than in the summer hemisphere, where they only reach up to -5 DU or -0.5 K. Ozone anomalies in the summer hemisphere also extend to higher latitudes. Size, up to 25 DU or 2.5 K, and annual variation, much larger signal in the winter/spring hemisphere, of the QBO response found here agree well with previous results summarized by Baldwin et al. (2001).
- The tropical and subtropical anomalies can be explained by a residual circulation with descending/ascending motion in the tropics and ascending/descending motion in the subtropics (Andrews et al., 1987; Baldwin et al., 2001). According to the thermal wind relation, these vertical motions are induced by the westerly/easterly vertical wind shear associated with the QBO. For the case of westerly wind shear, e.g. when max-
- imum westerly winds are found at or above 30 hPa, descending motion occurs below 30 hPa. This brings ozone rich air down in the tropical lower stratosphere, thereby increasing tropical total ozone and adiabatically warming the tropical lower stratosphere. This explains the yellow and red band of positive anomalies around the Equator in Fig. 4. In the subtropics, ascending motion brings ozone poor air up, thereby decreasing the point of the subtropics.
- ing subtropical total ozone and adiabatically cooling the subtropical lower stratosphere. These regions appear as blue bands in Fig. 4. In the winter hemisphere, the subtropical anomaly is enhanced by the strong residual Brewer-Dobson circulation in winter and spring (Baldwin et al., 2001).

ACPD

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.

| Title Page | | |
|------------------------|--------------|--|
| Abstract | Introduction | |
| Conclusions | References | |
| Tables | Figures | |
| | | |
| | ►I | |
| • | ► | |
| Back | Close | |
| Full Screen / Esc | | |
| | | |
| Print Version | | |
| Interactive Discussion | | |

While the tropical and subtropical QBO response is zonally symmetric, QBO-related fluctuations at higher latitudes in the winter hemispheres, between 40° and 80°, are not zonally symmetric: High 50 hPa temperatures, i.e. the red or yellow regions in Fig. 4 occur during QBO-westerlies at 30 hPa from Europe to Northeastern Asia in 5 DJF (and MAM, not shown), north of North-America in SON (not shown), and above the

- Southern Atlantic Ocean in JJA (and SON as well, not shown). Where TOMS/SBUV data are available, the total ozone maps also indicate a response in these regions. These higher latitude features are also out of phase with the subtropical and in phase with the tropical signal. Phase changes in QBO-response at latitudes around 40° N or
- ¹⁰ 50° N have been noted previously (Bojkov and Fioletov, 1995; Yang and Tung, 1995). Yang and Tung argued that phase changes appear because high-latitude total ozone anomalies in some years have the same sign as anomalies in mid-latitudes. In other years they have the opposite sign. When projected onto any QBO-predictor, these "missing" beats produce an apparent phase change. Irregular "missing" beats, not accounted for by the regression, would also create a band of low R^2 , as seen in Fig. 1.
- Because of the inclusion of zonal wind anomalies at 60°, 50 hPa, as one explanatory variable in the regression (see Figs. 9 and 10), QBO modulations related to polar vortex strength do not appear in Fig. 4. Therefore, Fig. 5 shows results of a regression where trend, solar cycle and QBO were used as the only explanatory variables. For most seasons and latitudes, the QBO signal for this reduced set of explanatory variables is very similar to the QBO signal obtained with the full set of explanatory variables in Fig. 4. The main difference between the two sets of explanatory variables is a dipole between low polar temperatures north of 60° N and high temperatures in a band around 40° N, that are seen in Fig. 5 for the reduced set of explanatory variables in DJF and SON (not shown), during QBO-westerlies at 30 hPa. Also, high temperatures and high total ozone around 40° are more pronounced and more zonally symmetric in Fig. 5,
- compared to Fig. 4. As shown later in Fig. 9, such a see-saw between high and lower latitudes is the signature of polar vortex strength. Because the QBO modulates the strength of the polar vortex, parts of the effects of polar vortex strength on total ozone

ACPD

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



and temperature are taken over by the QBO explanatory variable, when no explanatory variable for polar vortex strength is included in the regression. This is the case for Fig. 5.

- Colder polar temperatures and a stronger polar vortex during times of QBO⁵ westerlies (e.g. at 30 hPa) are well documented (Holton and Tan, 1980). A strong polar vortex goes hand in hand with a weaker meridional circulation, which means lower temperature and low total ozone inside the polar vortex and higher temperature and high total ozone at mid-latitudes. All these features of the *Holton-Tan* relation between polar vortex strength and the QBO can be seen in Fig. 5. However, the *Holton-Tan*relation is true in a statistical sense only. It does not hold for every winter. As pointed out by Labitzke and van Loon (2000) the *Holton-Tan* relation seems to be valid mainly for minima of the 11-year solar cycle. The reverse, i.e. warm polar temperatures for QBO-westerlies, seems to occur during solar maxima. The QBO is therefore only a
- weak predictor for polar vortex strength. The full regression correctly attributes the
 high-latitude anomalies to the more direct predictor polar vortex strength (zonal wind at 60°), not to the QBO.

An interesting feature of Fig. 4 is the zonally asymmetric QBO signal (red and yellow areas) over Northeastern Asia. It can be interpreted as a strengthening or westward extension of the stratospheric Aleutian anti-cyclone correlated with QBO-westerlies at

- 30 hPa. The feature starts in SON (not shown) as an eastward shift to Northern North-America. In DJF the stratospheric Aleutian anti-cyclone has become stronger above Eastern Asia and this anomaly persists into MAM (not shown). A similar, but weaker feature appears over the Southern Atlantic in Southern Hemisphere winter (JJA), where a stratospheric anti-cyclone is usually found south of Australia and New-Zealand. This
- ²⁵ anti-cyclone also seems to become stronger in its western part, or extending westwards, during QBO-westerlies at 30 hPa.

ACPD

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.

| Title Page | | |
|-------------------|--------------|--|
| Abstract | Introduction | |
| Conclusions | References | |
| Tables | Figures | |
| | | |
| | | |
| • | ► | |
| Back | Close | |
| Full Screen / Esc | | |
| | | |
| Print Version | | |
| | | |

3.4. Solar cycle effects

As pointed out by Labitzke and van Loon (2000), or Ruzmaikin and Feynman (2002), the atmospheric response to the 11-year solar cycle differs between westerly and easterly phases of the QBO. To account for this, regression was done separately for two

- ⁵ subsets of the data. One subset contains only data during times of westerly equatorial zonal wind, the other subset only during easterly winds (at 50 hPa). Because of splitting into two subsets, no QBO explanatory variable is included in the regression. Note that following Labitzke and van Loon, 50 hPa winds were used to separate easterly and westerly phases of the QBO, whereas throughout the rest of the paper 30 hPa
- winds, typically 4 months ahead of 50 hPa winds, are used as the reference. For improved clarity, we also decided to lower the confidence level to 80%, and to show only results for a regression where polar vortex strength was not included as an explanatory variable. Thus, the results presented here are compatible with most of Labitzke's investigations. Note that results outside of the polar regions change very little, when the
- full set of explanatory variables, including zonal winds at 60° N and 60° S, is used. Also, results for 50 hPa temperature change very little when the full 1958 to 2001 period is used, instead of the normal 1979 to 2001 period of TOMS/SBUV data.

Figures 6 and 7 show regression results for the two QBO sub-sets. As usual, total ozone and 50 hPa temperature show very similar features. This is more pronounced

- than for the complete data-set, without grouping. Results for the complete data-set without grouping are similar to the average from both sub-sets. Generally, total ozone and 50 hPa temperature are higher during solar maxima and lower during solar minima. The differences in Figs. 6 and 7 typically reach 5 to 25 DU or 0.5 to 2.5 K. The total ozone and temperature responses look very similar. Both responses are usually
- smaller in the tropics than in the extra tropics. From the tropics to mid-latitudes, up to 40° or 50°, and for both QBO subsets, the solar cycle signals are more or less zonally symmetric. The solar signal is stronger during easterly QBO phases, weaker during westerly phases. At low to mid-latitudes, the solar cycle response is similar in both

ACPD

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.

| Title Page | | | |
|------------------------|-------------------|--|--|
| Abstract | Introduction | | |
| Conclusions | References | | |
| Tables | Figures | | |
| 14 | | | |
| | | | |
| • | • | | |
| Back | Close | | |
| Full Scre | Full Screen / Esc | | |
| | | | |
| Print Version | | | |
| Interactive Discussion | | | |

hemispheres during QBO-East, whereas it is substantially larger in the summer hemisphere during QBO-West. While SON (not shown) shows a similar picture as DJF, the solar cycle response during MAM (not shown) is quite different from both DJF and JJA. The MAM response is quite weak and mostly restricted to high latitudes, during
 ⁵ QBO-East, whereas during QBO-West it is strong. and similar to, but weaker than, the DJF response for QBO-East.

Similar to the findings for the QBO in Figs. 4 and 5, the high latitude response to the solar cycle in Figs. 6 and 7 is not zonal, but localized to certain longitudes. Where total ozone data are available, they show similar high latitude features as 50 hPa tem-¹⁰ perature. Between QBO-East and QBO-West, the high latitude responses appear at complementary longitudes: For QBO-West and DJF, 50 hPa temperatures are higher during solar maxima over Europe, and lower over the Northern Pacific (red and blue regions in Fig. 7). However, for QBO-East and DJF, 50 hPa temperatures are lower during solar maxima over parts of Northern Russia and higher in the North Pacific Re-¹⁵ gion (blue and red areas in Fig. 6). A complementary picture is also seen for JJA in the

Southern Hemisphere. There, 50 hPa temperatures are higher during solar maxima in a region south of South America during QBO-East, but are lower during solar maxima over the Southern Pacific during QBO-West.

These results for total ozone and 50 hPa temperature are broadly consistent with ²⁰ previous results for stratospheric temperatures or total ozone (Zerefos et al., 1997; Hood, 1997; Labitzke and van Loon, 2000). In particular it is confirmed that solar effects on the atmosphere differ from season to season and differ between westerly and easterly phases of the QBO (Labitzke and van Loon, 2000; Ruzmaikin and Feynman, 2002). As pointed out by Lee and Smith (2003), regression on data spanning only

²⁵ a few solar cycles probably cannot separate QBO and solar cycle effects completely. Conceptual models indicate that QBO and solar cycle can interact with atmospheric noise to change the probability for warm or cold stratospheric winters (Lawrence et al., 2000). According to Hood (1997), observed changes in lower stratospheric dynamics can explain the total ozone fluctuations and their similarity to the temperature 3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



fluctuations. However, general circulation models even including a representation of atmospheric chemistry, largely have not been able to reproduce the full size of the variations observed in the lower stratosphere (Labitzke et al., 2002; Tourpali et al., 2003), particularly at higher latitudes. A deeper understanding of the underlying physical mechanisms is still lacking.

3.5. Stratospheric aerosol effects

5

Since the beginning of regular stratospheric weather records, three major volcanic eruptions, Mt. Pinatubo in 1991, El Chichón in 1982, and Agung in 1963, have resulted in large increases of stratospheric aerosol. Most of the volcanic aerosol leaves
the stratosphere through sedimentation within about 2 to 3 years. Light absorption and scattering by the additional particles tends to warm the lower latitude stratosphere. Particularly at cold temperatures and higher latitudes, the particles also provide surface area for chemical reactions, which result in enhanced ozone destruction by chlorine. This was most pronounced for the Pinatubo episode, when stratospheric chlorine
loading was higher than at the time of the other eruptions (Solomon, 1999; Robock, 2000).

Based on the 1979 to 1995 total ozone record from SBUV, McCormack et al. (1997) pointed out that it may be difficult to separate between effects of the 11-year solar cycle and effects from volcanic aerosol, since both major volcanic eruptions of El Chichón in

- ²⁰ 1982 and Pinatubo in 1991 occurred near solar maximum. Extension of the data record to 2001, almost to the next solar maximum and without another volcanic eruption, seems to have alleviated this problem. In the present study nearly the same results are obtained for stratospheric aerosol effects on total ozone and 50 hPa temperature, whether the solar-cycle is included in the regression or not. Similarly, results for the
- stratospheric solar-cycle predictor do not depend much on whether the aerosol term is included or not. Also, influence of the stratospheric aerosol term or of the solar cycle term on 50 hPa temperature change very little when the much longer 1958 to 2001 data-set is used.

ACPD

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



The maximum ozone and 50 hPa temperature changes attributed to stratospheric aerosol loading are shown in Fig. 8 for northern spring (MAM) and fall (SON). Since the Mt. Pinatubo eruption resulted in the highest stratospheric aerosol loading for over 50 years, the values in Fig. 8 largely reflect anomalies from the years 1991 to 1993.

- ⁵ Different from most other explanatory variables, there is little geographical correlation between the aerosol responses for total ozone and 50 hPa temperature. Temperatures at 50 hPa responded by an increase, typically by 2 to over 3 K, in a broad belt between 40° S and 40° N. The response is very similar for all seasons. At higher latitudes, red areas in Fig. 8 above North America and south-east of Australia in SON give in-
- dications for strengthening and/or eastward movement of the Aleutian and Australian stratospheric anti-cyclones. Similar high-latitude features are also seen in winter in the respective hemispheres (DJF and JJA, both not shown). There is no indication of a significant negative temperature response at mid-latitudes.

In contrast to 50 hPa temperature, the ozone response is usually negative. It is ¹⁵ largest at high latitudes, smallest in the tropics. Ozone decreases exceeding 40 DU are found at northern mid to high latitudes in spring (MAM) and over parts of Antarctica in southern spring (SON). The ozone response for DJF (not shown) is very similar to MAM. In JJA (not shown) losses are found at lower latitudes like in SON, and the large northern high-latitude losses from MAM remain, but are substantially reduced.

- The aerosol responses found here, a 50 hPa temperature increase by up to 3 K at low latitudes, and total ozone depletions by up to 40 DU, mostly at northern high-latitudes, are consistent with previous studies (Angell , 1997; McCormack et al., 1997; Robock, 2000).
 - 3.6. Zonal wind anomalies at 60° N and 60° S, 50 hPa
- The intensity of the meridional residual Brewer-Dobson circulation in the winter stratosphere is highly correlated with the strength of the polar winter vortex (Staehelin et al., 2001; Salby and Callaghan, 2002). A stable winter vortex is also a prerequisite for large-scale chemical processing on polar stratospheric cloud particles leading to

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.

| Title Page | | |
|------------------------|--------------|--|
| Abstract | Introduction | |
| Conclusions | References | |
| Tables | Figures | |
| | | |
| | P 1 | |
| • | • | |
| Back | Close | |
| Full Screen / Esc | | |
| | | |
| Print Version | | |
| Interactive Discussion | | |
| | | |

"ozone-hole" conditions (Solomon, 1999). Zonal wind anomalies at 60° latitude at the 50 hPa level are a good indicator of vortex strength and were added to the set of explanatory variables. The size of total ozone and 50 hPa temperature fluctuations attributed to zonal wind anomalies is shown in Figs. 9 and 10 for northern and southern 5 spring.

For high winds and a strong polar vortex, total ozone and temperatures are low in the polar region. The polar response is very strong and reaches up to 50 DU or 5 K. Polar total ozone and 50 hPa temperature are inversely correlated to zonal wind. An opposite response is found at mid- to low latitudes. Temperature responds stronger and at higher latitudes than total ozone. The temperature response peaks with up to 2.5 K around 30° to 40°. The total ozone response is more restricted to the tropics and reaches up to 5 DU. The Southern Hemisphere response (SON) is slightly weaker, but more spread out, than the Northern Hemisphere response. Results for northern fall and winter (SON and DJF, both not shown) are very similar to northern spring (MAM).

¹⁵ However, effects of southern vortex strength in southern fall and winter (MAM and JJA, both not shown), are much less pronounced than in southern spring (SON). This reflects the much stronger and less variable Antarctic vortex and the generally weaker Brewer-Dobson circulation in the Southern Hemisphere during winter.

The responses of total ozone and 50 hPa temperature reflect the close coupling of mean meridional Brewer Dobson circulation and polar vortex strength. A strong vortex goes hand in hand with a weak meridional circulation. Less ozone and heat are transported polewards. Total ozone and temperatures remain high at lower latitudes and low at higher latitudes. A strong and cold vortex also favors rapid chemical ozone destruction in spring ("ozone hole"), which will result in low total ozone and, through less ozone radiative heating, also lower 50 hPa temperatures. This effect will enhance the transport effect. It is more relevant for the Northern Hemisphere where "ozone hole" conditions only occur in some years, than for the Southern Hemisphere, where a strong and stable vortex is established in almost every year.

ACPD

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.

| Title Page | | |
|------------------------|--------------|--|
| Abstract | Introduction | |
| Conclusions | References | |
| Tables | Figures | |
| | | |
| | | |
| • | • | |
| Back | Close | |
| Full Screen / Esc | | |
| | | |
| Print Version | | |
| Interactive Discussion | | |

3.7. ENSO

The El Niño/Southern-Oscillation phenomenon (*ENSO*) is a coupled fluctuation of the tropical Pacific Ocean and the atmosphere (e.g., Coghlan, 2002). During the El Niño warm-phase, ocean surface temperature in the Eastern Tropical Pacific is 1 to 2 K

- ⁵ above normal, during the La Niña cold-phase it is 1 to 2 K below normal. These changes in sea surface temperature are accompanied by changes in many atmospheric parameters, e.g. tropospheric winds and rainfall patterns. During El Niño, tropical easterlies weaken, dry conditions are found in the Western Pacific (Australia, Indonesia), wet conditions extend further into the Eastern Pacific (South America).
- The warmer sea surface temperatures also lead to a warmer troposphere and a higher tropopause throughout most of the tropics, but particularly over the Eastern Tropical Pacific. In the extra tropics, particularly over regions in Eastern Asia and the Pacific, temperatures and tropopause height are lower (Kiladis et al., 2001). The opposite is the case for La Niña. *ENSO* is accompanied by residual circulation cells with descending
- and ascending branches in the troposphere. These are compensated by opposing motions in the lower stratosphere (Gettelman et al., 2001; Kiladis et al., 2001). ENSO thus leads to lower stratospheric temperature and total ozone anomalies, that are opposite to the temperature anomalies in the troposphere.

Figure 11 shows the typical size of total ozone and 50 hPa temperature changes between La Niñas and El Niños for DJF. Note that the *ENSO* explanatory variable used here (NCEP's Southern Oscillation Index) is positive for La Niña events and negative during El Niños. Thus, Fig. 11 shows that total ozone and 50 hPa temperature are high in DJF during La Niña throughout parts of the tropics, particularly above the Eastern Pacific and the tropical Atlantic for total ozone. In the tropics, the difference between

La Niña and El Niño is typically about 5 to 10 DU or 0.5 to 1 K. The signal is strongest in DJF and above the Eastern Pacific. SON and MAM (both not shown) exhibit a similar, generally weaker and less significant, response. A similar tropical response to *ENSO*, i.e. zonal changes up to 10 DU with a superimposed see-saw between the

ACPD

3, 3411–3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



Eastern and Western Pacific, has been reported, e.g. by Shiotani (1992). Outside of the tropics, total ozone and 50 hPa temperature are generally low during La Niña, and high during El Niño. This is most pronounced in the summer hemisphere (DJF and JJA, the latter is not shown).

- A striking, zonally asymmetric, feature are the substantially colder 50 hPa temperatures found during La Niña over Eastern Asia (-3 K, blue region in Fig. 11). They seem to be coupled with warmer temperatures over Europe (+3 K, red region over Europe). A positive anomaly over Europe is also found for total ozone. However, the strong temperature anomaly over Eastern Asia does not appear in the total ozone response, or is not significant there. The temperature anomalies indicate a weakening
- of the westerly part of the Aleutian anti-cyclone. This might be related to a generally stronger and more stable Arctic polar vortex during La Niña (Labitzke and van Loon, 1999). As before for QBO and solar cycle, a strong polar signal from *ENSO* is likely taken over by the zonal wind at 60° explanatory variable.
- As mentioned, tropospheric temperature, e.g. at 400 hPa, and El Niño are highly correlated in the tropics and some other regions. When both 400 hPa temperature and an El Niño explanatory variable are used in the regression, one, the other, or both may take over part of the signal. To give an indication how serious this dependence might be, Fig. 11 contrasts *ENSO* results for the full regression according to Eq. (1) in
- ²⁰ the upper panels, with results where T(400) was not included as an explanatory variable in the lower panels. Differences between the two sets of explanatory variables are generally small and occur mainly in the tropics. When 400 hPa temperature is included (upper panels) the total ozone and temperature response to the *ENSO* predictor is weaker in the tropics, particularly over the Eastern Pacific, *ENSO*'s main center of
- ²⁵ action. In this region, 400 hPa temperature takes over part of the response from the *ENSO* explanatory variable. Similar but smaller changes are found for the other seasons (not shown). Although T(400) and *ENSO* are an example of highly correlated explanatory variables, the results do not depend very much on whether the other explanatory variable is included or not. For other explanatory variables the effects are

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.

| Title | Title Page | | |
|------------------------|--------------|--|--|
| Abstract | Introduction | | |
| Conclusions | References | | |
| Tables | Figures | | |
| 14 | | | |
| | | | |
| • | ► | | |
| Back | Close | | |
| Full Screen / Esc | | | |
| | | | |
| Print Version | | | |
| Interactive Discussion | | | |

usually even smaller.

4. Conclusions

Multiple linear regression analysis provides a good near global picture of natural and anthropogenic influences on interannual variations of total ozone and lower stratospheric temperature, e.g. at 50 hPa. For most influences the geographical distribution of the total ozone and 50 hPa temperature responses is very similar. It appears that interannual fluctuations of both parameters are controlled to a large degree by the same physical processes. As a rule of thumb, a 10 DU change in total ozone corresponds to a 1 K change of 50 hPa temperature. The length of the 1979 to 2001 TOMS/SBUV data record, and the even longer 1958 to 2001 NCEP reanalysis, make it possible to get a fairly good separation of different influences. There may still be difficulties, e.g. separating solar cycle and QBO influences, or QBO and *ENSO* influences (Lee and Smith, 2003). However, our sensitivity test indicate no major problems. Although new data have been added and the picture presented here has become more comprehensive, findings are not fundamentally different from earlier studies where the various

influences were studied separatly (Shiotani, 1992; McCormack et al., 1997; Zerefos et al., 1997; Labitzke and van Loon, 2000; Baldwin et al., 2001).

For the different explanatory variables, ozone and temperature fluctuations range from about 5 DU or 0.5 K (two standard deviations of the corresponding time series
term in Eq. (1)) to 50 DU or 5 K, respectively. Large influences come from the linear trend term, up to -30 DU or -1.5 K/decade, from terms related to polar vortex strength, up to 50 DU or 5 K, from tropospheric meteorology, up to 30 DU or 3 K, or from the QBO, up to 25 DU or 2.5 K. The 11-year solar cycle, up to 25 DU or 2.5 K, or *ENSO*, up to 10 DU or 1 K, are somewhat smaller influences. A large but sporadic influence comes from stratospheric aerosol, which leads to warming up to 3 K at low latitudes and ozone depletion up to 40 DU at high latitudes. Response to QBO, polar vortex strength, and to a lesser degree, to *ENSO* exhibit a clear inverse correlation between

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



low latitudes and higher latitudes. Responses to solar cycle or 400 hPa temperature, however, have the same sign over most of the globe. The obtained quantification of the various influences should help in the detection of the expected recovery of total ozone. However, given the long stratospheric residence time of atmospheric chlorine and uncertainties in the various influences, not discussed much in this paper, it is unlikely that unambiguous signs of a recovery will be seen before 2015 or 2020 (see

unlikely that unambiguous signs of a recovery will be seen before 2015 or 2020 (see also WMO, 2003).

For most explanatory variables, the total ozone and 50 hPa temperature response is zonally symmetric at low and mid-latitudes. At high latitudes, however, zonal symmetry is more the exception than the rule. For example, QBO, solar-cycle or *ENSO*, all related

- to polar vortex strength and stratospheric warmings (Labitzke and van Loon, 1999), seem to influence position and strength of the stratospheric anti-cyclones over the Aleutians and south of Australia. This indicates that latitudinal effects need to be considered, when trying to understand the physical mechanisms leading to the observed
- ¹⁵ connections at higher latitudes (see also Callaghan and Salby, 2002). Two-dimensional models cannot resolve latitudinal structures, and this may be a major drawback. Recently, several three-dimensional global circulation models include a representation of stratospheric chemistry with feedback of the chemical species on radiative heating and cooling (Austin et al., 2003). Two such models, MA-ECHAM and ECHAM-DLR, are
- ²⁰ currently used for transient simulations. These simulations account for source gas changes over the last 40 years, for solar-flux changes, for changing sea-surface temperatures, QBO and volcanic aerosol. It is planned to analyze these simulations in the same fashion as has been done for the observations in this paper. Comparison between simulated and observed effects of the various influence, should help towards
- ²⁵ a much better understanding of the underlying mechanisms.

Acknowledgements. This investigation is part of the KODYACS project, funded under AF02000 (www.afo2000.de) by the German Ministry for Education and Research, grant ATF43.

ACPD

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.

| Title Page | | |
|------------------------|--------------|--|
| Abstract | Introduction | |
| Conclusions | References | |
| Tables | Figures | |
| | | |
| | | |
| • | • | |
| Back | Close | |
| Full Screen / Esc | | |
| Drink) (aming | | |
| Print Version | | |
| Interactive Discussion | | |

References

5

20

Andrews, D. G., Holton, J. R., and Leovy, C. B., Middle Atmosphere Dynamics, Academic Press, San Diego, USA, 481, 1987. 3422

Angell, J. K.: Stratospheric warming due to Agung, El Chichón, and Pinatubo taking into account the Quasi-Biennial Oscillation, J. Geophys. Res., 102, 9479–9485, 1997. 3428

- Appenzeller, C., Weiss, A. K., and Stähelin, J.: North Atlantic Oscillation modulates total ozone winter trends, Geophys. Res. Lett., 27, 1134–1138, 2000. 3413, 3417, 3420
- Austin, J. A., Shindell, D., Beagley, S. R., Brühl, C., Dameris, M., Manzini, E., Nagashima, T., Newman, P., Pawson, S., Pitari, G., Rozanov, E., Schnadt, C., and Shepherd, T. G.:
- ¹⁰ Uncertainties and assessments of chemistry-climate models of the stratosphere, Atm. Chem. Phys., 3, 1–27, 2003. 3433
 - Baldwin, M. P. and Dunkerton, T. J.: Propagation of the Arctic Oscillation from the stratosphere to the troposphere, J. Geophys. Res., 104, 30 937–30 946, 1999. 3417

Baldwin, M. P., Gray, L. J., Dunkerton, T. J., Hamilton, K., Haynes, P. H., Randel, W. J., Holton,

- J. R., Alexander, M. J., Hirota, I., Horinouchi, T., Jones, D. B. A., Kinnersley, J. S., Marquardt, C., and Takahashi, M.: The Quasi-Biennial Oscillation, Rev. Geophys., 39, 179–229, 2001.
 3413, 3422, 3432
 - Bojkov, R. D., Bishop, L., Hill, W. J., Reinsel, G. C., and Tiao, G. C.: A statistical trend analysis of revised Dobson Total Ozone data over the Northern hemisphere, J. Geophys. Res., 95, 9785–9807, 1990. 3413, 3414, 3418
 - Bojkov, R. D., and Fioletov, V. E.: Estimating the global ozone characteristics during the last 30 years, J. Geophys. Res., 100, 16,537–16,551, 1995. 3423
 - Callaghan, P. F. and Salby, M. L.: Three-Dimensionality and forcing of the Brewer-Dobson circulation, J. Atmos. Sci., 59, 976-991, 2002.
- ²⁵ Coughlan, C.: El Niño causes, consequences and solutions, Weather, 57, 209–215, 2002. 3430
 - Draper, N. R. and Smith, H.: Applied Regression Analysis, Wiley-Interscience, New York, 706, 1998. 3415, 3417

Engel, A., Schmidt, U., and McKenna, D.: Stratospheric trends of CFC-12 over the past two

30 decades: recent observational evidence of declining growth rates, Geophys. Res. Lett., 25, 3319–3322, 1998. 3413

Fioletev, V. E., Bodeker, G. E., Miller, A. J., McPeters, R. D., and Stolarski, R.: Global and zonal

3, 3411–3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.

| Title Page | | | | |
|-------------------|----------------------|--|--|--|
| Abstract | Introduction | | | |
| Conclusions | References | | | |
| Tables | Figures | | | |
| | | | | |
| I 4 | ▶ | | | |
| • | • | | | |
| Back | Close | | | |
| Full Screen / Esc | | | | |
| Full Scr | een / Esc | | | |
| Full Scr | een / Esc | | | |
| Full Scr Print | een / Esc Version | | | |

total ozone variations estimated from ground-based and satellite measurements: 1964-2000, J. Geophys. Res., 107, 4647, doi:10.1029/2001JD001350, 2002. 3420

- Gettelman, A., Randel, W. J., Massie, S., Wu, F., Read, W. G., and Russell III, J. M.: El Niño as a natural experiment for studying the Tropical Tropopause Region, J. Clim., 14, 3375–3392, 2001. 3430 5
 - Holton, J. R. and Tan, H.-C.: The influence of the equatorial guasi-biennial oscillation on the global circulation at 50 mb, J. Atmos. Sci., 37, 2200-2208, 1980. 3417, 3424
 - Hood, L. L.: The solar cycle variation of total ozone: Dynamical forcing in the lower stratosphere, J. Geophys. Res., 102, 1355–1370, 1997. 3413, 3426
- 10 Hood, L. L., Rossi, S., and Beulen, M.: Trends in lower stratospheric zonal winds, Rossby wave breaking behavior, and column ozone at northern mid latitudes, J. Geophys. Res., 104, 24 321-24 339, 1999, 3413
 - Jäger, H., Uchino, O., and Nagai, T.: Ground-based remote sensing of the decay of the Pinatubo eruption cloud at three Northern Hemisphere sites, Geophys. Res. Lett., 22, 607-610, 1995. 3416

15

Kiladis, G. B., Straub, K. H., Reid, G. C., and Gage, K. S.: Aspects of interannual and intra seasonal variability of the tropopause and lower stratosphere, Q. J. Roy. Met. Soc., 127, 1961–1983, 2001. 3430

Kistler, R., Kalnay, E., Collins, W., Saha, S., White, G., Woollen, J., Chelliah, M., Ebisuzaki, W.,

- Kanamitsu, M., Kousky, V., van den Dool, H., Jenne, R., and Fiorino, M.: The NCEP-NCAR 20 50-Year Reanalysis: Monthly Means, CD-ROM and Documentation, Bull. Am. Met. Soc., 82, 247-267, http://wesley.wwb.noaa.gov/reanalysis.html 2001.
 - Labitzke, K. and van Loon, H.: The stratosphere: phenomena, history, and relevance, Springer Verlag, Berlin, 197, 1999. 3421, 3431, 3433
- Labitzke, K. and van Loon, H.: The QBO effect on the solar signal in the global stratosphere in the winter of the Northern Hemisphere, J. Atmos. Solar-Terr. Phys., 62, 621-628, 2000. 3413, 3417, 3424, 3425, 3426, 3432
 - Labitzke, K., Austin, J., Butchart, N., Knight, J., Takahashi, M., Nakamoto, M., Nagashima, T., Haigh, J., and Williams, V.: The global signal of the 11-year solar cycle in the stratosphere:
- observations and models, J. Atmos. Solar-Terr. Phys., 64, 203–210, 2002. 3427 30
 - Lawrence, J. K., Cavidad, A. C., and Ruzmaikin, A.: The response of atmospheric circulation to weak solar forcing J. Geophys. Res., 105, 24839–24848, 2000. 3426

Lee, H. and Smith, A. K.: Simulation of the combined effects of solar cycle, guasi-biennial oscil-

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.

| Title Page | | | |
|------------------------|--------------|--|--|
| Abstract | Introduction | | |
| Conclusions | References | | |
| Tables | Figures | | |
| | | | |
| | ►I | | |
| • | • | | |
| Back | Close | | |
| Full Screen / Esc | | | |
| | | | |
| Print Version | | | |
| Interactive Discussion | | | |

lation, and volcanic forcing on stratospheric ozone changes in recent decades, J. Geophys. Res., 108, ACH 4, doi:10.1029/2001JD001503, 2003. 3413, 3426, 3432

- McCormack, J. P., Hood, L. L., Nagatani, R., Miller, A. J., Planet, W. G., and McPeters, R. D.: Approximate separation of volcanic and 11-year signals in the SBUV-SBUV/2 total ozone record over the 1979–1995 period, Geophys. Res. Lett., 24, 2729–2732, 1997. 3427, 3428,
- 3432 Perlwitz, J. and Graf, H.-F.: The statistical connection between tropospheric and stratospheric circulation of the Northern Hemisphere in winter, J. Clim., 8, 2281–2295, 1995. 3417
- Ramaswamy, V., Chanin, M. L., Angell, J., Barnett, J., Gaffen, D., Gelman, M., Keckhut, P.,
- Koshelkov, Y., Labitzke, K., Lin, J. J. R., O'Neil, A., Nash, J., Randel, W., Rood, R., Shiotani, M., Swinbank, R., and Shine, K.: Stratospheric temperature trends: observations and model simulations, Rev. Geophys., 39, 71–122, 2001. 3413, 3420, 3421

Randel, W. J. and Cobb, J. B.: Coherent variations of monthly mean total ozone and lower stratospheric temperature, J. Geophys. Res., 99, 5433–5447, 1994. 3413

- Reed, R. J.: The role of vertical motions in ozone-weather relationship, J. Meteor., 7, 263–267, 1950. 3421
 - Reid, G. C.: Seasonal and interannual temperature variations in the tropical stratosphere, J. Geophys. Res., 99, 18923–18932, 1994. 3413

Robock, A.: Volcanic eruptions and climate, Rev. Geophys., 38, 191–219, 2000. 3416, 3427, 3428

- Ruzmaikin, A. and Feynman, J.: Solar influence on a major mode of atmospheric variability, J. Geophys. Res., 107, ACL 7, doi:10.1029/2001JD001239, 2002. 3425, 3426
- Sato, M., Hansen, J. E., Lacis, A., and Thomason, L. W.: Stratospheric aerosol optical thickness, http://www.giss.nasa.gov/data/strataer/, 2003. 3416
- Salby, M. L., and Callaghan, P. F.: Interannual changes of the stratospheric circulation: relationship to ozone and tropospheric structure, J. Clim., 24, 3673–3685, 2002. 3413, 3417, 3428

Shiotani, M.: Annual, Quasi-Biennial and El Niño-Southern Oscillation *ENSO* time-scale variations in Equatorial Total Ozone, J. Geophys. Res., 97, 7625–7633, 1992. 3413, 3431, 3432

30

20

5

Solomon, S.: Stratospheric ozone depletion: A review of concepts and history, Rev. Geophys., 37, 275–316, 1999. 3413, 3416, 3417, 3427, 3429

SPARC/IOC/GAW: Assessment of Trends in the Vertical Distribution of Ozone, SPARC Report

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.

| Title Page | | | |
|-------------|----------------------|--|--|
| Abstract | Introduction | | |
| Conclusions | References | | |
| Tables | Figures | | |
| | ÞI | | |
| | | | |
| • | • | | |
| Back | Close | | |
| | 01000 | | |
| Full Scre | een / Esc | | |
| Full Scre | een / Esc | | |
| Full Scre | een / Esc /ersion | | |

No. 1, WMO Global Ozone Research and Monitoring Project-Report No. 44, 1998, http://www.aero.jussieu.fr/~sparc/SPARCReport1/. 3413, 3414, 3418

- Staehelin, J., Harris, N. R. P., Appenzeller, C., and Eberhard, J.: Ozone trends: A review, Rev. Geophys., 39, 231–290, 2001. 3413, 3419, 3428
- Steinbrecht, W., Claude, H., Köhler, U., and Hoinka, K. P.: Correlations between tropopause height and total ozone: Implications for long-term changes, J. Geophys. Res., 103, 19183– 19192, 1998. 3417, 3420, 3421
 - Steinbrecht, W., Claude, H., Köhler, U., and Winkler, P.: Interannual changes of total ozone and Northern Hemisphere circulation patterns, Geophys. Res. Lett., 28, 1191–1194, 2001.
- ¹⁰ **3413, 3414, 3416**
 - Stolarski, R. S. and Hollandsworth Frith, S.: Combined total ozone record from TOMS and SBUV instruments, http://code916.gsfc.nasa.gov/Data_services/merged/mod_data.public. html 2002. 3414

Tourpali, K., Schuurmans, C. J. E., van Dorland, R., Steil, B., and Brühl, C.: Stratospheric and

- tropospheric response to enhanced solar UV-radiation: A model study, Geophys. Res. Lett., 30, doi: 10.1029/2002GL016650, 2003. 3427
 - WMO (World Meteorological Organization): Scientific Assessment of Ozone Depletion: 2002, Global Ozone Research and Monitoring ProjectReport No. 47, 498, Geneva, http://www.wmo.ch/web/arep/ozone.html, 2003. 3413, 3414, 3416, 3420
- Yang, H. and Tung, K. K.: On the phase propagation of extra tropical Quasi-Biennial Oscillation in observational data, J. Geophys. Res., 100, 9091–9100, 1995. 3413, 3423
 - Zerefos, C. S., Bais, A. F., Ziomas, I. C., and Bojkov, R. D.: On the relative importance of Quasi-Biennial Oscillation and El Niño Southern Oscillation in the revised Dobson Total Ozone records, J. Geophys. Res., 97, 10135–10144, 1992. 3413
- Zerefos, C. S., Tourpali, K., Bojkov, B. R., Balis, D. S., Rognerund, B., and Isaksen, I. S. A.: Solar activity-total column ozone relationships: Observations and model studies with heterogeneous chemistry, J. Geophys. Res., 102, 1561–1569, 1997. 3413, 3416, 3426, 3432

ACPD

3, 3411–3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.

| Title Page | | | |
|-------------------|--------------|--|--|
| Abstract | Introduction | | |
| Conclusions | References | | |
| Tables | Figures | | |
| | | | |
| | ▶ | | |
| • | • | | |
| Back | Close | | |
| Full Screen / Esc | | | |
| | | | |
| Print Version | | | |
| | | | |

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.

| Title Page | | | |
|-------------------|--------------|--|--|
| Abstract | Introduction | | |
| Conclusions | References | | |
| Tables | Figures | | |
| | | | |
| I | ►I | | |
| • | • | | |
| Back | Close | | |
| Full Screen / Esc | | | |
| | | | |
| Print Version | | | |
| | | | |

© EGU 2003

Table 1. Two standard deviations of seasonal means for the various explanatory variables X. These values can be used to estimate regression coefficients c_X from the plots showing two standard deviations of $c_X X$. For T(400), the values depend on latitude and longitude, but generally increase with latitude. Approximate ranges are given for the Northern (upper line) and Southern Hemisphere (lower line)

| X | 2σ(DJF) | 2σ (MAM) | $2\sigma(JJA)$ | $2\sigma(SON)$ | unit |
|--------------------|-----------------------|-------------------------|-----------------------|-----------------------------------|--|
| <i>QBO</i> (10/30) | 37 | 37 | 35 | 36 | m/s |
| FS | 1010 | 980 | 990 | 1030 | 10 ⁻²³ J/s/m ² /Hz |
| u(60N) | 10.3 | 6.9 | 0.9 | 3.0 | m/s |
| u(60S) | 4.4 | 3.9 | 5.5 | 9.5 | m/s |
| ENSO | 1.9 | 1.5 | 1.4 | 1.9 | |
| <i>T</i> (400) NH | 0.6 (0°) to 6 (60° N) | 0.6 (0°) to 4 (60° N) | 0.6 (0°) to 3 (60° N) | 0.6 (0°) to 5 (60° N) | Κ |
| <i>T</i> (400) SH | 0.6 (0°) to 4 (80° S) | 0.6 (0°) to 3.5 (70° S) | 0.6 (0°) to 4 (80° S) | 0.6 (0°) to 3 (60 ⁰ S) | K |

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



© EGU 2003



Fig. 1. Global maps of R^2 from regression of 1978 to 2001 TOMS/SBUV total ozone anomalies on a 5° latitude by 10° longitude grid (left panels), or of 1978 to 2001 50 hPa temperature anomalies from NCEP reanalyses (right panels). No TOMS/SBUV observations are available in polar regions in winter.

3, 3411-3449, 2003

Interannual variations of total ozone and

temperature

W. Steinbrecht et al.



Fig. 2. Left panels: Linear trend, in DU per decade, from multiple regression of 1978 to 2001 TOMS/SBUV total ozone anomalies on a 5° latitude by 10° longitude grid. Right panels: Linear trend, in K per decade, from multiple regression of 1978 to 2001 temperature anomalies at 50 hPa from NCEP reanalyses on a 5° latitude by 10° longitude grid. In the white regions the linear trend is not statistically significant at the 90% confidence level. Also note that no TOMS/SBUV observations are available in polar regions in winter.



3, 3411–3449, 2003



Fig. 3. Size of fluctuations attributed to temperature anomalies at 400 hPa. The size is defined as 2 standard deviations, in DU or K, of the $c_{T(400)}T(400)$ time series term in Eq. (1). In this plot and the following, blue colors (negative values) indicate that ozone or temperature are inversely (negatively) correlated to the explanatory variable, whereas yellow and red colors (positive values) denote positive correlation. In the white regions the explanatory variable is not statistically significant at the 90% confidence level. Also note that no TOMS/SBUV observations are available in polar regions in winter.

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



© EGU 2003



Fig. 4. Same as Fig. 3 but for fluctuations attributed to the QBO (2σ of term $c_{QBO(10)} QBO(10) + c_{QBO(30)} QBO(30)$ in Eq. (1)). Here, blue colors indicate inverse (negative) correlation with westerly zonal wind at 30 hPa at the Equator, whereas yellow and red colors indicate positive correlation.

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



© EGU 2003



Fig. 5. Same as Fig. 4 but for regression with a reduced set of explanatory variables, containing only QBO, solar cycle and a linear trend.

3, 3411-3449, 2003



Fig. 6. Same as Fig. 4 but for total ozone fluctuations associated with the 11-year solar cycle. Yellow and red colors indicate positive correlation between total ozone and solar flux. Only data during the easterly phase of the QBO (at 50 hPa) are used. To improve clarity for this figure, as well as for Fig. 7, the confidence level was reduced to 80%, and zonal winds at 60° N and 60° S were not included as explanatory variables.

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



3, 3411-3449, 2003

O3 sol_cyc DJF (QBO West,ST) T50 sol_cyc DJF (QBO West,ST) T50 sol_cyc DJF (QBO West,ST) T50 sol_cyc DJF (QBO West,ST) O3 sol_cyc JJA (QBO West,ST) T50 sol_cyc JJA (QBO West,ST) T5

Fig. 7. Same as Fig. 6 but for the westerly phase of the QBO (at 50 hPa). To improve clarity for this figure, as well as for Fig. 6, the confidence level was reduced to 80%, and zonal winds at 60° N and 60° S were not included as explanatory variables.

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



© EGU 2003



Fig. 8. Same as previous figures, but showing the ozone and 50 hPa temperature response to stratospheric aerosol from the El Chichón and Pinatubo eruption. Here the maximum positive or negative response is shown, not two standard deviations as in the other plots.

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



© EGU 2003



Fig. 9. Same as previous figures, but showing the influence of polar vortex strength, represented by zonal wind anomalies at 60° N, 50 hPa, on total ozone and 50 hPa temperature.

3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



Fig. 10. Same as Fig. 9, but for zonal wind anomalies at 60° S, 50 hPa.



3, 3411-3449, 2003

Interannual variations of total ozone and temperature

W. Steinbrecht et al.



© EGU 2003



Fig. 11. Same as previous Figures, but showing the influence of El Niño/Southern-Oscillation on total ozone and 50 hPa temperature. Top panels: Obtained with full set of explanatory variables, including 400 hPa temperature. Bottom panels: Reduced set of explanatory variables, without 400 hPa temperature.