

**DMS oxidation**

R. von Glasow and P. J.  
Crutzen

# Model study of multiphase DMS oxidation with a focus on halogens

R. von Glasow<sup>1</sup> and P. J. Crutzen<sup>1,2</sup>

<sup>1</sup>Center for Atmospheric Sciences, Scripps Institution of Oceanography, University of California, San Diego, La Jolla, CA, 92093-0221, USA

<sup>2</sup>Atmospheric Chemistry Division, Max-Planck-Institut für Chemie, PO Box 3060, 55020 Mainz, Germany

Received: 26 October 2003 – Accepted: 12 December 2003 – Published: 18 December 2003

Correspondence to: R. von Glasow (roland@fiji.ucsd.edu)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

## Abstract

We studied the oxidation of dimethylsulfide (DMS) in the marine boundary layer (MBL) with a one-dimensional numerical model and focused on the influence of halogens. Our model runs show that there is still significant uncertainty about the end products of the DMS addition pathway, which is especially caused by uncertainty in the product yield of the reaction  $\text{MSIA} + \text{OH}$ . Under cloud-free conditions a MSA yield of only 5% in this reaction could make the addition pathway in the gas phase dominant for MSA formation whereas a yield of 0% would make the gas phase unimportant. Under cloudy conditions the uptake of DMSO and MSIA to droplets results in a contribution of the gas phase of only about 2% to the total formation rate of MSA. The aqueous phase reaction  $\text{MSIA} + \text{OH}$  is the main source for total MSA when gas phase production of MSA is unimportant.  $\text{BrO}$  strongly increases the importance of the addition branch in the oxidation of DMS even when present at mixing ratios smaller than  $0.5 \text{ pmol mol}^{-1}$ . The inclusion of halogen chemistry leads to higher DMS oxidation rates and smaller DMS to  $\text{SO}_2$  conversion efficiencies. The DMS to  $\text{SO}_2$  conversion efficiency is also drastically reduced under cloudy conditions. In clouds especially during winter the aqueous phase reaction  $\text{DMS} + \text{O}_3$  contributes 4–18% to total DMS oxidation. In cloud-free model runs between 5 and 15% of the oxidized DMS reacts further to particulate sulfur, in cloudy runs this fraction is almost 100%. In general, more particulate sulfur is formed when halogen chemistry is included. A possible enrichment of  $\text{HCO}_3^-$  in fresh sea salt aerosol would increase pH values enough to make the reaction of  $\text{S(IV)}^*$  with  $\text{O}_3$  dominant for sulfate production. It leads to a shift from MSA to  $\text{nss-SO}_4^{2-}$  production but increases the total  $\text{nss-SO}_4^{2-}$  only somewhat because almost all available sulfur is already oxidized to particulate sulfur in the base scenario. We discuss how realistic this is for the MBL. We found the reaction  $\text{MSA}_{\text{aq}} + \text{OH}$  to contribute about 10% to the production of  $\text{nss-SO}_4^{2-}$  in clouds. It is unimportant for cloud-free model runs. Sulfate production by  $\text{HOCl}_{\text{aq}}$  and  $\text{HOBr}_{\text{aq}}$  is important in cloud droplets even for small  $\text{Br}^-$  deficits and related small gas phase halogen concentrations. We found differences in

## DMS oxidation

R. von Glasow and P. J. Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

the diurnal variation of the  $\text{Br}^-$  in sea salt aerosol with a peak in the morning when the loss of  $\text{Br}^-$  from the sea salt is small and a peak during day when the loss is almost complete. Overall we find that the presence of halogens lead to processes that decrease the albedo of stratiform clouds in the MBL.

## 1. Introduction

Clouds are essential for our climate, not only for their role in the hydrological cycle but also for their influence on the global albedo, i.e. the amount of sunlight that is reflected back to space. The albedo of the ocean is between 0.06 and 0.1 (depending on wind speed), the type of clouds that we focus on in this study – stratocumulus clouds – has an albedo of about 0.6 (e.g. [Barry and Chorley, 1998](#)), implying a tenfold change from cloud-free to cloudy conditions. Therefore knowledge about the processes that lead to cloud formation and that influence the albedo is essential to understand short term forcing of climate. Clouds grow on cloud condensation nuclei (CCN) and their number, size, and composition determines cloud albedo (e.g. [Twomey, 1974](#)). Over clean oceans CCN are mainly sea salt and sulfate aerosols, with sulfate aerosols usually being dominant in number. Stratiform clouds (Sc, St) in clean regions with low CCN concentrations are most susceptible to changes in size and number of CCN because of small vertical velocities and supersaturations compared to more convective clouds. This type of clouds covers most of the time between 20% and 40% of the oceans at all latitudes (see [Lelieveld et al., 1989](#), for seasonal and latitudinal data).

The most important gaseous precursor for sulfate aerosol over the oceans is dimethyl sulfide (DMS), (see e.g. [Charlson et al., 1987](#)) which is produced by organisms in the oceans and subsequently emitted to the atmosphere where it is oxidized via two main reaction pathways: addition of an O atom and abstraction of an H atom. OH both adds to and abstracts from DMS. Contrary to abstraction, addition has a negative activation energy implying higher rates at lower temperatures.  $\text{NO}_3$  abstracts, but is mainly important in winter under clean marine conditions because of the small  $\text{NO}_x$

### DMS oxidation

R. von Glasow and P. J. Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

mixing ratios. Barnes et al. (1991) found in laboratory experiments that BrO also adds to DMS. Based on this, Toumi (1994) investigated in a model study the importance of BrO as a sink for DMS:



He found that for BrO mixing ratios of  $0.5 \text{ pmol mol}^{-1}$  the effect was already significant compared to the OH reactions. The rate coefficient for this reaction was measured again by Bedjanian et al. (1996), Ingham et al. (1999), and Nakano et al. (2001). The latter two groups extended the measurements to higher pressures and found the reaction rate coefficient to be about 60% faster than measured by Barnes et al. (1991). We use the value of Ingham et al. (1999) with  $k(\text{BrO} + \text{DMS}) = 2.54 \times 10^{-14} \exp(850/T)$ . The ultimate fate of the sulfur atom of the dimethyl sulfoxide (DMSO) that is formed in the initial addition reaction is still not completely known (see Sect. 4).

Under atmospheric conditions sulfuric acid ( $\text{H}_2\text{SO}_4$ ) is the only final product of the DMS oxidation that can form new aerosol particles, all other products may only condense onto (and thereby enlarge) existing particles (see Fig. 1 for an overview of the DMS oxidation reactions in the gas phase that are used in this study). The distinction between  $\text{H}_2\text{SO}_4$  and other final products is important for chemistry-cloud-climate feedback considerations because of changes in the number and size of CCN. Figure 2 shows the aqueous phase reaction mechanism used in this study. According to Ross et al. (1998), the product of the reaction  $\text{DMS}_{\text{aq}} + \text{OH}_{\text{aq}}$  is the adduct. We assumed that addition and abstraction (the latter with the product  $\text{HSO}_4^-$ ) take place with equal yields. We could show, however, that this reaction is negligible.

In this paper we discuss the results from our numerical model calculations of the oxidation of DMS in the MBL. We investigate several scenarios (different chemical initial conditions; summer vs. winter; cloud-free vs. cloudy) and focus on the differences between the two main oxidation branches, the production of particulate sulfur, and point to possible links between the natural cycles of sulfur and halogens. The halogens chlo-

## DMS oxidation

R. von Glasow and P. J. Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

rine and bromine in the marine boundary layer (MBL) originate from sea salt aerosol and can be released to the gas phase where they also play a role in the destruction of ozone (Sander and Crutzen, 1996; Vogt et al., 1996); see also von Glasow and Crutzen (2003, in press) for an overview.

In Sect. 2 we describe possible links between the cycles of sulfur and halogens and the potential climate importance. In Sect. 3 we give a description of the numerical model that we used for this study and discuss our findings in detail in Sect. 4.

## 2. Possible feedbacks between sulfur and halogens

Halogens interact with sulfur in the MBL in two ways: in the gas phase, mainly by the addition reaction  $\text{DMS} + \text{BrO} \longrightarrow \text{DMSO}$  and in the aqueous phase by production of sulfate:  $\text{HOX}_{\text{aq}} + \text{S(IV)} \longrightarrow \text{S(VI)}$  ( $\text{X}=\text{Cl}, \text{Br}$ , Vogt et al., 1996). The yield of  $\text{SO}_2$  in the addition branch of the DMS oxidation is smaller than in the abstraction branch, especially under cloudy conditions (see Sect. 4). Reduced gas phase concentrations of  $\text{SO}_2$  and  $\text{H}_2\text{SO}_4$  lead to a reduction of the precursors for new particle formation because all other intermediate products from the addition branch (dimethylsulfoxide (DMSO), dimethylsulfone ( $\text{DMSO}_2$ ), methyl sulfinic acid (MSIA,  $\text{CH}_3\text{S(O)OH}$ ), methyl sulfonic acid (MSA,  $\text{CH}_3\text{S(O)(O)OH}$ )) do not produce new particles, they can only condense on preexisting ones thereby increasing the size and, for example for dust, organic, or soot aerosol, the hygroscopicity of the CCN.

The additional S(IV) oxidation reactions by  $\text{HOX}_{\text{aq}}$  in the aqueous phase (aerosols and droplets) increase the uptake of  $\text{SO}_2$  from the gas phase, consequently reducing the production of gaseous  $\text{H}_2\text{SO}_4$  and subsequent new particle formation. On the other hand it leads to the growth of already existing particles. Therefore the net effect of the halogen - sulfur interactions are i) decreased precursors for new particle formation and ii) increased size and hygroscopicity of the CCN. This constitutes a natural indirect aerosol effect which would decrease the cloud albedo as opposed to the anthropogenic indirect aerosol effect which increases cloud albedo. The feedbacks are schematically

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGU 2003

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

shown in Fig. 3. Under cloud-free conditions this idea is to some extent dependent on the rate coefficient and products of the reaction MSIA + OH as explained in Sect. 4.1.

As the activation of sea salt bromide is acid catalyzed, one could hypothesize that the importance of the proposed climate feedback process has increased - at least in the regions that are influenced by anthropogenic pollution - since human activity raised the gas phase acidity of the atmosphere through SO<sub>2</sub> and NO<sub>x</sub> emissions leading to H<sub>2</sub>SO<sub>4</sub> and HNO<sub>3</sub>; it might also be more important in regions with a lot of ship traffic.

So far, no global information about BrO concentrations in the MBL is available. Significant depletion of bromide from sea salt aerosol, however, was found in many locations (see Sander et al., 2003, for an overview) which is an indicator for active halogen chemistry in the MBL.

We want to stress that the implications of lower SO<sub>2</sub> (and therefore H<sub>2</sub>SO<sub>4</sub>) yield from the addition path are valid for all addition reactions.

### 3. Model description and setup

We used the one-dimensional model of the marine boundary layer MISTRA-MPIC as described in von Glasow et al. (2002b,a). In addition to dynamics and thermodynamics, it includes a detailed microphysical module that calculates particle growth explicitly and accounts for interactions between radiation and particles. Chemical reactions in the gas phase are considered in all model layers. In cloud-free layers with relative humidity greater than the crystallization humidity, aerosol chemistry is considered in “sulfate” aerosol (dry radius less than 0.5 μm, initial composition 32% (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 64% NH<sub>4</sub>HSO<sub>4</sub>, 4% NH<sub>4</sub>NO<sub>3</sub>) and “sea salt” aerosol (dry radius greater than 0.5 μm, initial composition like sea water) bins. In cloud layers two additional chemical aqueous phase bins are considered: droplets that grew on sulfate and on sea salt aerosols.

The model includes 138 gas phase (H-O-S-C-N-Br-Cl) reactions that are important for the chemistry of the MBL including halogen chemistry. A comprehensive reaction set with 143 aqueous phase reactions, 12 heterogeneous reactions and 21 equilibria

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

is used for each of the four particulate bins. We updated the DMS oxidation mechanism and treat it in more detail now (see reaction table in electronic attachment to this paper or <http://www.rolandvonglasow.de>). We assumed that  $\text{H}_2\text{SO}_4$ , MSA, MSIA, and  $\text{DMSO}_2$  have infinite effective Henry's law constant which we implemented by using  $H^* = 10^{16} \text{ M atm}^{-1}$ . If we assume smaller  $H^*$ , MSA degasses from the sulfate aerosol during night. The formation of  $\text{DMSO}_2$  in the reaction  $\text{DMSO} + \text{OH}$  is uncertain and reaction of  $\text{DMSO}_2$  with  $\text{OH}$ ,  $\text{NO}_3$ ,  $\text{O}_3$ , and  $\text{Cl}$  were found to be not of importance (Falbe-Hansen et al., 2000). Our assumption of infinite solubility might overestimates its aqueous fraction which is no major shortcoming in our mechanism, because we assume it to be stable both in the gas and in the aqueous phase, so it only affects the gas-aqueous distribution of an unreactive species.

The accommodation coefficients for  $\text{H}_2\text{SO}_4$ , MSA, MSIA, and  $\text{DMSO}_2$  are taken from De Bruyn et al. (1994). The initial gas phase concentrations for the different runs are listed in Table 1. In the remainder of this text we will use the generic term “particles” to refer to both aerosols and cloud droplets.

We will discuss detailed process studies of the multiphase chemistry of the MBL under idealized conditions to investigate the reaction mechanism and then show results for “mean” conditions for the tropics and Cape Grim, Tasmania ( $41^\circ\text{S}$ ).

The latitude chosen for the process studies is  $\varphi = 30^\circ \text{N}$  at the end of July for the summer runs and at the end of January for the winter runs. In the cloud-free runs, the boundary layer height is roughly 700 m, moisture and heat fluxes from the sea surface are adjusted to yield a stable boundary layer. The relative humidity at the sea surface is roughly 65%, increasing with height to around 90% below the inversion that caps the MBL. After a spin-up of the dynamical part of the model for 2 days the complete model was integrated for 5 days (the model runs start at midnight). The surface temperature is about  $14^\circ\text{C}$  and  $3^\circ\text{C}$  in the summer and winter runs, respectively. According to Peixoto and Oort (1991) (their figure 7.4) the average seasonal difference in surface air temperature over the oceans is less than  $10^\circ\text{C}$  except for very high latitudes and usually less than  $5^\circ\text{C}$ . Differences between the winter and summer runs are only due

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

to different meteorology, the chemical initial and boundary condition have not been changed to be able to concentrate on the influence of temperature and insolation. A change in these two parameters, however, does reduce the OH concentration by 70% which has major implications for the photochemistry.

5 For the runs with a stratiform cloud we chose the value of the large scale subsidence such that the cloud top remains stable at 775 m. The cloud base varies between 500 m during the day and 300 m during the night. The maximum liquid water content (LWC) is roughly  $0.75 \text{ g/m}^3$  before sunrise and  $0.45 \text{ g/m}^3$  in the afternoon. In the winter runs the cloud base is fairly constant around 300 m and the maximum LWC is roughly  $0.5 \text{ g/m}^3$ .

For both the cloud-free and cloudy process studies we tested different reaction rates and product yields for the reaction  $\text{MSIA} + \text{OH}$  that is critical for the yields of  $\text{SO}_2$ ,  $\text{MSA}$  and  $\text{H}_2\text{SO}_4$  in the addition pathway (see Sect. 4.1).

15 The MBL height for the “tropics” runs is about 850 m with a surface temperature of  $28^\circ\text{C}$ . The chemical initial conditions are taken from Wagner et al. (2002). We used three scenarios for “Cape Grim”, cloudy winter, cloudy summer, and cloud-free summer trying to get close to observed meteorological conditions but are aware of the fact that this is still only a rough approximation and certainly cannot account for the variability encountered. Like all other scenarios they are mainly meant as sensitivity studies to point to some specific findings. The cloud top in the winter run is at 710 m and at about 900 m in the summer run, the cloud base remains roughly around 350 m in the winter run, whereas it is highest during day with about 650 m and lowest during night (around 550 m) in the summer run. The maximum LWC for both seasons is about  $0.4 \text{ g/m}^3$ . The surface temperature is about  $8^\circ\text{C}$  and  $14^\circ\text{C}$  for the winter and summer runs, respectively. In the cloud-free run the inversion of the MBL is around 700 m, the relative humidity near the sea surface is about 65% it increases to 95% at the top of the MBL. Chemical initial conditions for these three runs are from Ayers et al. (1995, 1997a,b, 1999) and Monks et al. (1998, 2000).

In the cloud-free runs we initialized  $\text{SO}_2$  in the free troposphere with 50% of the MBL



## DMS oxidation

R. von Glasow and P. J. Crutzen

value, whereas we chose 25% for the cloudy runs. This reduces the downflux of  $\text{SO}_2$  in cloudy runs which is both a source of sulfur and of gas phase acidity. Downward transport across the inversion that caps the MBL is small in cloud-free runs.

We included a  $\text{NO}_2$  source of  $10 \text{ pmol mol}^{-1} \text{ day}^{-1}$  ( $2 \text{ pmol mol}^{-1} \text{ day}^{-1}$  for the Cape Grim runs) in order to account for uptake losses and maintain  $\text{NO}_x$  mixing ratios of about 3 and  $15 \text{ pmol mol}^{-1}$  for runs with and without halogen chemistry, respectively. Sources for  $\text{NO}_x$  in the clean MBL can, for example, be decomposition of ocean derived organic nitrates (e.g. Atlas et al., 1993; Chuck et al., 2002) or direct release from the ocean (Zafiriou et al., 1980).

We did all runs with and without inclusion of halogen chemistry in order to demonstrate the effect of halogens on the chemistry. Under the initial conditions that we chose for the process study runs almost all bromide gets released from the sea salt aerosol resulting in  $\text{BrO}$  mixing ratios around  $1.5 \text{ pmol mol}^{-1}$  during noon and  $3\text{--}4 \text{ pmol mol}^{-1}$  during the morning and afternoon peaks (see von Glasow et al., 2002b, for a discussion of the diurnal cycle of  $\text{BrO}$ ). In the cloudy process study runs the  $\text{BrO}$  mixing ratio is about  $1\text{--}1.5 \text{ pmol mol}^{-1}$ . It is less for the Cape Grim and tropics scenarios and will be discussed later.

#### 4. Results and discussion

The results of the model runs that are discussed in this section are summarized in Tables 2, 2, and 3 which list the DMS oxidation rates, non-sea salt sulfate ( $\text{nss-SO}_4^{2-}$ ) and MSA production rates,  $\text{DMS} \rightarrow \text{SO}_2$  conversion efficiencies, and MSA to  $\text{nss-SO}_4^{2-}$  ratios (as explained below). Figures 4 and 5 show the temporal evolution of  $\text{SO}_2$ , DMS, DMSO, MSA, and  $\text{DMSO/DMS}$ . If not explicitly stated otherwise all numbers that we mention in this section and in the tables refer to mean values over the depth of the MBL and the second to fifth model day for cloud-free runs and second and third day for cloudy runs.

This section is subdivided into four parts. First we will discuss the likely final products

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

of the addition pathway based on laboratory, field and modeling studies. Then we will present results from our sensitivity studies for the importance of different pathways in the oxidation of DMS, focussing on halogens. This is followed by a discussion of the particulate final products and related particle growth and a discussion of the conditions encountered in the remote clean marine boundary layer based on Cape Grim, Tasmania and Baring Head, New Zealand conditions. Finally we will discuss differences in the diurnal variation of particulate and gaseous bromine that could help explain differences that were so far encountered when comparing model and field results.

## 4.1. Final products in the addition pathway

As already mentioned, knowledge about the pathways and product yields of the DMS oxidation is essential to estimate the potential for new particle formation and for growth of existing aerosol particles. Usually it is assumed that MSA is only formed in the abstraction pathway as a product of the reaction  $\text{CH}_3\text{SO}_3 + \text{HO}_2$ . However, MSA might also be formed in the addition pathway. The initial addition reactions produce DMSO which reacts with OH (almost 100% yield, Urbanski et al., 1998; Arsene et al., 2002; Wang and Zhang, 2002, to MSIA). Lucas and Prinn (2002) suggested that MSIA might react with  $\text{O}_3$  to MSA. MSA could also be a product of the reaction  $\text{MSIA} + \text{OH}$  but the main product of the reaction  $\text{MSIA} + \text{OH}$  is  $\text{CH}_3\text{SO}_2$  (Kukui et al., 2003), a main intermediate in the abstraction pathway. The reaction  $\text{MSIA} + \text{OH}$  could therefore be regarded as a “crossover point” between the addition and abstraction pathway (see also Fig. 1). As will be shown below, the yield of MSA in this reaction and the rate coefficient are of major importance for the determination of the final products of the addition pathway under cloud-free conditions. If most DMSO that is formed in this reaction is transformed via MSIA to  $\text{CH}_3\text{SO}_2$  then there would be no significant differences between the end products of the addition and abstraction pathways in the gas phase. It is important to note that the intermediate products DMSO and MSIA are taken up by particles, especially in the cloudy MBL.

To investigate the possible end products of DMS oxidation and the implications for

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

the MBL, we performed sensitivity studies with different rate constants for the reaction MSIA + OH. Furthermore we investigate different yields of MSA in this reaction and differences between cloud-free and cloudy model runs.

According to the qualitative results of the smog chamber study by [Arsene et al. \(2002\)](#) the overall yield of SO<sub>2</sub> in the addition branch should be small, whereas [Kukui et al. \(2003\)](#) found that the yield of CH<sub>3</sub>SO<sub>2</sub> in the reaction MSIA + OH is 0.9 ± 0.2 with indications for a yield of unity. We used their measurement and previous estimates of the rate constant for the reaction MSIA + OH for sensitivity runs:  $k=1.0 \times 10^{-12} \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$  ([Lucas and Prinn \(2002\)](#)), runs labeled “Luc”),  $k=1.6 \times 10^{-11} \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$  ([Yin et al. \(1990\)](#)), runs labeled “Yin”),  $k=9.0 \times 10^{-11} \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$  ([Kukui et al. \(2003\)](#)), runs labeled “Kuk”). In the cases “Luc” and “Yin” we assumed a 100% yield of CH<sub>3</sub>SO<sub>2</sub> from this reaction whereas we tested yields of 0, 5 and 10% MSA for the “Kuk” cases. The “Luc” scenarios also include the reaction MSIA + O<sub>3</sub>.

For the “Cape Grim” and “tropics” scenarios we used a yield of 5% MSA from the reaction MSIA + OH which is well within the experimental uncertainty and increases gas phase MSA mixing ratios.

As already mentioned, we did a model run for each scenario with and without halogen chemistry. In the runs including halogen chemistry the reaction DMS + BrO makes the addition pathway be a lot more important than in runs without halogen chemistry (see Sect. 4.2).

#### 4.1.1. Model runs for the cloud-free MBL

In the “Luc” scenario the gas phase concentration of SO<sub>2</sub> is reduced by about 30% when halogen chemistry is considered (see Fig. 4). The production of SO<sub>2</sub> is smaller than in the other runs because the “crossing” from the addition to the abstraction pathway via the reaction MSIA + OH is unimportant. In the other runs, especially in the “Kuk” runs, this “crossing” is very efficient and the products SO<sub>2</sub>, H<sub>2</sub>SO<sub>4</sub> and MSA are formed as a consequence in the gas phase. In these runs the SO<sub>2</sub> mixing ratio is higher if halogen chemistry is included because of the production of MSIA following

## DMS oxidation

R. von Glasow and P. J. Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

the reaction  $\text{DMS} + \text{BrO}$ . Only on the last model day, the  $\text{SO}_2$  mixing ratios in the non-halogen runs exceed those of the halogen runs because OH is needed as reactant which is smaller by roughly 20–25% on the last model day in the halogen runs due to halogen-catalyzed  $\text{O}_3$  destruction (see Fig. 5).

The MSA production rate in “Luc” is about five times as high as in the “Yin” runs and ten times greater than in the “Kuk 0%” runs which is mainly caused by the reaction  $\text{MSIA} + \text{O}_3$ . In the “Luc” runs between 61% and 77% of  $\text{MSA}_{\text{tot}}$  is formed via the reaction  $\text{MSIA} + \text{O}_3$ . Formation of MSA in the reaction  $\text{CH}_3\text{SO}_3 + \text{HO}_2$  is less than 2.5% in the summer and about 10% in the winter runs of the total MSA production for all scenarios. The rest of the  $\text{MSA}_{\text{tot}}$  production occurs in the aerosols.

We tested how sensitive the results are to different yields of MSA in the reaction  $\text{MSIA} + \text{OH}$  (only for “Kuk” scenarios). Even a small yield of 5% leads to a more than sevenfold increase in the formation of MSA because of the high rate constant for  $\text{MSIA} + \text{OH}$  making the gas phase dominant for the formation of total MSA. Using a 10% yield the MSA production is fourteen times higher.

Reactions of the MSIA-precursor DMSO with compounds other than OH or strong uptake by particles could make the system less sensitive to the reaction  $\text{MSIA} + \text{OH}$ . We investigated the potential of a surface reaction of OH with  $\text{DMSO}_{\text{surf}}$  because Allen et al. (1999) found that DMSO partitions to the surface in DMSO - water mixtures. We found, however, that almost every OH molecule that hits the aerosols would have to react with DMSO to lead to a detectable increase in MSIA- formation, which is impossible. An upper limit for the aerosol surface fraction being covered with DMSO (based on our size distribution and  $\text{DMSO}_{\text{aq}}$  concentrations) is  $10^{-4}$ . We also tested if aqueous phase reactions of DMSO other than with OH could lead to an increase in aerosol phase MSA production but both  $\text{Cl}_2^-$  and  $\text{NO}_{3,\text{aq}}$  (which have the highest potential to be of importance) could not compete with the  $\text{OH}_{\text{aq}}$  reaction. We further tested if increasing the measured Henry’s law constant of DMSO (Watts and Brimblecombe, 1987) would lead to an increased production of  $\text{MSIA}^-$  via reaction with  $\text{OH}_{\text{aq}}$  but this was only the case when we increased it by 3 orders of magnitude, which is unrealistic.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

In the “Luc” runs that include MSIA + O<sub>3</sub>, this reaction is dominant for the production of total MSA (including aerosol). In all cloud-free runs that we discuss in this paper the gas phase production of MSA via CH<sub>3</sub>SO<sub>3</sub> + HO<sub>2</sub> contributes less than 10% of the total MSA production. For the “Kuk” runs with 5% yield of MSA the reaction MSIA + OH accounts for more than 70% of the total MSA production. If these reactions occur as assumed then MSA formation in the DMS abstraction pathway would be only of minor importance unlike usually assumed in the literature so far. The aqueous phase reaction MSIA + OH is the main source for total MSA when gas phase production of MSA is unimportant.

The peak mixing ratios of gas phase MSA occur before noon if halogen chemistry is included and at noon for non-halogen model runs (see Sect. 4.2 for an explanation for this). Peak values are less than 1 × 10<sup>5</sup> molec cm<sup>-3</sup> for the runs with 0% MSA yield, less than 1.0 × 10<sup>7</sup> molec cm<sup>-3</sup> for the runs with 5% MSA yield in the reaction MSIA + OH and about 2 × 10<sup>7</sup> molec cm<sup>-3</sup> for the runs with 10% MSA yield. In our “tropics” scenario with a 5% yield of MSA the peak concentrations of MSA are about 2.5 × 10<sup>6</sup> molec cm<sup>-3</sup> and 1.5 × 10<sup>6</sup> molec cm<sup>-3</sup>, with and without halogen chemistry, respectively.

So far only few measurements of gas phase MSA have been made. Published concentrations range from 1 - 9 × 10<sup>5</sup> molec cm<sup>-3</sup> in the Antarctic (Mauldin et al., 2001; Jefferson et al., 1998), to 10<sup>5</sup> - 10<sup>6</sup> molec cm<sup>-3</sup> for the tropics (Davis et al., 1999), and < 3 × 10<sup>4</sup> - 3.7 × 10<sup>7</sup> molec cm<sup>-3</sup> for the Eastern Mediterranean (Bardouki et al., 2003). These measurements indicate that there has to be at least some formation of gas phase MSA in the atmosphere which we could not satisfactorily reproduce in the model without assuming that MSA is produced in the addition pathway.

Davis et al. (1999) estimated that during flight 7 of the PEM Tropics-A campaign the gas phase contributed 1% to total MSA production. They assumed a MSA yield of 3% in the reaction MSIA + OH and 0.35% in the abstraction reaction of OH with DMS. Lucas and Prinn (2002) concluded that reactions in addition to those proposed by Yin et al. (1990) are needed to reproduce observed MSA(g) during ACE 1. Assuming only

the abstraction reaction DMS + OH as precursor for MSA(g), [Bardouki et al. \(2003\)](#) estimated a contribution of the gas phase to total MSA production of 0.1% for the Eastern Mediterranean Sea.

#### 4.1.2. Model runs for the cloudy MBL

5 Under cloudy conditions the picture changes drastically. The uptake of the MSIA-precursor DMSO into droplets is fast. This leads to a strongly decreased reaction rate of MSIA + OH  $\rightarrow$  CH<sub>3</sub>SO<sub>2</sub> and no significant contribution of the addition pathway to the formation of SO<sub>2</sub>. Therefore the concentration of SO<sub>2</sub> is smaller in all runs including halogen chemistry than in the runs without because of the predominance of addition  
10 reactions in the halogen runs. In the “Kuk” scenarios the mean SO<sub>2</sub> concentrations are about 12% smaller in summer in halogen runs than in runs without halogens and 64% in winter. One has to keep in mind that due to strong uptake by the cloud droplets the gas phase concentrations of SO<sub>2</sub> in the cloud runs is more than an order of magnitude smaller than in cloud-free runs. The importance of gas phase reactions for the forma-  
15 tion of MSA in the cloudy runs is minimal, they never contribute more than 2% to the total MSA formation.

Due to the fast uptake by droplets of the intermediate products in the addition pathway the production of MSA is strongly increased in the runs with halogen chemistry because of the dominance of addition as loss for DMS in these cases.

#### 20 4.1.3. Conclusions

Our model runs show that there is still significant uncertainty about the end products of the DMS addition pathway. Especially the sensitivity in the product yield of the reaction MSIA + OH makes it hard to come to final conclusions. If no MSA is formed in this reaction and if the reaction MSIA + O<sub>3</sub> is slower than assumed by [Lucas and Prinn](#)  
25 [\(2002\)](#), than the gas phase production of MSA under cloud-free conditions is about 2% in summer and 10% in winter of the total MSA formation. Or, phrased differently,

### DMS oxidation

R. von Glasow and P. J. Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**DMS oxidation**R. von Glasow and P. J.  
Crutzen[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGU 2003

gas-phase production of MSA would be of minor importance compared to production in aerosols. If, however, the reaction  $\text{MSIA} + \text{OH}$  had a MSA yield of only 5% it could become the dominant MSA production pathway (for both halogen and no halogen conditions). Then the DMS-abstraction pathway would be only of minor importance for MSA production.

Under cloudy conditions the gas phase contributes only about 2% to the total formation rate of MSA and is insignificant for winter runs. The sensitivities to the gas phase reactions of MSIA are very weak due to the uptake of the MSIA precursor DMSO and the uptake of MSIA itself.

The aqueous phase reaction  $\text{MSIA} + \text{OH}$  is the main source for total MSA when gas phase production of MSA is unimportant. Field measurements indicate that there has to be at least some formation of gas phase MSA in the atmosphere which we could not satisfactorily reproduce in the model without assuming that MSA is produced in the addition pathway.

## 4.2. Oxidation of DMS

### 4.2.1. Model runs for cloud-free MBL

The DMS oxidation rates are the same for the previously discussed scenarios “Luc”, “Yin”, and “Kuk” because differences between these runs only affect the products of DMS. Inclusion of halogens increases the DMS destruction by about 25% in summer and by 100% in winter. The contribution of BrO to the destruction of DMS is about 64% in the summer runs and about 84% for the winter runs. It increases with time even though BrO concentrations stay constant because the concentration of the next most important oxidant OH decreases with time due to decreasing  $\text{O}_3$  which gets photochemically destroyed, especially by halogens. The contribution of BrO to total DMS destruction on the second and third day is about 56% which is still a very high number. The addition pathway dominates in halogen runs with about 76% compared to 44% in the runs without halogens (89% vs. 58% in winter).

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

In the runs without halogen chemistry about 5-6% and 11.5% in summer and winter, respectively, of the total DMS destruction is by  $\text{NO}_3$ , the rest by OH. This number is strongly dependent on  $\text{NO}_x$  mixing ratios which were about 2–3 and 13  $\text{pmol mol}^{-1}$  for runs with and without halogen chemistry, respectively. For the winter runs the mixing ratios are about 3 and 20  $\text{pmol mol}^{-1}$  for runs with and without halogen chemistry, respectively.  $\text{NO}_3$  is relatively more important under winter than summer conditions because of longer nights and roughly 70% smaller concentrations of OH in winter. This is caused by an adjustment of the solar elevation and temperature to winter conditions; we did not change the chemical initial conditions. In summer  $\text{NO}_3$  contributes about 1% to the total DMS oxidation rate in the halogen runs and OH and Cl about 27% and 8%, respectively (see also the discussion of seasonal differences in von Glasow et al., 2002b). The source for  $\text{NO}_x$  in the model is a prescribed flux of 10  $\text{pmol mol}^{-1} \text{ day}^{-1}$  (see Sect. 3). Furthermore, the photolysis of  $\text{NO}_3^-$  in sea salt aerosol leads to an additional flux of  $\text{NO}_2$  to the gas phase of about 0.7  $\text{pmol mol}^{-1}$  per day.

We do not expect the contribution of BrO to be this high over the remote oceans but discuss these cases to illustrate the chemistry. In the scenarios “tropics” and “Cape Grim” (see Sect. 4.2.3) we will discuss smaller influence of halogen chemistry. To our knowledge there have been only two measurements of BrO reported in the MBL: Leser et al. (2003) report about 0.7  $\text{pmol mol}^{-1}$  from the North Atlantic whereas Saiz-Lopez et al. (2003) measured up to 6  $\text{pmol mol}^{-1}$  at Mace Head, Ireland. The presence of BrO has also been inferred from differential optical absorption spectrometry (DOAS) measurements in open-ocean surface air at Hawaii (Pszenny et al., 2003). Due to the vertical profile in sea salt aerosol acidity with a maximum near the inversion that caps the MBL (von Glasow and Sander, 2001), BrO mixing ratios show a vertical profile in the model with maxima at the top of the MBL (see also von Glasow et al., 2002b). Around noon, they are about 1.5  $\text{pmol mol}^{-1}$  at the surface and 2.5  $\text{pmol mol}^{-1}$  near the inversion that caps the MBL. The morning peaks are between 4.5 and 6.2  $\text{pmol mol}^{-1}$ . BrO is never smaller than 1.5  $\text{pmol mol}^{-1}$  during day.



## DMS oxidation

R. von Glasow and P. J.  
Crutzen

## 4.2.2. Model runs for cloudy MBL

In cloud runs BrO is relatively less important as oxidant for DMS than in cloud-free runs but it is still a major oxidant if halogen chemistry is considered in the model. The addition pathway dominates in halogen runs with about 62% compared to 40% in the runs without halogens (80% vs. 40% in winter). The importance of OH as oxidant increased in the halogen runs to about 55% and that of Cl decreased to 3%.

Based on the measurements of Gershenson et al. (2001), Boucher et al. (2003) included the aqueous phase oxidation of DMS by O<sub>3</sub> in a global model of the sulfur cycle and found it to contribute about 6% to the overall destruction of DMS. In our cloud runs including halogen chemistry it contributes about 4% and in runs without halogen chemistry 7% of the total DMS oxidation in the MBL. Under winter conditions, however, the contribution is 6% and 18.6%, with and without halogen chemistry, respectively. It is unimportant under cloud-free conditions. These numbers suggest that this reaction should be included in models of the sulfur cycle, especially when looking at high latitudes or winter conditions.

## 4.2.3. Model runs for special environments

The BrO mixing ratios for the “tropics” scenario are small with 0.3 pmol mol<sup>-1</sup> around noon and peak values of about 1.5 pmol mol<sup>-1</sup>. Nevertheless, the relative contribution of BrO to DMS oxidation is very significant with almost 17%.

In the “Cape Grim” scenarios with clouds, the BrO contribution is very strong for summer with 38% but only about 14% during winter. We did additional runs for the “Cape Grim summer” scenario where we enhanced the sea salt alkalinity by increasing the bicarbonate (HCO<sub>3</sub><sup>-</sup>) concentrations in fresh sea salt aerosol. These studies are based on field observations (Sievering et al. (1999), Sievering et al., 2003, submitted to J. Geophys. Res.) that enrichment of HCO<sub>3</sub><sup>-</sup> could occur at the ocean’s surface where sea salt aerosol is produced from bubble bursting. They refer to this as “biological alkalinity”. In sea salt aerosol HCO<sub>3</sub><sup>-</sup> acts as a buffer via the equilibrium

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGU 2003

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

$\text{CO}_2 + \text{H}_2\text{O} \rightleftharpoons \text{HCO}_3^- + \text{H}^+$ . It titrates most of the acidity in sea salt until the  $\text{HCO}_3^-$  reservoir is used up and only then the particle pH drops.  $\text{HCO}_3^-$  is therefore of major importance for the regulation of the pH of sea salt aerosol. We tested a 100% (“2x  $\text{HCO}_3^-$ ” run), a 150% (“2.5x  $\text{HCO}_3^-$ ” run), and a 350% (“4.5x  $\text{HCO}_3^-$ ” run) increase in  $\text{HCO}_3^-$ . In these cases BrO mixing ratios were even smaller than in the “tropics” cases: 0.1 - 0.2 pmol mol<sup>-1</sup> in the “2x  $\text{HCO}_3^-$ ” run and less than 0.1 pmol mol<sup>-1</sup> in the “2.5x  $\text{HCO}_3^-$ ” and “4.5x  $\text{HCO}_3^-$ ” runs. For comparison: in the “Cape Grim, summer” run the BrO mixing ratios are 0.5 - 1.5 pmol mol<sup>-1</sup>. The relative contribution of BrO to DMS is still significant although drastically reduced in the runs with increased alkalinity: about 38% in the “Cape Grim summer” run, about 4.6% in the “2x  $\text{HCO}_3^-$ ” run, 2.5% in the “2.5x  $\text{HCO}_3^-$ ” run, and 1.3% in the “4.5x  $\text{HCO}_3^-$ ”. The bromine deficit (calculated from the  $\text{Br}^- : \text{Na}^+$  ratio in the sea salt aerosol as compared to that in seawater) is with 13 - 33% in the “Cape Grim summer” run smaller than the observed summer values of about 60 - 80% (Ayers et al., 1999). The deficit is only 3.5% in the “2x  $\text{HCO}_3^-$ ” run and less than 1% in the “2.5x  $\text{HCO}_3^-$ ” and “4.5x  $\text{HCO}_3^-$ ” scenarios. This, however, and the very high seasalt pH values of 8 - 10 are unrealistic if compared with measurements that show pH values between 3.5 and 5.1 for clean to moderately polluted conditions (Keene and Savoie, 1998, 1999; Keene et al., 2002; Pszenny et al., 2003). Caused by this strong reduction of the activation of  $\text{Br}^-$ , the development of gas phase chemistry in the scenarios with  $\text{HCO}_3^-$  enrichment had a greater similarity with the non-halogen run than with the halogen run.

In the base cloud-free model run for “Cape Grim” conditions the BrO mixing ratio is with 5 - 15 pmol mol<sup>-1</sup> unrealistically high, the bromine deficit of 30-85%, however, is similar to the measured values of Ayers et al. (1999). One way to explain this discrepancy would involve organic bromine compounds that “sequester” bromine (Toyota et al., 2003). In this study, however, we did not investigate this possibility. We also did runs with a 2.5x and 4.5x enrichment of  $\text{HCO}_3^-$  and found the expected increase in sea salt aerosol pH and smaller BrO mixing ratios. On the other hand, the  $\text{Br}^-$  deficits of

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

20 and 3% were too small compared with measurements. The contribution of BrO to the DMS oxidation decreased from 78% in the run without increased sea salt alkalinity to 16% and 2.5% with a 2.5 and 4.5 fold enrichment in  $\text{HCO}_3^-$ , respectively.

In the “Cape Grim, winter” scenarios the oxidation of DMS is reduced due to the reduction by more than a factor of twenty in the DMS emission and to smaller oxidant concentrations. BrO mixing ratios are less than  $0.5 \text{ pmol mol}^{-1}$  but it still plays an important role for the oxidation of DMS with 16% of the total. The most important oxidant is the  $\text{NO}_3$  radical with a contribution of 40% including halogen chemistry and 77% without it.  $\text{NO}_x$  mixing ratios are 6 - 12  $\text{pmol mol}^{-1}$  and the nighttime  $\text{NO}_3$  mixing ratios are up to  $0.5 \text{ pmol mol}^{-1}$ . The main reason for the importance of  $\text{NO}_3$  are the small OH and BrO concentrations.

#### 4.2.4. DMS $\rightarrow$ $\text{SO}_2$ conversion efficiency

The DMS  $\rightarrow$   $\text{SO}_2$  conversion efficiency is a measure for the importance of gas phase reactions and for the production of new particle formation precursor gases. Many estimates of the DMS conversion efficiency have been published and cover a wide range roughly from 0.2 to 1. Usually measurements of DMS,  $\text{SO}_2$  and sometimes OH are used together with photochemical models to derive the conversion efficiencies. As discussed in detail by [de Bruyn et al. \(2002\)](#), these calculations are very dependent on the reaction mechanism that was chosen and can even yield different results for the same field campaign. It is also not yet established if there is a latitude dependence of the conversion efficiency but measurements indicate that it is higher at low latitudes (see [de Bruyn et al., 2002](#), and references therein).

The DMS  $\rightarrow$   $\text{SO}_2$  conversion efficiencies were calculated for all runs (see Tables 2 and 2), they range from 0.140 to 0.957 and showed an expected rise with increase in the rate constant of  $\text{MSIA} + \text{OH}$ . The conversion efficiency is reduced by only 2-3% if we use a yield of 5% for MSA in the “Kuk” scenario and by 4-7% for the “Kuk 10%” run. It is strongly reduced under colder temperatures and with lower OH (see “winter”

cases).

The SO<sub>2</sub> yield from the oxidation of DMS is always smaller when halogen chemistry is included. This is caused by the higher DMS destruction rates in halogen runs and slower conversion of CH<sub>3</sub>S to CH<sub>3</sub>SO<sub>2</sub> that involves two reactions with O<sub>3</sub> which gets depleted by halogen chemistry.

When halogen chemistry is accounted for, the destruction of DMS starts earlier in the morning and has a more abrupt onset (see Figs. 4 and 5) because of the early morning peak in BrO. This propagates into DMSO and MSA(g) (if the MSA yield in the reaction MSIA + OH is greater than zero) which also shows a peak before noon. These diurnal variation might be used for a qualitative test during field campaigns to see if halogen chemistry was important. The diurnal variation of BrO will only show the morning and afternoon peaks as long as the NO<sub>x</sub> concentrations are not too high to make NO<sub>2</sub> be the main sink (instead of HO<sub>2</sub>) for BrO (see also discussion in von Glasow et al., 2002b).

In cloudy runs, the difference in the DMS → SO<sub>2</sub> conversion efficiency between runs with and without halogen chemistry is even higher than under cloud-free conditions and especially low under winter conditions because of the uptake of DMS and intermediate products like DMSO and MSIA by the cloud droplets.

#### 4.2.5. Conclusions

We confirmed the influence of BrO in the oxidation of DMS. It strongly increases the importance of the addition branch in the oxidation of DMS even when present at mixing ratios smaller than 0.5 pmol mol<sup>-1</sup>. The general trend is that the inclusion of halogen chemistry leads to higher DMS oxidation rates and smaller DMS → SO<sub>2</sub> conversion efficiencies. The DMS → SO<sub>2</sub> conversion efficiency is also drastically reduced under cloudy conditions due to uptake of the intermediate products in the addition pathway. The presence of halogens changes the diurnal cycle sulfur species: the destruction of DMS starts earlier and the mixing ratios of the oxidation products (DMSO, MSIA, MSA, SO<sub>2</sub>, and H<sub>2</sub>SO<sub>4</sub>) peak earlier when halogen chemistry is included. We also

### DMS oxidation

R. von Glasow and P. J. Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

confirmed the importance of the reaction  $\text{DMS}_{\text{aq}} + \text{O}_3$  for the oxidation of DMS under cloudy conditions, especially in winter.

#### 4.3. Sulfate vs. MSA and aerosol size

Sulfur is present in the particles as S(IV) (sum of  $\text{SO}_2$ ,  $\text{HSO}_3^-$ ,  $\text{SO}_3^{2-}$ , and MSA) and S(VI) (sum of  $\text{H}_2\text{SO}_4$ ,  $\text{HSO}_4^-$ , and  $\text{SO}_4^{2-}$ ). In many studies MSA is not included in the definition of S(IV); in the following we refer to the sum of  $\text{SO}_2$ ,  $\text{HSO}_3^-$ , and  $\text{SO}_3^{2-}$  as S(IV)\*, which is a very small fraction of the particulate sulfur and is rapidly oxidized to S(VI). Seawater contains sulfate as well, which gets incorporated into sea salt aerosols. In order to distinguish between the additional sulfate that is taken up, usually the sea salt fraction of sulfate is subtracted in measurements (based on the constant ratios of sulfate and sodium in seawater) and this property is called non-sea-salt  $\text{SO}_4^{2-}$  (nss- $\text{SO}_4^{2-}$ ). We use this term nss- $\text{SO}_4^{2-}$  to refer to the sum of  $\text{H}_2\text{SO}_4$ ,  $\text{HSO}_4^-$ , and  $\text{SO}_4^{2-}$  not originating directly from the sea water. In the following sections we will discuss the production of particulate sulfur, and the MSA to nss- $\text{SO}_4^{2-}$  ratio.

##### 4.3.1. Production of particulate sulfur

In cloudy runs the conversion of the DMS-derived sulfur into particulate sulfur (MSA and nss- $\text{SO}_4^{2-}$ ) is almost complete. In the cloud-free runs only 5–15% is taken up by particles, the rest remains in the gas phase mainly as  $\text{SO}_2$  and eventually gets deposited. The deposition rates of  $\text{SO}_2$  in the cloud-free runs are about ten times higher than in the cloudy runs.

Under **cloud-free** conditions the production of particulate sulfur is always higher when halogen chemistry is included. In the “Luc” scenarios the difference is about 19% whereas it is 26% in the “Yin” and about 45–48% in the “Kuk” scenarios, respectively. In winter the differences increase to 65–94%.

In the cloud-free runs 61–93% of the total MSA production occurs in the gas phase in summer and 70–88% in winter, respectively. If neither of the reactions  $\text{MSIA} + \text{O}_3$

### DMS oxidation

R. von Glasow and P. J. Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

(tested only for “Luc” scenarios) nor MSIA + OH are included as sources for MSA(g) than the contribution of the gas phase is reduced to about 2% in “Yin”, 10% in “Yin, winter”, and 5–6% in “Kuk 0%”. A large fraction of the  $\text{nss-SO}_4^{2-}$  production takes place in the gas phase but it is less important in runs with halogen chemistry. The contribution by HOX (X=Cl, Br) in the sea salt aerosol is 42–51% in summer and 71–75% in winter, respectively, but it is only about 9% in the run for tropical conditions. Neglecting halogen chemistry reduces the production of  $\text{nss-SO}_4^{2-}$  in all cloud-free runs (except for the “tropics” scenarios where it remains about the same), the change in the production of MSA (=P(MSA)) is not uniform. In the runs with a significant fraction of MSA formation in the gas phase (“Luc”, “Kuk 5%”, and “tropics”) P(MSA) is higher with halogen chemistry.

The oxidation of  $\text{S(IV)}^*$  in aerosol particles by  $\text{O}_3$  is only important for high pH values. Due to the vertical decrease in seasalt aerosol pH (see von Glasow and Sander, 2001) this is the case only at the bottom of the MBL. Most of the sulfate production via this pathway occurs in the lowest 50 - 100 m and is rapidly deposited to the ocean. For the cloud-free model runs the contribution of  $\text{O}_3$  to the total  $\text{nss-SO}_4^{2-}$  production rate is less than 2% in summer and about 7% in winter. For the runs without halogen chemistry, however, the relative contribution increased to 8–12% in summer and around 50% in winter. The contribution of HOCl and HOBr is about 42–54% in summer and up to 76% in winter.  $\text{H}_2\text{O}_2$  is also an important oxidant in aerosols with 5–10% in halogen runs and 16–22% in runs without halogen chemistry (but see the “Cape Grim” scenarios with increased alkalinity below).

In the runs with **stratiform clouds** the total  $\text{nss-SO}_4^{2-}$  production rate is higher for the runs without halogen chemistry. This can mainly be explained by greater concentrations of  $\text{S(IV)}^*$  due to a greater importance of the DMS - abstraction pathway. Furthermore higher ozone concentrations in the gas phase lead to higher OH and  $\text{H}_2\text{O}_2$  concentrations in both gas and aqueous phase and therefore greater importance of the reaction  $\text{H}_2\text{O}_{2,\text{aq}} + \text{S(IV)}$ .

Under cloudy conditions  $\text{S(IV)}$  oxidation by  $\text{O}_3$  is of minor importance due to low pH

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

values of the order of 4–5. In summer it is less than 1.5% and in winter less than 4%. As already discussed in von Glasow et al. (2002a) the oxidation of S(IV) by HOCl and HOBr contributes significantly to the total nss-SO<sub>4</sub><sup>2-</sup> production rate with about 40%. H<sub>2</sub>O<sub>2</sub> is the most important oxidant with 50–55% in halogen and 86–91% in cloud free runs. The total production of particulate sulfur is higher if halogen chemistry included because of the increase in MSA production in these cases. This is caused by the production of DMSO in the reaction DMS + BrO and consecutive uptake of DMSO to the particles and oxidation of it in the aqueous phase. More than 98% of the MSA is produced in the particles via MSIA<sub>aq</sub> + OH<sub>aq</sub>, mainly (90–97%) in sulfate droplets.

The results were different for the **Cape Grim** scenarios. Due to higher windspeeds the flux of sea salt aerosols and therefore of alkalinity is greater than in the previously discussed process study runs and consequently the sea salt aerosol and sea salt cloud droplet pH values are higher. This is especially true for the “2x HCO<sub>3</sub><sup>-</sup>”, “2.5x HCO<sub>3</sub><sup>-</sup>”, and “4.5x HCO<sub>3</sub><sup>-</sup>” runs where we increased the bicarbonate content of each aerosol particle. The pH of sea water is about 8 but due to shrinking and loss of water the pH increases in sea salt aerosols near the surface to 9 in the “Cape Grim, summer” run and to 10 in the runs with increased sea salt alkalinity. In sea salt cloud droplets the pH is about 5–5.5 in the “Cape Grim, summer” run and up to 7.3 in the runs with enhanced HCO<sub>3</sub><sup>-</sup>. Therefore the sea salt droplets but especially aerosol pH is high enough to make O<sub>3, aq</sub> an important oxidant. The total contribution of O<sub>3</sub> to nss-SO<sub>4</sub><sup>2-</sup> production increased from 15.9% to 39.2%, 45.5%, and 51.9% in the runs “Cape Grim, summer”, “2x HCO<sub>3</sub><sup>-</sup>”, “2.5x HCO<sub>3</sub><sup>-</sup>”, and “4.5x HCO<sub>3</sub><sup>-</sup>”, respectively. The total production rate of nss-SO<sub>4</sub><sup>2-</sup>, however, increased only insignificantly among the runs with different HCO<sub>3</sub><sup>-</sup> enrichment. This indicates that the oxidation of S(IV)<sup>\*</sup> reached a saturation point and that merely redistribution among the different pathways occurs. Virtually all nss-SO<sub>4</sub><sup>2-</sup> production via O<sub>3</sub> occurred in sea salt particles. The partitioning between sea salt aerosol and droplets changes with increasing buffer capacity. In the “Cape Grim, summer” scenario 96% of the O<sub>3</sub> related nss-SO<sub>4</sub><sup>2-</sup> production is in sea salt aerosol and 3.7% in sea salt droplets. The sea salt aerosol contribution to total

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

nss-SO<sub>4</sub><sup>2-</sup> production decreases to 85%, 78.6%, and 70.7% but the sea salt droplet contribution increases to 15.3%, 20.4%, and 28.9% with increasing HCO<sub>3</sub><sup>-</sup> enrichment in the three cases with increased alkalinity. The shift of nss-SO<sub>4</sub><sup>2-</sup> production from sulfate droplets to sea salt particles leads to an increase in nss-SO<sub>4</sub><sup>2-</sup> deposition of 55% in the “2x HCO<sub>3</sub><sup>-</sup>” run and 70% in the runs with higher HCO<sub>3</sub><sup>-</sup> enrichment.

Even though the gas phase halogen chemistry lost significantly in importance with HCO<sub>3</sub><sup>-</sup> enrichment, the nss-SO<sub>4</sub><sup>2-</sup> production by HOCl and HOBr still played an important role with 40.9%, 22.9%, 21.3%, and 17.9% of the total in the runs without HCO<sub>3</sub><sup>-</sup> enrichment, with 2x, 2.5x, and 4.5x HCO<sub>3</sub><sup>-</sup> enrichment, respectively.

Reduced DMS emissions result in a strong decrease in the production of particulate sulfur in the “Cape Grim, winter” runs. Due to the absence of significant sources of acidity in winter the Br<sup>-</sup> deficit is similar to observations with a maximum of 7% leading to the already mentioned small BrO mixing ratios. But again even small concentrations of HOCl and HOBr in the droplets lead to a contribution of 22% of these to species to total nss-SO<sub>4</sub><sup>2-</sup> production.

In the cloud-free “Cape Grim, summer” run the nss-SO<sub>4</sub><sup>2-</sup> production rate increases from 23.3 mol m<sup>-3</sup> d<sup>-1</sup> in the run without enrichment of alkalinity to 31.7 mol m<sup>-3</sup> d<sup>-1</sup> in the 2.5x HCO<sub>3</sub><sup>-</sup> run but it decreases again to 30.9 mol m<sup>-3</sup> d<sup>-1</sup> in the 4.5x HCO<sub>3</sub><sup>-</sup> run. This is due to a reduction in the oxidation of DMS caused by less BrO and therefore less precursors for particulate sulfur. The fraction of oxidized DMS that ends up in particles, however, increases from 39% to 62% and 64% in the three runs it therefore does not show the saturation effect that we observed in the cloudy “Cape Grim” runs. This fraction is also significantly higher than in the other cloud-free runs that we discussed.

The aqueous phase reaction of MSA with OH (forming sulfate) is unimportant both as sink for MSA and as source of nss-SO<sub>4</sub><sup>2-</sup> in cloud-free runs except for the “tropics” scenario where it accounts for 0.2% and 4.4% of the total nss-SO<sub>4</sub><sup>2-</sup> production with and without halogens, respectively. In cloudy runs, however, it contributes about 8–10% in no-halogen runs and 11–15% in halogen runs to the total oxidation rate but

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGU 2003



in our “Cape Grim, winter” scenarios the contribution is 27% and 10%, for runs with and without halogen chemistry, respectively. These rates indicate that the reaction  $\text{MSA}_{\text{aq}} + \text{OH}$  is a major contributor to  $\text{nss-SO}_4^{2-}$  production in the cloudy MBL. In our “Cape Grim, summer” cases it contributed only 2–3%.

#### 5 4.3.2. $\text{MSA}:\text{nss-SO}_4^{2-}$

In field campaigns often the ratio of MSA to  $\text{nss-SO}_4^{2-}$  is measured in aerosol particles. The MSA to  $\text{nss-SO}_4^{2-}$  ratio is dependent not only on processes occurring during a certain time period that is being studied but also on the particle history. In the field these effects cannot be separated but in a model they can. In the model we initialized our sulfate aerosol without MSA and therefore have small absolute MSA to  $\text{nss-SO}_4^{2-}$  ratios even after strong uptake of MSA. To avoid this problem we discuss “ $\text{P}(\text{MSA}):\text{P}(\text{nss-SO}_4^{2-})$ ” values, using the rates of MSA and  $\text{nss-SO}_4^{2-}$  production within the particles including uptake from the gas phase. Especially in the sulfate particles this changes the ratios drastically. To illustrate this we list two examples from the “Kuk 5%” scenarios for sulfate aerosol: cloud-free  $\text{MSA}:\text{nss-SO}_4^{2-} = 0.013$ ,  $\text{P}(\text{MSA}):\text{P}(\text{nss-SO}_4^{2-}) = 0.331$ ;  
15 cloudy:  $\text{MSA}:\text{nss-SO}_4^{2-} = 0.089$ ,  $\text{P}(\text{MSA}):\text{P}(\text{nss-SO}_4^{2-}) = 0.472$ .

The  $\text{P}(\text{MSA})$  to  $\text{P}(\text{nss-SO}_4^{2-})$  ratios are only then the same quantities as measured in the field when the ocean was the only source for sulfur and when no significant collision-coalescence of particles with different origin occurred. If part of the aerosol population has been advected from the continent with pollution source for sulfate, the MSA to  $\text{nss-SO}_4^{2-}$  ratio is smaller. Therefore one might expect that the modeled ratios are higher than the measured ones except for very clean regions without any advection of pollution. Values for  $\text{P}(\text{MSA}):\text{P}(\text{nss-SO}_4^{2-})$  are listed in Table 3.

25 Measured MSA to  $\text{nss-SO}_4^{2-}$  ratios span a wide range depending on the importance of pollution, which decreases the value, and latitude: generally higher values are measured at higher latitudes. Reported values of  $\text{MSA}:\text{nss-SO}_4^{2-}$  are between 0.002 (Kouvarakis and Mihalopoulos, 2002) and 0.44 (Sciare et al., 2001).

### DMS oxidation

R. von Glasow and P. J. Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

In the model runs most of the  $\text{nss-SO}_4^{2-}$  and MSA are in the sulfate particles which is qualitatively confirmed by Saltzman et al. (1986) who found that most of these species can be found in small particles.

The  $P(\text{MSA})$  to  $P(\text{nss-SO}_4^{2-})$  ratio is reduced in sea salt particles in all cloud-free runs when halogen chemistry is included, indicating that in these cases the increase in MSA production is less important than increase due to production of  $\text{nss-SO}_4^{2-}$  in the particles.

For the sulfate aerosols in the cloud-free runs the ratios are very similar in halogen and no-halogen runs for the “Yin” and the “Kuk 0%” scenarios but the ratio is higher in the halogen runs for the other scenarios. This is caused by the high fraction of MSA being produced in the aerosols of about 33% and 45% in “Yin” and the “Kuk 0%”, respectively and very little uptake of MSA from the gas phase.

Neglecting halogen chemistry reduces the  $P(\text{MSA})$  to  $P(\text{nss-SO}_4^{2-})$  ratio in the droplets under cloudy conditions, indicating that in these cases the increase in MSA production is more important than increase due to production of  $\text{nss-SO}_4^{2-}$  in the particles.

#### 4.3.3. Seasonal variation of $\text{MSA}:\text{nss-SO}_4^{2-}$ at Cape Grim

For temperatures approximately below  $8^\circ\text{C}$  the addition reaction of OH with DMS becomes faster than abstraction. This implies higher yields of MSIA which is either taken up by particles and reacts to MSA or reacts in the gas phase to MSA and/or produces  $\text{CH}_3\text{SO}_2$  which reacts to  $\text{CH}_3\text{SO}_3$ . Under cold conditions the formation of MSA from  $\text{CH}_3\text{SO}_3$  is more important so the general expectation is that  $\text{MSA}:\text{nss-SO}_4^{2-}$  is higher in winter than in summer.

The field data of Ayers et al. (1991) and Andreae et al. (1999) from Cape Grim, Tasmania, show higher DMS, particulate MSA, and  $\text{nss-SO}_4^{2-}$  in summer than in winter but also a peak of the MSA to  $\text{nss-SO}_4^{2-}$  ratio in summer. This implies that other processes than just the previously mentioned temperature dependence must play a

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

role or that the assumption of MSA production only from the reaction  $\text{CH}_3\text{SO}_3 + \text{HO}_2$  (see e.g. Yin et al. (1990) mechanism) is wrong.

Ayers et al. (1996) compared field data with a photochemical model that uses a chemical mechanism which is mainly based on Yin et al. (1990), assuming that MSA is only formed in the abstraction pathway. To fit their model to data observed at Cape Grim, however, they have to assume a very high rate coefficient for the thermal decomposition of  $\text{CH}_3\text{SO}_3$  ( $k = 1.1 \times 10^{17} \exp(-12057/T)$ ). Andreae et al. (1999) proposed that  $\text{SO}_2$  sources other than the oxidation of DMS play a role in the seasonal variation of  $\text{MSA:nss-SO}_4^{2-}$ .

Another possibility is to assume that MSA formation is also occurring in the addition pathway. Koga and Tanaka (1996) assume that MSA is predominantly produced in the addition pathway via assumed reactions of the  $\text{DMS-OH}$  adduct. They pointed to the importance of the  $\text{NO}_3 + \text{DMS}$  reaction that according to their box-model runs explained the difference between winter and summer. In their model the reaction  $\text{DMS} + \text{NO}_3$  is the main sink for  $\text{NO}_3$  and a smaller ocean-flux of DMS in winter implies a smaller sink for  $\text{NO}_3$ . This results in  $\text{NO}_3$  concentrations that are a factor of 4–5 higher in winter than in summer. They calculate a reduction in OH concentrations of a factor of 8 from summer to winter. The net result is a decrease in  $\text{MSA:nss-SO}_4^{2-}$  in winter compared to summer.

A greater importance of halogens in summer than in winter would be an additional explanation for higher  $\text{MSA:nss-SO}_4^{2-}$  in summer than in winter if clouds are present and/or if MSA is produced in the gas phase from MSIA (see numbers in Table 3). This idea is supported by the measurements of Ayers et al. (1999) who showed that at Cape Grim, the loss of bromine from sea salt aerosol was strongest in summer.

#### 4.3.4. Conclusions

In cloud-free model runs between 5 and 15% of the oxidized DMS reacts further to particulate sulfur, in cloudy runs this fraction is almost 100%. In general more particulate sulfur is formed when halogen chemistry is included. If  $\text{HCO}_3^-$  enrichment is prescribed

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

in sea salt aerosol the sea salt aerosol and cloud droplet pH values are high enough to make the reaction of  $S(IV)^*$  with  $O_3$  dominant. It leads to a shift from MSA to  $nss-SO_4^{2-}$  production and increases the total  $nss-SO_4^{2-}$  only somewhat because almost all available sulfur is already oxidized to particulate sulfur in the base scenario. It would also lead to an increase in sulfate deposition by 55–70%. High enrichments of  $HCO_3^-$  in fresh sea salt lead to bromide deficits in the aerosol that are too small and sea salt aerosol pH values that are too high compared with measurements, so that additional acidity sources would have to be active if these enrichments were widespread over the oceans. Without increased alkalinity the gas phase mixing ratios of BrO are too high which could be due to neglecting reactions with organic compounds that can sequester bromine.

Including halogen chemistry leads to greater formation rates of particulate sulfur under most conditions. HOCl and HOBr are important for the oxidation of  $S(IV)^*$  in particles even when the gas phase halogen concentrations and bromide release from the sea salt aerosol are small.

We found the reaction  $MSA_{aq} + OH_{aq}$  to contribute about 10–15% to the production of  $nss-SO_4^{2-}$  in clouds. It is unimportant for cloud-free model runs.

An additional explanation for the higher MSA: $nss-SO_4^{2-}$  ratio in aerosols at Cape Grim in summer than winter (contrary to what is expected from the temperature dependence of the involved reactions) might be the greater influence of halogens in summer in the production of MSA.

#### 4.4. Diurnal variation of $Br^-$

Our “generic” cloud-free model runs (e.g. “kuk, 5%”) have a small  $Br^-$  to  $Na^+$  in the sea salt aerosol or in other words a very high bromine deficit, implying that most of the bromine is in the gas phase. Then the diurnal variation of sea salt  $Br^-$  is mainly determined by exchange processes between particulate phases and the gas phase and  $Br^-$  shows a peak during day and a minimum during night (see also von Glasow et al.,

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

2002b). Under conditions when the multiphase catalytic cycles are not as efficient as in the base scenario due to lower gas phase acidity, lower gas phase  $O_3$ , and/or higher temperature, then the additional bromide from the emission of fresh sea salt is not as rapidly released and accumulates over night in the sea salt aerosol and is depleted during day. This leads to a reversed diurnal variation in sea salt aerosol  $Br^-$  and bromide deficit as is the case in the “tropics” and “Cape Grim” scenarios. Figure 6 shows the  $Br^-$  deficit for the “kuk, 5%” and the “tropics” scenarios.

This is in accordance with field studies (Rancher and Kritz, 1980; Sander et al., 2003; Pszenny et al., 2003) that found under conditions with bromine deficits of less than 90% that sea salt  $Br^-$  has a maximum in the early morning and is being depleted during the day.

In von Glasow et al. (2002a) we showed that the presence of a cloud also changes the diurnal variation of the  $Br^-$  concentration and brings it into accordance with measurements.

## 5. Climate relevance

We could show that halogens play an important role in the cycle of sulfur in the atmosphere.  $BrO$  increases the oxidation rate of DMS even if it is present in small concentrations only and  $HOCl$  and  $HOBr$  increase the oxidation of  $S(IV)^*$  even if gas phase halogen concentrations are very small (below the detection limit of current instruments). In most cases the presence of halogens leads to an increase of the production rate of particulate sulfur.

Under cloud-free conditions the products of the reaction  $MSIA + OH$  are important to estimate the final products of DMS addition. If MSA is formed in this reaction - even with a very small yield - than most MSA is being formed in the gas phase, if not, then most of it is formed in the aerosol phase. If no MSA is formed in the gas phase then more  $SO_2$ , a precursor for new particle formation, is formed through the addition pathway. Here, important uncertainties still exist.

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

So far we mentioned only the effects of chemical reactions on the production of particulate sulfur and the precursors of new particle formation that lead to less but bigger aerosol particles. These effects combined would lead to a reduction of cloud albedo, which is often referred to as “first indirect aerosol effect”. Fewer and bigger aerosol particles also leads to another effect: a more efficient initiation of collision-coalescence processes among droplets, the main precipitation formation mechanism in warm (ie ice-free) clouds. This would cause an increase of the wash-out of particles and reduce cloud lifetime thereby even further reducing the cloud albedo. The cloud lifetime effect is often referred to as “second indirect effect”.

Note that these effects all lead to a reduction of cloud albedo as opposed to the anthropogenic indirect aerosol effects that lead to an increase in cloud albedo (e.g. [Intergovernmental Panel on Climate Change, 2001](#)).

*Acknowledgements.* We want to thank I. Barnes, A. Kukui, G. Le Bras, J. Caine, H. Sievering, and D. Rosenfeld for helpful discussions and making their work available to us before publication.

## References

- Allen, H. C., Gragson, D. E., and Richmond, G. L.: Molecular structure and adsorption of dimethyl sulfoxide at the surface of aqueous solutions, *J. Phys. Chem. B*, 103, 660–666, 1999. [6744](#)
- Andreae, M. O., Elbert, W., Cai, Y., Andreae, T. W., and Gras, J.: Non-sea-salt sulfate, methanesulfonate, and nitrate aerosol concentrations and size distributions at cape grim, tasmania, *J. Geophys. Res.*, 104, 21 695–21 706, 1999. [6758](#), [6759](#)
- Arsene, C., Barnes, I., Becker, K. H., Schneider, W. F., Wallington, T. T., Mihalopoulos, N., and Patroescu-Klotz, I. V.: Formation of methane sulfinic acid in the gas-phase OH-radical initiated oxidation of dimethyl sulfoxide, *Environ. Sci. Technol.*, 36, 5155–5163, 2002. [6742](#), [6743](#)
- Atlas, E., Pollock, W., Greenberg, J., Heidt, L., and Thompson, A. M.: Alkyl nitrates, non-

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGU 2003

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

- methane hydrocarbons, and halocarbon gases over the equatorial pacific-ocean during saga-3, *J. Geophys. Res.*, 98, 16 933–16 947, 1993. [6741](#)
- Ayers, G. P., Ivey, J. P., and Gillett, R. W.: Coherence between seasonal cycles of dimethylsulphide during a phytoplankton bloom, *Nature*, 349, 404–406, 1991. [6758](#)
- 5 Ayers, G. P., Bartley, S. T., Ivey, J. P., and Forgan, B. W.: Dimethylsulfide in marine air at cape grim, 41's, *J. Geophys. Res.*, 100, 21 013–21 021, 1995. [6740](#), [6768](#)
- Ayers, G. P., Cainey, J. M., Granek, H., and Leck, C.: Dimethylsulfide oxidation and the ratio of methanesulfonate to non sea-salt sulfate in the marine aerosol, *J. Atmos. Chem.*, 25, 307–325, 1996. [6759](#)
- 10 Ayers, G. P., Cainey, J. M., Gillett, R. W., Saltzman, E. S., and Hooper, M.: Sulfur dioxide and dimethyl sulfide in marine air at cape grim, tansania, *Tellus*, 49B, 292–299, 1997a. [6740](#), [6768](#)
- Ayers, G. P., Granek, H., and Boers, R.: Ozone in the marine boundary layer at cape grim: Model simulation, *J. Atmos. Chem.*, 27, 179–195, 1997b. [6740](#), [6768](#)
- 15 Ayers, G. P., Gillett, R. W., Cainey, J. M., and Dick, A. L.: Chloride and bromide loss from sea-salt particles in southern ocean air, *J. Atmos. Chem.*, 33, 299–319, 1999. [6740](#), [6750](#), [6759](#), [6768](#)
- Bardouki, H., Berresheim, H., Vrekoussis, M., Sciare, J., Kouvarakis, G., Oikonomou, K., Schneider, J., and Mihalopoulos, N.: Gaseous (dms, msa, SO<sub>2</sub>, H<sub>2</sub>SO<sub>4</sub> and dmsO) and particulate (sulfate and methanesulfonate) sulfur species over the northeastern coast of crete, *Atmos. Chem. Phys. Discuss.*, 3, 3869–3906, 2003. [6745](#), [6746](#)
- 20 Barnes, I., Bastian, V., Becker, K. H., and Overath, R. D.: Kinetic studies of the reactions of IO, BrO and ClO with DMS, *Int. J. Chem. Kinet.*, 23, 579–591, 1991. [6736](#)
- Barry, R. G. and Chorley, R. J.: *Atmosphere, Weather & Climate*, Routledge, 1998. [6735](#)
- 25 Bedjanian, Y., Poulet, G., and Bras, G. L.: Kinetic study of thereaction of bro radicals with dimethylsulfide, *Int. J. Chem. Kinet.*, 28, 383–389, 1996. [6736](#)
- Boucher, O., Moulin, C., Belviso, S., Aumont, O., Bopp, L., Cosme, E., von Kuhlmann, R., Lawrence, G. G., Pham, M., Redyy, M. S., Sciare, J., and Venkataraman, C.: Sensitivity study of dimethylsulphide (dms) atmospheric concentrations and sulphate aerosol indirect radiative forcing to the dms source representation and oxidation, *Atmos. Chem. Phys.*, 3, 49–65, 2003. [6749](#)
- 30 Charlson, R. J., Lovelock, J. E., Andreae, M. O., and Warren, S. G.: Oceanic phytoplankton, atmospheric sulphur, cloud albedo and climate, *Nature*, 326, 655–661, 1987. [6735](#)

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGU 2003

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

- Chuck, A. L., Turner, S. M., and Liss, P. S.: Direct evidence for a marine source of C<sub>1</sub> and C<sub>2</sub> alkyl nitrates, *Science*, 297, 1151–1154, 2002. [6741](#)
- Davis, D., Chen, G., Bandy, A., Thornton, D., Eisele, F., Mauldin, L., Tanner, D., Lenschow, D., Fuelberg, H., Huebert, B., Heath, J., Clarke, A., and Blake, D.: Dimethyl sulfide oxidation in the equatorial pacific: Comparison of model simulations with field observations for dms, SO<sub>2</sub>, H<sub>2</sub>SO<sub>4</sub>(g), msa(g), ms, and nss, *J. Geophys. Res.*, 104, 5765–5784, 1999. [6745](#)
- De Bruyn, W. J., Shorter, J. A., Davidovits, P., Worsnop, D. R., Zahniser, M. S., and Kolb, C. E.: Uptake of gas phase sulfur species methanesulfonic acid, dimethylsulfoxide, and dimethyl sulfone by aqueous surface, *J. Geophys. Res.*, 99, 16 927–16 932, 1994. [6739](#)
- de Bruyn, W. J., Harvey, M. and E. S. Saltzman: DMS and SO<sub>2</sub> at baring head, new zealand: Implications for the yield of SO<sub>2</sub> from DMS, *J. Atmos. Chem.*, 41, 189–209, 2002. [6751](#)
- Falbe-Hansen, H. S. S., Jensen, N. R., Pedersen, T., and Hjorth, J.: Atmospheric gas-phase reactions of dimethylsulphoxide and dimethylsulphone with OH and NO<sub>3</sub> radicals, Cl atoms and ozone, *Atmos. Environ.*, 34, 1543–1551, 2000. [6739](#)
- Gershenzon, M., Davidovits, P., Jayne, J. T., Kolb, C. E., and Worsnop, D. R.: Simultaneous uptake of DMS and ozone on water, *J. Phys. Chem. A*, 105, 7031–7036, 2001. [6749](#)
- Ingham, T., Bauer, D., Sander, R., Crutzen, P. J., and Crowley, J. N.: Kinetics and products of the reactions BrO + DMS and Br + DMS at 298 k, *J. Phys. Chem. A*, 103, 7199–7209, 1999. [6736](#)
- Intergovernmental Panel on Climate Change, *Climate Change 2001 - The Scientific Basis*, Cambridge University Press, 2001. [6762](#)
- Jefferson, A., Tanner, D. J., Eisele, F. L., Davis, D. D., Chen, G., Crawford, J., Huey, J. W., Torres, A. L., and Berresheim, H.: Oh photochemistry and methane sulfonic acid formation in the coastal antarctic boundary layer, *J. Geophys. Res.*, 103, 16 476–16 566, 1998. [6745](#)
- Keene, W. C. and Savoie, D. L.: The ph of deliquesced sea-salt aerosol in polluted marine air, *Geophys. Res. Lett.*, 25, 2181–2184, 1998. [6750](#)
- Keene, W. C. and Savoie, D. L.: Correction to "the ph of deliquesced sea-salt aerosol in polluted marine air", *Geophys. Res. Lett.*, 26, 1315–1316, 1999. [6750](#)
- Keene, W. C., Pszenny, A. A. P., Maben, J. R., and Sander, R.: Variation of marine aerosol acidity with particle size, *Geophys. Res. Lett.*, 29, doi: 10.1029/2001GL013 881, 2002. [6750](#)
- Koga, S. and Tanaka, H.: Simulation of seasonal variations of sulfur compounds in the remote marine atmosphere, *J. Atmos. Chem.*, 23, 163– 192, 1996. [6759](#)
- Kouvarakis, G. and Mihalopoulos, N.: Seasonal variation of dimethylsulfide in the gas phase



## DMS oxidation

R. von Glasow and P. J.  
Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

and of methanesulfonate and non-sea-salt sulfate in the aerosols phase in the eastern mediterranean atmosphere, *Atmos. Environ.*, 36, 929–938, 2002. [6757](#)

Kukui, A., Borissenko, D., Laverdet, G., and Bras, G. L.: Gas phase reactions of OH radicals with dimethyl sulfoxide and methane sulfonic acid using turbulent flow reactor and chemical ionization mass spectrometry, *J. Phys. Chem. A*, 107, 5732–5742, 2003. [6742](#), [6743](#), [6769](#)

Lelieveld, J., Crutzen, P. J., and Rodhe, H.: Zonal average cloud characteristics for global atmospheric chemistry modelling, Tech. Rep. CM-76, Dep. of Meteorology, University of Stockholm (MISU), 1989. [6735](#)

Leser, H., Hönninger, G., and Platt, U.: Max-doas measurements of BrO and NO<sub>2</sub> in the marine boundary layer, *Geophys. Res. Lett.*, 30, pages=1537, doi:10.1029/2002GL015811, [6748](#)

Lucas, D. D. and Prinn, R. G.: Mechanistic studies of dimethylsulfide oxidation products using an observationally constrained model, *J. Geophys. Res.*, 107, doi: 10.1029/2001JD000843, 2002. [6742](#), [6743](#), [6745](#), [6746](#), [6769](#)

Mauldin, R. L., Eisele, F. L., Tanner, D. J., Kosciuch, E., Shetter, R., Lefer, B., Hall, S. R., Nowak, J. B., Buhr, M., Chen, G., Wang, P., and Davis, D.: Measurements of OH, H<sub>2</sub>SO<sub>4</sub>, and msa at the south pole during iscat, *Geophys. Res. Lett.*, 28, 3629–3632, 2001. [6745](#)

Monks, P. S., Carpenter, L. J., Penkett, S. A., Ayers, G. P., Gillet, R. W., Galbally, I. E., and Meyer, C. P.: Fundamental ozone photochemistry in the remote marine boundary layer: The SOAPEX experiment, measurements and theory, *Atmos. Environ.*, 32, 3647–3664, 1998. [6740](#), [6768](#)

Monks, P. S., Salisbury, G., Holland, G., Penkett, S. A., and Ayer, G. P.: A seasonal comparison of ozone photochemistry in the remote marine boundary layer, *Atmos. Environ.*, 34, 2547–2561, 2000. [6740](#), [6768](#)

Nakano, Y., Goto, M., Hashimoto, S., Kawasaki, M., and Wallington, T. J.: Cavity ring-down spectroscopic study of the reactions of Br atoms and BrO radicals with dimethyl sulfide, *J. Phys. Chem. A*, 105, 11045–11050, 2001. [6736](#)

Peixoto, J. P. and Oort, A. H.: *Physics of Climate*, American Institute of Physics, New York, 1991. [6739](#)

Pszenny, A. A. P., Moldanova, J., Keene, W. C., Sander, R., Maben, J. R., Martinez, M., Crutzen, P. J., Perner, D., and Prinn, R. G.: Halogen cycling and aerosol ph in the hawaiian marine boundary layer, *Atmos. Chem. Phys. Discuss.*, 3, 4701–4753, 2003. [6748](#), [6750](#), [6761](#)

Rancher, J. and Kritz, M. A.: Diurnal fluctuations of Br and I in the tropical marine atmosphere,

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

J. Geophys. Res., 85, 5581– 5587, 1980. [6761](#)

Ross, A. B., Mallard, W. G., Helman, W. P., Buxton, G. V., Huie, R. E., and Neta, P.: NDRL-NIST Solution kinetics database: Version 3.0, Notre Dame Radiation Laboratory, Notre Dame, and National Institut of Standards and Technology, Gaithersburg, 1998. [6736](#)

5 Saiz-Lopez, A., Plane, J. M. C., and Shillito, J. A.: Bromine oxide in the mid-latitude marine boundary layer, submitted to Science, 2003. [6748](#)

Saltzman, E. S., Savoie, D. L., Prospero, J. M., and Zika, R. G., Methane sulfonic acid and non-sea-salt sulphate in pacific air: Regional and seasonal variations, J. Atmos. Chem., 4, 227–240, 1986. [6758](#)

10 Sander, R. and Crutzen, P. J.: Model study indicating halogen activation and ozone destruction in polluted air masses transported to the sea, J. Geophys. Res., 101, 9121–9138, 1996. [6737](#)

Sander, R., Keene, W. C., Pszenny, A. A. P., Arimoto, R., Ayers, G. P., Baboukas, E., Cainey, J. M., Crutzen, P. J., Duce, R. A., Hönninger, G., Huebert, B. J., Maenhaut, W., Mihalopoulos, N., Turekian, V. C., and Dingenen, R. V.: Inorganic bromine compounds in the marine boundary layer: A critical review, Atmos. Chem. Phys., 3, 1301–1336, 2003. [6738](#), [6761](#)

15 Sciare, J., Baboukas, E., and Mihalopoulos, N.: Short-term variability of atmospheric dms and its oxidation products at amsterdam island during summer time, J. Atmos. Chem., 39, 281–302, 2001. [6757](#)

20 Sievering, H., Lerner, B., Slavich, J., Anderson, J., M. Posfai, M., and Cainey, J.: O<sub>3</sub> oxidation of SO<sub>2</sub> in sea-salt aerosol water: Size distribution of non-sea-salt sulfate during the first aerosol characterization experiment (ace 1), J. Geophys. Res., 104, 21 707–21 717, 1999. [6749](#)

Toumi, R.: BrO as a sink for dimethylsulphide in the marine atmosphere, Geophys. Res. Lett., 21, 117–120, 1994. [6736](#)

25 Toyota, K., Kanaya, Y., Takahashi, M., and Akimoto, H.: A box model study on photochemical interactions between vocs and reactive halogen species in the marine boundary layer, Atmos. Chem. Phys. Discuss., 3, 4549 –4632, 2003. [6750](#)

Twomey, S.: Pollution and the planetary albedo, Atmos. Environ., 8, 1251–1256, 1974. [6735](#)

30 Urbanski, S., Stickel, R. E., and Wine, P. H.: Mechanistic and kinetic study of the gas-phase reaction of hydroxyl radical with dimethyl sulfoxide, J. Phys. Chem. A, 102, 10 522–10 529, 1998. [6742](#)

Vogt, R., Crutzen, P. J., and Sander, R.: A mechanism for halogen release from sea-salt aerosol in the remote marine boundary layer, Nature, 383, 327–330, 1996. [6737](#)

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

- von Glasow, R. and Crutzen, P. J.: Tropospheric halogen chemistry, in *Treatise on Geochemistry*, 2003, in press. [6737](#)
- von Glasow, R. and Sander, R.: Variation of sea salt aerosol pH with relative humidity, *Geophys. Res. Lett.*, 28, 247–250, 2001. [6748](#), [6754](#)
- 5 von Glasow, R., Sander, R., Bott, A., and Crutzen, P. J.: Modeling halogen chemistry in the marine boundary layer. 2. interactions with sulfur and cloud-covered mbl, *J. Geophys. Res.*, 107, 4323, doi: 10.1029/2001JD000943, 2002a. [6738](#), [6755](#), [6761](#)
- von Glasow, R., Sander, R., Bott, A., and Crutzen, P. J.: Modeling halogen chemistry in the marine boundary layer. 1. cloud-free mbl, *J. Geophys. Res.*, 107, 4341, doi: 10.1029/2001JD000942, 2002b. [6738](#), [6741](#), [6748](#), [6752](#), [6760](#)
- 10 Wagner, V., von Glasow, R., Fischer, H., and Crutzen, P. J.: Are CH<sub>2</sub>O measurements in the marine boundary layer suitable for testing the current understanding of CH<sub>4</sub> photooxidation?, *J. Geophys. Res.*, 107, 4029, doi: 10.1029/2001JD000722, 2002. [6740](#), [6768](#)
- Wang, L. and Zhang, J.: Ab initio study of reaction of dimethyl sulfoxide (DMSO) with OH radical, *Chem. Phys. Lett.*, 356, 490–496, 2002. [6742](#)
- 15 Watts, S. F. and Brimblecombe, P.: The Henry law constant of dimethylsulfoxide, *Environ. Technol. Lett.*, 8, 483–486, 1987. [6744](#)
- Yin, F., Grosjean, D., and Seinfeld, J. H.: Photooxidation of dimethyl sulfide and dimethyl disulfide. i: Mechanism development, *J. Atmos. Chem.*, 11, 309–364, 1990. [6743](#), [6745](#), [6759](#), [6769](#)
- 20 Zafiriou, O. C., McFarland, M., and Bromund, R. H.: Nitric oxide in seawater, *Science*, 207, 637–639, 1980. [6741](#)

## DMS oxidation

R. von Glasow and P. J.  
Crutzen**Table 1.** Initial mixing ratios of gas phase species at the surface (in nmol mol<sup>-1</sup>) and DMS emissions in molec cm<sup>-2</sup>.

species	remote MBL	Cape Grim (summer)	Cape Grim (winter)	tropics
O <sub>3</sub>	20.0	15.0	32.0	10.3
NO <sub>2</sub>	0.02	0.01	0.005	0.02
HNO <sub>3</sub>	0.01	0.01	0.01	0.01
PAN	0.01	0.01	0.01	0.01
NH <sub>3</sub>	0.08	0.04	0.02	0.08
H <sub>2</sub> O <sub>2</sub>	0.6	0.7	0.1	0.6
SO <sub>2</sub>	0.09	0.01	0.01	0.02
DMS	0.06	0.12	0.02	0.04
CH <sub>4</sub>	1800.0	1670.0	1690.0	1698.0
CO	70.0	44.0	66.0	54.8
C <sub>2</sub> H <sub>6</sub>	0.5	0.25	0.25	0.0
HCHO	0.3	0.3	0.3	0.2
HCl	0.04	0.01	0.005	0.02
DMS emission	2 × 10 <sup>9</sup>	3.5 × 10 <sup>9</sup>	1.5 × 10 <sup>8</sup>	8 × 10 <sup>8</sup>

Values labeled “remote MBL” are for the “generic” runs and are the same for summer and winter runs whereas different values are used for the Cape Grim runs in summer and winter. Initial conditions for the Cape Grim runs are taken from [Ayers et al. \(1995, 1997a,b, 1999\)](#); [Monks et al. \(1998, 2000\)](#) and initial conditions for the “tropics” run are from [Wagner et al. \(2002\)](#).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

**Table 2.** Overview of oxidation rates.

run	DMS oxidation rate	relative contribution to total oxidation rate				nss-SO <sub>4</sub> <sup>2-</sup> production rate	relative contribution to total production rate				MSA production rate	DMS → SO <sub>2</sub> conversion efficiency
	[mol m <sup>-3</sup> d <sup>-1</sup> ]	BrO	OH(ad)	OH(ab)	NO <sub>3</sub>		OH(g)	HOX <sub>aq</sub>	O <sub>3,aq</sub>	H <sub>2</sub> O <sub>2,aq</sub>		
cloud-free		%	%	%	%	[mol m <sup>-3</sup> d <sup>-1</sup> ]	%	%	%	%	[mol m <sup>-3</sup> d <sup>-1</sup> ]	
Luc	4.38 × 10 <sup>-9</sup>	63.4	12.7	14.7	1.2	3.31 × 10 <sup>-10</sup>	38.8	50.8	1.8	8.5	1.80 × 10 <sup>-10</sup>	0.284
Luc, no hal.	3.49 × 10 <sup>-9</sup>	-	44.3	50.2	5.4	2.97 × 10 <sup>-10</sup>	69.0	-	10.2	20.5	1.37 × 10 <sup>-10</sup>	0.587
Yin	4.37 × 10 <sup>-9</sup>	63.6	12.6	14.6	1.2	4.91 × 10 <sup>-10</sup>	43.0	45.4	1.4	10.1	0.33 × 10 <sup>-10</sup>	0.597
Yin, no hal.	3.49 × 10 <sup>-9</sup>	-	44.3	50.2	5.5	3.71 × 10 <sup>-10</sup>	69.3	-	9.2	21.3	0.42 × 10 <sup>-10</sup>	0.789
Yin, winter	5.00 × 10 <sup>-9</sup>	83.5	5.5	3.1	1.2	2.19 × 10 <sup>-10</sup>	11.8	74.8	7.0	6.2	0.13 × 10 <sup>-10</sup>	0.221
Yin, winter, no hal.	2.54 × 10 <sup>-9</sup>	-	57.7	30.8	11.5	1.23 × 10 <sup>-10</sup>	30.6	-	52.0	16.7	0.17 × 10 <sup>-10</sup>	0.513
Kuk, 0% MSA	4.37 × 10 <sup>-9</sup>	63.8	12.5	14.5	1.3	5.87 × 10 <sup>-10</sup>	46.2	41.9	1.2	10.7	0.18 × 10 <sup>-10</sup>	0.789
Kuk, 0% MSA, no hal	3.48 × 10 <sup>-9</sup>	-	44.3	50.2	5.6	4.18 × 10 <sup>-10</sup>	69.1	-	8.4	21.3	0.19 × 10 <sup>-10</sup>	0.905
Kuk, 5% MSA	4.37 × 10 <sup>-9</sup>	63.8	12.5	14.5	1.3	5.70 × 10 <sup>-10</sup>	46.2	41.9	1.2	10.6	1.38 × 10 <sup>-10</sup>	0.762
Kuk, 5% MSA, no hal.	3.48 × 10 <sup>-9</sup>	-	44.3	50.2	5.6	4.15 × 10 <sup>-10</sup>	68.6	-	8.2	21.2	0.79 × 10 <sup>-10</sup>	0.888
Kuk, 5% MSA, winter	5.01 × 10 <sup>-9</sup>	83.8	5.3	2.9	1.2	2.65 × 10 <sup>-10</sup>	14.1	71.4	7.0	7.3	0.89 × 10 <sup>-10</sup>	0.406
Kuk, 5% MSA, winter, no hal.	2.52 × 10 <sup>-9</sup>	-	57.6	30.8	11.6	1.36 × 10 <sup>-10</sup>	31.6	-	49.7	17.7	0.44 × 10 <sup>-10</sup>	0.653
Kuk, 10% MSA	4.38 × 10 <sup>-9</sup>	63.8	12.5	14.5	1.3	5.54 × 10 <sup>-10</sup>	46.2	41.9	1.1	10.6	2.58 × 10 <sup>-10</sup>	0.734
Kuk, 10% MSA, no hal.	3.48 × 10 <sup>-9</sup>	-	44.3	50.2	5.6	4.09 × 10 <sup>-10</sup>	68.5	-	8.0	21.1	1.39 × 10 <sup>-10</sup>	0.871
tropics	1.17 × 10 <sup>-9</sup>	16.9	19.6	51.0	9.5	1.77 × 10 <sup>-10</sup>	89.0	8.9	0.3	1.7	0.20 × 10 <sup>-10</sup>	0.929
tropics, no hal.	1.13 × 10 <sup>-9</sup>	-	24.1	62.7	13.1	1.79 × 10 <sup>-10</sup>	93.3	-	0.3	1.8	0.14 × 10 <sup>-10</sup>	0.957

See text for explanation of the different model runs. The numbers are means over the MBL from the 2nd until 5th model day. The contribution of the individual path is given as relative contribution to the total oxidation rate. The main difference between the runs is the rate constant of the reaction MSIA + OH. The following values were chosen: runs labeled "Luc" -  $k=1.0 \times 10^{-12} \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$  (Lucas and Prinn (2002), also including the reaction MSIA + O<sub>3</sub>), runs labeled "Yin" -  $k=1.6 \times 10^{-11} \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$  (Yin et al., 1990), runs labeled "Kuk" -  $k=9.0 \times 10^{-11} \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$  (Kukui et al., 2003). The percentages given for the "Kuk" runs are the yield of MSA in the reaction MSIA + OH.

5

**DMS oxidation**

R. von Glasow and P. J. Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

Table 2. Continued.

run	DMS oxidation rate	relative contribution to total oxidation rate				nss-SO <sub>4</sub> <sup>2-</sup> production rate	relative contribution to total production rate				MSA production rate	DMS → SO <sub>2</sub> conversion efficiency
	[mol m <sup>-3</sup> d <sup>-1</sup> ]	BrO %	OH(ad) %	OH(ab) %	NO <sub>3</sub> %	[mol m <sup>-3</sup> d <sup>-1</sup> ]	OH(g) %	HOX <sub>aq</sub> %	O <sub>3,aq</sub> %	H <sub>2</sub> O <sub>2,aq</sub> %	[mol m <sup>-3</sup> d <sup>-1</sup> ]	
Cape Grim												
summer	6.45 × 10 <sup>-9</sup>	77.8	7.1	10.7	0.8	23.33 × 10 <sup>-10</sup>	10.7	40.8	47.2	1.2	2.11 × 10 <sup>-10</sup>	0.700
summer, no hal.	4.95 × 10 <sup>-9</sup>	-	39.4	56.1	4.5	21.53 × 10 <sup>-10</sup>	10.5	-	86.8	2.0	0.94 × 10 <sup>-10</sup>	0.892
summer, 2.5x HCO <sub>3</sub> <sup>-</sup>	5.26 × 10 <sup>-9</sup>	15.9	32.9	47.2	2.7	31.65 × 10 <sup>-10</sup>	3.2	17.6	78.5	0.4	1.08 × 10 <sup>-10</sup>	0.827
summer, 4.5x HCO <sub>3</sub> <sup>-</sup> cloudy	4.97 × 10 <sup>-9</sup>	2.5	38.4	54.7	4.0	30.90 × 10 <sup>-10</sup>	2.3	22.4	74.8	0.3	0.95 × 10 <sup>-10</sup>	0.804
Yin												
Yin, no hal.	3.43 × 10 <sup>-9</sup>	35.4	26.6	28.4	1.6	15.94 × 10 <sup>-10</sup>	1.4	39.4	0.7	41.0	17.66 × 10 <sup>-10</sup>	0.406
Kuk, 0% MSA	3.43 × 10 <sup>-9</sup>	35.4	26.6	28.4	1.6	18.24 × 10 <sup>-10</sup>	1.7	39.0	0.7	44.6	15.12 × 10 <sup>-10</sup>	0.493
Kuk, 0% MSA, no hal	3.08 × 10 <sup>-9</sup>	-	40.5	43.2	9.7	20.74 × 10 <sup>-10</sup>	1.6	-	1.3	82.5	9.85 × 10 <sup>-10</sup>	0.652
Kuk, 5% MSA	3.43 × 10 <sup>-9</sup>	35.4	26.6	28.4	1.6	18.03 × 10 <sup>-10</sup>	1.6	38.9	0.7	44.2	15.39 × 10 <sup>-10</sup>	0.485
Kuk, 5% MSA, no hal.	3.08 × 10 <sup>-9</sup>	-	40.5	43.2	9.7	20.60 × 10 <sup>-10</sup>	1.6	-	1.3	82.3	10.04 × 10 <sup>-10</sup>	0.646
Kuk, 5% MSA, winter	3.44 × 10 <sup>-9</sup>	64.0	15.5	8.6	2.1	6.65 × 10 <sup>-10</sup>	0.1	54.4	1.5	27.9	21.18 × 10 <sup>-10</sup>	0.140
Kuk, 5% MSA, winter, no hal.	2.01 × 10 <sup>-9</sup>	-	39.6	22.0	18.8	9.96 × 10 <sup>-10</sup>	0.2	-	4.0	68.3	10.60 × 10 <sup>-10</sup>	0.419
Cape Grim												
summer	4.01 × 10 <sup>-9</sup>	38.2	29.2	29.3	1.2	15.56 × 10 <sup>-10</sup>	1.0	40.9	15.9	38.1	15.37 × 10 <sup>-10</sup>	0.451
summer, no hal.	3.61 × 10 <sup>-9</sup>	-	46.0	46.4	5.8	22.07 × 10 <sup>-10</sup>	1.2	-	33.8	57.9	10.65 × 10 <sup>-10</sup>	0.644
summer, 2x HCO <sub>3</sub> <sup>-</sup>	3.63 × 10 <sup>-9</sup>	4.6	44.2	44.5	4.5	17.57 × 10 <sup>-10</sup>	0.9	22.9	39.3	31.9	11.32 × 10 <sup>-10</sup>	0.614
summer, 2.5x HCO <sub>3</sub> <sup>-</sup>	3.61 × 10 <sup>-9</sup>	2.5	45.0	45.4	4.9	17.73 × 10 <sup>-10</sup>	0.9	21.3	45.5	27.1	11.19 × 10 <sup>-10</sup>	0.625
summer, 4.5x HCO <sub>3</sub> <sup>-</sup>	3.60 × 10 <sup>-9</sup>	1.3	45.5	45.9	5.2	17.88 × 10 <sup>-10</sup>	0.8	17.9	51.9	23.9	11.19 × 10 <sup>-10</sup>	0.630
winter	0.13 × 10 <sup>-9</sup>	15.8	13.6	8.9	40.4	0.81 × 10 <sup>-10</sup>	0.1	22.2	38.2	1.6	0.59 × 10 <sup>-10</sup>	0.499
winter, no hal.	0.18 × 10 <sup>-9</sup>	-	7.7	5.0	77.1	1.46 × 10 <sup>-10</sup>	0.0	-	78.5	2.6	0.30 × 10 <sup>-10</sup>	0.815

See text and caption of Table 2 for explanation of the different model runs. The numbers are means over the MBL over the 2nd and 3rd model day. The contribution of the individual path is given as relative contribution to the total oxidation rate.

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**Table 3.** Modeled P(MSA) - P(nss-SO<sub>4</sub><sup>2-</sup>) ratios.

run	aerosol		droplets	
	sulfate	sea salt	sulfate	sea salt
<b>cloud-free</b>				
Luc	0.958	0.316	-	-
Luc, no hal.	0.406	0.534	-	-
Yin	0.064	0.071	-	-
Yin, no hal.	0.068	0.187	-	-
Yin, winter	0.174	0.043	-	-
Yin, winter, no hal.	0.143	0.131	-	-
Kuk, 0% MSA	0.036	0.026	-	-
Kuk, 0% MSA, no hal.	0.037	0.059	-	-
Kuk, 5% MSA	0.402	0.125	-	-
Kuk, 5% MSA, no hal.	0.201	0.167	-	-
Kuk, 5% MSA, winter	1.744	0.118	-	-
Kuk, 5% MSA, winter, no hal.	0.668	0.167	-	-
Kuk, 10% MSA	0.789	0.229	-	-
Kuk, 10% MSA, no hal.	0.370	0.279	-	-
<hr/>				
tropics	0.115	0.115	-	-
tropics, no hal.	0.016	0.157	-	-
CG, summer	0.632	0.033	-	-
CG, summer, no hal.	0.285	0.017	-	-
CG, summer, 2.5x HCO <sub>3</sub> <sup>-</sup>	0.742	0.013	-	-
CG, summer, 4.5x HCO <sub>3</sub> <sup>-</sup>	0.873	0.011	-	-
<b>cloudy</b>				
Yin	0.289	0.252	1.153	0.786
Yin, no hal.	0.629	0.399	0.612	0.456
Kuk, 0% MSA	0.209	0.136	0.868	0.560
Kuk, 0% MSA, no hal.	0.540	0.273	0.478	0.363
Kuk, 5% MSA	0.472	0.386	0.882	0.569
Kuk, 5% MSA, no hal.	0.643	0.389	0.484	0.366
Kuk, 5% MSA, winter	1.429	1.062	3.307	1.178
Kuk, 5% MSA, winter, no hal.	0.775	0.442	1.098	0.614
<hr/>				
CG, summer	0.813	0.040	1.601	0.108
CG, summer, no hal.	0.553	0.049	0.785	0.173
CG, summer, 2x HCO <sub>3</sub> <sup>-</sup>	0.857	0.017	1.672	0.050
CG, summer, 2.5x HCO <sub>3</sub> <sup>-</sup>	0.894	0.016	1.878	0.044
CG, summer, 4.5x HCO <sub>3</sub> <sup>-</sup>	0.952	0.018	2.048	0.050
CG, winter	0.105	0.032	1.200	0.090
CG, winter, no hal.	1.174	0.006	0.062	0.015

See text for explanation of the different model runs and P(MSA):P(nss-SO<sub>4</sub><sup>2-</sup>). The values are averaged over the MBL for model days 2–5 for the cloud-free runs and days 2–3 for cloudy runs.

**DMS oxidation**

R. von Glasow and P. J.  
Crutzen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Print Version

Interactive Discussion

© EGU 2003

## DMS oxidation

R. von Glasow and P. J. Crutzen



**Fig. 1.** Schematic of the DMS oxidation reactions in the gas phase as used in this study. Uncertain reactions are denoted with a question mark, dashed pathways are given only for reference and are not included in the model.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

⏴

⏵

Back

Close

Full Screen / Esc


Print Version

Interactive Discussion



## DMS oxidation

R. von Glasow and P. J. Crutzen



**Fig. 2.** Schematic of the DMS oxidation reactions in the aqueous phase as used in this study.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

DMS oxidation

R. von Glasow and P. J. Crutzen





Fig. 3. Schematic of proposed halogen - sulfur - climate links in the MBL.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

## DMS oxidation


R. von Glasow and P. J.  
Crutzen

**Fig. 4.** Temporal evolution in 50 m altitude of some sulfur species for the runs: “Luc” (black, solid line), “Luc, nh” (red, dotted line), “Yin” (green, dashed line), and “Yin, nh” (blue, dash-dotted line).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGU 2003

## DMS oxidation


R. von Glasow and P. J.  
Crutzen

**Fig. 5.** Temporal evolution in 50 m altitude of some sulfur species for the runs: “Kuk 0%” (black, solid line), “Kuk 0% nh” (red, dotted line), “Kuk 5%” (green, dashed line), and “Kuk 5%, nh” (blue, dash-dotted line).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGU 2003

## DMS oxidation

R. von Glasow and P. J.  
Crutzen

**Fig. 6.** Temporal evolution of the bromide deficit in 50 m height in the sea salt aerosol for the cloud-free scenarios “kuk, 5%” (solid line) and “tropics” (dashed line).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGU 2003