Atmos. Chem. Phys. Discuss., 5, 7929–7964, 2005 www.atmos-chem-phys.org/acpd/5/7929/ SRef-ID: 1680-7375/acpd/2005-5-7929 European Geosciences Union



ACPD

5, 7929-7964, 2005

Intercomparison exercise between radiative transfer models

F. Hendrick et al.



EGU

Intercomparison exercise between different radiative transfer models used for the interpretation of ground-based zenith-sky and multi-axis DOAS observations

F. Hendrick¹, M. Van Roozendael¹, A. Kylling^{2,*}, A. Petritoli³, A. Rozanov⁴, S. Sanghavi⁵, R. Schofield^{6,**}, C. von Friedeburg⁵, T. Wagner⁵, F. Wittrock⁴, D. Fonteyn¹, and M. De Mazière¹

¹Institut d'Aéronomie Spatiale de Belgique, Brussels, Belgium

²Norwegian Institute for Air Research, Kjeller, Norway

³Institute of Atmospheric Science and Climate, Bologna, Italy

⁴Institute of Environmental Physics, University of Bremen, Bremen, Germany

⁵Institute of Environmental Physics, University of Heidelberg, Heidelberg, Germany

⁶National Institute of Water and Atmospheric Research, Omakau, Central Otago, New Zealand

 $\ensuremath{\textcircled{O}}$ 2005 Author(s). This work is licensed under a Creative Commons License.

5, 7929-7964, 2005

Intercomparison exercise between radiative transfer models

F. Hendrick et al.

Title Page		
Abstract	Introduction	
Conclusions	References	
Tables	Figures	
•	►	
Back	Close	
Full Screen / Esc		
Print Version		
Interactive Discussion		

* now at: St. Olavs University Hospital, Trondheim, Norway ** now at: NOAA Aeronomy Laboratory, Boulder, Colorado, USA

Correspondence to: F. Hendrick (franch@oma.be)

Received: 30 June 2005 - Accepted: 23 August 2005 - Published: 2 September 2005

Abstract

We present the results of an intercomparison exercise between six different radiative transfer (RT) models carried out in the framework of QUILT, an EU funded project based on the exploitation of the Network for the Detection of Stratospheric Change (NDSC).

- ⁵ RT modeling is an important step in the interpretation of Differential Optical Absorption Spectroscopy (DOAS) observations. It allows the conversion of the slant column densities (SCDs) into vertical column densities (VCDs) using calculated air mass factors (AMFs). The originality of our study resides in comparing SCD simulations in multi-axis (MAX) geometry (trace gases: NO₂ and HCHO) and in taking into account the photo-
- ¹⁰ chemical enhancement for calculating SCDs of rapidly photolysing species (BrO, NO₂, and OCIO) in zenith-sky geometry. Concerning the MAX simulations, good agreement is observed between the different models with the calculated NO₂ and HCHO SCDs differing by no more than 5% in the elevation and solar zenith angles (SZA) ranges investigated (5°-20° and 35°-85°, respectively). The impacts of aerosol scattering,
- ¹⁵ ground albedo, and relative azimuth on MAX simulations have also been tested. Large discrepancies appear for the aerosol effect, suggesting differences between models in the treatment of the aerosol scattering. A better agreement is obtained in the case of the ground albedo and relative azimuth effects. In zenith-sky geometry, the different models agree generally well, especially below 90° SZA. At higher SZA, larger discrepancies are observed with relative difference values between 2% and 14% in some
- cases. All the initialization data and results have been made publicly available through the QUILT project web site (http://nadir.nilu.no/quilt/), enabling the testing of other RT codes designed for the calculation of SCDs/AMFs.

1. Introduction

²⁵ Since the middle of the 1970s, stratospheric ozone and several trace gases directly or indirectly involved in the ozone depletion like NO₂, BrO, and OCIO have been moni-

ACPD

5, 7929–7964, 2005

Intercomparison exercise between radiative transfer models

Title Page			
Abstract	Introduction		
Conclusions	References		
Tables	Figures		
Back	Close		
Full Scre	Full Screen / Esc		
Print Version			
Interactive Discussion			

tored from the ground using the Differential Optical Absorption Spectroscopy (DOAS)-technique (Platt, 1994). A significant part of this monitoring effort has been carried out through the framework of the Network for the Detection of Stratospheric Change (NDSC). The NDSC consists of about 20 sites combining complementary observations
 techniques distributed in five Primary Stations (Arctic, Alpine, Hawaii, New Zealand, and Antarctic) together with many globally distributed complementary sites. Network operations started in January 1991 and have provided a consistent, standardised set of long-term measurements of stratospheric and, more recently, tropospheric trace gases, particles, and physical parameters for detecting atmospheric change, validating

- space-borne sensors, and testing and improving multidimensional models of both the stratosphere and troposphere (further information at http://www.ndsc.ncep.noaa.gov/). Concerning the scattered sunlight DOAS instruments, they operated until recently almost only in zenith-sky geometry, thus probing mainly the altitude region corresponding to the stratosphere, especially at large solar zenith angles (SZAs). Over the last
- decade, new instruments pointing not only at zenith but also towards the horizon (off-axis geometry) have been developed (e.g., Heckel et al., 2005; Hönninger et al., 2004; Wagner et al., 2004; Wittrock et al., 2004). Pointing at an elevation angle close to the horizon provides sensitivity to the lowest absorption layers and therefore enables the measurement of tropospheric species. The Multi-AXis (MAX-) DOAS technique
 combines different elevation angles. Due to the variation of the light paths in the tropo-
- ²⁰ combines different elevation angles. Due to the variation of the light paths in the troposphere with the elevation angle, the different viewing directions have maximum sensitivity at different heights, thus providing some information on the vertical distribution of the absorber.

An important step in the interpretation of DOAS observations is the conversion of the slant column densities (SCDs) – which are the direct product of the DOAS analysis – into vertical column densities (VCDs) using calculated air mass factors (AMFs). The AMF represents the ratio of the effective optical path through the atmosphere to the vertical optical path and is given by the ratio SCD/VCD (Noxon et al., 1979; Solomon et al., 1987). In practice, AMFs are calculated using a radiative transfer (RT) model. In the

ACPD

5, 7929-7964, 2005

Intercomparison exercise between radiative transfer models



case of short lived species such as BrO, OCIO, and NO₂, the calculation of SCDs and therefore AMFs is complicated by the variation of the concentration of these species along a given light path due to the fast diurnal variation of these radicals coupled to the local SZA variation along the light path (Fish et al., 1995). The so-called photochemical
 ⁵ enhancement effect is taken into account by initializing RT models with a table containing the concentrations of the absorber for different heights and SZAs. This table is then interpolated to determine the trace gas concentration at the altitudes and local SZAs of the different scatter points considered along a given sun ray path. Such a concentration table is usually generated by running a stacked box photochemical model initialized with the output of a 3-D chemical transport model (CTM) corresponding to the day and

- Notice of the observed of the boundary interface (of his) corresponding to the day and location of interest. SCDs calculated by coupling a 3-D CTM to a photochemical box-/RT models interface can also be directly compared to observed SCDs. For example, Sinnhuber et al. (2002) compared ground-based zenith-sky UV-visible observations of BrO obtained at 11 sites with simulations from the 3-D CTM SLIMCAT (Chipperfield,
- 15 1999). The RT model used was based on the single scattering ray tracing scheme of Solomon et al. (1987). Recently, several groups have initiated AMF calculations in order to interpret MAX-DOAS measurements. Hönninger et al. (2004) have studied the behaviour of AMFs as a function of the solar zenith, elevation, and relative azimuth angles, ground albedo, and aerosol loading for several idealized trace gas profiles. For
- ²⁰ this purpose, they have used the RT model TRACY based on a Monte Carlo approach (von Friedeburg, 2003). This model has been also used by Wagner et al. (2004) to perform RT modeling for MAX-DOAS O_4 observations at different aerosol conditions. Sensitivity tests on O_4 AMFs were carried out by Wittrock et al. (2004) with SCIATRAN, a RT code based on a combined differential-integral approach involving the Picard iter-
- ative approximation (CDIPI) (Rozanov et al., 2000, 2001). In this work, SCIATRAN was also used to derive NO₂ profile information from MAX-DOAS observations. In Heckel et al. (2005), this model generated appropriate AMFs for the conversion of MAX-DOAS measurements of HCHO into VCDs. These examples show that RT modeling plays a central role in the interpretation of ground-based DOAS observations and that different

ACPD

5, 7929-7964, 2005

Intercomparison exercise between radiative transfer models

Title Page			
Abstract	Introduction		
Conclusions	References		
Tables	Figures		
•	•		
Back	Close		
Full Scr	Full Screen / Esc		
Print Version			
Interactive Discussion			

RT computation schemes are available for this purpose.

The QUILT (Quantification and Interpretation of Long-Term UV-Vis Observations of the Stratosphere) project is an EU funded project based on the exploitation of the NDSC and aimed at quantifying ozone loss and investigating its relation to active halo-

- ⁵ gen and nitrogen species using the existing ground-based, satellite and balloon borne UV-visible data as well as 3-D atmospheric modelling tools. One of the tasks of this project has been to test the consistency between the different RT models existing within the consortium and used to interpret the ground-based and satellite DOAS data. This task has been achieved through several SCD simulations comparison tests performed
- ¹⁰ using identical settings for all the models. Both ground zenith-sky (trace gases: BrO, NO₂, and OCIO) and MAX geometries (trace gases: NO₂ and HCHO) have been considered for these tests, the photochemical enhancement being taken into account only in zenith-sky geometry.
- Here we report on the results of this intercomparison exercise. It should be noted that
 our study does not address the issue of the absolute accuracy of the SCD calculations, the consistency between simulated and measured SCDs having already been tested in several papers (e.g., Sinnhuber et al., 2002; Tørnkvist et al., 2002). It can be considered to a certain extent as the continuation of Sarkissian et al. (1995) who calculated O₃ and NO₂ AMFs with different RT models in zenith-sky geometry but without taking
 into account the photochemical enhancement effect (use of a single profile for the initialization of the models). Our paper is divided into five parts. In Sect. 2, we describe the different RT models involved in the intercomparison exercise. The comparison tests in zenith-sky and MAX geometries are described in Sects. 3 and 4, respectively, and their corresponding results are discussed therein. Sects. 5 and 6 are dedicated to the
- ²⁵ impact of the aerosols and ground albedo on MAX simulations, respectively.

ACPD

5, 7929-7964, 2005

Intercomparison exercise between radiative transfer models

Title Page		
Abstract	Introduction	
Conclusions	References	
Tables	Figures	
I ◀	►I	
•	►	
Back	Close	
Full Screen / Esc		
Print Version		
Interactive Discussion		

2. Description of the RT models

The groups contributing RT calculations to the intercomparison exercise were the remote sensing groups at the Universities of Bremen (UBRE) and Heidelberg (UHEI), Norwegian Institute for Air Research (NILU), Institute of Atmospheric Science and Cli-

- mate (ISAC-CNR), National Institute of Water and Atmospheric Research (NIWA), and Institut d'Aéronomie Spatiale de Belgique (IASB). A summary of the characteristics of the models used by these groups is given in Table 1. From now on, each model will be referred to by the name of its corresponding group (in brackets here above). All the models include the possibility of taking into account the photochemical enhancement.
 The solution approaches to the RT equation (RTE) used in the different RT codes are as follows:
 - The Finite Difference method (UBRE model): uses the integro-differential equations obtained by the expansion of the RTE in a Fourier cosine series in azimuth. The application of vertical and angular discretizations produces a matrix equation for each Fourier component which can be solved by using a recursion method (Lenoble, 1985; Rozanov et al., 2000 and 2001).
 - The Discrete Ordinate method (NILU and IASB models): as in the Finite Difference method, the azimuth dependence of the radiation field is expressed as a Fourier cosine series in azimuth. However, the solution of the Fourier components is obtained in this case using a numerical quadrature scheme, allowing to replace the integrals by sums and thus reducing the RTE to the solution of a set of coupled linear first-order differential equations (Lenoble, 1985; Stamnes et al., 1988; Spurr, 2001).
 - The backward Monte Carlo method (UHEI model): a photon emerges from a detector in an arbitrary line of sight direction and is followed in the backward direction along the path towards the sun. The various events which may happen

ACPD

5, 7929–7964, 2005

Intercomparison exercise between radiative transfer models

F. Hendrick et al.

Title	Title Page		
Abstract	Introduction		
Conclusions	References		
Tables	Figures		
14			
•	►		
Back	Close		
Full Scr	Full Screen / Esc		
Print	Print Version		
Interactive Discussion			

EGU

25

15

20

to the photon at various heights in the atmosphere are defined by suitable probability distributions. Random numbers decide on the occurrence of events. At the location of the last scattering event prior to leaving the atmosphere, the impinging radiance is calculated and weighted with the value of the scattering phase function and with the attenuation of the complete path. A large number of such random photon paths will reproduce the light contributing to the simulated measurement (von Friedeburg, 2003; Lenoble, 1985).

 Single scattering ray tracing (ISAC and NIWA models) method: in zenith-sky geometry, it consists of computing the attenuation from the sun through discrete spherical atmospheric shells to "points" where the light beams are scattered into the detector (Solomon et al., 1987).

3. Zenith-sky simulations of BrO, NO₂, and OCIO SCDs including photochemical enhancement

3.1. Comparison test description

5

10

In this test, BrO, OCIO, and NO₂ SCDs have been calculated in zenith-sky geometry in single scattering (SS) and when possible, in multiple scattering (MS) modes, taking into account the photochemical enhancement effect. The settings imposed for all models are summarized in Table 2. Color plots of the diurnal variation tables (concentration of BrO, OCIO, and NO₂ as a function of the altitude and SZA) are shown in Figs. 1, 2, and 3, respectively. They are the output of the stacked box photochemical model PSCBOX (Errera and Fonteyn, 2001; Hendrick et al., 2000 and 2004) initialized with the 3-D-CTM SLIMCAT chemical and meteorological fields. Two scenarios have been considered: Harestua (60° N, Norway) at sunset in summer for NO₂ SCD calculations and Harestua at sunset under chlorine activated conditions for BrO and OCIO.
 All calculations included absorption by O₃ and in the case of BrO and OCIO, also the

ACPD	4	С	Ρ	D
------	---	---	---	---

5, 7929-7964, 2005

Intercomparison exercise between radiative transfer models

Title	Title Page		
Abstract	Introduction		
Conclusions	References		
Tables	Figures		
•	Þ		
Back	Close		
Full Sc	reen / Esc		
Print	Print Version		
Interactive	Interactive Discussion		

3.2. Results

The results of the zenith-sky simulations of BrO, NO₂, and OCIO SCDs are presented in Fig. 4. We first concentrate on the general behaviour of the simulated SCDs. Figure 4 shows that the impact of MS is more important for BrO than for NO₂, and for OCIO, it is mostly significant at large SZA (>92°). The different behaviours of BrO and NO₂ regarding MS effects can be mainly attributed to the different wavelengths used for both calculations (352 nm for BrO and 422 nm for NO₂). In the case of OCIO, the large impact of MS at SZA larger than ~92° is related to the photochemistry of this species combined with geometrical considerations. Figure 2 shows that OCIO is situated in the $\sim 17-22$ km altitude range and only appears at SZA>92°. At these SZAs, the absorption due to OCIO along a given line of sight occurs where the local SZA is the largest, i.e., near the vertical above the observing point. Given that, a significant part of the OCIO layer is likely to be in the earth's shadow region and therefore can only be probed by multiply scattered light. This explains the large impact of MS on OCIO SCDs in these conditions. The differences between the photochemical behaviours of BrO, NO₂, and OCIO are also manifest from Fig. 4. BrO SCDs display a sharp decrease above ~92° SZA (see Fig. 1), consistent with the rapid conversion of BrO into its night-time reservoirs (mostly BrCl in the present case) in the absence of sunlight. In contrast, the concentrations of OCIO and NO₂ increase with the SZA (see Figs. 2 20

and 3, respectively), which explains the persistence of relatively large SCD values at corresponding SZAs, especially for MS mode.

Concerning the comparison of simulated SCDs, the relative differences between the results of the different models and the IASB ones arbitrarily taken as reference appear ²⁵ in the lower plots of Fig. 4 and a summary of the maximum difference values obtained is presented in Table 3. In SS mode, good agreement is observed below 90° SZA between the NIWA, IASB, NILU, and UBRE models: for the three species, the relative difference between the NIWA, NILU, and UBRE models and the IASB one (reference

ACPD

5, 7929-7964, 2005

Intercomparison exercise between radiative transfer models



model) is smaller than 1%. In the case of the ISAC-CNR model, simulated SCDs are systematically larger and the relative difference rises up to 4% for BrO, 5% for NO₂ and 7.5% for OCIO. Above 90° SZA, agreement between the NIWA, IASB, UBRE, and NILU models is still very good with relative difference values smaller than 1.8% for BrO,

- ⁵ 1.3% for NO₂, and 2.1% for OCIO. A maximum spread between models of about 2% is a better agreement than the one obtained by Sarkissian et al. (1995) without taking into account the photochemical enhancement (from Sarkissian et al. (1995), the maximum spread values rise up to 13% for O₃ and 8% for NO₂ at 94° SZA). Concerning the ISAC-CNR simulations above 90° SZA, the discrepancies with the IASB
- ¹⁰ model are larger than the ones observed for the NIWA and NILU models: the relative differences are smaller than 4% for BrO and rise up to 7.1% for NO₂, and 14.3% for OCIO. Although these maximum relative difference values are consistent with those of Sarkissian et al. (1995), they suggest that the ISAC-CNR model differs from the other models in the way the optical thickness of each atmospheric shell is calculated.
- ¹⁵ Sarkissian et al. (1995) investigated the influence of the different methods used to interpolate the concentration and to calculate the geometrical path in individual shells. They found that the impact of these parameters increases with increasing SZA and can reach 4% above 90° SZA. Since the photochemical enhancement is included in the present test, we cannot neglect the impact of differences in interpolating the tables
- of photochemical species concentrations. This effect is consistent with the observed increase of the relative differences between models with increasing SZA since the step between the SZA values corresponding to the concentration tables is increasing at large SZA. Concerning the calculations in MS mode, agreement below 90° SZA between the NILU, IASB, UBRE, and UHEI models is good since the relative differences
- with the IASB models are smaller than 4.4%. Above 90° SZA, slightly larger discrepancies are generally observed as in SS mode and the maximum relative differences with the IASB model rise up to 6.6% for the NILU model (obtained for OCIO SCD calculations), 5.5% for the UHEI model (obtained for NO₂ SCD calculations), and 4.2% for the UBRE model (obtained for OCIO SCD calculations). It should be noted that UHEI

5, 7929-7964, 2005

Intercomparison exercise between radiative transfer models



OCIO data are absent from the comparison. This is caused by the fact that too many photons (around 10⁵) would have been required to calculate each SCD with sufficient accuracy and precision. In contrast to the NO₂ or BrO profile, the stratospheric OCIO is usually confined to a layer of only a few kilometres thickness, and the concentration ⁵ gradient is very high at its lower and upper ends (see Fig. 2). The Monte Carlo character of the UHEI model causes the scattering point altitude to vary around a given value. For a profile of the shape described, the SCD does react very sensitively to even small scattering altitude variation, i.e., a scattering location just below the profile yields a low SCD, while a slight shift upwards causes the photon to continue the flight
¹⁰ on a slant path within the profile, leading to a sharp increase of the SCD. The Monte Carlo approach induced a variation of the scattering altitude, which in the case of OCIO results in a strong variation of the SCD itself. This can only be overcome with a very large number of modelled photons, which causes the UHEI model to be inefficient for

this specific kind of scenario. With the expected increase in computational power, this

disadvantage is likely to become insignificant in a near future.

4. Multi-axis simulations of NO₂ and HCHO SCDs

4.1. Comparison test description

In contrast to the zenith-sky simulations, the photochemical enhancement has not been taken into account in the MAX simulations. The reason is that this effect becomes significant only during twilight, i.e. above 85° SZA, and we have limited our comparison in the 38°–85° SZA range because the sensitivity of the MAX-DOAS observations to the troposphere is largest during daytime, i.e. at small SZA. The NO₂ and HCHO profiles used for the initialization of the models are shown in Fig. 5. All calculations included absorption by O₃ and in the case of HCHO, also the absorption by NO₂. SCD simulations have been performed for 5°, 10°, and 20° of elevation above the horizon and for each elevation angle, four values of the azimuth angle of line of sight (AzLOS) have 5, 7929–7964, 2005

Intercomparison exercise between radiative transfer models



been considered: 30° , 60° , 90° , and 120° . The AzLOS is the relative azimuth angle between the pointing direction of the instrument and the sun azimuth. The other settings appear in Table 4. Concerning the HCHO SCD calculations, the layer thickness in the altitude range where the concentration of this species changes rapidly with the altitude (0–3 km) has been fixed to 0.2 km instead of the 1 km thickness used in the case of NO₂, reducing interpolation effects. Only models enabling calculations in MS mode have been involved in this test since MS is the only acceptable scattering mode in MAX geometry (Wittrock et al., 2004). These models are: IASB, NILU, UBRE and UHEI.

10 4.2. Results

Figure 6 shows the results of NO₂ and HCHO SCD simulations for 5°, 10°, and 20° of elevation and 90° of AzLOS. A summary of the maximum difference values between the different models and the IASB one (again taken arbitrarily as reference model) is also presented in Table 5. The agreement between IASB, NILU, UBRE and UHEI is
¹⁵ good: when comparing the results for both species over the whole SZA range and for all the elevation angles, the relative differences between the NILU, UBRE, and UHEI models and the IASB one are smaller than 5.1%. Concerning the UBRE model, it underestimates the IASB model results by maximum 4.5% for NO₂ and 5.1% for HCHO. The relative differences are also almost constant over the whole SZA range. This is in

- ²⁰ contrast with the relative differences between the UHEI and IASB models which show a SZA dependence, especially at low elevation angles: e.g., for NO₂ at 5° of elevation, the UHEI model overestimates the IASB one by 3% at 40° SZA and underestimates it by 3.6% at 85° SZA. However, with observed maximum relative difference of 3.8% for NO₂ and 4.5% for HCHO (see Table 5), the agreement between both models is good.
- As expected since they are both based on the UVspec/DISORT package, the NILU and the IASB models agree very well: for both NO₂ and HCHO, the relative difference between the results of both models is smaller than 1.3% over the entire SZA range and for all elevation angles.

5, 7929–7964, 2005

Intercomparison exercise between radiative transfer models



The impact of the relative azimuth has been tested through the calculation of NO_2 and HCHO SCDs for different AzLOS values (30°, 60°, 90°, and 120°). Wittrock et al. (2004) have shown in their RT studies on O₄ AMF in MAX geometry that the AzLOS effect can be very large, especially for AzLOS pointing towards the sun (impact on O_4 ⁵ AMF larger than 10% for AzLOS smaller than 90° compared to the 90° AzLOS case). The AzLOS effect on NO₂ SCDs is illustrated in Fig. 7 where the relative differences between the NO₂ SCDs at 30°, 60°, and 120° AzLOS and the SCDs at 90° AzLOS (reference) are plotted as a function of the SZA for 5° and 20° of elevation. As in Wittrock et al. (2004), the largest AzLOS effect is observed at small AzLOS and high elevation angle. However, its magnitude is significantly smaller than for O_4 : about 7% on NO₂ SCDs at 50° SZA for 30° AzLOS and 20° of elevation instead of 45% for O₄ in similar conditions. It is also observed from Fig. 7 that the consistency between the IASB, NILU, UBRE, and UHEI models with respect to the impact of the AzLOS is reasonably good. In the case of HCHO (not shown here), the AzLOS effect is smaller

by 1–2% than the one observed for NO_2 . 15

5. Impact of aerosol scattering on multi-axis simulations of NO₂ and HCHO **SCDs**

The aerosols have a strong impact on MAX-DOAS observations (Wagner et al., 2004; Wittrock et al., 2004; and Heckel et al., 2005). The most important effect of the aerosol extinction is a reduction of the visibility of the atmosphere and thus a limitation by 20 scattering of the light path of the lowest viewing directions, reducing the difference in tropospheric absorption path between the viewing directions. Given this effect and its impact on MAX-DOAS O₄ observations, Wagner et al. (2004) have also shown that O_4 observations can provide a new method for deriving information on atmospheric aerosols. 25

In the present study, MAX NO₂ and HCHO SCDs have been simulated with and without aerosol scattering in order to test the consistency between the different RT 5, 7929-7964, 2005

Intercomparison exercise between radiative transfer models



models regarding the impact of the aerosols in this geometry. The aerosol extinction coefficient profiles used for initializing the models are shown in Fig. 8. They have been constructed from the aerosol model of Shettle (1989) included in the IASB and NILU RT models. Profiles of aerosol absorption coefficient and asymmetry factor have been also imposed in this test. The other settings are the same as the ones described in Sect. 4.1, except that HCHO and NO₂ SCDs have been calculated only for 90° of AzLOS.

Figure 9 shows the impact of the aerosol scattering on simulated MAX NO_2 and HCHO SCDs. Large discrepancies are observed between the UBRE and UHEI models and the IASB and NILU ones regarding the relative differences between simulations with and without aerosol scattering. In the case of HCHO, the impact of the aerosol scattering decreases and the agreement between the different models becomes better when the elevation angle increases. Similar features are observed for NO_2 but to a lesser extent. Figure 9 also shows that the impact of the aerosol scattering is smaller on

- ¹⁵ HCHO than NO₂ SCDs: the relative differences between calculations with and without aerosol scattering are comprised for all models, elevation angles, and SZAs between -13% and +1.5% for HCHO and between -21% and +4% for NO₂. These discrepancies could not be resolved satisfactorily so far. Their origin is unlikely to be found in the aerosol settings since all models have been initialized with the same aerosol profiles
- (extinction and absorption coefficients and asymmetry factor). In order to progress further in our understanding of these persistent discrepancies, a thorough examination of individual aerosol routines and additional comparison tests appear to be needed, which is beyond the scope of the present intercomparison exercise. This issue, and more generally the impact of the aerosol scattering on MAX DOAS AMF simulations, will be
- addressed more thoroughly as part of a new exercise, currently led by the University of Heidelberg in the framework of the European Network of Excellence on Atmospheric Composition Change ACCENT.

ACPD

5, 7929-7964, 2005

Intercomparison exercise between radiative transfer models

Title Page			
Abstract	Introduction		
Conclusions	References		
Tables	Figures		
•	•		
Back	Close		
Full Scre	Full Screen / Esc		
Print Version			
Interactive Discussion			

6. Impact of ground albedo on multi-axis simulations of NO₂ and HCHO SCDs

Ground albedo has a significant impact on the radiative transfer close to the ground (Hönniger et al., 2004; Wittrock et al., 2004). It is particularly important to properly estimate this parameter for observational sites displaying large albedo changes depending

on the season or viewing direction. The main effect of an increase of the albedo is an increase of the number of scattering events in the layers close to the surface, resulting in longer absorption paths at these altitudes and therefore to higher absolute AMFs. However, the difference in AMF between horizon and zenith-sky viewing directions is reduced, due to the fact that the enhancement of the optical path with increasing albedo is the largest for zenith-sky observations (Wittrock et al., 2004).

In order to test the consistency between the different RT models regarding the impact of the ground albedo in MAX geometry, NO_2 and HCHO SCDs have been simulated with ground albedo values fixed to 0 and 0.9. The other model settings are the same as those described in Sect. 4.1. As for the test on the impact of aerosol scattering, simulations have been performed only for 90° AzLOS.

Figure 10 illustrates the ground albedo effect on simulated MAX HCHO and NO₂ SCDs. The UBRE, NILU, and IASB models show excellent consistency with similar HCHO and NO₂ SCD increases being observed with the three models. An increase of the ground albedo value from 0.0 to 0.9 leads to an increase of the HCHO SCD of about 20% and 55% at 80° SZA for 5° and 20° of elevation, respectively, whereas the NO₂ SCD increases by 5% and 12% in the same conditions. The fact that the HCHO SCDs are more sensitive to the ground albedo is expected since for the present simulations, HCHO, in contrast to NO₂, is mostly located in the lower part of the troposphere (between 0 and 5 km). The corresponding increase of the SCD values are systematically larger using the UHEI model: 32% and 84% for HCHO SCDs, and 12% and 35%

²⁵ cally larger using the UHEI model: 32% and 84% for HCHO SCDs, and 12% and 35% for NO₂ SCDs. This behaviour is likely to be attributed to the different concepts used in the Monte Carlo model as compared to the analytical codes and further tests would be needed to find out the detailed reasons of this.

ACPD

5, 7929-7964, 2005

Intercomparison exercise between radiative transfer models



7. Conclusions

In the present intercomparison exercise, we have tested the consistency between six RT models used for interpreting ground-based zenith-sky and MAX-DOAS observations in the QUILT EU project. In the context of this project based on the exploitation

of the NDSC, the comparison and optimization of these RT models is of central importance in order to provide accurate time-series of ground-based DOAS observations. This study represents a step forward with respect to previously published work in that it compares RT models in MAX geometry and takes into account the photochemical enhancement effect for calculating SCDs of rapidly photolysing species in zenith-sky
 geometry.

Comparisons of NO₂ and HCHO SCDs in MAX geometry and multiple scattering mode show good agreement between all involved models: the calculated NO₂ and HCHO SCDs differ generally by no more than 5% in the SZA and elevation angle ranges investigated (35° - 85° and 5° - 20° , respectively). The impacts of the relative az-

- ¹⁵ imuth, aerosol scattering, and ground albedo on NO₂ and HCHO SCDs have been also quantitatively determined. As for O₄ in the Wittrock et al. (2004) study, the azimuth effect is found to be the largest at small AzLOS and high elevation angles. However, this effect is much smaller for NO₂ and HCHO than for O₄ (maximum differences relative to 90° AzLOS in the 30°–120° AzLOS range and for 20° of elevation rising up to 7% and
- ²⁰ 5% for NO₂ and HCHO SCDs, respectively, instead of 45% for O₄). The models have also shown reasonably good consistency concerning this effect. This is in contrast to the aerosol scattering effect for which large discrepancies still persist. Since all codes have been initialized in a common way, this result suggests that significant differences exist between the models regarding the treatment of the aerosol scattering. This issue,
- and more generally the impact of the aerosol scattering on the MAX DOAS AMF simulations, will be addressed more thoroughly as part of a new exercise, currently led by the University of Heidelberg in the framework of the European Network of Excellence ACCENT. Concerning the impact of the ground albedo, very good agreement has been

ACPD

5, 7929–7964, 2005

Intercomparison exercise between radiative transfer models



observed between the IASB, NILU, and UBRE models whereas the albedo effect is significantly larger using the UHEI model, which is based on a Monte Carlo approach. It should be noted that the conclusions drawn here on the levels of agreement between models in the different comparison tests depend on the assumptions made for the vertical profiles. In the case of NO₂ and HCHO, very different vertical profiles are indeed possible and could lead to different levels of agreement.

The comparisons of zenith-sky BrO, NO₂, and OCIO SCDs calculated in both single and multiple scattering modes show good overall agreement to a level that is consistent and in some cases better than in the Sarkissian et al. (1995) work. In single scattering

5

- ¹⁰ mode, the relative difference between the models is smaller than 1% below 90° SZA. In the case of the ISAC-CNR model, relative differences rising up to 7.5% have been observed in this SZA range. Above 90° SZA, agreement between the IASB, NILU, UBRE, and NIWA models is still very good with a maximum spread value of about 2%. Larger discrepancies have been observed with the ISAC-CNR model, especially
- ¹⁵ in the OCIO SCD calculations (relative difference with the IASB model up to 14%). These discrepancies could be partly due to differences in calculating the density in individual atmospheric shells. In multiple scattering mode, the differences between the NILU, UHEI, and UBRE models and the IASB one are smaller than 4.4% below 90° SZA for the three species. Above 90° SZA, slightly larger discrepancies have been
- observed with a maximum relative difference with the IASB model of about 6.5%. As in MAX geometry, the assumptions made on the vertical profiles used for the zenith-sky simulations can have an impact on the agreement observed between the models. This is particularly true for OCIO, for which a small change in the vertical profiles could change the behaviour of the different models quite a bit.
- All the initialization data (profiles and cross sections sets) and results of this intercomparison exercise have been gathered in a "RT model validation package" enabling the testing of other RT codes aiming to the calculation of SCDs/AMFs. This validation package has been made publicly available through the QUILT project web site (http://nadir.nilu.no/quilt/).

5, 7929-7964, 2005

Intercomparison exercise between radiative transfer models

Title Page			
Abstract	Introduction		
Conclusions	References		
Tables	Figures		
•	•		
Back	Close		
Full Scr	Full Screen / Esc		
Print Version			
Interactive Discussion			

Acknowledgements. This research was financially supported by the European Commission (contract QUILT, EVK2-2000-00545) and the Belgian Federal Science Policy Office (contracts ESAC II EV/35/3A and MO/35/006 & 012). M. P. Chipperfield (University of Leeds) is acknowledged for providing us with the SLIMCAT data. We would like also to thank the coordinators of the QUILT EC project, B. Arlander and G. O. Braathen (NILU).

References

5

20

- Barbe, A., Marché, P., Secroun, C., and Jouve, P.: Measurements of tropospheric and stratospheric H₂CO by an infrared high resolution technique, Geophys. Res. Lett., 6(6), 463–465, 1979.
- ¹⁰ Burrows, J. P., Dehn, A., Deters, B., Himmelmann, S., Richter, A., Voigt, S., and Orphal, J.: Atmospheric remote-sensing reference data from GOME: Part 1. Temperature dependent absorption cross sections of NO₂ in the 231–794 nm range, J. Quant. Spectrosc. Radiat, Transfer, 60, 1025–1031, 1998.

Burrows, J. P., Richter, A., Dehn, A., Deters, B., Himmelmann, S., Voigt, S., and Orphal, J.: At-

mospheric remote-sensing reference data from GOME: Part 2. Temperature dependent absorption cross sections of O₃ in the 231–794 nm range, J. Quant. Spectrosc. Radiat. Transfer, 61, 509-517, 1999.

Cantrell, C. A., Davidson, J. A., McDaniel, A. H., Shetter, R. E., and Calvert, J. G.: Temperaturedependent formaldehyde cross sections in the near-ultraviolet region, J. Phys. Chem., 94, 3902–3908, 1990.

Chipperfield, M. P.: Multiannual simulations with a three-dimensional chemical transport model, J. Geophys. Res., 104(D1), 1781–1805, 1999.

Errera, Q. and Fonteyn, D.: Four-dimensional variational chemical assimilation of CRISTA stratospheric measurements, J. Geophys. Res., 106(D11), 12253–12265, 2001.

Ehhalt, D. H. and Tönnißen, A.: Hydrogen and carbon compounds in the stratosphere, Proceedings of the NATO Advanced Study Institute on Atmospheric Ozone: Its Variation and Human Influences, edited by: Aikin, A. C., Report No. FAA-EE-80-20, U.S. Federal Aviation Administration, 1979.

Fish, D. J., Jones, R. L., and Strong, E. K.: Midlatitude observations of the diurnal variation of stratospheric BrO, J. Geophys. Res., 100(D9), 18863–18871, 1995.

ACPD

5, 7929–7964, 2005

Intercomparison exercise between radiative transfer models

Title Page			
Abstract	Introduction		
Conclusions	References		
Tables Figures			
	►I		
•	►		
Back	Close		
Full Scre	Full Screen / Esc		
Print Version			
Interactive Discussion			

- Hendrick, F., Mueller, R., Sinnhuber, B.-M., Bruns, M., Burrows, J. P., Chipperfield, M. P., Fonteyn, D., Richter, A., Van Roozendael, M., and Wittrock, F.: Simulation of BrO Diurnal Variation and BrO Slant Columns: Intercomparison Exercise Between Three Model Packages, Proceedings of the 5th European Workshop on Stratospheric Ozone, Saint Jean de
- Luz, France, 27 Sept.–1 Oct. 1999, Air Pollution Research Report no. 73, European Commission – DG XII, Brussels, 2000.

Hendrick, F., Barret, B., Van Roozendael, M., Boesch, H., Butz, A., De Mazière, M., Goutail, F., Hermans, C., Lambert, J.-C., Pfeilsticker, K., and Pommereau, J.-P.: Retrieval of nitrogen dioxide stratospheric profiles from ground-based zenith-sky UV-visible observations: validational for the stratespheric profiles from ground-based zenith-sky UV-visible observations: validational for the stratespheric profiles from ground-based zenith-sky UV-visible observations: validational for the stratespheric profiles from ground-based zenith-sky UV-visible observations: validational for the stratespheric profiles from ground-based zenith-sky UV-visible observations: validational for the stratespheric profiles from ground-based zenith-sky UV-visible observations: validational for the stratespheric profiles from ground-based zenith-sky UV-visible observations: validational for the stratespheric profiles from ground-based zenith-sky UV-visible observations: validational for the stratespheric profiles from ground-based zenith-sky UV-visible observations: validational for the stratespheric profiles from ground-based zenith-sky UV-visible observations: validational for the stratespheric profiles from ground-based zenith-sky UV-visible observations: validational for the stratespheric profiles from ground-based zenith-sky UV-visible observations: validational for the stratespheric profiles from ground-based zenith-sky UV-visible observations: validational for the stratespheric profiles from ground-based zenith-sky UV-visible observations: validational for the stratespheric profiles from ground-based zenith-sky UV-visible observations: validational for the stratespheric profiles from ground-based zenith-sky UV-visible observations: validational for the stratespheric profiles from ground-based zenith-sky UV-visible observations: validational for the stratespheric profiles for the stratespheric profiles from ground-based zenith-sky UV-visible observations: validational for the stratespheric pr

tion of the technique through correlative comparisons, Atmos. Chem. Phys., 4, 2091–2106, 2004,

SRef-ID: 1680-7324/acp/2004-4-2091.

Heckel, A., Richter, A., Tarsu, T., Wittrock, F., Hak, C., Pundt, I., Junkermann, W., and Burrows, J. P.: MAX-DOAS measurements of formaldehyde in the Po-Valley, Atmos. Chem. Phys., 5, 909–918, 2005.

15

SRef-ID: 1680-7324/acp/2005-5-909.

- Hönninger, G., Friedeburg, C. V., and Platt, U.: Multi Axis Differential Optical Absorption Spectroscopy (MAX-DOAS), Atmos. Chem. Phys., 4, 231–254, 2004, SRef-ID: 1680-7324/acp/2004-4-231.
- ²⁰ Lenoble, J.: Radiative Transfer in Scattering and Absorbing Atmospheres: Standard computational procedures, A. Deepak Publishing, Hampton, Virginia USA, 1985.
 - Mayer, B. and Kylling, A.: Technical note: The libRadtran software package for radiative transfer calculations description and examples of use, Atmos. Chem. Phys., 5, 1855–1877, 2005, SRef-ID: 1680-7324/acp/2005-5-1855.
- Noxon, J. F., Whipple, E. C., and Hyde, R. S.: Stratospheric NO₂. 1. Observational method and behaviour at midlatitudes, J. Geophys. Res., 84, 5047–5076, 1979.
 - Petritoli, A., Ravegnani, F., Giovanelli, G., Bortoli, D., Bonafè, U., Kostadinov, I., and Oulanovsky, A.: Off-axis measurements of atmospheric trace gases by use of an airborne ultraviolet-visible spectrometer, Appl. Opt., 41(27), 5593–5599, 2002a.
- Petritoli, A., Giovanelli, G., Kostadinov, I., Ravegnani, F., Bortoli, D., Bonasoni, P., Evangelisti, F., Bonafè, U., and Calzolari, F.: Tropospheric and stratospheric NO₂ amount deduced by slant column measurements at Mt. Cimone station, Adv. Space Res., 29(11), 1691–1695, 2002b.

5, 7929-7964, 2005

Intercomparison exercise between radiative transfer models

Title Page		
Abstract	Introduction	
Conclusions	References	
Tables	Figures	
	►I	
•	►	
Back	Close	
Full Screen / Esc		
Print Version		
Interactive Discussion		

- Platt, U: Differential optical absorption spectroscopy (DOAS), in Air Monitoring by Spectroscopic Techniques, Chem. Anal. Ser., edited by: Sigrist, M. W., 127, 27–84, John Wiley, New York, 1994.
- Rozanov, A., Rozanov, V.-V., and Burrows, J.-P.: Combined differential-integral approach for the
- radiation field computation in a spherical shell atmosphere: Nonlimb geometry, J. Geophys. Res., 105, D18, 22937, 2000.
 - Rozanov, A., Rozanov, V.-V., and Burrows J.-P.: A numerical radiative transfer model for a spherical planetary atmosphere: combined differential-integral approach involving the Picard iterative approximation, J. Quant. Spectrosc. Radiat. Transfer, 69, 491, 2001.
- ¹⁰ Sarkissian, A., Roscoe, H. K., Fish, D., Van Roozendael, M., Gil, M., Chen, H. B., Wang, P., Pommereau, J.-P., and Lenoble, J.: Ozone and NO₂ air-mass factors for zenith-sky spectrometers: Intercomparison of calculations with different radiative transfer models, Geophys. Res. Lett., 22(9), 1113–1116, 1995.

Schofield, R.: The vertical distribution of atmospheric BrO from ground-based measurements, PhD thesis, University of Auckland, Auckland, 2003.

Schofield, R., Connor, B. J., Kreher, K., Johnston, P. V., and Rodgers, C. D.: The retrieval of profile and chemical information from ground-based UV-visible spectroscopic measurements, J. Quant. Spectrosc. Radiat. Transfer, 86, 115–131, 2004.

15

Shettle, E. P.: Models of aerosols, clouds, and precipitation for atmospheric propagation stud-

- ²⁰ ies, AGARD Conference Proceedings No. 454: Atmospheric propagation in the UV, visible, IR and mm-region and related system aspects, 1989.
 - Sinnhuber, B.-M., Arlander, D. W., Bovensmann, H., Burrows, J. P., Chipperfield, M. P., Enell, C.-F., Frieß, U., Hendrick, F., Johnston, P. V., Jones, R. L., Kreher, K., Mohamed Tahrin, N., Müller, R., Pfeilsticker, K., Platt, U., Pommereau, J.-P., Pundt, I., Richter, A., South,
- A. M., Tørnkvist, K.K., Van Roozendael, M., Wagner, T., and Wittrock, F.: Comparison of measurements and model calculations of stratospheric bromine monoxide, J. Geophys. Res., 107(D19), 4398, doi:10.1029/2001JD000940, 2002.
 - Solomon, S., Schmeltekopf, A. L., and Sanders, R. W.: On the Interpretation of Zenith sky Absorption Measurements, J. Geophys. Res., 92(D7), 8311–8319, 1987.
- Spurr, R. J. D.: Linearized Radiative Transfer Theory: A general discrete ordinate approach to the calculation of radiances and analytic weighting functions, with application to atmospheric remote sensing, PhD thesis, Technische Universiteit Eindhoven, 2001.

Stamnes, K., Tsay, S.-C., Wiscombe, W., and Jayaweera, K.: Numerically stable algorithm for

5, 7929-7964, 2005

Intercomparison exercise between radiative transfer models

Title Page		
Abstract	Introduction	
Conclusions	References	
Tables	Figures	
	►I	
•	•	
Back	Close	
Full Screen / Esc		
Print Version		
Interactive Discussion		

discrete-ordinate-method radiative transfer in multiple scattering and emitting layered media, Appl. Opt., 27, 2505–2509, 1988.

- Tørnkvist, K. K., Arlander, D. W., and Sinnhuber, B.-M.: Ground-based UV measurements of BrO and OCIO over Ny-Ålesund during Winter 1996 and 1997 and Andøya during Winter 1998/99, J. Atmos. Chem., 43(2), 75–106, 2002.
- ⁵ 1998/99, J. Atmos. Chem., 43(2), 75–106, 2002. von Friedeburg, C.: Derivation of Trace Gas Information combining Differential Optical Absorption Spectroscopy with Radiative Transfer Modelling, PhD thesis, University of Heidelberg, 2003.

Wagner, T., Dix, B., Friedburg, C. v., Frieß, U., Sanghavi, S., Sinreich, R., and Platt,

¹⁰ U.: MAX-DOAS O₄ measurements: A new technique to derive information on atmospheric aerosols-Principles and information content, J. Geophys. Res., 109, D22205, doi:10.1029/2004JD004904, 2004.

Wahner, A., Ravishankara, A. R., Sander, S. P., and Friedl, R.R.: Absorption cross-section of BrO between 312 and 385 nm at 298 and 223 K, Chem. Phys. Lett., 152, 507–512, 1988.

¹⁵ Wahner, A., Tyndall, G. Y., and Ravishankara, A. R.: Absorption cross-section for OCIO as a function of temperature in the wavelength range from 240–490 nm, J. Phys. Chem., 91, 2735–2738, 1987.

Wittrock, F., Oetjen, H., Richter, A., Fietkau, S., Medeke, T., Rozanov, A., and Burrows, J. P.: MAX-DOAS measurements of atmospheric trace gases in Ny-Ålesund – Radiative transfer

²⁰ studies and their application, Atmos. Chem. Phys., 4, 955–966, 2004, SRef-ID: 1680-7324/acp/2004-4-955.

ACPD

5, 7929–7964, 2005

Intercomparison exercise between radiative transfer models

Title Page			
Abstract	Introduction		
Conclusions	References		
Tables Figures			
I ∢ ►I			
•	•		
Back Close			
Full Screen / Esc			
Print Version			
Finit Version			
Interactive Discussion			

Table 1. Short description of the RT models involved in the intercomparison exercise. Note that both IASB and NILU models are based on the UVspec/DISORT package. However, they are not strictly identical since they have been adapted separately by each group in order to allow SCD/AMF calculations taking into account the photochemical enhancement.

Group	Model main features	Scattering mode(s)	Reference(s)
UBRE	SCIATRAN model: -Combining Differential-Integral approach using the Picard-Iterative approximation (CDIPI) -Treatment of MS in a full spherical geometry including refraction -Chemistry included (2-D array of profile variation with SZA) -Treatment for aerosol and cloud scattering, and ground albedo -Raman scattering included	SS+MS	Rozanov et al. (2000, 2001); Wittrock et al. (2004)
UHEI	TRACY model: -Backward Monte Carlo approach -Treatment of MS in a full spherical geometry including refraction -Chemistry included (2-D array of profile variation with SZA) -Treatments for aerosol and cloud scattering, and ground albedo -Raman scattering included	SS+MS (only MS in this study)	von Friedeburg (2003); Hönninger et al. (2004)
NILU and IASB	UVspec/DISORT package: -Discrete ordinate method -Chemistry included (2-D array of profile variation with SZA) -Treatment of MS and refraction in a pseudo-spherical geometry (direct beam only) -Treatment for aerosol and cloud scattering, and ground albedo	SS+MS	Mayer and Kylling (2005)
ISAC-CNR	AMEFCO model: -Single scattering model in a 2-D atmosphere (profile variation with SZA) -Treatment in full spherical geometry including refraction and aerosol scattering	SS	Petritoli et al. (2002a, b)
NIWA	 Single scattering model in a 2-D atmosphere (profile variation with SZA) Treatment in full spherical geometry including refraction and aerosol scattering 	SS	Schofield (2003); Schofield et al. (2004)

ACPD

5, 7929–7964, 2005

Intercomparison exercise between radiative transfer models



Table 2. Model settings for the comparison test in zenith-sky geometry. All the initialisation data are available on the QUILT web site (http://nadir.nilu.no/quilt/).

Altitude grid	0–120 km/1 km thick	radiative	transfer
Wavelength	352 nm (BrO)	moo	dels
	368 nm (OCIO)	E Hand	
	422 nm (NO ₂)	F. Hendr	ick et al.
Diurnal variation table (see Figs. 1, 2, and 3)	From PSCBOX model		
p, T profiles	From SLIMCAT 3-D-CTM; interpolation		
	using AFGL 1976 outside the SLIMCAT	Title	Page
	altitude range		
O ₃ profile	From PSCBOX model; interpolation	Abstract	Introduction
	using AFGL 1976 outside the SLIMCAT	Conclusions	References
	altitude range	Contractione	
NO_2 profile (for BrO and OCIO SCDs calculations)	From PSCBOX model; interpolation	Tables	Figures
	using AFGL 1976 outside the SLIMCAT		
MP	altitude range	14	
Mie scattering	Not included		
Retraction	Not included	•	•
Cross sections sets	BrO: Wahner et al. (1988)		
	OCIO: Wahner et al. (1987)	Back	Close
	NO ₂ : Burrows et al. (1998)		
	O ₃ : Burrows et al. (1999)	Full Scre	en / Esc
Ground albedo	0.20		
Scattering mode	SS+if possible MS	Print V	/ersion

Interactive Discussion

ACPD

5, 7929-7964, 2005

Intercomparison exercise between

5, 7929-7964, 2005

Intercomparison exercise between radiative transfer models

F. Hendrick et al.

Title Page				
Abstract	Introduction			
Conclusions	References			
Tables	Figures			
•	►			
Back	Close			
Full Screen / Esc				
Print Version				
Interactive Discussion				

Table 3. Maximum relative difference observed between the different RT models and the IASB one – arbitrarily taken as reference – for the comparison test in zenith-sky geometry. Note that the SS and MS models have been compared to the IASB model in SS and MS modes, respectively.

	BrO [%]	NO ₂ [%]	OCIO [%]
NILU SS	+1.0	+0.5	+2.1
ISAC SS	+4.1	+7.1	+14.3
NIWA SS	+1.5	+0.7	+1.6
UBRE SS	+1.8	+1.3	-1.4
NILU MS	+5.8	+2.7	+6.6
UHEI MS	-5.0	+5.5	_
UBRE MS	-1.0	-0.6	-4.2

5, 7929-7964, 2005

Intercomparison exercise between r

Title Page				
Abstract	Introduction			
Conclusions	References			
Tables	Tables Figures			
•	•			
■ Back	► Close			
 Back Full Scr 	Close een / Esc			
	Close een / Esc Version			

Table 4. Model settings for the comparison test in MAX geometry. All the initialisation data are available on the QUILT web site (http://nadir.nilu.no/quilt/).

			EXELCISE	DEIMEE
Altitude grid	NO_2 : 0–120 km/1 km thick		radiative	transfe
-	HCHO: 0–120 km/0.2 km thick between		mou	اماد
	0 and 3 km and 1 km thick above 3 km		mot	
Wavelength	356 nm (HCHO)		F Hendr	ick et al
	$422 \text{ nm} (NO_2)$		1. Hond	ion of an
p. T profiles	From SLIMCAT 3-D-CTM: interpolation using	-		
F) . F	AFGL 1976 outside the SLIMCAT altitude range			_
O ₂ profile	From PSCBOX model: interpolation using		Title	Page
- 3 F	AFGL 1976 outside the SLIMCAT altitude range		Abotropt	Introduct
NO ₂ profile (for HCHO SCDs calculations)	From PSCBOX model: interpolation using		ADSITACI	Introduc
	AFGL 1976 outside the SLIMCAT altitude range		Conclusions	Referen
HCHO and NO ₂ profiles (see Fig. 5)	HCHO: Barbe et al. (1979):			
······································	Ehhalt and Tönnißen (1979)		Tables	Figure
	NO ₂ : from PSCBOX model			
Mie scattering	Not included			
Refraction	Not included			
Cross sections sets	HCHO: Cantrell et al. (1990)		4	Þ
	NO _o : Burrows et al. (1998)	-		
	O_{c} : Burrows et al. (1999)		Back	Close
Ground albedo	0.20	-		
Elevation angle	5° 10° and 20°		Full Scre	en / Esc
Azimuth angle of line of sight	$30^{\circ} 60^{\circ} 90^{\circ}$ and 120°			
Scattering mode	MS		Print V	/ersion

5, 7929-7964, 2005

Intercomparison exercise between radiative transfer models

F. Hendrick et al.

Title Page				
Abstract Introduction				
Conclusions	References			
Tables	Figures			
•	• •			
Back	Back Close			
Full Screen / Esc				
Print Version				
Interactive Discussion				

Table 5. Maximum relative difference observed between the different RT models and the IASB one – arbitrarily taken as reference – for the comparison test in MAX geometry.

		5° of elevation [%]	10° of elevation [%]	20° of elevation [%]
	NO ₂	-0.2	+0.2	+0.3
NILU MS	HCHO	+1.0	+1.3	+1.1
	NO_2	-3.6	-3.2	-3.8
UHEI MS	HCHO	-3.0	-3.0	-4.5
	NO_2	-4.5	-3.7	-3.3
OBRE MS	HCHO	-5.0	-5.0	-5.1



Fig. 1. Color plot of the BrO profiles table used to initialize the models in the comparison test in zenith-sky geometry. The variation of the BrO profiles as a function of the SZA has been calculated for Harestua (60° N, Norway) at sunset under chlorine activated conditions.

models F. Hendrick et al. **Title Page** Abstract Introduction Conclusions References Tables Figures ÞI ► Back Close Full Screen / Esc **Print Version** Interactive Discussion



Intercomparison exercise between





radiative transfer models F. Hendrick et al. **Title Page** Abstract Introduction Conclusions References Tables Figures ∎∎ ÞI ► Back Close Full Screen / Esc **Print Version** Interactive Discussion



Fig. 3. Color plot of the NO_2 profiles table used to initialize the models in the comparison test in zenith-sky geometry. The variation of the NO_2 profiles as a function of the SZA has been calculated for Harestua (60° N, Norway) at sunset in summer.

EGU

Print Version

Interactive Discussion







5, 7929–7964, 2005





Fig. 5. NO_2 (left) and HCHO (right) profiles in concentration (upper plots) and VMR (lower plots) used to initialize the RT models in the comparison tests in MAX geometry.

Print Version

Full Screen / Esc

4

Back

►

Close

Interactive Discussion

5, 7929-7964, 2005

Intercomparison exercise between radiative transfer models

F. Hendrick et al.



EGU

Fig. 6. NO_2 and HCHO SCDs calculated in MS mode in the comparison test in MAX geometry. For each species, the upper plots correspond to the SCDs and the lower plots to the relative differences between the different models and the IASB one arbitrarily chosen as reference. Results for 90° AzLOS and for 5°, 10°, and 20° of elevation are plotted here.

SZA [°]

NO,

90° AzLOS - 10° elev

60 SZA [°]

нсно

90° AzLOS - 10° elev

60 SZA [°] 80

40

40 60 80

80

40 60 80 SZA [°]

40

90° AzLOS - 20° elev

IASB MS NILU MS

- UBRE MS UHEI MS

SZA [°]

60 SZA [°]

90° AzLOS - 20° elev

IASB MS NILU MS

UBRE MS UHEI MS

60 SZA [°]

SZA [°]

80

NILU MS-IASB MS

UBRE MS-IASB MS

UHEI MS-IASB MS

40

40 60 80

NILU MS-IASB MS

UBRE MS-IASB MS UHEI MS-IASB MS

80

40 60 80

40

90° AzLOS - 5° elev

SZA [°]

60 80

SZA [°]

90° AzLOS - 5° elev

60 80 SZA [°]

SZA [°]

40

SCD [x10¹⁶ molec/cm²]

Q^N 40 60 80

Relative difference [%]

-10 40

HCHO SCD [x10¹⁶ molec/cm²]

Relative difference [%]

-10 40 60 80

F





Fig. 7. Impact of the relative azimuth on the NO₂ SCDs calculated in MAX geometry. Here are plotted the relative differences between the NO₂ SCDs calculated at 30° (upper plots), 60° (middle plots), and 120° (lower plots) AzLOS and those calculated at 90° AzLOS (reference case). Left and right plots correspond to results for 5° and 20° of elevation, respectively. Note that the IASB blue lines are superimposed to the NILU red lines.

5, 7929-7964, 2005



Intercomparison exercise between radiative transfer models F. Hendrick et al. **Title Page** Abstract Introduction Conclusions References Tables Figures ∎∎ ÞI ► Back Close Full Screen / Esc **Print Version**

Fig. 8. Profiles of the aerosol extinction coefficient at 356 nm and 422 nm used for testing the impact of the aerosols on MAX simulations of HCHO and NO_2 SCDs, respectively.

Interactive Discussion









Fig. 10. Impact of the ground albedo on simulated MAX NO₂ and HCHO SCDs. For each species, the upper plots correspond to the SCDs and the lower plots to the relative differences between SCDs calculated with ground albedo values of 0.9 and 0.0 (reference: albedo=0.0). Results for 5°, 10°, and 20° of elevation and 90° AzLOS are plotted here.

5, 7929-7964, 2005 Intercomparison exercise between radiative transfer models F. Hendrick et al. **Title Page** Abstract Introduction Conclusions References Figures Tables ∎∎ ► 4 Back Close Full Screen / Esc **Print Version** Interactive Discussion

ACPD