

**Tropical analyses
compared to
long-duration balloon
data**

B. M. Knudsen et al.

Accuracy of analyzed temperatures, winds and trajectories in the Southern Hemisphere tropical and midlatitude stratosphere as compared to long-duration balloon flights

B. M. Knudsen¹, T. Christensen¹, A. Hertzog², A. Deme², F. Vial², and J.-P. Pommereau³

¹Danish Meteorological Institute, Copenhagen, Denmark

²Laboratoire de Météorologie Dynamique du Centre National de la Recherche Scientifique, France

³Service d'Aéronomie du Centre National de la Recherche Scientifique, France

Received: 10 May 2006 – Accepted: 13 June 2006 – Published: 7 August 2006

Correspondence to: B. M. Knudsen (bk@dmi.dk)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Eight super-pressure balloons floating at constant level between 50 and 80 hPa and three Infra-Red Montgolfier balloons of variable altitude (15 hPa daytime, 40–80 hPa night time) have been launched at 22° S from Brazil in February–May 2004 in the frame of the HIBISCUS project. The flights lasted for 7 to 79 days residing mainly in the tropics, but some of them passed the tropical barrier and went to southern mid-latitudes. Compared to the balloon measurements just above the tropical tropopause the ECMWF operational temperatures show a systematic cold bias of 0.9 K and the easterly zonal winds are too strong by 0.7 m/s. This bias in the zonal wind adds to the ECMWF trajectory errors, but they still are relatively small with e.g. about an error of 700 km after 5 days. The NCEP/NCAR reanalysis trajectory errors are substantially larger (1300 km after 5 days). In the southern midlatitudes the cold bias is the same, but the zonal wind bias is almost zero. The trajectories are generally more accurate than in the tropics, but for one balloon a lot of the calculated trajectories end up on the wrong side of the tropical barrier and this leads to large trajectory errors.

1 Introduction

The accuracy of analyzed temperatures in the tropical tropopause layer has been studied extensively, due to e.g. its influence on the stratospheric humidity. Simmons et al. (1999) found the operational ECMWF analyses from 1996–1998 to have a standard-level bias of the order of 0.5°C or less compared to radiosondes. The temperature minima were, however, substantially overestimated, partially due to a 20 hPa vertical resolution of the model back then.

From three long-duration super-pressure balloon flights launched at 0.1° N from Ecuador, Vial et al. (2001) studied the accuracy of both ECMWF temperatures and winds at around 60 hPa from late August to mid October 1998 in the equatorial region. They found a warm bias of the ECMWF temperatures of about 0.5 K compared with

ACPD

6, 7499–7518, 2006

Tropical analyses compared to long-duration balloon data

B. M. Knudsen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

long-duration balloons. The easterly zonal winds were too strong by 2.4 m/s, but this could be explained by the balloon Stokes drift due to a Rossby-gravity wave near the equator.

In the SPARC Intercomparison of Middle Atmosphere Climatologies (SPARC, 2002; Randel et al., 2004a) several analyses and reanalyses were studied. Noteworthy is a 1–3 K warm bias of the tropical tropopause for the (UK)MO, CPC, and NCEP analyses from 1992–1997. Randel et al. (2004b) found this still to be true for (UK)MO and NCEP in 2001–2002. Using GPS-derived temperatures Gobiet et al. (2005) found an ECMWF cold bias of 1–2 K at the tropical tropopause in 2003–2004.

While past results for reanalyses still could be valid depending on changes in the observing system, the operational analyses are also subject to continuous model developments and past results may thus not reflect the current status. In this paper we compare operational ECMWF analyses to long-duration balloon launched from Bauru (22° S, 49° W) in Brazil in February–May 2004 during the HIBISCUS campaign (J.-P. Pommereau et al., 2006¹). We compare temperatures, horizontal winds, and trajectories. We primarily analyze data in the region just above the tropical tropopause (50–80 hPa). Trajectories based on NCEP/NCAR reanalyses are also compared. A paper on the ERA-40 reanalyses compared to long-duration balloon flights back to 1988 will also be submitted to the HIBISCUS special issue (Christensen et al., 2006²).

¹Pommereau, J.-P., Garnier, A., Goutail, F., et al.: An overview of the HIBISCUS campaign, *Atm. Chem. Phys. Discuss.*, HIBISCUS special issue, in preparation, 2006.

²Christensen, T., Knudsen, B. M., Pommereau, J.-P., Letrenne, G., Hertzog, A., and Vial, F.: Validation of ECMWF ERA-40 tropical lower stratosphere temperatures and winds with long-duration balloon data, *Atmos. Chem. Phys. Discuss.*, HIBISCUS special issue, in preparation, 2006.

Tropical analyses compared to long-duration balloon data

B. M. Knudsen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2 Long-duration balloon flights

Eight super-pressure (BP) and three Infra-Red Montgolfier (MIR) long duration balloons have been flown from Brazil during the HIBISCUS campaign (Pommereau et al., this special issue). One of the balloons (BP2) failed and a second one (BP3) experienced some problems of transmission which make the data useless. The BPs are spherical constant volume and therefore constant density (isopycnic) balloons made of trilaminated polyester. They were of two sizes: 10 m diameter flying around 55 hPa (19 km) varying a little with the load and 8.5 m diameter at 75 hPa (18 km) that is at or a little above the cold point tropopause. Two of the 6 HIBISCUS BP flights stay in the tropics, while the 4 others drifted to the southern-hemisphere mid-latitudes.

The scientific payload on BP flights, called Rumba, carries a GPS for location (± 10 m) and wind (± 0.01 m/s), a pressure (± 0.6 hPa) and two temperature sensors. The data were sampled every 15' and transmitted by the ARGOS satellite data collection system. The temperature sensors are small thermistors (YSI microbeads), with an accuracy of 0.25 K. The sensors are mounted 180° apart on a 1-m boom, hanging 5 m below the gondola. The thermistors are heated by the sun during day and consequently the daytime temperature observations exhibit a warm bias. This bias has been corrected as in Hertzog et al. (2004). On the Hibiscus flights, two kinds of thermistors were used: $120 \mu\text{m}$ diameter and $240 \mu\text{m}$ diameter. As expected, the temperatures measured with the largest thermistors have a larger bias than those measured with the smallest one (Fig. 1). Night time temperatures measured with the largest sensor are also colder than those measured with the smallest one. This is due to the sensor radiative cooling, which scales as the square of the sensor diameter. With these two sizes, it is possible to roughly estimate the cold bias of the small thermistors night time temperatures to 0.1 K. This small bias is neglected hereinafter and the night time temperatures measured with the small sensors are used in this paper as a reference.

The MIR balloon (Pommereau and Hauchecorne, 1979) is a hot air balloon heated by solar radiation during day-time and infrared radiation from the Earth during night-time.

ACPD

6, 7499–7518, 2006

Tropical analyses compared to long-duration balloon data

B. M. Knudsen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Tropical analyses
compared to
long-duration balloon
data**

B. M. Knudsen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Therefore its altitude varies from around 15 hPa (27 km) during the day to 40–80 hPa (22–18 km) at night depending on the cloud cover, except during the first 3 days before the complete escape of helium used for the initial ascent, when the MIR could be as high as 4 hPa. The position and thus the wind are obtained from GPS just like for the BP balloons. Because of the higher altitude, only night time temperatures are meaningful, but will not be used in the present study limited to altitudes close to the tropopause.

Table 1 gives some information on the flights. Among the 6 BPs, one (BP1) was leaking and fell after 13 days. All others stayed in flight until the end of their batteries between 27 and 80 days depending on the energy consumption of the additional passenger payload flown. The three flights BP5, BP7, and BP8 are closest to the tropical tropopause and their average tropical temperatures are 198–199 K. Among the 3 MIR flown, 2 dropped after 7 and 9 days over the South Pacific Convergence Zone. The third flew for 39 days for one and a half circumnavigation of the earth between 20 and 10° S.

3 Analyses

The ECMWF operational analysis in 2004 are produced by a 4-D variational analysis (Rabier et al., 2000) and is used at 6 hourly resolution. ECMWF T511 fields were extracted at the 60 model levels (spacing ~ 1.4 km) in a $1.5^\circ \times 1.5^\circ$ latitude-longitude grid (from a T79 truncation) and interpolated linearly in between. The top level is 0.1 hPa (~ 60 km). The integration scheme is a 2nd order Runge-Kutta scheme with a time step of 30 min (BP) or 10 min (MIR). Such an integration should give rise to much smaller errors than other errors connected to trajectory calculations such as analyses or interpolation errors (Knudsen and Carver, 1994).

The 6-hourly NCEP/NCAR reanalyses (Kalnay et al., 1996; Kistler et al., 2000) were used in a T62 truncation with 28 levels in the vertical up to 10 hPa. Contrary to the ECMWF trajectory calculation the NCEP/NCAR trajectories use cubic spline interpolation in space and time and a fourth order Runge-Kutta scheme with a time step of half

an hour.

4 Temperatures and winds

In this section, we compare the ECMWF analysis with the observations gathered during the BP flights by the Rumba gondola. The Rumba temperature measurements on the BP balloons shows a warm bias during daytime, which has been corrected. Figure 2 shows the histogram of differences between ECMWF and observed temperature, zonal and meridional wind. The mean bias of ECMWF relative to the balloon measurements is -0.9 K and the standard deviation is 1.3 K independent of whether it is from day or night and tropics or mid-latitudes. This cold bias of the ECMWF temperatures is in agreement with comparisons to radiosondes at 100 hPa and started already in 1998 (Simmons, 2003). The bias is also seen in comparisons with GPS-derived temperatures (Gobiet et al., 2005). First results from the new ECMWF T799 model, which became operational on 1 February 2006, indicate that the cold bias is reduced (Krüger et al., 2006). Vial et al. (2001) found a warm bias of about 0.5 K around 1 September 1998, at 60 hPa (20 km). This does not disagree with the present results since the flights they studied were closer to the equator. Actually Gobiet et al. (2005), who used GPS-derived temperatures, also indicate a warm bias at 20 km close to the equator.

The zonal (meridional) velocity has a mean bias of -0.4 (0.0) m/s and a standard deviation of 3.1 (3.5) m/s. If the comparison is limited to the data collected in the tropics (north of 30° S), the bias of the zonal velocity is -0.7 m/s , indicating that the ECMWF tropical easterlies are too strong. A spectral analysis (not shown) has shown that much of the scatter in these comparisons is due to meso- and short-scale inertia-gravity waves, which the ECMWF analysis has difficulties in capturing. These waves are more frequent at the tropical tropopause than above.

Tropical analyses compared to long-duration balloon data

B. M. Knudsen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

5 Trajectories

To assess how accurate calculated trajectories are, trajectory calculations (Knudsen et al., 2001; Knudsen and Carver, 1994; Hertzog et al., 2004) were started every 2 h along the flight track. In each time step the trajectories calculated for the BP flights were forced to a pressure lying on the same isopycnic (constant density) surface as the balloon. Thereby the vertical motion of the trajectories is handled. The trajectories for the MIR flights were forced to balloon pressure. Figure 3 shows the flight track for the 79 day long BP4 flight, and every 12th of the calculated trajectories. Most calculated trajectories stay in the tropical reservoir except for a few. One of the trajectories even moves to the Antarctic, just like some of the other BP balloons did. This does not necessarily indicate that the analyzed tropical barrier is leakier than the real, since trajectory errors could bring the calculated trajectories to regions where transport out of the tropical reservoir does occur. In agreement with the previous section, this figure also shows that the major part of the wave perturbations seen on the BP4 trajectory is not caught by the ECMWF analyses.

The horizontal balloon velocity is a very good approximation of the horizontal air velocity (Vial et al., 2001). In order to mimic the balloon behaviour in the vertical, isopycnic trajectories were computed for BP, while MIR trajectories were obtained by forcing the pressure to the observed balloon pressure. With this method only horizontal trajectory errors can be addressed, but these are in fact the most important ones if the vertical transport is calculated with a state-of-the-art radiation code (Knudsen et al., 2001).

Special attention has been attributed to the 79 day long BP4 flight, which remained in the tropics. Figure 4 shows the average spherical distance between the calculated and observed trajectory as a function of trajectory duration. The standard error on the average is calculated with lag 2 h autocorrelations taken into account and are indicated by the shading. The errors for the NCEP/NCAR reanalyses are based on trajectories started every 12 h and are calculated under the assumption that the autocorrelations

Tropical analyses compared to long-duration balloon data

B. M. Knudsen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

are the same as for the ECMWF trajectories. The NCEP/NCAR reanalyses trajectory errors are larger than the ECMWF errors at the 68% confidence level except for durations of 5.25–8.25 days, even though the shadings overlap. The difference is only significant at the 95% confidence level for a duration of 12 h. The coarse vertical resolution of the NCEP/NCAR reanalyses could be responsible for the differences. Another contributing factor could be increased dynamical consistency of the ECMWF 4-D variational data assimilation, where departures of observations from a forecast in a 12 h time window are minimized.

Figure 4 also shows the trajectory errors for all other flights in 2004. The balloons leaving the tropics were only used until the zero wind line was crossed at about 27° S. The MIR data are only used after the time when the pressure permanently is below 10 hPa (except that the MIR mLidar does reach a minimum pressure of 9.2 hPa). The MIR balloons run at higher altitudes than the BP. In the Arctic trajectory errors increase with altitude (Knudsen et al., 2001) due to e.g. the decreasing number of radiosondes and their increasing errors. Judging from the trajectory errors on the longest MIR and BP flights there is no increase with height in the tropics. This result could, however, be influenced by the reduced occurrence of atmospheric waves, which ECMWF has difficulties in catching, along the higher-altitude MIR flights. The longest MIR and BP flights together outnumber the other flights by a factor of 3.7 for a trajectory duration of 5 days. This means that their trajectory errors should be used and the trajectory errors on the other flights are presumably consistent with them. The trajectory error in the zonal (meridional) direction is defined as the spherical distance at fixed balloon latitude (longitude). For the BP4 flight the trajectory errors are a factor of 2.4 larger in the zonal direction than in the meridional direction after 5 days. This is partially caused by the bias in the zonal wind.

In the Arctic the trajectory errors are approximately halved (Hertzog et al., 2004). This is primarily due to the bias in the meridional wind. This bias may be due to the reduced number of radiosondes and the break-down of geostrophy in the Tropics, which makes it difficult to transform satellite observations of temperature related quantities to

Tropical analyses compared to long-duration balloon dataB. M. Knudsen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

winds. In principle it should be possible to correct for such a bias and thereby reduce the trajectory errors. In the Arctic wind speeds are much larger, so trajectory errors relative to the trajectory length (Relative Spherical Distance = RSD) are much smaller.

The trajectory errors after 2 h are a good measure of the errors of the vector wind.

5 For the tropical part of the BP flights it can be transformed to an average vector wind difference of -0.86 m/s, which agrees quite well with the results found in section 4. For the MIR flights the result is -0.58 m/s, indicating a slightly reduced wind error at higher altitudes in the lower stratosphere.

10 Figure 5 shows the spherical distance between the calculated and observed trajectories for the extratropical part of the flights BP5-8 (i.e. after the time when the zonal wind turned zero). The calculated trajectories were cut-off at 79.5° S. In case of missing trajectories due to this cut-off uncertainties were not calculated. The last 12.75 days of the BP8 flight were removed to avoid trajectories to be cut-off. For BP6 and BP8 the trajectory errors are in fact close to the errors in the Arctic. However, the wind speed is lower so that the RSD is larger than in the Arctic. For the BP5 flight the trajectory errors are comparable to the tropical errors. Much larger errors occur for the BP7 flight. As seen on Fig. 6 this is due to a substantial part of the calculated trajectories staying in the tropics, whereas the balloon moves towards the South Pole. These trajectories of course have very large errors, but the average error is in fact not significantly larger than the errors of the other flights at 95% confidence due to very large error bars.

20 Trajectories close to a flow barrier like the tropical barrier can thus have large errors, because unavoidable small errors can push the calculated trajectory to the wrong side of the barrier or the barrier can be misplaced in the analyses. In this case the trigger was probably a situation with very low wind speeds as depicted in Fig. 7. New trajectories were started every 12 h from the red crosses. The balloon makes a loop, which is not caught by the ECMWF trajectories. The arrows show the 80 hPa wind field at the time when the balloon passes the cross in the loop. Situations of low wind speeds are critical for trajectory calculations since the errors on the analyzed wind do not go to zero as the wind speed does (Knudsen et al., 2001). Most of the trajectories started

**Tropical analyses
compared to
long-duration balloon
data**

B. M. Knudsen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

before or during the loop thus take a more northerly course and end up in the tropics, while most of the trajectories started afterwards move towards the South Pole. Most of the other balloon flights also encountered low wind speeds, but this did not lead to so large trajectory errors because it did not push such a large number of trajectories to the wrong side of a flow barrier.

6 Conclusions

The ECMWF operational temperatures in 2004 show a systematic cold bias of 0.9 K just above the tropical tropopause (50–80 hPa). The easterly zonal winds in this region are too strong by 0.7 m/s. In the southern extratropics the temperature bias is the same and the zonal wind has almost no bias. After 5 days the average trajectory error is about 500 km, when discarding one balloon flight with very large errors. This is true for both the tropics and southern extratropics and also for tropical flights at about 13–45 hPa. The absolute trajectory errors are not much larger than the errors in the Arctic, but there the wind speeds are much larger.

The ECMWF reanalysis, ERA-40, temperatures show a smaller systematic cold bias around 60 hPa of 0.5 K compared to long-duration balloon flights close to the equator in 1998 (Christensen et al., 2006²). For these flights the trajectory errors are about 1000 km after 5 days, which is possibly due to Rossby-gravity waves which ERA-40 is unable to catch. Flights at higher altitudes have trajectory errors of about 500 km after 5 days, which is more in line with the results for the ECMWF operational analyses shown in this paper.

Acknowledgements. ECMWF and NCAR/NCEP are thanked for supplying meteorological data. The balloons were operated by the Centre National d'Etudes Spatiales in cooperation with the Institute of Meteorology of the State University of Sao Paulo in Brazil, which are gratefully acknowledged. This work was done within the HIBISCUS and SCOUT-O3 projects funded by the European Commission.

Tropical analyses compared to long-duration balloon data

B. M. Knudsen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

References

- Gobiet, A., U. Foelsche, A. K. Steiner, M. Borsche, G. Kirchengast, and Wickert, J.: Climatological validation of stratospheric temperatures in ECMWF operational analyses with CHAMP radio occultation data, *Geophys. Res. Lett.*, 32, L12806, doi:10.1029/2005GL022617, 2005.
- 5 Hertzog, A., Basdevant, C., Vial, F., and Mechoso, C. R.: Some results on the accuracy of stratospheric analyses in the Northern Hemisphere inferred from long-duration balloon flights, *Q. J. R. Meteorol. Soc.*, 129, 607–626, 2004.
- Kalnay, E., Kanamitsu, M., Kistler, R., et al.: The NCEP/NCAR 40-year reanalysis project, *Bull. Am. Meteorol. Soc.*, 77, 437–471, 1996.
- 10 Kistler, R., Kalnay, E., Collins, W., et al.: The NCEP-NCAR 50-year reanalysis: Monthly means CD-ROM and documentation, *Bull. Am. Meteorol. Soc.*, 82, 247–267, 2000.
- Knudsen, B. M., Pommereau, J.-P., Garnier, A., Nunez-Pinharanda, M., Denis, L., Letrenne, G., Durand, M., and Rosen, J. M.: Comparison of stratospheric air parcel trajectories based on different meteorological analyses, *J. Geophys. Res.*, 106, 3415–3424, 2001.
- 15 Knudsen, B. M. and Carver, G. D.: Accuracy of the isentropic trajectories calculated for the EASOE campaign, *Geophys. Res. Lett.*, 21, 1199, 1994.
- Krüger, K., Tegtmeier, S., Schöllhammer, K., and Rex, M.: Multi-year calculations of transport processes in the TTL, 2nd SCOUT-O3 Annual Conference, Jülich, Germany 20–23 March, 2006.
- 20 Pommereau, J.-P. and Hauchecorne, A.: A new atmospheric vehicle: la Mongolfière infrarouge, *Adv. Space Res.*, 55, 1979.
- Rabier, F., Järvinen, H., Klinker, E., Mahfouf, J.-F., and Simmons, A.: The ECMWF operational implementation of four dimensional variation assimilation. I: Experimental results with simplified physics, *Q. J. R. Meteorol. Soc.*, 126, 1143–1170, 2000.
- 25 Randel, W. J., Wu, F., Oltmans, S. J., Rosenlof, K., and Nedoluha, G. E.: Interannual Changes of Stratospheric Water Vapor and Correlations with Tropical Tropopause Temperatures, *J. Atmos. Sci.*, 61, 2133–2148, 2004b.
- Randel, W., Udelhofen, P., Fleming, E., Geller, M., Gelman, M., Hamilton, K., Karoly, D., Orland, D., Pawson, S., Swinbank, R., Wu, F., Baldwin, M., Chanin, M.-L., Keckhut, P., Labitzke, K., Remsberg, E., Simmons, A., and Wu, D.: The SPARC Intercomparison of Middle-Atmosphere Climatologies, *J. Clim.*, 17, 986–1003, 2004a.
- 30 Simmons, A.: Representation of the stratosphere in ECMWF operations and ERA-40, Work-

ACPD

6, 7499–7518, 2006

Tropical analyses compared to long-duration balloon data

B. M. Knudsen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

shop on modelling and assimilation for the stratosphere and troposphere, Reading, 23–26 June, 2003.

5 Simmons, A. J., Untch, A., Jacob, C., Kållberg, P., and Undén, P.: Stratospheric water vapour and tropical tropopause temperatures in ECMWF analyses and multi-year simulations, Q. J. R. Meteorol. Soc., 125, 353–386, 1999

SPARC Intercomparison of Middle Atmosphere Climatologies. SPARC Rep. 3, 96 pp., 2002.

10 Vial, F., Hertzog, A., Mechoso, C. R., Basdevant, C., Cocquerez, P., Doubough, V., and Nouel, F.: A study of the dynamics of the equatorial lower stratosphere by the use of ultra long duration balloons. Part I: Planetary scales, J. Geophys. Res., 106, 22 725–22 744, 2001.

ACPD

6, 7499–7518, 2006

**Tropical analyses
compared to
long-duration balloon
data**

B. M. Knudsen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Tropical analyses compared to long-duration balloon data

B. M. Knudsen et al.

Table 1. Bauru 2004 long-duration balloon flight data: Start and end day and duration of the part used in the calculations is given. 10–90% quantiles of the pressures and latitude ranges are shown. Mean wind speed for the BP flights is given in last column. Values for the tropical part of BP5–8 are in parenthesis.

Flight	Start	End	Duration (days)	Pressure (hPa) (10–90% quantile)	Latitude (° S)	Mean wind (m/s)
BP1	6 Feb	19 Feb	12.42	54–55	20–24	10.5
BP4	29 Feb	19 May	79.00	52–54	12–24	11.2
BP5	6 March	14 April (10 April)	38.67 (4.25)	73–84(73–75)	21–79(27)	11.9(9.5)
BP6	7 March	2 May (17 March)	55.58 (10.17)	58–64(57–59)	21–76(28)	14.8(9.0)
BP7	10 March	6 April (18 March)	26.92 (8.58)	74–85(73–75)	20–77(27)	14.2(8.5)
BP8	11 March	5 April (18 March)	24.50 (6.58)	69–79(68–71)	18–(27)	14.8(10.0)
MIR SAOZ1	6 Feb	14 Feb	8.67	15–45	18–23	26.0
MIR SAOZ2	27 Feb	05 April	37.92	16–41	8–22	15.4
MIR mLidar	11 March	19 March	7.25	13–29	21–24	19.4

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 1. Temperature variations on super-pressure balloon flights (at 50–80 hPa) as a function of the sun zenith angle. Measured temperatures (noisy curves) exhibit a warm bias during day, which is larger for the largest thermistors (grey) than for the smallest ones (black). The temperatures measured with the largest thermistors also show a cold bias during night. The variations shown here are computed with respect to the night time temperatures measured with the smallest sensor (see text). (smooth curves) Correction applied to the raw measurements.

**Tropical analyses
compared to
long-duration balloon
data**

B. M. Knudsen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Tropical analyses
compared to
long-duration balloon
data**

B. M. Knudsen et al.




Fig. 2. Histograms of differences between ECMWF analyses and Rumba observations (analyzed – observed): (left) temperature, (center) zonal velocity, (right) meridional velocity.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Tropical analyses
compared to
long-duration balloon
data**

B. M. Knudsen et al.




Fig. 3. BP4 flight track (thick red: 1st revolution, blue: 2nd revolution) along with 24 hourly calculated trajectories up to 20 days duration (orange: 1st revolution, cyan: 2nd revolution).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Tropical analyses compared to long-duration balloon data

B. M. Knudsen et al.




Fig. 4. Mean tropical trajectory errors for the 2004 flights. Dark and light grey shading gives 68% confidence limits on the means for BP4 using ECMWF and NCEP/NCAR reanalysis, respectively. The BP flight level usually is 52–75 hPa, whereas the MIR flight level usually is 13–45 hPa.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Tropical analyses
compared to
long-duration balloon
data**

B. M. Knudsen et al.




Fig. 5. Mean extra-tropical trajectory errors for the 2004 flights. Thin dashed lines give 68% confidence limits on the means. For comparison the tropical flight BP4 is included.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Tropical analyses
compared to
long-duration balloon
data**

B. M. Knudsen et al.




Fig. 6. BP7 extratropical flight track (thick red line) along with 12 hourly calculated trajectories (thin orange lines).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Tropical analyses
compared to
long-duration balloon
data**

B. M. Knudsen et al.




Fig. 7. Close-up of the BP7 flight track when passing South America (thick red line) and 12 hourly calculated trajectories (thin orange lines) started from the red crosses. Horizontal winds are shown as arrows with a scale such that the north-south distance between the arrow centres equals 20 ms^{-1} .

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)