

Interactive comment on “Contribution of fungi to primary biogenic aerosols in the atmosphere: active discharge of spores, carbohydrates, and inorganic ions by Asco- and Basidiomycota” by W. Elbert et al.

W. Elbert et al.

Received and published: 17 November 2006

Corrigenda to Elbert et al., Atmos. Chem. Phys. Discuss., 6, 11317-11355, 2006 (to be implemented upon revision):

p. 11322, l. 22: diameters

p. 11322, l. 22: 10 μm (Ingold, 2001b).

p. 11325, l. 26: and 4-97 ng m^{-3}

p. 11326, l. 13: In contrast to mannitol, the average glucose

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

p. 11326, l. 24: of 18-220 ng m⁻³

p. 11330, l. 21: of ~60 m⁻² s⁻¹

p. 11345, Tab. 3, c. 4, l. 23: Balbina, Amazonas (2001)

p. 11345, Tab. 3, c. 4, l. 29: Balbina, Amazonas (2001)

p. 11346, Tab. 4, c. 2, l. 23: 12.6^a

p. 11346, Tab. 4, c. 2, l. 29: 76.7^a

p. 11346, Tab. 4, c. 5, l. 23: Balbina, Amazonas (2001)

p. 11346, Tab. 4, c. 5, l. 29: Balbina, Amazonas (2001)

p. 11346, Tab. 4, footnote: ^(a) average of campaign (19-28 July 2001)

p. 11352, Fig. 1, capt., l. 4: concentration range (instead of “range of diurnal average”)

p. 11352, Fig. 2, capt., l. 4: concentration range (instead of “range of diurnal average”)

Interactive comment on Atmos. Chem. Phys. Discuss., 6, 11317, 2006.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper