

**ISORROPIA II: a new
aerosol equilibrium
model**

C. Fountoukis and
A. Nenes

**ISORROPIA II: a computationally efficient
thermodynamic equilibrium model for K^+ -
 Ca^{2+} - Mg^{2+} - NH_4^+ - Na^+ - SO_4^{2-} - NO_3^- - Cl^- - H_2O
aerosols**

C. Fountoukis¹ and A. Nenes^{1,2}

¹School of Chemical and Biomolecular Engineering, Georgia Institute of Technology, 311
Ferst Drive, Atlanta, GA 30332-0100, USA

²School of Earth and Atmospheric Sciences, Georgia Institute of Technology, 311 Ferst Drive,
Atlanta, GA 30332-0100, USA

Received: 24 January 2007 – Accepted: 31 January 2007 – Published: 7 February 2007

Correspondence to: A. Nenes (nenes@eas.gatech.edu)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

This study presents ISORROPIA II, a thermodynamic equilibrium model for the $K^+ - Ca^{2+} - Mg^{2+} - NH_4^+ - Na^+ - SO_4^{2-} - NO_3^- - Cl^- - H_2O$ aerosol system. A comprehensive evaluation of its performance is conducted against the thermodynamic module SCAPE2 over a wide range of atmospherically relevant conditions. The two models overall agree well, to within 13% for aerosol water content and total PM mass, 16% for aerosol nitrate and 6% for aerosol chloride and ammonium. Largest discrepancies were found under conditions of low RH, primarily from differences in the treatment of water uptake and solid state composition. In terms of computational speed, ISORROPIA II was always found to be more than an order of magnitude faster than SCAPE2, with robust and rapid convergence under all conditions. The addition of crustal species does not slow down the thermodynamic calculations (compared to the older ISORROPIA code) because of optimizations in the activity coefficient calculation algorithm. Based on its computational rigor and performance, ISORROPIA II appears to be a highly attractive alternative for use in large scale air quality and atmospheric transport models.

1 Introduction

Aerosols, or airborne particulate matter (PM), play a central role in atmospheric processes. They reflect a significant amount of radiation back to space, thus enhancing the planetary albedo. Atmospheric aerosols can cause visibility impairment in highly polluted areas (Altshüller, 1984) through their interactions with electromagnetic radiation. By acting as cloud condensation nuclei (CCN), they affect cloud droplet number concentration, cloud droplet effective radius, and cloud reflectivity. Changes in aerosol concentrations also affect droplet size distribution affecting precipitation frequency and cloud lifetime. Aerosols can also be responsible for acid rain production, which can adversely affect soil and water quality, especially in environments rich in SO_2 and NO_x . It has been established that inhaled aerosol particles are detrimental to human health;

ACPD

7, 1893–1939, 2007

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ISORROPIA II: a new aerosol equilibrium modelC. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

as particles can contain toxic inorganic and organic substances that are often correlated with asthma and chronic obstructive pulmonary disease (Zanobetti et al., 2000; Ramachandran and Vincent, 1999; Brauer and Brook, 1997; Schwartz, 1994). Recent studies (Kaiser, 2005) have suggested that fine particles ($PM_{2.5}$) are more effective in causing respiratory illness and premature death than larger particles due to their ability to penetrate deeper into the lung. Dockery et al. (1993), who conducted a survey on six cities over 16 years, found that people living in areas with higher aerosol concentrations had a lifespan two years less than those living in cleaner areas. The knowledge of the chemical composition and physical state of atmospheric particles may be a critical link between toxicity and particulate matter.

Atmospheric aerosols are composed of water, inorganic salts, crustal material, organics and trace metals. A large part of the particle (dry) mass is inorganic (25–50%) with ammonium (NH_4^+), sodium (Na^+), sulfate (SO_4^{2-}), bisulfate (HSO_4^-), nitrate (NO_3^-) and chloride (Cl^-) being the most important contributors to the dry inorganic $PM_{2.5}$ (Heitzenberg, 1989). Crustal species, such as Ca^{2+} , K^+ , Mg^{2+} are a major component of dust, hence an important constituent of ambient particles. These inorganic species may be in the form of aqueous ions, or in the form of precipitated solids, in thermodynamic equilibrium with atmospheric gases and humidity.

To compute the composition and phase state of aerosols, every atmospheric gas/aerosol model requires knowledge of the thermodynamic equilibrium state because the driving force for mass transfer of species between gas and aerosol phases is the departure from equilibrium. Performing thermodynamic equilibrium calculations for aerosol systems is a demanding computational task (e.g., Nenes et al., 1999) because it involves the global optimization of a nonlinear convex problem, or, the solution of numerous nonlinear equations. The non-linearity is stronger under conditions of low relative humidity, where aqueous aerosol solutions are strongly non-ideal and require the use of activity coefficients (which increases computational cost). Therefore, efficient and accurate solution algorithms are highly needed.

Numerous aerosol inorganic equilibrium models have been developed over the

years, differing in the chemical species that they can treat, the solution method used and the type of input they can accept. Recent examples include AIM2 (Clegg and Pitzer, 1992; Clegg et al., 1992, 1994, 1995, 1998a, b; Wexler and Clegg, 2002), SCAPE2 (Kim et al., 1993a,b; Kim and Seinfeld, 1995; Meng et al., 1995), EQUISOLV II (Jacobson et al., 1996; Jacobson, 1999a,b), ISORROPIA (Nenes et al., 1998; Nenes et al., 1999), GFEMN (Ansari and Pandis, 1999a,b), EQSAM2 (Metzger et al., 2002a, b; Metzger et al., 2006), HETV (Makar et al., 2003), MESA (Zaveri et al., 2005a, b) and UHAERO (Amundson et al., 2006). AIM2 and GFEMN use the iterative Gibbs free energy minimization method to solve equilibrium problems for $\text{NH}_4^+/\text{Na}^+/\text{NO}_3^-/\text{SO}_4^{2-}/\text{Cl}^-$ systems. UHAERO uses the Gibbs free energy minimization method (using a primal-dual method, coupled to a Newton iteration method) and offers a choice of the Pitzer, Simonson, Clegg (PSC) mole fraction-based model (Pitzer and Simonson, 1986; Clegg and Pitzer, 1992; Clegg et al., 1992) or the ExUNIQUAC model (Thomsen and Rasmussen, 1999) for the activity coefficient calculations. These models have treat either the ammonium – nitrate – sulfate system or the ammonium – sodium – nitrate – chloride – sulfate system. Even though direct minimization usually gives the most accurate results, it comes at a large computational cost (Ansari and Pandis, 1999b). MESA simultaneously iterates for all solid-liquid equilibria using a pseudo-transient continuation method and solves for the $\text{NH}_4^+/\text{Na}^+/\text{NO}_3^-/\text{SO}_4^{2-}/\text{Cl}^-$ system of species with the addition of calcium cations. EQUISOLV II sequentially solves for the root of each equation in the system of equilibrium reactions and then iterates over the entire domain until convergence. This method is ideal for the incorporation of new reactions and species with minimal programming effort, but optimal computational performance is obtained only on vectorized computational platforms (Zhang et al., 2000). EQSAM2 uses a simplified parameterization of the nonideal solution properties based on the relationship between activity coefficients and the relative humidity (Metzger et al., 1999) to solve the $\text{NH}_4^+/\text{Na}^+/\text{NO}_3^-/\text{SO}_4^{2-}/\text{Cl}^-/\text{Ca}^{2+}/\text{Mg}^{2+}/\text{K}^+/\text{RCOO}^-$ system. SCAPE2 divides the problem into several subdomains based on major species that impact equilibrium partitioning and water uptake. By always attempting to solve for a liquid phase, SCAPE2

ISORROPIA II: a new aerosol equilibrium modelC. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

predicts the presence of water even at very low ambient relative humidities (<10%), and for this reason often does not predict the presence of a crystalline phase (solid precipitate).

Similar to SCAPE2, ISORROPIA determines the subsystem set of equilibrium equations and solves for the equilibrium state using the chemical potential method. The code solves analytically as many equations as possible through successive substitutions; remaining equilibrium reactions are solved numerically with bisection for stability. ISORROPIA also offers the choice of using precalculated tables of binary activity coefficients and water activities of pure salt solutions, which speeds up calculations. Another important feature of the model is the use of mutual deliquescence of multi-component salt particle solutions, which lowers the deliquescence point of the aerosol phase. All the previously described models solve the “forward” problem, in which total (gas + aerosol) concentrations of chemical species along with ambient temperature and relative humidity are used as input. Besides the forward problem, ISORROPIA also offers the ability to solve for the “reverse problem”, in which known quantities are the concentrations of sodium, ammonium, nitrate and sulfate in the aerosol phase together with the ambient temperature and relative humidity. Being computationally efficient, ISORROPIA has proved to be the model of choice for many three-dimensional air quality models (CMAQ, CAMx, etc.), chemical transport and general circulation models (Ansari and Pandis, 1999b; Yu et al., 2005). HETV is based on the algorithms of ISORROPIA for sulfate, nitrate and ammonium aerosol systems and is optimized for running on vectorized computational architectures.

An important drawback of the above codes (with the exception of SCAPE2, EQUISOLV II and EQSAM2) is lack of treatment of crustal species (Ca, K, Mg). It has been shown (Jacobson, 1999b; Moya et al., 2001b) that the inclusion of crustal species in a thermodynamic equilibrium framework can be important in modeling size/compositional distribution of inorganic aerosols. An attempt to treat crustals as “equivalent sodium” was met with modest success (Moya et al., 2001a) provided that Ca was a relatively small fraction of aerosol dry mass.

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

In the current study, we present a new model, “ISORROPIA II”, in which the thermodynamics of the crustal elements of calcium, potassium and magnesium have been added to the preexisting suite of components of the computationally efficient ISORROPIA. The new model, combining the computational advances with the explicit thermodynamics of crustals, is compared against the predictions of SCAPE2, both in terms of speciation and computational requirements.

2 Thermodynamic equilibrium calculations

2.1 Equilibrium constants

In a closed (aerosol-gas phase) system composed of i chemical species and j reactions at constant temperature T , and pressure P , the Gibbs free energy of the system, G , is minimum at chemical equilibrium. This condition is equivalent to stating that the system of reactants is equal to that of products, which can be written as (Nenes et al., 1998):

$$\prod_i a_i^{v_{ij}} = K_j(T) \quad (1)$$

where a_i is the activity of species i , v_{ij} is the stoichiometric coefficient of species i participating in the reaction j , and K_j is the equilibrium constant of the j -th reaction at temperature T ,

$$K_j(T) = \exp \left[-\frac{\sum_i v_{ij} \mu_i^0(T)}{RT} \right] \quad (2)$$

where R is the universal gas constant and $\mu_i^0(T)$ is the standard chemical potential of species i at 1 atm pressure and temperature T (in K).

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

K_j is a function of temperature according to the Van't Hoff equation:

$$\frac{d \ln K(T)}{dT} = \frac{\Delta H^{\circ}(T)}{RT^2} \quad (3)$$

where $\Delta H^{\circ}(T)$ is the enthalpy change of the reaction at temperature T (Denbigh, 1981). For a small temperature range, $\Delta H^{\circ}(T)$ can be approximated by:

$$\Delta H^{\circ}(T) = \Delta H^{\circ}(T_o) + \Delta c_p^{\circ}(T - T_o) \quad (4)$$

where $\Delta c_p^{\circ}(T)$ is the change of molar heat capacity of products minus reactants. By substituting Eq. (4) into Eq. (3) and integrating from a reference temperature T_o (typically at 298.15 K) to T , we obtain:

$$K(T) = K_o \exp \left[-\frac{\Delta H^{\circ}(T_o)}{RT_o} \left(\frac{T_o}{T} - 1 \right) - \frac{\Delta c_p^{\circ}}{R} \left(1 + \ln \left(\frac{T_o}{T} \right) - \frac{T_o}{T} \right) \right] \quad (5)$$

where K_o is the equilibrium constant at T_o .

2.2 Activity of species

The activity of species i , a_i , if an ideal gas, is equal to its partial pressure ($a_i = p_i$) (Seinfeld and Pandis, 1998). If i is an electrolyte dissolved in water, $a_i = \gamma_i^{(\nu_+ + \nu_-)} m_+^{\nu_+} m_-^{\nu_-}$, where γ_i is the activity coefficient of i in water, ν_+ and ν_- are the moles of cations and anions, respectively, released per mole of electrolyte and m_+ , m_- are their molalities, respectively. The activity of each solid phase species is assumed to be unity.

2.3 Activity coefficients

In ISORROPIA II, the multicomponent activity coefficients, γ_{12} for each ionic pair 1–2 are computed using Bromley's formula (Bromley, 1973),

$$\log \gamma_{12} = -A_{\gamma} \frac{z_1 z_2 I^{1/2}}{1 + I^{1/2}} + \frac{z_1 z_2}{z_1 + z_2} \left[\frac{F_1}{z_1} + \frac{F_2}{z_2} \right] \quad (6)$$

1899

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

where γ_{12} is the mean activity coefficient of cation 1 and anion 2 at 298.15 K, A_γ is the Debye-Hückel constant ($0.511 \text{ kg}^{0.5} \text{ mol}^{-0.5}$ at 298.15 K) and,

$$F_1 = Y_{21} \log \gamma_{12}^o + Y_{41} \log \gamma_{14}^o + Y_{61} \log \gamma_{16}^o + \dots + \frac{A_\gamma I^{1/2}}{1+I^{1/2}} [z_1 z_2 Y_{21} + z_1 z_4 Y_{41} + z_1 z_6 Y_{61} + \dots] \quad (7)$$

$$F_2 = X_{12} \log \gamma_{12}^o + X_{32} \log \gamma_{32}^o + X_{52} \log \gamma_{52}^o + \dots + \frac{A_\gamma I^{1/2}}{1+I^{1/2}} [z_1 z_2 X_{12} + z_3 z_2 X_{32} + z_5 z_2 X_{52} + \dots] \quad (8)$$

5 where odd subscripts refer to cations and even subscripts refer to anions, $Y_{21} = \left(\frac{z_1+z_2}{2}\right)^2 \frac{m_2}{I}$, $X_{12} = \left(\frac{z_1+z_2}{2}\right)^2 \frac{m_1}{I}$, z_i is the absolute charge of ionic species i , and γ_{ij}^o is the mean ionic activity coefficient of the binary pair $i-j$ (“binary” activity coefficient) computed at the ionic strength of the multicomponent solution, I , $I = \frac{1}{2} \sum_i m_i z_i^2$.

Following the recommendations of Kim et al. (1993), binary activity coefficients, γ_{12}^o ,
10 are calculated using the Kusik-Meissner relationship (Kusik and Meissner, 1978),

$$\log \gamma_{12}^o = z_1 z_2 \log \Gamma^o \quad (9)$$

where

$$\Gamma^o = [1 + B(1 + 0.1I)^q - B] \Gamma^* \quad (10)$$

$$B = 0.75 - 0.065q \quad (11)$$

$$15 \log \Gamma^* = \frac{-0.5107I^{1/2}}{1 + CI^{1/2}} \quad (12)$$

$$C = 1 + 0.055q \exp(-0.023I^3) \quad (13)$$

and q is a parameter specific for each binary pair (Table 4).

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The effect of temperature on multicomponent activity coefficients is described by (Meissner and Peppas, 1973):

$$\log \gamma_{ij}(T) = [1.125 - 0.005(T - 273.15)] \log \gamma_{ij}(T_0) - [0.125 - 0.005(T - 273.15)] A \quad (14)$$

where $\gamma_{ij}(T)$ is the multicomponent activity coefficient of the ionic pair i - j at temperature

$$T, \text{ and, } A = -\frac{0.41/T^{0.5}}{1+T^{0.5}} + 0.039/T^{0.92}.$$

2.4 Aerosol water content

During the calculation of aerosol water content, it is assumed that the ambient water vapor pressure is unaffected by the aerosol water uptake (Seinfeld and Pandis, 1998).

Therefore, if ambient relative humidity is known, phase equilibrium between gas and aerosol-phase (Seinfeld and Pandis, 1998) gives that the water activity, a_w , is equal to the ambient fractional relative humidity, RH, (i.e., expressed on a 0.0 to 1.0 scale):

$$a_w = \text{RH} \quad (15)$$

Instead of determining water content that satisfies the constraint of Eq. (15) from explicit calculations of water activity (which would require an iterative, hence computationally expensive procedure, (Stelson and Seinfeld, [1982]), the water uptake of aerosols is approximated through the ZSR relationship (Robinson and Stokes, 1965),

$$W = \sum_i \frac{M_i}{m_{oi}(a_w)} \quad (16)$$

where W is the mass concentration of aerosol water (kg m^{-3} air), M_i is the molar concentration of species i (mol m^{-3} air), and $m_{oi}(a_w)$ is the molality of an aqueous binary solution of the i -th electrolyte with the same a_w (i.e., relative humidity) as in the multicomponent solution. The water activities used and their corresponding sources are given in Table 6.

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2.5 Deliquescence relative humidity (DRH)

For each salt, there is a characteristic relative humidity, known as the deliquescence relative humidity (DRH), above which a phase transition from solid to saturated aqueous solution occurs. The DRH varies with temperature and for small T changes is given by (Wexler and Seinfeld, 1991):

$$\ln \frac{DRH(T)}{DRH(T_o)} = -\frac{M_w m_s L_s}{1000R} \left(\frac{1}{T} - \frac{1}{T_o} \right) \quad (17)$$

where M_w is the molar mass of water and m_s is the molality of the saturated solution at temperature T_o . L_s is the latent heat of fusion for the salt from a saturated solution given by $L_s = \Delta H_{cr} - \Delta H_{aq}$; ΔH_{cr} , ΔH_{aq} are the molar enthalpies of formation of the crystalline phase and the species in aqueous solution, respectively.

2.6 Mutual deliquescence relative humidity (MDRH)

In every multicomponent mixture there exists a characteristic relative humidity (known as mutual deliquescence relative humidity, MDRH, Wexler and Seinfeld, 1991), for which all salts are saturated with respect to their aqueous solution. The MDRH is a eutectic point so it is below the DRH of all the pure solids composing the system and is the minimum RH for which a stable aqueous phase exists (Wexler and Seinfeld, 1991). When $MDRH < RH < RH_{wet}$ (where RH_{wet} is the DRH of the salt with the lowest DRH in the mixture under consideration) the solution is said to be in the mutual deliquescence region (MDR, Nenes et al., 1998). Computing the aerosol composition in the MDR is a computationally demanding task (e.g., Potukuchi and Wexler, 1995a, b) which we seek to avoid. Given that the MDR corresponds usually to a narrow RH range, we approximate composition in a MDR using the Nenes et al. (1998) method, which involves computing the weighted average of a “dry” and “wet” solution:

$$W = (1 - c) W_{wet} \quad (18)$$

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

$$G = cG_{\text{dry}} + (1 - c)G_{\text{wet}} \quad (19)$$

$$S = cS_{\text{dry}} + (1 - c)S_{\text{wet}} \quad (20)$$

$$D = (1 - c)D_{\text{wet}} \quad (21)$$

The weighting factor, c , is given by (Nenes et al., 1998):

$$c = \frac{RH - RH_{\text{wet}}}{MDRH - RH_{\text{wet}}} \quad (22)$$

and G , S , D are the concentrations of gaseous, solid and dissolved species, respectively. The subscripts “wet” and “dry” in Eqs. (18)–(22) denote the two solutions which are weighted. In the above equations, we assume that gases and solids are linearly weighted according to their proximity to RH_{wet} and $MDRH$ (as expressed by c), while dissolved species are scaled to the amount of water. $MDRH$ points for the new mixtures in ISORROPIA II are shown in Table 5. Although Eqs. (18)–(22) are an approximation of the thermodynamic solution, they qualitatively follow the RH-dependence of speciation and conserve aerosol dry mass.

3 ISORROPIA II: species considered and general solution procedure

The system modeled by ISORROPIA II consists of the following potential components (species in bold are new in ISORROPIA II):

Gas phase: NH_3 , HNO_3 , HCl , H_2O

Liquid phase: NH_4^+ , Na^+ , H^+ , Cl^- , NO_3^- , SO_4^{2-} , $\text{HNO}_{3(\text{aq})}$, $\text{NH}_{3(\text{aq})}$, $\text{HCl}_{(\text{aq})}$, HSO_4^- , OH^- , H_2O , **Ca^{2+}** , **K^+** , **Mg^{2+}**

Solid phase: $(\text{NH}_4)_2\text{SO}_4$, NH_4HSO_4 , $(\text{NH}_4)_3\text{H}(\text{SO}_4)_2$, NH_4NO_3 , NH_4Cl , NaCl , NaNO_3 , NaHSO_4 , Na_2SO_4 , **CaSO_4** , **$\text{Ca}(\text{NO}_3)_2$** , **CaCl_2** , **K_2SO_4** , **KHSO_4** , **KNO_3** , **KCl** , **MgSO_4** , **$\text{Mg}(\text{NO}_3)_2$** , **MgCl_2**

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1 shows thermodynamic properties for all species considered. Table 2 displays all the equilibrium reactions used in ISORROPIA II along with values for their equilibrium constants. When the concentration of crustals (Ca, K, Mg) is zero, routines of ISORROPIA are used, which since its original release (Nenes et al., 1998) has been substantially improved for robustness, speed and expanded to solve a wider range of problems (updates can be obtained from <http://nenes.eas.gatech.edu/ISORROPIA>).

3.1 Solution procedure

The number of species and equilibrium reactions is determined by the relative abundance of each aerosol precursor (NH_3 , Na, Ca, K, Mg, HNO_3 , HCl, H_2SO_4) and the ambient relative humidity and temperature. The major species potentially present are determined from the value of the following ratios:

$$R_1 = \frac{[\text{NH}_4^+] + [\text{Ca}^{2+}] + [\text{K}^+] + [\text{Mg}^{2+}] + [\text{Na}^+]}{[\text{SO}_4^{-2}]}$$

$$R_2 = \frac{[\text{Ca}^{2+}] + [\text{K}^+] + [\text{Mg}^{2+}] + [\text{Na}^+]}{[\text{SO}_4^{-2}]}$$

$$R_3 = \frac{[\text{Ca}^{2+}] + [\text{K}^+] + [\text{Mg}^{2+}]}{[\text{SO}_4^{-2}]}$$

where $[X]$ denotes the concentration of an aerosol precursor X (mol m^{-3} of air). R_1 , R_2 and R_3 are termed “total sulfate ratio”, “crustals and sodium ratio” and “crustals ratio” respectively; based on their values, 5 aerosol composition regimes are defined, the possible species for which are listed in Table 3.

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ISORROPIA II: a new aerosol equilibrium modelC. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The DRH at $T_o=298.15\text{ K}$, the thermodynamic data for the L_s (Eq. 17) as well as the Kussik-Meissner activity coefficient parameters (Eq. 13) are shown in Table 4. Table 6 displays the polynomial fit parameters for computing the molalities of binary solutions as a function of water activity (obtained from Kim and Seinfeld, 1995; Ha and Chan, 1999; and Kelly and Wexler, 2006) for CaSO_4 , $\text{Ca}(\text{NO}_3)_2$, CaCl_2 , K_2SO_4 , KHSO_4 , KNO_3 , KCl , MgSO_4 , $\text{Mg}(\text{NO}_3)_2$ and MgCl_2 . For $(\text{NH}_4)_2\text{SO}_4$, NH_4HSO_4 , $(\text{NH}_4)_3\text{H}(\text{SO}_4)_2$, NH_4NO_3 , NH_4Cl , NaCl , NaNO_3 , NaHSO_4 and Na_2SO_4 , the water activity database was updated since the original release of ISORROPIA, using the output from the AIM model (<http://www.hpc1.uea.ac.uk/~e770/aim.html>).

As in ISORROPIA, ISORROPIA II solves two classes of problems:

1. Forward (or “closed”) problems, in which known quantities are T , RH and the total (gas + aerosol) concentrations of NH_3 , H_2SO_4 , Na, HCl, HNO_3 , Ca, K, and Mg.
2. Reverse (or “open”) problems, in which known quantities are T , RH and the precursor concentrations of NH_3 , H_2SO_4 , Na, HCl, HNO_3 , Ca, K, and Mg in the aerosol phase.

Below the MDRH of an aerosol mixture, the particle is a solid if the aerosol is following its deliquescence branch. However, when the RH over a wet particle is decreasing, the wet aerosol may not crystallize below the MDRH but instead remain in a metastable state, where it is composed of an aqueous supersaturated solution (Seinfeld and Pandis, 1998). ISORROPIA II can address both states (termed “stable” where salts precipitate once the aqueous phase becomes saturated with respect to them, and, “metastable”, if the aerosol is composed only of an aqueous phase which can be supersaturated with respect to dissolve salts).

Depending on the three sulfate ratios and the relative humidity, ISORROPIA II solves the appropriate set of equilibrium equations and together with mass conservation, electroneutrality, water activity equations and activity coefficient calculations, the final concentrations at thermodynamic equilibrium are obtained. Figure 1 illustrates a general description of the solution procedure.

3.2 Important issues

- When calculating species concentration, the stable state solution algorithm of ISORROPIA II starts with assuming a completely dry aerosol. As the ambient relative humidity increases (or decreases) ISORROPIA II dissolves each of the salts present (depending on their DRH) and calculates solid and ion concentrations and water uptake. The exact opposite methodology is adopted by other models. For example, SCAPE2 initially assumes that all salts present are completely dissolved and based on the ambient relative humidity and DRH of each salt calculates solid concentration if a precipitate is assumed to form. Differences in the “solution dynamics” may lead to differences in water content and speciation, especially at low RH, and are further analyzed in Sect. 4.
- ISORROPIA II uses the principle of “compositional invariance with RH cycling” to determine the aerosol composition at low RH (i.e., when the aerosol is solid). This is done because aerosol cycles RH many times in nature throughout its lifetime and the invariant solution will in general represent its composition more accurately in the atmosphere. Compositional invariance is applied when the aerosol contains volatile anions, sulfate and non-volatile univalent cations (Na, K). In such cases, Na and K preferentially associate with SO_4 to form Na_2SO_4 and K_2SO_4 before they are bound with NO_3 and Cl to form NaNO_3 , KNO_3 , NaCl, and KCl. Other models may not adopt this approach and may lead to differences in predicted water uptake, especially at low RH. For example, ISORROPIA II predicts that potassium will preferentially neutralize sulfate to form K_2SO_4 . Then excess potassium reacts with available HNO_3 and HCl to form KNO_3 and KCl. Therefore, in the above example ISORROPIA II assumes that potassium mainly binds with sulfate since sulfate is less volatile than nitrate or chloride when exposed to RH cycling, thus more likely to stay in the aerosol phase and form K_2SO_4 .

ACPD

7, 1893–1939, 2007

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3.3 Simplifications and assumptions in ISORROPIA II

Numerous simplifying assumptions are taken to increase computational speed and numerical stability without compromising rigor substantially. These are:

- Sulfuric acid, sodium and crustals have a very low vapor pressure and can safely be assumed that they exclusively reside in the aerosol phase.
- The first dissociation of sulfuric acid ($\text{H}_2\text{SO}_{4(\text{aq})} \rightarrow \text{H}^+ + \text{HSO}_4^-$) is assumed to be complete and not considered in the equilibrium calculations.
- For a wide range of ionic strengths (0–30 M), typical of ambient aerosols, the solubility product of magnesium sulfate (MgSO_4) was found to be always less than its equilibrium constant. Therefore, ISORROPIA II assumes MgSO_4 is always deliquesced when an aqueous phase is present, avoiding any computations for precipitating MgSO_4 out of solution.
- Calcium sulfate (CaSO_4) is assumed completely insoluble.
- For sulfate rich cases ($R_1 < 2$) $\text{NH}_{3(\text{g})}$, $\text{NO}_{3(\text{aq})}^-$ and $\text{Cl}_{(\text{aq})}^-$ are assumed minor species that do not significantly perturb the equilibrium through the $\text{NH}_{3(\text{g})} + \text{H}_2\text{O}_{(\text{aq})} \leftrightarrow \text{NH}_{4(\text{aq})}^+ + \text{OH}_{(\text{aq})}^-$, $\text{HNO}_{3(\text{g})} \leftrightarrow \text{H}_{(\text{aq})}^+ + \text{NO}_{3(\text{aq})}^-$ and $\text{HCl}_{(\text{g})} \leftrightarrow \text{H}_{(\text{aq})}^+ + \text{Cl}_{(\text{aq})}^-$ reactions, respectively. The code solves the appropriate set of equilibrium reactions (for the major species) and then the three gases ($\text{NH}_{3(\text{g})}$, $\text{HNO}_{3(\text{g})}$, $\text{HCl}_{(\text{g})}$) are subsequently dissolved through the equilibria described above. The same is assumed for the dissolved undissociated ammonia, nitric and hydrochloric acid in the aqueous phase ($\text{NH}_{3(\text{aq})}$, $\text{HNO}_{3(\text{aq})}$, $\text{HCl}_{(\text{aq})}$).
- For sulfate poor cases ($R_1 > 2$) bisulfate ion (HSO_4^-) is considered a minor species from the reaction $\text{HSO}_{4(\text{aq})}^- \leftrightarrow \text{H}_{(\text{aq})}^+ + \text{SO}_{4(\text{aq})}^{2-}$ (see Table 3).

ACPD

7, 1893–1939, 2007

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- 5
- MDRH points for multicomponent mixtures containing crustals are not known; They are approximated instead with data for known mixtures with as similar as possible composition (Table 5). For example, the MDRH point for a $(\text{NH}_4)_2\text{SO}_4$ – NH_4NO_3 – NH_4Cl – Na_2SO_4 – K_2SO_4 – MgSO_4 mixture is (not known and) assumed to be the same as for the $(\text{NH}_4)_2\text{SO}_4$ – NH_4NO_3 – NH_4Cl – Na_2SO_4 mixture. The absence of crustals in the consideration of the MDRH points of those mixtures is expected to introduce small underprediction of water, since *i*) both potassium and magnesium have similar deliquescence properties with sodium (Moya et al., 2001a), and, *ii*) highly insoluble salts (i.e., CaSO_4) do not significantly impact water activity, hence do not significantly contribute to DRH depression.

10

 - OH^- is assumed a minor species.
 - When crustals are in excess compared to all the anions, ISORROPIA II assumes that the solution is close to neutral ($\text{pH} \approx 7$). This is consistent with a presence of excess carbonate in the aerosol phase, which has a pK_a of ~ 6.4 (Meng et al., 1995).

15

 - The DRH of NH_4NO_3 is strongly dependent on temperature. Under low temperature conditions ($T < 270 \text{ K}$), this changes the order (starting from low to high RH) with which salts deliquesce (Fig. 2). For these cases the DRH of NH_4NO_3 in ISORROPIA II is assumed to not “cross over” the DRH of the other salts present in the solution, especially since thermodynamic data for supercooled NH_4NO_3 solutions are not known. The same is assumed for NH_4Cl and NaNO_3 which exhibit similar behavior with NH_4NO_3 (Fig. 2).

20

 - γ_{OH^-} and γ_{H^+} are assumed equal to unity, as the activity coefficient routines cannot explicitly calculate them.

ISORROPIA II: a new aerosol equilibrium modelC. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3.4 ISORROPIA II: new features

The main improvements to the original ISORROPIA release (Nenes et al., 1998) which are included in ISORROPIA II (and in the latest release of ISORROPIA version 1.7, <http://nenes.eas.gatech.edu/ISORROPIA>) are:

- Gas/liquid/solid partitioning has been extended to include crustal elements which resulted in 10 more salts in the solid phase and 3 more ions in the aqueous phase (Table 3).
- In addition to a thermodynamically stable state the aerosol can also be in a metastable state where no precipitate is formed (always an aqueous solution).
- The water activity database has been updated, using the output from the AIM model (<http://www.hpc1.uea.ac.uk/~e770/aim.html>).
- Temperature dependency of the activity coefficients is included. This has been done for both pre-calculated tables and online calculations of activity coefficients.
- The MDRH points for all the systems considered have been calculated using the GFEMN model of Ansari and Pandis (1999b).
- The activity coefficient calculation algorithm has been optimized to increase computational speed and avoid numerical errors.
- The tabulated Kusk-Meissner binary activity coefficient data have been recomputed through the online calculations for the midpoint of each ionic strength interval.
- A new subroutine has been added to provide the user with the option to “force” ISORROPIA II to conserve mass up to machine precision.

ACPD

7, 1893–1939, 2007

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

4 Evaluation of ISORROPIA II

ISORROPIA II is evaluated against the predictions of SCAPE2 for a wide range of conditions characteristic of urban, remote continental, non-urban continental and marine aerosol (Heitzenberg, 1989; Fitzgerald, 1991; Ansari and Pandis, 1999a). For urban and non-urban continental aerosol, sulfates, nitrates and ammonium are usually dominant inorganic species. Sodium and chloride often compose the majority of the marine particulate matter (usually with some crustals and sulfates present). This classification is mainly qualitative, as mixing between aerosol types often occurs in the atmosphere.

Table 7 lists the 16 different sets of precursor concentrations that were used in the intercomparison study. Sulfuric acid concentrations range between 1.0–5.7 $\mu\text{g m}^{-3}$ for marine and non-urban continental and 10.0–15.0 $\mu\text{g m}^{-3}$ for urban and remote continental aerosol. For the 16 cases considered, conditions 3, 4, 15 and 16 are sulfate-rich ($R_1 < 1$ or $1 < R_1 < 2$), conditions 1, 2, 13 and 14 represent sulfate near-neutral ($R_1 \approx 2$) aerosol and cases 5–12 are sulfate-poor ($R_1 > 2$), (Table 7). For each set of precursor concentrations, composition at thermodynamic equilibrium was calculated for 11 different RHs ranging from 10–98%; temperature was kept fixed at 298.15 K. In the evaluation study both the thermodynamically stable and metastable state solutions of ISORROPIA II are computed.

For the intercomparison study we calculate the normalized mean error (NME) defined as

$$\text{NME} = \frac{\sum_i^n |I_i - S_i|}{\sum_i^n S_i},$$

where I_i represents predictions of ISORROPIA II for case i , S_i predictions of SCAPE2 and n is the total number of cases considered.

Finally we compare the CPU time requirements between SCAPE2 and ISORROPIA II, stable and metastable solution of ISORROPIA II, as well as between ISORROPIA II and ISORROPIA for all the simulation conditions of Table 7.

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

4.1 Overall assessment of ISORROPIA II vs. SCAPE2

For all the simulations performed in this work, the water activity database of SCAPE2 was replaced with the one used in ISORROPIA II (reflecting the most updated water activity database). Activities of aqueous species in SCAPE2 were computed using Bromley's formula for multicomponent activity coefficients and the Kussik-Meissner method for binary coefficients.

In Fig. 3 we compare predictions of aerosol water, nitrate, chloride, ammonium, total PM and hydrogen concentrations between ISORROPIA II (stable solution, forward problem solved), and SCAPE2 for the conditions specified in Table 7. Both models predict similar amount of aerosol water content (Fig. 3a) with a normalized mean error of 13.5%. Most of this discrepancy is found in the low RH regimes ($RH < 60\%$) where SCAPE2 predicts higher water concentration compared to ISORROPIA II. This discrepancy is attributed to a) non-convergence of SCAPE2, which is corroborated by the large CPU time required for obtaining a solution (see Table 9), and, b) errors in the calculations of activity coefficients (both binary and multicomponent). At low RH (i.e., low liquid water content), the aqueous solution is highly non-ideal (hence the solution highly non-linear), consequently small changes in activity coefficients may result in large changes in the dissolved species concentrations and the predictions of liquid water content. A few cases exist (for $RH > 65\%$) for which ISORROPIA II predicts less aerosol water than SCAPE2 (Fig. 3a); this originates from differences in aerosol nitrate which then affects water uptake. For a few marine cases, SCAPE2 predicts negligible water due to non-convergence (Fig. 3a).

In Fig. 3b, total aerosol nitrate concentrations are compared for all the input conditions of Table 7. Overall, the agreement is very good with a mean error of 16.5%. ISORROPIA II predicts non-negligible amount of nitrate for some urban cases while SCAPE2 does not. For a few non-urban continental cases ISORROPIA II underpredicts aerosol nitrate compared to SCAPE2. The sources of these discrepancies are further investigated through specific examples in Sect. 4.2.

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Aerosol chloride concentration predictions are shown in Fig. 3c where both models show similar results (NME=6.5%) with small discrepancies for a few marine cases (due to non-convergence of SCAPE2 solution) in which chloride exists in significant amount due to significant presence of sea salt particles.

For aerosol ammonium predictions (Fig. 3d), no substantial differences between the two models were found (NME=2.1%). Discrepancies were primarily found in some non-urban continental cases which represent a sulfate-poor, ammonium-rich environment and are further analyzed in Sect. 4.2. Even though a few differences exist in the predicted concentrations of semi-volatile species, the total PM composition (Fig. 3e) shows very good agreement (NME=13.0%). The worst agreement between the two models was seen for H^+ predictions (Fig. 3f) with the normalized mean error significantly higher than for any other component (NME=64%). The discrepancy occurs at low RH (as it scales with water content).

SCAPE2 predictions are also compared against the metastable state solution of ISORROPIA II (Fig. 4). Table 8 shows normalized mean errors between ISORROPIA II (both stable and metastable solutions) and SCAPE2 for the simulations of Table 7. As can be seen in Figs. 3, 4 and Table 8, the stable state predictions of ISORROPIA II are closer to SCAPE2 predictions. This is expected since for low RHs (<40%) SCAPE2 solution, by always attempting to solve for a liquid phase, deviates from the stable state behavior, moving towards the metastable state (see Sect. 3.2).

4.2 Understanding the discrepancies between ISORROPIA II and SCAPE2

The previous discussion provides an overall intercomparison of the two models for a broad RH and composition domain; some cases are further examined to gain more insight as to the cause of discrepancies. In Fig. 5 we compare aerosol water content, aqueous phase potassium, aqueous phase ammonium and aqueous phase nitrate concentration predictions for case 3 (see Table 7) which produced the largest discrepancy in aerosol water and total PM concentrations. This case represents an urban type aerosol with the solution being highly acidic ($R_1=1.27$). Under such conditions, the wa-

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ter content discrepancy between the models is largest for low RHs for the reasons outlined in Sect. 3.2. This is clearly shown in Fig. 5a where only for $RH > 70\%$ SCAPE2 and ISORROPIA II closely follow each other. SCAPE2 predicts significant amount of aqueous phase potassium (Fig. 5b) and ammonium (Fig. 5c) at low relative humidities, while ISORROPIA II predicts gradual deliquescence of K_2SO_4 from 65% to 85% RH. However, SCAPE2 predicts complete deliquescence of K_2SO_4 at $RH=55\%$ which may be due to non-convergence of its numerical solution. Particulate phase ammonium is mainly present as ammonium bisulfate (NH_4HSO_4) and letovicite ($(NH_4)_3H(SO_4)_2$) in ISORROPIA II. However, SCAPE2 predicts the formation of $(NH_4)_3H(SO_4)_2$ only, throughout the whole RH regime. This can also be seen in Fig. 5c where ISORROPIA II predicts a two-step dissolution of ammonium; one at $RH=40\%$ from the deliquescence of $(NH_4)_3H(SO_4)_2$ and one at $RH=70\%$ from the deliquescence of NH_4HSO_4 . Water uptake with SCAPE2 exhibits deliquescence only of $(NH_4)_3H(SO_4)_2$ at $RH=40\%$. Both models predict similar amounts of aqueous phase nitrate for all RHs (Fig. 5d) which shows that the assumption of ISORROPIA II for $NO_{3(aq)}^-$ being a minor species for sulfate – rich cases is a good assumption.

Figure 6 shows comparison of aerosol water, $NaCl_{(s)}$ and $K_{(aq)}$ and $Mg_{(aq)}$ predictions for case 12 (see Table 7), which is a sulfate poor, sodium and crustal species rich aerosol ($R_1=5.1$). The two models agree well (mean error of 5.1%) in aerosol water content predictions (Fig. 6a). SCAPE2, however, predicts significantly higher aqueous potassium for $RH < 40\%$ (Fig. 6b). This is mainly due to different approaches used to associate K with NO_3^- and SO_4^{2-} at low RH. ISORROPIA II uses the principle of “compositional invariance” (Sect. 3.2), hence it preferentially associates K with SO_4^{2-} to form K_2SO_4 , and then KNO_3 and KCl . SCAPE2 tends to partition first as KNO_3 and KCl and then as K_2SO_4 . Since the DRH of KCl is lower than K_2SO_4 , SCAPE2 deliquesces aerosol potassium at a lower RH than ISORROPIA II. Unlike potassium, both models predict the partitioning of sodium between sodium nitrate ($NaNO_3$) and sodium chloride ($NaCl$) in a similar way. This is shown in Fig. 6c where the dissociation of $NaCl_{(s)}$ as a function of RH is similar between both models (NME=12.1%). Aqueous

ISORROPIA II: a new aerosol equilibrium modelC. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

magnesium is the same in both models (Fig. 6d), supporting the postulation (Sect. 3.3) that MgSO_4 never precipitates out of solution.

In Fig. 7 we compare aerosol water, $\text{NO}_{3(\text{aq})}$ and $\text{NH}_{4(\text{aq})}$ as a function of RH for case 5, a sulfate poor, ammonium rich aerosol ($R_1=23.9$, $R_2=0.80$, $R_3=0.37$). Compared to SCAPE2, ISORROPIA II slightly underpredicts aerosol water, aqueous nitrate and ammonium. This difference is seen for RHs between 25–65%. That is because SCAPE2 predicts total deliquescence of sulfates at RH=40% while ISORROPIA II does at RH=70%. The increase of water content shifts the equilibrium of $\text{HNO}_{3(\text{g})} \longleftrightarrow \text{H}_{(\text{aq})}^+ + \text{NO}_{3(\text{aq})}^-$ and $\text{NH}_{3(\text{g})} + \text{H}_2\text{O}_{(\text{aq})} \leftrightarrow \text{NH}_{4(\text{aq})}^+ + \text{OH}_{(\text{aq})}^-$ to the right predicting more aqueous nitrate (Fig. 7b) and ammonium (Fig. 6c) for the same RH regime.

4.3 Metastable vs. stable solutions

The differences between metastable and stable thermodynamic solutions of ISORROPIA II are illustrated in Fig. 8; SCAPE2 is also included for comparison. Figure 8 shows aerosol water and aqueous potassium concentration as a function of relative humidity for a sulfate near-neutral aerosol (case 13 of Table 7). The thermodynamically stable solution of ISORROPIA II predicts deliquescence of the aerosol mixture at 60% RH (DRH of ammonium nitrate). The MDRH for this specific aerosol mixture is 0.46, which explains the aqueous phase potassium (and aerosol water) concentration predicted by the deliquescence solution of ISORROPIA II between 40 and 60% RH. As expected, the metastable solution predicts significant amounts of water below the MDRH (and by definition particulate potassium is deliquesced at all RHs). SCAPE2 yields a solution that is between the stable and metastable ISORROPIA II. Below 40% RH, the predicted concentration of aerosol water by SCAPE2 is slightly larger than the stable solution of ISORROPIA II (Fig. 8a). This results in partial dissolution of aerosol potassium (RH<40%) predicted by SCAPE2 as opposed to the stable solution of ISORROPIA II which does not predict deliquescence of aerosol potassium for this RH regime (Fig. 8b).

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

4.4 “Forward” vs. “Reverse” problem solution

ISORROPIA II is designed to solve both forward and reverse problems. It is useful to assess whether the two solution modes predict identical outputs for the same input. For this assessment, the output from the forward problem (particulate phase concentrations of NH_4 , SO_4 , Na, Cl, NO_3 , Ca, K, and Mg) has been used as input to the reverse problem. The two solution modes are assessed by comparing predictions of aqueous nitrate and sulfate. Since sulfate is only found in the aerosol phase, aqueous phase sulfate calculations are used to evaluate the solid/liquid partitioning behavior between the two solution algorithms, while aqueous nitrate is used as a proxy for gas-aerosol partitioning (for all the conditions specified in Table 7). The agreement between the two solutions was found to be excellent with the NME being $3.4 \pm 1.1\%$ for aqueous sulfate and $2.5 \pm 1.3\%$ for aqueous nitrate concentration.

4.5 Computational speed

The timing tests were performed on a Dell 8300 Intel Pentium 4 CPU 3.20 GHz, 512 MB of RAM workstation running Windows XP operating system. Both codes were compiled with Watcom FORTRAN compiler version 2.0 with full optimization options on. Table 9 shows the CPU time needed by the two models for the aerosol types described in Table 7. ISORROPIA II consumes much less CPU time compared to SCAPE2 with the difference being at least an order of magnitude for all aerosol cases. The amount of time required by ISORROPIA II for each aerosol case was found to be approximately the same even if the convergence criterion for solids and water was decreased down to 10^{-6} or 10^{-7} which is a proof of the rapid and robust convergence of the code. However, larger convergence criterion was used for the intercomparison study (see Table 9), to assure a quick and convergent solution from SCAPE2. For completeness we also compare the CPU time required by ISORROPIA (version 1.7, 03/15/2006) for all the simulation conditions of Table 7, but with crustals set to zero). Although ISORROPIA II solves for more species than ISORROPIA, it is not slower because of

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

optimizations in the activity coefficient calculation algorithm in ISORROPIA II. Finally, in Table 9 we compare the CPU time required by the stable and metastable solutions of ISORROPIA II. As expected, the metastable solution is slightly faster than the stable solution since the absence of solid species requires the solution of fewer equations.

5 Summary

A new model, ISORROPIA II, is developed which treats the thermodynamics of $K^+ - Ca^{2+} - Mg^{2+} - NH_4^+ - Na^+ - SO_4^{2-} - NO_3^- - Cl^- - H_2O$ aerosol systems. A comprehensive evaluation was conducted against the thermodynamic model SCAPE2 in terms of composition predicted and computational speed for a wide variety of aerosol conditions that cover typical urban, remote continental, marine and non-urban continental environments. The overall predictions of aerosol water, total PM and concentration of semi-volatile species were generally comparable between the two models under most conditions. For aerosol water content and total PM mass the two models agreed within approximately 13%. The normalized mean error for total aerosol nitrate predictions was 16% while for aerosol chloride and ammonium concentration the agreement was within 2–6%. Small discrepancies were found to exist between the two models under certain conditions, primarily for relative humidities between 40 and 70%. These discrepancies are mainly attributed to the solution dynamics treatment of water uptake in mutual deliquescence regions and the association of non-volatile cations with sulfate, nitrate and chloride. For all cases examined, ISORROPIA II is more than an order of magnitude faster than SCAPE2, showing robust and rapid convergence for all conditions examined, making it one of the most computationally efficient and comprehensive inorganic thermodynamic equilibrium modules available.

Acknowledgements. This work was supported by the National Oceanic and Atmospheric Administration under contract NMRAC000-5-04017. We would like to acknowledge the contributions and support of the whole community that lead to significant improvements of the original

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ISORROPIA code. In particular we would like to acknowledge the contributions of (in alphabetical order): A. Ansari, V. Bouchet, P. Bhave, S. Clegg, B. Hutzl, K. Kapaldo, B. Koo, S. Kreidenweis, P. Makar, F. San Martini, D. Mauzerall, C. Nolte, B. Pun, A. Russell, U. Shankar, D. Tong, J. West, D. Waldron, A. Zakey, and Y. Zhang. We also would like to thank C. Pilinis for his
5 comments on the manuscript.

References

- Altshüller, A. P.: Atmospheric particle sulfur and sulfur dioxide relationships at urban and nonurban locations, *Atmos. Environ.*, 18, 1421–1431, 1984.
- Amundson, N. R., Caboussat, A., He, J. W., Martynenko, A. V., Savarin, V. B., Seinfeld, J. H., and Yoo, K. Y.: A new inorganic atmospheric aerosol phase equilibrium model (UHAERO), *Atmos. Chem. Phys.*, 6, 975–992, 2006,
10 <http://www.atmos-chem-phys.net/6/975/2006/>.
- Ansari, A. S. and Pandis, S. N.: Prediction of multicomponent inorganic atmospheric aerosol behavior, *Atmos. Environ.*, 33, 745–757, 1999a.
- 15 Ansari, A. S. and Pandis, S. N.: An analysis of four models predicting the partitioning of semivolatile inorganic aerosol components, *Aerosol Sci. Technol.*, 31, 129–153, 1999b.
- Brauer, M. and Brook, J. R.: Ozone personal exposures and health effects for selected groups residing in the Fraser Valley, *Atmos. Environ.*, 31, 2113–2122, 1997.
- Bromley, L. A.: Thermodynamic properties of strong electrolytes in aqueous solutions, *AIChE J.*, 19, 313–320, 1973.
20
- Chan, M. N. and Chan, C. K.: Mass transfer effects in hygroscopic measurements of aerosol particles, *Atmos. Chem. Phys.*, 5, 2703–2712, 2005,
<http://www.atmos-chem-phys.net/5/2703/2005/>.
- Clegg, S. L. and Pitzer, K. S.: Thermodynamics of multicomponent, miscible, ionic solutions: generalized equations for symmetrical electrolytes, *J. Phys. Chem.*, 96, 3513–3520, 1992.
- 25 Clegg, S. L., Pitzer, K. S., and Brimblecombe, P.: Thermodynamics of multicomponent, miscible, ionic solutions. II. Mixture including unsymmetrical electrolytes, *J. Phys. Chem.*, 96, 9470–9479, 1992.
- Clegg, S. L., Pitzer, K. S., and Brimblecombe, P.: (additions and corrections for their published
30 papers), *J. Phys. Chem.*, 98, 1368, 1994.

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Clegg, S. L., Pitzer, K. S., and Brimblecombe, P.: (additions and corrections for their published papers), *J. Phys. Chem.*, 99, 6755, 1995.
- Clegg, S. L., Brimblecombe, P., and Wexler, A. S.: Thermodynamic model of the system $\text{H}^+ - \text{NH}_4^+ - \text{SO}_4^{2-} - \text{NO}_3^- - \text{H}_2\text{O}$ at tropospheric temperatures, *J. Phys. Chem., A*, 102, 2137–2154, 1998a.
- Clegg, S. L., Brimblecombe, P., and Wexler, A. S.: Thermodynamic model of the system $\text{H}^+ - \text{NH}_4^+ - \text{Na}^+ - \text{SO}_4^{2-} - \text{NO}_3^- - \text{Cl}^- - \text{H}_2\text{O}$ at 298.15 K, *J. Phys. Chem., A*, 102, 2155–2171, 1998b.
- Denbigh, K.: *The principles of chemical equilibrium*, Fourth Ed., Cambridge University Press, Cambridge, 1981.
- Dockery, D. W., Pope, C. A., Xu, X., Spengler, J. D., Ware, J. H., Fay, M. E., Ferris, B. G., and Speizer, F. E.: An Association between Air Pollution and Mortality in Six U.S. Cities, *The New England Journal of Medicine*, 329, 1753–1759, 1993.
- Fitzgerald, J. W.: Marine aerosols: A review, *Atmos. Environ.*, 25A, 533–545, 1991.
- Ha, Z. and Chan, C. K.: The water activities of MgCl_2 , $\text{Mg}(\text{NO}_3)_2$, MgSO_4 , and their mixtures, *Aerosol Sci. Technol.*, 31, 154–169, 1999.
- Heitzenberg, J.: Fine particles in the global troposphere: a review, *Tellus* 41B, 149–160, 1989.
- Intergovernmental Panel on Climate Change: *The scientific Basis*, Cambridge Univ. Press, New York, 2001.
- Jacobson, M. Z., Tabazadeh, A., and Turco, R. P.: Simulating equilibrium within aerosols and nonequilibrium between gases and aerosols, *J. Geophys. Res.*, 101, 9079–9091, 1996.
- Jacobson, M. Z.: *Chemical Equilibrium and dissolution processes*, *Fundamentals of Atmospheric Modeling*, Cambridge University Press, New York, 476–510, (Chapter 18), 1999a.
- Jacobson, M. Z.: Studying the effect of calcium and magnesium on size-distributed nitrate and ammonium with EQUISOLV II, *Atmos. Environ.*, 33, 3635–3649, 1999b.
- Kaiser, J.: Mounting evidence indicts fine-particle pollution, *Science*, 307, 1858–1861, 2005.
- Kelly, J. T. and Wexler, A. S.: Water uptake by aerosol: Water activity in supersaturated potassium solutions and deliquescence as a function of temperature, *Atmos. Environ*, 40, 4450–4468, 2006.
- Kim, Y. P., Seinfeld, J. H., and Saxena, P.: Atmospheric gas – aerosol equilibrium I. Thermodynamic model, *Aerosol Sci. Technol.*, 19, 157–181, 1993a.
- Kim, Y. P., Seinfeld, J. H., and Saxena, P.: Atmospheric gas – aerosol equilibrium II. Analysis of common approximations and activity coefficient calculation methods, *Aerosol Sci. Technol.*,

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

19, 182–198, 1993b.

Kim, Y. P. and Seinfeld, J. H.: Atmospheric gas - aerosol equilibrium III. Thermodynamics of crustal elements Ca^{2+} , K^+ , and Mg^{2+} , *Aerosol Sci. Technol.*, 22, 93–110, 1995.

Kusik, C. L. and Meissner, H. P.: Electrolyte activity coefficients in inorganic processing, *AIChE Symp. Series*, 173, 14–20, 1978.

Makar, P. A., Bouchet, V. S., and Nenes, A.: Inorganic chemistry calculations using HETV – a vectorized solver for the SO_4^{2-} - NO_3^- - NH_4^+ system based on the ISORROPIA algorithms, *Atmos. Environ.*, 37, 2279–2294, 2003.

Meng, Z. Y., Seinfeld, J. H., Saxena, P., and Kim, Y. P.: Atmospheric gas – aerosol equilibrium IV. Thermodynamics of carbonates, *Aerosol Sci. Technol.*, 23, 131–154, 1995.

Meissner, H. P., Kusik, C. L., and Tester, J. W.: Activity coefficients of strong electrolytes in aqueous solution – effect of temperature, *AIChE Journal*, 18(3), 661–662, 1972.

Meissner, H. P. and Peppas, N. A.: Activity coefficients - aqueous Solutions of polybasic acids and their salts, *AIChE Journal*, 19(4), 806–809, 1973.

Metzger, S. M., Dentener, F. J., and Lelieveld, J.: Aerosol multiphase chemistry – A parameterization for global modeling, *Int. Rep. 99-12, for Mar. and Atmos. Res.*, Utrecht, Netherlands, 1999.

Metzger, S. M., Dentener, F. J., Lelieveld, J., and Pandis, S. N.: Gas/aerosol partitioning I: a computationally efficient model, *J. Geophys. Res.*, 107(D16), 4312, doi:10.1029/2001JD001102, 2002a.

Metzger, S. M., Dentener, F. J., Jeuken, A., Krol, M., and Lelieveld, J.: Gas/aerosol partitioning II: global modeling results, *J. Geophys. Res.*, 107(D16), 4313, doi:10.1029/2001JD001103, 2002b.

Metzger, S., Mihalopoulos, N., and Lelieveld, J.: Importance of mineral cations and organics in gas-aerosol partitioning of reactive nitrogen compounds: case study based on MINOS results, *Atmos. Chem. Phys.*, 6, 2549–2567, 2006,

<http://www.atmos-chem-phys.net/6/2549/2006/>.

Moya, M., Ansari, A. S., and Pandis, S. N.: Partitioning of nitrate and ammonium between the gas and particulate phases during the 1997 IMADA-AVER study in Mexico City, *Atmos. Environ.*, 35, 1791–1804, 2001a.

Moya, M., Pandis, S. N., and Jacobson, M. Z.: Is the size distribution of urban aerosols determined by thermodynamic equilibrium? An application to Southern California, *Atmos. Environ.*, 36, 2349–2365, 2001b.

ACPD

7, 1893–1939, 2007

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ISORROPIA II: a new aerosol equilibrium modelC. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Nenes, A., Pandis, S. N., and Pilinis, C.: ISORROPIA: A new thermodynamic equilibrium model for multiphase multicomponent inorganic aerosols, *Aquatic Geochem.*, 4, 123–152, 1998.
- Nenes, A., Pilinis, C., and Pandis, S. N.: Continued development and testing of a new thermodynamic aerosol module for urban and regional air quality models, *Atmos. Environ.*, 33, 1553–1560, 1999.
- Pilinis, C., and Seinfeld, J. H.: Continued development of a general equilibrium model for inorganic multicomponent atmospheric aerosols, *Atmos. Environ.*, 21, 2453–2466, 1987.
- Pilinis, C. and Seinfeld, J. H.: Water content of atmospheric aerosols, *Atmos. Environ.*, 23, 1601–1606, 1989.
- Pitzer, K. S. and Simonson, J. M.: Thermodynamics of multicomponent, miscible, ionic systems: Theory and equations, *J. Phys. Chem.*, 90, 3005–3009, 1986.
- Potukuchi, S. and Wexler, A. S.: Identifying solid-aqueous phase transitions in atmospheric aerosols – I. Neutral-acidity solutions, *Atmos. Environ.*, 29, 1663–1676, 1995a.
- Potukuchi, S. and Wexler, A. S.: Identifying solid-aqueous phase transitions in atmospheric aerosols – II. Acidic solutions, *Atmos. Environ.*, 29, 3357–3364, 1995b.
- Ramachandran, G. and Vincent, J. H.: A Bayesian approach to retrospective exposure assessment, *Applied Occupational and Environmental Hygiene*, 14, 547–558, 1999.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, John Wiley & Sons, Inc., 1998.
- Stelson, A. W. and Seinfeld, J. H.: Relative humidity and temperature dependence of the ammonium nitrate dissociation constant, *Atmos. Environ.*, 16(5), 983–992, 1982.
- Thomsen, K. and Rasmussen, P.: Modeling of vapor-liquid-solid equilibrium in gas-aqueous electrolyte systems, *Chem. Eng. Sci.*, 54, 1787–1802, 1999.
- Wexler, A. S. and Seinfeld, J. H.: The distribution of ammonium salts among a size and composition dispersed aerosol, *Atmos. Environ.*, 24A, 1231–1246, 1990.
- Wexler, A. S. and Seinfeld, J. H.: Second – generation inorganic aerosol model, *Atmos. Environ.*, 25A, 2731–2748, 1991.
- Wexler, A. S. and Clegg, S. L.: Atmospheric aerosol models for systems including the ions H^+ , NH_4^+ , Na^+ , SO_4^{2-} , NO_3^- , Cl^- , Br^- , and H_2O , *J. Geophys. Res.*, 107, 4207, doi:10.1029/2001JD000451, 2002.
- Yu, S., Dennis, R., Roselle, S., Nenes, A., Walker, J., Eder, B., Schere, K., Swall, J., and Rorbarge, W.: An assessment of the ability of three-dimensional air quality models with current thermodynamic equilibrium models to predict aerosol NO_3^- , *J. Geophys. Res.*, 110, D07S13,

doi:10.1029/2004JD004718, 2005.

Zanobetti, A., Schwartz, J., and Dockery, D. W.: Airborne particles are a risk factor for hospital admissions for heart and lung disease, *Environmental Health Perspective*, 108, 1071–1082, 2000.

5 Zaveri, R. A., Easter, R. C., and Peters, L. K.: A computationally efficient multi-component equilibrium solver for aerosols (MESA), *J. Geophys. Res.*, 110, D24203, doi:10.1029/2004JD005618, 2005a.

Zaveri, R. A., Easter, R. C., and Wexler, A. S.: A new method for multicomponent activity coefficients of electrolytes in aqueous atmospheric aerosols, *J. Geophys. Res.*, 110, D02201, 10 doi:10.1029/2004JD004681, 2005b.

Zhang, Y., Seigneur, C., Seinfeld, J. H., Jacobson, M., Clegg, S. L., and Binkowski, F. S.: A comparative review of inorganic aerosol thermodynamic equilibrium models: similarities, differences, and their likely causes, *Atmos. Environ.*, 34, 117–137, 2000.

ACPD

7, 1893–1939, 2007

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. Thermodynamic Properties for all species in ISORROPIA II*.

Species	$\Delta\mu_i^0(298\text{K}), \text{kJ mol}^{-1}$	$\Delta H_i^0, \text{kJ mol}^{-1}$	$C_p^0, \text{J mol}^{-1} \text{K}^{-1}$
Ca(NO₃)₂(s)	-743.070	-938.390	149.370
CaCl₂(s)	-748.10	-795.800	72.590
CaSO₄(s)	-1798.280	-2022.630	186.020
KHSO₄(s)	-1031.300	-1160.600	87.160
K₂SO₄(s)	-1321.370	-1437.790	131.460
KNO₃(s)	-394.860	-494.630	96.400
KCl(s)	-409.140	-434.750	51.300
MgSO₄(s)	-1170.600	-1284.900	96.480
Mg(NO₃)₂(s)	-589.400	-790.650	141.920
MgCl₂(s)	-591.790	-641.320	71.380
Ca²⁺_(aq)	-553.580	-542.830	-
K⁺_(aq)	-283.270	-252.380	21.800
Mg²⁺_(aq)	-454.800	-466.850	-
NaCl(s)	-384.138	-411.153	50.500
NaNO ₃ (s)	-367.000	-467.850	92.880
Na ₂ SO ₄ (s)	-1270.160	-1387.080	128.200
NaHSO ₄ (s)	-992.800	-1125.500	85.000
NH ₄ Cl(s)	-202.870	-314.430	84.100
NH ₄ NO ₃ (s)	-183.870	-365.560	139.300
(NH ₄) ₂ SO ₄ (s)	-901.670	-1180.850	187.490
NH ₄ HSO ₄ (s)	-823.000	-1026.960	127.500
(NH ₄) ₃ H(SO ₄) ₂ (s)	-1730.000	-2207.000	315.000
HNO ₃ (g)	-74.720	-135.060	53.350
HCl(g)	-95.299	-92.307	29.126
NH ₃ (g)	-16.450	-46.110	35.060
NH ₃ (aq)	-26.500	-80.290	79.900
H ⁺ _(aq)	0.000	0.000	0.000
Na ⁺ _(aq)	-261.905	-240.120	46.400
NH ₄ ⁺ _(aq)	-79.310	-132.510	79.900
HSO ₄ ⁻ _(aq)	-755.910	-887.340	-84.000
SO ₄ ²⁻ _(aq)	-744.530	-909.270	-293.000
NO ₃ ⁻ _(aq)	-111.250	-207.360	-86.600
Cl ⁻ _(aq)	-131.228	-167.159	-136.400
OH ⁻ _(aq)	-157.244	-229.994	-148.500

* Compiled by: Kim and Seinfeld, (1995), Kim et al., (1993); Species in bold are new in ISORROPIA II.

- Data not available.

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. Equilibrium Relations and Temperature Dependence Constants used in ISORROPIA II*.

Reaction	Equilibrium Constant Expression	$K^0(298.15\text{ K})$	$\frac{\Delta H^0(T_0)}{RT_0}$	$\frac{\Delta C_p^0}{R}$	Units
$\text{Ca}(\text{NO}_3)_2(\text{s}) \leftrightarrow \text{Ca}_{(\text{aq})}^{2+} + 2\text{NO}_3^-(\text{aq})$	$[\text{Ca}^{2+}][\text{NO}_3^-]^2 \gamma_{\text{Ca}^{2+}} \gamma_{\text{NO}_3^-}^2$	6.067×10^5	9.549	–	$\text{mol}^3 \text{kg}^{-3}$
$\text{CaCl}_2(\text{s}) \leftrightarrow \text{Ca}_{(\text{aq})}^{2+} + 2\text{Cl}^-(\text{aq})$	$[\text{Ca}^{2+}][\text{Cl}^-]^2 \gamma_{\text{Ca}^{2+}} \gamma_{\text{Cl}^-}^2$	7.974×10^{11}	27.995	–	$\text{mol}^3 \text{kg}^{-3}$
$\text{CaSO}_4 \cdot 2\text{H}_2\text{O}(\text{s}) \leftrightarrow \text{Ca}_{(\text{aq})}^{2+} + \text{SO}_4^{2-}(\text{aq}) + 2\text{H}_2\text{O}$	$[\text{Ca}^{2+}][\text{SO}_4^{2-}] \gamma_{\text{Ca}^{2+}} \gamma_{\text{SO}_4^{2-}} a_w^2$	4.319×10^{-5}	–	–	$\text{mol}^2 \text{kg}^{-2}$
$\text{K}_2\text{SO}_4(\text{s}) \leftrightarrow 2\text{K}_{(\text{aq})}^+ + \text{SO}_4^{2-}(\text{aq})$	$[\text{K}^+]^2 [\text{SO}_4^{2-}] \gamma_{\text{K}^+}^2 \gamma_{\text{SO}_4^{2-}}$	1.569×10^{-2}	–9.589	45.807	$\text{mol}^3 \text{kg}^{-3}$
$\text{KHSO}_4(\text{s}) \leftrightarrow \text{K}_{(\text{aq})}^+ + \text{HSO}_4^-(\text{aq})$	$[\text{K}^+][\text{HSO}_4^-] \gamma_{\text{K}^+} \gamma_{\text{HSO}_4^-}$	24.016	–8.423	17.964	$\text{mol}^2 \text{kg}^{-2}$
$\text{KNO}_3(\text{s}) \leftrightarrow \text{K}_{(\text{aq})}^+ + \text{NO}_3^-(\text{aq})$	$[\text{K}^+][\text{NO}_3^-] \gamma_{\text{K}^+} \gamma_{\text{NO}_3^-}$	0.872	14.075	19.388	$\text{mol}^2 \text{kg}^{-2}$
$\text{KCl}(\text{s}) \leftrightarrow \text{K}_{(\text{aq})}^+ + \text{Cl}^-(\text{aq})$	$[\text{K}^+][\text{Cl}^-] \gamma_{\text{K}^+} \gamma_{\text{Cl}^-}$	8.680	–6.167	19.953	$\text{mol}^2 \text{kg}^{-2}$
$\text{MgSO}_4(\text{s}) \leftrightarrow \text{Mg}_{(\text{aq})}^{2+} + \text{SO}_4^{2-}(\text{aq})$	$[\text{Mg}^{2+}][\text{SO}_4^{2-}] \gamma_{\text{Mg}^{2+}} \gamma_{\text{SO}_4^{2-}}$	1.079×10^5	36.798	–	$\text{mol}^2 \text{kg}^{-2}$
$\text{Mg}(\text{NO}_3)_2(\text{s}) \leftrightarrow \text{Mg}_{(\text{aq})}^{2+} + 2\text{NO}_3^-(\text{aq})$	$[\text{Mg}^{2+}][\text{NO}_3^-]^2 \gamma_{\text{Mg}^{2+}} \gamma_{\text{NO}_3^-}^2$	2.507×10^{15}	36.677	–	$\text{mol}^3 \text{kg}^{-3}$
$\text{MgCl}_2(\text{s}) \leftrightarrow \text{Mg}_{(\text{aq})}^{2+} + 2\text{Cl}^-(\text{aq})$	$[\text{Mg}^{2+}][\text{Cl}^-]^2 \gamma_{\text{Mg}^{2+}} \gamma_{\text{Cl}^-}^2$	9.557×10^{21}	59.352	–	$\text{mol}^3 \text{kg}^{-3}$
$\text{HSO}_4^-(\text{aq}) \leftrightarrow \text{H}_{(\text{aq})}^+ + \text{SO}_4^{2-}(\text{aq})$	$\frac{[\text{H}^+][\text{SO}_4^{2-}]}{[\text{HSO}_4^-]} \gamma_{\text{H}^+} \gamma_{\text{SO}_4^{2-}} / \gamma_{\text{HSO}_4^-}$	1.015×10^{-2}	8.85	25.14	mol kg^{-1}
$\text{NH}_3(\text{g}) \leftrightarrow \text{NH}_3(\text{aq})$	$\frac{P_{\text{NH}_3(\text{aq})}}{P_{\text{NH}_3}} \gamma_{\text{NH}_3}$	5.764×10^1	13.79	–5.39	$\text{mol kg}^{-1} \text{atm}^{-1}$
$\text{NH}_4^+(\text{aq}) \cdot \text{H}_2\text{O}(\text{aq}) \leftrightarrow \text{NH}_4^+(\text{aq}) + \text{OH}^-(\text{aq})$	$\frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_4^+(\text{aq})] a_w} \gamma_{\text{NH}_4^+} \gamma_{\text{OH}^-} / \gamma_{\text{NH}_3}$	1.805×10^{-5}	–1.50	26.92	mol kg^{-1}
$\text{HNO}_3(\text{g}) \leftrightarrow \text{H}_{(\text{aq})}^+ + \text{NO}_3^-(\text{aq})$	$\frac{P_{\text{HNO}_3}}{[\text{H}^+][\text{NO}_3^-]} \gamma_{\text{H}^+} \gamma_{\text{NO}_3^-}$	2.511×10^6	29.17	16.83	$\text{mol}^2 \text{kg}^{-2} \text{atm}^{-1}$
$\text{HNO}_3(\text{g}) \leftrightarrow \text{HNO}_3^*(\text{aq})$	$\frac{P_{\text{HNO}_3(\text{aq})}}{P_{\text{HNO}_3}} \gamma_{\text{HNO}_3}$	2.1×10^5	29.17	16.83	$\text{mol kg}^{-1} \text{atm}^{-1}$
$\text{HCl}_{(\text{g})} \leftrightarrow \text{H}_{(\text{aq})}^+ + \text{Cl}^-(\text{aq})$	$\frac{P_{\text{HCl}}}{[\text{H}^+][\text{Cl}^-]} \gamma_{\text{H}^+} \gamma_{\text{Cl}^-}$	1.971×10^6	30.20	19.91	$\text{mol}^2 \text{kg}^{-2} \text{atm}^{-1}$
$\text{HCl}_{(\text{g})} \leftrightarrow \text{HCl}^\diamond(\text{aq})$	$\frac{P_{\text{HCl}}}{[\text{HCl}(\text{aq})]} \gamma_{\text{HCl}}$	2.5×10^3	30.20	19.91	$\text{mol kg}^{-1} \text{atm}^{-1}$
$\text{H}_2\text{O}_{(\text{aq})}^+ \leftrightarrow \text{H}_{(\text{aq})}^+ + \text{OH}^-(\text{aq})$	$\frac{[\text{H}^+][\text{OH}^-]}{a_w} \gamma_{\text{H}^+} \gamma_{\text{OH}^-}$	1.010×10^{-14}	–22.52	26.92	$\text{mol}^2 \text{kg}^{-2}$

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. Continued.

Reaction	Equilibrium Constant Expression	K^0 (298.15K)	$\frac{\Delta H^0(T_0)}{RT_0}$	$\frac{\Delta C_p^0}{R}$	Units
$\text{Na}_2\text{SO}_{4(s)} \leftrightarrow 2\text{Na}_{(\text{aq})}^+ + \text{SO}_{4(\text{aq})}^{2-}$	$[\text{Na}^+]^2 [\text{SO}_4^{2-}] \gamma_{\text{Na}^+}^2 \gamma_{\text{SO}_4^{2-}}$	4.799×10^{-1}	0.98	39.75	$\text{mol}^3 \text{kg}^{-3}$
$(\text{NH}_4)_2 \text{SO}_{4(s)} \leftrightarrow 2\text{NH}_{4(\text{aq})}^+ + \text{SO}_{4(\text{aq})}^{2-}$	$[\text{NH}_4^+]^2 [\text{SO}_4^{2-}] \gamma_{\text{NH}_4^+}^2 \gamma_{\text{SO}_4^{2-}}$	1.817×10^0	-2.65	38.57	$\text{mol}^3 \text{kg}^{-3}$
$\text{NH}_4\text{Cl}_{(s)} \leftrightarrow \text{NH}_{3(g)} + \text{HCl}_{(g)}$	$P_{\text{NH}_3} P_{\text{HCl}}$	1.086×10^{-16}	-71.00	2.40	atm^2
$\text{NaNO}_{3(s)} \leftrightarrow \text{Na}_{(\text{aq})}^+ + \text{NO}_{3(\text{aq})}^-$	$[\text{Na}^+] [\text{NO}_3^-] \gamma_{\text{Na}^+} \gamma_{\text{NO}_3^-}$	1.197×10^1	-8.22	16.01	$\text{mol}^2 \text{kg}^{-2}$
$\text{NaCl}_{(s)} \leftrightarrow \text{Na}_{(\text{aq})}^+ + \text{Cl}_{(\text{aq})}^-$	$[\text{Na}^+] [\text{Cl}^-] \gamma_{\text{Na}^+} \gamma_{\text{Cl}^-}$	3.766×10^1	-1.56	16.90	$\text{mol}^2 \text{kg}^{-2}$
$\text{NaHSO}_{4(s)} \leftrightarrow \text{Na}_{(\text{aq})}^+ + \text{HSO}_{4(\text{aq})}^-$	$[\text{Na}^+] [\text{HSO}_4^-] \gamma_{\text{Na}^+} \gamma_{\text{HSO}_4^-}$	2.413×10^4	0.79	14.75	$\text{mol}^2 \text{kg}^{-2}$
$\text{NH}_4\text{NO}_{3(s)} \leftrightarrow \text{NH}_{3(g)} + \text{HNO}_{3(g)}$	$P_{\text{NH}_3} P_{\text{HNO}_3}$	4.199×10^{-17}	-74.735	6.025	atm^2
$\text{NH}_4\text{HSO}_{4(s)} \leftrightarrow \text{NH}_{4(\text{aq})}^+ + \text{HSO}_{4(\text{aq})}^-$	$[\text{NH}_4^+] [\text{HSO}_4^-] \gamma_{\text{NH}_4^+} \gamma_{\text{HSO}_4^-}$	1.383×10^0	-2.87	15.83	$\text{mol}^2 \text{kg}^{-2}$
$(\text{NH}_4)_3 \text{H}(\text{SO}_4)_2(s) \leftrightarrow 3\text{NH}_{4(\text{aq})}^+ + \text{HSO}_{4(\text{aq})}^- + \text{SO}_{4(\text{aq})}^{2-}$	$[\text{NH}_4^+]^3 [\text{SO}_4^{2-}] [\text{HSO}_4^-] \gamma_{\text{NH}_4^+}^3 \gamma_{\text{SO}_4^{2-}} \gamma_{\text{HSO}_4^-}$	2.972×10^1	-5.19	54.40	$\text{mol}^5 \text{kg}^{-5}$

* Compiled by: Kim and Seinfeld (1995), Kim et al. (1993); Reactions with constants in bold are new in ISORROPIA II.

♣ The equilibrium constant K_{1b} of the reaction $\text{HNO}_{3(\text{aq})} \xrightleftharpoons{K_{1b}} \text{H}_{(\text{aq})}^+ + \text{NO}_{3(\text{aq})}^-$ is calculated from K_1 and K_{1a} of the reactions $\text{HNO}_{3(\text{g})} \xrightleftharpoons{K_1} \text{H}_{(\text{aq})}^+ + \text{NO}_{3(\text{aq})}^-$ and $\text{HNO}_{3(\text{g})} \xrightleftharpoons{K_{1a}} \text{HNO}_{3(\text{aq})}$, respectively: $K_{1b} = K_1 / K_{1a}$.

◇ The equilibrium constant K_{2b} of the reaction $\text{HCl}_{(\text{aq})} \xrightleftharpoons{K_{2b}} \text{H}_{(\text{aq})}^+ + \text{Cl}_{(\text{aq})}^-$ is calculated from K_2 and K_{2a} of the reactions $\text{HCl}_{(\text{g})} \xrightleftharpoons{K_2} \text{H}_{(\text{aq})}^+ + \text{Cl}_{(\text{aq})}^-$ and $\text{HCl}_{(\text{g})} \xrightleftharpoons{K_{2a}} \text{HCl}_{(\text{aq})}$, respectively: $K_{2b} = K_2 / K_{2a}$.

- Data not available.

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 3. Potential species for the five aerosol types¹.

R_1	R_2	R_3	Aerosol Type	Major Species			Minor Species
				Solid Phase	Aqueous Phase	Gas Phase	
$R_1 < 1$	any value	any value	Sulfate Rich (free acid)	NaHSO ₄ , NH ₄ HSO ₄ , KHSO₄ , CaSO₄	Na ⁺ , NH ₄ ⁺ , H ⁺ , HSO ₄ ⁻ , SO ₄ ²⁻ , NO ₃ ⁻ , Cl ⁻ , Ca²⁺ , K⁺ , H ₂ O	H ₂ O _(g)	NH _{3(g)} , NO _{3(aq)} ⁻ , Cl _(aq) ⁻ , NH _{3(aq)} , HNO _{3(aq)} , HCl _(aq)
$1 \leq R_1 < 2$	any value	any value	Sulfate Rich	NaHSO ₄ , NH ₄ HSO ₄ , Na ₂ SO ₄ , (NH ₄) ₂ SO ₄ , (NH ₄) ₃ H(SO ₄) ₂ , CaSO₄ , KHSO₄ , K₂SO₄ , MgSO₄	Na ⁺ , NH ₄ ⁺ , H ⁺ , HSO ₄ ⁻ , SO ₄ ²⁻ , NO ₃ ⁻ , Cl ⁻ , Ca²⁺ , K⁺ , Mg²⁺ , H ₂ O	H ₂ O _(g)	NH _{3(g)} , NO _{3(aq)} ⁻ , Cl _(aq) ⁻ , NH _{3(aq)} , HNO _{3(aq)} , HCl _(aq)
$R_1 \geq 2$	$R_2 < 2$	any value	Sulfate Poor, Crustals & Sodium Poor	Na ₂ SO ₄ , (NH ₄) ₂ SO ₄ , NH ₄ NO ₃ , NH ₄ Cl, CaSO₄ , K₂SO₄ , MgSO₄	Na ⁺ , NH ₄ ⁺ , H ⁺ , SO ₄ ²⁻ , NO ₃ ⁻ , Cl ⁻ , Ca²⁺ , K⁺ , Mg²⁺ , H ₂ O, NH _{3(aq)} , HNO _{3(aq)} , HCl _(aq)	HNO _{3(g)} , HCl _(g) , NH _{3(g)} , H ₂ O _(g)	HSO ₄ ⁻
$R_1 \geq 2$	$R_2 \geq 2$	$R_3 < 2$	Sulfate Poor, Crustals & Sodium Rich, Crustals Poor	Na ₂ SO ₄ , NaNO ₃ , NaCl, NH ₄ NO ₃ , NH ₄ Cl, CaSO₄ , K₂SO₄ , MgSO₄	Na ⁺ , NH ₄ ⁺ , H ⁺ , SO ₄ ²⁻ , NO ₃ ⁻ , Cl ⁻ , Ca²⁺ , K⁺ , Mg²⁺ , H ₂ O, NH _{3(aq)} , HNO _{3(aq)} , HCl _(aq)	HNO _{3(g)} , HCl _(g) , NH _{3(g)} , H ₂ O _(g)	HSO ₄ ⁻
$R_1 \geq 2$	$R_2 \geq 2$	$R_3 > 2$	Sulfate Poor, Crustals & Sodium Rich, Crustals Rich	NaNO ₃ , NaCl, NH ₄ NO ₃ , NH ₄ Cl, CaSO₄ , K₂SO₄ , MgSO₄ , Ca(NO₃)₂ , CaCl₂ , Mg(NO₃)₂ , MgCl₂ , KNO₃ , KCl	Na ⁺ , NH ₄ ⁺ , H ⁺ , SO ₄ ²⁻ , NO ₃ ⁻ , Cl ⁻ , Ca²⁺ , K⁺ , Mg²⁺ , H ₂ O, NH _{3(aq)} , HNO _{3(aq)} , HCl _(aq)	HNO _{3(g)} , HCl _(g) , NH _{3(g)} , H ₂ O _(g)	HSO ₄ ⁻

¹ Species in bold are new in ISORROPIA II.

Table 4. Deliquescence relative humidities, temperature dependence and parameter q values for all the salts modeled in ISORROPIA II².

Salt	DRH (298.15 K)	$-\frac{18}{1000R}L_s m_s$	q
Ca(NO₃)₂	0.4906	-430.5	0.93
CaCl₂	0.2830	-1121.0	2.40
CaSO₄	0.9700	–	–, ^a
KHSO₄	0.8600 [♣]	–	–, ^b
K₂SO₄	0.9751	35.6	-0.25
KNO₃	0.9248	–	-2.33
KCl_(s)	0.8426	158.9	0.92
MgSO₄	0.8613 [◇]	-714.5 *	0.15 *
Mg(NO₃)₂ [◇]	0.5400	–	2.32
MgCl₂	0.3284	-1860.2	2.90
NaCl [♣]	0.7528	25.0	2.23
Na ₂ SO ₄ [♣]	0.9300	80.0	-0.19
NaNO ₃ [♣]	0.7379	304.0	-0.39
(NH ₄) ₂ SO ₄ [♣]	0.7997	80.0	-0.25
NH ₄ NO ₃ [♣]	0.6183	852.0	-1.15
NH ₄ Cl [♣]	0.7710	239.0	0.82
NH ₄ HSO ₄ [♣]	0.4000	384.0	(+), ^c
NaHSO ₄ [♣]	0.5200	-45.0	(+), ^d
(NH ₄) ₃ H(SO ₄) ₂ [♣]	0.6900	186.0	(+), ^e
H ₂ SO ₄ [♣]	0.000	–	0.70
H-HSO ₄ [♣]	0.000	–	8.00
HNO ₃ [♣]	N/A	–	2.60
HCl [♣]	N/A	–	6.00

* Kim and Seinfeld (1995); [♣] Pilinis and Seinfeld (1989); [◇] Ha and Chan (1999); [♣] Kim et al. (1993); – Data not available. ^a $\gamma_{\text{CaSO}_4}=0$; ^b $\gamma_{\text{KHSO}_4}=\frac{\gamma_{\text{H-HSO}_4}\cdot\gamma_{\text{KCl}}}{\gamma_{\text{HCl}}}$; ^c $\gamma_{\text{NH}_4\text{HSO}_4}=\frac{\gamma_{\text{H-HSO}_4}\cdot\gamma_{\text{NH}_4\text{Cl}}}{\gamma_{\text{HCl}}}$;

^d $\gamma_{\text{NaHSO}_4}=\frac{\gamma_{\text{H-HSO}_4}\cdot\gamma_{\text{NaCl}}}{\gamma_{\text{HCl}}}$; ^e $\gamma_{(\text{NH}_4)_3\text{H}(\text{SO}_4)_2}=\left(\gamma_{(\text{NH}_4)_2\text{SO}_4}\cdot\gamma_{\text{NH}_4\text{HSO}_4}\right)^{0.2}$

² Species in bold are new in ISORROPIA II. 1926

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 5. Mutual deliquescence relative humidities, for the new salts modeled in ISORROPIA II³.

Salt Mixture	MDRH*
Ca(NO ₃) ₂ , CaCl ₂ , K ₂ SO ₄ , KNO ₃ , KCl, MgSO ₄ , Mg(NO ₃) ₂ , MgCl ₂ , <i>NaNO₃, NaCl, NH₄NO₃, NH₄Cl</i>	0.200
<i>(NH₄)₂SO₄, NH₄NO₃, NH₄Cl, Na₂SO₄, K₂SO₄, MgSO₄</i>	0.460
Ca(NO ₃) ₂ , K ₂ SO ₄ , KNO ₃ , KCl, MgSO ₄ , Mg(NO ₃) ₂ , MgCl ₂ , <i>NaNO₃, NaCl, NH₄NO₃, NH₄Cl</i>	0.240
<i>(NH₄)₂SO₄, NH₄Cl, Na₂SO₄, K₂SO₄, MgSO₄</i>	0.691
Ca(NO ₃) ₂ , K ₂ SO ₄ , KNO ₃ , KCl, MgSO ₄ , Mg(NO ₃) ₂ , <i>NaNO₃, NaCl, NH₄NO₃, NH₄Cl</i>	0.240
<i>(NH₄)₂SO₄, Na₂SO₄, K₂SO₄, MgSO₄</i>	0.697
K ₂ SO ₄ , MgSO ₄ , <i>KHSO₄, NH₄HSO₄, NaHSO₄, (NH₄)₂SO₄, Na₂SO₄, (NH₄)₃H(SO₄)₂</i>	0.240
<i>(NH₄)₂SO₄, NH₄NO₃, Na₂SO₄, K₂SO₄, MgSO₄</i>	0.494
K ₂ SO ₄ , KNO ₃ , KCl, MgSO ₄ , Mg(NO ₃) ₂ , <i>NaNO₃, NaCl, NH₄NO₃, NH₄Cl</i>	0.240
K ₂ SO ₄ , MgSO ₄ , KHSO ₄ , NaHSO ₄ , <i>(NH₄)₂SO₄, Na₂SO₄, (NH₄)₃H(SO₄)₂</i>	0.363
K ₂ SO ₄ , KNO ₃ , KCl, MgSO ₄ , <i>NaNO₃, NaCl, NH₄NO₃, NH₄Cl</i>	0.596
K ₂ SO ₄ , MgSO ₄ , KHSO ₄ , <i>(NH₄)₂SO₄, Na₂SO₄, (NH₄)₃H(SO₄)₂</i>	0.610
Ca(NO ₃) ₂ , K ₂ SO ₄ , KNO ₃ , KCl, MgSO ₄ , Mg(NO ₃) ₂ , <i>NaNO₃, NaCl, NH₄NO₃, NH₄Cl</i>	0.240
K ₂ SO ₄ , KNO ₃ , KCl, MgSO ₄ , Mg(NO ₃) ₂ , <i>NaNO₃, NaCl, NH₄NO₃, NH₄Cl</i>	0.240

* Obtained from Potukuchi and Wexler (1995a, b) for mixtures with closest composition (T=298.15 K).

³ Species in *italics* determine the mixture from which the MDRH value has been taken for each case.

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**ISORROPIA II: a new
aerosol equilibrium
model**

 C. Fountoukis and
A. Nenes

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)
Table 6. Coefficients of $m(a_w)$ from the polynomial fit $m(a_w) = k_0 + k_1 a_w + k_2 a_w^2 + \dots$

Species	k_0	k_1	k_2	k_3	k_4	k_5
$\text{Ca}(\text{NO}_3)_2^\clubsuit$	34.400	-155.360	438.210	-706.380	579.230	-190.270
CaCl_2^*	20.847	-97.599	273.220	-422.120	331.160	-105.450
CaSO_4	N/A	N/A	N/A	N/A	N/A	N/A
KHSO_4^\S	1.061	-0.101	1.579×10^{-2}	-1.950×10^{-3}	9.515×10^{-5}	-1.547×10^{-6}
$\text{K}_2\text{SO}_4^\clubsuit$	1061.51	-4748.97	8096.16	-6166.16	1757.47	0
KNO_3^*	1.2141×10^4	-5.1173×10^4	8.12524×10^4	-5.7527×10^4	1.5305×10^4	0
KCl^\clubsuit	179.721	-721.266	1161.03	-841.479	221.943	0
MgSO_4^\diamond	-0.778	177.740	-719.790	1174.600	-863.440	232.310
$\text{Mg}(\text{NO}_3)_2^\diamond$	12.166	-16.154	0	10.886	0	-6.815
MgCl_2^\diamond	11.505	-26.518	34.937	-19.829	0	0

\clubsuit The coefficients for $\text{Ca}(\text{NO}_3)_2$ given by Kim and Seinfeld (1995) contain a typographical error. The coefficients used here were obtained by fitting the equation $m(a_w) = k_0 + k_1 a_w + k_2 a_w^2 + k_3 a_w^3 + k_4 a_w^4 + k_5 a_w^5$ to their Fig. 15.

* source: Kim and Seinfeld (1995).

\clubsuit source: Kelly and Wexler (2006).

\S Same as NaHSO_4 .

\diamond source: Ha and Chan (1999).

**ISORROPIA II: a new
aerosol equilibrium
model**

 C. Fountoukis and
A. Nenes

Table 7. List of input conditions for model simulations^a.

Case	Aerosol Type	Na	H ₂ SO ₄	NH ₃	HNO ₃	HCl	Ca ²⁺	K ⁺	Mg ²⁺	R ₁ , R ₂ , R ₃
1	Urban (1)	0.000	10.000	3.400	2.000	0.000	0.400	0.330	0.000	2.14, 0.18, 0.18
2	Urban (2)	0.023	10.000	3.400	2.000	0.037	0.900	1.000	0.000	2.44, 0.48, 0.47
3	Urban (3)	0.000	15.000	2.000	10.000	0.000	0.900	1.000	0.000	1.27, 0.31, 0.32
4	Urban (4)	0.000	15.000	2.000	10.000	0.000	0.400	0.330	0.000	0.89, 0.12, 0.12
5	N-u Cont. ^b (1)	0.200	2.000	8.000	12.000	0.200	0.120	0.180	0.000	23.9, 0.80, 0.37
6	N-u Cont. (2)	0.100	4.000	10.000	7.000	0.100	0.120	0.180	0.050	14.8, 0.34, 0.24
7	N-u Cont. (3)	0.023	5.664	12.000	2.000	0.037	0.120	0.180	0.050	12.4, 0.18, 0.17
8	N-u Cont. (4)	0.023	5.664	20.400	0.611	0.037	0.120	0.180	0.000	20.9, 0.15, 0.13
9	Marine (1)	2.000	1.000	0.010	0.300	3.121	0.100	0.100	0.070	9.36, 9.30, 0.80
10	Marine (2)	1.500	1.000	0.010	1.500	2.500	0.360	0.450	0.050	8.66, 8.60, 2.21
11	Marine (3)	2.500	3.000	0.001	3.000	2.500	0.500	1.000	0.050	4.86, 4.86, 1.31
12	Marine (4)	3.000	3.000	0.020	2.000	3.121	0.360	0.450	0.130	5.14, 5.10, 0.84
13	Rem. Cont. ^b (1)	0.000	10.000	4.250	0.145	0.000	0.080	0.090	0.000	2.49, 0.04, 0.04
14	Rem. Cont. (2)	0.023	10.000	3.000	1.000	0.037	0.080	0.090	0.000	1.78, 0.05, 0.04
15	Rem. Cont. (3)	0.100	15.000	3.000	4.000	0.100	0.080	0.090	0.000	1.21, 0.06, 0.03
16	Rem. Cont. (4)	0.200	15.000	3.000	8.000	0.200	0.080	0.090	0.040	1.25, 0.10, 0.04

^a Simulations for each case were conducted for 10, 25, 40, 55, 65, 70, 75, 80, 85, 90 and 98% relative humidity. Temperature was set to 298.15 K. Concentration given in $\mu\text{g m}^{-3}$. ^b N-u Cont., non-urban continental; Rem. Cont., remote continental.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Table 8. Normalized mean errors between ISORROPIA II and SCAPE2 for the simulations in Table 7.

NME (%)	H₂O_(p)	NO_{3(p)}	Cl_(p)	NH_{4(p)}	Total PM	H⁺_(aq)
ISORROPIA-II (Stable)	13.5	16.5	6.5	2.1	13.0	64.9
ISORROPIA-II (Metastable)	14.7	23.7	6.6	6.7	14.3	68.0

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes

Table 9. CPU time required for the simulations in Table 7.

Aerosol Case	Convergence criterion	ISORROPIA II (stable) CPU time (msec)	SCAPE2 CPU time (msec)	$\frac{\text{CPU}_{\text{SCAPE2}}}{\text{CPU}_{\text{ISORROPIA-II}}}$	$\frac{\text{CPU}_{\text{ISORROPIA-II(stable)}}}{\text{CPU}_{\text{ISORROPIA-II(metastable)}}}$	$\frac{\text{CPU}_{\text{ISORROPIA-II}}}{\text{CPU}_{\text{ISORROPIA}}}$
Marine		30	50	16.7	1.16	1.0
Urban		20	210	10.5	1.09	1.0
Remote continental	10^{-3}	20	440	22.0	1.05	1.0
Non-urban continental		20	110	5.5	1.00	1.0
Marine		30	>1000	>1000	1.16	1.0
Urban		20	280	14.0	1.09	1.0
Remote continental	10^{-4}	20	>1000	>1000	1.05	1.0
Non-urban continental		20	420	21.0	1.00	1.0
Marine		30	>1000	>1000	1.16	1.0
Urban		20	>1000	>1000	1.09	1.0
Remote continental	10^{-5}	20	>1000	>1000	1.05	1.0
Non-urban continental		20	1250	62.50	1.00	1.0

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶


Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**ISORROPIA II: a new
aerosol equilibrium
model**C. Fountoukis and
A. Nenes**Fig. 1.** Generic solution procedure of ISORROPIA II.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**ISORROPIA II: a new
aerosol equilibrium
model**C. Fountoukis and
A. Nenes**Fig. 2.** DRH as a function of temperature for all ISORROPIA salts.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes




Fig. 3. Concentration of aerosol water (a), nitrate (b), chloride (c), ammonium (d), total PM (e), and hydrogen (f), as predicted by ISORROPIA II (thermodynamically stable solution) and SCAPE2 for all the conditions described in Table 7. Temperature is set to 298.15 K.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


**ISORROPIA II: a new
aerosol equilibrium
model**C. Fountoukis and
A. Nenes

Fig. 4. Same as Fig. 2 but using the metastable solution of ISORROPIA II.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


ISORROPIA II: a new aerosol equilibrium modelC. Fountoukis and
A. Nenes

Fig. 5. Concentration of aerosol water (a), aqueous potassium (b), aqueous ammonium (c), and aqueous nitrate (d) as predicted by ISORROPIA II (thermodynamically stable solution) and SCAPE2 for the urban (3) case (Table 7) corresponding to a sulfate rich aerosol behavior ($1 < R_1 < 2$). Temperature is set to 298.15 K.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ISORROPIA II: a new aerosol equilibrium model

C. Fountoukis and
A. Nenes




Fig. 6. Concentration of aerosol water (a), aqueous potassium (b), solid sodium chloride (c), and aqueous magnesium (d) as predicted by ISORROPIA II (thermodynamically stable solution) and SCAPE2 for the marine (4) case (Table 7) corresponding to a sulfate poor, crustals and sodium rich aerosol behavior ($R_1 > 2$, $R_2 > 2$). Temperature is set to 298.15 K.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion


ISORROPIA II: a new aerosol equilibrium modelC. Fountoukis and
A. Nenes

Fig. 7. Concentration of aerosol water (a), aqueous nitrate (b), and aqueous ammonium (c) as a function of relative humidity as predicted by ISORROPIA II (thermodynamically stable solution) and SCAPE2 for the non-urban continental (1) case (Table 7) corresponding to a sulfate poor, ammonium rich aerosol behavior ($R_1 > 2$, $R_2 < 2$). Temperature is set to 298.15 K.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


ISORROPIA II: a new aerosol equilibrium modelC. Fountoukis and
A. Nenes

Fig. 8. Concentration of aerosol water (a), and aqueous phase potassium (b) as a function of relative humidity as predicted by ISORROPIA II (using the thermodynamically stable and metastable solutions) and SCAPE2 for the remote continental (1) case (Table 7) corresponding to a sulfate near-neutral aerosol behavior. Temperature is set to 298.15 K.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)