

---

Interactive  
Comment

## ***Interactive comment on “The potential importance of frost flowers, recycling on snow, and open leads for Ozone Depletion Events” by M. Piot and R. von Glasow***

**M. Piot and R. von Glasow**

Received and published: 23 November 2007

### **Reply Part 2**

Technical suggestions made by the reviewer:

Reply: Thanks for the technical suggestions, we have followed most of them.

P4532, L17: “OL provide a source region for sea salt aerosols” Is this true? I would have thought that the associated fetch associated with a small Polyna or OL would not generate large enough waves for aerosol generation.”

Reply: This is an interesting question. In general, the sea salt aerosol production from an open lead depends on the size of the lead, the wind stress at the water surface, and

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

EGU

Interactive  
Comment

the morphology (presence of frazil ice, etc). Model simulations from [Gong et al.(1997)] and discussions from [Rankin et al.(2000)] suggest that open leads do contribute to the sea salt aerosol production in the Arctic. We estimate that the leads modeled in our simulations are large enough (several kilometers) to induce this production.

“General: Could the FF aerosols be replaced with blown snow with deposited bromide and still have the same impact?”

Reply: Yes. Please, see p. 4532, l. 11-13

“We don’t think that leapfrogging is the best description of the process of bromine movement.”

Reply: We do not see a difference between the terms “hopping” and “leapfrogging” and prefer our choice.

“if we understand correctly, all the calculations do not assume an initial presence of bromine along the trajectory”

Reply: We assume the reviewer meant “an initial presence of bromine in the snow-pack”. We discuss this in detail in our first “Reply” to the first reviewer.

“What is the initial temperature profile?”

Reply: The initial temperature profiles are shown in Fig. 1:

However, we do not want to add it to the paper. We believe the change from  $T$  to  $\theta$  profile in Fig. 2 (see “reply” to comment 3) is sufficient to better understand the temperature conditions in the model.

P4538, L6: “ID(Br) as defined, although accumulated deposition, this is not a local quantity as the air parcel continues to move. Rather it is the integral of deposition along the path. This should be made clearer” Reply: Done

P4542, L17: “I estimate the ozone loss due to surface deposition is about 10 ppbv in

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

---

Interactive  
Comment

4 days . Or assuming a 100 m MBL would imply a VD of about 0.02 cm/s. If the MBL remains at about 100 m this seems rather rapid as it would remove the ozone in a few weeks whereas 40 ppbv can persist in transit through the Arctic for a few weeks with no perceptible depletion.”

Reply: We have compared the modeled ozone deposition velocity with measured velocities over packed ice from [Strunin et al.(1997)]. In MISTRA the average deposition velocity is  $V_{O_3} = 0.04\text{--}0.05 \text{ cm s}^{-1}$  while it is found to be  $0.07 \text{ cm s}^{-1}$  by Strunin et al. In addition, recent observations [Peterson and Honrath(2001), Albert et al.(2002)] suggest that the ozone deposition on snow may be more important than believed in the past see [Bottenheim et al.(2002)]. Note that the polar boundary layer has an initial height of 300 m in the model, not 100 m, but it seems that the calculations made by the reviewer assume  $z=300 \text{ m}$  rather than 100. Also, the PBL has often been measured at heights above 500 m [Bottenheim et al.(2002)], which significantly reduces the loss rate on snow. We have included the reference of [Strunin et al.(1997)] into the paper, p. 4541, l. 20

“In general: OL. The size of the leads used sound rather large - what sort of statistics are available? Also what are the aspect ratios of the leads? Surely the wind doesn't always blow across the leads?”

Reply: We included in the term “open lead” open leads and polynyas. Polynyas range in size from 10 to  $10^5 \text{ km}^2$  [Morales Maqueda et al.(2004)]. Open leads are long, narrow ice openings. They are meters to kilometers wide and kilometers to tens of kilometers long; see, e.g., [Smith et al.(1990)]. As mentioned by the reviewer the wind does not always blow across a lead. Blowing in a direction different than  $90^\circ$  from the axis of a lead would increase the surface influence on the air parcel crossing it. Nevertheless, the extent of the modeled leads are based on a rather arbitrary choice.

“What about ubiquitous ice crystals?”

Reply: We believe the reviewer is referring to airborne ice crystals. Indeed,

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

these crystals provide an adequate surface for heterogeneous reactions; see [Adams et al.(2002)]. However, our model does not include reactions on frozen surfaces. Therefore, we cannot assess the relative importance of reactions on ice in comparison with reactions in/on liquid surfaces.

P4554: "Role of acids? SO<sub>2</sub> sources in Russia in the high Arctic appear to have been decreasing. This is likely to have affected the general acidity of aerosols over the last 20 years. Has it impacted ODEs?"

Reply: This is a very good question but it is beyond the scope of our paper and cannot be answered with certitude. The presented model results show that acidity (e.g., SO<sub>2</sub> or Arctic haze) impacts the strength of ODEs. Should the acidity have indeed decreased over the last 20 years, it seems very likely that reactions requiring H<sup>+</sup> to proceed would be reduced. The strength of the bromine explosion, e.g., is likely to be affected by less acidic transported pollutions.

## References

[Abbatt(1994)] Abbatt, J. P. D.: Heterogeneous reaction of HOBr with HBr and HCl on ice surfaces at 228K, Geophys. Res. Lett., 21 (8), 665–668, 1994.

[Adams et al.(2002)] Adams, J. W., Holmes, N. S., and Crowley, J. N.: Uptake and Reaction of HOBr on Frozen and dry Salt Surfaces, Atmos. Chem. Phys., 2, 79–91, 2002.

[Albert et al.(2002)] Albert, M. R., Grannas, A. M., Bottenheim, J., Shepson, P. B., and Perron, F. E.: Processes and properties of snow-air transfer in the high Arctic with applications to interstitial ozone at Alert, Canada, Atmos. Environ., 36, 2779–2787, 2002.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

[Argentini et al.(2003)] Argentini, S., Viola, A. P., Mastrantonio, G., Maurizi, A., Georgiadis, T., and Nardino, M.: Characteristics of the boundary layer at Ny-Ålesund in the Arctic during the ARTIST field experiment, *Annals Geophys.*, 46, 185–195, 2003.

[Bottenheim et al.(2002)] Bottenheim, J. W., Fuentes, J. D., Tarasick, D. W., and Anlauf, K. G.: Ozone in the Arctic Lower Troposphere During Winter and Spring 2000 (ALERT2000), *Atmos. Environ.*, 36, 2535–2544, 2002.

[Dominé et al.(2005)] Dominé, F., Taillandier, A. S., Simpson, W. R., and Severin, K.: Specific surface area, density and microstructure of frost flowers, *Geophys. Res. Lett.*, 32, L13502, doi:10.1029/2005GL023245, 2005.

[Fickert et al.(1999)] Fickert, S., Adams, J. W., and Crowley, J. N.: Activation of Br<sub>2</sub> and BrCl via uptake of HOBr onto aqueous salt solutions, *J. Geophys. Res.*, 104 (D19), 23 719–23 727, doi:10.1029/1999JD900359, 1999.

[Foster et al.(2001)] Foster, K. L., Plastridge, R. A., Bottenheim, J. W., Shepson, P. B., Finlayson-Pitts, B. J., and Spicer, C. W.: The Role of Br<sub>2</sub> and BrCl in Surface Ozone Destruction at Polar Sunrise, *Science*, 291, 471–474, doi:10.1126/science.291.5503.471, 2001.

[Georgiadis et al.(2000)] Georgiadis, T., Bonafé, U., Calzolari, F., Nardino, M., Orsini, A., Pirazzini, R., Ravagnani, F., Sozzi, R., Trivellone, G., and Argentini, S.: Study of the surface energy balance at Ny-Ålesund, Svalbard, Conference Proceedings-Italian Physical Society, 69, 163–174, 2000.

[Gong et al.(1997)] Gong, S. L., Barrie, L. A., Prospero, J. M., Savoie, D. L., Ayers, G. P., Blanchet, J.-P., and Spacek, L.: Modeling sea-salt aerosols in the atmosphere: 2. Atmospheric concentrations and fluxes, *J. Geophys. Res.*, 102, 3819–3830, 1997.

[Grannas et al.(2002)] Grannas, A. M., Shepson, P. B., Guimbaud, C., Sumner, A. L., Albert, M., Simpson, W., Dominé, F., Boudries, H., Bottenheim, J., Beine, H. J., Hon-

rath, R., and Zhou, X.: A study of photochemical and physical processes affecting carbonyl compounds in the Arctic atmospheric boundary layer, *Atmos. Environ.*, 36, 2733–2742, 2002.

ACPD

7, S7206–S7216, 2007

[Hafskjold et al.(1993)] Hafskjold, B., Ikeshojit, T., and Ratkjes, S. K.: On the molecular mechanism of thermal diffusion in liquids, *Mol. Phys.*, 80 (6), 1389–1412, 1993.

Interactive  
Comment

[Hansen et al.(1999)] Hansen, J. C., Li, Y., Francisco, J. S., and Li, Z.: On the Mechanism of the BrO + CH<sub>2</sub>O Reaction, *J. Phys. Chem. A*, 103, 8543–8546, 1999.

[Hartmann et al.(1999)] Hartmann, J., Albers, F., Argentini, S., Borchert, A., Bonafé, U., Cohrs, W., Conidi, A., Freese, D., Georgiadis, T., Ippoliti, A., Kaleschke, L., Lüpkes, C., Maixner, U., Mastrantonio, G., Ravagnani, F., Reuter, A., Trivellone, G., and Viola, A.: Reports on Polar Research. Arctic Radiation and Turbulence Interaction Study (ARTIST), Alfred-Wegener-Institute for Polar and Marine Res., Bremerhaven, Germany, 1999.

[Hoff and Trivett(1984)] Hoff, R. M. and Trivett, N. B. A.: Ground-based measurements of Arctic haze made at Alert, N.W.T., Canada, during the Arctic Gas and Aerosol Sampling Project (AGASP), *Geophys. Res. Lett.*, 11, 389–392, 1984.

[Hollwedel(2005)] Hollwedel, J. C.: Observations of Tropospheric and Stratospheric Bromine Monoxide from Satellite, Ph.D. thesis, Universität Heidelberg, 2005.

[Huff and Abbatt(2000)] Huff, A. K. and Abbatt, J. P. D.: Gas-Phase Br<sub>2</sub> Production in Heterogeneous Reactions of Cl<sub>2</sub>, HOCl, and BrCl with Halide-Ice Surfaces, *J. Phys. Chem. A*, 104, 7284–7293, 2000.

[Kalnajs and Avallone(2006)] Kalnajs, L. E. and Avallone, L. M.: Frost flower influence on springtime boundary-layer ozone depletion events and atmospheric bromine levels, *Geophys. Res. Lett.*, 33, 10 810–+, 10.1029/2006GL025809, 2006.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

---

Interactive  
Comment

[Keil and Shepson(2006)] Keil, A. D. and Shepson, P. B.: Chlorine and bromine atom ratios in the springtime Arctic troposphere as determined from measurements of halogenated volatile organic compounds, *J. Geophys. Res.*, 111, D17303, doi:10.1029/2006JD007119, 2006.

[Kempers(1989)] Kempers, L. J. T. M.: A thermodynamic theory of the Soret effect in a multicomponent liquid, *J. Chem. Phys.*, 90 (11), 6541–6548, 1989.

[Koop et al.(2000)] Koop, T., Kapilashrami, A., Molina, L. T., and Molina, M. J.: Phase transitions of sea-salt/water mixtures at low temperatures: Implications for ozone chemistry in the polar marine boundary layer, *J. Geophys. Res.*, 105 (D21), 26 393–26 402, doi:2000JD900413, 2000.

[Landgraf and Crutzen(1998)] Landgraf, J. and Crutzen, P.: An Efficient Method for 'On-Line' Calculations of Photolysis and Heating Rates, *J. Atmos. Sci.*, 55, 863–878, 1998.

[Li and Winchester(1989)] Li, S.-H. and Winchester, J. W.: Resolution of ionic components of late winter Arctic aerosols, *Atmos. Environ.*, 23 (11), 2387–2399, 1989.

[McElroy et al.(1999)] McElroy, C. T., McLinden, C. A., and McConnell, J. C.: Evidence for bromine monoxide in the free troposphere during the Arctic polar sunrise, *Nature*, 397, 338–340, 1999.

[Morales Maqueda et al.(2004)] Morales Maqueda, M. A., Willmott, A. J., and Biggs, N. R. T.: Polynyas dynamics: A review of observations and modeling, *Review of Geophysics*, 42, 1–37, 2004.

[Morin et al.(2005)] Morin, S., Hoenninger, G., Staebler, R. M., and Bottenheim, J.: A high time resolution study of boundary layer ozone chemistry and dynamics over the Arctic ocean near Alert, Nunavut, *Geophys. Res. Lett.*, 32, doi:10.1029/GL2004022098, 2005.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

Interactive  
Comment

[Orlando et al.(2000)] Orlando, J. J., Ramacher, B., and Tyndall, G. S.: Upper Limits for the Rate Coefficients for Reactions of BrO with Formaldehyde and HBr, *Geophys. Res. Lett.*, 27, 2633–2636, doi: 10.1029/2000GL011725, 2000.

[Oum et al.(1998)] Oum, K. W., Lakin, M. J., and Finlayson-Pitts, B. J.: Bromine activation in the troposphere by the dark reaction of O<sub>3</sub> with seawater ice, *Geophys. Res. Lett.*, 25 (21), 3923–3926, 1998.

[Peterson and Honrath(2001)] Peterson, M. C. and Honrath, R. E.: Observations of Rapid Photochemical Destruction of Ozone in Snowpack Interstitial Air, *Geophys. Res. Lett.*, 28 (3), 511–514, doi:2000GL012129, 2001.

[Rankin et al.(2000)] Rankin, A. M., Auld, V., and Wolff, E. W.: Frost flowers as a source of fractionated sea salt aerosol in the polar regions, *Geophys. Res. Lett.*, 27 (21), 3469–3472, doi:2000GL011771, 2000.

[Sander et al.(1997)] Sander, R., Vogt, R., Harris, G. W., and Crutzen, P. J.: Modeling the chemistry of ozone, halogen compounds, and hydrocarbons in the arctic troposphere during spring, *Tellus*, 49B (5), 522–532, 1997.

[Schnell et al.(1989)] Schnell, R. C., Barry, R. G., Miles, M. W., Andreas, E. L., Radke, L. F., Brock, C. A., McCormick, M. P., and Moore, J. L.: Lidar detection of leads in Arctic sea ice, *Nature*, 339, 530–532, doi:10.1038/339530a0, 1989.

[Serreze et al.(1992)] Serreze, M. C., Kahl, J. D., and Schnell, R. C.: Low-level temperature inversions of the Eurasian Arctic and comparisons with Soviet drifting station data, *J. Climate*, 5 (6), 1992.

[Simpson et al.(2005)] Simpson, W. R., Alvarez-Aviles, L., Douglas, T. A., Sturm, M., and Dominé, F.: Halogens in the coastal snow pack near Barrow, Alaska: Evidence for active bromine air-snow chemistry during springtime, *Geophys. Res. Lett.*, 32, L04811, doi: 10.1029/2004GL021748, 2005.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Interactive  
Comment

[Simpson et al.(2007)] Simpson, W. R., von Glasow, R., Riedel, K., Anderson, P., Ariya, P., Bottenheim, J., Burrows, J., Carpenter, L., Frieß, U., Goodsite, M. E., Heard, D., Hutterli, M., Jacobi, H.-W., Kaleschke, L., Neff, B., Plane, J., Platt, U., Richter, A., Roscoe, H., Sander, R., Shepson, P., Sodeau, J., Steffen, A., Wagner, T., and Wolff, E.: Halogens and their role in polar boundary-layer ozone depletion, *Atmos. Chem. Phys.*, 7, 4375–4418, 2007.

[Smith et al.(1990)] Smith, S. D., Muench, R. D., and Pease, C. H.: Polynyas and leads: An overview of physical processes and environment, *J. Geophys. Res.*, 95, 1990.

[Spicer et al.(2002)] Spicer, C. W., Plastridge, R. A., Foster, K. L., Finlayson-Pitts, B. J., Bottenheim, J. W., Grannas, A. M., and Shepson, P. B.: Molecular halogens before and during ozone depletion events in the Arctic at polar sunrise: concentrations and sources, *Atmos. Environ.*, 36 (15-16), 2721–2731, doi:10.1016/S1352-2310(02)00125-5, 2002.

[Strunin et al.(1997)] Strunin, M. A., Postnov, A. A., and Mezrin, M. Y.: Meteorological potential for contamination of arctic troposphere: Boundary layer structure and turbulent diffusion characteristics, *Atmos. Res.*, 44, 37–51, 1997.

[Timm and Janek(2005)] Timm, H. and Janek, J.: On the Soret effect in binary nonstoichiometric oxides - kinetic demixing of cuprite in a temperature gradient, *Solid State Ionics*, 176 (11-12), 1131–1143, doi:10.1016/j.ssi.2005.01.010, 2005.

[Toyota et al.(2004)] Toyota, K., Kanaya, Y., Takahashi, M., and Akimoto, H.: A box model study on photochemical interactions between VOCs and reactive halogen species in the marine boundary layer, *Atmos. Chem. Phys.*, 4, 1961–1987, 2004.

[von Glasow et al.(2002a)] von Glasow, R., Sander, R., Bott, A., and Crutzen, P. J.: Modeling halogen chemistry in the marine boundary layer. 2. Interactions

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

with sulfur and cloud-covered MBL, *J. Geophys. Res.*, 107 (D17), 4323, doi: 10.1029/2001JD000943, 2002a.

[von Glasow et al.(2002b)] von Glasow, R., Sander, R., Bott, A., and Crutzen, P. J.: Modeling halogen chemistry in the marine boundary layer 1. Cloud-free MBL, *J. Geophys. Res.*, 107 (D17), 4341, doi:10.1029/2001JD000942, 2002b.

[Wesely(1989)] Wesely, M. L.: Parameterization of surface resistances to gaseous deposition in regional-scale numerical models, *Atmos. Environ.*, 23, 1293–1304, 1989.

---

Interactive comment on *Atmos. Chem. Phys. Discuss.*, 7, 4521, 2007.

ACPD

7, S7206–S7216, 2007

---

Interactive  
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

Interactive  
Comment




Figure 1: Initial temperature and potential temperature profiles at the start of the one-dimensional model runs in MISTRAL.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper