

This discussion paper is/has been under review for the journal *Atmospheric Chemistry and Physics (ACP)*. Please refer to the corresponding final paper in *ACP* if available.

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Spatial heterogeneity of satellite derived land surface parameters and energy flux densities for LITFASS-area

A. Tittebrand^{1,*} and F. H. Berger²

¹Institute for Hydrology and Meteorology, Dresden, Germany

²Deutscher Wetterdienst, Meteorologisches Observatorium Lindenberg, Germany

* now at: Institute for Oceanography, Hamburg, Germany

Received: 6 May 2008 – Accepted: 2 August 2008 – Published: 26 August 2008

Correspondence to: A. Tittebrand (antje.tittebrand@zmaw.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Remote sensing data provide area integrated information of surface properties in different spatial or temporal resolutions according to different sensor features. Landsat ETM+, Terra MODIS and NOAA-AVHRR surface temperature and spectral reflectance were used to infer further surface parameters and radiant- and energy flux densities for LITFASS-area, a $20 \times 20 \text{ km}^2$ heterogeneous area in Eastern Germany, mainly characterized by the land use types forest, crop, grass and water. Based on the Penman-Monteith-approach the actual latent heat flux ($L.E$), as key quantity of the hydrological cycle, is determined for each sensor in the accordant spatial resolution with an improved parametrization. However, using three sensors, significant discrepancies between the inferred parameters can cause flux distinctions resultant from differences of the sensor filter response functions or atmospheric correction methods. The approximation of MODIS- and AVHRR- derived surface parameters to the reference parameters of ETM (via regression lines and histogram stretching, respectively), further the use of accurate land use classifications (CORINE and a new Landsat-classification), and a consistent parametrization for the three sensors were realized to obtain a uniform base for investigations of the spatial variability. For the target area the spatial heterogeneity is analysed investigating frequency distribution functions (PDF) for surface parameters and energy fluxes. PDF is the most promising way to describe subgrid heterogeneity due to the given data in different spatial resolution. Aim of this study is to find typical distribution pattern of parameters (albedo, NDVI) for the determination of $L.E$ determined from the highly resolved ETM data within pixel on coarser scale (MODIS, AVHRR). The analyses for 4 scenes in 2002 and 2003 showed that clear distribution-pattern for forest for NDVI and albedo are found. Grass and crop distributions show higher variability and differ significantly to each other in NDVI but only marginal in albedo. Regarding NDVI-distribution functions NDVI was found to be the key variable for $L.E$ -determination.

ACPD

8, 16219–16254, 2008

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

1 Introduction

The land surface interface is a key component of the climate system since it provides the coupling between the atmosphere and the land biosphere and hydrology (Giorgi, 1996). Up to the present remote sensing is the only observing method for mapping land surface characteristics or energy fluxes globally (Liang et al., 2002) with their known uncertainties.

The land surface variables and vegetation variables, such as surface temperature, surface reflectance, Normalised Difference Vegetation Index (NDVI), Leaf Area Index (LAI) and Soil Adjusted Vegetation Index (SAVI) can be derived directly from satellite observations (e.g. Wan and Dozier, 1989; Becker and Li, 1990; Li and Becker, 1993; Ma et al., 2006). The regional radiant and energy flux densities can be inferred only indirectly with the aid of these land surface variables and further vegetation parameters (Pinker, 1990).

The theoretical framework of a satellite based determination of radiant and energy flux densities with focus to the latent heat flux ($L.E$) based on Penman-Monteith for LITFASS-area (Lindenberg Inhomogeneous Terrain: fluxes between the Atmosphere and the Surface – a long term Study) was given in previous studies (Berger, 2001; Tittebrand et al., 2005) and is presented here with an improved parametrization for Landsat 7 ETM+ and NOAA 16-AVHRR 3 data and adapted for Terra MODIS data.

Parameter and flux distinctions between the sensors result from differences of the sensor filter response functions, correction methods of the atmospheric effect (van Leeuwen et al., 1999; Teillet et al., 1997) or they are based on different or outdated land use classifications (e.g. USGS land cover). Thus, it was necessary to realize comparable land surface parameters to allow a consistent parametrization scheme for the three sensors. An adaptation method for MODIS (albedo, NDVI and surface temperature) and AVHRR (NDVI) was carried out using the accordant ETM parameters as reference data. An improved and consistent parametrization for the three sensors led to satisfying findings confirmed by new validation results. The second source of

ACPD

8, 16219–16254, 2008

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



unmeant variability of the resultant fluxes arises from the land use classifications. Thus, after comparison to an in-situ land use classification (Beyrich et al., 2004) for LITFASS-area, the Coordinated Information on the Environment classification (CORINE) and a new Landsat-classification (Prechtel, 2005) were used for the study, accordant to their spatial resolution for AVHRR (1 km), MODIS (1 km) and ETM (30 m), respectively.

However, the monitoring of vegetation requires data of high temporal resolution which are up to now only provided by sensors with moderate spatial resolution. Their spatial resolution ranges from a few hundred meters (Envisat MERIS, Terra MODIS) up to one kilometer (NOAA AVHRR). While using such moderate resolution data, the heterogeneity of the observed surface could get lost mainly because the landscape is a mosaic of objects such as agricultural fields or vegetation patches that are often smaller than the pixel size at a moderate resolution. So the spatial heterogeneity on an intra-pixel scale will be lost (Garrigues et al., 2006). Thus, high spatially resolved data (e.g. Landsat ETM) are fine enough to describe sub-grid heterogeneity, but at the expense of a coarser repeat cycle of the satellite.

Numerous studies describe different ways of considering sub-grid scale land-surface heterogeneities especially in atmospheric models (Arain et al., 1996; Avissar, 1992; Chebouni et al., 1995; Koster and Suarez, 1992a, b; Lhomme et al., 1994; Li and Avissar, 1994; Noilhan et al., 1997; Sellers et al., 1997; Shuttleworth et al., 1997). Basically, three methods have been developed: The usage of effective parameters (Arain et al., 1996; Lhomme, 1992; Claussen, 1995a) have been proposed as a means for applying equations developed for homogeneous or point-scale processes, to processes occurring at larger spatial scales over heterogeneous surfaces. One method for deriving expressions for the effective parameters is to express the energy balance equation separately in terms of effective and component surface parameters, and then to match the expression term by term using some initial assumption about which fluxes will be “conserved” in going from the small to the large scale (Njoku et al., 1996). The “tile” approach (Li and Avissar, 1994) describes the heterogeneity inside a model grid cell in terms of a number of homogeneous “tiles” representing the dominant vegetation

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

and soil types. The grid fluxes are the averages of the tile fluxes weighted by their fractional area. For the “mosaic” approach (Koster and Suarez, 1992a; Mölders and Raabe, 1996) the model grid cell is subdivided into a regular smaller high-resolution grid. While the coarse grid for the host model is preserved, the SVAT scheme is used for the smaller grid. The averages of the sub-grid fluxes represent the fluxes on the coarse grid. In a SVAT scheme a number of parameters are used to describe the soil and vegetation characteristics inside the grid cell. These parameters are usually selected in correspondence to the most common vegetation and soil types within the grid square.

The usage of frequency distributions or generalized to probability density functions (PDF) offers another possibility to describe the variability of land-surface characteristics, at a resolvable scale of interest (Li and Avissar, 1994). This statistical-dynamical approach (Entekhabi and Eagleson, 1989; Avissar, 1992; Bonan et al., 1993; Giorgi 1996a, b) is needed because of the nonlinear relationship between most land-surface characteristics and the land surface heat fluxes. Therefore PDF could be used to describe spatial variability of a specific surface parameter for a satellite pixel or a model grid with coarser spatial resolution that is described by a higher resolved data set (MAGIM, 2006). The PDF can replace the use of effective parameters or other specific averaging strategies. Furthermore it considers the extreme values as well as the mode which can be completely different from the simple mean. This method was used by Giorgi (1996a, b) with the PDF of soil moisture and surface temperature to describe the representation of surface heterogeneity to be used in climate models. Li and Avissar (1994) demonstrated that the results of determined fluxes depend on the shape of the spatial distribution of the land surface parameters, used for the calculation. Typically, the more skewed the distribution within the range of values the larger the error (for nonlinear relationships) of the flux calculated using the mean instead of the distribution. Investigations in our study focus especially on the determination of latent heat and the distribution pattern of the therefore needed parameters (e.g. albedo, NDVI).

Here the spatial heterogeneity was analysed for parameters and fluxes in their given

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



spatial resolution, as well as averaged for comparison (using arithmetic mean and the dominant land use method) and enhanced with the characterisation of their frequency distributions

- (a) for the entire target area,
- (b) regarded for several land use types and
- (c) on pixel base.

Based on this work, typical pattern for the use of the PDF-method for the determination of $L.E$ (using the Penman-Monteith approach) could be provided determined from the highly resolved ETM data within pixel on coarser scale (MODIS, AVHRR).

2 Target area and data

2.1 Target area

The study focuses on a 20 km×20 km area around the Richard Aßmann Observatory (MOL-RAO) of the German Meteorological Service (Deutscher Wetterdienst, DWD) situated in North Eastern Germany 50 km Southeast of Berlin. The land use in this area is dominated by forest and agricultural fields with approximately 45% each, lake coverage of 7% and villages cover nearly 4%. The forest is dominated in the Western part of the area while agriculture is mainly situated in the Eastern part (Fig. 1). The target area is formed by inland glaciers, exhibiting a slightly surface with differences of only 80–100 m over distances of about 10–15 km with a number of small and medium-sized lakes (Mengelkamp et al., 2006)

2.2 Satellite data and time

The investigation is carried out with Landsat-7 ETM+ (Enhanced Thematic Mapper), Terra MODIS (Moderate Resolution Imaging Spectroradiometer) and NOAA 16 AVHRR

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 (Advanced Very High Resolution Radiometer) data. Landsat-7 ETM+ data, provided as Level 1B data, offer a spatial resolution of 30 m with a 16 day repeat cycle. The determination of brightness temperature and spectral reflectance on top of atmosphere (TOA) was applied after the Landsat User Handbook (2004).

5 From MODIS surface temperature and reflectance products from the Earth Observing System Data Gateway were used (EOS, 2007) for the determination of NDVI as well as for further parameters for the calculation of the energy fluxes. The reflectance product MOD09GHK was estimated for each band to produce a measurement equivalent to a ground-level measurement with no atmospheric scattering or absorption. The correction scheme identifies atmospheric gases, aerosols and thin cirrus clouds. These
10 MOD09GHK data are provided as a grid-level-2G, 500 m product in the sinusoidal projection. MODA11 temperature data were used, also given as a gridded product with a 1 km spatial resolution and corrected for the atmospheric effect.

In former studies of the institute NOAA 16-AVHRR 3 data were processed with the modularic scheme SESAT (Strahlungs- und Energieflüsse aus Satellitendaten, Berger, 2001; Tittebrand et al., 2005). For a comparable determination of the energy fluxes according to ETM and MODIS in this work only surface temperature (after Mc Clain, 1985; split window method), spectral reflectance and TOA-NDVI (Top of Atmosphere) of SESAT were used, given with a 1 km spatial resolution. For this case study four
20 cloud free scenes in spring and summer 2002 and 2003 were used: 9 May 2002, 28 July 2002, 20 August 2002 and 17 April 2003.

For evaluation purposes a field experiment (Beyrich et al., 2004) was planned and realized in May until June 2003. At that time two cloud free Landsat-overpasses were recorded. Unfortunately at the 31 May 2003 problems with Landsat's scan line corrector occurred (Landsat, 2003) thus these scenes were not available for the analysis. Only the cloud free scene a couple of weeks earlier (17 April 2003) could be used. Spring and summer 2003 were dry in contrast to the summer 2002, especially in August 2002, where the scene was taken only a few days after the flash-flood in Eastern Germany. So it was of interest to determine the latent heat flux under various soil moisture and
25

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



atmospheric conditions. For comparison half-hourly in-situ measurements of the routine observations in Lindenberg (Beyrich et al., 2004) were used but with absence of data of the turbulent fluxes except for the April scene in 2003.

2.3 Land use classifications

5 To use the best appropriate land use classification two were selected each for a fixed resolution:

- 1 km: CORINE (Coordinated Information on the Environment) is a consistent, comparable land-use and land-cover data base for the European environmental policy. To meet the requirements a satellite based (Landsat-7 ETM+) pan-European initial data acquisition of land use/cover types has been carried out.

- 25 m: Landsat-classification 2002 was developed by the Institute of Cartography of the Technical University of Dresden based on two Landsat 7 ETM+ scenes of Saxony in Germany to support the satellite retrieval algorithm for parameter and flux determination.

15 A comparison of the percentage representation of the main land uses for LITFASS area is shown in Table 1 for the land use classifications compared to field data. Further information to these classifications is given in the appendix.

For the comparison of land use classifications consistent classes had to be found. Therefore classes containing for example different field fruits were summarized to one class: crop. Nevertheless there are bigger differences between not only the percental representation of each class but also for the classification. A good agreement can be found for the water classes compared to the reference classification. Actually in LITFASS-area there is no deciduous forest (Fig. 1). Both classifications match this well and give coniferous forest as dominant land use type for high vegetation. Nevertheless, regarding Table 1, coniferous forest is underestimated for CORINE and Landsat-classification with differences of 14% and 7% respectively compared to the reference.

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Most uncertainties can be outlined for crop and grass, probably because of their similar spectral behaviour. For both land use types their differences are higher to each other than to the reference classification which possesses values in between. As mentioned above and outlined in Keil et al. (2005) CORINE tends to overestimate agricultural areas. Compared to the reference classification of LITFASS-area (Beyrich et al., 2004) CORINE and the Landsat-classification showed nearly comparable results hence applying them to the satellite data due to their spatial resolution: CORINE for MODIS and AVHRR and the Landsat-classification for ETM data. For flux determination the classifications were reduced to four classes: forest, grass, crop and water. Remaining classes were assigned to one of the four classes, as e.g. urban areas to the forest class because of the comparable roughness length (Beyrich et al., 2004).

3 Satellite derived fluxes and adaptation due to sensor differences

3.1 Satellite based parameters

To solve the Penman-Monteith equation the only satellite based quantities were surface spectral reflectance (and hence broadband albedo), NDVI and surface temperature.

The ETM-broadband albedo was inferred using channels 1, 3, 4, 5 and 7 after Liang et al. (2002) after the correction of atmospheric effects using 6S (Vermote et al., 1995). Surface temperature was determined after Sobrino et al. (2004) using a single windows channel method for channel 6. The $NDVI = (\rho_{NIR} - \rho_{VIS}) / (\rho_{NIR} + \rho_{VIS})$ for ETM is based on reflectance in channel 3 and in channel 4.

MODIS-broadband albedo was consistently computed after Liang et al. (2002) using channels 1 to 5 and 7. NDVI can be derived combining channel 1 and 2. For determination of the MODIS-fluxes the spectral reflectance data were averaged to the 1 km (interpolated 800 m) resolution allowing the combination of reflectance and temperature (both products provided), NDVI and surface temperature derived as mentioned above are the base for the determination of further parameters (e.g. LAI) and their

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



resulting energy fluxes.

For AVHRR albedo and TOA-NDVI and surface temperature were used from the SESAT scheme (Berger, 2001) as base for energy fluxes.

3.2 Determination of surface fluxes

- 5 As described in Tittebrand et al. (2005) the Penman-Monteith method was used to infer the actual latent heat fluxes on satellite-pixel base for the LITFASS-area assuming constant boundary layer conditions. Therefore the difference of net radiation (R_n) and soil heat flux at the surface (G), to be termed available energy, and a couple of further parameters (LAI, roughness length, wind speed, relative humidity and others) are
10 needed. The approach is given as follows:

$$L.E_{PM} = \frac{s}{s + \gamma \left(1 + \frac{r_c}{r_a}\right)} (R_n - G) + \frac{1}{s + \gamma \left(1 + \frac{r_c}{r_a}\right)} \frac{\rho c_p}{r_a} (e_s(T_{2m}) - e(T_{2m})) \quad (1)$$

- here $R_n - G$ is the available energy from net radiation R_n and surface soil heat flux G , s is the slope of the saturation vapor versus temperature curve, ρ the air density in kg m^{-3} , c_p the specific heat of air at constant pressure, γ the psychrometric constant,
15 and $e_s(T_{2m}) - e(T_{2m})$ the saturation deficit (e_s according to Magnus-Equation, Sonntag, 1982). The temperature in 2 m height is determined from $T_{\text{sur}} - T_{2m} = G/A_G$ after De Rooy and Holtslag (1999) with $A_G = 9 \text{ W m}^{-2} \text{ K}^{-1}$. The aerodynamic resistance r_a is derived from the logarithmic wind profile:

$$r_a = \frac{0.74}{k^2 u_{2m}} \left(\ln \frac{(z-d)}{z_0} \right)^2 \quad (2)$$

- 20 with the Karman constant $k=0.4$, $z-d$ the measurement height above the zero plane displacement, z_0 the roughness length and u_{2m} the wind speed at 2 m. Because these quantities are not available for each single pixel the parameterisation of r_a is realized with land use dependent values for z_0 and vegetation height according to

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hagemann (2002). Wind speed and relative humidity were included from in-situ tower measurements of the Observatory Lindenberg (MOL-RAO) for the respective time of overpass. In case of no data the boundary layer conditions are assumed to be constant (relative humidity $r_H=55\%$, visibility=23 km, wind speed=3 m/s) and they are used for the entire area.

The canopy resistance r_c is given by Allen (1986) and Allen et al. (1998):

$$r_c = \frac{r_s}{LAI_{\text{active}}} \quad (3)$$

with LAI_{active} the fraction of leaf area which actively contributes to vapour transport and here defined as $LAI_{\text{active}}=0.5 \text{ LAI}$.

The stomatal resistance r_s is a plant specific parameter depending to some extend on light intensity, temperature, leaf-air vapor pressure difference and ambient CO_2 concentration. Canopy resistance describes the resistance of water vapor transport through the entire transpiring vegetation canopy. For lower vegetation without water stress Allen et al. (1998) give a value of about 100 s m^{-1} . Minimum r_s of forest stands show quite a range. Kelliher et al. (1995) give as maximum leaf stomatal conductance $5.7 \pm 2.4 \text{ mm s}^{-1}$ (equivalent to $r_{s\text{min}}=123 \dots 303 \text{ s m}^{-1}$) for conifer forest and $4.6 \pm 1.7 \text{ mm s}^{-1}$ (equivalent to $r_{s\text{min}}=158 \dots 345 \text{ s m}^{-1}$) for deciduous forest. Breuer et al. (2003) show even broader spans with $r_{s\text{min}}$ well beyond 700 s m^{-1} for species of both types. In this study for forest 400 s m^{-1} , for low vegetation 100 s m^{-1} and as an approximation for mixed lower/forest vegetation 250 s m^{-1} were used.

In the study the LAI is computed according to the land use information of the LITFASS-area. To determine the LAI, for certain land use classes empirical relations from field studies are assigned based on the given land use as mentioned above. Grass and the different forms of crop are summarised in one group with lower vegetation. Therefore the approach of Bach et al. (2000) is used with:

$$\text{LAI} = (1.86 \cdot \text{NDVI})^{6.06} \quad (4)$$

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



for coniferous forest LAI is given by Bach et al. (2000) with:

$$\text{LAI} = (1.6 \cdot \text{NDVI})^3 \quad (5)$$

The sensitivity of the used parameters and their influence on the determination of $L.E$ is described in detail in Tittebrand et al. (2005).

5 Providing all needed parameters for the Penman-Monteith approach the radiation and energy fluxes can be inferred: The net radiation at surface is computed as the sum of the individual short- and longwave flux densities. Partly these components are achieved using an inverse remote sensing approach, so called look-up tables (LUT). Insolation at surface R_g and incoming longwave radiation emitted by the atmosphere
10 are calculated with the radiation transfer code Streamer (Key, 1999) over the entire short- and longwave spectra for the conditions of the specific date. Reflected shortwave radiation is calculated from R_g and the surface reflectance of each pixel. Outgoing longwave radiation is computed from surface temperature using Stefan-Boltzmann law.

15 Soil heat flux G for lower vegetation (grass and crop) is calculated using the approach by Gao (1998):

$$G = 0.539SR^{-0.53}e^{-K/\cos\theta_0}R_n \quad (6)$$

With the simple ratio $SR = (1 - \text{NDVI}) / (1 + \text{NDVI})$, the constant K and the sun zenith angle. For forest stands the canopy storage is added to the soil heat flux, based on field studies in Tharandt (near Dresden) for spruce measurements and here used for
20 satellite derived fluxes:

$$Gg = ag + bg \cdot R_n \quad (7)$$

$$Gj = aj + bj \cdot R_n \quad (8)$$

$$G = Gg + Gj \quad (9)$$

25 with Gg the soil heat flux, Gj the canopy heat flux and G the resultant entire surface soil heat flux for forest. The coefficients ag , bg , aj and bj resultant on linear regressions and are time dependent and used due to the time of satellite overpass (Berger, 2001).

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



To complete the energy balance equation ($R_n = L.E + H + G$), the sensible heat flux H is calculated and not inferred as residuum with the variables as mentioned above:

$$H = \rho c_p \frac{T_s - T_{2m}}{r_a} \quad (10)$$

3.3 Sensor-adaptation

5 Differing in range, shape and different influences by water vapour, the differences in the filter response functions of the three sensors have a strong influence on the determination of spectral reflectance and can result in a sensor-dependent NDVI (van Leeuwen et al., 1999; Teillet et al., 1997; Batra et al., 2006) and hence in differing results concerning the energy fluxes.

10 In this work the ETM results are treated as reference concerning the best validation results of the three satellites in comparison to grass-in-situ measurements of Lindenberg for all four scenes. Realizing comparable fluxes besides the mentioned differences in sensor features and filter response functions the AVHRR and MODIS fluxes should be computed using the same parameters and assumptions as used for the ETM data.

15 Comparing ETM- (arithmetically averaged to MODIS-resolution) and MODIS-albedo, NDVI and surface temperature for each pixel high correlation coefficients for the four investigated scenes 9 May 2002, 17 April 2003, 20 August 2002 and 28 July 2002 could be found: temperature ($r=0.66, 0.79, 0.78$ and 0.75 respectively), albedo ($r=0.95, 0.97, 0.90$ and 0.93) and NDVI ($r=0.79, 0.92, 0.89$ and 0.89). That allowed the usage

20 of the regression lines for correction/shifting (approximate to the 1:1 line) and therefore to create comparable MODIS-parameters for the according ETM-parameters. Especially the MODIS surface temperatures showed much lower values before and could be seen as the reason for the deviations e.g. in R_n between ETM and MODIS for all scenes.

25 For AVHRR surface temperature, albedo and TOA-NDVI were used from the SESAT scheme as base for the new calculation of the energy fluxes. Taking into account the

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2 h delay of the AVHRR overpass (AVHRR measurements are closed to the noon maximum in contrast to MODIS measurements at 10:00 UTC) the regression-adaptation-method should not be used for these sensor differences. The data analysis showed that an adaptation of NDVI – which is significantly lower for AVHRR (Buheaosier et al., 2003; van Leeuwen et al., 2006) than for ETM and MODIS (Fig. 2), is sufficient to infer comparable energy fluxes. NDVI adaptation was realized by stretching the NDVI-distribution histograms. Energy and radiant fluxes are then calculated according to the ETM algorithm, because of the comparable NDVI. Surface temperature was in good agreement compared to in situ data and therefore not changed. The resultant $L.E$ fluxes for the three sensors are shown in Fig. 3 in their given spatial resolution and for MODIS and AVHRR (smaller plots middle and respectively right) in the former computation for 20 August 2002. Comparing the original fluxes to the fluxes with parameter adaptation a significant improvement could be achieved not only for the range of $L.E$ but also for the spatial distribution of the values especially for AVHRR. A confinement according to water values had to be found using constant values $L.E = 125 \text{ Wm}^{-2}$ because the Penman-Monteith method was not applicable for water bodies. These plots are treated as the base for further analysis concerning spatial heterogeneity including investigations according to standard deviations, averaging methods and analysis of frequency distributions of fluxes and parameters.

Table 2 shows exemplarily the validation results from the 20 August 2002 compared to the grass-station of Lindenberg for ETM (original), MODIS (corrected) and AVHRR (corrected). Due to uncertainties in the geolocation especially of the AVHRR scenes also ambient pixels around the target pixel are extracted and averaged to achieve a relative statistic certainty to encounter the location of ground measurements. The results, given in Table 2 show an excellent agreement for R_n and surface temperature. Indeed ETM- R_n seems to overestimate at about 30 Wm^{-2} but taking the standard deviation into account the in-situ measurement lies within the error bars. Soil heat flux G remains an uncertain and hard to derive quantity with overestimations of all sensors of $40\text{--}50 \text{ Wm}^{-2}$. Values for the turbulent fluxes are missing for this day but taking the

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


other quantities into account – the evaporation can be assumed to fit well, especially for these very wet conditions in August 2002, although LITFASS-area is dominated by sandy soils with high infiltration. H is computed using Eq. (10) and not the remaining quantity of the energy balance. So the system is open, but with only small gaps in the energy balance (ETM: 23.22 Wm^{-2} , MODIS: 19.32 Wm^{-2} and AVHRR: 11.04 Wm^{-2}).

Validation data from the MOL-RAO for the turbulent fluxes $L.E$ and H are only available for 17 April 2003 and in comparison to the ETM data the satellite derived fluxes are in good agreement for latent heat with 93 Wm^{-2} (in-situ measurement) versus 72.25 Wm^{-2} (standard deviation=20.58) inferred from the satellite data, but for H with 48 Wm^{-2} (in situ) versus 118.96 Wm^{-2} (standard deviation=12.79), with higher deviations. The comparison between in-situ data and MODIS- as well as AVHRR results for the turbulent fluxes shows significant higher deviations than for ETM.

To gain comparable energy fluxes of the three sensors they had to be averaged onto the same spatial resolution. Figure 4 shows the LITFASS-mean inferred from ETM and MODIS data that were averaged to a comparable AVHRR-resolution (1.1 km). Two averaging strategies are used: a simple mean (arithmetically, left) and the average based on the dominant land use-method (right). Additionally standard deviation (red bar) is given. The sensor-means differ only marginal compared to each other as well as for the averaging methods in each of the Figures.

Between the two averaging-methods a correlation of $r=0.94$ could be found for $L.E$. The distributions of the values (Fig. 5) show obvious differences regarding the shape while the range mainly remains. For both methods the range is narrower compared to the original ETM- $L.E$ distribution. Weighting with the area of a special (dominant) land use, the shape better agree with the reference. However, by the usage of the PDF approach improvements according to the results are promised taking the entire range and shape of the distribution of the parameter for flux determination into account.

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Density functions – results and discussion

As briefly shown in the previous chapter the distribution of values can give further information concerning heterogeneity. LITFASS-mean is nearly the same for some fluxes between the different sensors. But the question is whether it is possible to find typical PDF of parameters and fluxes for selected land use classes?

We split here the overall frequency distribution into land use classes, resulting in individual PDF of Albedo, NDVI and $L.E$. The focus is given to the question whether it is necessary to differentiate between crop and grass or if it is enough to have one “low vegetation” class. Therefore a land use dependent distribution was realized based on the Landsat-classification for ETM and on CORINE for MODIS and AVHRR. For the analysis the left panel of Fig. 5 was investigated according to the land use types grass, crop and forest with respect to the question: Which land use class provides which pattern? Figure 6 shows exemplarily the frequency distribution for albedo and NDVI (as example for satellite derived parameters) and the resultant $L.E$ for ETM on the 20 August 2002.

First: for albedo (left side of Fig. 6), a completely different pattern can be found for forest compared to crop and grass, which show very similar distributions as expected. Highest frequency for forest is around 11–12% according to the dark color of the forest including nearly 100% *Pinus Sylvestris*. Crop and grass values rise slowly from 10% on, finding the peak around 17–19% (more obvious for crop) and then descending again. Second: NDVI (middle) and $L.E$ (right side) show a strong agreement in their distribution-pattern, so the key parameter NDVI can be found clearly in the $L.E$ results. For August it can be pointed out that NDVI differs clearly for crop and grass. Indeed there is a comparable frequency maximum at $NDVI=0.78$ and $L.E=320\text{--}340\text{ Wm}^{-2}$ but for crop a second peak is found for $NDVI=0.3$, which is a typical value for old vegetation or a significant soil part, and with $L.E=40\text{--}60\text{ Wm}^{-2}$. Depending on the type of crop or time of the harvest this peak is significant or not and can not be found for grasslands. For pine the frequency maximum can be found for 0.8 and 320 Wm^{-2} for NDVI and

ACPD

8, 16219–16254, 2008

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$L.E$, respectively. The classes with smaller NDVI and $L.E$ are less represented and only caused by the understory. Analysis (distribution functions and areal plots) of NDVI and $L.E$ show a direct relationship for all land use types. Thus it can be pointed out the higher the NDVI (and thus the LAI) the higher the $L.E$ inferred with Penman-Monteith.

5 With exception for crop- and grass-albedo which are more or less regularly distributed, the mean (given within the diagrams in Fig. 6) and mode of the distribution functions are unequal especially for NDVI and $L.E$ of grass and crop noticeable by the considerable skew of the functions. Sensitivity studies show that a change in LAI from 1 to 3 (e.g. caused by an NDVI increase from 0.51 to 0.78 (mean and mode respectively, here for
10 crop from Fig. 6 middle) and in dependence on the used NDVI-LAI relationship, can results in an increase of $L.E$ up to 162 Wm^{-2} (Berger and Schwiebus, 2004; Tittebrand et al., 2005).

Using different land use classifications for ETM and MODIS/AVHRR data can result in uncertainties according to disagreement of the corresponded classes. Thus a pixel
15 investigation should show whether the entirety of all ETM pixels within one MODIS-pixel show the same dominant land use compared to the accordant MODIS pixel itself. Ten randomized MODIS pixels for each land use class (grass, forest and crop) are collected for the investigation including approximately 600 ETM pixels for comparison. If at least 80% of the ETM pixels provide the same land use type than the MODIS pixels,
20 statements according to the representativeness of the distribution-pattern should be allowed. A good agreement can be found for the forest class whereas it was difficult to differentiate between crop and grass classes that are often interchanged between Landsat-classification (ETM) and CORINE (MODIS). Figure 7a shows the frequency distributions of R_n for grass (left), crop (middle) and forest (right) exemplarily for the 28
25 July 2002.

The variability of the R_n distributions representing the same land use is highest for crop followed by grass and forest. While for grass and forest there is one significant peak at about 500 Wm^{-2} and about 600 Wm^{-2} respectively, for crop different frequency peak positions (dominating at about 440 Wm^{-2} and 550 Wm^{-2}) occur. The range is

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



much broader and the distributions differ from each other. This variability is explicable by the different kinds of crop that are summarised within this group providing different reflectance or vegetation indices due to their appearance.

Regarding $L.E$ in Fig. 7b for all land use types a broad range occur but also with observable peak positions. Comparable higher variability of the ETM distributions for grass and crop is found whereas for forest the distributions are closed to each other providing one significant peak for all at about 350 Wm^{-2} .

These analyses are further repeated for early summer (9 May 2002) and late summer (20 August 2002). Summarizing, it can be outlined that each land use type provides typical distribution patterns both for the selected parameters of the Penman-Monteith approach and the resultant fluxes. Changes of their peak-positions are according to seasonal changes. The typical shapes of the density functions are seen to be stable and therefore useful for PDF-applications. In general PDF for NDVI, Albedo, R_n and further parameter could be used to generate PDF of latent heat. The advantage is then the consideration of the natural heterogeneity of the surface by taking the entire range of the distributions of the parameters into account. However the $L.E$ determination is based on a lot of variables and assumptions that can be tampered by missing satellite measured parameters.

5 Conclusions and outlook

In this study an investigation of the heterogeneity of land surface parameters and energy fluxes for LITFASS-area (Germany, 50 km southeast of Berlin) was carried out.

Therefore some preliminary work had to be done to ensure the comparability of the results including the elimination of sensor differences between ETM, MODIS and AVHRR. This was obtained through adaptation, using regression lines of surface temperature, albedo and NDVI for MODIS and histogram stretching of the NDVI for AVHRR, with reference to ETM. Thus the same $L.E$ -determination method for all sensors could be realized. The results show reasonable agreement with in-situ data of the observa-

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tions in Lindenberg.

Due to the land use based parametrization for flux determination new land use classifications were used: the Landsat-classification (TU-Dresden, $25 \times 25 \text{ m}^2$) was applied for ETM data and CORINE ($1 \times 1 \text{ m}^2$) was used for MODIS and AVHRR data showing good agreement in comparison to in-situ data for LITFASS-area. Nevertheless differences remain, e.g. an overestimation of the crop classes of CORINE or an underestimation of water areas (Keil et al., 2005). The quality criterion for the Landsat-classification was an accuracy of 5% which had to be reached. Based on the same determination method and comparable land use classifications for the different sensors, a variability analysis has been carried out.

The LITFASS-mean and standard deviations were investigated for the data averaged (arithmetically and using the dominant land use method) to the AVHRR resolution. It was demonstrated that the LITFASS-mean of the fluxes is nearly comparable irrespective whether it is arithmetically or dominant averaged. Compared to the original resolution: averaging e.g. $L.E$ from the ETM sensor with both methods (e.g. for 20 August 2002) causes a decrease in the distribution range from $0\text{--}380 \text{ Wm}^{-2}$ with the maximum frequency at 320 Wm^{-2} (original resolution) to a range of $60\text{--}340 \text{ Wm}^{-2}$ with the maximum frequency for arithmetically averaged $L.E$ at 310 Wm^{-2} and for dominant land use at 320 Wm^{-2} . For all fluxes the standard deviation for arithmetically averaged data is lower than for dominant land use averaged data.

The heterogeneity analysis was carried out with respect to frequency distributions for the land use classes grass, crop and forest to find typical pattern of their distribution functions and the comparability and variability of ETM parameters or flux distributions on MODIS-pixel base.

The analysis with respect to land use classes show differences in the distribution pattern, not only between grass and forest but also between grass and crop. One question of the study was to investigate whether it is possible to summarize both classes to one group e.g. lower vegetation. Thus, for further studies using the PDF e.g. for $L.E$ -determination it is suggested to differentiate between crop and grass because a second

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



peak for crop showing the bare soil conditions in the distributions of NDVI and thus $L.E$ appears (in three different seasonal states of summer) that can not be neglected.

Selecting MODIS pixels with known land use classes and analyzing therefore their included ETM-distributions (consisting of approximately 600 pixels each), allowed conclusions with respect to pixel-heterogeneity, assumed that they offer the same land use type (at least 80%). The distribution variability of $L.E$ is shown to be higher than for R_n except for the crop class where an evident difference in the distribution pattern can be found. The best distribution agreement can be outlined for forest. According to spatial variability of $L.E$ former studies also showed that the latent heat flux was the most sensitive parameter for that issue (Li and Avissar, 1994). Thus the results indicate that it is very important to consider the spatial variability of NDVI and therefore LAI, stomatal resistance and further important parameters of the Penman-Monteith approach.

Averaging highly resolved satellite data arithmetically or using dominant land use gives only marginal differences, but it is assumed that using PDF and thus the entire range of the surface characteristics, including extremes as well as the mode instead of the mean, should provide more realistic conditions for the determination of energy fluxes. Li and Avissar (1994) confirmed that the results of determined fluxes depend on the shape of the spatial distribution of the land surface parameters, used for the calculation. Typically, the more skewed the distribution within the range of values the larger their error (for nonlinear relationships between parameter and flux e.g.: for LAI, inferred nonlinear from NDVI). They found largest differences for lognormal distributed land surface parameters and showed that the shape of the distribution is essential for the determination of accurate fluxes. Our preliminary results (MAGIM, 2006) confirm the possible application of the PDF-approach for the determination of latent heat flux with Penman-Monteith using NDVI, albedo, relative humidity and wind speed for grassland and coniferous forest. Assuming χ^2 -distributions with a different range and mode for grass and forest the method was applicable for these two land use types as a first approximation. The approach can now be improved by the new distribution pattern of the parameters for the two land use types and by the enhancement for the

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

often represented land use type “crop” with a clearly different distribution function compared to grass - especially for NDVI as the key variable for the $L.E$ -determination using Penman-Monteith.

Finally, realizing the $L.E$ -determination with the PDF approach offers new interpretation of validation results. Until now pixel mean and standard deviation are compared to point measurements often resulting in significant differences. Taking the entire PDF of the resultant latent heat flux for comparison into account, the in situ measurement can be different from mean or mode and nevertheless be representative for the pixel.

Appendix

CORINE: CORINE is structured in 3 layers, referring to 5 main groups in the first layer and specifying the areas in layer 2 and 3. So 44 land use or land cover units that can be differentiated. These classes are strongly affected by nature conservation and environmental background. A verification in technical form and content supports the target accuracy. 2006 CORINE was compared to LUCAS-data (land use/cover area frame statistical survey, EEA, 2006) The accuracy of CORINE 2000 was found to be $87\% \pm 0.8\%$ and highest accuracy was given for water classes as well as for industrial areas (EEA, 2006). Also large areas of agriculture and coniferous forests offered an accuracy between 90 and 95%. In contrast Keil et al. (2005) found an overestimation of crop areas and an underestimation of water bodies. One reason can be seen in the minimum area, which is 25 ha, to acquire areas of interest. Thus areas smaller than 25 ha are left out and dominant areas tend to overestimate.

Landsat-classification 2002: 15 classes could be identified by using a hierarchical system similar to CORINE. Using defined spectral thresholds preliminary classes of different land use could be selected. These classes then could be obtained by applying classification algorithms as maximum-likelihood, cluster analysis or neighborhood analysis (prior verified with training areas, Prechtel, personal communication, 2007) to differentiate the classes. The target accuracy of the resultant classes was 5% and was

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tested by drawing a sample.

Reference classification: For comparison an additionally reference classification, based on field observations was used for comparison (DWD, 2003). This classification is given with 25 m spatial resolution and was observed during the field experiment in 2003. It was not possible to acquire the entire terrain, so nearly 9% are not classified and dedicated to the other classes (Beyrich et al., 2004). The classification is given in 32 or summarized to 9 classes. Changes are according to agriculture and differences in the kind of crop. Actually for the satellite analysis in this study all kinds of crop are summarized to one “crop-class” thus the differences between the years 2003 and 2002 are negligible (for the analysis).

Acknowledgements. This study was founded by the Federal Ministry for Education and Research (BMBF) within the framework of EVA-GRIPS (Grant Number 01 LD 0103), which is a contribution to the climate research programme DEKLIM. Thanks to the German Weather Service for providing in-situ data for the validation.

References

- Allen, R. G.: A Penman for all seasons, J. Irrig. Drain Eng., 112/4, 348–368, 1986.
- Allen, R. G., Pereira, L. S., Raes, D., and Smith, M.: Crop evapotranspiration – guidelines for computing crop water requirements, FAO Irrigation and Drainage paper, 56, Rome, Italy, 1998.
- Arain, A. M., Miachaud, J. D., Shuttleworth, W. J., and Dolman, A. J.: Testing of vegetation parameter aggregation rules applicable to the Biosphere-Atmosphere Transfer Scheme (BATS) at the Fife site, J. Hydrol., 177, 1–22, 1996.
- Avissar, R.: Conceptual aspects of a statistical-dynamical approach to represent landscape subgrid-scale heterogeneities in atmospheric models, J. Geophys. Res., 97, 2729–2742, 1992.
- Bach, H., Braun, M., Lampart G., and Ranzi, R.: Estimates of surface parameters via remote sensing for the Toce watershed, in: RAPHAEL – Runoff and Atmospheric Processes for Flood Hazard Forecasting and Control, edited by: Bacchi, B. and Ranzi, R., EC, Directorate

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



General XII, Programme Environment and Climate 1994–1998, Contract no. ENV4-CT97-0552, Final Report, 2000.

Batra, N., Islam, S., Venturini, V., Bisht G., and Jiang, L.: Estimation and comparison of evapotranspiration from MODIS and AVHRR sensors for clear sky days over the Southern Great Plains, *Remote Sens. Environ.*, 103(1), 1–15, 2006.

Becker, F. and Li, Z.: Towards a local split window method over land surfaces, *Int. J. Remote Sens.*, 11, 369–393, 1990.

Becker, F. and Li, Z.: Surface temperature and emissivity at various scales: Definition, measurement and related problems, *Remote Sens. Rev.*, 12, 225–253, 1995.

Berger, F. H.: Bestimmung des Energiehaushaltes am Erdboden mit Hilfe von Satellitendaten, *Tharandter Klimaprotokolle*, Band 5, 206 pp., 2001.

Berger, F. H. and Schwiebus, A.: Energieflüsse heterogener Landoberflächen, abgeleitet aus Satellitendaten, *Schlussbericht Projekt VERTIKO, TUD2, FK 07 AFT37-TUD2, Technische Universität Dresden*, 27 pp., 2004.

Beyrich, F., Adam, W. K., Bange, J., Behrens, K., Berger, F. H., Bernhofer, C., Bösenberg, J., Dier, H., Foken, T., Gödecke, M., Görsdorf, U., Güldner, J., Hennemuth, B., Heret, C., Huneke, S., Kohsiek, W., Lammert, A., Lehmann, V., Leiterer, U., Leps, J.-P., Liebethal, C., Lohse, H., Lüdi, A., Mauder, M., Meijnger, W. M. L., Mengelkamp, H.-T., Queck, R., Richter, S. H., Spieß, T., Stiller, B., Tittebrand, A., Weisensee, U., and Zittel, P.: Verdunstung über einer heterogenen Landoberfläche – das LITFASS-2003 Experiment: ein Bericht, *Arbeitsergebnisse Nr. 79, Deutscher Wetterdienst, Offenbach, Germany*, 100, 2004.

Bonan, G. B., Pollard, D., and Thompson, S. L.: Influence of Subgrid-Scale Heterogeneity in Leaf Area Index, Stomatal Resistance, and Soil Moisture on Grid-Scale Land–Atmosphere Interactions, *J. Climate*, 6(10), 1882–1897, 1993.

Breuer, L., Eckhardt, K., and Frede, H.-G.: Plant parameter values for models in temperate climates, *Ecol. Model.*, 169, 237–293, 2003.

Buheaosier, K., Tsuchiya, K., Keneko, M., and Sung, S. J.: Comparison of image data acquired with AVHRR, MODIS, ETM+ and ASTER over Hokkaido, Japan, *Adv. Space Res.*, 32, 11, 2211–2216, 2003.

Chehbouni, A., Njoku, E. G., Lhomme, J.-P., and Kerr, Y. H.: An approach for averaging surface temperature and surface fluxes over heterogeneous surfaces, *J. Climate*, 5, 1386–1393, 1995.

Claussen, M.: Estimation of regional heat and momentum fluxes in homogenous terrain with

ACPD

8, 16219–16254, 2008

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- bluff roughness elements, *J. Hydrol.*, 166, 353–369, 1995a.
- De Rooy, W. C. and Holtslag, A. A. M.: Estimation of surface radiation and energy flux densities from single-level weather data, *J. Appl. Meteorol.*, 38, 526–540, 1999.
- EEA, European Environment Agency: The thematic accuracy of CORINE land cover 2000, assessment using LUCAS (Land Use/Cover Area frame statistical Survey), Technical report No. 7, European Environment Agency, Copenhagen, 85 pp., 2006.
- EOS: <http://delenn.gsfc.nasa.gov/~imswwww/pub/imswelcome/index.html>, 5 October 2007.
- Entekhabi, D. and Eagleson, P. S.: Land Surface Hydrology Parametrization for atmospheric general circulation models including subgrid scale spatial variability, *J. Climate*, 2, 816–831, 1989.
- Gao, W., Coulter, R. L., Lesht, B. M., Qiu, J., and Wesely, M. L.: Estimating clear-sky regional surface fluxes in the Southern Great Plains Atmospheric Radiation Measurement Site with ground measurements and satellite observations, *J. Appl. Meteorol.*, 37, 5–22, 1998.
- Garrigues, S., Allard, D., Baret, F., and Weiss, M.: Quantifying spatial heterogeneity at the landscape scale using variogram models, *Remote Sens. Environ.*, 103, 81–96, 2006.
- Giorgi, P.: An approach for representation of surface heterogeneity in land surface models – Part 1: Theoretical framework, *Mon. Weather Rev.*, 125, 1885–1899, 1996a.
- Giorgi, P.: An approach for representation of surface heterogeneity in land surface models – Part 2: Validation and sensitivity experiments, *Mon. Weather Rev.*, 125, 900–1919, 1996b.
- Hagemann, S.: An improved land surface parameter dataset for global and regional climate models, Report No. 336, MPI for Meteorology, Hamburg, Germany, 2002.
- Keil, M., Kiefl, R., and Strunz, G.: CORINE Land Cover 2000 – Europaweit harmonisierte Aktualisierung der Landnutzungsdaten für Deutschland, Abschlussbericht zum F+E Vorhaben, UBA FKZ 201 12 209 (1 May 2001–31 December 2004), 2005.
- Kelliher, F. M., Leuning, R., Raupach, M. R., and Schulze, E.-D.: Maximum conductances for evaporation from global vegetation types, *Agr. Forest Meteorol.*, 73, 1–16, 1995.
- Key, J. R.: Streamer – user's guide, Technical report 96-01, Department of Geography, Boston University, Boston, MA, USA, 1999.
- Koster, R. D. and Suarez, M. J.: A comparative analysis of two land surface heterogeneity representations, *J. Climate*, 5, 12, 1379–1390, 1992a.
- Koster, R. D. and Suarez, M. J.: Modeling the land surface boundary in climate models as a composite of independent vegetation stands. *J. Geophys. Res.*, 97(D3), 2697–2715, 1992b.
- Landsat: http://landsat.usgs.gov/project_facts/project_news/June_2003.php, 10 September

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2003.

Lhomme, J.-P.: Energy balance of heterogeneous terrain: averaging the controlling parameters, *Agr. Forest Meteorol.*, 61, 11–21, 1992.

Lhomme, J.-P., Chehbouni, A., and Monteny, B.: Effective parameters of surface energy balance in heterogeneous landscape, *Bound.-Lay. Meteorol.*, 71, 297–309, 1994.

Li, B. and Avissar, R.: The Impact of Spatial Variability of Land-Surface-Characteristics on Land-Surface Heat Fluxes, *J. Climate*, 7, 527–537, 1994.

Liang, S., Shuey, C. J., Russ, A. L., Fang, H., Chen, M., Walthall, C. L., Daughtry, C. S. T., and Hunt Jr., R.: Narrowband to Broadband conversion of land surface albedo – II: Validation, *Remote Sens. Environ.*, 84, 25–41, 2002.

Ma, Y., Zhong, L., Su, Z., Ishikawa, H., Meneti, M., and Koike, T.: Determination of regional distributions and seasonal variations of land surface heatfluxes from Landsat-7 enhanced Thematic Mapper data over the central Tibetan Plateau area, *J. Geophys. Res.*, 111, D10305, doi:10.1029/2005JD006742, 2006.

MAGIM: Forschergruppe 536 der Deutschen Forschungsgemeinschaft: Matter fluxes in grasslands of Inner Mongolia as influenced by stocking rate (MAGIM), Fortsetzungsantrag MAGIM-Phase II: 04/2007-03/2010, Institut für Meteorologie und Klimaforschung (IMK-IFU), Forschungszentrum Karlsruhe, Garmisch-Partenkirchen, Germany, 2006.

McClain, E. P., Pichel, W. G., and Walton, C. C.: Comparative performance of AVHRR-based multichannel sea surface temperatures, *J. Geophys. Res.*, 90, 11 587–11 601, 1985.

Mengelkamp, H.-T., Beyrich, F., Heinemann, G., Ament, F., Bange, J., Berger, F. H., Bösenberg, J., Foken, T., Hennemuth, B., Heret, C., Huneke, S., Johnsen, K.-P., Kohsiek, W., Leps, J.-P., Liebenthal, C., Lohse, H., Mauder, M., Meijninger, W., Raasch, S., Simmer, C., Spieß, T., Tittebrand, A., Uhlenbrock, J., and Zittel, P.: Evaporation over a heterogeneous land surface: the EVA-GRIPS project, *B. Am. Meteorol. Soc.*, 775–786, 2006.

Mölders, N. and Raabe, A.: Numerical investigations on the influence of subgrid-scale surface heterogeneity on evapotranspiration and cloud processes, *J. Appl. Meteor.*, 35, 782–795, 1996.

NASA Goddard Space Flight Center: LANDSAT 7 Science Data User Handbook, Landsat Project Science Office, NASA Goddard Space Flight Center, MD, USA, 2004.

Njoku, E. G., Hook, S. J., and Chehbouni, A.: Effects of surface heterogeneity on thermal remote sensing of land parameters, in: *Scaling up in Hydrology using Remote Sensing*, edited by: Stewart, J. B., Engmann, E. T., Feddes, R. A., and Kerr, Y., John Wiley and Sons,

ACPD

8, 16219–16254, 2008

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



hichester, 1996.

Noilhan, J., Lacarrere, P., Dolman, A. J., and Blyth, E. M.: Defining area-average parameters in meteorological models for land surfaces with mesoscale heterogeneity, *J. Hydrol.*, 190, 302–316, 1997.

5 Pinker, R. T.: Satellites and our understanding of the surface energy balance, *Palaeogeogr. Palaeoclimatol.*, 82, 321–342, 1990.

Sellers, P. J., Heiser, M. D., Hall, F. G., Verma, S. B., Desjardins, R. L., Schuepp P. M., and MacPherson, J. I.: The impact of using area-averaged land surface properties – topography, vegetation condition, soil wetness – in calculations of intermediate scale (approximately 10 km²) surface-atmosphere heat and moisture fluxes, *J. Hydrol.*, 190, 269–301, 1997.

10 Shuttleworth, W. J., Yang, Z.-L., and Arain, M. A.: Aggregation rules for surface parameters in global models, *Hydrol. Earth Syst. Sci.*, 1, 217–226, 1997, <http://www.hydrol-earth-syst-sci.net/1/217/1997/>.

Sobrino J. A., Jimenez-Munoz, J. C., and Paolini, L.: Land surface temperature retrieval from LANDSAT TM 5, *Remote Sens. Environ.*, 90, 434–440, 2004.

15 Sonntag, D. and Heinze, D.: Sättigungsdampfdruck- und Sättigungsdampfdichtetafeln für Wasser und Eis, 1. Ed., VEB Deutscher Verlag für Grundstoffindustrie, Leipzig, 1982.

Teillet, P. M., Staenz, K., and Williams, D. J.: Effects of spectral, spatial, and radiometric characteristics on remote sensing vegetation indices of forested regions, *Remote Sens. Environ.*, 61, 139–149, 1997.

20 Tittebrand, A., Schwiebus, A., and Berger, F. H.: The influence of land surface parameters on energy flux densities derived from remote sensing data, *Meteorol. Z.*, 14(2), 227–236, 2005.

Van Leeuwen, W. J. D., Huete, A. R., and Laing, T. W.: MODIS Vegetation Index Compositing Approach: a prototype with AVHRR data, *Remote Sens. Environ.*, 69, 264–280, 1999.

25 Van Leeuwen, W. J. D., Orr, B. J., Marsh, S. E., and Herrmann, S. M.: Multi-sensor NDVI data continuity: uncertainties and implications for vegetation monitoring applications, *Remote Sens. Environ.*, 100, 67–81, 2006.

Vermote, E., Tandre, D., Deutze, J. L., Herman, M., Morcrette, J. J.: Second Simulation of the Satellite Signal in the Solar Spectrum (6S), 6S User Guide, NASA Goddard Space Flight Center, Greenbelt, 54 pp., 1997.

30 Wan, Z. and Dozier, J.: Land surface temperature measurement from space: Physical principles and inverse modelling, *IEEE T. Geosci. Remote*, 27, 268–278, 1989.

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Spatial heterogeneity
for LITFASS-area**A. Tittebrand and
F. H. Berger

Table 1. Comparison of the land-use classifications CORINE and Landsat-classification that are used for the satellite data for LITFASS-area with the reference classification (Beyrich et al., 2004) based on a field study in 2003.

Classification/Satellite	25 m_Landsat % -classification	CORINE %	Field study Reference %
water	7	8	7
deciduous forest	4	–	–
mixed forest	7	2	–
coniferous forest	29	36	43
grass	24	8	13
crop	27	43	32
urban areas	2	3	5

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger

Table 2. Validation results for corrected (corr) energy flux densities (net radiation (R_n), sensible heat flux (H), latent heat flux ($L.E$) and soil heat flux (G)) and surface temperature (T_s) for 20 August 2002.

	in-situ measurement	ETM 09:45 UTC	Standard deviation ETM results	in-situ measurement	MODIS-corr 10:00 UTC	standard deviation for MODIS results
R_n [Wm^{-2}]	440	473.93	31.6	460	458.06	40.13
H [Wm^{-2}]	–	160.76	21.9	–	165.17	45.3
$L.E$ [Wm^{-2}]	–	206.90	53	–	187.15	94.5
G [Wm^{-2}]	30	83.05	14.28	34	86.42	22.31
T_s [K]	307.55	308.66	2.68	309.45	309.59	2.97

	in-situ measurement	AVHRR-corr 11:45 UTC	standard deviation for AVHRR results
R_n [Wm^{-2}]	461	476.66	27.11
H [Wm^{-2}]	–	153.42	44.07
$L.E$ [Wm^{-2}]	–	235.23	94.17
G [Wm^{-2}]	37	76.97	25.5
T_s [K]	310.55	307.72	2.04

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Spatial heterogeneity
for LITFASS-area**

A. Tittebrand and
F. H. Berger




Fig. 1. Land use classification of LITFASS-area from in-situ reference data 2003.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Spatial heterogeneity
for LITFASS-area**

A. Tittebrand and
F. H. Berger




Fig. 2. Differences in NDVI determined with the three sensors in their given spatial resolution, note that AVHRR-NDVI is uncorrected for atmospheric effects (TOA).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Spatial heterogeneity for LITFASS-area

A. Tittebrand and
F. H. Berger




Fig. 3. recalculated $L.E$ fluxes for 20 August 2002 for MODIS (middle) and AVHRR (right) in their given spatial resolution, based on the Penman-Monteith approach, using the same parametrization as for ETM (left). (Small plots show the determined preliminary fluxes before, without adaptation to the ETM-parameters).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


**Spatial heterogeneity
for LITFASS-area**A. Tittebrand and
F. H. Berger

Fig. 4. Mean and standard deviation (std) of the energy fluxes and surface temperature for LITFASS-area for 20 August 2002, arithmetic mean (left), dominant land use method (right), both averaged to AVHRR resolution (1.1 km).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




**Spatial heterogeneity
for LITFASS-area**A. Tittebrand and
F. H. Berger

Fig. 5. Frequency distribution of ETM- $L.E$ for 20 August 2002 with original resolution (left), arithmetic mean (middle), dominant land use method (right), both averaged to MODIS resolution (800 m).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


**Spatial heterogeneity
for LITFASS-area**A. Tittebrand and
F. H. Berger

Fig. 6. Frequency distributions of albedo (left column), NDVI (middle column) and $L.E.$ (right column) for 20 August 2002 for different land use types (grass, crop, forest).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


**Spatial heterogeneity
for LITFASS-area**A. Tittebrand and
F. H. Berger

Fig. 7a. Frequency distribution of ETM- R_n for 28 July 2002 for grass (left), crop (middle) and forest (right), within the accordant MODIS pixels.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


**Spatial heterogeneity
for LITFASS-area**A. Tittebrand and
F. H. Berger

Fig. 7b. Frequency distribution of ETM-L.E for 28 July 2002 for grass (left), crop (middle) and forest (right), within the accordant MODIS pixels.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)