

This discussion paper is/has been under review for the journal *Atmospheric Chemistry and Physics (ACP)*. Please refer to the corresponding final paper in *ACP* if available.

**ECMWF vs. AIRS
UTH and CALIPSO
clouds**

N. Lamquin et al.

Evaluation of upper tropospheric humidity forecasts from ECMWF using AIRS and CALIPSO data

N. Lamquin¹, K. Gierens², C. J. Stubenrauch¹, and R. Chatterjee³

¹Laboratoire de Météorologie Dynamique, Ecole Polytechnique, Palaiseau, France

²Deutsches Zentrum für Luft- und Raumfahrt, Institut für Physik der Atmosphäre, Oberpfaffenhofen, Germany

³National Institute of Technology, Tiruchchirappalli, Tamil, Nadu, India

Received: 28 July 2008 – Accepted: 21 August 2008 – Published: 1 October 2008

Correspondence to: N. Lamquin (nicolas.lamquin@lmd.polytechnique.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

An evaluation of the upper tropospheric humidity from the European Centre of Medium-Range Weather Forecasts (ECMWF) Integrated Forecast System (IFS) is presented. We first make an analysis of the spinup behaviour of ice supersaturation in weather forecasts. It shows that a spinup period of at least 12 h is necessary before using forecast humidity data from the upper troposphere. We compare the forecasted upper tropospheric humidity with coincident relative humidity fields retrieved from the Atmospheric InfraRed Sounder (AIRS) and with cloud vertical profiles from the Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation (CALIPSO). The analysis is made over one year, from October 2006 to September 2007, and we discuss how relative humidity and cloud features appear both in the IFS and in the observations. In a last part, we investigate the presence of ice supersaturation within low vertical resolution pressure layers by comparing the IFS outputs for high-resolution and low-resolution humidity profiles and by simulating the interpolation of humidity over radiosonde data. A new correction method is proposed and tested with these data.

1 Introduction

Ice supersaturation in the upper troposphere is an explicit feature in the Integrated Forecast System (IFS) of the European Centre of Medium-Range Weather Forecasts (ECMWF), operational since 13 September 2006 (IFS cycle 31r1). This new feature, introduced by Tompkins et al. (2007), has produced some changes in the statistics of upper tropospheric humidity and cloud fraction in the IFS. In particular, there is an increase in upper-tropospheric humidity, a decrease in high-level cloud cover and, to a much lesser extent, cloud ice amounts. First analyses of the frequency distribution of the modelled supersaturation values showed good agreement with a climatology derived from in situ aircraft observations (Tompkins et al., 2007). The global distribution of frequency of occurrence of supersaturated regions in the new scheme compares

ECMWF vs. AIRS UTH and CALIPSO clouds

N. Lamquin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



well with remotely sensed microwave limb sounder (MLS) data, with the most marked underprediction occurring in regions where the model is known to underpredict deep convection (Tompkins et al., 2007).

A major inconsistency in the IFS is that the tangent linear and adjoint models used in the data assimilation scheme do not involve ice supersaturation (Janisková et al., 2002), while the forecast model does. The only way an analysis (i.e. the initial state for a new forecast run) can obtain supersaturation is during the final trajectory integration which uses the full physics of the forecast model. In other words, forecasts are initialized with states that usually have less ice supersaturation than the forecast for the same time from the day before. Hence, there is a supersaturation spinup in the forecast runs, and we will first present an analysis of it.

The present paper also aims at further tests of the new supersaturation scheme regarding the simulated upper tropospheric humidity. The ECMWF tropical water vapour has been assessed by Luo et al. (2007) with the help of Measurements of Ozone and water vapour by in-service Airbus airCRAFT (MOZAIC) data, but the forecast data did not include the new ice supersaturation scheme. For our purpose we first compare the IFS forecasts with collocated Atmospheric InfraRed Sounder (AIRS) retrievals of relative humidity within pressure layers of 200–250, 250–300, 300–400, and 400–500 hPa for clear sky and cloudy situations. The influence of the vertical extent of clouds has been investigated by relating upper tropospheric humidity of the IFS forecasts to collocated Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation (CALIPSO) cloud vertical profiles. The comparisons are made over Europe (latitudes $\in[32^\circ, 74^\circ]$, longitudes $\in[-27^\circ, 45^\circ]$) with one year of data, from October 2006 to September 2007.

For the comparisons, we mostly use data on standard pressure levels. These data are interpolated from much higher resolved model level data, and it is conceivable that some supersaturation events will get lost in the interpolation procedure. To explore the effect of the interpolation further and to assess the occurrence of ice supersaturation using data with low vertical resolution, we compare relative humidity using ECMWF forecasts of high and low resolution, and we also simulate the averaging effect by using

ECMWF vs. AIRS UTH and CALIPSO clouds

N. Lamquin et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

radiosonde data from Lindenberg, Germany.

2 Data handling

2.1 ECMWF forecasts

ECMWF provides global weather forecasts twice daily (00:00 and 12:00 UTC) with output time steps of 3 h in the first 3 days and 6 h from then on to day 10. The model uses a horizontal spectral resolution of T511 (corresponding to a resolution of about 25 km at the equator) and 91 layers in the vertical (corresponding to a resolution of about 15 hPa). For our studies we extracted temperature, humidity and cloud cover data from the upper troposphere (500 to 200 hPa) in T213 resolution (about 65 km at the equator) and remapped them onto a regular $0.5^\circ \times 0.5^\circ$ grid. We analyzed one year (from October 2006 to September 2007) of data on the standard pressure levels (200, 250, 300, 400, 500 hPa) and January and July 2007 on model levels which we interpolated on a vertical grid with higher resolution of 25 hPa. We first computed relative humidity from the specific humidity and temperature on model levels and then made the vertical interpolation. For the comparison with the satellite data layer values instead of level values of relative humidity are obtained by averaging the lower and upper edges for each standard pressure layer. This procedure is summarized in Fig. 1.

Figure 2 shows the difference between relative humidity with respect to ice obtained at high vertical resolution and the one obtained by interpolation to lower resolution as a function of relative humidity at high vertical resolution. The figure shows that data produced with different interpolation procedures differ typically by a few percent (in RHi-units).

The new supersaturation scheme is only in effect at temperatures lower than 250 K. Relative humidity is calculated with respect to the ice phase (RHi) in this temperature regime. Thus comparisons are only made for temperatures lower than 250 K.

ECMWF vs. AIRS UTH and CALIPSO clouds

N. Lamquin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2 A-Train data and collocations with ECMWF

The A-Train satellite constellation (Stephens et al., 2002) provides a large panel of new instruments and offers increased possibilities to understand the Earth's atmosphere and climate. Among these instruments AIRS provides temperature and humidity profiles since May 2002, and CALIPSO provides cloud vertical profiles since July 2006. For our purposes we will use observations made by these two instruments to evaluate the ECMWF forecasts. Differences in spatial resolution and sampling of both instruments have led to different collocation schemes which are described in the following.

Onboard the Earth Observing System (EOS) satellite Aqua, AIRS provides very high resolution measurements of Earth emitted radiation in three spectral bands from 3.74 to 15.40 μm , using 2378 channels, at 01:30 and 13:30 local time (LT). The spatial resolution of these measurements is 13.5 km at nadir. Nine AIRS measurements (3 \times 3) correspond to one footprint of the Advanced Microwave Sounder Unit (AMSU). AIRS level 2 (L2) standard products include temperature at 28 pressure levels from 0.1 hPa to the surface and water mixing ratios at 14 pressure layers from 50 hPa to the surface (Suskind et al., 2003, 2006). These atmospheric profiles were retrieved from cloud-cleared AIRS radiances (Chahine et al., 2006) within each AMSU footprint, at a spatial resolution of about 45 km, which is close to the one of ECMWF. Validations with radiosonde data from the NOAA NESDIS operational meteorological database archive (Divakarla et al., 2006) and with Atmospheric Radiation Measurement (ARM) data (Tobin et al., 2006) have shown that the accuracy is close to 1 K in 1 km layers for temperature and better than 15% in 2 km layers for water vapour. The nominal vertical resolution of temperature and humidity profiles is about 2 to 3 km in the upper troposphere (Gettelman et al., 2004; Read et al., 2007). We use version 5 of AIRS L2 data and retrieve RH_i as in Lamquin et al. (2008). Specific humidity is obtained as an integral over an atmospheric layer, whereas temperatures are given at top and bottom of each layer. Therefore RH_i is determined over the atmospheric layer, as in (Stubenrauch and Schumann, 2005). The saturation specific humidity with respect

ECMWF vs. AIRS UTH and CALIPSO clouds

N. Lamquin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**ECMWF vs. AIRS
UTH and CALIPSO
clouds**

N. Lamquin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

to ice integrated over the pressure layer is obtained from the saturation water vapour pressure with respect to ice (Sonntag, 1990). The latter is determined in steps of 1 hPa from the linearly interpolated temperature profiles within the pressure layer. A selection is made over AIRS L2 profiles of best and good quality using the conditions $Qual.H_2O \neq 2$ and $P_{Good} > 600$ hPa from AIRS L2 quality flags (Suskind et al., 2006; Tobin et al., 2006). If the atmosphere is too cloudy, the atmospheric profile may be of bad quality. In addition, as suggested in Gettelman et al. (2004) and Read et al. (2007), layers for which the water vapour content is lower than the nominal instrument sensitivity ($q = 20$ ppmv) are rejected. RH_i is kept for four pressure layers between 200 and 500 hPa: 200–250, 250–300, 300–400, and 400–500 hPa. Our analysis is made over Europe, a region where the tropopause is situated around 200–300 hPa, mostly depending on the season. AIRS L2 also provides the air pressure at the tropopause and a description of clouds in terms of cloud pressure and effective cloud cover for up to two cloud layers. Pressure layers are discarded when the tropopause lays inside the layer to avoid biases due to the inversion of the temperature gradient at the tropopause.

Aqua overflies at 01:30 and 13:30 LT, and the AIRS large swath (about 1700 km) makes it possible to find numerous events which are close in terms of time and location to ECMWF forecasts for 00:00 and 12:00 UTC. The geographical proximity is realized by associating events with a centre-to-centre distance $\sqrt{(\Delta lat)^2 + (\Delta lon)^2}$ smaller than 0.25° . Events are rejected when the time interval is larger than 30 min. One year of collocations then leads to a total amount of 325 851, 527 561, 377 856, and 69 754 events for the pressure layers 400–500, 300–400, 250–300 and 200–250 hPa, respectively.

The CALIPSO mission is a collaboration between National Aeronautics and Space Administration (NASA) and the French National Center of Space Studies (Centre National d'Etudes Spatiales, CNES) (Winker et al., 2003). CALIPSO contains the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) which can discriminate the vertical distribution of water and ice clouds as well as aerosol masses (Sassen, 1991). The performance of CALIOP is summarized in Winker et al. (2007). The instrument provides an accurate vertical profile of backscattered radiation at 532 nm and 1064 nm at

**ECMWF vs. AIRS
UTH and CALIPSO
clouds**N. Lamquin et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

a vertical resolution of 60 m for altitudes between 8.2 and 20.2 km, where high clouds are situated. CALIOP is a nadir viewing instrument, and the width of each shot is about 70 m, sampled every 333 m along the track. The 5 km CALIPSO L2 cloud products provide the number of cloud layers as well as their vertical extent (top and base altitudes) averaged over 5 km, as long as the signal is not totally attenuated by thick clouds (with τ larger than 5; Winker et al., 2003). We use version 2 of CALIPSO L2 data, with geometrical cloud height transformed to pressure using meteorological atmospheric profiles of the Global Modeling and Assimilation Office (GMAO) which are available in the CALIPSO L1 data. The collocation of the ECMWF forecasts with CALIPSO does not provide as many events as the collocation with AIRS, because CALIOP only measures at nadir. Nevertheless, a sufficient number of collocated profiles (75 280) was found. Again the time interval has to be smaller than 30 min, and an IFS grid box is associated with a CALIPSO pixel only when the centre-to-centre distance is smaller than 0.15° . This takes into account the fact that CALIPSO pixels are much smaller than the IFS grid boxes and that we only keep events close to the centre. One to four CALIPSO pixels fall within an IFS box. We will, however, consider these cases as independent. In Lamquin et al. (2008) collocations of AIRS and CALIPSO considered five pixels with an averaging of their vertical profiles and no large difference was inferred by considering the pixels independently.

2.3 Radiosonde data

We use relative humidity data obtained from radiosonde measurements made at the meteorological observatory Lindenberg near Berlin, Germany, between February 2000 and April 2001. Usually, radiosonde humidity measurements in the upper troposphere have problems; in particular, they are subject to a dry bias. Several correction methods exist in the literature (e.g. Soden and Lanzante, 1996; Wang et al., 2002), and our data have been corrected, too. The data and their handling as well as the correction method have been described in detail in (Nagel et al., 2001; Spichtinger et al., 2003a). The vertical resolution of the data is approximately 50 m. Ice supersaturation can be

detected and is present in 28% of the measured profiles, but it cannot be decided whether a data point was taken within a cloud or in clear sky. Most ice supersaturated layers are located between 200 and 450 hPa, the mean altitude is 300 hPa in summer and fall, and 340 hPa in winter and spring. For more details see Spichtinger et al. (2003a).

3 Supersaturation spinup

ECMWF uses an assimilation scheme that does not account for ice supersaturation in the upper troposphere. Hence, data assimilation leads to analyses that severely underestimate the true occurrence and range of ice supersaturation. Since the analyses serve as initial conditions for the forecast runs, the forecast model needs some time for spinup of the supersaturation field. Studies of upper tropospheric humidity should not use forecast humidity data from the spinup phase since they are unreliable.

We have investigated the spinup using forecasts from two months, namely October 2006 and April 2007. For every day, and for the noon and midnight forecast runs, we used forecasts up to day 3 and counted at each 3-h forecast step the number of grid boxes (in $0.5^\circ \times 0.5^\circ$ resolution) on the 250 hPa level where $RHi \geq 100\%$. This number, divided by the total number of grid boxes per level and averaged over all forecast runs for that month yields the fraction of supersaturated grid boxes, shown in the left hand panels of Fig. 3. The corresponding right hand panels show the average supersaturation in the grid boxes with supersaturation. The upper and lower curves in each panel are the mean plus/minus one standard deviation. We only studied global means, i.e. we did not look at single regions as e.g. northern midlatitudes, as we think one should use the humidity data only when the model has left the spinup phase everywhere. Figure 3 shows that the investigated quantities display similar spinup behaviour in the two months. There is a steep rise of the fraction of supersaturated grid boxes in about the first half day into the forecast. During that period the average supersaturation also shows an increase and a more noisy behaviour than later. Analogous behaviour was

ECMWF vs. AIRS UTH and CALIPSO clouds

N. Lamquin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



found for April 2007 on the 200 and 300 hPa levels (only one month studied with these levels, not shown) and for January 2007 on the 250 hPa level (not shown). Hence a spinup period of at least 12 h is necessary before using forecast humidity data from the upper troposphere. For the following investigations we use 24-h forecasts.

5 We note that the mean fraction of gridboxes with ice supersaturation is slightly higher than 10%, consistent with findings from airborne in situ data (Gierens et al., 1999). The mean supersaturation is of the order 6 to 10%, which is slightly lower than airborne and satellite measurements (e.g. Spichtinger et al., 2003b) indicate. This underestimation will be discussed in Sect. 4.3.

10 4 Evaluation of UTH

In the following, we first compare RH_i between ECMWF (RH_{iE}) and AIRS (RH_{iA}) in the upper troposphere for clear and cloudy situations. CALIPSO is used to investigate the effect of clouds in more detail. At last, the effect of vertical resolution on ice supersaturation occurrence is studied with the help of radiosonde data.

15 4.1 Comparison with AIRS relative humidity fields

This comparison involves the use of AIRS L2 cloud pressure and effective cloud cover. A layer is declared cloudy when the pressure of the highest cloud layer is between the pressures at the layer's top and bottom. Recent studies by Kahn et al. (2007) and Stubenrauch et al. (2008) have revealed an underestimation of AIRS L2 cloud pressure for clouds with pressure larger than 440 hPa. Figure 4 presents distributions of differences between AIRS cloud pressure and CALIPSO pressure of the middle of the cloud for two retrievals: the L2 standard product and the LMD retrieval presented in Stubenrauch et al. (2008), separately for high clouds ($p_{\text{AIRS}} < 400$ hPa) and clouds with p_{AIRS} between 400 and 500 hPa. Statistics is shown for AIRS–CALIPSO collocated data in the Northern Hemisphere (NH) midlatitudes for January and July 2007. Due

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to the slight underestimation of the AIRS L2 cloud pressure in the layer between 400 and 500 hPa we only keep data with clouds having L2 cloud pressure between 400 and 450 hPa in this specific pressure layer.

Figure 5 presents two-dimensional histograms of RH_{iA} from AIRS (RH_{iA}) versus RH_{iE} from ECMWF (RH_{iE}), separately for four pressure layers in the upper troposphere. The statistics is divided into three scene types, distinguished by AIRS L2: clear sky and two degrees of effective cloudiness (threshold at 30%) in the upper troposphere.

The frequencies of occurrence are normalized by the maximum value for each plot. This representation reveals two distinct modes: 1) a linear correlation between RH_{iA} and RH_{iE} for clear sky and low effective cloudiness and 2) a large range of RH_{iA} values when the RH_{iE} is around 100% and AIRS L2 effective cloudiness is larger than 30%. We analyze the two modes separately.

The first (“dry”) mode (with RH_{iE} lower than 80%) shows a good agreement with mean differences of $RH_{iE}-RH_{iA}$ found to be 2.7, 1.7, 3.4, 1.4% with standard deviations 10.4, 13.2, 15.1, 15.3%, respectively for the layers 200–250, 250–300, 300–400, and 400–500 hPa.

The second (“moist”) mode (with RH_{iE} around 100%) of Fig. 5 occurs when AIRS detects clouds. These situations ($RH_{iE} \in [90\%, 110\%]$) reveal a U-shape distribution of ECMWF cloud cover (see Fig. 6), with maxima at low and at large cloud cover. The fact that homogeneous ice nucleation needs high supersaturation (as implemented in the cloud scheme) explains that there are clear sky situations in spite of saturated or supersaturated air. When the cloud cover within an IFS grid box approaches 100%, then the relative humidity approaches 100% because in-cloud supersaturation (as observed for instance by Comstock et al., 2004; Lee et al., 2004; Ovarlez et al., 2002) cannot be represented in the current cloud scheme (Tompkins et al., 2007). This is highlighted in Fig. 7, presenting overall distributions of RH_{iE} and RH_{iA} in the pressure layer 300–400 hPa.

The IFS cloud cover distribution in Fig. 7 has larger values than the AIRS L2 effective cloud cover distribution, because high RH_i can be forecasted related to high cloud

**ECMWF vs. AIRS
UTH and CALIPSO
clouds**

N. Lamquin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cover, whereas an infrared sounder can determine RH_i (using cloud-cleared radiances) only when the cloudiness is not too high. Also, the effective cloud cover (weighted by cloud emissivity) of an optically thin cloud is much smaller than its cloud cover. Therefore, also the IFS cloud cover distribution for lower RH_i is broader than the AIRS effective cloud cover distribution.

To investigate RH_i of the moist mode further in relationship with clouds, we use in addition collocated AIRS – CALIPSO data. Figure 8 presents distributions of RH_{iA} when RH_{iE} > 80% compared to distributions of RH_{iA} in the presence of CALIPSO high clouds (as in Lamquin et al., 2008), separately for the same pressure layers as in Fig. 5. In all cases the distributions compare well, suggesting that the conditions “humidity around ice saturation” in the IFS and “cloud present” in CALIPSO lead to similar distributions of RH_{iA} which peak at about 80% and are very broad.

RH_{iA} appears on average drier than RH_{iE} in this second mode. One arguable reason is the limitation to scenes with relatively low effective cloudiness for AIRS humidity and temperature retrievals. Another reason could be the vertical resolution of the AIRS weighting functions of the channels used for the humidity retrieval, as the retrieval of humidity considers parts of the vertical profile both over and under the standard pressure levels. The effect of this has been demonstrated by Lamquin et al. (2008), showing that the mean relative humidity for cirrus scenes with clouds extending over the whole pressure layer is on average only about 70%. This value is slightly lower than the peak value of the distributions in Fig. 8, since clouds are on average geometrically thicker than the pressure layers (Lamquin et al., 2008).

Despite this dry AIRS bias of about 20%, on a statistical basis, the moist mode represents well how clouds are expected to appear in both datasets. This is encouraging. The comparison of AIRS and ECMWF humidities indicates problems with the vertical resolution of satellite data. A fairer comparison may be possible by computing the radiance fields from the model profiles and comparing these to the measured radiances. However, it is not obvious how histograms of radiance differences could be used for model improvements. For this purpose one may prefer to use instruments with higher

**ECMWF vs. AIRS
UTH and CALIPSO
clouds**

N. Lamquin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



vertical resolution. Towards this end we now use CALIPSO to relate RH_{iE} to a very vertically-resolved description of clouds.

4.2 Comparison with CALIPSO cloud vertical profiles

The use of CALIPSO data will give insights on how well cloud formation is represented in the model. All results are presented in Fig. 9.

In a first step (top left) we relate the high cloud (with apparent middle of the cloud $\rho_{\text{cloud}} = (\rho_{\text{top}} + \rho_{\text{base}}) / 2 < 500 \text{ hPa}$) occurrence as seen by CALIPSO with the maximum RH_{iE} forecasted between 200 and 500 hPa by the IFS (data in low vertical resolution). High cloud occurrence is declared as soon as one cloud layer from CALIPSO is observed at altitudes higher than 500 hPa. The maximum humidity is better suited than the average over the layer, because cloud formation is related to the humidity peak rather than to the average. To study the effect of temperature on this relationship we divided the data into two temperature intervals of roughly equal data amounts, $T < 220 \text{ K}$ and $T > 220 \text{ K}$, T corresponding to the level of maximum RH_{iE} . The figure shows that CALIPSO high cloud occurrence increases with maximum RH_{iE} , for both temperature intervals. Note that there are a few cases of clouds for which the model predicts low humidity values. This might be very optically thin (but nevertheless geometrically thick, see below) clouds with low ice concentration or outflows from larger cloud complexes that protrude into a dry environment. The cloud occurrence at $\text{RH}_{iE} = 100\%$ is about 65%, and 100% cloud occurrence is only reached at certain supersaturations that reflect the supersaturation thresholds for homogeneous nucleation together with the $\pm 20\%$ humidity fluctuations in the model. Larger maximum RH_{iE} values are reached in the cold temperature interval than in the warmer interval, since the humidity threshold for homogeneous nucleation increases with decreasing temperature.

In a second step (top right) the total vertical extent of high clouds from CALIPSO is related to the maximum RH_{iE} . We define the total vertical extent as the sum of the vertical extent (in m) of all high cloud layers detected by CALIPSO. In both temperature intervals the total extent increases with the maximum RH_{iE} . This agrees qualitatively

ECMWF vs. AIRS UTH and CALIPSO clouds

N. Lamquin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with the results of Lamquin et al. (2008). One observes that the average thickness of the clouds detected by CALIPSO in spite of a dry forecast is of the order 1500 m.

For single-layer high clouds detected by CALIPSO, Fig. 9 presents the distribution of the difference between their altitude and the altitude of the middle of the layer in which the maximum RH_{iE} is found (bottom left). The distribution is symmetric with a peak at zero. The standard deviation of the distribution reflects rather the resolution of the standard pressure levels than the actual data scatter. The model predicts the highest humidity values mostly at the correct altitude, namely at that altitude where CALIPSO finds a cloud.

Finally (bottom right), a distribution of the maximum of RH_{iE} between 200 and 500 hPa is shown for scenes of high clouds detected by CALIPSO. The peak probability is around 100% which shows that this particular value in the model is well representative for the presence of high clouds. The distribution is strongly left-skewed because of dry cases for which a high cloud was detected by CALIPSO. The standard deviation is 19%, comparable to the distribution widths for in-cloud humidity distributions reported by Ovarlez et al. (2002) and Immler et al. (2007).

Summarising, the IFS model predicts high RH_i where CALIPSO actually detects high clouds. This lends credence to the good quality of the ECMWF upper tropospheric cloud forecast. However, sometimes clouds are observed where ECMWF predicts dry air. Such cases need further consideration.

4.3 Influence of vertical resolution on occurrence of ice supersaturation

In this section we investigate effects of vertical resolution on the apparent frequency of ice supersaturation. This problem is important especially for satellite IR sounders, since ice supersaturated layers are frequently much shallower (see Spichtinger et al., 2003a) than the weighting functions of satellite instrument channels (Gierens et al., 2004).

Since ECMWF humidity data exist in both high and low vertical resolution we will first investigate the effect with these. Generally we expect that interpolation leads to a loss

ECMWF vs. AIRS UTH and CALIPSO clouds

N. Lamquin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of detail in the interpolated profiles, single shallow supersaturated layers will not be detected at low vertical resolution, resulting in an underestimated overall occurrence of ice supersaturation. We expect the same effect to occur when using satellite IR sounder data, because the broad weighting functions average thin supersaturated layers within thicker subsaturated layers (Gierens et al., 2004; Kahn et al., 2008; Lamquin et al., 2008).

The two upper panels of Fig. 10 present the probability of occurrence of ice supersaturation within a layer (determined from ECMWF data with high vertical resolution) as a function of RH_{iE} averaged over this layer. Results are shown for layers 400–500, 300–400, and 250–300 hPa in January and for layers 300–400, 250–300, and 200–250 hPa in July (as the tropopause is higher than in January). To avoid difficulties inferred by the tropopause eventually lying inside the pressure layer we only select cases for which the temperature gradient is constantly negative throughout the layer. One observes that supersaturation at the high resolution vertical levels can be present as soon as the data on the standard levels indicate $RH_{iE} \gtrsim 50\%$. In all cases the probability to have a supersaturated layer at 25 hPa resolution follows an s-shaped function of RH_{iE} . The curves increase sharply just below $RH_{iE} = 100\%$ and reach 100% probability when the low resolution RH_{iE} is slightly supersaturated, around 105–110%. That 100% probability is not already reached at $RH_{iE} = 100\%$ is certainly an effect of the two different interpolation schemes used to compute the low- and high-resolution profiles. This is supported by Fig. 2 when the high-resolution relative humidity is at 100%: low-resolution relative humidity is on average slightly higher than 100%.

If one can find a general expression for this s-shaped behaviour of the probability function due to low vertical resolution, it could be used for correcting apparent frequencies of supersaturation obtained from satellite IR sounder instruments. A common way to estimate the frequency of occurrence of ice supersaturation is to select a threshold lower than 100% (as in Stubenrauch and Schumann, 2005 and Rädcl and Shine, 2008) and to count all cases with RH_i exceeding the threshold as supersaturated. The threshold can be obtained by considering the distribution of RH_i for cloudy cases, which

**ECMWF vs. AIRS
UTH and CALIPSO
clouds**

N. Lamquin et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

is biased dry, because the value of RHi at the peak of this distribution should in reality correspond to 100%. However, Rädcl and Shine (2008) use a method based on hit and rejection rates as well as Peirce skill scores (Peirce, 1884). The use of such threshold is subject to the risk of false alarms.

To study this behaviour further, we also analyzed the radiosonde data from Lindenberg described in Sect. 2.4. We determine RHi at each standard pressure level by averaging the specific humidity inside the layer and by integrating the saturation specific humidity (as in Lamquin et al., 2008) between the edges of the layer. The high-resolution radiosonde levels give the opportunity to detect local ice supersaturation within the boundaries of a layer, and its probability of occurrence is displayed as a function of RHi averaged over the whole layer. The plots are shown on bottom of Fig. 10; they show a similar s-function as before even though the scheme and the source of the data employed are different. This confirms that, on a statistical basis, ice supersaturation may be present even when small low-resolution values of RHi are given. It seems possible to replace the use of a fixed threshold for the determination of ice supersaturation frequency by a probability function. Its shape may vary from one type of data to another but in any case may be close to an s-function.

The empirical s-functions start at zero at low RHi and reach unity at saturation (or slightly above). Furthermore, they are monotonically increasing, that is, they have the mathematical properties of a cumulative distribution function, $F(r)$. $F(r)$ is the probability that the high-resolution humidity profile exceeds saturation when the low-resolution humidity is r . The derivative of F is a probability density function, $f(r)$ and we have $\int_0^r f(r')dr' = F(r)$. We are now seeking a suitable interpretation, which then might help to simulate the s-function for later applications. A suitable interpretation is the following: when scanning data from small to higher values of r , $f(r)dr$ is the probability that the high-resolution humidity profile exceeds saturation for the first time in the infinitesimal interval $[r, r+dr]$. For values of r exceeding 100% this probability tends to zero because then the high-resolution humidity profile always exceeds saturation.

Finally, we test the application of the s-function to determine the occurrence of ice

**ECMWF vs. AIRS
UTH and CALIPSO
clouds**N. Lamquin et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

supersaturation in the upper troposphere by comparing the result to the one obtained by applying a simple threshold. Since the empirical s-function is not easy to relate to a simple function (it is not symmetrical enough to be modelled by a logistic function), we use such a function empirically obtained for all layers combined. We determine the frequency of ice supersaturation for all pressure layers for the year of ECMWF low resolution data over Europe by using the s-function or thresholds at 80% (as in Rädcl and Shine, 2008) and 100%: the use of the s-function leads to about 20% supersaturation occurrence while the use of thresholds at 80% or 100% lead, respectively to 25% and 9% supersaturation occurrence. In a recent work (Burkhardt et al., 2008) ice supersaturation occurrence in the midlatitudes upper troposphere is found to be around 20% with large seasonal variations using MOZAIC and AIRS data. As expected, the 100% threshold leads to a severe underestimation of the true probability of ice supersaturation. Using a fixed threshold of 80% leads to a slightly higher value than using the s-function. One can argue that ice supersaturation occurrence from AIRS in Burkhardt et al. (2008) does not use any correction algorithm but still compares well to MOZAIC in the midlatitudes. However, the seasonal variabilities between the two datasets do not fit perfectly as the peak occurrence is found during winter for MOZAIC and during spring for AIRS. We then underline that our estimation from ECMWF using the s-function compares better to MOZAIC, all the more because we observe a comparable seasonal cycle of ice supersaturation occurrence. Burkhardt et al. (2008) find (DJF) 26–27% (MAM) 18–19% (JJA) 17–18% (SON) 23–24% with MOZAIC while we find (DJF) 23% (MAM) 19% (JJA) 15% (SON) 22%.

5 Conclusions

Upper tropospheric humidity and cloudiness forecasts from ECMWF's Integrated Forecast model (IFS) including the new ice supersaturation feature (Tompkins et al., 2007) have been compared with collocated humidity and cloud retrievals from AIRS and CALIPSO over Europe. An initial study of the global supersaturation spinup behaviour

ECMWF vs. AIRS UTH and CALIPSO clouds

N. Lamquin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



shows that at least 12 h are necessary before using forecast humidity data from the upper troposphere. Relative humidity of IFS and AIRS was compared in different pressure layers, separately for clear and cloudy situations distinguished by AIRS. Two modes were detected: 1) a dry mode in which IFS predicts $RH_{iE} < 80\%$ (and mainly cloud free) and in which the relative humidities show a good agreement (with standard deviations of the order 10 to 15%, the order of humidity measurement uncertainties) and 2) a moist mode in which IFS predicts values around ice saturation and AIRS provides a range of RH_i values from about 50% up to and exceeding 150% with a peak probability around 80–90%. It may be noted that AIRS always detects clouds in these cases. A comparison of IFS relative humidities with the cloud products from CALIPSO showed a strong positive correlation between RH_{iE} and the probability that CALIPSO detects a cloud in the respective layer. The peak relative humidity from IFS is mostly located in the pressure layer where CALIPSO indeed detects a cloud. The CALIPSO cloud probability reaches 100% when the IFS humidity approximately reaches the threshold for homogeneous nucleation; the cloud probability at ice saturation is 65%. The comparison uncovers that occasionally CALIPSO finds clouds (of geometrical thickness exceeding 1 km) where IFS predicts dry air. These cases need further consideration.

Finally we tested the dependence of vertical resolution on the reported frequency of ice supersaturation. This is a problem in particular for satellite sounder observations, but also for the forecast data when only the standard pressure levels are retained. We compared the IFS data on the standard pressure levels with those of higher vertical resolution and high-resolution radiosonde data with layer averages obtained in a way close to how upper tropospheric humidity is determined for AIRS. These exercises demonstrated that the true frequency of occurrence as a function of the low-resolution relative humidity follows a s-shaped function, that can be used for a correction algorithm. Application of such a correction yielded a slightly lower values than the method using a fixed RH_i threshold which corresponds to the maximum of the RH_i distribution of cloudy scenes. The s-function correction should be further tested with more data and its possible connection to extreme-value theory should be explored as well. It

**ECMWF vs. AIRS
UTH and CALIPSO
clouds**N. Lamquin et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

might turn out as a valuable tool for predicting the true number of exceedances over a threshold when only coarse-resolution data profiles are available for analysis.

Acknowledgements. The first author thanks his colleagues at Deutsches Zentrum für Luft- und Raumfahrt (DLR) for their warm and friendly welcome, as this article is the outcome of a collaboration in the framework of the Monge PhD grant of Ecole Polytechnique. The AIRS version 5 data were obtained through the Goddard Earth Sciences Data and Information Services (<http://daac.gsfc.nasa.gov/>) and the CALIPSO data were obtained through the Atmospheric Sciences Data Center (ASDC) at NASA Langley Research Center by the ICARE Thematic Center created by CNES (<http://www-icare.univ-lille1.fr/>) and its interface ClimServ created for Institut Pierre Simon de Laplace (<http://climserv.ipsl.polytechnique.fr/>). ECMWF data have been retrieved within the framework of a special project on “Ice Supersaturation and Cirrus Clouds (SPDEISSR)”. The authors thank all the corresponding teams for assistance and public release of data products. The work contributes to COST Action ES-0604 “Water Vapour in the Climate System (WaVaCS)” and to the aforementioned SPDEISSR.



The publication of this article is financed by CNRS-INSU.

References

- Burkhardt, U., Kärcher, B., Ponater, M., Gierens, K., and Gettelman, A.: Contrail cirrus supporting areas, *Geophys. Res. Lett.*, 35, L16808, doi:10.1029/2008GL034056, 2008. 17922
- Chahine, M. T. and Coauthors: AIRS: Improving weather forecasting and providing new data on greenhouse gases, *B. Am. Meteorol. Soc.*, 87, 911–926, 2006. 17911
- Comstock, J., Ackerman, T. P., and Turner, D. D.: Evidence of high ice supersaturation in cirrus clouds using ARM Raman lidar measurements, *Geophys. Res. Lett.*, 31, L11106, doi:10.1029/2004GL019705, 2004. 17916
- Divakarla, M. G., Barnet, C. D., Goldberg, M. D., McMillin, L. M., Maddy, E., Wolf, W., Zhou, L., and Liu, X.: Validation of Atmospheric Infrared Sounder temperature and water vapour

17924

ACPD

8, 17907–17937, 2008

ECMWF vs. AIRS UTH and CALIPSO clouds

N. Lamquin et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- retrievals with matched radiosondes measurements and forecasts, *J. Geophys. Res.*, 111, D09S15, doi:10.129/205JD006116, 2006. 17911
- Gettelman, A., Weinstock, E. M., Fetzner, E. J., et al.: Validation of Aqua satellite data in the upper troposphere and lower stratosphere with in situ aircraft instruments, *Geophys. Res. Lett.*, 31, L22107, doi:10.1029/2004GL020730, 2004. 17911, 17912
- Gierens, K., Schumann, U., Helten, M., Smit, H., and Marenco, A.: A distribution law for relative humidity in the upper troposphere and lower stratosphere derived from three years of MOZAIC measurements, *Ann. Geophys.*, 17, 1218–1226, 1999, <http://www.ann-geophys.net/17/1218/1999/>. 17915
- Gierens, K., Kohlhepp, R., Spichtinger, P., and Schroedter-Homscheidt, M.: Ice supersaturation as seen from TOVS, *Atmos. Chem. Phys.*, 4, 539–547, 2004, <http://www.atmos-chem-phys.net/4/539/2004/>. 17919, 17920
- Immler, F., Treffeisen, R., Engelbart, D., Krüger, K., and Schrems, O.: Cirrus, contrails, and ice supersaturated regions in high pressure systems at northern mid latitudes, *Atmos. Chem. Phys.*, 8, 1689–1699, 2008, <http://www.atmos-chem-phys.net/8/1689/2008/>. 17919
- Janisková, M., Mahfouf, J.-F., Morcrette, J.-J., and Chevallier, F.: Linearized radiation and cloud schemes in the ECMWF model: Development and evaluation, *Q. J. Roy. Meteor. Soc.*, 128, 1505–1527, 2002. 17909
- Kahn, B. H., Chahine, M. T., Stephens, G. L., Mace, G. G., Marchand, R. T., Wang, Z., Barnet, C. D., Eldering, A., Holz, R. E., Kuehn, R. E., and Vane, D. G.: Cloud type comparisons of AIRS, CloudSat, and CALIPSO cloud height and amount, *Atmos. Chem. Phys.*, 8, 1231–1248, 2008, <http://www.atmos-chem-phys.net/8/1231/2008/>. 17915
- Kahn, B. H., Liang, C. K., Eldering, A., Gettelman, A., Yue, Q., and Liou, K. N.: Tropical thin cirrus and relative humidity observed by the Atmospheric Infrared Sounder, *Atmos. Chem. Phys.*, 8, 1501–1518, 2008, <http://www.atmos-chem-phys.net/8/1501/2008/>. 17920
- Kärcher, B. and Lohmann, U.: A parametrization of cirrus cloud formation: Homogeneous freezing of supercooled aerosols, *J. Geophys. Res.*, 107(D2), 4010, doi:10.1029/2001JD000470, 2002.
- Lamquin, N., Stubenrauch, C. J., and Pelon, J.: Upper tropospheric humidity and cirrus geometrical and optical thickness: Relationships inferred from 1 year of collocated AIRS and

**ECMWF vs. AIRS
UTH and CALIPSO
clouds**N. Lamquin et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- CALIPSO data, *J. Geophys. Res.*, 113, D00A08, doi:10.1029/2008JD010012, 2008. 17911, 17913, 17917, 17919, 17920, 17921
- Lee, S.-H., Wilson, J. C., Baumgardner, D., Herman, R. L., Weinstock, E. M., LaFleur, B. G., Kok, G., Anderson, B., Lawson, P., Baker, B., Strawa, A., Pittman, J. V., Reeves, J. M., and Bui, T. P.: New particle formation observed in the tropical/subtropical cirrus clouds, *J. Geophys. Res.*, 109, D20209, doi:10.1029/2004JD005033, 2004. 17916
- Luo, Z., Kley, D., Johnson, R. H., and Smit, H. G. J.: Ten years of measurements of tropical upper-tropospheric water vapor by MOZAIC. Part II: assessing the ECMWF humidity analysis, *J. Climate*, 21, 1449–1466, 2007. 17909
- Nagel, D., Leiterer, U., Dier, H., Kats, A., Reichardt, J., and Behrendt, A.: High accuracy humidity measurements using the standardized frequency method with a research upper-air sounding system, *Meteorol. Z.*, 10, 395–405, 2008. 17913
- Ovarlez, J., Gayet, J.-F., Gierens, K., Ström, J., Ovarlez, H., Auriol, F., Busen, R., and Schumann, U.: Water vapor measurements inside cirrus clouds in northern and southern hemispheres during INCA, *Geophys. Res. Lett.*, 29(16), 1813, doi:10.1029/2001GL014440, 2002. 17916, 17919
- Peirce, C.: The numerical measure of the success of predictions, *Science*, 12, 453–545, 1884. 17921
- Rädel, G. and Shine, K. P.: Radiative forcing by persistent contrails and its dependence on cruise altitudes, *J. Geophys. Res.*, 113, D07105, doi:10.1029/2007JD009117, 2008. 17920, 17921, 17922
- Read, W. G., Lambert, A., Bacmeister, J., et al.: Aura Microwave Limb Sounder upper tropospheric and lower stratospheric H₂O and relative humidity with respect to ice validation, *J. Geophys. Res.*, 112, D24S35, doi:10.1029/2007JD008752, 2007. 17911, 17912
- Sassen, K.: The polarization lidar technique for cloud research: A review and current assessment, *B. Am. Meteorol. Soc.*, 72, 1848–1866, 1991. 17912
- Soden, B. J. and Lanzante, J. R.: An assessment of satellite and radiosonde climatologies of upper-tropospheric water vapor, *J. Climate*, 9, 1235–1250, 1996. 17913
- Sonntag, D.: Important new values of the physical constants of 1986, vapor pressure formulation based on the ITS-90 and psychrometer formulae, *Meteorol. Z.*, 70, 340–344, 1990. 17912
- Spichtinger, P., Gierens, K., Leiterer, U., and Dier, H.: Ice supersaturation in the tropopause region over Lindenberg, Germany, *Meteorol. Z.*, 12, 143–156, 2003. 17913, 17914, 17919

**ECMWF vs. AIRS
UTH and CALIPSO
clouds**N. Lamquin et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Spichtinger, P., Gierens, K., and Read, W.: The global distribution of ice-supersaturated regions as seen by the Microwave Limb Sounder, *Q. J. Roy. Meteor. Soc.*, 129, 3391–3410, 2003. 17915
- 5 Stephens, G., Vane, D. G., Boain, R. J., et al.: The CloudSat mission and the A-train, *B. Am. Meteorol. Soc.*, 83, 1771–1790, 2002. 17911
- Stubenrauch, C. J. and Schumann, U.: Impact of air traffic on cirrus coverage, *Geophys. Res. Lett.*, 32, L14813, doi:10.1029/2005GL022707, 2005. 17911, 17920
- Stubenrauch, C. J., Cros, S., Lamquin, N., Armante, R., Chédin, A., Crevoisier, C., and Scott, N. A.: Cloud properties from AIRS and evaluation with CALIPSO, *J. Geophys. Res.*, in press, 10 2008. 17915
- Susskind, J., Barnet, C., and Blaisdell, J.: Retrieval of atmospheric and surface parameters from AIRS/AMSU/HSB data in the presence of clouds, *IEEE Trans. Geosci. Remote Sens.*, 41, 390–409, 2003. 17911
- Susskind, J., Barnet, C., Blaisdell, J., Iredell, L., Keita, F., Kouvaris, L., Molnar, G., and Chahine, 15 M.: Accuracy of geophysical parameters derived from AIRS/AMSU as a function of fractional cloud cover, *J. Geophys. Res.*, 111, D09S17, doi:10.1029/2005JD006272, 2006. 17911, 17912
- Tobin, D. C., Revercomb, H. E., Knuteson, R. O., et al.: Atmospheric radiation measurement site atmospheric state best estimates for Atmospheric Infrared Sounder temperature and wa- 20 ter vapor retrieval validation, *J. Geophys. Res.*, 111, D09S14, doi:10.1029/2005JD006103, 2006. 17911, 17912
- Tompkins, A. M., Gierens, K., and Rädel, G.: Ice supersaturation in the ECMWF Integrated Forecast System, *Q. J. Roy. Meteorol. Soc.*, 133, 53–63, 2007. 17908, 17909, 17916, 17922
- Wang, J. H., Cole, H. L., Carlson, D. J., Miller, E. R., Beierle, K., Paukkunen, A., and Laine, 25 T. K.: Corrections of humidity measurement errors from the Vaisala RS80 radiosonde – Application to TOGA COARE data, *J. Atmos. Ocean. Tech.*, 19, 981–1002, 2002. 17913
- Winker, D. M., Pelon, J., and McCormick, M. P.: The CALIPSO mission: Spaceborne lidar for observation of aerosols and clouds, *Proceedings of SPIE*, 4893, 1–11, 2003. 17912, 17913
- Winker, D. M., Hunt, W. H., and McGill, M. J.: Initial performance assessment of CALIOP, 30 *Geophys. Res. Lett.*, 34, L19803, doi:10.1029/2007GL030135, 2007. 17912

**ECMWF vs. AIRS
UTH and CALIPSO
clouds**N. Lamquin et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ECMWF vs. AIRS UTH and CALIPSO clouds

N. Lamquin et al.




Fig. 1. Sketch of the different relative humidity products, AIRS provides humidity fields integrated over pressure layers and ECMWF humidity fields are given at pressure levels either at low or high vertical resolution. For comparison with AIRS the relative humidities from ECMWF are averages between edges of the layers.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**ECMWF vs. AIRS
UTH and CALIPSO
clouds**

N. Lamquin et al.




Fig. 2. Two-dimensional histogram of the difference between the interpolated low-resolution relative humidity and the original high-resolution relative humidity (y-axis) against the high-resolution relative humidity (x-axis) at standard pressure levels 200, 250, 300, 400, and 500 hPa. The probability is normalized by the maximum value (in percent) and the colour scale is limited to 50% with values higher than 50% represented by the same colour. All levels combined, January and July 2007 combined. The solid line represents binned mean, dashed lines represent binned mean \pm standard deviation.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ECMWF vs. AIRS
UTH and CALIPSO
clouds

N. Lamquin et al.




Fig. 3. Fraction of grid boxes with ice supersaturation (left panels) and mean supersaturation in these grid boxes (right panels), averaged over all forecast runs (noon and midnight each day) of ECMWF for October 2006 and April 2007, in $0.5^\circ \times 0.5^\circ$ resolution globally. The upper and lower curve in each panel are the respective mean plus/minus one standard deviation. All data refer to the 250 hPa pressure level.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ECMWF vs. AIRS
UTH and CALIPSO
clouds

N. Lamquin et al.




Fig. 4. Difference in cloud pressure between AIRS and CALIPSO for clouds higher than 400 hPa (top) and clouds between 400 and 500 hPa (bottom). Cloud pressures from AIRS determined from LMD retrieval (plain) and from AIRS level 2 (dashed). NH midlatitudes, January and July 2007.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ECMWF vs. AIRS
UTH and CALIPSO
clouds

N. Lamquin et al.




Fig. 5. Two-dimensional histograms of RH_A (y-axis) versus RH_E (x-axis) for four pressure layers: 200–250 hPa (top), 250–300 hPa, 300–400 hPa, and 400–500 hPa (bottom) in clear sky (left) and with average AIRS L2 effective cloud cover $<30\%$ (middle) and $>30\%$ (right) and cloud pressure within the pressure layer. A 1-to-1 line or 0-line are also shown for comparison.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ECMWF vs. AIRS UTH and CALIPSO clouds

N. Lamquin et al.




Fig. 6. Left: distributions of effective cloud cover from AIRS level 2 for high clouds ($p_{\text{cld}} < 500$ hPa) for three categories of retrieved relative humidity wrt ice: $\text{RH}_{iA} < 50\%$, $50\% < \text{RH}_{iA} < 80\%$ and $\text{RH}_{iA} > 80\%$. Right: distribution of cloud cover from ECMWF for two categories of relative humidity wrt ice $\text{RH}_{iE} < 50\%$ and $\text{RH}_{iE} \in [90\%, 110\%]$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**ECMWF vs. AIRS
UTH and CALIPSO
clouds**

N. Lamquin et al.




Fig. 7. Overall distributions of relative humidity wrt ice from AIRS (plain) and ECMWF (dashed) for the layer 300–400 hPa

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ECMWF vs. AIRS
UTH and CALIPSO
clouds

N. Lamquin et al.




Fig. 8. Distributions of relative humidity wrt ice from AIRS in two cases: collocated clouds from CALIPSO (plain) and collocated with $\text{RH}_{\text{I}_E} > 80\%$ (dashed). Four pressure layers: 200–250 hPa (top left), 250–300 hPa, 300–400 hPa, and 400–500 hPa (bottom right).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ECMWF vs. AIRS
UTH and CALIPSO
clouds

N. Lamquin et al.




Fig. 9. Results from the CALIPSO-ECMWF collocation. Top left: occurrence of high clouds ($p_{\text{clid}} < 500$ hPa) seen by CALIPSO as a function of the maximum relative humidity wrt ice seen between 200 and 500 hPa by ECMWF for two temperature ranges $T < 220$ K and $T > 220$ K. Top right: same but showing the total thickness of clouds seen by CALIPSO. Bottom left: distribution of the distance (in hPa) between the apparent middle of CALIPSO single-layer clouds and the apparent middle of the pressure layer where the maximum relative humidity wrt ice (between 200 and 500 hPa) seen from ECMWF is located. Bottom right: distribution of the maximum relative humidity wrt ice from ECMWF in the presence of collocated CALIPSO high clouds.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ECMWF vs. AIRS UTH and CALIPSO clouds

N. Lamquin et al.




Fig. 10. Top: probability of occurrence of ice supersaturation on the high-resolution vertical levels as a function of the relative humidity wrt ice on the low-resolution vertical levels, January and July 2007, IFS data. Bottom: probability of occurrence of ice supersaturation within thin (≈ 50 m) layers seen from corrected radiosondes as a function of the relative humidity computed for the whole layer.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

