

This discussion paper is/has been under review for the journal *Atmospheric Chemistry and Physics (ACP)*. Please refer to the corresponding final paper in *ACP* if available.

**Contribution of
processes affecting
atmospheric
dynamics**

M. Gonçalves et al.

Contribution of atmospheric processes affecting the dynamics of air pollution in south-western Europe during a typical summertime photochemical episode

M. Gonçalves¹, P. Jiménez-Guerrero², and J. M. Baldasano^{1,2}

¹Projects Engineering Department, Technical University of Catalonia, Barcelona, Spain

²Barcelona Supercomputing Center, Barcelona, Spain

Received: 15 July 2008 – Accepted: 25 August 2008 – Published: 23 October 2008

Correspondence to: J. M. Baldasano (jose.baldasano@bsc.es)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

The southern Mediterranean region frequently experiences critical levels of photochemical pollutants during summertime. In order to account for the contribution of different atmospheric processes during this type of episodes, the WRF-ARW/HERMES/CMAQ modelling system was applied with high resolution (1 km², 33 sigma vertical layers, 1 h) to assess the different dynamics in a coastal environment and an inland-continental zone: the North-eastern and Central Iberian Peninsula (NEIP and CIP, respectively). The former is characterized by a very complex terrain, while the latter behaves as a flat area, which clearly affects the pattern of local flows. A representative type of photochemical pollution episode (occurring over 78% of summer days) which occurred during 17–18 June 2004 is selected as the study period. The CMAQ Integrated Process Rate provides the hourly contributions of atmospheric processes to net O₃, NO_x and NMVOCs concentrations. The O₃ photochemical formation occurs mainly in downwind areas from the main NO_x emission sources during midday. At surface level it accounts for 50 to 75 μg m⁻³ h⁻¹. The urban areas and main roads, as main sources of NO_x emissions, act as O₃ sinks, quenching up to -200 μg m⁻³ per hour during the traffic circulation peaks. The O₃ concentration gradient generated, larger during daytime, increases the contribution of diffusion processes to ground-level O₃ (up to 200 μg m⁻³ h⁻¹ fluxes, mainly from upper vertical layers). The maximum positive contributions of gas-phase chemistry to O₃ occur in the coastal domain at high levels, while in the continental domain they take place in the whole atmospheric column under the PBL. The transport of ozone precursors by advective flows sets the location of the maximum O₃ surface concentrations. The O₃ chemical formation involves the oxidation of less NMVOCs in the NEIP than in the CIP domains, due to differences in chemical sensitivity between these areas. The dry deposition is an important sink in the lowest layer of the model, together with vertical diffusion flows. Finally, the contributions from cloud processes, wet deposition and heterogeneous chemistry are negligible during the whole episode, characterized by a high solar radiation and neither precipitation

ACPD

8, 18457–18497, 2008

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

nor cloudiness. This process analysis provides new quantitative information about the origin of the peaks of O_3 and its precursors, aiding the design of abatement strategies in south-western Europe.

1 Introduction

5 Atmospheric chemistry transport model simulations suggest that summertime O_3 is enhanced in the entire Mediterranean troposphere, contributing substantially to the radiative forcing of climate and air quality issues (Lawrence et al., 1999; Hauglustaine and Brasseur, 2001; Jiménez et al., 2006). Specifically the south-western Europe and the Iberian Peninsula (IP) frequently experience photochemical pollution episodes
10 during summertime, in which O_3 and PM_{10} exceedances of the European air quality targets are registered. Atmospheric modelling is a fundamental tool to assess the air quality levels in these situations. Nevertheless, the air quality models usually provide the net concentration of pollutants, without the capabilities of understanding and isolating the atmospheric processes involved, which would explain the reasons for a model's
15 predictions and show the relative importance of each process (Jeffries and Tonnesen, 1994; Jang et al., 1995a, b). For the Mediterranean area, these issues have been studied by measurement campaigns, but also combining both modelling and experimental techniques (e.g. Millán et al., 1996; 1997; Lelieveld et al., 2002; Lawrence et al., 2003; Roelofs et al., 2003; Cros et al., 2004; Dufour et al., 2005; Jonson et al., 2006, among
20 others). Currently, some Eulerian grid models, such as the Models-3 Community Multiscale Air Quality Model – CMAQ (Gipson, 1999), permits to assess the contribution of different atmospheric processes to the net concentrations of pollutants.

This analysis is useful not only for a better understanding of the atmospheric behaviour of an specific area, i.e. New York (Jang et al., 1995a, b) or the Puget Sound
25 Region, USA (Jiang et al., 2002); but also for assessing which are the models or the models' configurations presenting better skills (Jeffries and Tonnesen, 1994; O'Neil and Lamb, 2005). The peer-reviewed works applying the process analysis focus mainly

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



on episodic events (San José et al., 2002; O’Neil and Lamb, 2005), although annual (Zhang et al., 2006) and even climatologic simulations (Hogrefe et al., 2005) have been also performed.

This work pretends to assess and quantify the contribution of different atmospheric processes to the concentration of O₃ and its precursors in south-western Europe by means of the Integrated Process Rate (IPR) analysis tool available in the CMAQv4.6 model. In order to highlight the different behaviour of atmospheric dynamics in a coastal environment and an inland-continental zone, the North-eastern and Central Iberian Peninsula domains (NEIP and CIP, respectively) have been selected. These areas show different topographic patterns, the former being a coastal region characterized by a very complex terrain and the latter a continental inner region with a much simpler topography, which brings different locally-driven flows. Moreover, they house the largest Spanish urban centres, Barcelona and Madrid, and therefore photochemical pollution episodes are of special concern by their direct effects on population.

2 Methods

2.1 Modeling system

The WRF-ARW/HERMES-2004/CMAQ modelling system is applied with high spatial (1 km²) and temporal resolution (1 h) to the study areas. The fine scale used is essential in complex terrains such as those studied (Jiménez et al., 2005, 2006).

The Weather Research and Forecasting (WRF) Model (Michalakes et al., 2005; Skamarock et al., 2005) provides the meteorology parameters as inputs to CMAQ. The initial and boundary conditions are obtained from the National Centers for Environmental Prediction (NCEP) reanalysis data (available at the standard pressure levels for every 6 h with 0.5×0.5 degree resolution). Four nested domains are defined over each study area (Table 1), covering the final domains (D4; Fig. 1): the NEIP (322×259 km²) and the CIP (181×214 km²), respectively. 33 sigma vertical layers cover the troposphere

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

(up to 50 hPa \approx 20 km), with 12 layers under the PBL.

The High Elective Resolution Emission Modelling System (HERMES) has been developed for Spain with a high spatial (1 km²) and temporal (1 h) resolution (Baldasano et al., 2008). This model focuses on the estimation of gas and particulate matter pollutants, including the ozone precursors. HERMES considers the emissions from the following sources:

1. power generation plants,
2. industrial installations,
3. domestic and commercial fossil fuel use,
4. domestic and commercial solvents use,
5. road transport,
6. ports,
7. airports and
8. biogenic emissions;

using a bottom-up approach except for the domestic and commercial fossil fuel use, where a top-down approach was adopted and regional emissions were allocated to fine grid cells by surrogate indexes. It follows the methodologies and criteria of previous emission models developed for the eastern Iberian Peninsula: EMICAT2000 (Parra et al., 2004, 2006) and EMIVAL (Arévalo et al., 2004). The reference year chosen is 2004, since it is the most recent year in which all the needed data for the HERMES development are available. HERMES generates results according to the European Environmental Agency's Selected Nomenclature for Air Pollution (SNAP). Furthermore, HERMES has the capacity of presenting results according to individual installations, industrial activities, land use classification or type of pollutant or process (fugitive, evaporative, hot or cold emissions).

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Contribution of
processes affecting
atmospheric
dynamics**M. Gonçalves et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The chemistry transport model used to compute the concentrations of photochemical pollutants is CMAQ (Byun and Schere, 2006). The initial and boundary conditions were derived from a one-way nested simulation covering a domain of $1392 \times 1104 \text{ km}^2$ centred in the Iberian Peninsula, that used EMEP emissions for 2004, disaggregated to 18 km. A 48-h spin-up was performed to minimize the effects of initial conditions for the final domains. The chemical mechanism selected for the simulations following the criteria of Jiménez et al. (2003) was Carbon Bond IV (CBM-IV) (Gery et al., 1989) including aerosols and heterogeneous chemistry. The CMAQ model configuration uses the Yamartino-Blackman Cubic scheme (YAM) for the advection and convection transport and the Eddy diffusion scheme for the vertical and horizontal diffusion. NO_x and volatile organic compounds (VOC) speciation of HERMES emissions are detailed in Parra et al. (2006). The vertical and horizontal resolution of the final domains (D4; Fig. 1) are the same as those used in the meteorological simulation (1 km^2 , 33 vertical layers) and the species concentration and atmospheric processes contributions are estimated hourly.

The high resolution employed and the huge number of variables involved in the atmospheric integrations (not only the pollutants concentrations, but also the process analysis) require high-performance computing. The availability of the MareNostrum supercomputer hold in the BSC-CNS, together with the advances in the parallelization of air quality codes, has allowed these simulations.

2.2 Integrated process rate (IPR)

The contributions of different processes are assessed by means of the Integrated Process Rate (IPR) analysis available in the CMAQv4.6 model. It provides the effects of all the physical processes and the net effect of chemistry on model predictions, which is the fraction of the mass or concentration in each model cell (or group of cells) in terms of the process that gave rise to this portion of the mass or concentration. More details can be found in Jeffries and Tonnesen (1994); Jang et al. (1995a, b) and Gipson (1999). The contributions are estimated for each cell (1 km^2) on a hourly basis, providing the

gas-phase chemistry, the net transport (horizontal and vertical advection), horizontal and vertical diffusion, emissions, dry and wet deposition and clouds interactions contributions to the net concentration of pollutants. The main assumptions of the modelling system are summarized below in order to clarify the subsequent discussion of results (Gipson, 1999):

1. Net transport: It considers horizontal and vertical advection sum, which is the transport of pollutants due to the mean wind fields. The advection scheme used is globally mass-conserving. The horizontal advection is estimated by the YAM scheme (Yamartino, 1993), deriving a vertical velocity component at each grid cell that satisfies the mass continuity equation using the driving meteorology model's air density. The vertical advection module works with no mass-exchange boundary conditions at the bottom or top of the model. In this work the net effect of horizontal and vertical advection is considered to obtain comparable contributions with the rest of the processes. As already pointed out by Jiang et al. (2003), in some cases it could be more meaningful to aggregate the contributions of several processes instead of considering each of them separately.
2. Gas phase chemistry: CBM-IV mechanism is used with the Euler Backward Iterative (Hertel et al., 1993) solver.
3. Diffusion: The diffusion involves sub-grid scale turbulent mixing of pollutants. Horizontal diffusion is estimated by a diffusion coefficient based on local wind deformation. Vertical diffusion is estimated using the Eddy diffusivity theory.
4. Dry deposition: The deposition process is simulated as a flux boundary condition that affects the concentration in the lowest layer.
5. Clouds chemistry and wet deposition: Clouds play a key role in aqueous chemical reactions, vertical mixing of pollutants and removal of pollutants by wet deposition. They also indirectly affect the concentration of pollutants by altering the solar radiation, which, in turn, affects photochemical pollutants, such as ozone, and the flux

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of biogenic emissions. CMAQ models sub-grid convective precipitating clouds, sub-grid non-precipitating clouds, and grid-resolved clouds. The cloud module vertically redistributes pollutants for the sub-grid clouds, calculates in-cloud and precipitation scavenging, performs aqueous chemistry, and accumulates wet deposition amounts. The used scheme is a RADM based cloud processor that uses the asymmetric convective model to compute convective mixing.

6. Aerosols: The third generation aerosol module takes chemical species concentrations and reactivity rates from the chemistry solvers and primary particulate concentrations from the emissions processor to compute fine and coarse particulate concentrations. The primary aerosols emissions are estimated by HERMES, not taking into account marine aerosols. The deposition velocity for particles is estimated from the aerosol size distribution, which is calculated from the mass and number concentration for each of the modes considered: Aitken (0–0.1 μm), accumulation (0.1–2.5 μm), and coarse (2.5–10 μm)

The contribution of processes in this work is analyzed in several ways:

2.2.1 Analysis focused on model evaluation

The contribution of processes to the concentration of gaseous pollutants in the lowest vertical level is assessed for specific points in the domains, coincident with the location of characteristic air quality stations (AQS): rural background AQS and urban background AQS. The contributions are obtained as concentration ($\mu\text{g m}^{-3}$) variation for the surface cell. The weighted contributions are also obtained taking into account the Eq. (1), where: $\%PC_i$ is the relative contribution of the i process to the net fluxes from or to the specific cell or domain (including chemical contributions) and PC_j is the contribution of each process, being j =gas-phase chemistry, transport (horizontal plus vertical advection), emissions, horizontal and vertical diffusion, dry deposition, clouds

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

processes and aerosols processes.

$$\%PC_i = \frac{PC_i}{\sum_j \text{abs}(PC_j)} \cdot 100 \quad (1)$$

Note that the sum of $\%PC_i$ for all i is not 100%, being the sum of $\text{abs}(\%PC_i)$ exactly 100%.

5 2.2.2 Analysis of the dynamics of pollutants during the episode

The processes are analyzed for the selected domains and up to the selected top of the atmosphere (33 sigma vertical layers up to 50 hPa) depicting contribution maps.

2.2.3 Analysis of specific locations with high O₃ levels during the episode

10 Finally, the contribution of the different processes is assessed for four selected subdomains (20×10 km²) in which maximum O₃ concentrations occur during the episode. Positive and negative contributions expressed as pollutants density under the PBL (height obtained from the model) in g m⁻² are assessed for O₃, NO_x and NMVOCs in each subdomain.

2.3 Domains description

15 The coastal domain (NEIP) covers 83 398 km² and is located in the Mediterranean littoral (Fig. 1 – left). It is characterized by a very complex terrain, dominated in the northern part by the Pyrenees mountains (up to 3400 m), and with two mountain chains parallel to the coast: the Pre-coastal (1000–1500 m height) and Coastal chains (500 m average height). The Ebro valley is located in the Southern region and acts as an important channel for local wind flows. The Central Plateau covers the inland area. 20 The major sources of pollutants emission are the urban areas of Barcelona, accounting for 3.1 million inhabitants, and Tarragona, which is located in a densely industrialized

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



area; and the road network connecting the Iberian Peninsula with France, all of them located along the coastal axis. Also inland some industrial areas have an important contribution to pollutants emissions (i.e. the Cercs power plant). The HERMES model estimates 299 t d^{-1} of NO_x emitted during the episode, being 57% produced by on-road traffic, which constitutes the main source of primary pollutants in the region (Costa and Baldasano, 1996; Parra et al., 2006; Baldasano et al., 2008), contributing also by 24% of the 692 t d^{-1} of NMVOCs emitted in the coastal domain. In this particular case 48% comes from biogenic sources. The industrial and power generation sectors are the main emitters of SO_2 and primary particles, accounting for 94% and 71% of the mass emissions (125 t d^{-1} , 45 t d^{-1}), respectively.

The continental (CIP) domain covers an area of $38\,734 \text{ km}^2$ in the centre of the Iberian Peninsula (Fig. 1 – right). The main topographic features in the region are the Central System located in the north-western area of the domain, with summits reaching 2500 m, and the Tajo valley in the southern area. Both clearly affect the wind flows during the episode, characterized by the dominance of mesoscale phenomena. The main emission sources in the region are the Madrid urban area, the Spanish administrative capital, which accounts for 5.8 million inhabitants; the road network that connects Madrid with surrounding conurbations and several industrial sources, mainly located in the south-western part of the domain. The HERMES emissions model indicates that on-road traffic accounts for 75% of total NO_x emissions within the region (231 t d^{-1}), and 19% of NMVOCs (790 t d^{-1} in the whole area), being the biogenic sources the main contributors to these primary pollutants (64%). Industrial and power generation emissions of SO_2 and primary particulate matter account for 90% and 65% of the total emissions (66 t d^{-1} and 36 t d^{-1} , respectively).

2.4 Episode selection

The 17–18 June 2004 episode is characterized by a western recirculation in the synoptic scale, a typical summertime situation in south-western Europe since these condi-

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

tions dominate 45% of the annual and 78% of the summertime transport patterns over the coastal Mediterranean areas (Jorba et al., 2004) and 36% of the annual and 45% of the summertime situations in the central-continental areas of the Iberian Peninsula. They are frequently associated with local-to-regional episodes of air pollution related to high levels of O₃ during summer (e.g. Toll and Baldasano, 2000; Barros et al., 2003; Ortega et al., 2004; Taghavi et al., 2004; Cousin et al., 2005; Coll et al., 2005; Jiménez et al., 2006, among others), being the study case one of the most polluted episodes of 2004 in the considered areas. The weak synoptic forcing and the stagnant conditions dominate the Iberian Peninsula. Therefore the mesoscale phenomena induced by the particular topography of the regions control the superficial wind flows. In these conditions the sea breezes and mountain-valley winds, and the development of the Iberian thermal low are characteristic features of the region (Jorba et al., 2004). The main processes inland are the convective circulations developed by the surface heating and the formation of compensatory subsiding flows in coastal areas (Millán et al., 1997; Pérez et al., 2004).

3 Results and discussion

3.1 Summary of the model evaluation for the 17–18 June 2004 episode

The surface concentrations of pollutants predicted by the WRF-ARW/HERMES/CMAQ model are evaluated against hourly data from 45 AQS (provided by the Environmental Departments of the Catalonia and Madrid Governments, Spain). The evaluation results for O₃ accomplish the recommendations of the European Union, set by the European Directives 1999/30/EC and 2002/3/EC (uncertainty of 50% for the air quality objective for modelling assessment methods), and the US Environmental Protection Agency guidelines (US-EPA, 1991, 2005), which suggest values of ±10–15% for the mean normalized bias error (MNBE), ±15–20% for the unpaired peak prediction accuracy (UPA) and 30–35% for the mean normalized gross error (MNGE) for concentrations above a

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

prescribed threshold to be met by simulated O_3 to be considered for regulatory applications. The averaged MNBE for O_3 is 7.95%; the MNGE, 25.89% and the UPA, -0.39% (Table 2). Moreover, the modelled NO_2 , SO_2 and PM_{10} surface levels agree with observations, with values for these statistical parameters lower or around 50%. The largest deviations are observed in the NO_2 predictions accounting for 50.92% MNGE, nevertheless the reliability of the modelling system is acceptable being MNBE and the UPA -12.68% and -28.35% , respectively; and the average absolute error and the mean bias accounting for $27 \mu\text{g m}^{-3}$ and $-16 \mu\text{g m}^{-3}$, respectively.

The O_3 concentrations are overpredicted by the model both in the NEIP and CIP domains (Table 2), being the average normalized bias 5.85% in the former and 12.98% in the latter. Previous studies (Jiménez et al., 2006) point that under low pressure gradient situations mesoscale models tend to underestimate daytime wind flows, which would favour the O_3 accumulation. The O_3 peaks are overestimated in the CIP (UPA: 11.02%), while an underprediction is assessed for the NEIP domain (UPA: -5.18%). The flow patterns in both domains differ, presenting the NEIP a very complex behaviour, with layering of pollutants and multiday accumulations that may be not accurately reproduced by the model, causing this underprediction in peak concentrations.

NO_2 concentrations are overpredicted in the CIP domain (positive bias of 5.89%), which can be attributed to the weaknesses of the model to represent wind flows under this low pressure gradient situation. The underprediction in the coastal domain (MNBE: -27.12%) could be related with inaccuracies in the modelled chemical behaviour, linked to O_3 chemistry that may be causing also the high levels estimated for this pollutant. The NO_2 peak concentrations are underestimated in both domains (UPA: -26.3% in the CIP domain and -23.73% in the NEIP domain). Jiménez-Guerrero et al. (2008) run the WRF-ARW/HERMES/CMAQ model for the NEIP indicating an underprediction of NO_2 due to the relatively low vertical resolution in the lower troposphere that could generate an artificial vertical exchange between the boundary layer and the free troposphere, enhancing the NO_x venting. In this case, doubling the vertical resolution (33 vertical layers versus 16 in the mentioned study) leads to the same problem.

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

This fact suggests that discrepancies between modelled and measured levels do not only depend on transport patterns, but also on the chemical behaviour represented by the model.

The underestimation in SO₂ concentrations in both domains (MNBE of -32.77% and -41.73% in the NEIP and CIP domains, respectively) and in peak levels (UPA of -29.23% and -47.53%) could reflect a slight underestimation on sulphur oxides emissions.

On average, the PM₁₀ concentrations are also slightly underpredicted (MNGE of -15.86% and UPA -48.42%), which could be related not only to the inaccuracies in representing accumulation and transport patterns during this low gradient pressure situation, but also to the limitations related to the emissions model (HERMES) that does not take into account natural sources of primary particulates such as erosive or saltation processes or marine aerosols (Vautard et al., 2005).

The contribution of processes in the first simulated layer (0–38 m a.g.l.) provides useful information about the performance of the model for predicting surface O₃ concentrations. The surface O₃ concentration in rural background stations of the coastal domain (such as Agullana, Ponts and Rubí) and the continental domain (i.e. SM Valdeiglesias, Guadarrama and Buitrago de Lozoya) is overestimated during the morning and underestimated in the afternoon (the average bias is negative for the costal domain stations, -8.1%, -7.1% and -11.8%, respectively, and -8.01%, 10.27% and 2.21% for the continental domain AQS aforementioned). The representative sites selected reflect a common behaviour (see Fig. 2) independently of their location, which may indicate slight deviations in modelled advection fluxes. The vertical diffusion accounts for 33% of the net processes contribution to the surface cells, but is compensated by 32% removal of dry deposition. The processes controlling the total change in these cells' concentration are the net transport term (horizontal + vertical advection, accounting on average for 8%) and chemical production (7%). If the vertical transport (diffusion + dry deposition) is considered as a whole, the transport weight in the change of concentration involves 48% and chemistry 37% of positive contribution. The largest de-

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

5 viations in model predictions on the coastal domain stations occur specifically in the 16:00 to 20:00 UTC period, when the daytime breezes decrease and the land breezes and night-time winds start developing. The surface winds during this period are over-estimated by the meteorological model which in turn favours pollutants venting in the chemistry transport model simulation. In the CIP concentrations are underestimated from 10:00 UTC, but maximum deviations are found in the 19:00–22:00 UTC period.

10 Previous works confirm the trend of the mesoscale meteorological models, and specifically the WRF-ARW, to underpredict the land breezes and underestimate the advective flows in the latter hours of the day (Jiménez-Guerrero et al., 2008), which causes overprediction in O_3 concentrations during night-time. The average processes contributions to net O_3 concentrations assessed during this period indicates that net transport accounts for 56% and chemical destruction for –26%, when considering the combined effect of vertical processes (vertical diffusion + dry deposition).

15 In the locations of urban background stations (such as Barcelona-Ciutadella and Tarragona in the NEIP, or Aranjuez and Chapineria in the CIP) the modelled O_3 has a similar behaviour (Fig. 3). Albeit during the morning there is an overestimation trend, during the afternoon and night-time the model predictions are frequently below the measured levels. The chemical regime in the NEIP domain involves O_3 photochemical destruction in the first layer accounting for –46% of net night-time fluxes and –44% of net daytime fluxes on average, when considering the vertical fluxes as a whole (being the sum of vertical diffusion and dry deposition a positive term that accounts for 10% and 13%, respectively). In the CIP the periods of O_3 net formation occur in the first vertical layer during daytime, not existing in the urban background stations of the NEIP. On average the net effect of daytime gas phase chemistry accounts for
20 –5% of total fluxes of O_3 . The main positive contributions come from net transport, involving 86% of net concentration variation in the surface cells. During night-time main sinks of O_3 are chemical destruction (–27%) and the vertical flux term (vertical diffusion plus dry deposition, that combined remove –12% of net O_3 fluxes). In the coastal domain stations the chemical destruction of O_3 is the main removal process
25

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

even during daytime, probably due to the O_3 titration by NO_x emissions. On-road traffic plays an important role in this sense being the major NO_x emitter the Barcelona city area.

This behaviour is also reproduced by the model in urban stations such as Barcelona-Eixample, Constantí or L'Hospitalet (NEIP) and Majadahonda, Coslada or Leganés (CIP) where main contributions to surface O_3 come from transport and vertical diffusion, and the main sinks are gas-phase chemistry and dry deposition.

3.2 Dynamics of pollutants in the coastal and central south-western Mediterranean

The results of the photochemical simulations for the coastal and central-continental domains show the maximum O_3 concentrations occurring in downwind areas from Barcelona (NEIP) and Madrid (CIP) conurbations after the maximum photochemical activity hours (Fig. 4). During the day the increase of the solar radiation and the temperature (reaching 30–35°C in both domains) promote the high levels of O_3 , exceeding in some cases the population information threshold ($180 \mu\text{g m}^{-3}$). The gas-phase chemistry accounts in these cases for 50 to $75 \mu\text{g m}^{-3} \text{h}^{-1}$ in the first layer cells. In the urban areas and main roads, as main sources of NO_x , chemical destruction of O_3 occurs up to $-200 \mu\text{g m}^{-3}$ per hour during the traffic circulation peaks. The O_3 concentration gradient generated in these areas, larger during daytime, increases the contribution of diffusion processes to ground-level O_3 (up to $200 \mu\text{g m}^{-3} \text{h}^{-1}$ fluxes, mainly from upper vertical layers). Surface O_3 concentrations are larger in the continental than in the coastal domain. While in the NEIP the breezes and mountain-valley winds regime involves the accumulation and recirculation of pollutants aloft, in the CIP the main flows are dominated by thermal phenomena that transport pollutants within the convective recirculation cell. The PBL height reaches its maximum at midday, being higher in the central-continental domain – around 3200 m a.g.l. – than in the coastal domain – maximum around 2000 m a.g.l. –, due to the lamination of the PBL growth by the Mediterranean Sea breezes.

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Contribution of
processes affecting
atmospheric
dynamics**M. Gonçalves et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The emitted NO_x and NMVOCs in the central-continental domain during the morning react forming O_3 , the surface heating generates a convective cell and the photochemical pollution plume rises, reaching 2500–3000 m a.g.l. (Figs. 5–6) and moves downwind, affecting the southern area during the 17 June and the eastern area on the 18 June. The maximum surface O_3 concentrations occur at midday (from 13:00 to 16:00 UTC of 17 June and 12:00 to 14:00 UTC of 18 June). During the afternoon and dusk the convective cell weakens and falls over the Madrid downwind areas, the PBL height reduction involves high O_3 surface concentrations even when the photochemical production diminishes. At night-time, the O_3 generation ceases, and the concentration of photochemical pollutants decays. The katabatic winds dominant in the Central system area enhance the effect of the wind shift, being the main flows channelled by the Tajo valley towards the south-western area (hourly maximum concentrations in this period are located in this area, see Fig. 4).

The coastal domain behaviour differs, being more favourable to several days photochemical episodes, due to the reservoir layers formed over the Mediterranean Sea during night-time (Figs. 5–6). Specifically, the main emission sources are located in the coastal area. When the mesoscale phenomena dominate the pollutants transport, the breeze regime development in during the morning transports the primary pollutants inland. The littoral mountain chain acts as a barrier at dawn, recirculating the pollutants towards the Mediterranean Sea; these return flows are enhanced by the development of anabatic winds. As the day advances the sea breeze regime develops and reaches the Pre-littoral chain, higher than the previous (1000–1500 m), producing a second recirculation flow aloft. On the other hand, the river valleys and main roads act channelling the pollutants flow inland. The pollution plume reaches flat inland areas, where it accumulates during the afternoon and dusk, because of the Pyrenees barrier cutting the flows to the northern area. In fact maximum O_3 surface concentrations during the episode are reached in the 12:00–14:00 UTC period in north-eastern areas from the Barcelona conurbation and maximum hourly concentrations in the 18:00–21:00 UTC period are found in the northern domain (in the Pre-Pyrenees region) (Fig. 4). When

the mesoscale wind regime shifts, katabatic winds and land breezes involve a return flow of the pollutants plume over the Mediterranean Sea. Moreover, previous studies (e.g. Jiménez et al., 2006) have detected a layer of pollutants outbreak in altitude with peninsular origin (3000–3500 m a.g.l.) that contributes in this kind of episodes to the high concentration of pollutants in the area.

On the other hand, the urban plume reactivity differs, being the Madrid airshed more favourable to O₃ formation in situ, while in Barcelona O₃ titration by NO_x emissions involves a sink of O₃ in the surface layer. Formation occurs mainly in upper levels and in areas located around 100 km away from the city (Fig. 5).

In summary, the main contributors to O₃ surface concentrations are the net transport and vertical diffusion, while the main sink is dry deposition. The wet deposition and cloud processes are negligible during the episode, due to the low cloudiness and the absence of precipitation, typical in a summertime period such as the one studied. The aerosols interaction with photochemical gaseous pollutants is also negligible. The chemical contribution differs depending on the location and the NO_x-VOCs ratio. The process analysis indicates that maximum chemical production of O₃ does not occur in the first vertical layer of the model, but in layers aloft during the central hours of the day (Fig. 5). This effect will be deeply discussed in the next section. The horizontal and vertical advective transport determine the areas that will be affected by these pollutants, being the transport patterns much simpler in the central domain than in the coastal area, where stratified layers and accumulation of pollutants are frequent during summertime episodes with stagnant conditions, as shown in Fig. 6. The process analysis confirms the results of previous experimental and modelling studies carried out in these areas (e.g. Baldasano et al., 1994; Millán et al., 1997; Soriano et al., 2001; Pérez et al., 2004; Jiménez et al., 2006).

3.3 Processes contribution to O₃-NO_x-NMVOCs concentrations

In order to define the origins of maximum O₃ concentrations in the coastal and continental domains, two subdomains (20×10 km²) – MAX1 and MAX2 – are defined con-

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sidering the areas where the maximum concentrations are estimated by the model on the first and second day of the episode, respectively (Fig. 7). The O_3 , NO , NO_2 and NMVOCs density ($g\ m^{-2}$) in the atmospheric column under the PBL is assessed together with the contributions of different processes leading to these levels. All values are obtained as an hourly average for the whole subdomain areas ($200\ km^2$). The night-time modelled PBL height (Fig. 7) is near surface level (around $20\ m\ a.g.l.$), then the pollutants density and processes contribution are negligible versus those of daytime periods.

The O_3 maximum concentrations during the 17 June occur in the MAX1 domains. In the coastal-MAX1 the horizontal advective transport to the area constitutes the major positive contribution to net O_3 until 16:00 UTC ($0.042\ g\ m^{-2}\ h^{-1}$ on average from 09:00 to 16:00 UTC). There is also a significant vertical advective transport removing O_3 from the PBL ($-0.052\ g\ m^{-2}\ h^{-1}$). Later on, until 19:00 UTC, the horizontal advective flows remove O_3 from this area ($0.057\ g\ m^{-2}\ h^{-1}$ on average), due to the transport of the pollutants plume towards the Mediterranean Sea. The O_3 chemical production from 07:00 UTC to 17:00 UTC is the second source of importance ($0.019\ g\ m^{-2}\ h^{-1}$). The continental-MAX1 domain presents a similar behaviour, being the horizontal advective transport the most important positive contribution to net O_3 until 15:00 UTC, on average $0.042\ g\ m^{-2}\ h^{-1}$ (Fig. 8). Then, the wind shift involves a negative contribution of horizontal advection until 19:00 UTC ($-0.079\ g\ m^{-2}\ h^{-1}$). In this case there are no important vertical injections, being the contributions of vertical advection almost negligible compared to horizontal advection and chemical production (i.e. vertical transport causes a removal of $-0.006\ g\ m^{-2}\ h^{-1}$ on average in the 11:00 to 15:00 UTC period). The complex topography in the coastal area explains, as already commented, the vertical transport flows of pollutants caused by the breeze regimes mechanically enhanced by the mountain chains located along the coast. Meanwhile, in the continental domain the vertical flows promote the transport of pollutants under the PBL; these are thermally produced and they have no orographic forcing involving injections in upper levels.

On the other hand, the MAX2 subdomains stand for the locations of maximum O_3

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Contribution of
processes affecting
atmospheric
dynamics**

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

concentrations during the second day of the episode. As discussed for the first day, the contributions to net O_3 density are mainly transport; the continuous horizontal advective flows involve high O_3 densities in the coastal-MAX2 subdomain (horizontal transport contributes on average with $0.064 \text{ g m}^{-2} \text{ h}^{-1}$ during the 08:00 to 19:00 UTC period). Once again, the vertical injections are the major sink of O_3 in the region ($-0.062 \text{ g m}^{-2} \text{ h}^{-1}$ on average). Chemical production in this case is lower than in areas nearer from the Barcelona urban area, such as MAX1 domain ($0.009 \text{ g m}^{-2} \text{ h}^{-1}$ on average), and the O_3 concentrations estimated by the model are clearly affected by horizontal advection. The continental-MAX2 domain is characterized by an important horizontal advection during the central hours of the day, from 12:00 to 15:00 UTC contributing with $0.062 \text{ g m}^{-2} \text{ h}^{-1}$ on average; then the horizontal flows remove O_3 from the domain ($-0.078 \text{ g m}^{-2} \text{ h}^{-1}$ until 18:00 UTC). The relatively high O_3 density observed from 14:00 to 16:00 UTC is due to the chemical production ($0.015 \text{ g m}^{-2} \text{ h}^{-1}$ from 13:00 to 18:00 UTC) and the larger integration height (PBL height 3134 to 3220 m a.g.l. as observed in Fig. 7) considered.

The NEIP subdomains (coastal) are characterized by the low weight of emissions in the NO_x budget under the PBL (Fig. 9) when compared to CIP (continental). Specifically in the NEIP-MAX1, the horizontal advection is the main source of NO - NO_2 until 16:00 UTC (3.4×10^{-4} and $0.0022 \text{ g m}^{-2} \text{ h}^{-1}$, respectively) and chemical reaction involves the main sink of these oxides (-4.4×10^{-4} and $-0.0016 \text{ g m}^{-2} \text{ h}^{-1}$, respectively). From then on the horizontal transport removes NO_2 and vertical transport from layers above the PBL contributes positively until 18:00 UTC. While NO continues oxidizing and chemical destruction is still the main sink of this compound, from 18:00 to 20:00 UTC net chemical production of NO_2 in situ is observed ($2.07 \times 10^{-4} \text{ g m}^{-2} \text{ h}^{-1}$). The minimum NO_x density observed on 17 June around 15:00–16:00 UTC reflects the direct effect of the peaks of traffic emissions during the early morning and the afternoon in closer areas (such as Barcelona city), which is the main origin of the advected NO_x .

The CIP subdomains present a similar behaviour concerning NO_x (Fig. 9), being the major source the horizontal advection from surrounding areas ($2.0 \times 10^{-4} \text{ g m}^{-2} \text{ h}^{-1}$ for

**Contribution of
processes affecting
atmospheric
dynamics**

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

NO and $0.0011 \text{ g m}^{-2} \text{ h}^{-1}$ for NO_2 until 19:00 UTC) and the main sinks the gas-phase chemical reactions ($-5.810^{-4} \text{ g m}^{-2} \text{ h}^{-1}$ and $-4.2 \times 10^{-4} \text{ g m}^{-2} \text{ h}^{-1}$, respectively). Nevertheless the emissions of NO have limited importance in these subdomains, and specifically in the CIP-MAX1 domain, where NO emissions contribute with an approximately constant density, from 04:00 UTC to 19:00 UTC ($4.9 \times 10^{-4} \text{ g m}^{-2} \text{ h}^{-1}$). The increase in NO_2 density during the 15:00–18:00 UTC period due to the advective transport and the lower chemical destruction may be related to the high O_3 density observed. The NO_x levels in the central-MAX2 domain are low during the 17 June and increase considerably by horizontal advection during the 18 June. In the early morning (until 10:00 UTC) losses of NO_x by vertical diffusion ($-2.8 \times 10^{-5} \text{ g m}^{-2} \text{ h}^{-1}$ for NO and $-1.2 \times 10^{-4} \text{ g m}^{-2} \text{ h}^{-1}$ for NO_2) to upper layers are estimated by the model.

In the coastal-MAX1 domain the NMVOCs levels are relatively high (Fig. 10). Horizontal advection until 17:00 UTC (contributing on average with $7.5 \times 10^{-5} \text{ g m}^{-2} \text{ h}^{-1}$) and vertical advection from upper layers until 19:00 UTC ($3.6 \times 10^{-5} \text{ g m}^{-2} \text{ h}^{-1}$) are their main sources. The daytime average contribution of emissions is low and accounts for $1.2 \times 10^{-5} \text{ g m}^{-2} \text{ h}^{-1}$. During the morning, until 10:00 UTC, some NMVOCs are injected in layers above the PBL by vertical diffusion ($-1.5 \times 10^{-5} \text{ g m}^{-2} \text{ h}^{-1}$) and vertical advective processes ($-2.5 \times 10^{-5} \text{ g m}^{-2} \text{ h}^{-1}$ on average until 16:00 UTC), involving a sink of these organic compounds, together with chemical reaction ($-1.3 \times 10^{-5} \text{ g m}^{-2} \text{ h}^{-1}$ on average during daytime). Nevertheless, the net transport (advection and diffusion) involves a positive contribution during daytime on average of $3.8 \times 10^{-5} \text{ g m}^{-2} \text{ h}^{-1}$. From 11:00 UTC the vertical diffusion contributes positively to the NMVOCs density under the PBL, nevertheless its contribution is almost negligible compared to the net transport by advection. The NEIP-MAX2 behaviour is more homogeneous, being the horizontal advective flows the main contributors to net NMVOCs density under the PBL ($5.6 \times 10^{-5} \text{ g m}^{-2} \text{ h}^{-1}$) and vertical advection the main sink ($-3.8 \times 10^{-5} \text{ g m}^{-2} \text{ h}^{-1}$), together with chemical reaction ($-6.5 \times 10^{-6} \text{ g m}^{-2} \text{ h}^{-1}$) and vertical diffusion specially during the morning ($-1.3 \times 10^{-5} \text{ g m}^{-2} \text{ h}^{-1}$ until 11:00 UTC).

The horizontal advection affecting the CIP-MAX1 domain during 17 June involves the

largest contribution to net NMVOCs density under the PBL until 16:00 UTC (Fig. 10, average contribution of $1.27 \times 10^{-4} \text{ g m}^{-2} \text{ h}^{-1}$), then the winds shift removes these compounds by the same process. The chemical destruction reaches its maximum during the 14:00–15:00 UTC, accounting for $-2.0 \times 10^{-5} \text{ g m}^{-2} \text{ h}^{-1}$. Vertical advection and diffusion remove NMVOCs from the PBL, but in a lower extent. The specific study area is not characterized by a high emissions rate; therefore the contribution of this process is almost negligible compared to transport processes. The CIP-MAX2 presents a similar behaviour during the 18 June, when it is directly affected by the horizontal advective flows. The winds shift occurs one hour earlier, being the horizontal transport the main sink of NMVOCs in the area from 16:00 UTC ($-6.5 \times 10^{-5} \text{ g m}^{-2} \text{ h}^{-1}$). The maximum removal by chemical destruction also occurs in the 14:00–15:00 UTC period ($-1.7 \times 10^{-5} \text{ g m}^{-2} \text{ h}^{-1}$).

Differences in chemical sensitivity are observed for the studied domains (Fig. 11). The coastal domains present a similar chemical behaviour; the NMVOCs are the only locally emitted precursors, which react with transported NO_x generating O_3 by gas-phase chemistry. The CIP-MAX1 domain includes emitters of NO_x and NMVOCs, which react forming O_3 , nevertheless the O_3 chemistry does not involve destruction of NMVOCs in the same magnitude as the coastal domains. This is due to the more NO_x sensitive regime of this area. In absence of solar radiation, when the O_3 stopped forming, the NO oxidation continues producing net NO_2 (18:00–19:00 UTC period). In the CIP-MAX2 domain the NMVOCs emissions are an important source, which from 10:00 to 19:00 UTC on 17 June are chemically destroyed, being the generation of O_3 by chemistry relatively low. It could be explained by the absence of locally emitted NO_x and the slight contribution of net transport of these oxides in this specific day to the area. On the second day of the episode the horizontal advection directly affect the MAX2 involving a much higher net chemical production of O_3 , being the chemical oxidation of NMVOCs lower.

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

4 Conclusions

The Integrated Process Rate implemented in the CMAQ model was applied to obtain quantitative information about atmospheric processes affecting the concentration of pollutants in typical coastal and continental environments located in south-western Europe: the North-eastern Iberian Peninsula (NEIP) and Central Iberian Peninsula (CIP) domains, respectively. A representative summertime photochemical pollution episode characterized by stagnant conditions over the area was selected.

The model performance agrees with US-EPA and European Directives recommendations, nevertheless specifically in background air quality stations tends to overestimate the O_3 morning concentrations and underestimate the O_3 levels during the afternoon. The chemical destruction may be underpredicted during night-time, which favours the high estimated morning concentrations, while the overpredicted flows during the last hours of the day and night-time could cause an enhanced venting of pollutants. The mesoscale meteorological models have shown inaccuracies in predicting wind flows during the dominant stagnant conditions; they play a fundamental role in these deviations. NO_2 is clearly linked to O_3 , with both inaccuracies in transport and chemical behaviour being the main cause of the errors in the simulated concentrations. On the other hand, O_3 peaks and NO_2 underprediction in the coastal domain may be related to difficulties for the model to define the accumulation layers formed and characteristic recirculation of pollutants due to the very complex terrain. In the case of SO_2 and PM_{10} a slight underestimation of emissions could be the main cause for the underestimation of these pollutants.

Applying the Integrated Process Rate tool to the first vertical layer simulated provides information about the surface concentration of pollutants estimated by the model and permits to test the mass consistency. In order to perform a deeper study of the contributions of main atmospheric processes leading to the levels of these pollutants the vertical component has to be considered. In this work the whole atmospheric column under the PBL is selected, reflecting that chemistry and transport patterns vary with

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

height.

The process analysis indicates that the maximum concentration of photochemical pollutants occur due to transport phenomena. Specifically in the coastal domain (NEIP), the high surface O_3 levels are not produced in situ, but come from horizontally advected flows during the morning and gas-phase chemical contributions occurring aloft. During the afternoon and dusk vertical advective flows inject the pollutants in layers above the PBL, which accumulate due to the stagnant conditions over the region and contribute positively to surface concentrations the next simulated day. The central-continental domain (CIP) behaviour slightly differs, with horizontal advection being also the main contributor to O_3 surface concentrations, but having the chemical formation some importance at low levels. The transport patterns differ between the coastal and the continental area, where the O_3 precursors are homogeneously distributed in the whole atmospheric column under the PBL and vertical injections of pollutants above the PBL are less frequent.

The dry deposition is an important sink for pollutants in the lowest layer of the model, coupled in the present simulation with vertical diffusion flows. In spite of the stagnant conditions dominating during the episode, the diffusive processes contributions to net pollutants concentrations under the PBL are relatively low, and in particular the horizontal diffusion is negligible compared to other atmospheric processes. Vertical diffusion compensates the loss of O_3 in surface layers due to NO titration, contributing positively to net O_3 concentrations in the road network and urban areas. The O_3 peaks at surface level are higher in the CIP domain, mainly due to the much simpler transport pattern compared to the coastal region, which involves that almost all the chemically produced O_3 under the PBL contributes to surface concentrations when the convective cell weakens. Controlling emissions of precursors in both domains is a decisive factor in order to abate photochemical pollution episodes during summertime. The emitted NMVOCs have mainly a biogenic origin and therefore policies focused on reducing NO_x emissions must be addressed in these regions. Finally, the cloud processes, wet deposition and heterogeneous chemistry contributions are negligible during the whole

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

episode, characterized by a high solar radiation and no precipitation or cloudiness.

This work explores the possibilities of applying process analysis to high resolution simulations proving that it could be useful not only to better evaluate the simulation results, but also to perform more accurate source apportionment of pollutants over a region.

Acknowledgements. The authors gratefully acknowledge E. López for the implementation of HERMES, O. Jorba for the meteorological input data and T. Loricán for his help in generating 3-D images. This work was funded by the projects CICYT CGL2006-08903 and CGL2006-11879 of the Spanish Ministry of Education and Science and CALIOPE project 441/2006/3-12.1 of the Spanish Ministry of the Environment. Simulations were carried out in the MareNostrum supercomputer: 94.21 Teraflops of theoretical performance peak; 10240 processors PowerPC 970MP (2560 JS21 blades); Fig. 7 was obtained from Google Maps, © 2007 – Images Digital-Globe, Terrametrics, NASA.

References

- Arévalo G., Salvador, R., Gassó, S., Millan, M., and Baldasano, J. M.: Application of a high-resolution emission model in Valencia Community (Spain), in: Air Pollution 2004, Ed. WITpress, Rhodes, Greece, 31–40, 2004.
- Baldasano, J. M., Cremades, L., and Soriano, C.: Circulation of air pollutants over the Barcelona geographical area in summer, in: Proceedings of Sixth European Symposium Physico-Chemical Behaviour of Atmospheric Pollutants, Varese, Italy, 18–22 October 1993, Report EUR 15609/1 EN, 474-479, 1994.
- Baldasano, J. M., Güereca, P., López, E., Gassó, S., and Jiménez-Guerrero, P.: Development of a high resolution (1 km×1 km, 1 h) emission model for Spain: the High-Selective Resolution Modelling Emission System (HERMES), Atmos. Environ, 42, 7215–7233, 2008.
- Barros, N., Borrego, C., Toll, I., Soriano, C., Jiménez, P., and Baldasano, J. M.: Urban photochemical pollution in the Iberian Peninsula: Lisbon and Barcelona airsheds, J. Air Waste Manage., 53, 347–359, 2003.
- Byun, D. W. and Schere, K. L.: Review of the governing equations, computational algorithms

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- and other components of the Models-3 Community Multiscale Air Quality (CMAQ) Modeling System, *Appl. Mech. Rev.*, 59(2), 51–77, 2006.
- Coll, I., Pinceloup, S., Perros, P. E., Laverdet, G., and Le Bras, G.: 3D analysis of high ozone production rates observed during the ESCOMPTE campaign, *Atmos. Res.*, 74, 477–505, 2005.
- Cousin, F., Tulet, P., and Rosset, R.: Interaction between local and regional pollution during ESCOMPTE 2001: impact on surface ozone concentrations (IOP2a and 2b), *Atmos. Res.*, 74, 117–137, 2005.
- Cros, B., Durand, P., Cachier, H., Drobinski, P., Fréjafon, E., Kottmeier, C., Perros, P. E., Peuch, V.-H., Ponche, J.-L., Robin, D., Saïd, F., Toupance, G., and Wortham, H.: The ESCOMPTE program: an overview, *Atmos. Res.*, 69, 241–279, 2004.
- Dufour, A., Amodei, M., Ancellet, G., and Peuch, V.-H.: Observed and modelled chemical weather during ESCOMPTE, *Atmos. Res.*, 74, 161–189, 2005.
- Gipson, L. G.: Science Algorithms of the EPA Models-3 Community Multiscale Air Quality (CMAQ) Modeling System: process analysis, EPA/600/R-99/030, US-EPA, <http://www.epa.gov/asmdnerl/CMAQ/CMAQscienceDoc.html>, 37 pp., 1999.
- Gery, M. W., Whitten, G. Z., Killus, J. P., and Dodge, M. C.: A photochemical kinetics mechanism for urban and regional scale computer modeling, *J. Geophys. Res.*, 94, 12, 925–956, 1989.
- Hauglustaine, D. A. and Brasseur, G. P.: Evolution of tropospheric ozone under anthropogenic activities and associated radiative forcing on climate, *J. Geophys. Res.*, 106, 32 337–32 360, 2001.
- Hertel O., Berkowicz, R., Christensen, J., and Hov, O.: Test of two numerical schemes for use in atmospheric transport-chemistry models, *Atmos. Environ.*, 27A, 2591–2611, 1993.
- Hogrefe, C., Lynn, B., Rosenzweig, C., Goldberg, R., Civerolo, K., Ku, J.-Y., Rosenthal, J., Knowlton, K., and Kinney, P. L.: Utilizing CMAQ process analysis to understand the impacts of climate change on ozone and particulate matter, in: 4th Annual CMAS Models-3 Users' Conference, 26–28 September 2005, Chapel Hill, NC, USA, 2005.
- Huang, J. P., Fung, J., Zhang, Y., Lau, A., and Qin, Y.: Process analysis of different synoptic patterns of O₃ episodes in Hong Kong, in: 86th Annual Meeting of AMS, 28 January–3 February, 2006, Atlanta GA, 2006.
- Jang, J. C., Jeffries, H. E., Byun, D., and Pleim, J. E.: Sensitivity of ozone to model grid resolution – I: Application of high-resolution regional acid deposition model, *Atmos. Environ.*,

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

21, 3085–3100, 1995a.

Jang, J. C., Jeffries, H. E., and Tonnesen, S.: Sensitivity of ozone to model grid resolution – II: Detailed process analysis for ozone chemistry, *Atmos. Environ.*, 29, 3101–3114, 1995b.

Jeffries, H. E. and Tonnesen, S.: A comparison of two photochemical reaction mechanisms using mass balance and process analysis, *Atmos. Environ.*, 28, 2991–3003, 1994.

Jiang, G., Lamb, B., and Westberg, H.: Using back trajectories and process analysis to investigate photochemical ozone production in the Puget Sound region, *Atmos. Environ.*, 37, 1489–1502, 2003.

Jiménez, P., Baldasano, J. M., and Dabdub, D.: Comparison of photochemical mechanisms for air quality modelling, *Atmos. Environ.*, 37, 4179–4194, 2003.

Jiménez, P., Jorba, O., Parra, R., and Baldasano, J. M.: Evaluation of MM5-EMICAT2000-CMAQ performance and sensitivity in complex terrain: high-resolution application to the northeastern Iberian Peninsula, *Atmos. Environ.*, 40, 5056–5072, 2005.

Jiménez, P., Lelieveld, J., and Baldasano, J. M.: Multiscale modelling of air pollutants dynamics in the northwestern Mediterranean basin during a typical summertime episode, *J. Geophys. Res.*, 111, D18306, doi:10.1029/2005JD006516, 2006.

Jiménez-Guerrero, P., Jorba, O., Baldasano, J. M., and Gassó, S.: The use of a modeling system as a tool for air quality management: annual high-resolution simulations and evaluation, *Sci. Total Environ.*, 390, 323–340, 2008.

Jonson, J. E., Simpson, D., Fagerli, H., and Solberg, S.: Can we explain the trends in European ozone levels?, *Atmos. Chem. Phys.*, 6, 51–66, 2006, <http://www.atmos-chem-phys.net/6/51/2006/>.

Jorba, O., Pérez, C., Rocadenbosch, F., and Baldasano, J. M.: Cluster Analysis of 4-Day Back Trajectories Arriving in the Barcelona Area (Spain) from 1997 to 2002, *J. Appl. Meteorol.*, 43(6), 887–901, 2004.

Lawrence, M. G., Crutzen, P. J., Rasch, P. J., Eaton, B. E., and Mahowald, N. M.: A model for studies of tropospheric photochemistry: description, global distributions and evaluation, *J. Geophys. Res.*, 104, 26 245–26 277, 1999.

Lawrence, M. G., Rasch, P. J., von Kuhlmann, R., Williams, J., Fischer, H., de Reus, M., Lelieveld, J., Crutzen, P. J., Schultz, M., Stier, P., Huntrieser, H., Heland, J., Stohl, A., Forster, C., Elbern, H., Jakobs, H., and Dickerson, R. R.: Global chemical weather forecasts for field campaign planning: predictions and observations of large-scale features during MINOS, CONTRACE, and INDOEX, *Atmos. Chem. Phys.*, 3, 267–289, 2003,

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

<http://www.atmos-chem-phys.net/3/267/2003/>.

Lelieveld, J., Berresheim, H., Borrmann, S., et al.: Global air pollution crossroads over the Mediterranean, *Science*, 298, 794–799, 2002.

Michalakes, J., Dudhia, J., Gill, D., Henderson, T., Klemp, J., Skamarock, W., and Wang, W.: The Weather Research and Forecasting Model: software architecture and performance, Proceedings of the Eleventh ECMWF Workshop on the Use of High Performance Computing in Meteorology, edited by: Zwiefhofer, W. and Mozdzynski, G., World Scientific, European Centre for Medium-Range Weather Forecasts, UK, 156–168, 2005.

Millán, M., Salvador, R., Mantilla, E., and Artíñano, B.: Meteorology and photochemical air pollution in southern Europe: experimental results from EC research projects, *Atmos. Environ.*, 30, 1909–1924, 1996.

Millán, M., Salvador, R., Mantilla, E., and Kallos, G.: Photo-oxidant dynamics in the Mediterranean basin in summer: results from European research projects, *J. Geophys. Res.*, 102, 8811–8823, 1997.

O’Neil, S. and Lamb, B.: Intercomparison of the Community Multiscale Air Quality Model and CALGRID using Process Analysis, *Environ. Sci. Technol.*, 39, 5742–5753, 2005.

Ortega, S., Soler, M. R., Beneito, J., and Pino, D.: Evaluation of two ozone air quality modelling systems, *Atmos. Chem. Phys.*, 4, 1389–1398, 2004, <http://www.atmos-chem-phys.net/4/1389/2004/>.

Parra, R., Gassó, S., and Baldasano, J. M.: Estimating the biogenic emissions of non-methane volatile organic compounds from the North western Mediterranean vegetation of Catalonia, Spain, *Sci. Total Environ.*, 329, 241–259, 2004.

Parra, R., Jiménez, P., and Baldasano, J. M.: Development of the high spatial resolution EMI-CAT2000 emission model for air pollutants from the north-eastern Iberian Peninsula (Catalonia, Spain), *Environ. Pollut.*, 140, 200–219, 2006.

Pérez, C., Sicard, M., Jorba, O., Comerón, A., and Baldasano, J. M.: Summertime recirculations of air pollutants over the north-eastern Iberian coast observed from systematic EARLINET lidar measurements in Barcelona, *Atmos. Environ.*, 38, 3983–4000, 2004.

Roelofs, G. J., Scheeren, H. A., Heland, J., Ziereis, H., and Lelieveld, J.: A model study of ozone in the eastern Mediterranean free troposphere during MINOS (August 2001), *Atmos. Chem. Phys.*, 3, 1199–1210, 2003, <http://www.atmos-chem-phys.net/3/1199/2003/>.

San José, R., Pérez, J. L., Pleguezuelos, C., Camacho, F., and González, R. M.: MM5-CMAQ air quality modeling process analysis: Madrid case, *Air pollution X: ecology and the environ-*

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ment, 53, C. A. BREBBIA, Wessex Institute of Technology, UK, and University of the West of England, Bristol, UK, 2002.

Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Wang, W., and Powers, J. G.: A Description of the Advanced Research WRF Version 2, NCAR Technical note NCAR/TN-468+STR, NCAR, 2005.

Sillman, S. and He, D.: Some theoretical results concerning O_3 - NO_x -VOC chemistry and NO_x -VOC indicators, *J. Geophys. Res.*, 107(D22), 4659, doi:10.1029/2001JD001123, 2002.

Soriano, C., Baldasano, J. M., Buttler, W. T., and Moore, K.: Circulatory Patterns of air pollutants within the Barcelona air basin in a summertime situation: lidar and numerical approaches, *Bound.-Lay. Meteorol.*, 98(1), 33–55, 2001.

Taghavi, M., Cautenet, S., and Foret, G.: Simulation of ozone production in a complex circulation region using nested grids, *Atmos. Chem. Phys.*, 4, 825–838, 2004, <http://www.atmos-chem-phys.net/4/825/2004/>.

Toll, I. and Baldasano, J. M.: Modeling of photochemical air pollution in the Barcelona area with highly disaggregated anthropogenic and biogenic emissions, *Atmos. Environ.*, 34, 3060–3084, 2000.

US EPA: Guideline for regulatory application of the urban airshed model, Office of Air and Radiation, Office of Air Quality Planning and Standards, Technical Support Division, Research Triangle Park, North Carolina, USA, US EPA Report No. EPA-450/4-91-013, 1991.

US EPA: Guidance on the use of models and other analyses in attainment demonstrations for the 8-hour ozone NAAQS, Office of Air Quality Planning and Standards, Research Triangle Park, North Carolina, USA, US EPA Report No. EPA-454/R-05-002, 128 pp., 2005.

Vautard, R., Bessagnet, B., Chin, M., and Menut, L.: On the contribution of natural Aeolian sources to particulate matter concentrations in Europe: Testing hypotheses with a modelling approach, *Atmos. Environ.*, 39, 3291–3303, 2005.

Yamartino, R. J.: Nonnegative, conserved scalar transport using grid-cell-centered spectrally constrained Blackman cubics for applications on a variable-thickness mesh, *Mon. Weather Rev.*, 121, 753–763, 1993.

Zhang, Y., Vijayaraghavan, K., Huang, J., and Jacobson, M. Z.: Probing into Regional O_3 and PM pollution: a 1-year CMAQ simulation and process analysis over the United States, in: 14th Joint Conference on the Applications of Air Pollution Meteorology with the Air and Waste Management Association, 28 January–2 February 2006, Atlanta, GA, USA, 2006.

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Table 1. Summary of the WRF-ARW meteorological model configuration.

Domains definition	Physical parameterizations
Four one way nested domains	ARW dynamical core
D1-European domain 55×55 cells of 54 km	Yonsei University PBL scheme
D2-Iberian Peninsula domain 94×82 cells of 18 km	Kain-Fritsch (D1–D2) and explicit
D3-Iberian Peninsula Zone: 104×103 cells of 6 km	(D3–D4) cumulus scheme
D4a-Northeastern IP domain 322×259 cells of 1 km	Single Moment 3 class
	microphysics' scheme
D4b-Central IP domain 181×214 cells of 1 km	RRTM for long-wave radiation scheme
	and
(Schematic representation in Fig. 1)	Dudhia for short-wave scheme
	Noah Land Surface Model.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. Summary of validation results for the 17–18 June 2004.

45 air quality stations (AQS) data	Bias ($\mu\text{g m}^{-3}$)	Error ($\mu\text{g m}^{-3}$)	MNBE (%)	MNGE (%)	UPA (%)
O ₃	4.9	25.6	7.95%	25.89%	-0.39%
NO ₂	-16.0	27.0	-12.68%	50.92%	-28.35%
SO ₂	-8.9	11.0	-35.76%	47.95%	-35.33%
PM ₁₀	-12.1	16.9	-15.86%	38.59%	-48.42%
Coastal Domain (NEIP) (33 AQS)	Bias ($\mu\text{g m}^{-3}$)	Error ($\mu\text{g m}^{-3}$)	MNBE (%)	MNGE (%)	UPA (%)
O ₃	2.3	25.7	5.85%	26.65%	-5.18%
NO ₂	-26.7	34.5	-27.12%	51.82%	-31.87%
SO ₂	-10.0	12.9	-32.77%	49.95%	-29.23%
PM ₁₀	-3.3	16.1	15.19%	50.65%	-14.82%
Central Domain (CIP) (12 AQS)	Bias ($\mu\text{g m}^{-3}$)	Error ($\mu\text{g m}^{-3}$)	MNBE (%)	MNGE (%)	UPA (%)
O ₃	11.1	25.4	12.98%	24.06%	11.02%
NO ₂	-2.2	17.3	5.89%	49.78%	-23.83%
SO ₂	-6.8	7.1	-41.73%	43.94%	-47.53%
PM ₁₀	-14.4	17.1	-23.62%	35.57%	-56.82%

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.




Fig. 1. Selected domains and topography of the final study areas (D4): North-eastern IP – $322 \times 259 \text{ km}^2$ – (left) and Central IP – $181 \times 214 \text{ km}^2$ – (right) domains.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.




Fig. 2. Atmospheric processes contribution to net O₃ concentration in the 1st simulated vertical layer (0–38 m a.g.l.), O₃ hourly average concentration and O₃ measured in rural background air quality stations of the NEIP (up-Agullana AQS) and the CIP domain (down-Guadarrama AQS) during 18 June 2004.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[⏴](#)[⏵](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.




Fig. 3. Atmospheric processes contribution to net O₃ concentration in the first simulated vertical layer (0–38 m a.g.l.), O₃ hourly average concentration and O₃ measured in urban background air quality stations of the NEIP (BCN-Ciudadella AQS) and the CIP domain (down-Aranjuez AQS) during 18 June 2004.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[⏴](#)[⏵](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.




Fig. 4. Hourly maximum O₃ surface concentration location (circles) in the coastal (up) and the central (down) domains during the 17–18 June 2004 episode, simulated by WRF-ARW/HERMES-2004/CMAQ. Squares: concentrations above the EU human health protection limit (180 µg m⁻³, Dir 2002/3/EC).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Contribution of
processes affecting
atmospheric
dynamics**

M. Gonçalves et al.




Fig. 5. $10 \mu\text{g m}^{-3}$ isosurface of chemical production (red) and destruction (blue) of O_3 for the CIP (up) and NEIP domains (down) at 10:00 and 18:00 UTC of 17–18 June 2004.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Contribution of
processes affecting
atmospheric
dynamics**

M. Gonçalves et al.




Fig. 6. $10 \mu\text{g m}^{-3}$ isosurface of O_3 net transport (horizontal + vertical advection) – positive contribution in red and negative contribution in blue – for the CIP (up) and NEIP domains (down) at 10:00 and 18:00 UTC of 17–18 June 2004.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.




Fig. 7. Subdomains MAX1 and MAX2 (in yellow) selected in the NEIP (up) – MAX1: LAT: 41.63–41.72, LON: 2.08–2.32; MAX2: LAT: 41.67–41.85, LON: 2.42–2.54 – and CIP domains (down) – MAX1: LAT: 40.17–40.35, LON: –3.55–(–3.43); MAX2: LAT: 40.47–40.65; –3.36–(–3.24) –. Maximum hourly concentrations above $180 \mu\text{g m}^{-3}$ in red, concentrations below this limit in purple. Evolution of the modelled PBL height in the NEIP and CIP subdomains.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.




Fig. 8. Atmospheric processes contribution to net O_3 density ($g\ m^{-2}$) under the modelled PBL during 17–18 June 2004. Averaged contributions for the NEIP-MAX1 and MAX2 domains, and CIP-MAX1 and MAX2 domains.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.




Fig. 9. Atmospheric processes contribution to net NO_x density (g m⁻²) under the modelled PBL during 17–18 June 2004. A veraged contributions for the NEIP-MAX1 and MAX2 domains, and CIP-MAX1 and MAX2 domains. 18495

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.




Fig. 10. Atmospheric processes contribution to net NMVOCs density (g m^{-2}) under the modelled PBL during 17–18 June 2004. Averaged contributions for the NEIP-MAX1 and MAX2 domains, and CIP-MAX1 and MAX2 domains.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Contribution of processes affecting atmospheric dynamics

M. Gonçalves et al.




Fig. 11. Gas-phase chemistry and emissions contributions to net $\text{NO}_x\text{-NMVOCs-O}_3$ density (g m^{-2}) under the modelled PBL height (in red) during 17–18 June 2004 for the NEIP-MAX1 and MAX2 domains, and the CIP-MAX1 and MAX2 domains.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion