

This discussion paper is/has been under review for the journal *Atmospheric Chemistry and Physics (ACP)*. Please refer to the corresponding final paper in *ACP* if available.

**Satellite-derived
radiative impact of
tropospheric NO₂**

A. P. Vasilkov et al.

Impact of tropospheric nitrogen dioxide on the regional radiation budget

**A. P. Vasilkov¹, J. Joiner², L. Oreopoulos², J. F. Gleason², P. Veeffkind³,
E. Bucselá⁴, E. A. Celarier⁵, R. J. D. Spurr⁶, and S. Platnick²**

¹Science Systems and Applications Inc., Lanham, MD, USA

²Goddard Space Flight Center, Greenbelt, MD, USA

³Royal Netherlands Meteorological Institute (KNMI), de Bilt, The Netherlands

⁴SRI International, Menlo Park, CA, USA

⁵University of Maryland, Baltimore County, USA

⁶RT Solutions, Cambridge, MA, USA

Received: 11 March 2009 – Accepted: 23 May 2009 – Published: 3 June 2009

Correspondence to: A. P. Vasilkov (alexander.vassilkov@ssaihq.com)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Following the launch of several satellite ultraviolet and visible spectrometers including the Ozone Monitoring Instrument (OMI), much has been learned about the global distribution of nitrogen dioxide (NO_2). NO_2 , which is mostly anthropogenic in origin, absorbs solar radiation at ultraviolet and visible wavelengths. We parameterized NO_2 absorption for fast radiative transfer calculations. Using this parameterization with cloud, surface, and NO_2 information from different sensors in the NASA A-train constellation of satellites and NO_2 profiles from the Global Modeling Initiative (GMI), we compute the global distribution of net atmospheric heating due to tropospheric NO_2 for January and July 2005. We assess the impact of clouds and find that because most of NO_2 is contained in the boundary layer in polluted regions, the cloud shielding effect can significantly reduce the net atmospheric heating due to NO_2 . We examine the effect of diurnal variations in NO_2 emissions and chemistry on net atmospheric heating and find only a small impact of these on the daily-averaged heating. While the impact of NO_2 on the global radiative forcing is small, locally it can produce instantaneous net atmospheric heating of $2\text{--}4\text{ W/m}^2$ in heavily polluted areas. We also examine the sensitivity of NO_2 absorption to various geophysical conditions. Effects of the vertical distributions of cloud optical depth and NO_2 on net atmospheric heating and downwelling radiance are simulated in detail for various scenarios including vertically-inhomogeneous convective clouds observed by CloudSat. The maximum effect of NO_2 on downwelling radiance occurs when the NO_2 is located in the middle part of the cloud where the optical extinction peaks.

1 Introduction

The absorption of incoming solar and outgoing terrestrial radiation by trace gases is one of the key factors in the Earth's radiative budget and plays a leading role in climate change (Hansen et al., 2007). Carbon dioxide (CO_2) is the largest anthropogenic cli-

Satellite-derived radiative impact of tropospheric NO_2

A. P. Vasilkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Satellite-derived
radiative impact of
tropospheric NO₂**

A. P. Vasilkov et al.

mate forcing gas, but other trace gases are also important including methane (CH₄), tropospheric ozone (O₃), nitrous oxide (N₂O), and halocarbons such as chlorofluorocarbons (CFCs) (Forster et al., 2007). Little attention has been paid to nitrogen dioxide (NO₂), because its effect on the global radiative budget, particularly the radiative forcing, is small. However, on a local scale, enhanced NO₂ in polluted areas can change the partitioning of absorbed solar radiation between the atmosphere and surface; it produces atmospheric heating in the troposphere and contributes to dimming (reduction of solar radiation) at the surface.

Solomon et al. (1999) showed that NO₂ can produce significant absorption of solar radiation locally under some conditions. During several days of ground-based observations in a moderately polluted area (Boulder, Colorado, USA), the peak absorption of downwelling radiation by NO₂ was 5–12%. The estimated peak net heating was in the range of 5–30 W/m². The high peak values appeared to be influenced by lightning-generated NO_x associated with thunderstorms.

NO₂ trends spanning a decade have been derived with data from the Global Ozone Monitoring Experiment (GOME), the Scanning Imaging Absorption Chartography (SCIAMACHY), and the Ozone Monitoring Instrument (OMI) (e.g., Richter et al., 2005; van der A et al., 2006, 2008). In some areas, NO₂ has increased significantly. For instance in the Shanghai region, tropospheric NO₂ increased ~29%/year since 1996. In other areas, NO₂ has been decreasing. Reductions of up to 7% per year have been observed in Europe and parts of the eastern United States.

In this paper, we compute the global distribution of the net atmospheric heating (NAH) due to tropospheric NO₂ with an emphasis on polluted areas. We also examine the sensitivity of the NO₂ NAH to various geophysical parameters including the vertical cloud distribution. Here, we use several data sets from the NASA A-train constellation of afternoon satellites flying within fifteen minutes of each other near 13:30 local solar time. These include cloud and surface properties from the NASA Aqua Moderate Resolution Imaging Spectroradiometer (MODIS) on the Auqua platform and tropospheric NO₂ and cloud pressure from OMI, which flies on the NASA Aura satellite.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Satellite-derived radiative impact of tropospheric NO₂

A. P. Vasilkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The organization of the paper is as follows: in Sect. 2, we develop a parameterization of NO₂ absorption that is used in a radiative transfer scheme to compute NO₂ NAH. In Sect. 3, we show how the vertical profiles of NO₂ and cloud optical depth affect NAH and downwelling surface radiances for clear-sky and cloudy conditions. In Sect. 4, we present computed global and regional distributions of the tropospheric NO₂ NAH based on NO₂ amounts derived from OMI observations and cloud data taken from MODIS and OMI retrievals. Conclusions are given in Sect. 5.

2 Parameterization of NO₂ absorption

NO₂ absorbs solar radiation throughout the spectral range 300–650 nm with a peak near 400 nm. The temperature dependence of the NO₂ absorption is weak, so we neglect it and use the cross section at 294 K from Vandaele et al. (1998). In this work, we neglect a small portion (<1%) of absorbed energy that goes to NO₂ fluorescence (Thornton et al., 2000).

Figure 1 shows the optical thickness, τ , for a vertical path through an NO₂ column amount of 0.5 Dobson Units (DU, 1 DU = 1.343 × 10¹⁶ mol/cm²), that is typical of heavily polluted areas (e.g., Cede et al., 2005) and the approximate spectral absorption of solar irradiance $A = F\tau \approx F(1 - \exp(-\tau))$, where F is the extraterrestrial solar irradiance. The low values of optical thickness even for high NO₂ amounts indicate that NO₂ is a relatively weak absorber. According to the data in Fig. 1, most solar absorption takes place in the spectral range of 350–500 nm.

To reduce the computational burden for heating calculations, we parametrize the spectral NO₂ absorption coefficients following Chou and Lee (1996). The transmission function averaged over a spectral band is first calculated from

$$T(w/\mu) = \frac{\int_{\lambda_1}^{\lambda_2} F(\lambda) \exp[-a(\lambda)w/\mu] d\lambda}{\int_{\lambda_1}^{\lambda_2} F(\lambda) d\lambda}, \quad (1)$$

where w is the NO_2 column amount, F is the extraterrestrial solar flux, a is the NO_2 absorption coefficient, and μ is the cosine of the solar zenith angle (SZA). An effective absorption coefficient, a_{eff} , for a given spectral band is then derived by inverting the transmittance function:

$$a_{\text{eff}} = \ln[T(w/\mu)]\mu/w. \quad (2)$$

Variations of a_{eff} with w/μ are relatively small and do not exceed 5% for realistic values of w/μ . Table 1 gives the mean values of a_{eff} for the two spectral bands in the Chou and Suarez (2002) algorithm where NO_2 absorption is significant.

To assess the overall efficiency of the parameterization, we compute NO_2 NAH using the high spectral resolution and effective absorption coefficients in clear skies for different SZAs. The calculations are performed with the Linearized Discrete-Ordinate Radiative Transfer-Rotational Raman Scattering (LIDORT-RRS) code that does not account for polarization (Spurr et al., 2008). In general, the NAH is defined as

$$Q = (F_d - F_u)_{\text{TOA}} - (F_d - F_u)_{\text{surface}}, \quad (3)$$

where F_d and F_u are the downward and upward irradiances, respectively, and the subscripts TOA and surface refer to the top-of-atmosphere and ground, respectively. All NAHs reported here are those due to tropospheric NO_2 , i.e., $Q = Q_{\text{trop NO}_2} - Q_{\text{no trop NO}_2}$.

Figure 2 shows that the difference between the NAH computed at high spectral resolution and with the parameterization does not exceed 5% for $\text{SZA} < 60^\circ$ in either spectral band. To compute the NAH in Fig. 2, we use a total NO_2 column amount of 0.5 DU distributed uniformly in the lowest km of the atmosphere. The surface albedo is set to 0.05.

It should be noted that no other atmospheric absorbers are included in this computation and computations carried out for sensitivity studies described in Sect. 3. An important absorber in the spectral range of 320–700 nm is ozone. Ozone absorption can affect NO_2 NAH because ozone absorption in the stratosphere reduces downwelling flux thus decreasing NO_2 heating in the planetary boundary layer where most NO_2 is

Satellite-derived radiative impact of tropospheric NO_2

A. P. Vasilkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



confined in polluted land regions. However, this effect is small. We performed computations of NAH for three cases: ozone and NO₂ $Q_{O_3+NO_2}$, ozone alone, Q_{O_3} , and NO₂ alone, Q_{NO_2} , in the spectral range of 320–400 nm. The difference $Q_{O_3+NO_2} - Q_{O_3}$, that characterizes NO₂ heating in the presence of ozone absorption, appears to be less than Q_{NO_2} just by 1.7% (at SZA=45°). In computations of global and regional NAH described in Sect. 4, we accounted for ozone absorption and calculated the NO₂ NAH as the difference $Q_{O_3+NO_2} - Q_{O_3}$.

3 Sensitivity studies

To better understand the global calculations of NO₂ NAH that are shown in the next section, we first perform a series of sensitivity tests. These tests are conducted at a SZA of 45° with the high spectral resolution absorption coefficients.

Our calculations show that the dependence of the NAH on the vertical distribution of tropospheric NO₂ in clear skies is relatively small. For instance, when the top of a 0.5 DU 1 km thick layer of NO₂ increases from 1 to 3 km, the NAH increases ~8% in the 400–700 nm band and ~15% in 320–400 nm band. The total increase of the NAH in the two bands is ~10%.

Figure 3 shows the dependence of the NAH on surface albedo for clear skies with 0.5 DU of NO₂ placed in the lowest km. The significant increase of NAH with surface albedo is a result of enhanced photon path lengths through the NO₂ layer at higher albedos owing to an increase in photons reflected from the surface.

We next investigate the impact of clouds on the tropospheric NO₂ NAH. We take the phase function to have the Henyey-Greenstein form with an asymmetry parameter of 0.85. This corresponds to water clouds with equivalent droplet radius of 6 μm (Hu and Stamnes, 1993).

The impact of clouds on the NAH is significant and depends on both the vertical cloud structure and the placement of NO₂ with respect to the cloud. For NO₂ above a cloud layer, the cloud increases the tropospheric (or stratospheric) NO₂ NAH similar to

Satellite-derived radiative impact of tropospheric NO₂

A. P. Vasilkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the effect of a higher surface albedo.

We next consider an NO₂ layer that lies entirely below a cloud. Here, we use a vertically uniform cloud with a geometrical thickness of 1 km. The cloud bottom was placed just above the NO₂ layer located in the lowest km. In this situation, the cloud reduces the tropospheric NO₂ NAH. The NAH dependence on cloud optical depth, τ , is shown in Fig. 4.

In this work, we define the NAH for a partly cloudy scene as $Q=Q_{\text{clr}}(1-f)+Q_{\text{cld}}f$, where f is the geometrical cloud fraction, and Q_{clr} and Q_{cld} are the NAH for clear and overcast skies, respectively. This definition follows the so-called independent pixel approximation and assumes there is no radiative interaction between clear and cloudy pixels. For the above scenario with the cloud placed just above the NO₂ layer (in the lowest km), NAH decreases linearly with increasing cloud fraction. The slope of the NAH dependence on cloud fraction is dependent on τ .

We next perform radiative transfer simulations in realistic vertically extended clouds to study the effect of the vertical distribution of NO₂ within a cloud. We use the two cloud optical extinction profiles shown in the left panel of Fig. 5. They are derived from CloudSat radar reflectivities (Stephens et al., 2002) and MODIS TOA reflectances. We use the CloudSat standard product 2B-TAU that provides cloud optical depth by layer (Stephens et al., 2008). Vasilkov et al. (2008) and Ziemke et al. (2009) used these profiles to examine ozone absorption and rotational-Raman scattering within clouds and showed that they are typical for tropical deep convection. The total cloud optical depths are 32 and 66 for profiles 1 and 2, respectively. The cloud tops as detected by both MODIS and CloudSat are within 150–250 hPa for both profiles. The profiles show the clouds to be optically thin near the tops.

The middle panel of Fig. 5 shows the NAH (300–700 nm) as a function of NO₂ layer midpoint for the two CloudSat profiles. The NO₂ layer is 0.5 DU distributed uniformly within 1 km. Also shown is the NAH computed for a vertically uniform distribution of NO₂ throughout the clouds (0.5 DU). The NAH is minimal when the NO₂ is located in the lower-most portions of the clouds. For profile 1, the NAH increases rapidly for NO₂

Satellite-derived radiative impact of tropospheric NO₂

A. P. Vasilkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



layer heights between ~ 3 and 7 km as the in-cloud multiple scattering increases. The NAH then decreases with NO_2 layer height for layers near the cloud top. The NAH for profile 2 exhibits a plateau when the NO_2 is located in between the two distinct cloud layers. The NO_2 layer heights at which the NAH are equal to values for the vertically uniform NO_2 distribution throughout the clouds differ noticeably for two cloud profiles.

To simulate the Solomon et al. (1999) measurements of downwelling zenith surface radiance, we also calculate the fractional change of surface zenith radiances due to tropospheric NO_2 : $I/I_0 - 1$, where I and I_0 are the zenith radiances at 400 nm calculated with and without NO_2 , respectively. Solomon et al. (1999) performed calculations assuming a vertically uniform cloud and suggested that the cloud effect on NO_2 absorption would be greatest if the NO_2 is placed near the cloud top.

The right panel of Fig. 5 shows the changes of the zenith radiance as a function of the NO_2 layer height for the two CloudSat profiles along with that computed for a vertically uniform distribution of NO_2 throughout the clouds. It is interesting to note that the maximum effect of NO_2 absorption on the zenith radiance occurs when the NO_2 is located near the peak of the cloud extinction. The maximum photon path enhancement takes place in the most scattering portion of clouds, and that is where the NO_2 absorption effect on the downwelling radiance is also at a maximum.

4 Global NO_2 net atmospheric heating calculations

In this section, we perform global radiative transfer (RT) calculations with a stand-alone version of the algorithm developed by Chou and Suarez (2002) (henceforth referred to as CS). The algorithm has been used in various atmospheric models in the NASA Goddard Laboratory for Atmospheres, We first provide brief descriptions of the CS algorithm and input data sets.

Satellite-derived radiative impact of tropospheric NO_2

A. P. Vasilkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.1 Chou-Suarez (CS) radiative transfer algorithm

In the CS algorithm, the short-wave spectrum is divided into 8 bands in the ultraviolet (UV) and photo-synthetically active (PAR) regions with a single ozone absorption coefficient and Rayleigh scattering coefficient in each band. NO₂ absorption is similarly accounted for in two bands: 320–400 and 400–700 nm using the effective absorption coefficients listed in Table 1.

Reflection and transmission of cloud layers are calculated using the δ -Eddington approximation, and fluxes are computed with a two-stream model. Data provided by Yang et al. (2000) for ice clouds and Tsay et al. (1989) for water clouds are used to derive an effective band-mean single scattering albedo, extinction coefficient, and asymmetry factor. Aerosol is not included in our computations. Roughly, aerosol acts like a cloud of low optical depth and the OMI cloud algorithm treats it as such.

4.2 Input data sets

4.2.1 OMI NO₂ retrievals

Here we briefly describe the standard OMI NO₂ algorithm and retrievals. Additional details can be found in Bucsela et al. (2006, 2008) and Celarier et al. (2008). OMI provides daily global coverage with a swath width of 2600 km (Levelt et al., 2006). The NO₂ level 2 retrievals are reported at the nominal OMI spatial resolution (13 km×24 km at nadir).

NO₂ total column amounts are retrieved with a differential optical absorption spectroscopy (DOAS) technique using OMI TOA reflectance spectra. First, NO₂ slant column amounts are estimated by spectral fitting within the wavelength region 415–465 nm (Bucsela et al., 2006; Celarier et al., 2008). This spectral range falls within OMI's visible detector that has a spectral resolution of about 0.6 nm. Initial vertical column amounts are calculated using air mass factors (AMFs) for a stratospheric NO₂ distribution, and are approximately geometric. Tropospheric AMFs are derived with

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



gridded annual mean NO₂ profiles from the GEOS-Chem model (Bey et al., 2001).

The NO₂ stratosphere-troposphere separation is based on the fact that the stratospheric field has small gradients, particularly in the zonal direction. The slowly varying spatial component of the total NO₂ column is identified as the stratospheric field and the rest is assigned to the troposphere. The initial tropospheric column amount is then multiplied by the ratio of the stratospheric-to-tropospheric AMFs.

The OMI NO₂ retrievals have been compared with in situ, ground-based, and aircraft measurements. Using in situ NO₂ aircraft measurements, Bucselá et al. (2008) conclude that the OMI algorithm yields NO₂ amounts lower than the in situ columns by a factor of 0.86. Similarly, Celarier et al. (2008) report that the OMI tropospheric columns are underestimated by 15–30%. Comparing the total NO₂ amounts with those derived from ground-based direct-Sun Brewer measurements, Wenig et al. (2008) show that monthly averages of coinciding measurements correlate well, but OMI retrievals are about 25% lower than the Brewer data.

Because tropospheric NO₂ has significant diurnal variations (Cede et al., 2005; Boersma et al., 2008), it is important to note that the OMI equator crossing time is 13:30 local time. The diurnal cycle of tropospheric NO₂ is governed by emissions that are mostly due to human activities in urban regions and chemical loss that is strongest at midday. Therefore, the NO₂ concentration exhibits a minimum near the afternoon OMI overpass time in urban regions (Boersma et al., 2008). As expected, OMI retrieves lower NO₂ column amounts (20–30%) over urban areas than SCIAMACHY, which has an equator crossing time of 10:00 (Boersma et al., 2008).

To compute global NAH, we use monthly mean values of the OMI NO₂ tropospheric column gridded to a resolution of 1° latitude × 1° longitude. We use the following filters to compute the gridbox means:

1. View zenith angles < 50°; this removes large-footprint pixels at the swath edge.
2. Radiative cloud fractions (percent of pixel radiance contributed by cloudy portion of the scene) < 30%. This removes pixels where significant amounts of NO₂ are

Satellite-derived radiative impact of tropospheric NO₂

A. P. Vasilkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Satellite-derived radiative impact of tropospheric NO₂

A. P. Vasilkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



shielded by clouds. The use of a threshold less than 30% gives similar results but worse cover.

3. South Atlantic anomaly (SAA) flag bit set to zero or SAA flag > 0 and the fitting $\chi^2 < 1500$.
- 5 4. Processing quality flag bits 0 and 1 equal to zero (for information about the flags, see http://toms.gsfc.nasa.gov/omi/no2/OMNO2_readme.pdf).

Figure 6 shows the well-known seasonal variability in tropospheric NO₂ column amounts as observed by OMI. In industrial regions, such as the eastern US, Europe, and Southeast Asia, the dominant source of NO₂ emissions is anthropogenic. These areas have a column NO₂ maximum in the winter due to the longer lifetime of NO_x at that time of year (van der A et al., 2008). The effects of seasonal changes in biomass burning in Central Africa are also reflected in NO₂ columns.

We use monthly-mean NO₂ vertical distributions based on a simulation from the Global Modeling Initiative (GMI) chemistry-transport model (Duncan et al., 2007). We find that in polluted regions, most of the NO₂ is contained within a layer about 200 hPa thick above the surface. Rather than scaling the entire NO₂ profile to give the OMI-derived NO₂ column, we adjust the model NO₂ profiles such that the total column amounts are equal to the gridded OMI monthly mean value by adding or subtracting an appropriate amount in a 200 hPa thick layer just above surface. While there is little sensitivity to the NO₂ profile shape in clear skies, the profile shape in cloudy conditions significantly affects the computed NAH. The adjustment of NO₂ profiles provides a more accurate estimate of the cloud effect.

4.2.2 Cloud data sets

We use retrievals of cloud optical depth, effective radius, and phase from MODIS daily gridded 1° × 1° level 3 (L3) data sets (Platnick et al., 2003) as described in detail by Joiner et al. (2009). The dataset includes separate statistical descriptions of parame-

ters for ice and water clouds. We use the gridbox-mean effective radii and cloud optical depths for ice and water clouds, along with their associated cloud fractions. Joiner et al. (2009) showed that the use of daily gridbox-mean values, as opposed to the full gridbox statistical distributions that are provided in the MODIS data set, provides sufficiently good accuracy for RT calculations related to trace-gas absorption.

To compute NAH, we also need to specify the cloud extent. As in Joiner et al. (2009), we use optical centroid cloud pressures from the OMI rotational-Raman algorithm (Joiner et al., 2004; Joiner and Vasilkov, 2006). The optical centroid cloud pressures from this algorithm are similar to those derived from oxygen dimer and oxygen A-band absorption (Sneep et al., 2008) and are significantly different from the cloud-top pressure (Vasilkov et al., 2008). The optical centroid cloud pressure is more appropriate than cloud-top pressure for radiative calculations in the visible and ultraviolet because it accounts for photon penetration into the cloud (Ziemke et al., 2009).

We distribute the cloud optical depth uniformly in a 1 km layer centered at the OMI centroid cloud pressure. This provides a good first order approximation for cloud effects on trace-gas absorption at visible wavelengths when the absorber is well mixed (Ziemke et al., 2009). However, errors will occur in more complex situations such as when the absorber and cloud distributions are non-uniform as shown in Sect. 3. It is difficult to quantify the errors as the exact distributions in such situations are unknown.

4.2.3 Surface albedo and other ancillary data sets

As in Joiner et al. (2009), we specify the surface albedo separately for direct and diffuse fluxes using 16-day gridded albedos from the MODIS filled-land surface albedo product (MOD43B3) (Lucht et al., 2000). Over ocean, we use a model of the surface albedo from Jin et al. (2004) with the 2 m wind speed from the Goddard Earth Observing System 5 Data Assimilation System (GEOS-5 DAS) (Rienecker et al., 2007). Temperature, water vapor, and ozone profiles are also taken from the GEOS-5 DAS.

Satellite-derived radiative impact of tropospheric NO₂

A. P. Vasilkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.3 Results

4.3.1 Diurnally-averaged net atmospheric heating

Here, we compute monthly-mean NO_2 NAH by averaging values calculated every two hours. We fix the NO_2 column amount at the OMI gridded monthly mean value, i.e., the value near 13:30 local time. This will produce systematic errors in the diurnally-averaged NAH that will depend upon the diurnal variations in NO_2 . The error magnitude is estimated in Sect. 4.3.3. Although we use monthly mean values of the tropospheric NO_2 column for computing NAH, we use daily cloud data from MODIS and OMI to properly account for the high variability of cloud optical thickness and pressure and the non-linearity of these cloud parameters with respect to the NO_2 NAH.

Figure 7 shows the global distribution and seasonal variations in the NAH along with the cloud effect. The spatial distribution of the NAH primarily reflects the global distribution of NO_2 . However, the effect of SZA is also evident in the NAH. For example, NO_2 column amounts in European urban and industrial areas are somewhat higher than over Central Africa in January. The NAH over Europe, however, is lower than over Central Africa owing to the reduced amount of solar irradiance at high latitudes in January. The seasonal maximum of diurnally-averaged NAH for polluted extra-tropical regions is in summer owing to the greater amount of solar irradiance.

The cloud effect is defined as $Q_{\text{allsy}}/Q_{\text{clearsky}} - 1$. In polluted regions, large values of this parameter are mainly due to shielding of the planetary boundary layer (PBL) where most of the NO_2 resides. This decreases atmospheric heating as shown in Fig. 7. The decrease (in terms of percent) is larger in winter than summer for most polluted areas.

Over the ocean, NO_2 is not so restricted to the PBL as it is in polluted land regions. Most of the tropospheric NO_2 column over ocean results from pollution transport, NO_2 brought down from the stratosphere through tropopause folds, and lightning. In this situation clouds can increase heating when NO_2 is present above them as shown in Sect. 3. However, the total amount of NO_2 and associated heating over oceans is

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



relatively small.

On the west coast US and in southern Europe during July, the net cloud effect is close to zero. In these areas there is a compensation of the cloud shielding and cloud enhancement effects.

5 4.3.2 Instantaneous net atmospheric heating

Figure 8 shows detailed maps of maximum daily (instantaneous) NAH over North America and Southeast Asia for January and July 2005. As noted above, the computed daily maxima assume a fixed NO_2 column amount equal to the OMI-derived value. The winter instantaneous maximum NAH values are higher than summer values in many polluted areas in contrast with the diurnally-averaged NAH values. The instantaneous maximum NAH values are approximately a factor of 2 higher than the diurnally-averaged.

4.3.3 Effect of time variations in sources and chemistry

In the above calculations, we assumed a constant NO_2 amount equal to the OMI-derived value near 13:30 local time. Here, we examine the effect of diurnal variations in NO_2 on the calculated NAH. Figure 9 shows diurnal variations in the tropospheric NO_2 column for two geographic regions in August 2006 as given by Boersma et al. (2008):

1. An urban area in the northeastern United States where NO_2 amounts are dominated by fossil fuel burning
2. southern Africa, where NO_2 is produced primarily from biomass burning

Three scenarios are considered:

1. constant NO_2 amounts (equal to the OMI amount);
2. constant NO_x emissions;

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3. diurnally varying NO_x emissions.

NO₂ columns for the latter two cases were derived with the GEOS-Chem model (Boersma et al., 2008). To calculate SZAs needed for the NAH computations, we used 40° N and 20° S latitude for the US and African areas, respectively, for mid-August.

The instantaneous NAH values for these different scenarios are shown in Fig. 9 for cloud-free conditions. For the case of constant NO₂, the variation of NAH is determined only by SZA changes. There are significant differences (up to ~30%) in the instantaneous NAH values calculated for the various scenarios and similar differences in the peak values for the urban case. However, daily-integrated and averaged NAHs differ less. Daily-averaged NAHs are lower for the constant NO₂ amount as compared with the diurnally varying NO₂ columns (11–14% for the urban case and 1–6% for the biomass burning scenario). These differences are relatively small and represent an upper bound on the error in the computed diurnally-averaged NAH shown in Sect. 4.3.1. Note that the cloud effect reduces the error for an all-sky monthly mean.

4.3.4 Effect of spatial resolution

The effect of spatial resolution on the computed mean NO₂ column is well known; averaging the data on a finer grid will produce larger peak values in polluted areas. The data may be averaged at a finer resolution than the actual pixel sizes. Because of the 32 day repeat cycle of the Aura spacecraft, the OMI pixel centers vary from day to day. Therefore, by spatial oversampling, some information about NO₂ distributions may be obtained at a higher resolution than the individual pixel sizes. In Fig. 10, we compute the maximum daily NAH as in Fig. 8, but using NO₂ columns gridded to 0.1°×0.1° resolution for a region encompassing Los Angeles. In order to maximize the coverage, we averaged all data from 1 January 2005–31 December 2007 and included cloud fractions up to 100%. At this resolution, instantaneous NAHs are approximately a factor of 2 higher than at 1°×1° resolution and the sub-grid variability is significant.

Satellite-derived radiative impact of tropospheric NO₂

A. P. Vasilkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 Discussion and conclusions

We showed that absorption by NO_2 can be accurately accounted for in standard radiative transfer algorithms using parametrized band-averaged absorption coefficients. Our sensitivity studies showed that NO_2 net atmospheric heating in cloudy conditions strongly depends upon the vertical distributions of both NO_2 and cloud optical depth in realistic deep convective clouds. Effects of the vertical distributions of cloud optical depth and NO_2 on downwelling radiance were simulated for vertically-inhomogeneous convective clouds observed by CloudSat. The simulations showed the maximum effect of NO_2 on downwelling radiance occurred when the NO_2 was located in the middle part of the cloud where the optical extinction peaked. Note that the net heating dependence on layer height that we computed would be virtually the same for any visible or near-IR absorber.

We computed the global distribution of daily maximum and diurnally-averaged NO_2 net heating using cloud, surface, and NO_2 information from different sensors in the A-train constellation of satellites and NO_2 profiles from the GMI. While the impact of NO_2 on the global radiative forcing was small, locally it could produce instantaneous NAH of $2\text{--}4\text{ W/m}^2$ in heavily polluted regions. We assessed the impact of clouds and found that because most of NO_2 was contained in the boundary layer in polluted regions, the cloud shielding effect significantly reduced the NAH due to NO_2 . We also examined the effect of diurnal variations in NO_2 emissions and chemistry on NAH and found only a small impact of these on the daily-averaged heating. Uncertainties in the NAH are due mostly to the uncertainties in the NO_2 satellite observations. For clear skies, NAH is practically proportional to NO_2 amount. We show that the dependence of NAH on the vertical profile of tropospheric NO_2 is small and the effect of NO_2 diurnal variations on the daily-integrated NAH is small as well. Thus, the NO_2 amount uncertainties directly propagate into NAH and relative error in NAH is determined by relative error in the NO_2 observations. Using the estimates of the NO_2 observations given in Sect. 4.2.1, we can state that NAH is underestimated by 20–25% for clear skies. The error estimate

Satellite-derived radiative impact of tropospheric NO_2

A. P. Vasilkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



is more complicated for cloudy skies. NAH nonlinearly depends on cloud optical depth, cloud height, NO_2 amount, and NO_2 vertical distribution. However, since clouds reduce the heating for most of the important areas, the clear sky error can be considered as the upper limit of the NAH uncertainties.

5 Our computed NAHs are somewhat lower than peak values reported by Solomon et al. (1999). However, it should be noted that the Solomon et al. (1999) measurements were highly localized in both time and space and were likely influenced by thunderstorms that may have produced small-scale NO_2 plumes. For our global calculations, we used monthly-averaged NO_2 column amounts in gridboxes of $\sim 100 \text{ km} \times 100 \text{ km}$.
10 Averaging at higher spatial resolutions results in higher peak NO_2 column amounts, greater heating, and peak values approaches those estimated by Solomon et al. (1999).

The radiative impact of NO_2 at the surface is dimming (a reduction in solar radiation). Dimming has been increasing over the last several decades primarily due to aerosol indirect effects (e.g., Stanhill and Cohen, 2001). Various estimates of dimming have
15 been reported, and they vary with location. Stanhill and Cohen (2001) estimate a reduction of 20 W/m^2 during the past 50 years (2.7% per decade) while Liepert (2002) reports a decline of 7 W/m^2 or 4% from 1961 to 1990. Our calculations of the surface irradiance reduction by NO_2 is $\sim 1 \text{ W/m}^2$ in the most polluted areas (Southeast Asia).
20 Therefore, although NO_2 may be of importance locally in radiative forcing, the trace gas is not likely contributing significantly to the observed global dimming.

NO_2 absorption is currently not included in most general circulation model radiative transfer codes. We have shown that it could be easily incorporated using effective absorption coefficients and distributions derived from satellite data or carried on line. In
25 this way, heating from stratospheric NO_2 , which is comparable to that from tropospheric NO_2 in slightly to moderately polluted conditions, can also be included.

Acknowledgements. The authors thank the OMI and MODIS science teams for the processing and distribution of data sets used here. We also thank Ronald Cohen, Richard Stolarski, and Arlindo da Silva for helpful comments. This material is based upon work supported by the Na-

**Satellite-derived
radiative impact of
tropospheric NO_2** A. P. Vasilkov et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

tional Aeronautics and Space Administration under agreement NNG06HX18C issued through the Science Mission Directorate for the Aura Science Team. Lazaros Oreopoulos gratefully acknowledges support for this work by the US Department of Energy, Office of Science, Office of Biological and Environmental Research, Environmental Sciences Division as part of the ARM program under grant DE-FG02-07ER64354.

References

- Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B. D., Fiore, A. M., Li, Q., Liu, H. Y., Mickley, L. J., and Schultz, M. G.: Global modeling of tropospheric chemistry with assimilated meteorology: Model description and evaluation, *J. Geophys. Res.*, 106, 23073–23095, doi:10.1029/2001JD001480, 2001. 12684
- Boersma, K. F., Eskes, H. J., and Brinksma, E. J.: Error analysis for tropospheric NO₂ retrievals from space, *J. Geophys. Res.*, 109, D04311, doi:10.1029/2003JD003962, 2004.
- Boersma, K. F., Jacob, D. J., Eskes, H. J., Pinder, R. W., Wang, J., and van der A, R. J.: Intercomparison of SCIAMACHY and OMI tropospheric NO₂ columns: Observing the diurnal evolution of chemistry and emissions from space, *J. Geophys. Res.*, 113, D16S26, doi:10.1029/2007JD008816, 2008. 12684, 12688, 12689
- Bucsela, E. J., Celarier, E. A., Wenig, M. O., Gleason, J. F., Veefkind, J. P., Boersma, K. F., and Brinksma, E. J.: Algorithm for NO₂ vertical column retrieval from the Ozone Monitoring Instrument, *IEEE Trans, Geosci. Remote Sens.*, 44, 1245–1258, doi:10.1109/TGRS.2005.863715, 2006. 12683
- Bucsela, E. J., Perring, A. E., Cohen, R. C., Boersma, K. F., Celarier, E. A., Gleason, J. F., Wenig, M. O., Bertram, T. H., Wooldridge, P. J., Dirksen, R., and Veefkind, J. P.: Comparison of tropospheric NO₂ from in situ measurements with near-real-time and standard product data from OMI, *J. Geophys. Res.*, 113, D16S31, doi:10.1029/2007JD008838, 2008. 12683, 12684
- Cede, A., Herman, J., Richter, A., Krotkov, N., and Burrows, J.: Measurements of nitrogen dioxide total column amounts using a Brewer double spectrophotometer in direct Sun mode, *J. Geophys. Res.*, 113, D05304, doi:10.1029/2005JD006585, 2005. 12678, 12684
- Celarier, E. A., Brinksma E. J., Gleason J. F., et al.: Validation of Ozone Monitoring Instrument

ACPD

9, 12675–12706, 2009

Satellite-derived radiative impact of tropospheric NO₂

A. P. Vasilkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



nitrogen dioxide columns, *J. Geophys. Res.*, 113, D15S15, doi:10.1029/2007JD008908, 2008. 12683, 12684

Chou, M.-D. and Lee, K.-T.: Parameterizations for the absorption of solar radiation by water vapor and ozone, *J. Atmos. Sci.*, 53, 1203–1208, 1996. 12678

Chou, M.-D. and Suarez, M. J.: A solar radiation parameterization for atmospheric studies, NASA Tech. Memo 104606, Vol. 15, 40 pp., 2002. 12679, 12682

Duncan, B. N., Strahan, S. E., Yoshida, Y., Steenrod, S. D., and Livesey, N.: Model study of the cross-tropopause transport of biomass burning pollution, *Atmos. Chem. Phys.*, 7, 3713–3736, 2007,

<http://www.atmos-chem-phys.net/7/3713/2007/>. 12685

Forster, P., Ramaswamy, V., Artaxo, P., et al.: in: *The Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M. et al., Cambridge Univ. Press, Cambridge, 2007. 12677

Hansen, J., Sato, M., Kharecha, P., Russell, G., Lea, D. W., and Siddall, M.: Climate change and trace gases, *Philos. T. Roy. Soc. A*, 365, 1925–1954, doi:10.1098/rsta.2007.2052, 2007. 12676

Hu, Y. X. and Stamnes, K.: An accurate parameterization of the radiative properties of water clouds, *J. Climate*, 6, 728–742, 1993. 12680

Jin, Z., Charlock, T., Smith, W., and Rutledge, K.: A parameterization of ocean surface albedo, *Geophys. Res. Lett.*, 31, L22301, doi:10.1029/2004GL021180, 2004. 12686

Joiner, J., Vasilkov, A. P., Flittner, D. E., Gleason, J. F., and Bhartia, P. K.: Retrieval of cloud pressure and oceanic chlorophyll content using Raman scattering in GOME ultraviolet spectra, *J. Geophys. Res.*, 109, D01109, doi:10.1029/2003JD003698, 2004. 12686

Joiner, J. and Vasilkov, A. P.: First results from the OMI Rotational Raman Scattering Cloud Pressure Algorithm, *IEEE T. Geosci. Remote*, 44, 1272–1282, 2006. 12686

Joiner, J., Schoeberl, M. R., Vasilkov, A. P., Oreopoulos, L., Platnick, S., Livesey, N. J., and Levelt, P. F.: Accurate satellite-derived estimates of the tropospheric ozone impact on the global radiation budget, *Atmos. Chem. Phys. Discuss.*, 9, 5505–5547, 2009, <http://www.atmos-chem-phys-discuss.net/9/5505/2009/>. 12685, 12686

Levelt, P. F., van der Oord, G. H. J., Dobber, M. R., et al.: The Ozone Monitoring Instrument, *IEEE T. Geosci. Remote*, 44, 1093–1101, 2006. 12683

Liepert, B. G.: Observed reductions of surface solar radiation at sites in the United States and worldwide from 1961 to 1990, *Geophys. Res. Lett.*, 29, 1421, doi:10.1029/2002GL014910,

**Satellite-derived
radiative impact of
tropospheric NO₂**

A. P. Vasilkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2002. 12691

Lucht, W., Schaaf, C. B., and Strahler, A. H.: An Algorithm for the retrieval of albedo from space using semiempirical BRDF models, *IEEE T. Geosci. Remote*, 38, 977–998, 2000. 12686

Platnick, S., King, M. D., Ackerman, S. A., Menzel, W. P., Baum, B. A., Riédi, J. C., and Frey, R. A.: The MODIS cloud products: algorithms and examples from Terra, *IEEE T. Geosci. Remote*, 41, 459–473, 2003. 12685

Richter, A., Burrows, J. P., Nuss, H., Granier, C., and Niemeier, U.: Increase in tropospheric nitrogen dioxide over China observed from space, *Nature*, 437, 129–132, doi:10.1038/nature04092, 2005. 12677

Rienecker, M. M., Suarez M. J., Todling R., et al.: The GEOS-5 data assimilation system - Documentation of versions 5.0.1, 5.1.0, and 5.2.0. NASA Tech. Memo. 2007-104606, vol. 27, edited by: Suarez, M. J., 2007. 12686

Sneep, M., De Haan, J., Stammes, P., Wang, P., Vanbauce, C., Joiner, J., Vasilkov, A. P., and Levelt, P. F.: Three way comparison between OMI/Aura and POLDER/PARASOL cloud pressure products, *J. Geophys. Res.*, 113, D15S23, doi:10.1029/2007JD008694, 2008. 12686

Solomon, S., Portmann, R. W., Sanders, R. W., and Daniels, J. S.: On the role of nitrogen dioxide in the absorption of solar radiation, *J. Geophys. Res.*, 104(D10), 12047–12058, 1999. 12677, 12682, 12691

Spurr, R. J. D., de Haan, J., van Oss, R., and Vasilkov, A. P.: Discrete ordinate radiative transfer in a stratified medium with first order rotational Raman scattering, *J. Quat. Spectrosc. Ra.*, 109(3), 404–425, 2008. 12679

Stanhill, G. and Cohen, S.: Global dimming: a review of the evidence for a widespread and significant reduction in global radiation with discussion of its probable causes and possible agricultural consequences, *Agr. Forest Meteorol.*, 107, 255–278, 2001. 12691

Stephens, G. L., Vane, D. G., Boain, R. J., et al.: The CloudSat mission and the A-Train: A new dimension of space-based observations of clouds and precipitation, *B. Am. Meteorol. Soc.*, 83, 1771–1790, 2002. 12681

Stephens, G. L., Vane, D. G., Tanelli, S., et al.: CloudSat mission: Performance and early science after the first year of operations, *J. Geophys. Res.*, 113, D00A18, doi:10.1029/2008JD009982, 2008. 12681

Thornton, J. A., Wooldridge, P. J., and Cohen, R. C.: Atmospheric NO₂: In situ laser-induced fluorescence detection at parts per trillion mixing ratios. *Anal. Chem*, 72, 528–539, 2000. 12678

Satellite-derived radiative impact of tropospheric NO₂

A. P. Vasilkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Tsay, S. C., Stamnes, K., and Jayaweera, K.: Radiative energy balance in the cloudy and hazy Arctic, *J. Atmos. Sci.*, 46, 1002–1018, 1989. 12683
- Vandaele, A. C., Hermans, C., Simon, P. C., Carleer, M., Colin, R., Fally, S., McCrieune, M. F., Jenouvrier, A., and Coquart, B.: Measurements of the NO₂ absorption cross-section from 42000 cm⁻¹ to 10000 cm⁻¹ (238–1000 nm) at 220 K and 294 K, *J. Quat. Spectrosc. Ra.*, 59, 171–184, 1998. 12678
- van der A, R. J., Peters, D. H. M. U., Eskes, H. J., Boersma, K. F., Van Roozendael, M., De Smedt, I., and Kelder, H. M.: Detection of the trend and seasonal variation in tropospheric NO₂ over China, *J. Geophys. Res.*, 111, D12317, doi:10.1029/2005JD006594, 2006. 12677
- van der A, R. J., Eskes, H. J., Boersma, K. F., van Noije, T. P. C., Van Roozendael, M., De Smedt, I., Peters, D. H. M. U., and Meijer, E. W.: Trends, seasonal variability and dominant NO_x sources derived from a ten year record of NO₂ measured from space, *J. Geophys. Res.*, 113, D04302, doi:10.1029/2007JD009021, 2008. 12677, 12685
- Vasilkov, A. P., Joiner, J., Spurr, R., Bhartia, P. K., Levelt, P. F., and Stephens, G.: Evaluation of the OMI cloud pressures derived from rotational Raman scattering by comparisons with other satellite data and radiative transfer simulations, *J. Geophys. Res.*, 113, D15S19, doi:10.1029/2007JD008689, 2008. 12681, 12686
- Wenig, M. O., Cede, A. M., Bucsela, E. J., Celarier, E. A., Boersma, K. F., Veeffkind, J. P., Brinksma, E. J., Gleason, J. F., and Herman, J. R.: Validation of OMI tropospheric NO₂ column densities using direct-Sun mode Brewer measurements at NASA Goddard Space Flight Center, *J. Geophys. Res.*, 113, D16S45, doi:10.1029/2007JD008988, 2008. 12684
- Yang, P., Liou, K.-N., Wyser, K., and Mitchell, D.: Parameterization of the scattering and absorption properties of individual ice crystals, *J. Geophys. Res.*, 105, 4699–4718, 2000. 12683
- Ziemke, J. R., Joiner, J., Chandra, S., Bhartia, P. K., Vasilkov, A., Haffner, D. P., Yang, K., Schoeberl, M. R., Froidevaux, L., and Levelt, P. F.: Ozone mixing ratios inside tropical deep convective clouds from OMI satellite measurements, *Atmos. Chem. Phys.*, 9, 573–583, 2009, <http://www.atmos-chem-phys.net/9/573/2009/>. 12681, 12686

**Satellite-derived
radiative impact of
tropospheric NO₂**

A. P. Vasilkov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Satellite-derived
radiative impact of
tropospheric NO₂**

A. P. Vasilkov et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)**Table 1.** Fractional solar flux (F/F_0) and effective NO₂ absorption coefficients (a_{eff}).

Spectral band (nm)	F/F_0	a_{eff} (cm ²)
320–400	0.0591	4.872×10^{-19}
400–700	0.3908	1.992×10^{-19}

Satellite-derived
radiative impact of
tropospheric NO₂

A. P. Vasilkov et al.




Fig. 1. Thick line: spectral optical thickness, τ (left scale) for 0.5 DU of NO₂. Thin line: corresponding approximate absorption, A , of solar irradiance by NO₂ (right scale).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Satellite-derived
radiative impact of
tropospheric NO₂**

A. P. Vasilkov et al.





Fig. 2. NO₂ net atmospheric heating computed with the high spectral resolution (solid line) and the effective (dashed line) absorption coefficient.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Satellite-derived
radiative impact of
tropospheric NO₂**

A. P. Vasilkov et al.

**Fig. 3.** NO₂ net atmospheric heating dependence on surface albedo.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Satellite-derived
radiative impact of
tropospheric NO₂**

A. P. Vasilkov et al.




Fig. 4. NO₂ net atmospheric heating dependence on cloud optical depth, τ , when the NO₂ layer is below the cloud.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Satellite-derived
radiative impact of
tropospheric NO₂

A. P. Vasilkov et al.




Fig. 5. Left: cloudSat/MODIS cloud extinction profiles used in radiative transfer simulations; middle: net heating dependence on the 1 km NO₂ layer midpoint. Dashed(solid) lines are for CloudSat profiles 1(2); vertical lines are for uniform distributions of NO₂ inside the clouds; right: as in middle but for zenith radiances.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Satellite-derived
radiative impact of
tropospheric NO₂**

A. P. Vasilkov et al.




Fig. 6. Monthly-mean tropospheric NO₂ column amounts for January (left) and July (right) 2005 from OMI. White areas correspond to pixels not meeting the cloud fraction criterion.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Satellite-derived
radiative impact of
tropospheric NO₂

A. P. Vasilkov et al.




Fig. 7. Clear-sky daily-averaged NAH (left) and cloud effect in % of clear-sky NAH (right) for January (top) and July (bottom) 2005.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Satellite-derived
radiative impact of
tropospheric NO₂

A. P. Vasilkov et al.




Fig. 8. Clear-sky daily maximum net atmospheric heating for January (left) and July (right) 2005 over the US (top) and Southeast Asia (bottom).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Satellite-derived
radiative impact of
tropospheric NO_2

A. P. Vasilkov et al.




Fig. 9. Diurnal variations in tropospheric NO_2 columns (thin lines) and NAH (thick lines) for the Northeastern US (left) and African regions (right). Diamonds: OMI-derived NO_2 column; solid lines: correspond to simulated diurnal variations in NO_x emissions (see text for details); dashed lines: correspond to constant NO_x emissions; dot-dash line: NAH computed for a constant NO_2 column corresponding to the OMI derived value.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Satellite-derived
radiative impact of
tropospheric NO₂**

A. P. Vasilkov et al.




Fig. 10. Annual average NO₂ columns (left) and corresponding maximum daily NAH for 01 July (right) over Los Angeles computed with a 0.1°×0.1° grid.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)