

This discussion paper is/has been under review for the journal *Atmospheric Chemistry and Physics (ACP)*. Please refer to the corresponding final paper in *ACP* if available.

**SVOCs in
Singapore's coastal
environment**

J. He and
R. Balasubramanian

The exchange of SVOCs across the air-sea interface in Singapore's coastal environment

J. He and R. Balasubramanian

Division of Environmental Science and Engineering, National University of Singapore, Engineering Drive 4, 117576, Singapore

Received: 17 May 2009 – Accepted: 1 June 2009 – Published: 16 June 2009

Correspondence to: R. Balasubramanian (eserbala@nus.edu.sg)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Coastal areas are vulnerable to the accumulation of semi-volatile organic compounds such as PAHs, OCPs and PCBs from atmospheric inputs. Dry particulate and wet depositions, and air-water diffusive exchange in the Singapore's south coastal area, where most of chemical and oil refinery industries are situated in, were estimated. Based on a yearly dataset, the mean annual dry particulate deposition fluxes of \sum_{16} PAHs, \sum_7 OCPs and \sum_{21} PCBs were $1328.8 \pm 961.1 \mu\text{g m}^{-2} \text{y}^{-1}$, $5421.4 \pm 3426.7 \text{ ng m}^{-2} \text{y}^{-1}$ and $811.8 \pm 578.3 \text{ ng m}^{-2} \text{y}^{-1}$, and the wet deposition of \sum_{16} PAHs and \sum_7 OCPs were 6667.1 ± 1745.2 and $115.4 \pm 98.3 \mu\text{g m}^{-2} \text{y}^{-1}$, respectively. Seasonal variation of atmospheric depositions was influenced by meteorological conditions. Air-water gas exchange fluxes had negative values for PAHs, HCHs and DDXs, indicating Singapore's south coast as a sink for the above-mentioned SVOCs. The relative contribution of each depositional process to the total atmospheric input was assessed by annual fluxes. The profile of dry particulate deposition, wet deposition and gas exchange fluxes seemed to be correlated with individual pollutant's properties such as molecular weight and Henry's law constant, etc. For the water column partitioning, the organic carbon-normalized partition coefficients between particulate and dissolved phases (K_{OC}) for both PAHs and OCPs were obtained. The relationships between K_{OC} of PAHs and OCPs and their respective octanol-water partition coefficient (K_{OW}) were examined. In addition, both adsorption onto combustion-derived soot carbon and absorption into natural organic matter for PAHs in marine water column were investigated. Enrichment factors in the sea-surface microlayer (SML) of the particulate phase were 1.2~7.1 and 3.0~4.9 for PAHs and OCPs, and those of dissolved phase were 1.1~4.9 and 1.6~4.2 for PAHs and OCPs, respectively. These enrichment factors are relatively higher than those reported for nearby coastal areas, which are most likely due to more organic surfactants floating in the south coastal surface of Singapore.

ACPD

9, 13235–13269, 2009

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

1 Introduction

Semi-volatile organic compounds (SVOCs) such as polycyclic aromatic hydrocarbons (PAHs), organo-chlorine pesticides (OCPs) and polychlorinated biphenyls (PCBs) can be transported widely from their original sources. These three groups of pollutants, PAHs, OCPs and PCBs, have been studied for decades due to their inherent toxicity, persistence and ability to disperse in the environment (Iwata et al., 1993; Jones and De Voogt, 1999; Oskam et al., 2004). These compounds can be introduced into the environment via the atmosphere, for example, in the emissions of incomplete combustion of fossil fuels (e.g. PAHs), out-gassing from pollutant-containing media such as paints, transformers and capacitors etc. (e.g. PCBs) and by spraying onto soils and vegetation (e.g. OCPs). Major urban and industrial centers increase loadings of SVOCs to proximate sea waters through riverine transport, atmospheric deposition via dry particle deposition, wet deposition, and air-sea gas exchange (McVeety and Hites, 1988; Bidleman et al., 1995; Park et al., 2002; Fang et al., 2008). In addition to acting as sinks for SVOCs, oceans can act as sources of SVOCs to coastal atmospheres and play important roles in the global biogeochemistry of SVOCs (Hinckley et al., 1991; Dachs et al., 1999, 2002). Particle-sorbed SVOCs can settle to the ocean surface by dry particle deposition, a uni-directional advective transport process from the atmosphere to the water, the removal rate by which is a function of the physical and chemical properties of the aerosols and bound pollutants, meteorological conditions and surface characteristics (Wania et al., 1998; Odabasi et al., 1999). In addition, SVOCs are removed from the atmosphere and transported to the waters by precipitation scavenging of atmospheric vapors and particles, which are incorporated into the rain within or below the clouds. After SVOCs are deposited into the bulk seawater, water-column partitioning can affect the distribution of pollutants between the dissolved aqueous and the solid phases and eventually impact the fate of these compounds in oceans (Luo et al., 2004). Other than the abovementioned processes, air-sea exchange can make SVOCs diffuse across the air-sea interface; however, the sea surface microlayer (SML), a unique compartment

ACPD

9, 13235–13269, 2009

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

at the air-sea boundary defined operationally as the upper millimeter (1~1000 μm) of the sea surface, has large storage capacity to delay the transport of SVOCs across the interface. This interfacial effect has been reported as the enrichment of contaminants with different physicochemical properties (Hardy, 1982; Chernyak et al., 1996; Wurl et al., 2006a, c).

Although a considerable number of studies as mentioned above have been conducted to assess the SVOCs exchange between air-sea interface across wide geographical areas, little work has been done to determine the significance of these fate and transfer processes of SVOCs in Southeast Asia. The tropical regions of Asia are considered to be an important source of many SVOCs (Iwata et al., 1994). Those SVOCs with a log K_{OA} of around 8 and a log K_{OW} of around -2 such as PCBs and OCPs can be classified as "multi-hoppers", which can volatilize effectively after deposition to either soils and waters and thus undergo "multiple hops" to migrate from tropical to temperate and even polar regions (Wania, 2003). In Southeast Asia, Basheer et al. (2003) have characterized persistent organic pollutants (PAHs, OCPs and PCBs) in Singapore's northeastern and southwestern coastlines surface seawater, but have not investigated any air-sea exchange processes mentioned above. Wurl et al. (2006a) have studied the distribution of organochlorine compounds (OCs, including OCPs and PCBs) in the SML and subsurface water (SSW) in Hong Kong. Wurl and Obbard (2006b) have also determined the occurrence of OCs in the water column of shallow Johor Strait, between Malaysia and northeastern Singapore, by short-term measurements; in addition, Wurl et al. (2006c) have investigated the role of the SML in air-sea gas exchange of OCs by snapshot sampling in the same sites of Johor Straits. However, the latter work by Wurl et al. (2006c) was just focused on diffusive exchange, one of the air-sea exchange processes only.

In this study, dry particle deposition and wet deposition of selected SVOCs were investigated based on an extensive set of data collected over a year. In addition to receiving riverine inflows and direct emissions, coastal areas tend to experience higher atmospheric deposition than the rest of the oceans due to the proximity to the sources

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

which are predominantly land-based. Singapore, a representative country of South-east Asia, is a small but highly developed island with dense industrial parks in the Southwestern part, where the terrestrial sources affect the surrounding coasts. In this study, Singapore's Southern coastline was chosen during the Northeast monsoon season to evaluate if this coastal area is a sink or source for selected SVOCs via air-sea diffusive exchange as well as to investigate the SML enrichment effect. Furthermore, the water-column partitioning was assessed by a short-term measurement on different aquatic particulate matrices in the ocean.

2 Theoretical approach

Atmospheric sampling was conducted in this study for each event in a consistent manner (24 h at 250 L min⁻¹); weekly concentrations of SVOCs in the atmosphere were computed as arithmetic mean values. In the case of rainwater, the rainfall amount for each event varied. Hence, those concentrations were calculated as volume-weighted mean (VWM) values.

Dry particulate and wet deposition flux can be calculated according to

$$F_{\text{dry}}(\mu\text{g m}^{-2} \text{yr}^{-1}) = \sum_{i=1}^{12} (C_A^i \times V_d \times 2592) \quad (1)$$

$$F_{\text{wet}}(\mu\text{g m}^{-2} \text{yr}^{-1}) = \sum_{i=1}^{12} (C_{\text{VWM}}^i \times P_W^i \times 10^{-3}) \quad (2)$$

where C_A^i , C_{VWM}^i are the monthly particulate concentration of pollutant in air (ng m⁻³) and monthly total (dissolved plus particulate) concentration of pollutant in rain (ng L⁻¹), respectively; V_d is the deposition velocity (ms⁻¹) and P_W^i is the rainfall (mm) for sampling month (i). V_d at a particular vertical height is calculated as the inverse of the sum

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of a number of resistances (Seinfeld and Pandis, 2006) using the following equation.

$$V_d = \frac{1}{R_a + R_b + R_a R_b v_s} + v_s \quad (3)$$

where v_s is the gravitational settling velocity (a function of particle size, density, and viscosity), R_a is aerodynamic resistance refers to turbulent transport from the free atmosphere down to the receptor surface ($R_a = 9/[u\sigma_\theta^2]$, which is a function of wind speed, u and standard deviation of wind speed, σ_θ (Turner, 1986) and R_b is quasi laminar layer resistance, which is a function of particle size, Brownian diffusion, interception and impaction; R_b on water surface is given by van den Berg et al. (2000). Each of R_a and R_b is site specific and determined to a large extent by atmospheric properties. Meteorological parameters such as ambient temperature (K) of 299.6~303.2, wind speed (m s^{-1}) of 1.1~3.6, and standard deviation of wind direction σ_θ (rad) of 0.13~0.78 were observed during the sampling period. In SEA air, a majority of aerosols observed in the size range of 0.1–20 μm by the optical counter were in the fine mode (less than 1 μm) (Kondo et al., 2004).

SVOCs in the water column partition into three phases: the truly dissolved phase, the particulate phase and the colloidal organic material (Totten et al., 2001). The truly dissolved and colloidal phases can be categorized as apparent dissolved phase, which refers to the non-filter retained fraction. The partitioning of SVOCs onto particulate and colloidal phases results in the removal of SVOCs from surface layers via gravitational settling and consequently reduces their volatilization rates (Wania et al., 1998). Based on the empirical estimation of sorption of hydrophobic pollutants onto natural sediments and soils (Karickhoff, 1981) and its application into water-column by Totten et al. (2001), the relationship between partition coefficient K_{OC} and K_{OW} can be derived as follows:

$$\log K_{OC} = \log\left(\frac{K_P}{f_{OC}}\right) = a \log K_{OW} + b \quad (4)$$

where K_{OC} equals partition coefficient K_P normalized by organic carbon fraction (f_{OC}) in particulate phase. K_P refers to the partitioning in the water column between the appar-

SVOCs in
Singapore's coastal
environment

J. He and
R. Balasubramanian

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ent dissolved and particulate phase (Totten et al., 2001), calculated as $C_P/(C_{d,a} \times \text{TSM})$ (C_P : the concentration of SVOCs associated with the particulate phase; $C_{d,a}$: the concentration in the apparent dissolved phase; and TSM: the concentration of total suspended matter). K_{OW} (L kg^{-1}) was obtained from the best available experimental data and predictive models with careful evaluations (de Lima Ribeiro and Ferreira, 2003; Xiao et al., 2004; Shen and Wania, 2005; Zhou and Zhu, 2005; Lü et al., 2007). Since the temperature dependence of K_{OW} for selected SVOCs is not significant (Bahadur et al., 1997; Lei et al., 2000) and the temperature of the seawater varied by no more than 4°C in this study, K_{OW} has not been corrected for temperature.

To accommodate additional sorption to soot phase, Accardi-dey and Gschwend (Accardi-dey and Gschwend, 2002) modified Gustafsson's model (Gustafsson et al., 1997) to yield a dual OC/BC model:

$$K_P = f_{OC}K_{OC} + f_{SC}K_{SC}C_{d,a}^{n-1} \quad (5)$$

where f_{SC} , K_{SC} are soot carbon fraction in particulate phase of seawater and soot-water partition coefficient, respectively, and n is Freundlich constant. If $n=1$, this model assumes that soot-water partition was a linear isotherm, otherwise it is a nonlinear process. This model has been applied to multi media and especially the sediment-water (porewater) partitioning (Gustafsson et al., 1997; Cornelissen and Gustafsson, 2005; Prevedouros et al., 2008). Considering the similarity of distribution between sediment/water and between particulate/water in sea water column, Eq. (5) can be reasonably applied to the latter process.

The apparent dissolved concentrations of SVOCs ($C_{d,a}$) is the sum of the truly dissolved (C_{truly}) and colloidal (C_{DOC} , DOC-dissolved organic carbon, mg L^{-1}) phases, onto both of which the fractions of SVOCs partitioned can be estimated as described in Eq. (6) by Totten et al. (2001) and Wurl et al. (2006a)

$$C_{d,a} = C_{\text{truly}} + C_{\text{DOC}} = C_{\text{truly}}(1 + K_{OW} \times \text{DOC} \times 10^{-7}) \quad (6)$$

A classical two-layer model, which has been previously applied to the air-sea exchange of SVOCs (Achman et al., 1993; Zhang et al., 2007; Li et al., 2009), assumes that the

rate of gas transfer is controlled by the pollutant's ability to diffuse across the air layer and sea surface water on either side of the air-water interface. The molecular diffusivity of the pollutant, dependent on the amount of resistance encountered in the liquid and gas films, describes the rate of transfer while the concentration gradient drives the direction of transfer (Totten et al., 2001). The flux F ($\text{ng m}^{-2} \text{day}^{-1}$) is calculated by

$$F = K_{OL}(C_{\text{truly}} - \frac{C_a}{H'}) \quad (7)$$

where K_{OL} (m day^{-1}) is the overall mass transfer coefficient, and $(C_{\text{truly}} - \frac{C_a}{H'})$ describes the concentration gradient (ng m^{-3}) where C_a is the gas-phase concentration of the compound in air that is divided by the dimensionless Henry's law constant (H) with $H = H/RT$ where R is the universal gas constant ($8.315 \text{ Pa m}^3 \text{ K}^{-1} \text{ mol}^{-1}$), H is Henry's law constant ($\text{Pa m}^3 \text{ mol}^{-1}$) corrected by the temperature and salinity, and T is the temperature at the air-water interface (K). A positive flux indicates net volatilization out of the seawater and a negative flux indicates net absorption into the seawater. The overall mass transfer coefficient (K_{OL}) comprises resistances to mass transfer across the air layer and the water layer.

$$\frac{1}{K_{OL}} = \frac{1}{K_w} + \frac{1}{K_a H'} \quad (8)$$

where K_a and K_w (m day^{-1}) are the mass transfer coefficients across the air layer and water layer, respectively. The mass transfer coefficient $K_{a,\text{comp}}$ (cm s^{-1}) for a compound in air was calculated using the following equation (Schwarzenbach et al., 2003):

$$K_{a,\text{comp}} = K_{a,\text{H}_2\text{O}} \left(\frac{M_{\text{H}_2\text{O}}}{M_{\text{comp}}} \right)^{-0.5 \times 0.67} \quad (9)$$

where $K_{a,\text{H}_2\text{O}} = 0.2 \times u_{10} + 0.3$ and M (g mol^{-1}) is molar mass of the compound. Wanninkhoff's quadratic equation (Wanninkhof, 1992) for the mass transfer coefficient K_w

**SVOCs in
Singapore's coastal
environment**

J. He and
R. Balasubramanian

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



has been applied in a number of studies (Totten et al., 2001; Wanninkhof et al., 2004; Wurl et al., 2006a), but this relationship was a semi-empirical estimation with low correlation ($r^2=0.66$) only at steady winds. A more robust model was thus used in this study to calculate $K_{w,comp}$ (cm s^{-1}) for compounds of interest (Schwarzenbach et al., 2003):

$$K_{w,comp} = \begin{cases} 0.65 \times 10^{-3} \left(\frac{Sc_{comp}}{600} \right)^{-0.67} & \text{for } u_{10} \leq 4.2 \text{ m s}^{-1} \text{ (smooth surface regime)} \\ (0.79 u_{10} - 2.68) \times 10^{-3} \left(\frac{Sc_{comp}}{600} \right)^{-0.5} & \text{for } 4.2 < u_{10} \leq 13 \text{ m s}^{-1} \text{ (rough surface regime)} \\ (1.64 u_{10} - 13.69) \times 10^{-3} \left(\frac{Sc_{comp}}{600} \right)^{-0.5} & \text{for } u_{10} > 13 \text{ m s}^{-1} \text{ (breaking wave regime)} \end{cases} \quad (10)$$

where the Schmidt number, Sc , is the ratio of kinematic viscosity ν_{kin} ($\text{cm}^2 \text{s}^{-1}$) and diffusivity D ($\text{cm}^2 \text{s}^{-1}$) by $Sc = \nu_{kin}/D$. The kinematic viscosity of seawater is obtained using the method recommended by Chen et al. (1973). The diffusivities of compounds of interest in seawater were calculated using the Eq. (10) developed by Wilke and Chang (1955).

$$D = 7.4 \times 10^{-8} \frac{(2.6 M_{comp})^{0.5} T}{\eta V_{comp}^{0.6}} \quad (11)$$

where T is the average temperature during each sampling event, V_{comp} is the molar volume of compound investigated (Mackay et al., 1992, 1996) and η is equal to the kinematic viscosity of solution (here referring to seawater). The important parameters involved in the abovementioned equations are listed in Table 1.

3 Experimental

3.1 Sampling

Dry and wet deposition sampling. For the study of depositional fluxes of selected SVOCs, both the atmospheric and precipitation samples were collected simultaneously at the National University of Singapore (NUS) atmospheric research station.

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Altogether, 37 atmospheric particulate samples and 32 rainwater samples were collected under different weather conditions between June 2007 and May 2008. The sampling site was located 67 m a.s.l. and was approximately 1 km away from the open sea. Details about the sampling instrument and the geography of the area are reported elsewhere (He and Balasubramanian, 2009a).

SSW and SML Sampling. The SSW samples were collected at the depth of 1 m at a southern coastal area of Singapore with a 2 L glass amber bottle inserted in a stainless-steel sampling device. The SML samples (approximately 1 L) were collected simultaneously using the glass plate technique (Harvey and Burzell, 1972) and stored in a amber glass container. Eight samples were collected for SSW and SML, respectively.

Air Sampling. Atmospheric samples were collected simultaneously with SSW by a high volume sampler (Model: TE-1000PUF, Tisch Environmental, Inc., USA) with quartz microfiber filters (Whatman, Tisch Environmental, Inc., USA) to capture particles and polyurethane foam plugs (PUF, TE-1012, Tisch Environmental, Inc., USA) in series (7.62 cm) at ca. 250 L min⁻¹ to collect gaseous phase. Particles were collected by a pre-combusted (at 450°C for 24 h) circular quartz filter of 4" diameter, followed by a combination of 3 PUF plugs connected in series (7.62 cm) to trap gas phase compounds. Eight paired samples (SSW+atmospheric gas) were collected.

Meteorological Conditions. The relevant meteorological parameters (air temperature, rainfall, wind speed, etc.) were obtained from an automated NUS weather station located in the same building where the atmospheric station is located. This meteorological station has been in operation to provide real-time meteorological data to the NUS community.

After collection, all water samples were filtered with pre-cleaned (heated at 450°C for 24 h) Whatman GF/F filters (0.7 μm, 47 mm i.d.). The actual volume of each sample collected was measured and recorded after filtration. Before and after filtration, GF/F filters were conditioned in a dry box (maintained at 22°C and 30% RH), and then weighed to calculate the mass of suspended particles.

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

3.2 Materials

All solvents used for this study were of pesticide grade: n-hexane (HEX), acetone (ACE), dichloromethane (DCM) and methanol (METH) (Tedia, USA). The standard mixture USEPA 610 including 16 priority PAHs (naphthalene (Naph), acenaphthylene (Acy), acenaphthene (Ace), fluorene (Flu), phenanthrene (Phe), anthracene (Ant), fluoranthene (Flt), pyrene (Pyr), benz[a]anthracene (B(a)A), chrysene (Chry), benzo[b]fluoranthene (B(b)F), benzo[k]fluoranthene (B(k)F), benzo[a]pyrene (B(a)P), indeno[1,2,3-cd]pyrene (Ind), dibenz[a, h]anthracene (DB(ah)A) and benzo[ghi]perylene (B(ghi)P)), in methanol: methylene chloride and the standard mixture of OCPs included in Kit 608-S were purchased from Supelco (USA); standard for PCBs (C-QME-01) was obtained from AccuStandard (New Haven, USA). Surrogate standards were used as follows: anthracene-d₁₀, fluoranthene-d₁₀ and benzo[e]pyrene-d₁₂ (Sigma-Aldrich, St. Louis, MO, USA) for PAHs; 13C₁₂-PCB 8 and 13C₆-Hexachlorobenzene (Cambridge Isotope Laboratories, MA, USA) for OCPs; 13C₁₂-PCBs congeners CB 77, 101, 141, 178 (Cambridge Isotope Laboratories, MA, USA) for PCBs. In addition, internal standards were applied as below: phenanthrene-d₁₀, pyrene-d₁₀ and benzo[a]pyrene-d₁₂ (Sigma-Aldrich, St. Louis, MO, USA) for PAHs; 13C₆-Tetrachlorobenzene and 13C₁₂-4,4'-DDT (Cambridge Isotope Laboratories, MA, USA) for OCPs; and 13C₁₂-PCBs congeners CB 8 and 206 (Cambridge Isotope Laboratories, MA, USA) for PCBs. Silica gel and florisil, obtained from Merck (Germany), were heated at 600°C for 24 h to remove any organic contamination.

3.3 Sample preparation and analysis

Sample Extraction and GC-MS Analysis. Prior to extraction, surrogate compounds were added to all samples. Water filters, aerosol filters and PUF samples were extracted separately by Dionex ASE 200. All filtered water samples were extracted on the same day as collection via liquid-liquid extraction using 3×50 mL DCM aliquots in a two liter separatory funnel (EPA method 3510C). Details of extraction for both air and

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



water samples and purification have been reported somewhere (He and Balasubramanian, 2009c). All extracts were finally blown by gentle nitrogen stream and reduced to 50 μL with internal standards, and kept in sealed vials at -20°C prior to GC-MS analysis.

5 Chemical analysis was performed using a QP2010 GC-MS equipped with a Shimadzu AOC-5000 auto injector and a DB-5 fused silica capillary column (30 m length and 0.25 mm I.D.; film thickness 0.25 μm) with purified helium as carrier gas. 4 μL of the sample was injected into the GC-MS in splitless mode with a sampling time of 1.5 min using high-pressure injection mode (pressure 2.47 atm maintained for 2.0 min, initial column flow 3.17 mL min^{-1}) as recommended in the GC-MS manual (Shimadzu).
10 PAHs, OCPs and PCBs standards as well as extracted samples were measured separately in selective ion monitoring mode (SIM) with a detector voltage 700 V. The most abundant ions were selected for quantification, and then one to four reference ions were used for confirmation of each analyte in SIM mode.

15 Measurement of OC and EC. Organic carbon (OC) and elemental carbon (EC) were determined by using a laboratory two-step thermal procedure (He and Balasubramanian, 2009b). Carbon contents were obtained by means of the 2400 series II CHNS/O analyzer (Perkin-Elmer Life and Analytical Sciences Inc.), which was operated in CHN mode with acetanilide (71.09% C, 6.71% H, 10.36% H) as a calibration standard and
20 with helium plus 8% oxygen as carrier gas. When the combustion furnace temperature was set at 450°C , the content of OC in a sample aliquot put in a combustion boat could be determined directly; when the combustion temperature is set as high as 950°C , the total of OC and EC was determined. Prior to analysis, carbonates are removed by adding diluted HCl. The EC content can then be obtained by subtracting OC from the
25 total of OC and EC.

3.4 Quality control

The analytical quality of the data obtained was determined using limits of detection (LOD), recovery, linearity, and by checking sampling artifacts, etc. During each set of

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



extractions, the field blanks (filter and PUF plugs) were included and the mean blank value was subtracted from the measured levels. The reliability of the whole analytical procedure was assessed by using surrogate standards for PAHs, OCPs and PCBs. The mean recoveries obtained per sample were greater than 80% and concentrations of all real-world samples were corrected accordingly. The linearity of calibration standards was calculated by regression analysis with values ranging from 0.99–1.00 (r^2) for SVOCs. Breakthrough was evaluated under field conditions by connecting three one-inch plugs in series and analyzing them separately. For 24 h air samples, measured organic compounds at the third plug were in the range of blank values. Three one-inch plugs used in series with the hi-volume PUF sampler could therefore trap gas-phase target compounds effectively. Control calibration standards spiked with internal standards were measured regularly to check instrument performance during analysis.

4 Results and discussion

4.1 Dry and wet depositions of SVOCs

By Eqs. (1) and (2), both dry and wet depositions were estimated on an annual basis based on the monthly concentrations of SVOCs as shown in Fig. 1. V_d (0.003 m s^{-1}) was not derived from sampling with dry deposition plates in this study but calculated by Eq. (3), which is also in the range of $0.001\text{--}0.01 \text{ m s}^{-1}$ for PAHs, OCPs and PCBs as reported in the literature (Eisenreich et al., 1981; McVeety and Hites, 1988; Hoff et al., 1996). Concentrations of PCBs in both dissolved and particulate phases of all rainwater samples were below the limits of detection in this study.

In Singapore's coastal area, the monthly dry deposition fluxes for PAHs, OCPs and PCBs were in the range of $46.0\text{--}275.6 \mu\text{g m}^{-2} \text{ month}^{-1}$, $60.7\text{--}906.1 \text{ ng m}^{-2} \text{ month}^{-1}$ and $3.1\text{--}93.1 \text{ ng m}^{-2} \text{ month}^{-1}$, respectively. A significant increase in the dry deposition fluxes of most investigated pollutants was observed during the pre-NE monsoon (Oct-Nov 2007) as shown in Fig. 1a, especially for PAHs. In this area, periodical mon-

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

soon winds could assist in dispersing aerosol particles during NE and SW monsoon seasons; the lighter winds during the pre-monsoon seasons (April–May and October–November) might strengthen the accumulation of particulate SVOCs. The monthly wet deposition flux for PAHs and OCPs ranged from 70.0~363.5 $\mu\text{g m}^{-2} \text{month}^{-1}$ and 4.7~39.9 $\mu\text{g m}^{-2} \text{month}^{-1}$, respectively. High wet deposition fluxes were observed from December 2007 to March 2008 for both PAHs and OCPs (Fig. 1a in sharp contrast to the low dry deposition fluxes during the same period (Fig. 1b). The rainy season that took place from December 2007 to March 2008 might have contributed to pronounced wet scavenging of aerosols containing SVOCs from the atmosphere. On the other hand, the peak in the dry deposition flux observed during Oct and Nov 2007 could likely be due to the typical low wind speeds during pre-monsoon seasons and the relatively less rainfall.

The mean annual atmospheric fluxes of selected SVOCs are summarized in Table 2. Particulate dry deposition fluxes of \sum_{16} PAHs, \sum_7 OCPs and \sum_{21} PCBs were 1328.8±961.1 $\mu\text{g m}^{-2} \text{y}^{-1}$, 5421.4±3426.7 $\text{ng m}^{-2} \text{y}^{-1}$ and 811.8±578.3 $\text{ng m}^{-2} \text{y}^{-1}$, respectively. The dry particulate deposition fluxes of \sum_7 OCPs and \sum_{21} PCBs were much lower than that of \sum_{16} -PAHs, most likely due to the local heavy PAH emissions into the atmosphere. In addition, the wet deposition fluxes of \sum_{16} PAHs and \sum_7 OCPs were 6667.1±1745.2 and 87.5±72.1 $\mu\text{g m}^{-2} \text{y}^{-1}$, respectively, which were much more dominant over their dry particulate deposition fluxes. However, it has to be noted that only particles were considered for dry deposition, but both particulate and dissolved phases were included for wet deposition in this study.

The profile of particulate \sum_{16} PAHs fluxes was dominated by B(ghi)P (12.8%), Ind (11.4%), Pyr (10.7%), Phe (10.0%) and Chry (9.3%) on an annual basis; and in wet precipitation, the dominant pollutants were Naph, Phe, Flt and Pyr, accounting for 12.8%, 8.3%, 2.5% and 2.4% of the \sum_{16} PAHs fluxes, respectively. Previous studies indicated that volatile organic compounds could be redistributed onto larger particles via gas-particle partitioning, while the less volatile compounds tend to stay on the particles on which they were emitted (Sahu et al., 2004; He and Balasubramanian, 2009c). One

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

can see in Table 2 that the dry deposition fluxes of high molecular weight (HMW) PAHs are comparable to those of low molecular weight (LMW) PAHs. HMW PAHs have been reported to contribute more to the dry deposition (Gigliotti et al., 2002). This anomaly in the present study may be because the particulate fractions of LMW PAHs in aerosols which prefer to be in coarse-mode readily settled. Table 2 clearly shows that the contribution of the LMW PAH pollutants in wet deposition was increased as compared to those in dry deposition. The main processes contributing to wet deposition fluxes of SVOCs are precipitation scavenging of particle-sorbed and gaseous chemicals, but particle scavenging, rather than the gas scavenging, is the dominant removal mechanism (Offenberg and Baker, 2002; Sahu et al., 2004). The below-cloud scavenging rates increase with particle size because the collision efficiency of particle-droplet encounter increases with increasing particle diameters (Calderón et al., 2008). In this study, it is understandable that the washout of those volatile pollutants attached with coarse particles dominated the wet depositional flux of SVOCs such as LMW PAHs and HCHs, considering the high rainfall amount and precipitation intensities in Singapore. However, scavenging of fine particles needs to be considered and could contribute as well via nucleation, Brownian diffusion, etc (Perry and Hobbs, 1994; Chate, 2005).

4.2 Water column partitioning

In sea subsurface water (SSW), the apparent dissolved and particulate occurrence levels of SVOCs are reported in Table 3. The mean concentrations of PAHs were 43.9 ± 35.8 and $131.4 \pm 101.8 \text{ ng L}^{-1}$ for both phases, respectively, showing a 3-fold enrichment of PAHs in the suspended particulate-phase in this marine environment. For HCHs and DDXs, the total mean concentrations were 1147.8 ± 898.5 and $361.9 \pm 271.8 \text{ pg L}^{-1}$ in the dissolved phase, and 728.9 ± 622.0 and $560.3 \pm 501.5 \text{ pg L}^{-1}$ in the particulate phase, respectively. The levels of both HCHs and DDXs were comparable to those reported for Johor strait located between Singapore and Malaysia previously reported by Wurl et al. (2006b).

4.2.1 Relationship between K_{OC} and K_{OW}

The organic carbon-normalized partition coefficients between particulate and dissolved phases (K_{OC}) for both PAHs and OCPs were obtained based on the data obtained from field measurements. The relationships between K_{OC} of PAHs and OCPs and their respective K_{OW} were investigated with the inclusion of predicted K_{OC} values by Karickhoff's model (Karickhoff, 1981) as in Fig. 2. In Fig. 2a, the observed K_{OC} values, especially for those PAHs with smaller K_{OW} , are much larger than predictions and the log K_{OC} is poorly correlated with log K_{OW} with low slope of 0.13 only. It has been suggested that the slope of the log $K_{OC}/\log K_{OW}$ relation should be equal to 1 when partitioning is in equilibrium (Chiou et al., 1983; Gschwend and Wu, 1985). Researchers initially hypothesized that the significant deviation from 1 may mainly be due to the non-equilibrium or the sorption of compounds to colloids but not particulate phase (Gschwend and Wu, 1985). Till now, a number of studies have reported that the field measured K_{OC} values were higher than the theoretically predicted ones in many different settings such as lake, sediments, and rain water (Baker et al., 1991; Poster and Baker, 1996; Accardi-dey and Gschwend, 2002). These results suggest the presence of particulate phase like soot (black carbon), to which the parent PAHs are more strongly associated than with natural organic matter/carbon. In contrast, field-obtained K_{OC} values for OCPs are correlated better with K_{OW} with a slope of 0.79 closer to theoretical value and greater $R^2=0.66$ (Fig. 2b). In addition, these K_{OC} for OCPs are typically lower, but agree better with the model simulations, indicating the dominant influence of organic-carbon on water column partitioning for OCPs. Similar observations for hydrophobic polychlorinated biphenyls (PCBs) have been widely reported as well (Baker et al., 1991; McGroddy et al., 1996; Poster and Baker, 1996).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

4.2.2 Sorption of PAHs to soot carbon

The observed partition coefficients (K_p) between particulate and dissolved phases are compared with predicted values by Eq. (5) as shown in Fig. 3 for lower molecular weight (LMW) PAHs such as Flu, Phe, Ant, and Pyr and higher molecular weight (HMW) PAHs such as B(a)A, Chry, B(b)F, B(k)F, B(a)P, and B(ghi)P. The use of Freundlich exponent $n=0.62$, which was derived for pyrene in sediment by Accardi-dey and Gschwend (2002), over-estimates the sorption of both LMW (Fig. 3a) and HMW (the predicted line not shown in Fig. 3b) to the particulates in Singapore's coastal sub-surface seawater using the measured f_{OC} and f_{SC} values, indicating the sorption dynamics may be different due to variations in specific surface areas and surface chemistries of soot carbons of various origins. Freundlich exponent n was varied to achieve the best fit at $n=0.80$ for LMW PAHs and at $n=1.2$ for HMW compounds. Both optimized Freundlich exponents are not equal to one, implying that K_{BC} is dependent on sorbate levels and the sorption of PAHs to soot carbon is a nonlinear process. For LMW PAHs such as Phe, Ant, Pyr, $n < 1$ has been reported (Cornelissen and Gustafsson, 2005; Prevedouros et al., 2008), but for HMW compounds such as B(a)P, K_p was over-predicted by a factor of 2~5 by using $n < 1$ (Prevedouros et al., 2008). For the optimum Freundlich exponents, K_p values increased by 1.4~1.6 log units and 0.7~1.3 log units for LMW and HMW PAHs, respectively. Soot carbon was the dominant sorbent here since its sorption contributed ~90% of the total partition coefficients, while the OC fraction was still an important sorbent and contributed to ~10% of the overall K_p values. It is hypothesized that sorptions onto both combustion-derived soot carbon and natural organic matter act in parallel to bind PAHs to particulate phase in marine water column. Additionally, it can be seen that the inclusion of soot carbon sorption reduced the dissolved water concentrations and increase the particulate enrichment of PAHs. The soot carbon fraction would lower the chemical availability of PAHs as compared to that expected from $f_{OC}K_{OC}$ alone so that the bioavailability may be reduced accordingly.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

4.3 Air-water diffusive exchange

4.3.1 Truly dissolved SVOCs

Those SVOCs sorbed onto colloidal matter (DOC) are not available for air-water exchange. The partitioning between truly dissolved and colloidal phases has to be investigated in order to obtain the truly dissolved concentrations of SVOCs. In this study, DOC concentrations are in the range of 0.6~7.8 mg L⁻¹. The predicted mean fractions adsorbed to DOC for individual PAH varied in the range of 0.1~40% as can be seen from Table 3. It clearly showed that HMW PAHs are more readily adsorbed to DOC (10~50%) than those LMW compounds (below 10%). α -, β -, and γ -HCH were adsorbed to DOC at a mean fraction of 0.26, 0.25 and 0.20%, consistent with the values (~0.4%) reported by Wurl et al. (2006b), while $F_{\text{DOC,C}}$ (%) for DDXs was much higher in the range of 30~40%. This distribution pattern suggests that the sorption of SVOCs onto DOC in the seawater is correlated with the chemical's K_{OW} , namely, pollutants with higher K_{OW} are more easily sorbed onto DOC.

4.3.2 Air-water gas exchange flux

Atmospheric gas-phase mean concentrations of Σ PAHs, Σ HCHs and Σ DDXs during November and December 2007 (Table 3) were 187.5±138.3 ng m⁻³, 958.9±336.7 pg m⁻³ and 39.5±7.7 pg m⁻³, respectively, which were in the range of levels reported previously (He et al., 2009). The mean annual fluxes of SVOCs across the air-water interface were estimated and are summarized in Table 2.

The fluxes of the individual PAH compound were in the range of -21.7±20.2~-1608.4±869.6 ng m⁻² day⁻¹ and Σ PAH flux reached up to -5884.0±4040.7 ng m⁻² day⁻¹. All PAHs showed negative fluxes indicating the tendency to transfer these contaminants from air to water (net absorption). The magnitude and direction of PAH fluxes varied widely on temporal and spatial scales all over the world. In Chesapeake Bay, individual fluxes ranged from 14 200 ng m⁻² y⁻¹

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**SVOCs in
Singapore's coastal
environment**J. He and
R. Balasubramanian

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

net volatilization of Flu to $11\,400\text{ ng m}^{-2}\text{ day}^{-1}$ net absorption of Phe on different sampling events (Bamford et al., 1999). Both New York harbor and Raritan Bay systems exhibited net volatilization fluxes for the majority of PAHs (Gigliotti et al., 2002), while LMW PAHs showed positive fluxes and HMW PAHs showed negative values in Mumbai harbor of India (Pandit et al., 2006). The magnitudes of net PAH fluxes are comparable to the reported values ($-0.01\sim-21.2\ \mu\text{g m}^{-2}\text{ day}^{-1}$) in an urban lake in Guangzhou of China (Li et al., 2009). The large net absorption fluxes in this island were likely a result of high gaseous concentrations of PAHs, attributed to the highly uniform ambient temperature and the strong terrestrial sources such as island-wide vehicular traffic, chemical industries, major power plants and oil refineries.

Air-water gas exchange fluxes were estimated in the range of $-44.4\pm 37.8\sim-73.7\pm 31.3\text{ ng m}^{-2}\text{ day}^{-1}$ and $-0.74\pm 0.67\sim-2.9\pm 1.8\text{ ng m}^{-2}\text{ day}^{-1}$ for HCHs and DDXs, respectively (Table 2), indicating Singapore's south coastal line as a sink for both HCHs and DDXs. The fluxes estimated here were comparable to those for the Johor strait located between Singapore and Malaysia with $-8.2\sim-66.8\text{ ng m}^{-2}\text{ day}^{-1}$ (Wurl et al., 2006). It has been reported that air-water gas exchange of SVOCs is a dynamic process that is especially sensitive to fluctuations in concentrations of both air and water phases (Wilkinson et al., 2005). Even though the Henry's law constants of DDXs are larger than those of HCHs, their net absorption was still much lower than that of HCHs, most likely due to much higher occurrence levels of gaseous HCHs, driving more HCHs transferred across the air-sea interface.

Annual air-sea gas exchange fluxes were calculated by multiplying the average daily fluxes by 365 days and the relative contribution of each depositional process to the total atmospheric deposition is shown in Fig. 4. For OCPs, most contribution to total OCP inputs was from wet deposition and air-sea exchange gas fluxes, while the dry particulate deposition dominated the total PAH inputs for Ind and B(ghi)P with $\sim 80\%$ of contribution. As molecular weight increases from the low- to medium-molecular weight PAHs, dry deposition fluxes contribution generally increased, reflecting more fractions on particles. Wet deposition dominated the total PAH inputs for LMW compounds from

Naph to Phe, and the profile of wet deposition contribution to total PAH inputs (Fig. 4) was similar to that of wet deposition fluxes for individual PAH compounds (Table 2). The contribution of air-sea gas absorption has been reported to decrease with the increasing of molecular weight (Gigliotti et al., 2002). However, in this study, it mostly prevailed for medium molecular weight compounds from Ant to B(a)A, but not for LMW PAHs. This may be because the volatile degassing of these LMW compounds from the seawater to air under this tropical climate offsets their transfer from gas to seawater and led to the low net absorption fluxes observed here.

4.4 Sea-surface microlayer enrichment

SML has an enrichment effect of SVOCs, most likely due to their hydrophobic character and great affinity for surfactants collected at the air-water interface (Hardy, 1982; Chernyak et al., 1996; Wurl et al., 2006a). This effect can be quantified by the enrichment factor EF, calculated as the ratio between SML and SSW concentrations (Manodori et al., 2006) as shown in Fig. 5. EFs in the SML of particulate phase were 1.2~7.1 and 3.0~4.9 for PAHs and OCPs, and those of dissolved phase were 1.1~4.9 and 1.6~4.2 for PAHs and OCPs, respectively. EFs in the SML for HCHs found in this study were relatively higher than those reported for the Johor Strait between Malaysia and Singapore (EF=1.1~1.4 for particulate phase and EF=3.3~4.4 for dissolved phase) by Wurl and Obbard (2006b). Similar profiles were found in the SML that EFs generally increased with an increase in molecular weight for both particulate and dissolved phases. In addition, the individual components exhibited similar trends in both phases, with relatively higher enrichments for particulate than for dissolved SVOCs. It is known that the major harbor, one of the busiest harbors in the world, is in the south coastal line of Singapore, and chemical industries as well as oil refineries situated in a group of small islands on the southwest coast of the Singapore Island. It is understandable that more organic film floats in the south coastal surface than in the north area (Johor Strait), resulting in more SVOCs enriched in the SML of the south coastal area. It also indicated that SML really has larger storage capacity (per volume) and may be able

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to delay the transport of SVOCs across the interface to SSW. In addition, the SML is a relatively dynamic compartment, highly influenced by changes of the meteorological and hydrographical conditions (Guitart et al., 2007). This may be the reason for high variation of EFs obtained in this study.

5 Conclusions

Both dry and wet depositions were estimated on the basis of the monthly concentrations of SVOCs. The dry particulate deposition showed lower fluxes during the monsoon seasons but a significant increase within the pre-NE monsoon. It was also observed that the high wet deposition fluxes took place during the rainy season (December 2007~March 2008) in sharp contrast to the low dry deposition fluxes during the same period. It has to be noted that the deposition profile was significantly influenced by meteorological/atmospheric conditions in this study. In SSW, the observed organic carbon-normalized partition coefficients between particulate and dissolved phases (K_{OC}) values for PAHs were much larger than predictions and the log K_{OC} is poorly correlated with log K_{OW} , while the field-obtained K_{OC} values for OCPs are correlated better with K_{OW} . The study of sorption of PAHs to soot carbon showed that soot carbon was the dominant sorbent for PAHs since its sorption contributed ~90% of the total partition coefficients, indicating soot carbon sorption reduced the dissolved water concentrations and possibly lowered the bio-availability of these pollutants in the sea water column. Other than the soot sorbent, colloidal matter (DOC) also played an important role in the partitioning of SVOCs in the water column, and reduced the availability of SVOCs for air-water exchange. In the air-water exchange study, all mean negative gas exchange fluxes showed net absorption of SVOCs onto the Singapore's south coastal sea-surface. The contribution of this process mostly prevailed for medium molecular weight compounds from Ant to B(a)A, but not for LMW PAHs as compared to dry particulate and wet depositions. Lastly, in the investigation of SML enrichment, a similar increasing trend of EFs for both particulate and dissolved phases

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

was found with an increase in molecular weight and the individual components had relatively higher enrichments for particulate than for dissolved SVOCs. In addition, this study also implied that SML may have the large storage capacity to delay the transport of SVOCs across the interface to SSW.

- 5 *Acknowledgements.* The authors gratefully acknowledge the financial support of this research project by the National University of Singapore.

References

- Accardi-dey, A. and Gschwend, P. M.: Assessing the combined roles of natural organic matter and black carbon as sorbents in sediments, *Environ. Sci. Technol.*, 36, 21–29, 2002.
- 10 Achman, D. R., Hornbuckle, K. C., and Eisenreich, S. J.: Volatilization of polychlorinated biphenyls from Green Bay, Lake Michigan, *Environ. Sci. Technol.*, 27, 75–87, 1993.
- Bahadur, N. P., Shiu, W. Y., Boocock, D. G. B., and Mackay, D.: Temperature dependence of octanol-water partition coefficient for selected chlorobenzenes, *J. Chem. Eng. Data*, 42, 685–688, 1997.
- 15 Baker, J. E., Eisenreich, S. J., and Eadie, B. J.: Sediment trap fluxes and benthic recycling of organic carbon, polycyclic aromatic hydrocarbons, and polychlorobiphenyl congeners in Lake Superior, *Environ. Sci. Technol.*, 25, 500–509, 1991.
- Bamford, H. A., Offenberg, J. H., Larsen, R. K., Ko, F. C., and Baker, J. E.: Diffusive exchange of polychclic aromatic hydrocarbons across the air-water interface of the Patapsco river, an urbanized subestuary of the Chesapeake Bay, *Environ. Sci. Technol.*, 33, 2138–2144, 1999.
- 20 Basheer, C., Obbard, J. P., and Lee, H. K.: Persistent organic pollutants in Singapore's coastal marine environment: part I, seawater, *Water Air Soil Poll.*, 149, 295–313, 2003.
- Bidleman, T. F., Jantunen, L. M., Falconer, R. L., Barrie, L. A., and Fellin, P.: Decline of hexachlorocyclohexane in the Arctic atmosphere and reversal of air-sea gas exchange, *Geophys. Res. Lett.*, 22, 219–222, 1995.
- 25 Calderón, S. M., Poor, N. D., Campbell, S. W., and Hartsell, B.: Rainfall scavenging coefficients for atmospheric nitric acid and nitrate in a subtropical coastal environment, *Atmos. Environ.*, 42, 7757–7767, 2008.

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Chate, D. M.: Study of scavenging of submicron-sized aerosol particles by thunderstorm rain events, *Atmos. Environ.*, 39, 6608–6619, 2005.
- Chen, S. F., Chan, R. C., Read, S. M., and Bromley, L. A.: Viscosity of sea water solutions, *Desalination*, 13, 37–51, 1973.
- 5 Chernyak, S. M., Rice, C. P., and McConnell, L. L.: Evidence of currently-used pesticides in air, ice, fog, seawater and surface microlayer in the Bering and Chukchi Seas, *Mar. Pollut. Bull.*, 32, 410–419, 1996.
- Chiou, C. T., Porter, P. E., and Schmedding, D. W.: Partition equilibria of non-ionic organic-compounds between soil organic-matter and water, *Environ. Sci. Technol.*, 17, 227–231, 1983.
- 10 Cornelissen, G. and Gustafsson, Ö.: Prediction of large variation in biota to sediment accumulation factors due to concentration-dependent black carbon adsorption of planar hydrophobic organic compounds, *Environ. Toxicol. Chem.*, 24, 495–498, 2005.
- Dachs, J., Lohmann, R., Ockenden, W. A., Mejanelle, L., Eisenreich, S. J., and Jones, K. C.: Oceanic biogeochemical controls on global dynamics of persistent organic pollutants, *Environ. Sci. Technol.*, 36, 4229–4237, 2002.
- 15 Dachs, J., Van Ry, D. A., and Eisenreich, S. J.: Occurrence of estrogenic nonylphenols in the urban and coastal atmosphere of the lower Hudson River estuary, *Environ. Sci. Technol.*, 33, 2676–2679, 1999.
- 20 de Lima Ribeiro, F. A. and Ferreira, M. M. C.: QSPR models of boiling point, octanol-water partition coefficient and retention time index of polycyclic aromatic hydrocarbons, *J. Mol. Struct.*, 663, 109–126, 2003.
- Eisenreich, S. J., Looney, B. B., and Thornton, J. D.: Airborne organic contaminants in the Great Lakes ecosystem, *Environ. Sci. Technol.*, 15, 30–38, 1981.
- 25 Fang, M., Ko, F., Baker, J. E., and Lee, C.: Seasonality of diffusive exchange of polychlorinated biphenyls and hexachlorobenzene across the air-sea interface of Kaohsiung Harbor, Taiwan, *Sci. Total Environ.*, 407, 548–565, 2008.
- Gigliotti, C. L., Brunciak, P. A., Dachs, J., Glenn, T. R., and Nelson, E. D.: Air-water exchange of polycyclic aromatic hydrocarbons in the New York – New Jersey, USA, harbor estuary, *Environ. Toxicol. Chem.*, 21, 235–244, 2002.
- 30 Gschwend, P. M. and Wu, S. C.: On the constancy of sediment water partition-coefficients of hydrophobic organic pollutants, *Environ. Sci. Technol.*, 19, 90–96, 1985.
- Guitart, C., García-Flor, N., Bayona, J. M., and Albaigés, J.: Occurrence and fate of polycyclic

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

aromatic hydrocarbons in the coastal surface microlayer, *Mar. Pollut. Bull.*, 54, 186–194, 2007.

Gustafsson, Ö., Haghseta, F., Chan, C. H., MacFarlane, J., and Gschwend, P. M.: Quantification of the dilute sedimentary soot phase: Implications for PAH speciation and bioavailability, *Environ. Sci. Technol.*, 31, 3341–3347, 1997.

Hardy, J. T.: The sea surface microlayer: biology, chemistry and anthropogenic enrichment, *Prog. Oceanogr.*, 11, 307–328, 1982.

Harvey, G. W. and Burzell, L. A.: A simple microlayer method for small samples, *Limnol. Oceanogr.*, 17, 156–157, 1972.

He, J. and Balasubramanian, R.: Rain-aerosol coupling in the tropical atmosphere of Southeast Asia: distribution and scavenging ratios of major ionic species, *J. Atmos. Chem.*, 60, 205–220, 2009a.

He, J. and Balasubramanian, R.: A study of gas/particle partitioning of SVOCs in the tropical atmosphere of Southeast Asia, *Atmos. Environ.*, in press, 2009b.

He, J. and Balasubramanian, R.: A study of precipitation scavenging of semi-volatile organic compounds (SVOCs) in a tropical area, *J. Geophys. Res.*, doi:10.1029/2008JD011685, in press, 2009c.

He, J., Balasubramanian, R., Karthikeyan, S., and Joshi, U. M.: Determination of semi-volatile organochlorine compounds in the atmosphere of Singapore using accelerated solvent extraction, *Chemosphere*, 75(5), 640–648, 2009.

Hinckley, D. A., Bidleman, T. F., and Rice, C. P.: Atmospheric organochlorine pollutants and air-sea exchange of hexachlorocyclohexane in the Bering and Chukchi Seas, *J. Geophys. Res.*, 96, 7201–7213, 1991.

Hoff, R. M., Strachan, W. M. J., Sweet, C. W., Chan, C. H., Shackleton, M., Bidleman, T. F., Brice, K. A., Burniston, D. A., Cussion, S., Gatz, D. F., Harlin, K., and Schroeder, W. H.: Atmospheric deposition of toxic chemicals to the Great Lakes: a review of data through 1994, *Atmos. Environ.*, 30, 3505–3527, 1996.

Iwata, H., Tanabe, S., Sakai, N., Nishimura, A., and Tatsukawa, R.: Geographical distribution of persistent organochlorines in air, water and sediments from Asia and Oceania, and their implications for global redistribution from lower latitudes, *Environ. Pollut.*, 85, 15–33, 1994.

Iwata, H., Tanabe, S., Sakai, N., and Tatsukawa, R.: Distribution of persistent organochlorines in the oceanic air and surface seawater and the role of ocean on their global transport and fate, *Environ. Sci. Technol.*, 27, 1080–1098, 1993.

**SVOCs in
Singapore's coastal
environment**

J. He and
R. Balasubramanian

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Jones, K. C. and De Voogt, P.: Persistent organic pollutants (POPs): state of science, *Environ. Pollut.*, 100, 209–221, 1999.
- Karickhoff, S. W.: Semi-empirical estimation of sorption of hydrophobic pollutants on natural sediments and soils, *Chemosphere*, 10, 833–846, 1981.
- 5 Kondo, Y., Morino, Y., Takegawa, N., et al.: Impacts of biomass burning in Southeast Asia on ozone and reactive nitrogen over the western Pacific in spring, *J. Geophys. Res.*, 109, D15S12, doi:10.1029/2003JD004203, 2004.
- Lü, W., Chen, Y., Liu, M., Chen, X., and Hu, Z.: OSPR prediction of n-octanol/water partition coefficient for polychlorinated biphenyls, *Chemosphere*, 69, 469–478, 2007.
- 10 Lei, Y. D., Wania, F., Shiu, W. Y., and Boocock, D. G. B.: HPLC-based method for estimating the temperature dependence of n-octanol-water partition coefficient, *J. Chem. Eng. Data*, 45, 738–742, 2000.
- Li, J., Cheng, H., Zhang, G., Qi, S., and Li, X.: Polycyclic aromatic hydrocarbon (PAH) deposition to and exchange at the air-water interface of Luhu, an urban lake in Guangzhou, China, *Environ. Pollut.*, 157, 273–279, 2009.
- 15 Luo, X., Mai, B., Yang, Q., Fu, J., Sheng, G., and Wang, Z.: Polycyclic aromatic hydrocarbons (PAHs) and organochlorine pesticides in water columns from the Pearl River and the Macao harbor in the Pearl River Delta in South China, *Mar. Pollut. Bull.*, 48, 1102–1115, 2004.
- Mackay, D., Shiu, W. Y., and Ma, K. C. (Eds.): *Illustrated Handbook of Physical-Chemical Properties and Environmental Fate for Organic Chemicals*, Lewis Publisher, Boca Raton, FL, 1992.
- 20 Mackay, D., Shiu, W. Y., and Ma, K. C. (Eds.): *Illustrated Handbook of Physical-Chemical Properties and Environmental Fate for Organic Chemicals* Lewis Publishers, Boca Raton, FL, 1996.
- 25 Manodori, L., Gambaro, A., Piazza, R., Ferrari, S., Stortini, A. M., Moret, I., and Capodaglio, G.: PCBs and PAHs in sea-surface microlayer and sub-surface water samples of the Venice Lagoon (Italy), *Mar. Pollut. Bull.*, 52, 184–192, 2006.
- McGroddy, S. E., Farrington, J. W., and Gschwend, P. M.: Comparison of the in situ and desorption sediment-water partitioning of polycyclic aromatic hydrocarbons and polychlorinated biphenyls, *Environ. Sci. Technol.*, 30, 172–177, 1996.
- 30 McVeety, B. D. and Hites, R. A.: Atmospheric deposition of polycyclic aromatic hydrocarbons to water surfaces: a mass balance approach, *Atmos. Environ.*, 22, 511–536, 1988.
- Odabasi, M., Sofuoglu, A., Tasdemir, Y., and Holsen, T. M.: Measurement of dry deposition

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- and air-water exchange of polycyclic aromatic hydrocarbons with the water surface sampler, *Environ. Sci. Technol.*, 33, 426–434, 1999.
- Offenberg, J. H. and Baker, J. E.: Precipitation scavenging of polychlorinated biphenyls and polycyclic aromatic hydrocarbons along an urban to over-water transect, *Environ. Sci. Technol.*, 36, 3763–3771, 2002.
- Oskam, I. C., Ropstad, E., Lie, E., Derocher, A. E., Wiig, Ø., Dahl, E., Larsen, S., and Skaare, J. U.: Organochlorines affect the steroid hormone cortisol in free-ranging polar bears (*Ursus maritimus*) at Svalbard, Norway, *J. Toxicol. Env. Health*, 67, 959–977, 2004.
- Pandit, G. G., Sahu, S. K., Puranik, V. D., and Venkat Raj, V.: Exchange of polycyclic aromatic hydrocarbons across the air-water interface at the creek adjoining Mumbai harbour, India, *Environ. Int.*, 32, 259–264, 2006.
- Park, J. S., Wade, T. L., and Sweet, S. T.: Atmospheric deposition of PAHs, PCBs and organochlorine pesticides to Corpus Christi Bay, Texas, *Atmos. Environ.*, 36, 1707–1720, 2002.
- Perry, K. D. and Hobbs, F. V.: Further evidence for particle nucleation in clear air adjacent to marine cumulus clouds, *J. Geophys. Res.*, 99, 22 803–22 818, 1994.
- Poster, D. L. and Baker, J. E.: Influence of submicron particles on hydrophobic organic contaminants and distributions of polycyclic aromatic hydrocarbons and polychlorinated biphenyls in rain water, *Environ. Sci. Technol.*, 30, 341–348, 1996.
- Prevedouros, K., Palm-Cousins, A., Gustafsson, Ö., and Cousins, I. T.: Development of a black carbon-inclusive multi-media model: Application for PAHs in Stockholm, *Chemosphere*, 70, 607–615, 2008.
- Sahu, S. K., Pandit, G. G., and Sadasivan, S.: Precipitation scavenging of polycyclic aromatic hydrocarbons in Mumbai, India, *Sci. Total Environ.*, 318, 245–249, 2004.
- Schwarzenbach, R. P., Geschwend, P. M., and Imboden, D. M. (Eds.): *Environmental organic chemistry*, Wiley-Interscience, New York, 2003.
- Seinfeld, J. H. and Pandis, S. N. (Eds.): *Atmospheric Chemistry and Physics: from Air Pollution to Climate Change*, John Wiley, New York, 2nd edn., 2006.
- Shen, L. and Wania, F.: Compilation, evaluation, and selection of physical-chemical property data for organochlorine pesticides, *J. Chem. Eng. Data*, 50, 742–768, 2005.
- Totten, L. A., Brunciak, P. A., Gigliotti, C. L., Dachs, J., Glenn, T. R., Nelson, E. D., and Eisenreich, S. J.: Dynamic air-water exchange of polychlorinated biphenyls in the New York – New Jersey harbor estuary, *Environ. Sci. Technol.*, 35, 3834–3840, 2001.

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Turner, D. B.: Comparison of Three Methods for Calculating the Standard Deviation of the Wind Direction, *J. Clim. Appl. Meteorol.*, 25, 703–707, 1986.
- van den Berg, A., Dentener, F., and Lelieveld, J.: Modelling the chemistry of the marine boundary layer: Sulfate formation and the role of sea-salt aerosol particles, *J. Geophys. Res.*, 105, 11 671–11 698, 2000.
- Wania, F.: Assessing the potential of persistent organic chemicals for long-range transport and accumulation in Polar regions, *Environ. Sci. Technol.*, 37, 1344–1351, 2003.
- Wania, F., Axelman, J., and Broman, D.: A review of processes involved in the exchange of persistent organic pollutants across the air-sea interface, *Environ. Pollut.*, 102, 3–23, 1998.
- Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean, *J. Geophys. Res.*, 97, 7373–7382, 1992.
- Wanninkhof, R., Sullivan, K. F., and Top, Z.: Air-sea gas transfer in the Southern Ocean, *J. Geophys. Res.*, 109, C08S19.01–C08S19.12, 2004.
- Wilke, C. R. and Chang, P.: Correlation of diffusion coefficients in dilute solutions, *AIChE J.*, 1, 264–270, 1955.
- Wilkinson, A. C., Kimpe, L. E., and Blais, J. M.: Air-water gas exchange of chlorinated pesticides in four lakes spanning, *Environ. Toxicol. Chem.*, 24, 61–69, 2005.
- Wurl, O., Obbard, J. P., and Lam, P. K. S.: Distribution of organochlorines in the dissolved and suspended phase of the sea-surface microlayer and seawater in Hong Kong, China, *Mar. Pollut. Bull.*, 52, 768–777, 2006a.
- Wurl, O. and obbard, J. P.: Distribution of organochlorine compounds in the sea-surface microlayer, water column and sediment of Singapore's coastal environment, *Chemosphere*, 62, 1105–1115, 2006b.
- Wurl, O., Karuppiyah, S., and Obbard, J. P.: The role of the sea-surface microlayer in the air-sea gas exchange of organochlorine compounds, *Sci. Total Environ.*, 369, 333–343, 2006c.
- Xiao, H., Li, N., and Wania, F.: Compilation, evaluation, and selection of physical-chemical property data for α -, β -, and γ -hexachlorocyclohexane, *J. Chem. Eng. Data*, 49, 173–185, 2004.
- Zhang, G., Li, J., Cheng, H., Li, X., Xu, W., and Jones, K. C.: Distribution of organochlorine pesticides in the Northern South China Sea: Implications for land outflow and air-sea exchange, *Environ. Sci. Technol.*, 41, 3884–3890, 2007.
- Zhou, W. and Zhu, L.: Distribution of polycyclic aromatic hydrocarbons in soil-water system containing a nonionic surfactant, *Chemosphere*, 60, 1237–1245, 2005.

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Table 1. Relevant parameters used in this study (source indicated in the text).

Compound	$\log K_{OW}$	H ($\times 10^{-4}$)	$K_{a,comp}$ (cm s^{-1})	D ($\times 10^{-5} \text{ cm}^2 \text{ s}^{-1}$)	Sc_{comp} ($\times 10^{-4}$)	V_{comp} ($\text{cm}^3 \text{ mol}^{-1}$)	$K_{w,comp}$ ($\times 10^{-5} \text{ cm s}^{-1}$)
Naph	3.37	1.28~1.48	0.99~1.96	1.95~2.00	4.46~4.72	169.8	3.49~3.62
Ace	3.92	79.5~88.8	1.06~2.08	2.12~2.17	4.12~4.36	173.0	3.68~3.82
Flu	4.18	45.4~50.4	1.09~2.13	2.09~2.14	4.17~4.42	188.0	3.65~3.79
Phe	4.57	19.7~21.7	1.11~2.19	2.09~2.14	4.17~4.41	199.0	3.65~3.79
Ant	4.54	23.9~26.5	1.11~2.19	2.10~2.16	4.14~4.39	197.0	3.67~3.81
Flt	5.22	8.17~8.84	1.16~2.28	2.11~2.17	4.12~4.37	217.3	3.68~3.82
Pyr	5.18	8.12~8.90	1.16~2.28	2.13~2.19	4.09~4.33	214.0	3.70~3.84
B(a)A	5.91	5.42~6.20	1.21~2.37	2.07~2.13	4.20~4.45	248.0	3.63~3.77
Chry	5.86	2.58~3.20	1.21~2.37	2.06~2.11	4.23~4.48	251.0	3.61~3.75
B(b)F	5.75	0.29~0.32	1.25~2.46	2.08~2.13	4.19~4.44	268.9	3.63~3.78
B(k)F	6.00	0.26~0.29	1.25~2.46	2.08~2.13	4.19~4.44	268.9	3.63~3.78
B(a)P	6.04	0.20~0.22	1.25~2.46	2.11~2.16	4.14~4.38	263.0	3.67~3.81
Ind	6.54	0.15~0.16	1.29~2.53	2.14~2.19	4.08~4.32	277.0 ^a	3.70~3.85
B(ghi)P	6.50	0.14~0.15	1.29~2.53	2.14~2.19	4.08~4.32	277.0	3.70~3.85
α -HCH	3.82	1.47~1.61	1.31~2.57	2.37~2.43	3.68~3.90	243.6	3.97~4.12
β -HCH	3.80	0.17~0.19	1.31~2.57	2.37~2.43	3.68~3.90	243.6	3.97~4.12
γ -HCH	3.72	1.10~1.15	1.31~2.57	2.37~2.43	3.68~3.90	243.6	3.97~4.12
4, 4'-DDD	6.33	4.80~5.22	1.35~2.66	2.14~2.19	4.08~4.32	312.6	3.70~3.85
4, 4'-DDE	6.93	29.0~33.1	1.35~2.65	2.16~2.22	4.03~4.27	305.2	3.73~3.88
4, 4'-DDT	6.39	5.05~5.73	1.40~2.75	2.16~2.22	4.03~4.26	333.5	3.73~3.88

^a The molar volume of Ind is not available yet to the best of our knowledge. Due to its similarity of molecular structure and weight to B(ghi)P, it is reasonably presumed that the molar volume of B(ghi)P could be used for Ind.

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. Annual mean atmospheric fluxes of selected SVOCs.

Compound	Dry deposition flux (Mean±S.D.)	Wet deposition flux (Mean±S.D.)	Air-sea exchange (Mean±S.D.)
PAHs	$\mu\text{g m}^{-2} \text{y}^{-1}$	$\mu\text{g m}^{-2} \text{y}^{-1}$	$\text{ng m}^{-2} \text{day}^{-1}$
Naph	65.3±56.0	850.7±428.8	-1608.4±869.6
Acy	28.0±13.1	62.9±60.9	-
Ace	32.7±20.5	102.7±106.5	-36.5±8.0
Flu	19.6±10.3	82.6±46.0	-27.3±22.0
Phe	130.6±78.4	554.4±395.6	-457.7±489.7
Ant	19.6±14.0	68.1±91.4	-210.6±53.9
Flt	112.0±88.6	165.6±133.2	-394.5±283.5
Pyr	140.0±13.1	162.2±156.5	-674.4±359.3
B(a)A	62.5±52.3	47.2±39.2	-192.2±137.4
Chry	121.3±102.6	108.3±95.2	-825.1±669.0
B(b)F	102.6±78.4	142.1±107.7	-383.1±131.7
B(k)F	62.5±72.8	12.4±14.3	-405.6±386.8
B(a)P	75.6±89.6	7.7±6.8	-630.0±563.0
Ind	149.3±168.0	11.9±7.8	-21.7±20.2
DB(ah)A	39.2±67.2	27.1±15.9	-
B(ghi)P	168.0±121.3	42.2±39.4	-17.0±6.6
Σ PAHs	1328.8±961.1	6,667.1±1,745.2	-5884.0±4040.7
OCPs	$\text{ng m}^{-2} \text{y}^{-1}$	$\mu\text{g m}^{-2} \text{y}^{-1}$	$\text{ng m}^{-2} \text{day}^{-1}$
α-HCH	1847.6±1136.8	45.5±36.8	-73.7±31.3
β-HCH	625.2±439.8	8.7±8.8	-44.4±37.8
γ-HCH	1390.3±711.5	24.4±19.1	-48.4±30.4
δ-HCH	933.1±765.2	5.8±4.6	-
4,4'-DDD	54.1±74.6	0.45±0.44	-2.9±1.8
4,4'-DDE	41.1±35.5	0.71±0.89	-0.74±0.67
4,4'-DDT	531.9±261.3	1.9±1.4	-1.1±0.85
Σ OCPs	5421.4±3426.7	87.5±72.1	-171.3±102.8
PCBs	$\text{ng m}^{-2} \text{y}^{-1}$		
CB 17+18	58.8±34.5	-	-
CB 28+31	14.9±10.9	-	-
CB 33	9.3±2.8	-	-
CB 52+49	34.5±22.4	-	-
CB 44	31.7±22.4	-	-
CB 74	29.9±23.3	-	-
CB 70+95	17.7±13.1	-	-
CB 101	39.2±46.7	-	-
CB 99	24.3±16.8	-	-
CB 110	34.5±28.0	-	-
CB 149	168.0±105.3	-	-
CB 138+158	121.3±79.3	-	-
CB 171	63.5±9.3	-	-
CB 177	43.9±21.5	-	-
CB 180	51.3±49.5	-	-
CB 183	72.8±9.3	-	-
Σ PCBs	811.8±578.3	-	-

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Concentrations of SVOCs in SSW, SML and atmospheric gas phases during November to December 2007 in Singapore's coastal area.

Compound	SSW			SML		Gas
	$C_{d,a}$	C_p	$F_{\text{DOC,C}} (\%)$	$C_{d,a}$	C_p	C_a
PAHs	ng L ⁻¹			ng L ⁻¹		ng m ⁻³
Naph	18.4±10.6	33.5±23.8	0.10±0.13	19.6±16.1	43.2±22.2	8.7±6.0
Ace	0.58±0.39	3.1±1.8	0.32±0.47	1.4±1.5	2.7±1.6	0.98±0.84
Flu	2.3±3.6	6.1±4.6	0.58±0.44	4.7±3.5	11.0±10.2	3.7±3.2
Phe	6.2±2.0	17.7±8.4	1.4±1.2	16.9±12.1	60.6±20.5	34.8±22.4
Ant	0.38±0.25	1.5±2.6	1.3±0.92	2.1±3.0	6.8±4.4	3.4±1.8
Flt	5.8±9.1	6.5±8.6	5.5±3.2	3.8±2.6	17.2±3.3	51.8±38.7
Pyr	1.1±0.43	3.9±2.5	5.1±5.7	7.4±5.9	19.6±4.5	56.3±44.1
B(a)A	1.3±1.2	9.2±5.9	18.7±14.8	2.5±2.1	20.0±8.5	15.1±25.9
Chry	1.5±1.4	6.8±4.1	17.3±12.0	3.2±2.1	26.5±14.7	8.0±5.9
B(b)F	0.74±0.50	5.6±5.2	14.5±13.2	4.2±2.2	25.1±15.3	1.9±0.96
B(k)F	0.58±0.36	8.4±6.1	21.4±19.1	1.7±1.3	27.9±18.5	0.61±0.56
B(a)P	1.2±0.72	15.7±11.0	22.7±16.3	4.5±4.2	47.6±27.9	0.85±0.74
Ind	0.91±1.6	5.1±8.4	42.5±23.9	3.1±1.5	12.7±8.7	0.026±0.014
DB(ah)A	2.1±2.8	0.97±0.72	51.9±24.7	1.9±1.3	3.0±2.7	0.010±0.007
B(ghi)P	0.87±0.79	7.3±8.1	40.7±23.8	3.8±2.0	26.3±12.5	0.028±0.025
∑PAHs	43.9±35.8	131.4±101.8		80.8±61.3	365.3±209.1	187.5±138.3
OCPs	pg L ⁻¹			pg L ⁻¹		pg m ⁻³
α-HCH	403.3±206.7	230.0±186.9	0.26±0.37	509.7±194.4	465.5±249.9	415.5±135.7
β-HCH	306.5±280.2	326.6±258.9	0.25±0.36	835.3±533.9	506.7±314.4	238.6±143.1
γ-HCH	257.4±181.2	112.7±131.4	0.20±0.30	232.2±240.2	230.5±145.9	195.9±193.6
δ-HCH	180.6±230.4	59.6±44.8	–	310.5±319.5	208.9±219.2	109.1±101.3
∑HCHs	1147.8±898.5	728.9±622.0		1887.8±1288.1	1411.7±929.5	958.9±336.7
4, 4'-DDD	28.6±22.4	8.7±7.9	33.4±22.7	39.3±36.9	26.3±32.5	7.5±5.9
4, 4'-DDE	91.2±50.5	10.3±16.1	40.1±33.9	165.9±50.6	34.6±20.3	3.3±2.4
4, 4'-DDT	242.1±198.9	541.3±477.5	35.9±23.1	290.6±130.2	1179.8±872.4	28.7±11.5
∑DDXs	361.9±271.8	560.3±501.5		493.4±214.1	1240.7±925.1	39.5±7.7

* $C_{d,a}$: concentration in apparent dissolved phase; C_p , concentration in suspended particulate phase in seawater; $F_{\text{DOC,C}}$: the fraction of SVOCs sorbed onto DOC; C_a , concentration in atmospheric gas.

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion


SVOCs in
Singapore's coastal
environmentJ. He and
R. Balasubramanian

Fig. 1. Seasonal variation in both dry and wet depositions of selected SVOCs between June 2007 and May 2008.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


SVOCs in
Singapore's coastal
environmentJ. He and
R. Balasubramanian

Fig. 2. Relationship between $\log K_{OC}$ (measured and predicted) and $\log K_{OW}$.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


SVOCs in
Singapore's coastal
environmentJ. He and
R. Balasubramanian

Fig. 3. Comparison of predicted and observed K_p (a) Flu, Phe, Ant, and Pyr (b) B(a)A, Chry, B(b)F, B(k)F, B(a)P, and B(ghi)P.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian




Fig. 4. The relative importance of dry particulate deposition, wet deposition, and air-sea gas exchange flux to total atmospheric deposition in the Singapore's south coastal area.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

SVOCs in Singapore's coastal environment

J. He and
R. Balasubramanian




Fig. 5. Enrichment factors (EF) of PAHs and OCPs in the sea-surface microlayer of Singapore's coastal line.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion