

This discussion paper is/has been under review for the journal *Atmospheric Chemistry and Physics (ACP)*. Please refer to the corresponding final paper in *ACP* if available.

**Ground-based lidar measurements from
Ny-Ålesund during
ASTAR 2007**

A. Hoffmann et al.

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007: a statistical overview

**A. Hoffmann¹, C. Ritter¹, M. Stock¹, M. Shiobara², A. Lampert¹, M. Maturilli¹,
T. Orgis¹, R. Neuber¹, and A. Herber³**

¹Alfred Wegener Institute for Polar and Marine Research in the Helmholtz Association, Telegrafenberg A 43, 14473 Potsdam, Germany

²National Institute of Polar Research, Kaga 1-9-10, Itabashi-ku, Tokyo, 190-8518 Japan

³Alfred Wegener Institute for Polar and Marine Research in the Helmholtz Association, Bürgermeister-Smidt-Straße 20, 27568 Bremerhaven, Germany

Received: 7 July 2009 – Accepted: 9 July 2009 – Published: 20 July 2009

Correspondence to: A. Hoffmann (anne.hoffmann@awi.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

During the Arctic Study of Tropospheric Aerosol, Clouds and Radiation (ASTAR) in March and April 2007, measurements obtained at the AWIPEV Research station in Ny-Ålesund, Spitsbergen (operated by the Alfred-Wegener-Institute for Polar and Marine Research and the Institut polaire français Paul-Emile Victor), supported the airborne campaign. This included Lidar data from the Koldewey Aerosol Raman Lidar (KARL) and the Micro Pulse Lidar (MPL), located in the atmospheric observatory as well as photometer data and the daily launched radiosonde. The MPL features nearly continuous measurements; the KARL was switched on whenever weather conditions allowed observations (145 h in 61 days). From 1 March to 30 April, 71 meteorological balloon soundings were performed and compared with the corresponding MPL measurements; photometer measurements are available from 18 March. For the KARL data, a statistical overview based on the optical properties backscatter ratio and volume depolarization can be given. The altitudes of the occurrence of the named features (subvisible and visible ice and water as well as mixed-phase clouds, aerosol layers) as well as their dependence on different air mass origins are analyzed. Although the spring 2007 was characterized by rather clean conditions, diverse case studies of cloud and aerosol occurrence during March and April 2007 are presented in more detail, including temporal development and main optical properties as backscatter, depolarization and extinction coefficients. Links between air mass origins and optical properties can be presumed but need further evidence.

1 Introduction

The Arctic is considered to be a sensitive indicator of climate change due to a large number of special interactions and feedback mechanisms (Curry et al., 1996). Especially aerosols and clouds have a significant influence on the solar and terrestrial radiation budget (Shupe and Intrieri, 2004; Quinn et al., 2007). Their evolution as well

ACPD

9, 15453–15510, 2009

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

as sources and sinks respectively formation and dissipation are not yet entirely understood. Therefore, data collection by ground-based and airborne experiments in this remote region is important (Solomon and Qin, 2007).

The ASTAR 2007 campaign was a follow-up of two aircraft campaigns in the Arctic in 2000 (Yamanouchi et al., 2005) and 2004 (Engvall et al., 2008), mainly focused on the measurement of aerosol and cloud properties in the polar troposphere. During ASTAR 2007, two research aircraft operating from the airport of Longyearbyen/Svalbard at 78° North were supported by ground based measurements from the AWIPEV Research station in Ny-Ålesund amongst others. The campaign was conducted from 26 March until 18 April to capture the end of the Arctic Haze season where pollution events are expected to occur in the Arctic (Herber et al., 2002).

Arctic Haze (Quinn et al., 2007) in early spring and forest fires during summer (Stohl et al., 2007) are considered to be the main sources of optically active aerosols entering the Arctic. Although the non-seasalt (nss) sulphate transport into the Arctic in spring time decreased significantly after the early nineties (Quinn et al., 2007), Haze events remain one of the most important air pollution events in this region for at least two reasons: first during snowmelt the Arctic is especially climatologically sensitive due to the surface albedo effect (Stone et al., 2002) and second, the strongly developing east Asian countries (and potentially Russia) will possibly contribute to relevant emissions in the future (Koch and Hansen, 2005) at least in higher tropospheric layers (Stohl, 2006). While our knowledge about the pollution pathways into the Arctic improved significantly in recent years (Law and Stohl, 2007) a realistic aerosol forecast in this remote region is still not possible as aerosol removal processes are insufficiently known and depend critically on the amount of water vapor and atmospheric trace gases. Particles around 0.2 μm size have the longest lifetime (Twomey, 1977) and can remain as long as 30 days in the dry wintertime Arctic troposphere with its inhibited turbulent mixing (Shaw, 1995).

Herber et al. (2002) presented a 9 year time series of aerosol optical depth (AOD) for Ny-Ålesund, Spitsbergen derived by sun and star photometer which showed a con-

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

stant level of aerosol contamination in time, despite of a general reduction of sulfur measured at the near by Zeppelin station (Quinn et al., 2007). However, a continuous time series of lidar observations for an analysis of the vertical distribution, size and phase of occurring aerosol is still missing for Spitsbergen as lidar observations can only be performed at clear sky conditions with thin clouds or aerosols that do not blind the photomultipliers. Additionally, continuous measurements require a manpower that is only available during campaigns like ASTAR.

While the ASTAR 2000 campaign was focused on tropospheric aerosol and radiation, for ASTAR 2004 and 2007 the research field was widened to cloud analysis. So far, Arctic clouds have mainly been investigated in different campaigns in the North American part of the Arctic e.g. during the Mixed-Phase Arctic Cloud Experiment (M-PACE 2004, Verlinde et al., 2007), the First International Satellite Cloud Climatology Project Regional Experiment Arctic Cloud Experiment (FIRE ACE 1998, Curry et al., 2000) or the Surface Heat Budget of the Arctic Ocean Project (SHEBA 1997–1998, Intrieri et al., 2002). As the Spitsbergen area is influenced by the Gulf Stream and hence warmer and characterized by different air mass pathways and a different orography than the same latitudes in the North American part of the Arctic, occurrence, properties and impact of clouds as well as aerosols can be very differing.

The total Arctic cloud cover varies seasonally with low cloud cover during winter (40–80%) and higher values during summer (80–90%) (Curry et al., 1996; Intrieri et al., 2002; Schweiger et al., 1999; Shiobara et al., 2003) as well as regionally e.g. due to local meteorological conditions. Even an annual variation with North Atlantic oscillation (NAO) phase is possible (Intrieri et al., 2002).

Clouds show a variety of different forms and therefore scattering properties. Within the Arctic troposphere one can distinguish ice clouds, water clouds and mixed-phase clouds whereas the latter ones are the most difficult to describe (Morrison et al., 2008). Characteristic features of mixed-phase clouds include the occurrence of a liquid layer on top of the cloud (Shupe et al., 2008) and the frequent existence of clouds within the boundary layer close to temperature inversions (Kahl, 1990).

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The challenges for precise cloud parameterization in the Arctic were recently pointed out by Wyser et al. (2008) in a comparison of 8 current regional climate models. The models were validated against cloud properties obtained during the SHEBA campaign (Curry et al., 1996). Currently, cloud parameterizations distinguish ice and liquid phase of mixed-phase clouds mainly as a function of temperature alone and are in need of improvement of Arctic cloud parametrization (Vavrus and Waliser, 2008). On the other hand measurements have shown that the liquid water content at low temperatures is sometimes higher than expected and that apart from temperature other parameters as cooling rates must be considered as well (Pinto et al., 2001). The influence of Arctic aerosol on climate and radiation budget has been discussed in Treffeisen et al. (2005) and Rinke et al. (2004). Even small and uniformly distributed aerosol significantly altered the atmospheric state. Hence, especially in this relatively pristine environment the precise description of aerosol properties and clouds in climate models is of importance.

The AWIPEV research station in Ny-Ålesund, Spitsbergen, a uniquely well equipped Arctic laboratory, provides long term remote sensing technologies. In this study we combined lidar, sun photometer and balloon sounding measurements to characterize the Arctic atmosphere in March and April 2007. As observational data in the Arctic are sparse and model parameterizations are usually tested against older campaign data, we used the special opportunity to run the KARL lidar (Ritter et al., 2004) between 1 March and 30 April at almost all times when there were no thick clouds, which would have affected the photomultipliers. Removing all data sets where multiple scattering occurred (predominantly at higher cirrus clouds), about 145 h of lidar data were gathered. This allows us to give a statistical overview of the Arctic spring time in the year 2007. Due to the high inter annual variability of the Arctic climate (Eckhardt et al., 2003) this analysis is still only a snapshot, observations of other years might differ substantially.

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2 Meteorological conditions in March and April 2007

In this chapter, an overview on the meteorological data obtained with radiosonde launches and photometer measurements as well as calculations of backward trajectories are presented.

2.1 Balloon soundings

Within the analyzed two months period, 71 weather balloons with Vaisala RS92 radiosondes (Vomel et al., 2007) have been launched at the AWIPEV base (38 in March and 33 in April). The soundings provide profiles of air temperature, relative humidity, wind speed and wind direction usually up to the mid stratosphere. The temperature T is measured using a capacitive wire and can be obtained with a resolution of 0.1 K and a total uncertainty of 0.2 K in the troposphere. The relative humidity RH can be measured between 0 and 100% with an accuracy of 5% and a resolution of 1% (Währn et al., 2004) but is considered to be less reliable at cold temperatures and above the troposphere (Treffeisen et al., 2007).

2.1.1 Temperatures

An overview of the temporal temperature evolution is given in Fig. 1. The temperature on the ground varied between 255 and 275 K. The tropopause was calculated using the World Meteorological Organization (WMO) definition (WMO, 2008) and is marked with red dots in Fig. 1. Its height varied between 7 and 10 km with temperatures of 200 to 225 K. In mid April stratospheric temperatures increased due to the the break-up of the polar vortex.

A frequent Arctic phenomenon are low level temperature inversions, which are complicated to describe inhibiting the mixing of surface-level air with that of the overlying free troposphere and thus playing an important role in the dynamics of the Arctic planetary boundary layer (Kahl, 1990). We analyzed the occurrence of inversions below 6 km

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

altitude using the 71 obtained temperature profiles in the original resolution of minimum 5 s read-out, which equates to a resolution of ca. 25 m. The algorithm added up the temperature difference between two adjacent height steps as long as it was positive. In Fig. 1 temperature inversions of more than 2 K are marked with blue dots. In March they were observed frequently, declining in April. The white dots mark the surface-based temperature inversions above 0.25 K, whose inversion base was below 25 m (Kahl, 1990), lower temperature differences were neglected due to signal noise. 16 out of the 20 surface-based inversions were observed in March, including the 4 events with surface-based inversions above 2 K. By beginning of March, the radiation deficit is reduced by a rapid increase in solar flux. Therefore the surface temperature rises in spring and the thermal stability near the surface decreases. Simultaneously the cloudiness also increases, weakening the inversion through the increased longwave radiative flux to the surface.

2.1.2 Relative humidity

Another quantity measured by the Vaisala sondes is the relative humidity. It can be seen that the relative humidity decreases with height as well (Fig. 2). Below 1 km the relative humidity was above 30%, decreasing towards higher altitudes and no higher values than 70% above 7 km. Also, it showed a very high day-to-day variability.

For comparison with lidar data, we calculated the relative humidity over ice using the Goff Gratch equation (Buck, 1981). It showed higher values of up to 120%, saturation also occurred in the upper troposphere and is often used for comparison with lidar data (see Sect. 4).

2.2 Aerosol optical depth and backward trajectories

The optical depth was measured using a sun photometer type SP1A produced by Dr. Schulz and Partner GmbH. The SP1A has a field of view of 1° and covers a spectral range from 350 nm to 1050 nm in 17 channels. A detailed description of the instru-

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ment and the performed measurements in Ny-Ålesund can be found in Herber et al. (2002). Due to the polar night conditions, first measurements could not be done before 18 March. The obtained aerosol optical depth (AOD) at 500 nm was determined to 0.05 ± 0.02 in March with an increasing tendency in April (0.08 ± 0.03). Compared to the mean values for the period from 1995 to 2008 of 0.1 this is comparably low. These low values already indicate that from the point of view of optical remote sensing instruments the spring 2007 was very clear with a paucity of obvious Arctic Haze events.

To determine the dominating weather patterns during March and April 2007, three-dimensional backward trajectories were calculated using the Pole-Equator-Pole-Tracer (PEP-Tracer) model (Orgis et al., 2009) with wind fields from the European Center for Medium Range Weather Forecast (ECMWF). For each day 5-day backward trajectories at 3 pressure levels (500, 700 and 850 hPa) and 4 starting times per day (00:00, 06:00, 12:00 and 18:00 UTC – coordinated universal time) were calculated using ensemble calculations of ca. 1000 trajectories. The statistical spread of the trajectory ensembles was determined to be ± 400 km. A use of longer trajectories is not reasonable as the cumulative errors in the particle location become very large. Cluster analysis with a total number of 8 clusters was performed to classify the trajectories into transport patterns, analyzing the spacial variance between different trajectories. The number of 8 clusters was found to be optimal for the Ny-Ålesund region by Eneroth et al. (2003), who did a similar analysis for the 10-year period from 1992–2001. For further analysis the trajectories arriving at 700 hPa were considered as they characterize the mid-troposphere and the main patterns were similar for the two other pressure levels. In Fig. 3 the trajectories for the 700 hPa level merged to 8 different clusters are shown. The cluster analysis in (Eneroth et al., 2003) identifies cluster 1, 2 and 4 (transport from Eurasia) to be more frequent in March and April than during the rest of the year (47.1% compared to 39.6%) for the 10-year-period 1992–2001. In our analysis for 2007, the value is with 30.0% even lower than the annual mean (see Table 1).

Comparing the ASTAR time period (26 March to 18 April) with the whole two months,

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

strong differences are apparent. During ASTAR, the clusters 6 and 7, representing winds from the Canadian Arctic and the Beaufort Sea are prevalent (40.2% compared to 31.1% in March and April), while cluster 2, 5 and 8, showing transport from the European continent and the central Arctic are underrepresented (20.8% compared to 49.6% in March and April). This documents the very clean atmospheric conditions during the ASTAR campaign possibly due to less pollution sources in northern Canada (Law and Stohl, 2007).

3 Cloud statistics from lidar measurements

For the cloud statistics, the complete available dataset of the two lidar systems was analyzed. The MPL data (1368 h) were used for a comparison with the temperature and relative humidity profiles obtained with the proximate balloon sounding (Sect. 3.1.1) and in a second analysis to retrieve a general overview in terms of cloud altitudes (Sect. 3.1.2). The KARL data (145 h) were examined to separate between aerosol and cloud events and to analyze trends in depolarization and backscatter values depending on the altitude and corresponding backward trajectory.

For the analysis of the retrieved signal profiles, the Klett algorithm (Klett, 1981) was used. Assuming a constant lidar ratio LR defined as the quotient between the aerosol backscatter coefficient β^{aer} (BC) and the aerosol extinction coefficient α^{aer} , profiles of β^{aer} were calculated. The backscatter ratio (BSR) for a given wavelength λ at range z is defined as

$$\text{BSR}(\lambda, z) = \frac{\beta^{\text{ray}}(\lambda, z) + \beta^{\text{aer}}(\lambda, z)}{\beta^{\text{ray}}(\lambda, z)} \quad (1)$$

where β^{ray} is the molecular Rayleigh backscatter coefficient. Note that aerosol backscattering refers to larger particles which also includes cloud droplets or crystals. If not further specified, BSR always corresponds to the particle/aerosol backscatter ratio in 532 nm (523 nm for the MPL).

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Furthermore, the volume depolarization (VD) can be estimated measuring the backscattered signal in parallel and perpendicular polarization to the polarization of the transmitted beam.

$$VD(\lambda, z) = \frac{P^{\text{perp}}(\lambda, z)}{P^{\text{par}}(\lambda, z)} \quad (2)$$

5 Assuming a background value of 1.4% which occurs due to Rayleigh scattering in the clear atmosphere, the quotient is normalized in the aerosol and cloud free atmosphere in 14 km altitude. The volume depolarization is an indicator for particle shapes and hence state of aggregation (Pal and Carswell, 1973) and can be measured with KARL for the elastic 532 nm channel.

10 Measurement of the backscatter ratio at different wavelengths allow the calculation of the color ratio CR, which is a measure of the particle size.

$$CR(\lambda_1, \lambda_2, z) = \frac{BSR(\lambda_1, z) - 1}{BSR(\lambda_2, z) - 1} \quad \text{with} \quad \lambda_1 > \lambda_2 \quad (3)$$

A color ratio close to unity indicates small particles, while large CR values indicate large particles compared to the wavelength. In this paper CR in the visible (532 nm) refers to $\lambda_1=1064$ nm and $\lambda_2=532$ nm, CR in the UV to $\lambda_1=532$ nm and $\lambda_2=355$ nm, respectively.

The spectral dependence of the AOD τ with the wavelength λ is expressed by the Ångström exponent α

$$\tau(\lambda) \propto \lambda^{-\alpha} \quad (4)$$

20 The Ångström exponent provides additional information on the particle size, the larger the exponent, the smaller the particles.

Multiple scattering causes the lidar return signal from clouds to increase with increasing receiver field of view (FOV) (Eloranta, 1998) and optical thickness of the cloud (Hu et al., 2006; Cho et al., 2008). For the MPL multiple scattering can be neglected as it

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

has no significant influence on the qualitative cloud altitude detection. KARL statistics refer to rather clear conditions, a few data sets with clouds showing multiple scattering (approx. at BSR>40 in 5 km altitude) have been removed.

3.1 MPL statistics

5 The micro pulse lidar (MPL) is a compact, continuously operating lidar system (Welton and Campbell, 2002), that is running on a twenty-four hour operation basis at AWIPEV research station and maintained by the base personnel since June 2003. It uses a Nd:YLF laser ($\lambda=523.5$ nm) and a Schmidt-Cassegrain telescope with 20 cm diameter for laser transmission and receiving. The main parameters of the system are listed in

10 Shiobara et al. (2003). The measured backscatter profiles cover a range of 60 km with a vertical resolution of 30 m and a temporal average of 1 min. The system was working continuously over the whole 2 months period with some data losses (14 days) due to snow on the window, which has to be removed manually. For further analysis the data from 1 March until 30 April was averaged to means of 10 min, background corrected

15 and cut above 21 km. For 10 altitude intervals of 1 km width, the retrieved BSR profiles were analyzed to find cloud structures within the interval using different thresholds for the difference between two adjacent BSR values. For each altitude interval beginning at the surface, values above 0.1 increasing for at least 3 height steps or a single BSR peak difference of minimal 0.2 were needed to detect a cloud. Furthermore, in case of low

20 cloud detection (below 5 km) the detection of clouds above 5 km was only accepted to be true if the signal to noise ratio (SNR) between 5.5 and 10 km was above a threshold value of 15. Therefore the data about the occurrence of clouds in the upper intervals is less reliable. Profiles with snow on the window can also be detected, looking for a strong backscatter peak at an altitude below 300 m.

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3.1.1 Comparison with radiosondes

As the balloons take about 30 min to reach the tropopause, the start time of the radiosonde was used to find the three temporally following MPL profiles. The profiles were averaged and the backscatter ratio was calculated with the Klett algorithm (Klett, 1981).

The following analysis was done for 57 balloon soundings as in the other cases the MPL window was covered with snow. For the analysis shown in Fig. 4 the 57 temperature profiles were averaged over the 10 altitude intervals, for each interval maximum relative humidity was listed. On the x-axis, the relative humidity and temperature are plotted respectively, on the y-axis the altitude interval, color coded is the frequency of occurrence of clouds in the particular interval. One has to be aware of the fact that due to the averaging process, peak values in temperature are suppressed. In addition the movement of the balloons in zonal and meridional direction can not be ignored. The comparison with the humidity data shows that below 7 km, no clouds at a mean relative humidity below 60% were observed. For the humidity over ice, clouds were only detected in altitude intervals with at least 50% relative humidity over ice (not shown). For the actual formation of clouds, the peak values are expected to be higher as our analysis uses averaged data and is only suitable for estimating minimal values. Although clouds mainly occur above 230 K, clouds were also found in higher altitudes at temperatures of 210 K. For the temperature analysis, the averaging error is small compared to the general lack of high cloud detection due to thicker low clouds (Shiobara et al., 2006). Further analyses with the KARL depolarization channel will be done in Sect. 3.2 to estimate the respective cloud phase.

3.1.2 Cloud altitudes and cloud fraction

As described in Lampert et al. (2009) the ASTAR 2007 cloud cover analysis over Ny-Ålesund showed an increasing fraction of low level and boundary layer clouds in the second half of April. Here, this analysis is redone using the algorithm above for the

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



whole set of 10-min profiles. The choice of 1-km-intervals within the troposphere can be justified with the vertical resolution of current climate models (Pfeiffer, 2006), which averages 9 to 10 levels between 850 and 300 hPa. The frequency of cloud occurrence in the different altitude intervals is marked with the dashed green curve in Fig. 5. Profiles with snow on the window (14 days total) are not considered. Low clouds between 1 and 4 km dominate, while there is another peak for the higher clouds between 7 and 8 km. The clear sky fraction, which is the fraction of time where the MPL did not detect any cloud in any altitude interval is estimated to be 33%. To calculate the total occurrence of cloud cover one would have to guess the cloudiness for the time fraction where the window was snow covered. Assuming this cloudiness to be 100% in the worst case and 67% as in the snow-free times in the best case, the total occurrence of clouds can be estimated to be between 67 and 78%. Other measurements on site carried out by Shiobara et al. (2003) showed about 85% cloud cover in March and April 2002. During the SHEBA campaign the cloud fraction in March/April 1998 was detected to be 83% while ISCCP satellite measurements for the Arctic regions in 1998 show lower values of 63% (Intrieri et al., 2002; Curry et al., 1996).

3.2 KARL statistics

The Koldewey Aerosol Raman Lidar (KARL) is a Nd:YAG based Raman lidar. It measures the elastically backscattered light in three wavelengths (355 nm, 532 nm and 1064 nm) as well as the N_2 shifted lines from the 2nd and 3rd harmonic. Additionally, a water vapor channel at 407 nm as well as depolarization at 532 nm are recorded. From the two N_2 shifted lines 387 nm and 607 nm, the extinction coefficient can be determined, making KARL a “3+2” Raman lidar. In the ASTAR 2007 configuration the field of view of the 30 cm mirror was 0.83 mrad. In November 2006 a new laser (Spectra Pro 290-50), which works at 50 Hz and yields more than 10 W at 355 nm and 532 nm and 20 W at 1064 nm, was installed in the lidar, increasing the energy output by a factor of 5 which significantly improved the data quality (Ritter et al., 2008). After beam widening the laser had an effective divergence of 0.5 mrad. With 10 min integration time and

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



60 m height resolution the elastic wavelengths are easily evaluable up to 25 km altitude in daylight conditions.

KARL can only be operated when the backscattered fraction of the light is not too strong in order not to damage the photomultipliers. This inhibits the evaluation of optically thick clouds with high backscatter, especially in the lower altitudes as the dynamic range of lidar return signals is a function of distance z^2 . Therefore, the available dataset is much smaller than the MPL's data set and a statistical evaluation of cloud occurrence in the strict sense is impossible.

However, we analyzed the frequency of occurrence of different atmospheric structures and classified them according to BSR and VD. For the following analysis all 798 10-min profiles were considered. Within this section, all values were calculated as means in the same altitude intervals used for the MPL cloud analysis in Sect. 3.1.2, while the case studies in Sect. 4 are conducted with the 10 min and 60 m resolution. The mean backscatter ratio at 532 nm and depolarization values depending on the altitude interval can be seen in Fig. 5. Less or thinner clouds were observed between 3 and 5 km, the strongest depolarization occurred between 6 and 10 km. This feature of height-increasing depolarization has been found in different studies (Sassen and Benson, 2001) but generally for higher tropospheric temperatures. A quantitative comparison with the MPL data is not possible due to the different data sets and their respective limitations.

Table 2 summarizes the occurrence of observed backscatter ratio at 532 nm for selected height intervals. The median value and the percentile distribution of the backscatter ratio for height intervals in the troposphere are given. The mean value would have been affected by the few strongest cloud cases. As expected, generally the backscatter ratio decreases with altitude. The strongest signals $BSR > 2$, astonishingly, are observed most frequently in a layer of 4–5 km altitude. We rate this as a real feature, because cases with $BSR > 10$ which might have led to a saturation of the photomultipliers in low, but not in higher altitudes, have not been considered in this table. These values are thought as a guideline for possible verification of climate modeling.

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.2.1 Scatterplots

For the analysis of the backscatter ratio BSR and the respective volume depolarization VD, Fig. 6 shows a scatter plot with different symbols for the different altitude intervals. High depolarization with low backscatter values indicating thin cirrus clouds is found between 4 and 8 km, while the opposite indicating water clouds can be found in the lowest 3 intervals. The interval between 0 and 1 km was neglected as the data are not reliable due to an incomplete overlap function. Mean values over the altitude intervals are shown, some water clouds or water layers with little VD and BSR average out and peak values can exceed the shown ones.

In Fig. 7 the characteristics for the two altitude intervals from 2 to 3 and 3 to 4 km depending on the origin of the air masses are plotted. Each 10-min interval was assigned to the temporally closest calculated backward trajectory in 700 hPa (see Sect. 2.2), color coded is the corresponding cluster number. Volume depolarization below 1.8% with increased BSR of more than 2 was only found for cluster 5, 7 and 8 which corresponds to air masses from Western Europe/East Atlantic, the Beaufort Sea and the central Arctic. This might be due to cyclones originating over the Atlantic and bringing warm air masses which leads to less formation of highly depolarizing ice particles. High volume depolarization above 10% with $BSR > 2$ occurred in air masses 3, 4, 6 and 7, originating over the Atlantic, Western Siberia and the Canadian Arctic. Medium VD between 1.8 and 10% was detected in air masses from the Canadian and the Central Arctic as well as from Europe (cluster 2, 6–8). The air masses originating from Europe might contain a considerable amount of aerosols (Stohl, 2006), but reliable correlations need a trajectory analysis over several years with corresponding lidar measurements.

3.2.2 Case classification

Despite overlapping properties of clouds and aerosols, we attempt a classification of different features depending on the measured volume depolarization and backscatter coefficient values (see Table 3). We distinguish clear sky conditions (C1), water clouds

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(C2), high BSR and low VD (C3) as well as the opposite (C4), thin and thick aerosols (C5/6), thin and thick ice clouds with low VD (C7/8) and thin and thick ice clouds with high depolarization (C9/10). Additionally, Fig. 8 gives an overview of the threshold values for the 10 cases. Note that C1 should only contain profiles with VD larger than 1.2 as there is no enhanced BSR from particles which could decrease the VD background value.

This classification is an attempt to distinguish different measured phenomena with a lidar according to their scattering properties. As these phenomena do not show up separately at all times, additional measurements are needed to successfully analyze lidar data, e.g. lidar data in different elastic and inelastic wavelengths, temperature or relative humidity as well as AOD measurements or trajectory analysis. For instance, some ice particles do not show any depolarization due to their special shape and orientation (Hu et al., 2007) and aerosols also can have a wide variety of sizes, shapes and materials which leads to a range of different scattering properties (Dubovik et al., 2002). Especially in the outer regions of the diagram the information is quite valuable, e.g. cases C1, C2, C4, C9, C10. For the central regions, superposition of different phenomena is possible. A layer showing up in area C5 might be aerosol or a water layer with a small fraction of ice particles. These mixed-phase clouds occur frequently in the Arctic (Intrieri et al., 2002).

The statistical results for the different altitude intervals are given in Table 4. Within the analyzed 145 h of data, the fraction where neither enhanced backscatter nor volume depolarization was observed (C1) increases with height as well as the ice cloud fraction (C7–C10) up to 8 km. Water clouds (C2), aerosols (C5/C6) and water clouds with a certain ice cloud fraction (C3–C6) decrease with height. With reference to the total number of detected clouds, pure water clouds (C2 1–7 km) account for 2.8%, mixed-phase clouds and aerosols (C3–C5 1–7 km) account for 70.0% and the ice cloud fraction (C3–C5 7–12 km and C7–C10 1–12 km) contributes to 27.2%. Although our system is not suitable for measuring thick clouds and hence they are not included in the statistics, this corresponds very well with the value of 73% of clouds containing water found by

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Intrieri et al. (2002) during the SHEBA experiment.

Table 5 shows the scattering characteristics according to different flow patterns. As shown in Sect. 2.2, the ASTAR campaign was characterized by air masses mainly coming from the Canadian Arctic, while within the whole two months period, flow from Eurasia was much more frequent. All KARL profiles for the corresponding clustered trajectories between 2 and 4 km altitude were analyzed with regard to their scattering characteristics. For trajectories originating in the Canadian Arctic, the fraction of clear sky measurements was significantly lower than for the Eurasian clusters, where the thin aerosol fraction was significantly increased, as expected in the Arctic Haze season. The ice cloud and water cloud fraction does not differ, while for Canadian air masses C4 is enhanced at the expense of C1, European air masses show opposite characteristics.

4 Case studies with KARL

Within this section case studies supporting our classification scheme are given using the KARL data of 5 particular days with interesting cloud and aerosol structures (cf. Table 3). Two more case studies observed with KARL during ASTAR 2007 are described in Lampert et al. (2009), an optically very thin low level liquid layer on 19 April and a multi layered ice cloud with varying LR on 21 April.

4.1 Low and medium depolarization C2–C6

Layers with comparably low depolarization were observed on 8 March in 10.5 km and 1.5 km, on 13 March and 15 March in 2.8 km and 3.2 km respectively and on 7 April at a temperature inversion in 0.6 km .

The lower layer on 8 March was observed from 14:45 to 16:50 UTC, the upper layer is presented within the same time frame although it persisted longer. Noticeable in the scatter plot Fig. 9 is the very low volume depolarization of below 1.8% for almost the whole altitude range, which, in combination with enhanced backscatter, suggests

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the existence of spherical scatterers. The balloon sounding at 11:00 UTC (see Fig. 10) showed enhanced relative humidity of 70 to 80% within the lower layer and around 50% for the upper layer (the relative humidity over ice was ca. 85%, respectively), which probably increased in the afternoon to enable cloud formation. The color ratio CR, defined as the quotient of backscatter ratios at different frequencies, was determined to be around 1 for the lower layer, which leads to the assumption of rather big particles compared to the wavelengths (Ansmann et al., 2003), indicating a subvisible water cloud layer. This independence of backscatter with wavelength is a typical result for cloud particles larger than $5\ \mu\text{m}$ (van de Hulst, 1981).

Within the higher layer the temperature during the radio sounding was below 210 K, permitting the existence of fluid water droplets. Very low depolarization values for ice particles can be observed for ice plates that are oriented horizontally (Hu et al., 2007; Reichardt et al., 2002). The lowest part of the ice cloud (10.4–10.5 km) showed a slightly higher depolarization (see Fig. 10). A feasible explanation would be a less perfect orientation of the plates than in the layer above. The lidar ratio of the ice cloud could be determined to 10 sr which leads to an optical depth of 0.001. The scatter close to our detection threshold of 1.2×10^{-4} .

The scatter plots for the low cloud layers on 13 and 15 March are shown in Fig. 11 and Fig. 13, respectively. The layer under consideration is marked in green.

On 13 March at 16:00 UTC the formation of a cloud layer could be observed. As can be seen in Fig. 12, at 15:47 UTC very little backscattering with no depolarization between 2.8 and 3.0 km occurred. At 16:00 UTC the layer intensified between 2.7 and 3.0 km while the depolarization increased to about 3% in a lower layer at 2.6 km. During the next 45 min, BC persisted at 3×10^{-5} between 2.6 and 3.0 km while the VD values between 2.2 and 2.6 km increase to above 15%. This cloud obviously evolved as a two-layer structure with spherical water droplets in the top layer and depolarizing ice crystals below. From 15:35 to 16:11 UTC, the aerosol optical depth in the visible, calculated with LR=18 increased gradually from 0.002 to 0.10 within the water layer while it only slightly increased within the ice layer (from 0.002 to 0.01). For the optically thicker

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

water layer, the evolution of the Ångström exponent for backscattering decreased from 1.3 at 15:35 to essentially 0 at 16:11 UTC. Hence, we were able to observe the growth of spherical water droplets in the top layer as have been reported by Pinto (1998) and Shupe et al. (2008).

5 The further temporal evolution shows the formation of a pure ice layer which is studied in further detail in Sect. 4.2.

On 15 March, during the radio sounding at 11:00 UTC 3 distinct layers with completely different characteristics could be observed (see Fig. 14). The two lower layers show similar values for BC (2×10^{-7}) with the middle layer featuring slightly more depolarization. Nevertheless, both layers are assumed to consist of spherical particles since depolarization is below 4%. The main difference between these two layers is the humidity. In Fig. 14 the radiosonde data is compared to the backscatter and depolarization profiles at 11:20 UTC. The relative humidity between 4 and 7 km is not higher than 30% which prohibits the formation of water clouds (see Sect. 3.1.2). Therefore this layer is assumed to consist of aerosol particles and will be further discussed in Sect. 4.3. Within the lower layer however, the relative humidity measured with the temporally close radio sounding was determined to be higher than 80% and a reduction in the temperature gradient just above the cloud in 3.4 to 3.6 km altitude occurred, probably separating two different air masses.

20 On 7 April, simultaneously to the launch of a RS-92 sonde, a short living low level subvisible cloud was detected in a temperature inversion around 600 m altitude. The boundary layer cloud occurred at 255.5 K in a temperature inversion of 1.5 K. The RS 92 measured 82% relative humidity, corresponding to 98% relative humidity over ice. Figure 16 depicts profiles of BC and VD and Fig. 15 the corresponding scatter plot. The low depolarization of some 2% is slightly higher than for a pure water cloud but too low for most ice crystals. As observed earlier, the layer of highest depolarization lies geometrically 100 m below the maximum backscatter. If we assume that some liquid water is mixed into this cloud which formed exactly at saturation then in the cloud the water mixing ratio must have been 22% higher or the temperature 2.35 K lower than

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

measured by the sonde (Fig. 16), which would be an unusually high deviation. As the cloud persisted for about 20 min (see Fig. 17) with a wind speed of 4 m/s this wet patch had an extent of ca. 5 km. Unfortunately, the low altitude of this subvisible cloud results in an incomplete overlap. Hence, it was not possible to calculate its extinction. The presented backscatter and volume depolarization values are ratios of lidar profiles and therefore not so strongly affected by incomplete overlap. Calculating BC for 355 nm and 532 nm by the ratio between the elastic and the N_2 Raman-shifted returns (Ansmann et al., 1992), the color ratio was determined to be unity. Therefore we assume rather large particles, which contrasts to an otherwise similar liquid precondensation layer on 19 April described in Lampert et al. (2009) in this issue.

4.2 High depolarization C7–C10

Highly depolarizing air masses were observed in the upper troposphere on 13 March and 15 March, on 13 March the formation of a low level, depolarizing ice cloud be observed as well (as indicated in Sect. 4.1).

From the start of the measurements on 13 March at 14:00 UTC until about 14:50 UTC an ice cloud structure between 6 and 8 km was observed. Its depolarization is up to 20% while the backscatter values are relatively low (see Fig. 11). With an estimated lidar ratio of 18 sr the optical depth at 14:17 UTC was calculated to 0.035, which is almost subvisible (Lynch et al., 2002).

Contemplating the temporal evolution of the lower layer, already described in Sect. 4.1, one observes an interesting transition between 17:30 and 17:50 UTC. As can be seen in Fig. 12, the backscatter coefficient decreases by a factor of about 4 while the depolarization increases to up to 18%. This can no longer be a water layer and thus, we observed a glaciation process. Supporting this theory, the temperature in this layer, measured during a radio sounding at 10:48 UTC, was about 250 K. Due to the decreasing backscatter coefficient the layer's AOD also decreases to about 0.02 at 17:55 UTC. The accumulated ice cloud layer has similar scattering characteristics as the first observed cirrus layer as can be seen in the scatter plot Fig. 11.

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Depolarization values of up to 12% were obtained for the high cloud layer on 15 March between 7.4 and 8.8 km altitude at 14:06 UTC when the observation started. During the radio sounding at 11:00 UTC the relative humidity over ice in this layer was well above 50%. Within the following one hour the ice cloud layer completely disappeared and the respective AOD decreased from 0.03 to 0.002, hence we observed a very thin, short living cloud structure.

4.3 Aerosols C5–C6

During March 2007, two weak aerosol events (C5) were observed, while in April not even weak aerosol could be definitely identified.

A persistent aerosol layer was observed from 14 March 20:40 UTC to the end of the observation at 15 March 12:40 UTC, due to the low cloud described in Sect. 4.1. The weak layer was observed in around 6 km altitude sinking to 5.3 km, and the corresponding air trajectories suggest an origin of the air masses originate from central Europe (see Fig. 19). It stands out due to its long lifetime, its considerable altitude – the majority of Arctic Haze events occur in the lower free troposphere over Spitsbergen (Scheuer et al., 2003) – its quite high depolarization of 5% and the considerable amount of precipitation of more than 20 mm which occurred in these air masses prior to their arrival. Nevertheless in the scatter plot in Fig. 13 the layer can clearly be distinguished from the cloud layer below. As can be seen in radiosonde data from 15 March 11:00 UTC in Fig. 14, the arriving air masses were with 35% relative humidity far too dry for cloud formation and the derived lidar ratios of 64.3 sr for the 532 nm channel and 39.6 sr for 355 nm seem to be too high for clouds. The maximum AOD of this layer of 0.025 for the UV and 0.01 for the visible occurred around 1:00 UTC at 15 March. As the VD values are still quite low we neglect the particles asphericity and perform an inversion of the lidar data with Mie theory according to Boeckmann (2001). As a result, an index of refraction of $n=1.6-0.011i$, a mono-modal log-normal size distribution of 93 particles per ccm ($\pm 50\%$), effective radius $r_{\text{eff}}=0.199 \mu\text{m}$ ($\pm 50\%$) and a distribution width of $\sigma=1.58$ ($\pm 10\%$) were found. This seems to be an appropriate value for a sulphate soot mixture,

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the main constituents of Arctic Haze (Yamanouchi et al., 2005).

Due to the insecurity of backward trajectories in the Arctic after more than 5 days it seems impossible to ascribe this event to a precise cause or location. From the wind speed of 4.5 m/s in 6 km altitude from the sounding on 15 March, we estimate that even in the unlikely case that the aerosol event stopped directly after the end of our lidar observations, it had a horizontal extent of at least 260 km. On the other hand its vertical extent never exceeded 900 m and over more than 50% in time was as small as 300 m.

Another weak Arctic Haze event occurred in the night from 18 to 19 March 2007. As can be seen from Fig. 20, the backscatter ratio around 3 km altitude is enhanced. At 2.5 km a clear maximum of the extinction can be seen in both N_2 Raman channels which yield to lidar ratios of 80 sr (± 12) for 532 nm and 60 sr (± 10) for 355 nm. These values are higher than the lidar ratios obtained at different Haze events over the site. Based on our (so far still partially unpublished) data the Arctic Haze typically has a lidar ratio of 40–60 sr at 532 nm and 30–50 sr at 355 nm. However, the lidar ratio seems to be almost always about 25% higher at 532 nm. The layer of increased extinction and backscatter coincides with an increase of depolarization from 2% to 2.5% .

The result of the inversion with Mie code gives a refractive index of $n=1.56-0.01i$ and the aerosol can best be described by a one-modal log-normal distribution with 343 particles per ccm ($\pm 30\%$), effective radius $r_{\text{eff}}=0.177 \mu\text{m}$ ($\pm 10\%$) and a distribution width of $\sigma=1.59$ ($\pm 10\%$) which is similar to the 15 March case. The corresponding HYSPLIT trajectories (Draxler and Hess, 1998) are shown in Fig. 21. The trajectories find a relation between European contamination and the arrival of air masses over Ny-Ålesund at 2400 m altitude – 100 m lower than our lidar observations suggest. Given the sparse meteorological data in the high Arctic we consider this mismatch as justifiable. No precipitation occurred out of the trajectory, so that an uptake of aerosol over the European continent seems possible.

On 7 April 2007 an increased AOD was measured by the photometer as summarized in Fig. 18. It shows the aerosol optical depth at 532.8 nm and the Ångström exponent.

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

During the time period observed with lidar, the optical depth grows without any change in the particle size, so no clear change of the air mass is seen, which is supported by the air back-trajectories which came from European Russia (but with heavy precipitation of more than 20 mm according to HYSPLIT). The value of the Ångström exponent is more “aerosol-like” the whole day and does not show any contamination with supermicron particles. The short living cloud at 11:00 UTC described in Sect. 4.1 is invisible in the sun photometer because lidar and photometer are looking in different directions at our polar site. An inversion of the photometer AOD as well as the phase function of scattering derived an index of refraction of 1.4 which is lower than expected for the sulfate-soot mixtures of Arctic Haze (Yamanouchi et al., 2005) which typically shows an refractive index of 1.5 or higher (Ritter et al., 2004). The low index of refraction was approved by an inversion of the lidar data at several times during that day, although, unfortunately the inversion of a size distribution turned out to be numerically unstable. This might at least partly be explained by the weakness of the N_2 Raman channel on that day. Moreover the scatter plot (Fig. 15) does not find any differences in the afternoon as the AOD increases above 0.1 compared to the cleaner morning. The radiosonde measured a slightly increased humidity in this 4 km altitude region and the HYSPLIT back-trajectories showed air masses coming from Russia (see Fig. 22) but with a precipitation of 20 mm. So at least some of the hydrophil aerosol should have been washed out. Therefore Arctic Haze as the cause of this increased optical depth seems to be unlikely. Probably small water droplets and ice crystals, which extend up to 6.4 km altitude in the afternoon (as can be seen in Fig. 17), possibly mixed with an insoluble aerosol component, gave rise to the AOD presented here.

This example clearly shows that “increased optical depth with low Ångström exponent and air mass from Russia” alone is not sufficient to adequately characterize the polar troposphere. Moreover it emphasizes the need for future combined in situ and remote sensing campaigns to fully describe aerosol cloud interactions especially in humid air.

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

5 Conclusions

In this work a statistics of the observations of (thin) clouds and aerosols at a site in the high European Arctic has been presented.

During March and April 2007, temperature and relative humidity were observed performing 71 balloon soundings, temperature inversion layers occurred frequently – surface-based as well as in up to 2.5 km altitude due to the orography of nearby mountains.

A statistics of the altitude dependent cloud cover from the MPL lidar has been presented, showing maximum cloud cover between 2 and 3 km. The overall cloud cover was determined to 73 ($\pm 5\%$) which is a little bit less than other ground based retrievals for the site (Shiobara et al., 2003) and other Arctic locations. From the comparison of MPL data with 54 radiosondes, we found some cases with relative humidity down to 60% sufficient for cloud existence especially in altitudes above 4 km. As never clear super saturation in clouds has been observed, a dry bias of the RS-92 in extreme environments must be considered, similar to the findings of Rowe et al. (2008).

The 145 h lidar data obtained with our the Raman lidar KARL refer to “clear conditions” with only thin cloud structures and aerosol layers. Case studies were performed for 5 different days, featuring the wide variety of Arctic clouds and aerosols. On 8 March a low (1.0–1.7 km), very weak and hence subvisible water cloud layer was observed simultaneously with a very high non-depolarizing ice cloud, presumably consisting of horizontally oriented ice plates. A boundary layer cloud at a temperature inversion was observed on 7 April 2007 while on 13 March 2007 the transformation of a low level mixed phase cloud with a liquid layer on top to a pure ice cloud be observed. Thin high level ice clouds with high volume depolarization as on 13 March and 15 March 2007 occurred frequently. On 15 and 18 March 2007, Arctic Haze layers with a refractive index of $n=1.6-0.01i$ were observed.

Being able to run the KARL whenever possible during March and April 2007, we successfully performed a statistics where we classified the different cloud and aerosol

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

structures according to their scattering properties VD and BSR. 10 case studies for 5 days have been described, leading to the conclusions that clouds with very strong VD and low BSR as well as low VD and strong BSR can be identified to be ice cloud layers or water layers respectively with a high certainty (with the exception of non-depolarizing ice structures). Layers with enhanced BSR and VD values between 1.2 and 5% can refer to aerosols but also to a superposition of different particles. Cloud classification is a powerful tool to have a first glimpse on the lidar data, further analysis however is necessary to fully identify and describe a phenomenon as can be seen for the ice cloud on 8 March and the assumed aerosol case on 7 April.

However, the year 2007 was characterized by a sparseness of aerosol events, as can be seen in the low photometer AOD in March (measurements started after the two haze events) and in the case of 7 April, where an increased AOD seems to be at least partially connected with water vapor and a low refractive index. Therefore, our results from Ny-Ålesund suggest, that the spring 2007 was clearer than average from the point of view of optical remote sensing instruments which allowed a statistical overview of the Arctic spring time in 2007.

These clean conditions during spring 2007 might be related to the fact that only 30% of all backward trajectories during the two months under consideration came from Europe, a lower value than typical. According to our results in Table 5, for our particular data set, air masses from Eurasia correspond to “clear” conditions which means a lack of both increased backscatter and any depolarization of the air. Our case C4 “depolarization without noticeable backscatter” – probably spurious ice crystals or, hypothetically, spurious Asian desert dust remnants which came via the Canadian Arctic towards Spitsbergen, are clearly underrepresented in the European trajectories. As expected, the stronger “aerosol-like” cases are overrepresented in the European air flow regime, but as discussed in the example of 7 April, not all increased AOD with air from Europe is actually Arctic Haze. Hence, comparing backward trajectories with our observed features, simple statements as “airflow from Eurasia especially with little precipitation, produces Arctic Haze” must be rejected.

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A precise description of the necessary conditions for Arctic Haze and cloud formation and its optical effect in the Arctic troposphere is beyond the scope of this paper. Such an intent would require additional long-term ground based data over at least a decade, as well as aircraft campaigns for a comparison with in situ and remote sensing instruments, which include measurements of both the gas phase and the particle content along the possible pollution pathways.

Acknowledgements. We are very thankful to the 2007 AWIPEV base personnel Mareike Peterson, Rainer Vockenroth and Cedric Couret, especially for the daily balloon soundings and data collection. The MPLNET project is funded by the NASA Earth Observing System and Atmospheric Radiation Sciences Program.

References

- Ansmann, A., Riebesell, M., Wandinger, U., Weitkamp, C., Voss, E., Lahmann, W., and Michaelis, W.: Combined Raman elastic-backscatter LIDAR for vertical profiling of moisture, aerosol extinction, backscatter, and LIDAR ratio, *Appl. Phys. B Photo.*, 55, 18–28, 1992. 15472
- Ansmann, A., Bösenberg, J., Chaikovsky, A., Comerón, A., Eckhardt, S., Eixmann, R., Freudenthaler, V., Ginoux, P., Komguem, L., Linné, H., Márquez, M. Á. L., Matthias, V., Mattis, I., Mitev, V., Müller, D., Music, S., Nickovic, S., Pelon, J., Sauvage, L., Sobolewsky, P., Srivastava, M. K., Stohl, A., Torres, O., Vaughan, G., Wandinger, U., and Wiegner, M.: Long-range transport of Saharan dust to northern Europe: The 11–16 October 2001 outbreak observed with EARLINET, *J. Geophys. Res.-Atmos.*, 108, 4783, doi:10.1029/2003JD003757, 2003. 15470
- Boeckmann, C.: Hybrid regularization method for the ill-posed inversion of multiwavelength lidar data in the retrieval of aerosol size distributions, *Appl. Optics* 40, 1329–1342, 2001. 15473
- Buck, A.: New Equations for Computing Vapor Pressure and Enhancement Factor, *J. Appl. Meteorol.*, 20, 1527–1532, 1981. 15459
- Cho, H.-M., Yang, P., Kattawar, G., Nasiri, S., Hu, Y., Minnis, P., Trepte, C., and Winker, D.: Depolarization ratio and attenuated backscatter for nine cloud types: analyses based on

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



collocated CALIPSO lidar and MODIS measurements, *Opt. Express*, 16, 3931, 1848–1866, 2008. 15462

Curry, J., Rossow, W., Randall, D., and Schramm, J.: Overview of Arctic Cloud and Radiation Characteristics, *J. Climate*, 9, 1731–1764, 1996. 15456, 15457, 15465

5 Curry, J., Hobbs, P., King, M., Randall, D., Minnis, P., Isaac, G., Pinto, J., Uttal, T., Bucholtz, A., Cripe, D., Gerber, H., Fairall, C., Garrett, T., Hudson, J., Intrieri, J., Jakob, C., Jensen, T., Lawson, P., Marcotte, D., Nguyen, L., Pilewskie, P., Rangno, A., Rogers, D., Strawbridge, K., Valero, F., Williams, A., and Wylie, D.: FIRE Arctic Clouds Experiment, *B. Am. Meteorol. Soc.*, 81, 5–30, 2000. 15456

10 Draxler, R. and Hess, G.: Description of the HYSPLIT-4 Modeling System, Air Resources Laboratory, 1998. 15474

Dubovik, O., Holben, B., Eck, T., Smirnov, A., Kaufman, Y., King, M., Tanré, D., and Slutsker, I.: Variability of Absorption and Optical Properties of Key Aerosol Types Observed in Worldwide Locations, *J. Atmos. Sci.*, 59, 590–608, 2002. 15468

15 Eckhardt, S., Stohl, A., Beirle, S., Spichtinger, N., James, P., Forster, C., Junker, C., Wagner, T., Platt, U., and Jennings, S. G.: The North Atlantic Oscillation controls air pollution transport to the Arctic, *Atmos. Chem. Phys.*, 3, 1769–1778, 2003, <http://www.atmos-chem-phys.net/3/1769/2003/>. 15457

Eloranta, E. W.: Practical Model for the Calculation of Multiply Scattered Lidar Returns, *Appl. Optics*, 37, 2464–2472, 1998. 15462

20 Eneroth, K., Kjellström, E., and Holmén, K.: A trajectory climatology for Svalbard; investigating how atmospheric flow patterns influence observed tracer concentrations, *Phys. Chem. Earth A/B/C*, 28, 1191–1203, 2003. 15460, 15484

Engvall, A., Krejci, R., Ström, J., Minikin, A., Treffeisen, R., Stohl, A., and Herber, A.: In-situ airborne observations of the microphysical properties of the Arctic tropospheric aerosol during late spring and summer, *Tellus B*, 60, 392–404, 2008. 15455

25 Herber, A., Thomason, L., and Gernandt, H.: Continuous day and night aerosol optical depth observations in the Arctic between 1991 and 1999, *J. Geophys. Res.-Atmos.*, 107, 4097, doi:10.1029/2001JD000536, 2002. 15455, 15460

30 Hu, Y., Liu, Z., Winker, D., Vaughan, M., Noel, V., Bissonnette, L., Roy, G., and McGill, M.: Simple relation between lidar multiple scattering and depolarization for water clouds, *Opt. Lett.*, 31, 1809–1811, 2006. 15462

Hu, Y., Vaughan, M., Liu, Z., Lin, B., Yang, P., Flittner, D., Hunt, B., Kuehn, R., Huang, J.,

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007A. Hoffmann et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Wu, D., Rodier, S., Powell, K., Trepte, C., and Winker, D.: The depolarization – attenuated backscatter relation: CALIPSO lidar measurements vs. theory, *Opt. Express*, 15, 5327–5332, 2007. 15468, 15470

Intrieri, J., Shupe, M., Uttal, T., and McCarty, B.: An annual cycle of Arctic cloud characteristics observed by radar and lidar at SHEBA, *J. Geophys. Res.-Oceans*, 107, 8030, doi:10.1029/2000JC000423, 2002. 15456, 15465, 15468, 15469

Kahl, J.: Characteristics of the low-level temperature inversion along the Alaskan Arctic coast, *Int. J. Climatol.*, 10, 537–548, 1990. 15456, 15458, 15459

Klett, J.: Stable analytical inversion solution for processing lidar returns, *Appl. Optics*, 20, 211–220, 1981. 15461, 15464

Koch, D. and Hansen, J.: Distant origins of Arctic black carbon: A Goddard Institute for Space Studies ModelE experiment, *J. Geophys. Res.-Atmos.*, 110, 4204, doi:10.1029/2004JD005296, 2005. 15455

Lampert, A., Ritter, C., Hoffmann, A., Gayet, J.-F., Mioche, G., Ehrlich, A., Dörnbrack, A., Wendisch, M., and Shiobara, M.: Observations of boundary layer, mixed-phase and multi-layer Arctic clouds with different lidar systems during ASTAR 2007, *Atmos. Chem. Phys. Discuss.*, 9, 15125–15179, 2009, <http://www.atmos-chem-phys-discuss.net/9/15125/2009/>. 15464, 15469, 15472

Law, K. S. and Stohl, A.: Arctic Air Pollution: Origins and Impacts, Science (New York, NY), 315, 1537–1540, 2007. 15455, 15461

Lynch, D. K., Sassen, K., Del Genio, A., Heymsfield, A. J., Minnis, P., Platt, M., Quante, M., Schumann, U., and Sundqvist, H.: *Cirrus: The Future*, Oxford University Press, USA, 2002. 15472

Morrison, H., Pinto, J., Curry, J., and McFarquhar, G.: Sensitivity of modeled arctic mixed-phase stratocumulus to cloud condensation and ice nuclei over regionally varying surface conditions, *J. Geophys. Res.-Atmos.*, 113, 5203, doi:10.1029/2007JD008729, 2008. 15456

Orgis, T., Brand, S., Schwarz, U., Handorf, D., Dethloff, K., and Kurths, J.: Influence of interactive stratospheric chemistry on large-scale air mass exchange in a global circulation model, *Eur. Phys. J. Special Topics*, in press, 174, 2009. 15460

Pal, S. and Carswell, A.: Polarization properties of lidar backscattering from clouds, *Appl. Optics*, 12, 1530–1535, 1973. 15462

Pfeiffer, S.: Modeling cold cloud processes with the regional climate model REMO, Ph.D. thesis, 2006. 15465

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Pinto, J.: Autumnal Mixed-Phase Cloudy Boundary Layers in the Arctic, *J. Atmos. Sci.*, 55, 2016–2038, 1998. 15471
- Pinto, J., Curry, J., and Intrieri, J.: Cloud-aerosol interactions during autumn over Beaufort Sea, *J. Geophys. Res.*, 106, 15077–15098, 2001. 15457
- 5 Quinn, P., Shaw, G., Andrews, E., Dutton, E., Ruoho-Airola, T., and Gong, S.: Arctic haze: current trends and knowledge gaps, *Tellus B*, 59, 99–114, 2007. 15454, 15455, 15456
- Reichardt, J., Reichardt, S., Behrendt, A., and McGee, T.: Correlations among the optical properties of cirrus-cloud particles: Implications for spaceborne remote sensing, *Geophys. Res. Lett.*, 29, 140000–140001, 2002. 15470
- 10 Rinke, A., Dethloff, K., and Fortmann, M.: Regional climate effects of Arctic Haze, *Geophys. Res. Lett.*, 31, L16202, doi:10.1029/2004GL020318, 2004. 15457
- Ritter, C., Kirsche, A., and Neuber, R.: Tropospheric Aerosol Characterized by a Raman LIDAR Over Spitsbergen, in: 22nd International Laser Radar Conference (ILRC 2004), edited by: Pappalardo, G. and Amodeo, A., vol. 561 of ESA Special Publication, 459 pp., 2004. 15457, 15475
- 15 Ritter, C., Hoffmann, A., Osterloh, L., and Boeckmann, C.: Estimation of the Liquid Water Content of a low level Arctic winter cloud, in: 24th International Laser Radar Conference (ILRC 2008), vol. 1, 597–582, 2008. 15465
- Rowe, P., Miloshevich, L., Turner, D., and Walden, V.: Dry Bias in Vaisala RS90 Radiosonde Humidity Profiles over Antarctica, *J. Atmos. Ocean. Tech.*, 25, 1529–1541 2008. 15476
- 20 Sassen, K. and Benson, S.: A Midlatitude Cirrus Cloud Climatology from the Facility for Atmospheric Remote Sensing. Part II: Microphysical Properties Derived from Lidar Depolarization, *J. Atmos. Sci.*, 58, 2103–2112, 2001. 15466
- Scheuer, E., Talbot, R., Dibb, J., Seid, G., DeBell, L., and Lefer, B.: Seasonal distributions of fine aerosol sulfate in the North American Arctic basin during TOPSE, *J. Geophys. Res.-Atmos.*, 108, 18–11, 2003. 15473
- 25 Schweiger, A., Lindsay, R., Key, J., and Francis, J.: Arctic clouds in multiyear satellite data sets, *Geophys. Res. Lett.*, 26, 1845–1848, 1999. 15456
- Shaw, G.: The Arctic Haze Phenomenon, *B. Am. Meteorol. Soc.*, 76, 2403–2414, 1995. 15455
- 30 Shiobara, M., Yabuki, M., and Kobayashi, H.: A polar cloud analysis based on Micro-pulse Lidar measurements at Ny-Ålesund, Svalbard and Syowa, Antarctica, *Phys. Chem. Earth A/B/C*, 28, 1205–1212, 2003. 15456, 15463, 15465, 15476
- Shiobara, M., Yabuki, M., Neuber, R., Spinhirne, J. D., Welton, E. J., Campbell, J. R., Hart,

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

W. D., and Berkoff, T. A.: Arctic experiment for ICESat/GLAS ground validation with a Micro-Pulse Lidar at Ny-Ålesund, Svalbard, *Polar Meteorology and Glaciology*, 20, 28–39, 2006. 15464

Shupe, M. and Intrieri, J.: Cloud Radiative Forcing of the Arctic Surface: The Influence of Cloud Properties, Surface Albedo, and Solar Zenith Angle, *J. Climate*, 17, 616–628, 2004. 15454

Shupe, M., Kollias, P., Persson, P., and McFarquhar, G.: Vertical Motions in Arctic Mixed-Phase Stratiform Clouds, *J. Atmos. Sci.*, 65, 1304–1322, 2008. 15456, 15471

Solomon, S. and Qin, D.: *Climate Change 2007 –The Physical Science Basis, Contribution of Working Group I to the Fourth Assessment Report of the IPCC*, <http://www.ipcc.ch/ipccreports/ar4-wg1.htm>, 2007. 15455

Stohl, A.: Characteristics of atmospheric transport into the Arctic troposphere, *J. Geophys. Res.-Atmos.*, 111, D11306, doi:10.1029/2005JD006888, 2006. 15455, 15467

Stohl, A., Berg, T., Burkhardt, J. F., Fjærraa, A. M., Forster, C., Herber, A., Hov, Ø., Lunder, C., McMillan, W. W., Oltmans, S., Shiobara, M., Simpson, D., Solberg, S., Stebel, K., Ström, J., Tørseth, K., Treffeisen, R., Virkkunen, K., and Yttri, K. E.: Arctic smoke – record high air pollution levels in the European Arctic due to agricultural fires in Eastern Europe in spring 2006, *Atmos. Chem. Phys.*, 7, 511–534, 2007, <http://www.atmos-chem-phys.net/7/511/2007/>. 15455

Stone, R., Dutton, E., Harris, J., and Longenecker, D.: Earlier spring snowmelt in northern Alaska as an indicator of climate change, *J. Geophys. Res.-Atmos.*, 107, 4089, doi:10.1029/2000JD000286, 2002. 15455

Treffeisen, R., Rinke, A., Fortmann, M., Dethloff, K., Herber, A., and Yamanouchi, T.: A case study of the radiative effects of Arctic aerosols in March 2000, *Atmos. Environ.*, 39(5), 899–911, 2005. 15457

Treffeisen, R., Krejci, R., Ström, J., Engvall, A. C., Herber, A., and Thomason, L. W.: Humidity observations in the Arctic troposphere over Ny-Ålesund, Svalbard based on 14 years of radiosonde data, *Atmos. Chem. Phys. Discuss.*, 7, 1261–1293, 2007, <http://www.atmos-chem-phys-discuss.net/7/1261/2007/>. 15458

Twomey, S.: *Atmospheric Aerosols*, Elsevier Scientific Publishing Co., New York, NY, USA, 1977. 15455

van de Hulst, H. C.: *Light scattering by small particles*, Wiley VCH, 1981. 15470

Vavrus, S. and Waliser, D.: An Improved Parameterization for Simulating Arctic Cloud Amount in the CCSM3 Climate Model, *J. Climate*, 21, 5673–5687, 2008. 15457

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007A. Hoffmann et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Table 1. Observed frequency of occurrence of the 8 different trajectory clusters, the * denotes cluster analysis from Eneroth et al. (2003).

	1	2	3	4	5	6	7	8
March/April 2007	7.0	10.7	8.6	12.3	20.5	10.7	11.9	18.4
ASTAR 2007	8.3	0	11.5	17.7	10.4	19.8	21.9	10.4
Annual Mean 92-01*	15.2	13.3	9.2	11.1	8.3	16.7	9.4	16.9
March/April 92-01*	17.6	17.5	10.0	12.0	7.0	13.0	7.9	15.2

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Table 2. KARL backscatter ratio (BSR at 532 nm) according to altitude intervals. Given is the median value as well as frequency of certain BSR values within the different altitude intervals.

H	median BSR	10–3	3–2	2–1.5	1.5–1.2	1.2–1.1	1.1–1.05
9–10	1.093	0	0	0	1.3	31.2	67.5
8–9	1.100	1.0	1.7	1.3	4.3	43.8	44.0
7–8	1.103	2.2	2.2	3.0	5.7	42.8	44.0
6–7	1.105	3.0	2.2	3.2	8.9	43.4	39.2
5–6	1.112	2.3	2.8	5.8	11.4	42.8	34.9
4–5	1.124	4.9	5.2	6.1	15.8	38.2	29.8
3–4	1.126	2.1	0.3	3.0	16.8	51.8	25.9
2–3	1.142	1.5	1.4	4.4	22.4	45.8	24.6
1–2	1.178	1.3	1.1	2.1	33.2	37.2	24.9

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Table 3. KARL data classification cases with respective VD and BSR thresholds and case studies.

		BSR	VD	case studies
C1:	clear sky	$BSR < 1.2$	$VD < 1.8$	no case studies
C2:	water	$BSR > 1.2$	$VD < 1.2$	15.03.
C3:	high BSR, low VD	$BSR > 1.2$	$1.2 < VD < 1.8$	08.03./07.04./13.03.
C4:	low BSR, high VD	$BSR < 1.2$	$VD > 1.8$	15.03.
C5:	thin aerosol	$1.2 < BSR < 2$	$1.8 < VD < 5$	15.03./18.03./ (07.04.)
C6:	thick aerosol	$BSR > 2$	$1.8 < VD < 5$	not observed
C7:	thin ice, low VD	$1.2 < BSR < 2$	$5 < VD < 16$	13.03./15.03.
C8:	thick ice, low VD	$BSR > 2$	$5 < VD < 16$	13.03./15.03.
C9:	thin ice, high VD	$1.2 < BSR < 2$	$VD > 16$	15.03.
C10:	thick ice, high VD	$BSR > 2$	$VD > 16$	15.03.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Table 4. KARL frequency of occurrence of cases C1 to C10 according to the data classification in Tab. 3 for different altitude intervals. Percentages are within an accuracy of 1% , peak values of VD and BSR average out.

H [km]	C1	C2	C3	C4	C5	C6	C7	C8	C9	C10
10–12	98	0	1	1	0	0	0	0	0	0
9–10	92	1	7	0	0	0	0	0	0	0
8–9	79	0	2	8	5	2	1	3	0	0
7–8	67	0	1	16	6	1	3	2	0	3
6–7	49	0	1	30	7	0	7	3	0	2
5–6	37	0	1	35	10	1	6	6	0	3
4–5	37	1	2	42	14	0	1	2	0	2
3–4	44	0	6	29	15	0	1	2	0	1
2–3	35	1	9	30	20	1	1	1	0	1
1–2	20	2	13	36	22	2	1	1	0	0

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Table 5. KARL frequency of occurrence of cases C1 to C10 according to the data classification in Table 3 for two different air mass origin areas. Considered are the means of the altitude intervals from 2 to 3 and 3 to 4 km as this corresponds to the 700 hPa trajectories used for the cluster analysis. Percentages are within an accuracy of 1%, peak values of VD and BSR average out.

	C1	C2	C3	C4	C5	C6	C7	C8	C9	C10
Eurasia (Cluster 2, 5, 8)	51	2	7	3	28	6	2	0	1	0
Canada (Cluster 6, 7)	13	3	11	37	28	1	5	2	0	0

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Ground-based lidar
measurements from
Ny-Ålesund during
ASTAR 2007**

A. Hoffmann et al.




Fig. 1. Temperature profiles obtained with 71 radio soundings from 1 March to 30 April 2007, red dots: tropopause, blue dots: temperature inversions above 2 K, white dots: surface-based temperature inversions above 0.25 K.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Ground-based lidar
measurements from
Ny-Ålesund during
ASTAR 2007**

A. Hoffmann et al.




Fig. 2. Relative humidity profiles obtained with 71 radio soundings from 1 March to 30 April 2007.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Ground-based lidar
measurements from
Ny-Ålesund during
ASTAR 2007**

A. Hoffmann et al.




Fig. 3. Cluster analysis for the 5-day backward trajectories (700 hPa level) performed with the PEP-Tracer model.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Ground-based lidar
measurements from
Ny-Ålesund during
ASTAR 2007**

A. Hoffmann et al.




Fig. 4. MPL cloud fraction in 10 altitude intervals of 1 km width depending on relative humidity and temperature, derived from the comparison of 57 radiosondes with respective MPL data.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Ground-based lidar
measurements from
Ny-Ålesund during
ASTAR 2007**

A. Hoffmann et al.




Fig. 5. 1-km-means of BSR and VD for the KARL data and frequency of cloud occurrence for the MPL data for 10 altitude intervals, MPL clear sky fraction was 33%.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Ground-based lidar
measurements from
Ny-Ålesund during
ASTAR 2007

A. Hoffmann et al.




Fig. 6. Scatter plot of volume depolarization VD vs. backscatter ratio BSR (532 nm) for 9 altitude intervals, the lowest interval was neglected due to high uncertainties. Dashed lines refer to BSR and VD thresholds defined in Sect. 3.2.2.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Ground-based lidar
measurements from
Ny-Ålesund during
ASTAR 2007**

A. Hoffmann et al.




Fig. 7. Scatter plot of volume depolarization VD vs. backscatter ratio BSR (532 nm) for the 8 trajectory clusters (see Fig. 3). Dashed lines refer to BSR and VD thresholds defined in Sect. 3.2.2.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.





Fig. 8. KARL data classification according to VD and BSR threshold values.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Ground-based lidar
measurements from
Ny-Ålesund during
ASTAR 2007**

A. Hoffmann et al.

**Fig. 9.** VD vs. BSR for different layers and time frames on 8 March 2007.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

KARL: BC 532 & VD – 8 March 2007




Fig. 10. Temporal evolution of BC (532 nm) and VD profiles from 15:43 to 16:17 UTC on 8 March 2007.

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Ground-based lidar
measurements from
Ny-Ålesund during
ASTAR 2007**

A. Hoffmann et al.

**Fig. 11.** VD vs. BSR for different layers and time frames on 13 March 2007.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[⏴](#)[⏵](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Ground-based lidar measurements from Ny-Ålesund during ASTAR 2007

A. Hoffmann et al.





Fig. 12. Temporal evolution of BC (532 nm) and VD profiles from 15:47 to 17:54 UTC on 13 March 2007.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Ground-based lidar
measurements from
Ny-Ålesund during
ASTAR 2007**

A. Hoffmann et al.

**Fig. 13.** VD vs. BSR for different layers and time frames on 15 March 2007.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Ground-based lidar
measurements from
Ny-Ålesund during
ASTAR 2007**

A. Hoffmann et al.




Fig. 14. Left: BC (532 nm) and VD profiles at 11:20 UTC on 15 March 2007, right: T and RH profiles from the radiosonde launch at 11:00 UTC on 15 March 2007.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Ground-based lidar
measurements from
Ny-Ålesund during
ASTAR 2007**

A. Hoffmann et al.




Fig. 15. VD vs. BSR for different layers and time frames on 7 April 2007.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Ground-based lidar
measurements from
Ny-Ålesund during
ASTAR 2007

A. Hoffmann et al.





Fig. 16. Left: BC (532 nm) and VD profiles at 11:06 UTC on 7 April 2007, right: T and RH profiles from the radiosonde launch at 11:00 UTC on 7 April 2007.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Ground-based lidar
measurements from
Ny-Ålesund during
ASTAR 2007**

A. Hoffmann et al.

**Fig. 17.** Temporal evolution of BC (532 nm) profiles on 7 April 2007.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Ground-based lidar
measurements from
Ny-Ålesund during
ASTAR 2007**

A. Hoffmann et al.




Fig. 18. AOD (532 nm) and Ångström exponent α derived from photometer measurements on 7 April 2007.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

NOAA HYSPLIT MODEL
 Backward trajectories ending at 1100 UTC 15 Mar 07
 GDAS Meteorological Data




Fig. 19. Backward trajectories for aerosol layers on 15 March 2007 at 11:00 UTC, calculated with the NOAA HYSPLIT model.

Ground-based lidar
 measurements from
 Ny-Ålesund during
 ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Ground-based lidar
measurements from
Ny-Ålesund during
ASTAR 2007**

A. Hoffmann et al.




Fig. 20. VD vs. BSR for different layers and time frames on 18 March 2007.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

NOAA HYSPLIT MODEL
 Backward trajectories ending at 2300 UTC 18 Mar 07
 GDAS Meteorological Data




Fig. 21. Backward trajectories for aerosol layers on 18 March 2007 at 23:00 UTC, calculated with the NOAA HYSPLIT model.

Ground-based lidar
 measurements from
 Ny-Ålesund during
 ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NOAA HYSPLIT MODEL
 Backward trajectories ending at 11 UTC 07 Apr 07
 GDAS Meteorological Data




Fig. 22. Backward trajectories for aerosol layers on 7 April 2007 at 11:00 UTC, calculated with the NOAA HYSPLIT model.

Ground-based lidar
 measurements from
 Ny-Ålesund during
 ASTAR 2007

A. Hoffmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion