

This discussion paper is/has been under review for the journal *Atmospheric Chemistry and Physics (ACP)*. Please refer to the corresponding final paper in *ACP* if available.

The ARCTAS aircraft mission: design and execution

**D. J. Jacob¹, J. H. Crawford², H. Maring³, A. D. Clarke⁵, J. E. Dibb⁴,
R. A. Ferrare², C. A. Hostetler², P. B. Russell⁶, H. B. Singh⁶, A. M. Thompson⁷,
G. E. Shaw⁸, E. McCauley⁹, J. R. Pederson⁹, and J. A. Fisher¹**

¹Harvard University, Cambridge, Massachusetts, USA

²NASA Langley Research Center, Hampton, Virginia, USA

³NASA Headquarters, Washington, DC, USA

⁴University of New Hampshire, Durham, New Hampshire, USA

⁵University of Hawaii, Honolulu, Hawaii, USA

⁶NASA Ames Research Center, Moffett Field, California, USA

⁷Pennsylvania State University, State College, Pennsylvania, USA

⁸University of Alaska, Fairbanks, Alaska, USA

⁹California Air Resources Board, Sacramento, California, USA

Received: 23 July 2009 – Accepted: 31 July 2009 – Published: 12 August 2009

Correspondence to: D. J. Jacob (djacob@fas.harvard.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The NASA Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS) mission was conducted in two 3-week deployments based in Alaska (April 2008) and western Canada (June–July 2008). The goal of ARCTAS was to better understand the factors driving current changes in Arctic atmospheric composition and climate, including (1) transport of mid-latitude pollution, (2) boreal forest fires, (3) aerosol radiative forcing, and (4) chemical processes. ARCTAS involved three aircraft: a DC-8 with detailed chemical payload, a P-3 with extensive aerosol payload, and a B-200 with aerosol remote sensing instrumentation. The aircraft augmented satellite observations of Arctic atmospheric composition, in particular from the NASA A-Train, by (1) validating the data, (2) improving constraints on retrievals, (3) making correlated observations, and (4) characterizing chemical and aerosol processes. The April flights (ARCTAS-A) sampled pollution plumes from all three mid-latitude continents, fire plumes from Siberia and Southeast Asia, and halogen radical events. The June–July flights (ARCTAS-B) focused on boreal forest fire influences and sampled fresh fire plumes from northern Saskatchewan as well as older fire plumes from Canada, Siberia, and California. The June–July deployment was preceded by one week of flights over California sponsored by the California Air Resources Board (ARCTAS-CARB). The ARCTAS-CARB goals were to (1) improve state emission inventories for greenhouse gases and aerosols, (2) provide observations to test and improve models of ozone and aerosol pollution. Extensive sampling across southern California and the Central Valley characterized emissions from urban centers, offshore shipping lanes, agricultural crops, feedlots, industrial sources, and wildfires.

1 Introduction

The Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS) mission was conducted in April and June–July 2008 by the Global Tropo-

ACPD

9, 17073–17123, 2009

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



spheric Chemistry Program and the Radiation Sciences Program of the US National Aeronautics and Space Administration (NASA). Its objective was to better understand the factors driving current changes in Arctic atmospheric composition and climate. It used chemical and radiative measurements from three research aircraft (DC-8, P-3, B-200) to interpret and augment the continuous observations of Arctic atmospheric composition from satellites. The aircraft were based in Alaska in April (ARCTAS-A) and in western Canada in June–July (ARCTAS-B). ARCTAS was part of the international POLARCAT program of field observations of Arctic atmospheric composition during the 2007–2008 International Polar Year (<http://www.polarcat.no/>) and was coordinated with several concurrent field programs. This paper provides an overview of the ARCTAS mission focusing on background, design, execution, and description of its components. It is intended as a general reference for future papers presenting ARCTAS results.

The Arctic has been warming rapidly over the past century, and the pace has accelerated in the past decades (Trenberth et al., 2007). Sea ice cover and permafrost show rapid declines (Lemke et al., 2007). General circulation models (GCM) simulations project strong 21st-century warming in the Arctic, reflecting in part the sea ice feedback (Christensen et al., 2007). Even so, these models underestimate the observed warming over the past decades (Shindell and Faluvegi, 2009), suggesting that they might underestimate future warming. One possible factor missing from current climate models is proper representation of the regional radiative forcing from aerosols and tropospheric ozone, which may play a particularly large role in the Arctic radiative budget and climate response (Shindell et al., 2007; Quinn et al., 2008). Decreasing the forcing from these short-lived agents could provide an opportunity to curb Arctic warming. The primary goal of ARCTAS was to better understand the sources of aerosols and tropospheric ozone in the Arctic, as well as the radiative properties of Arctic aerosols and of the underlying surface, in order to improve estimates of the associated radiative forcings.

Inflow of pollution from northern mid-latitudes is a well-known source of aerosols and ozone to the Arctic. It prompts not only climatic but air quality and ecological concerns.

The ARCTAS aircraft mission: design and executionD. J. Jacob et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The ARCTAS aircraft mission: design and executionD. J. Jacob et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

It has been known for decades that the Arctic is heavily affected by mid-latitude pollution in winter-spring, manifested by visible “Arctic haze” aerosol layers (Shaw, 1995). Arctic observations have been mainly at the surface, leading to the conventional view of a strong seasonal variation in mid-latitude pollution inflow with northern Europe as the dominant contributor. However, recent models and limited observations show transport from mid-latitudes taking place year-round through different processes at different altitudes and involving North America and Asia as well as Europe (Klonecki et al., 2003; Koch and Hansen, 2005; Stohl, 2006; Shindell et al., 2008). Sources at mid-latitudes are rapidly changing, with increases in Asia driven by industrialization and decreases in North America and Europe driven by air quality regulations. Better understanding of pollution transport to the Arctic and of the associated source-receptor relationships was a priority for ARCTAS.

Chemical processing plays a major role in determining the evolution of aerosols, ozone, and other pollutants in the Arctic. The chemical environment in the Arctic is characterized by remoteness, intense cold, and long periods of light and darkness. Photochemical reactions on snow and sea ice surfaces following the winter darkness provide a springtime source of halogen radicals that drive rapid ozone destruction and mercury oxidation (Simpson et al., 2007a). Halogen radical generation may be highly sensitive to climate change through sea ice cover (Simpson et al., 2007b). A goal of ARCTAS was to better understand the radical photochemistry of the Arctic atmosphere and its implications for the regional budgets of aerosols, ozone, and mercury.

Boreal forest fires in Siberia, Canada, and Alaska represent a major perturbation to the Arctic atmosphere in summer affecting the carbon cycle (Preston and Schmidt, 2006), climate (Amiro et al., 2001), air quality (Colarco et al., 2004), and land ecology (Soja et al., 2007). Long-term records for Canada show an increase in fires over the past 50 years correlated with rising temperatures (Gillett et al., 2004; Girardin, 2007), and this increase is expected to continue as a result of warming (Stocks et al., 1998; Flannigan et al., 2005). Aerosols emitted by the fires could represent a significant forcing or feedback to regional climate change in the Arctic. The summer deployment

of ARCTAS focused on better quantifying the fire emissions, the aerosol and chemical evolution in the fire plumes, and the associated radiative effects.

A central motivation for ARCTAS was to enhance the ability of satellite observations of atmospheric composition to address the above issues. The Arctic atmosphere is difficult to observe in situ because of its vastness, remoteness, and harshness. Polar-orbiting satellites have unique potential. They observe the long-range transport of mid-latitude pollution to the Arctic. They detect fires and the associated smoke plumes. They measure aerosol optical depths and other aerosol properties, and the concentrations of gas-phase species including ozone, CO, NO₂, and BrO that are important drivers of Arctic chemistry. NASA has a particular stake in satellite observations through its A-train suite of satellites (Aqua, Cloudsat, CALIPSO, PARASOL, Aura) following each other on the same polar orbit with total separation of 8 min. Table 1 lists the most important satellite measurements of atmospheric composition operational during ARCTAS.

Observation of the Arctic from space poses unique challenges including bright and cold surfaces, low sun angles, seasonal darkness, and extensive cloud cover. In addition, many species of environmental interest are not directly observable from space. ARCTAS aimed to use detailed observations from aircraft to provide the validation, retrieval constraints, correlative data, and process information needed to develop the potential of satellites for Arctic research. The combination of satellite and aircraft data can provide powerful information for constraining and evaluating models of Arctic atmospheric composition and climate, and thus improve model projections of future change.

The summer ARCTAS-B deployment was preceded by one week of flights over and around California to address regional issues of air quality and climate forcing. These flights were sponsored by the California Air Resources Board (CARB). Many areas of California exceed the state ozone and fine particulate matter (PM_{2.5}) air quality standards, and little improvement in ozone has been observed over the last decade despite continued emission reductions (Cox et al., 2008). Controlling emissions from heavy-duty diesel trucks has been far more difficult than from passenger cars. Shipping emis-

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sions have increased. Long-range transport of pollution from Asia or Mexico may be offsetting benefits from local pollution control measures. In addition to these air quality concerns, CARB is embarking on strategies to reduce state greenhouse gas emissions. The ARCTAS-CARB flights focused on collecting data to aid California improve its emission inventories and air quality models.

2 ARCTAS scientific themes and mission design

ARCTAS was organized around four main research themes: (1) long-range transport of pollution to the Arctic, (2) boreal forest fires, (3) aerosol radiative properties, and (4) ozone budget and chemical processes. We elaborate on each of these themes and how they defined the objectives and design of the mission. We also discuss the separate California air quality objectives of the ARCTAS-CARB component.

2.1 Long-range transport of pollution to the Arctic

Transport of pollution from northern mid-latitudes to the Arctic takes place throughout the year but the processes and source regions vary strongly with season (Shindell et al., 2008). Intense cooling in winter produces a “polar dome” that shields the Arctic lower troposphere from isentropic transport from lower latitudes (Klonecki et al., 2003). Low-altitude transport from cold northern Europe can still take place and provides then the dominant source of pollution measured at the surface (Stohl, 2006). Transport from North America and East Asia takes place mainly at higher altitudes, in particular through warm conveyor belts (WCBs) associated with storm tracks on the east coast of the continents (Koch and Hansen, 2005; Stohl, 2006). This cyclonic activity increases from winter to spring, resulting in a springtime peak of sulfate concentrations in the free troposphere (Scheuer et al., 2003). Mid-latitude pollution influence over the Arctic still takes place in summer and fall but is diluted by the weaker general circulation and the greater convective ventilation of the mid-latitude continents (Klonecki et al., 2003).

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Pollution transport to the Arctic in winter-spring is in general highly layered, which has led to the terminology “Arctic haze”. This layering reflects the stratification of the Arctic atmosphere, the lack of precipitation, and also possibly the local heating by light-absorbing aerosol (Quinn et al., 2008; Stone et al., 2008). Scavenging of aerosols and other water-soluble species takes place during WCB lifting of air masses, but this scavenging is not complete (Koike et al., 2003; Miyazaki et al., 2003) and preferentially removes sulfate relative to black carbon (BC) (Park et al., 2005).

The long-range transport theme of ARCTAS targeted the following questions:

- What are the transport pathways for different pollutants to the Arctic?
- What are the contributions from different source regions and what are the source-receptor relationships?

Satellites have obvious potential for observing the long-range transport of pollution to the Arctic. This potential is least in the winter when conditions are cold and dark, and when most of the transport takes place at low altitudes under a prevailingly cloudy atmosphere. We chose therefore to focus on spring as holding more promise for satellite observation, and also as offering more opportunity to observe the long-range transport from Asia which is less understood than transport from Europe and more likely to grow in the future. An attractive aspect of focusing on Asian influence in spring was the link to previous NASA aircraft campaigns along the Asian Pacific Rim in spring 1994 (PEM-West B; Hoell et al., 1997) and spring 2001 (TRACE-P; Jacob et al., 2003), and over the Northeast Pacific and southern Alaska in spring 2006 (INTEX-B; Singh et al., 2009). During those campaigns we observed frequent lifting of Asian pollution in WCBs followed by northeastward transport toward the Arctic in the general direction of Alaska (Fuelberg et al., 2003). This led to the selection of Fairbanks, Alaska as the main operational base for the ARCTAS-A spring deployment, and an overnight base in Thule, Greenland to achieve broader spatial coverage.

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2 Boreal forest fires

Wildfires in North America and Eurasia are important regional and hemispheric sources of aerosols, tropospheric ozone, and possibly mercury (Lavoue et al., 2000; Turetsky et al., 2006; Generoso et al., 2007; Leung et al., 2007). Fires are particularly active in the boreal forest zone in summer. Buoyancy generated by the fires can result in injection above the boundary layer (Kahn et al., 2008; McMillan et al., 2008) and in extreme cases in the stratosphere (Fromm and Servanckx, 2003). This lifting of emissions has important implications for the transport, evolution, and impacts of the fire plumes (Morris et al., 2006; Turquety et al., 2007; Hyer et al., 2007; Leung et al., 2007).

Emissions from boreal fires depend on burn conditions (Goode et al., 2000; Reid et al., 2005; Preston and Schmidt, 2006). Rapid aerosol evolution takes place in the fresh plumes through particle coagulation and condensation of organic gases, ammonia, and water (Fiebig et al., 2002; Colarco et al., 2004). This near-field evolution of the fire plumes determines their regional and global radiative impact (Westphal and Toon, 1991; Fiebig et al., 2003). Ozone production takes place in the fresh plumes from the precursor nitrogen oxide (NO_x) radicals, though this is limited by conversion to peroxyacetyl nitrates (PANs) (Jacob et al., 1992). Subsequent decomposition of PANs as the plumes age and sink yields additional ozone production on the hemispheric scale (Real et al., 2007).

The boreal forest fires theme of ARCTAS targeted the following questions:

- What are the emissions from boreal forest fires, and what is the near-field chemical evolution in the fire plumes?
- What are the optical properties of the aerosols emitted by boreal forest fires, and how do they evolve during aging of the fire plume?
- What are the injection heights of fire emissions and what processes determine these heights?

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- What are the implications of boreal forest fires for regional and global atmospheric composition?

Addressing these questions was the principal objective of the ARCTAS-B summer deployment. July was selected to maximize the probability of sampling large fires in boreal North America. The DC-8 and P-3 aircraft were based in Cold Lake, Alberta (54 N, 110 W), and the B-200 in Yellowknife, Northwest Territories (62 N, 114 W), on the basis of climatological fire data and available facilities. These bases also offered the opportunity to sample aged plumes from fires in Eurasia and the western US. An overnight base in Thule for the DC-8 was included in the summer deployment to increase spatial coverage and the opportunities for sampling aged biomass burning plumes.

2.3 Aerosol radiative forcing

Aerosol radiative forcing, both direct (clear-sky scattering and absorption) and indirect (aerosol effects on clouds), could be an important regional driver of Arctic climate change (Quinn et al., 2008). Major aerosol types include anthropogenic particles associated with Arctic haze, biomass burning particles from both boreal and lower-latitude fires, and mineral dust particles. These different origins can be separated on the basis of measured chemical, microphysical, and optical aerosol properties (Bergstrom et al., 2007; Clarke et al., 2007). Radiative effects of aerosols in the Arctic vary not only with aerosol composition but also with solar irradiance, surface albedo, and altitude (Pfister et al., 2008; Stone et al., 2008). Black carbon aerosol deposited to snow has a significant warming effect and may promote snow and sea ice melting (Clarke and Noone, 1985; Hansen and Nazarenko, 2004; Flanner et al., 2007).

Aerosol optical properties and the resulting radiative forcing evolve with atmospheric aging (Fiebig et al., 2003; Preston and Schmidt, 2006). This has been observed in particular for boreal forest fire plumes in Canada (O'Neill et al., 2002). Anthropogenic Arctic haze has a wide range of ages and scavenging histories, and this is expected

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in turn to yield a range of optical properties and radiative forcing (Quinn et al., 2007). Observations and models indicate a finer particle size mode in pollution layers in summer relative to spring, reflecting disappearance of the Arctic haze layers and more opportunity for nucleation of new particles (Engvall et al., 2008; Korhonen et al., 2008).

5 The aerosol radiative forcing theme of ARCTAS targeted the following questions:

- What is the regional radiative forcing from Arctic haze and fire plumes?
- How does this forcing evolve during plume aging?
- What are the major sources of black carbon aerosol to the Arctic?
- How does black carbon aerosol deposition affect ice albedo?

10 Characterizing the forcing associated with individual aerosol layers was the central task of the P-3 aircraft equipped with an ensemble of in situ aerosol and radiative observations. Validation and interpretation of the CALIOP satellite observations of altitude-resolved aerosol extinction was a primary charge of the B-200 aircraft equipped with a high spectral resolution lidar (HSRL). During the spring deployment, the B-200
15 operated out of Barrow, Alaska to coordinate with the DOE ISDAC aircraft mission focused on aerosol-cloud interactions and centered over the long-term Atmospheric Radiation Measurement (ARM) site at Barrow.

2.4 Ozone budget and chemical processes

20 Tropospheric ozone at high northern latitudes shows a pronounced spring maximum (Monks, 2000). The TOPSE aircraft campaign over the North American Arctic in February–May 2000 (Atlas et al., 2003) found that this maximum is driven by tropospheric production rather than by stratospheric input, and that the ozone budget in the region largely reflects a balance between in situ production and loss (Browell et al., 2003; Emmons et al., 2003). However, ozonesonde data from Canadian stations

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



show significant stratospheric impact on tropospheric ozone in both spring and summer (Tarasick et al., 2007; Thompson et al., 2007). Stratospheric influence could be particularly important in the upper troposphere where ozone is of most concern as a greenhouse gas, and this influence could increase in the future as stratospheric ozone recovers and the Brewer-Dobson circulation increases due to greenhouse forcing (Shindell et al., 2007). The Canadian ozonesonde data show a decrease in tropospheric ozone in the 1980s followed by an increase since then, for reasons that are unclear (Tarasick et al., 2005; Kivi et al., 2007).

Tropospheric ozone chemistry in the Arctic is largely determined by the supply of hydrogen oxide radicals (HO_x) and nitrogen oxide radicals (NO_x), but the processes controlling the abundances of these radicals are poorly understood. An additional feature of the Arctic boundary layer in spring is the presence of high concentrations of bromine and chlorine radicals produced by photochemical reactions involving sea salt particles deposited on sea ice. Bromine radicals can lead to rapid catalytic destruction of ozone in the boundary layer, and to oxidation and subsequent deposition of mercury. These occurrences are called ozone depletion events (ODEs) and mercury depletion events (MDEs) (Simpson et al., 2007b; Steffen et al., 2008). Chlorine atoms can also lead to rapid oxidation of volatile organic compounds (VOCs) (Zeng et al., 2006). Fresh sea ice and ice leads are thought to be preferential environments for generation of halogen radicals (Bottenheim and Chan, 2006; Simpson et al., 2007a), but the ODEs can persist for over 1000 km downwind (Ridley et al., 2003) and extend over vast tracts of the Arctic boundary layer (Zeng et al., 2006). Satellite observations of tropospheric BrO starting with the GOME instrument launched in 1995 (Chance, 1998) have provided detailed mapping of high-BrO events to infer preferential source regions, seasonal variations, and long-term trends of halogen radical generation (Hollwedel et al., 2004). However, these BrO observations had not been validated prior to ARCTAS.

The chemical processing theme of ARCTAS targeted the following questions:

- What controls HO_x and NO_x chemistry in the Arctic?

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- What drives halogen radical chemistry in the Arctic, and what is its regional extent?
- What are the regional implications for ozone, aerosols, and mercury?
- How does stratosphere-troposphere exchange affect tropospheric ozone in the Arctic?

The DC-8 included a detailed chemical payload to address these questions. One priority was to validate the satellite observations of high BrO events. These events are most frequent in March–April (Hollwedel et al., 2004). We chose April as more favorable in terms of photochemistry, satellite observation capabilities, and operations weather.

2.5 California air quality (ARCTAS-CARB)

The objectives of the California deployment (ARCTAS-CARB) were to: (1) improve state emission inventories for greenhouse gases and aerosols; (2) characterize off-shore emissions of sulfur and other pollutants from shipping and natural sources; and (3) characterize upwind boundary conditions for modeling local surface ozone and PM_{2.5}. Despite extensive abatement efforts, air quality standards are widely exceeded in southern California and the Central Valley (Cox et al., 2008). Little improvement in ozone has been observed in the past decade despite continued reduction in precursor emissions. The causes for this lack of progress are unknown but could reflect poorly understood non-linearities in the photochemical system. There is far more sulfate in southern California than can be accounted for in present emission inventories and ship emissions may be implicated. Another issue is the poorly quantified import of pollution from Asia and Mexico (Jacob et al., 1999; Oltmans et al., 2008; Wang et al., 2009). California is now embarking on synergistic strategies to control greenhouse gas emissions together with air pollutants. A first step is to improve inventories for these emissions.

The ARCTAS-CARB flights targeted the following questions:

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- How good is our current understanding of the HO_x-NO_x-O₃-aerosol photochemical system over the Los Angeles Basin as represented in air quality models?
- How should upwind boundary conditions for simulating air quality in California be specified?
- 5 – How do ship emissions and long-range transport affect the sulfur budget in southern California?
- What are the state emissions of VOCs and greenhouse gases from urban and industrial activities, agricultural operations, and wildfires?

The flights took advantage of the fully loaded DC-8 and P-3 aircraft based in California just before the ARCTAS-B summer deployment. In addition to the local air quality objectives, these California flights helped to address the ARCTAS Arctic objectives by providing contrasting data, expanding the ranges of the instruments, and characterizing emissions from California wildfires that would then be sampled as aged plumes from the aircraft based in Canada.

3 Mission execution

3.1 Aircraft payloads and support

Table 2 gives the characteristics of the DC-8, P-3, and B-200 aircraft as used in ARCTAS. Table 3a–c lists the payloads for each. The DC-8 had the largest range and payload. It was responsible for addressing the long-range transport and chemical processing themes of ARCTAS, and also contributed to the boreal fires theme. Its payload included measurements of ozone (in situ and remote); HO_x, NO_x, and BrO radicals; reservoirs for these radicals and related species; aerosols, carbon gases, and mercury; tracers of transport; and actinic fluxes. Measurements of HO_x radicals were made by two independent techniques (LIF and CIMS) aboard the DC-8, since previous

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



comparisons between data sets and with models in past aircraft missions had shown inconsistent results (Ren et al., 2008).

The P-3 and B-200 payloads focused on aerosol characterization and radiation measurements, targeting the aerosol radiative forcing theme of ARCTAS and also addressing aerosol source issues related to Arctic haze and boreal fires. The P-3 payload included in situ measurements of aerosol properties; radiative flux measurements; surface reflectivity measurements; and CO as a pollution tracer. The B-200 payload included a high-spectral resolution lidar (HSRL) and a scanning polarimeter.

Intensive ozonesonde launches during ARCTAS were made at the 17-site Arctic Intensive Ozonesonde Network Study (ARC-IONS) network (Table 3d). The spring sites were chosen to emphasize latitudinal gradients, opportunities to sample Arctic pollution, and coordination with aircraft and ground stations. The summer sites were aligned to capture cross-continental flow with Whitehorse, Yellowknife, and three eastern Canadian locations (Goose Bay, Egbert, Yarmouth) added to the network. Special launches were timed to coincide with TES satellite instrument overpasses for ozone validation.

ARCTAS execution was also supported by additional ground-based measurements (Table 3e) and various forecasts and analyses (Table 3f). The ground-based measurements included aerosol lidars, in situ measurements from an instrumented trailer, and multi-axis differential optical absorption spectroscopy (MAX-DOAS) measurements of BrO at Barrow. Meteorological and chemical transport model (CTM) forecasts, near-real-time satellite and CTM data, and fire reports were used to guide flights on a day-to-day basis.

3.2 Deployments

The spring ARCTAS-A deployment took place from 1 to 21 April, 2008. Flight tracks are shown in Fig. 1a. The DC-8 and P-3 were based in Fairbanks, Alaska (65 N, 148 W) and the B-200 in Barrow, Alaska (71 N, 157 W). The DC-8 made overnight flights to Thule, Greenland (77 N, 69 W) and Iqaluit, Nunavut (64 N, 69 W). Iqaluit was used as a contingency airport because of bad weather in Thule on the second planned

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The ARCTAS aircraft mission: design and executionD. J. Jacob et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

overnight. The P-3 also made an overnight flight to Thule. Flights were coordinated with the NOAA ARCPAC and DOE ISDAC aircraft missions concurrently based in Fairbanks. ARCPAC focused on the factors driving the radiative forcing by aerosols and tropospheric ozone in the Arctic, dovetailing with the objectives of ARCTAS. It used a NOAA P-3 aircraft with extensive in situ instrumentation for aerosols, ozone, related species, and radiation. ISDAC focused on aerosol-cloud interactions in the Arctic, including indirect forcing and scavenging processes. It used a Convair-580 aircraft with extensive in situ instrumentation for aerosol and cloud microphysical properties. Flight tracks were mainly to and over the ARM site at Barrow and involved coordination with the B-200.

The summer ARCTAS-B deployment took place from 26 June to 14 July, 2008. Flight tracks are shown in Fig. 1b. The DC-8 and P-3 were based in Cold Lake, Alberta (54 N, 110 W) and the B-200 in Yellowknife, Northwest Territories (62 N, 114 W). The DC-8 did a two-night stay in Thule to overfly Summit, Greenland (73 N, 39 W), where the US NSF led an intensive ground-based campaign. The Thule base also offered the opportunity to coordinate flights with the German DLR and French CNRS aircraft missions based in Kangerlussuaq, Greenland (67 N, 51 W) and focused on sampling North American biomass burning outflow.

The ARCTAS-CARB deployment took place from 18 to 24 June, 2008, immediately preceding the ARCTAS summer deployment. Flight tracks are shown in Fig. 1c. The DC-8 conducted four flights (33 h) from its base in Palmdale, California (35 N, 118 W). The P-3 conducted one flight (10 h) from its base in Moffett Field, California (37 N, 122 W). Additional data relevant to ARCTAS-CARB objectives were collected during the transit flights to and from Canada.

Figure 2 shows fire activity during the spring and summer deployments as indicated by MODIS fire counts. April 2008 saw typical seasonal biomass burning in Southeast Asia but unusually early fire activity in eastern Siberia due to low snow cover, as well as unusually large agricultural burning in Kazakhstan. April 2008 was the second highest fire month in Russia (behind May 2003) over the 2001–2008 duration of the MODIS

satellite record (Fisher et al., 2009). Flights from the NOAA ARCPAC mission out of Fairbanks observed strong and persistent influence from these Russian fires (Warneke et al., 2009). Figure 3 shows surface aerosol scattering measurements made at Poker Flats near Fairbanks in spring 2008. Biomass burning influence at the surface was anomalously large during ARCTAS (1–21 April) and even more so during ARCPAC (11–23 April). This unusual biomass burning influence must be recognized in the interpretation of the data.

Summer 2008 saw in general low fire activity in Canada (1.6 Mha burned, as compared to an average of 2.8 Mha for the 1990s, Flannigan et al., 2005) but there was unusually large fire activity in northern Saskatchewan within 1-hour flight time from Cold Lake, enabling sampling of fresh fire plumes and their evolution. Burning in northern Saskatchewan was four times the 20-year average (Soja et al., 2008). Summer 2008 also saw considerable fire activity in northern California and Siberia, and plumes from these fires were repeatedly sampled by the aircraft.

3.3 Flights

Tables 4 and 5 lists the research flights of the three aircraft during the spring and summer deployments, respectively. ARCTAS-CARB flights are included in Table 5. The Tables describe some specific features of each flight including coordination with other platforms. Coordination between aircraft involved vertically stacked flights to provide complementarity between in situ and remote measurements, and wingtip-to-wingtip legs at different altitudes to intercompare instruments. Vertical spirals were also conducted over several surface sites including Barrow (ARM site, tropospheric BrO), Summit (surface and sondes), and Eureka (vertical column measurements of trace gases).

A central component of ARCTAS was to enhance the value of satellite observations of Arctic atmospheric composition (Table 1), including validation under the unique Arctic conditions and acquisition of data to improve satellite retrievals. Coordination with satellites involved underpasses and vertical spirals along the satellite orbit tracks. Top priorities were validation of the CALIOP aerosol observations by the B-200 and P-3

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



aircraft, and of the OMI tropospheric BrO observations by the DC-8 aircraft.

All flights of the spring ARCTAS-A deployment targeted mid-latitude pollution layers transported to the Arctic. The P-3 and B-200 focused on aerosol layers, often in coordination with CALIOP. The P-3 also devoted significant flight time to measuring surface albedo and reflectance, including for sea ice at Elson Lagoon just north of Barrow and over the Arctic ice and clouds near Greenland. The DC-8 conducted extensive vertical profiling from its nominal floor (0.15 km over ocean, 0.3 km over land) to the stratosphere. To ensure that the boundary layer was properly sampled, the DC-8 occasionally descended to 0.1 km over land and made deliberate missed approaches of the Barrow and Prudhoe Bay airports.

All flights of the summer ARCTAS-B deployment targeted wildfire plumes. Again, the P-3 and B-200 focused on aerosol layers and coordination with CALIOP. The DC-8 focused on plume evolution, including sampling of fresh Canadian fire plumes as well as aged Asian and California fire plumes. A few flights also sampled plumes from oil sands recovery operations near Fort McMurray in northeast Alberta. On the outbound flight to Cold Lake, the DC-8 conducted an intercomparison with the NSF HIAPER aircraft as it completed its pre-HIPPO (HIAPER Pole-to-Pole Observations) mission out of Colorado. Pre-HIPPO focused on testing the HIAPER aircraft payload for global observations of greenhouse gases and BC aerosol. It included several flights to the Arctic in the May-June time frame, bridging the two ARCTAS deployments.

The ARCTAS-CARB flights focused on satellite validation, sampling of emissions, and providing upwind boundary conditions and atmospheric observations to improve air quality models. DC-8 satellite validation primarily focused on TES (ozone, CO), OMI (NO₂), and MOPITT (CO), while P-3 satellite validation focused on MISR and MODIS (aerosols). Plumes from large wildfires in northern California were repeatedly sampled. These plumes were transported from northern to southern California, allowing opportunities to study their interactions with urban pollution. Extensive low-level sampling was made of emissions from ships, population centers, industrial centers, agricultural crops, feedlots, and dairy farms. Flights were directed to investigate transboundary

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



pollution from Asia and Mexico.

4 Conclusions

The ARCTAS mission offers a rich data set for better understanding of Arctic pollution including long-range transport, boreal forest fires influences, aerosol radiative forcing, and chemical processes. The chemical payload aboard the DC-8 aircraft included ozone and aerosols, a comprehensive suite of their precursors and related species, greenhouse gases, and pollution tracers. The in situ and remote payloads aboard the P-3 and B-200 aircraft characterized aerosol properties and associated radiative forcing. The ensemble of data collected during ARCTAS is of great value for improving chemical transport and climate models for the Arctic. It should also be of value as a baseline against which to measure future change. All ARCTAS data are publicly available from the NASA web site <http://www-air.larc.nasa.gov/missions/arctas/arctas.html>.

Although the ARCTAS deployments covered only two 3-week periods in April and June–July 2008, the conditions encountered were highly favorable for addressing mission objectives. The aircraft observed springtime pollution transport events from Asia, North America, and Europe, springtime fire plumes from Siberia and Southeast Asia, and summertime fire plumes from Canada, California, and Siberia. They encountered a number of boundary layer ozone depletion events in the spring. Unusually large fire activity in northern Saskatchewan during the summer deployment allowed detailed study of fire emissions and the near-field evolution of fire plumes.

A central objective of ARCTAS was to increase the value of satellite observations of Arctic atmospheric composition. Satellites offer a unique resource for long-term Arctic monitoring. ARCTAS provided (1) direct validation of satellite measurements, (2) in situ and remote data to improve retrieval algorithms, (3) correlative measurements of species not observable from space, and (4) information on chemical and aerosol processes needed to connect the satellite observations to models. We expect that the

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ARCTAS data will pave the way for better use of satellite observations to monitor and understand changes in Arctic atmospheric composition and its environmental implications.

The ARCTAS-CARB flights conducted just before the ARCTAS-B summer deployment characterized atmospheric composition across large regions of California with unprecedented detail. Information on emissions from a large number of sources such as refineries, urban centers, ships, dairies, feedlots, and agricultural operations was obtained. The aircraft sampled plumes from wildfires in northern California and these were sampled again when the aircraft operated in Canada, providing information on plume aging as well as comparison with boreal fire plumes. We expect that the ARCTAS-CARB data will provide an important foundation for improving models of air quality and greenhouse gas emissions in California.

Acknowledgements. This work was funded by the NASA Global Tropospheric Chemistry Program, the NASA Radiation Sciences Program, and the California Air Resources Board. We thank Kelly Chance and Charles Gatebe for commenting on the draft and Javier Fochesatto for the graphics in Fig. 3.

References

- Adhikary, B., Kulkarni, S., Dallura, A., et al.: A regional scale chemical transport modeling of Asian aerosols with data assimilation of AOD observations using optimal interpolation technique, *Atmos. Environ.*, 37, 8600–8615, 2008.
- Anderson, B. E., Cofer, W. R., Bagwell, D. R., et al.: Airborne observations of aircraft aerosol emissions 1: total nonvolatile particle emission indices, *Geophys. Res. Lett.*, 25, 1689–1692, 1998.
- Apel, E. C, Hills, A. J., Lueb, R., Zindel, S., Eisele, S., and Riemer, D. D.: A fast-GC/MS system to measure C-2 to C-4 carbonyls and methanol aboard aircraft, *J. Geophys. Res.*, 108, 8794, doi:10.1029/2002JD003199, 2003.
- Amiro, B. D., Todd, J. B., Wotton, B. M., Logan, K. A., Flannigan, M. D., Stocks, B. J., Mason,

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The ARCTAS aircraft mission: design and executionD. J. Jacob et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

J. A., Martell, D. L., and Hirsch, K. G.: Direct carbon emissions from Canadian forest fires, 1959–1999, *Can. J. Forest Res.*, 31, 512–525, 2001.

Arellano Jr., A. F., Raeder, K., Anderson, J. L., Hess, P. G., Emmons, L. K., Edwards, D. P., Pfister, G. G., Campos, T. L., and Sachse, G. W.: Evaluating model performance of an ensemble-based chemical data assimilation system during INTEX-B field mission, *Atmos. Chem. Phys.*, 7, 5695–5710, 2007, <http://www.atmos-chem-phys.net/7/5695/2007/>.

Atlas, E. L., Ridley, B. A., and Cantrell, C. A.: The Tropospheric Ozone Production about the Spring Equinox (TOPSE) experiment: introduction, *J. Geophys. Res.*, 108, 8353, doi:10.1029/2002JD003172, 2003.

Bergstrom, R. W., Pilewskie, P., Russell, P. B., Redemann, J., Bond, T. C., Quinn, P. K., and Sierau, B.: Spectral absorption properties of atmospheric aerosols, *Atmos. Chem. Phys.*, 7, 5937–5943, 2007, <http://www.atmos-chem-phys.net/7/5937/2007/>.

Blake, N. J., Blake, D. R., Simpson, I. J., et al: NMHCs and halocarbons in Asian continental outflow during the Transport and Chemical Evolution over the Pacific (TRACE-P) field campaign: comparison with PEM-West B, *J. Geophys. Res.*, 108, 8806, doi:10.1029/2002JD003367, 2003.

Bottenheim, J. W. and Chan, E.: A trajectory study into the origin of spring time Arctic boundary layer ozone depletion, *J. Geophys. Res.*, 111, D19301, doi:10.1029/2006JD007055, 2006.

Browell, E. V., Hair, J. W., Butler, C. F., et al.: Ozone, aerosol, potential vorticity, and trace gas trends observed at high-latitudes over North America from February to May 2000, *J. Geophys. Res.*, 108, 8369, doi:10.1029/2001JD001390, 2003.

Brune, W. H., Tan, D., Faloon, I. F., et al.: OH and HO₂ chemistry in the North Atlantic free troposphere, *Geophys. Res. Lett.*, 26, 3077–3080, 1999.

Bucholtz, A., Bluth, R. T., Kelly, B., et al.: The Stabilized Radiometer Platform (STRAP)-An Actively Stabilized Horizontally Level Platform for Improved Aircraft Irradiance Measurements, *J. Atmos. Ocean Tech.*, 25, 2161–2175, 2008.

Cantrell, C. A., Edwards, G. D., Stephens, S., et al.: Peroxy radical behavior during the Transport and Chemical Evolution over the Pacific (TRACE-P) campaign as measured aboard the NASA P-3B aircraft (2003), *J. Geophys. Res.*, 108, 8797, doi:10.1029/2002JD002715, 2003.

Chance, K.: Analysis of BrO measurements from the Global Ozone Monitoring Experiment,

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Geophys. Res. Lett., 25, 3335–3338, 1998.

Choi, Y., Wang, Y., Zeng, T., Cunnold, D., Yang, E.-S., Martin, R., Chance, K., Thouret, V., and Edgerton, E.: Springtime transitions of O₃, NO₂, and CO over North America: Model evaluation and analysis, *J. Geophys. Res.*, 113, D20311, doi:10.1029/2007JD009632, 2008.

5 Christensen, J. H., Hewitson, B., Busuioc, A., et al.: Regional Climate Projections, in: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M., et al., Cambridge University Press, Cambridge, UK and New York, NY, USA, 2007.

10 Chu, D. A.: Evaluation of aerosol properties over ocean from Moderate Imaging Spectroradiometer (MODIS) during ACE-Asia, *J. Geophys. Res.*, 110, D07308, doi:10.1029/2004JD005208, 2005.

Clarke, A. D. and Noone, K. J.: Soot in Arctic snow: a cause for perturbations of radiative transfer, *Atmos. Environ.*, 19, 2045–2053, 1985.

15 Clarke, A., McNaughton, C., Kapustin, V., et al.: Biomass burning and pollution aerosol over North America: Organic components and their influence on spectral optical properties and humidification response, *J. Geophys. Res.*, 112, D12S18, doi:10.1029/2006JD007777, 2007.

20 Cleary, P.A., Wooldridge, P.J., and R.C. Cohen: Laser-induced fluorescence detection of atmospheric NO₂ using a commercial diode laser and a supersonic expansion, *Applied Optics*, 41, 6950–6956, 2002.

Colarco, P. R., Schoeberl, M. R., Doddridge, B. G., Marufu, L. T., Torres, O., and Welton, E. J.: Transport of smoke from Canadian forest fires to the surface near Washington, D.C.: injection height, entrainment and optical properties, *J. Geophys. Res.*, 109, D06203, doi:10.1029/2003JD004248, 2004.

25 Cox, P., Delao, A., Komorniczak, A., and Weller, R.: *The California Almanac of emissions and air quality-2008 edition*, California Air Resources Board, Sacramento, California, 2008.

Crouse, J. D., DeCarlo, P. F., Blake, D. R., Emmons, L. K., Campos, T. L., Apel, E. C., Clarke, A. D., Weinheimer, A. J., McCabe, D. C., Yokelson, R. J., Jimenez, J. L., and Wennberg, P. O.: Biomass burning and urban air pollution over the Central Mexican Plateau, *Atmos. Chem. Phys.*, 9, 4929–4944, 2009, <http://www.atmos-chem-phys.net/9/4929/2009/>.

30 Crouse, J., McKinney, K. A., Kwan, A. J., et al.: Measurement of gas-phase hydroperoxides



- by chemical ionization mass spectrometry, *Anal. Chem.*, 78, 6726–6732, 2006.
- DeCarlo, P. F., Dunlea, E. J., Kimmel, J. R., Aiken, A. C., Sueper, D., Crounse, J., Wennberg, P. O., Emmons, L., Shinozuka, Y., Clarke, A., Zhou, J., Tomlinson, J., Collins, D. R., Knapp, D., Weinheimer, A. J., Montzka, D. D., Campos, T., and Jimenez, J. L.: Fast airborne aerosol size and chemistry measurements above Mexico City and Central Mexico during the MILAGRO campaign, *Atmos. Chem. Phys.*, 8, 4027–4048, 2008, <http://www.atmos-chem-phys.net/8/4027/2008/>.
- Deeter, M. N., Emmons, L. K., Francis, G. L., et al.: Operational carbon monoxide retrieval algorithm and selected results for the MOPITT instrument, *J. Geophys. Res.*, 108, 4399, doi:10.1029/2002JD003186, 2003.
- Deeter, M. N., Edwards, D. P., Gille, J. C., and Drummond, J. R.: CO retrievals based on MOPITT near-infrared observations, *J. Geophys. Res.*, 114, D04303, doi:10.1029/2008JD010872, 2009.
- Diskin, G. S., Podolske, J. R., Sachse, G. W., and Slate, T. A.: Open-Path Airborne Tunable Diode Laser Hygrometer, in *Diode Lasers and Applications in Atmospheric Sensing*, SPIE Proceedings 4817, A. Fried, ed., 196–204, 2002.
- Emmons, L. K., Hess, P., Klonecki, A., et al.: Budget of tropospheric ozone during TOPSE from two chemical transport models, *J. Geophys. Res.*, 108, 8372, doi:10.1029/2002JD002665, 2003.
- Emmons, L. K.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geosci. Model Dev.*, to be submitted, 2009.
- Engvall, A. C., Krejci, R., Storm, J., et al.: In situ airborne observation of the microphysical properties of the Arctic tropospheric aerosol during late spring and summer, *Tellus*, 60B, 392–404, 2008.
- Fenn, M. A., Browell, E. V., Butler, C. F., et al.: Ozone and aerosol distributions and air mass characteristics over the South Pacific during the burning season, *J. Geophys. Res.*, 104, 16197–16212, 1999.
- Fiebig, M., Petzold, A., Wandinger, U., et al.: Optical closure for an aerosol column: method, accuracy, and inferable properties, applied to a biomass burning aerosol and its radiative forcing, *J. Geophys. Res.*, 107, 8133, doi:10.1029/2000JD000192, 2002.
- Fiebig, M., Stohl, A., Wendisch, M., Eckhardt, S., and Petzold, A.: Dependence of solar radiative forcing of forest fire aerosol on ageing and state of mixture, *Atmos. Chem. Phys.*, 3, 881–891, 2003,

The ARCTAS aircraft mission: design and executionD. J. Jacob et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

<http://www.atmos-chem-phys.net/3/881/2003/>.

Fisher, J. A., Jacob, D. J., Kopacz, M. A., et al.: Source attribution and interannual variability of Arctic pollution in spring constrained by aircraft (ARCTAS, ARCPAC) and satellite (AIRS) observations of carbon monoxide, *Atmos. Chem. Phys.*, submitted, 2009.

5 Flanner, M. G., Zender, C. S., Randerson, J. T., and Rasch, P. J.: Present-day radiative forcing and climate response from black carbon in snow, *J. Geophys. Res.*, 112, D11202, doi:10.1029/2006JD008003, 2007.

Flannigan, M. D., Logan, K. A., Amiro, B. D., Skinner, W. R., and Stocks, B. J.: Future area burned in Canada, *Climatic Change*, 72, 1–16, 2005.

10 Fried, A., Walega, J. G., Olson, J. R., et al.: Formaldehyde over North America and the North Atlantic during the summer 2004 INTEX campaign: methods, observed distributions, and measurement-model comparisons, *J. Geophys. Res.*, 113, D10302, doi:10.1029/2007JD009185, 2008.

Fromm, M. D. and Servranckx, R.: Transport of forest fire smoke above the tropopause by supercell convection, *Geophys. Res. Lett.*, 30, 1542, doi:10.1029/2002GL016820, 2003.

Fuelberg, H. E., Harrigan, D. L., and Sessions, W.: A meteorological overview of the ARCTAS 2008 mission, *Atmos. Chem. Phys.*, submitted, 2009.

Fuelberg, H. E., Kiley, C., Hannan, J. R., Westberg, D. J., Avery, M. A., and Newell, R. E.: Meteorological conditions and transport pathways during the transport and chemical evolution over the Pacific (TRACE-P) experiment, *J. Geophys. Res.*, 108, 8782, doi:10.1029/2002JD003092, 2003.

20 Gatebe, C. K., King, M. D., Platnick, S., Arnold, G. T., Vermote, E. F., and Schmid, B.: Airborne spectral measurements of surface-atmosphere anisotropy for several surfaces and ecosystems over southern Africa, *J. Geophys. Res.*, 108, 8489, doi:10.1029/2002JD002397, 2003.

25 Generoso, S., Bey, I., Attie, J.-L., and Breon, F.-M.: A satellite- and model-based assessment of the 2003 Russian forest fires: impact on the Arctic region, *J. Geophys. Res.*, 112, D15302, doi:10.1029/2006JD008344, 2007.

30 Gillett, N. P., Weaver, A. J., Zwiers, F. W., and Flannigan, M. D.: Detecting the effect of climate change on Canadian forest fires, *Geophys. Res. Lett.*, 31, L18211, doi:10.1029/2004GL020876, 2004.

Girardin, M. P.: Interannual to decadal changes in area burned in Canada from 1781 to 1982 and the relationship to Northern Hemisphere land temperatures, *Global Ecol. Biogeogr.*, 16, 557–566, 2007.

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The ARCTAS aircraft mission: design and executionD. J. Jacob et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Goode, J. G., Yokelson, R. J., Ward D. E., et al.: Measurements of excess O₃, CO₂, CO, CH₄, C₂H₄, C₂H₂, HCN, NO, NH₃, HCOOH, CH₃COOH, HCHO, and CH₃OH in 1997 Alaskan biomass burning plumes by airborne Fourier transform infrared spectroscopy (AFTIR), *J. Geophys. Res.*, 105, 22147–22166, 2000.
- 5 Hair, J. W., Hostetler, C. A., Cook, A. L., Harper, D. B., Ferrare, R. A., Mack, T. L., Welch, W., Izquierdo, L. R., and Hovis, F. E.: Airborne High Spectral Resolution Lidar for profiling aerosol optical properties, *Applied Optics*, 47, 6734–6752, doi:10.1364/AO.47.006734, 2008.
- Hansen, J. and Nazarenko, L.: Soot climate forcing via snow and ice albedos, *P. Natl. Acad. Sci.*, 101, 423–428, 2004.
- 10 Hirdman, D., Apsmo, K., Burkhardt, J. F., et al.: Transport of mercury in the Arctic atmosphere: Evidence for a springtime net sink and summer-time source, *Geophys. Res. Lett.*, 36, L12814, doi:10.1029/2009GL038345, 2009.
- Hoell, J. M., Davis, D. D., Jacob, D. J., et al.: Pacific Exploratory Mission – West Phase B: February–March 1994, *J. Geophys. Res.*, 102, 28223–28240, 1997.
- 15 Hollwedel, J., Wenig, M., Beirle, S., et al.: Year-ro-year variations of springtime polar tropospheric BrO as seen by GOME, *Adv. Space Res.*, 34, 804–808, 2004.
- Hyer, E. J., Allen, D. J., and Kasischke, E. S.: Examining injection properties of boreal forest fires using surface and satellite measurements of CO transport, *J. Geophys. Res.*, 112, D18307, doi:10.1029/2006JD008232, 2007.
- 20 Jacob, D. J., Wofsy, S. C., Bakwin, P. S., Fan, S.-M., Harriss, R. C., Talbot, R. W., Bradshaw, J., Sandholm, S., Singh, H. B., Gregory, G. L., Browell, E. V., Sachse, G. W., Blake, D. R., and Fitzjarrald, D. R.: Summertime photochemistry at high northern latitudes, *J. Geophys. Res.*, 97, 16421–16431, 1992.
- Jacob, D. J., Logan, J. A., and Murti, P. P.: Effect of rising Asian emissions on surface ozone in the United States, *Geophys. Res. Lett.*, 26(14), 2175–2178, 1999.
- 25 Jacob, D. J., Crawford, J. H., Kleb, M. M., Connors, V. S., Bendura, R. J., Raper, J. L., Sachse, G. W., Gille, J. C., Emmons, L., and Heald, C. L.: The Transport and Chemical Evolution over the Pacific (TRACE-P) aircraft mission: design, execution, and first results, *J. Geophys. Res.*, 108, 9000, doi:10.1029/2002JD003276, 2003.
- 30 Kahn, R., Gaitley, B., Martonchik, J., Diner, D., Crean, K., and Holben, B.: MISR global aerosol optical depth validation based on two years of coincident AERONET observations, *J. Geophys. Res.*, 110, D10S04, doi:10.1029/2004JD004706, 2005.
- Kahn, R. A., Li, W.-H., Moroney, C., Diner, D. J., Martonchik, J. V., and Fishbein, E.: Aerosol

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

source plume physical characteristics from space-based multiangle imaging, *J. Geophys. Res.*, 112, D11205, doi:10.1029/2006JD007647, 2007.

Kim, S., Huey, L. G., Stickel, R. E. J., Tanner, D. J., Crawford, J. H., Olson, J. R., Chen, G., Brune, W. H., Ren, X., Leshner, R., Wooldridge, P. J., Bertram, T. H., Perring, A., Cohen, R. C.,
5 Lefer, B., Shetter, R. E., Avery, M., Diskin, G., and Sokolik, I.: Measurement of pernitric acid in the free troposphere during the intercontinental chemical transport experiment – North America 2004, *J. Geophys. Res.*, 112, D12S01, doi:10.1029/2006JD007676, 2007.

Kivi, R., Kyro, E., Turunen, T., et al.: Ozone-sonde observations in the Arctic during 1989–2003: ozone variability and trends in the lower stratosphere and free troposphere, *J. Geophys. Res.*, 112, D08306, doi:10.1029/2006JD007271, 2007.

Klonecki, A., Hess, P., Emmons, L., Smith, L., and Orlando, J.: Seasonal changes in the transport of pollutants into the Arctic troposphere – model study, *J. Geophys. Res.*, 108, 8367, doi:10.1029/2002JD002199, 2003.

Koch, D. and Hansen, J.: Distant origins of Arctic black carbon: a Goddard Institute of Space Studies Model E experiment, *J. Geophys. Res.*, 110, D04204, doi:10.1029/2004JD005296, 2005.

Koike, M., Kondo, Y., Kita, K., et al.: Export of anthropogenic reactive nitrogen and sulfur compounds from the east Asia region in spring, *J. Geophys. Res.*, 108, 8789, doi:10.1029/2002JD003284, 2003.

Korhonen, H., Carslaw, K. S., Spracklen, D. V., Ridley, D. A., and Strom, J.: A global model study of processes controlling aerosol size distributions in the Arctic spring and summer, *J. Geophys. Res.*, 113, D08211, doi:10.1029/2007JD009114, 2008.

Kylling, A., Webb, A. R., Kift, R., Gobbi, G. P., Ammannato, L., Barnaba, F., Bais, A., Kazadzis, S., Wendisch, M., Jäkel, E., Schmidt, S., Kniffka, A., Thiel, S., Junkermann, W., Blumthaler, M., Silbernagl, R., Schallhart, B., Schmitt, R., Kjeldstad, B., Thorseth, T. M., Scheirer, R., and Mayer, B.: Spectral actinic flux in the lower troposphere: measurement and 1-D simulations for cloudless, broken cloud and overcast situations, *Atmos. Chem. Phys.*, 5, 1975–1997, 2005,
http://www.atmos-chem-phys.net/5/1975/2005/.

30 Labonne, M., Breon, F.-M., and Chevallier, F.: Injection height of biomass burning aerosols as seen from a spaceborne lidar, *Geophys. Res. Lett.*, 34, L11806, doi:10.1029/2007GL029311, 2007.

Lavoue, D., Liousse, C., Cachier, H., Stocks, B. J., and Goldammer, J. G.: Modeling of carbona-

ceous particles emitted by boreal and temperate wildfires at northern latitudes, *J. Geophys. Res.*, 105, 26871–26890, 2000.

Lemke, P., Ren, J., Alley, R. B., Allison, I., Carrasco, J., Flato, G., Fujii, Y., Kaser, G., Mote, P., Thomas, R. H., and Zhang, T.: in *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, UK and New York, NY, USA, 2007.

Leung, F.-Y. T., Logan, J. A., Park, R., et al.: Impacts of enhanced biomass burning in the boreal forests in 1998 on tropospheric chemistry and the sensitivity of model results to the injection height of emissions, *J. Geophys. Res.*, 112, D10313, doi:10.1029/2006JD007281, 2007.

Levelt, P. F., Hilsenrath, E., Leppenmeier, G. W., van den Oord, G. H. J., Bhartia, P. K., Tamminen, J., de Haan, J. F., and Veefkind, J. P.: Science objectives of the Ozone Monitoring Instrument, *IEEE T. Geosci Remote*, 44, 1199–1208, 2006.

Liang, Q., Douglass, A. R., Duncan, B. N., Stolarski, R. S., and Witte, J. C.: The governing processes and timescales of stratosphere-to-troposphere transport and its contribution to ozone in the Arctic troposphere, *Atmos. Chem. Phys.*, 9, 3011–3025, 2009, <http://www.atmos-chem-phys.net/9/3011/2009/>.

Livesey, N. J., Filipiak, M. J., Froidevaux, L., et al.: Validation of Aura Microwave Limb Sounder O₃ and CO observations in the upper troposphere and lower stratosphere, *J. Geophys. Res.*, 113, D15S02, doi:10.1029/2007JD008805, 2008.

McMillan, W. W., Warner, J. X., Corner, M. M., et al.: AIRS views transport from 12 to 22 July 2004 Alaskan/Canadian fires: correlation of AIRS CO and MODIS AOD with forward trajectories and comparison of AIRS CO retrievals with DC-8 in situ measurements during INTEX-A/ICARTT, *J. Geophys. Res.*, 113, D20301, doi:10.1029/2007JD009711, 2008.

McNaughton, C. S., Thornhill, L., Clarke, A. D., Howell, S. G., Pinkerton, M., Anderson, B., Winstead, E., Hudgins, C., Maring, H., Dibb, J. E., and Scheuer, E.: Results from the DC-8 inlet characterization experiment (DICE): Airborne versus surface sampling of mineral dust and sea salt aerosols, *Aerosol Sci. Tech.*, 40, 136–159, 2007.

Medina, J., Nenes, A., Sotiropolou, R. E. P., et al.: Cloud condensation nuclei closure during the International Consortium for Atmospheric Research on Transport and Transformation 2004 campaign: Effects of size-resolved composition, *J. Geophys. Res.*, 112, D10S31, doi:10.1029/2006JD007588, 2007.

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The ARCTAS aircraft mission: design and executionD. J. Jacob et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Morris, G. A., Hersey, S., Thompson, A. M., Stohl, A., Colarco, P. R., McMillan, W. W., Warner, J., Johnson, B. J., Witte, J. C., Kucsera, T. L., Larko, D. E., and Oltmans, S. J.: Alaskan and Canadian forest fires exacerbate ozone pollution in Houston, Texas, on 19 and 20 July 2004, *J. Geophys. Res.*, 111, D24S03, doi:10.1029/2006JD007441, 2006.

5 Moteki, N. and Kondo, Y.: Effects of mixing state on black carbon measurements by Laser-Induced Incandescence, *Aerosol Sci. Tech.*, 41, 398–417, 2007.

Moteki, N. and Kondo, Y.: Method to measure time-dependent scattering cross sections of particles evaporating in a laser beam, *J. Aerosol Sci.*, 39, 348–364, 2008.

10 Munro, R., Eisinger, M., Anderson, C., Callies, J., Corpaccioli, E., Lang, R., Lefebvre, A., Livschitz, Y., and Albiñana, A. P.: GOME-2 on MetOp, Proc. of The 2006 EUMETSAT Meteorological Satellite Conference, Helsinki, Finland, 12–16 June 2006, EUMETSAT, p. 48, 2006.

Miyazaki, Y., Kondo, Y., Koike, M., et al.: Synoptic-scale transport of reactive nitrogen over the western Pacific in spring, *J. Geophys. Res.*, 108, 8788, doi:10.1029/2002JD003248, 2003.

15 Monks, P. S.: A review of the observations and origins of the spring ozone maximum, *Atmos. Environ.*, 34, 3545–3561, 2000.

Olson, J. R., Crawford, J. H., Chen, G., Brune, W. H., Faloon, I. C., Tan, D., Harder, H., and Martinez, M.: A reevaluation of airborne HO_x observations from NASA field campaigns, *J. Geophys. Res.*, 111, D10301, doi:10.1029/2005JD006617, 2006.

20 Oltmans, S. J., Lefohn, A. S., Harris, J. M., and Shadwick, D. S.: Background ozone levels of air entering the west coast of the US and assessment of longer-term changes, *Atmos. Environ.*, 42, 6020–6038, 2008.

O'Neill, N. T., Eck, T. F., Holben, B. N., Smirnov, A., Royer, A., and Li, Z.: Optical properties of boreal forest fire smoke derived from Sun photometry, *J. Geophys. Res.*, 107, 4125, doi:10.1029/2001JD000877, 2002.

25 Park, R. J., Jacob, D. J., Palmer, P. I., Clarke, A. D., Weber, R. J., Zondlo, M. A., Eisele, F. L., Bandy, A. R., Thornton, D. C., Sachse, G. W., and Bond, T. C.: Export efficiency of black carbon aerosol in continental outflow: global implications, *J. Geophys. Res.*, 110, D11205, doi:10.1029/2004JD005432, 2005.

30 Pfister, G. G., Hess, P. G., Emmons, L. K., Rasch, P. J., and Vitt, F. M.: Impact of the summer 2004 Alaskan fires on top of the atmosphere clear-sky radiation fluxes, *J. Geophys. Res.*, 113, D02204, doi:10.1029/2007JD008797, 2008.

Preston, C. M. and Schmidt, M. W. I.: Black (pyrogenic) carbon: a synthesis of current knowl-

edge and uncertainties with special consideration of boreal regions, *Biogeosciences*, 3, 397–420, 2006,

<http://www.biogeosciences.net/3/397/2006/>.

Quinn, P. K., Shaw, G., Andrews, E., Dutton, E. G., Ruoho-Airola, T., and Gong, S. L.: Arctic haze: current trends and knowledge gaps. *Tellus*, 59B, 99–114, 2007.

Quinn, P. K., Bates, T. S., Baum, E., Doubleday, N., Fiore, A. M., Flanner, M., Fridlind, A., Garrett, T. J., Koch, D., Menon, S., Shindell, D., Stohl, A., and Warren, S. G.: Short-lived pollutants in the Arctic: their climate impact and possible mitigation strategies, *Atmos. Chem. Phys.*, 8, 1723–1735, 2008,

<http://www.atmos-chem-phys.net/8/1723/2008/>.

Real, E., Law, K., Weinzierl, B., et al.: Processes influencing ozone in Alaskan forest fire plumes during long-range transport over the North Atlantic, *J. Geophys. Res.*, 112, D10S41, doi:10.1029/2006JD007576, 2007.

Reid, J. S., Koppmann, R., Eck, T. F., and Eleuterio, D. P.: A review of biomass burning emissions part II: intensive physical properties of biomass burning particles, *Atmos. Chem. Phys.*, 5, 799–825, 2005,

<http://www.atmos-chem-phys.net/5/799/2005/>.

Ren, X., Olson, J. R., Crawford, J. H., et al.: HO_x chemistry during INTEX-A 2004: observation, model calculation, and comparison with previous studies, *J. Geophys. Res.*, 113, D05310, doi:10.1029/2007JD009166, 2008.

Ridley, B. A., Atlas, E. L., Montzka, D. D., et al.: Ozone depletion events in the high latitude surface layer during the TOPSE aircraft program, *J. Geophys. Res.*, 108, 8356, doi:10.1029/2001JD001507, 2003.

Russell, P. B., Livingston, J. M., Redemann, J., et al.: Multi-grid-cell validation of satellite aerosol property retrievals in INTEX/ITCT/ICARTT 2004, *J. Geophys. Res.*, 112, D12S09, doi:10.1029/2006JD007606, 2007.

Sachse, G. W., Hill, G. F., Wade, L. O., and Perry, M. G.: Fast-response, high-precision carbon monoxide sensor using a tunable diode laser absorption technique, *J. Geophys. Res.*, 92, 2071–2081, 1987.

Scheuer, E., Talbot, R. W., Dibb, J. E., Seid, G. K., deBell, L., and Lefer, G.: Seasonal distributions of fine aerosol sulfate in the North American Arctic Basin during TOPSE, *J. Geophys. Res.*, 108, 8370, doi:10.1029/2001JD001364, 2003.

Schoeberl, M. R.: A trajectory-based estimate of the tropospheric ozone column using the

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- residual method, *J. Geophys. Res.*, 112, D24S49, doi:10.1029/2007JD008773, 2007.
- Shaw, G. E.: The Arctic haze phenomenon, *B. Am. Meteorol. Soc.*, 2403–2413, 1995.
- Shetter, R. E. and Muller, M.: Photolysis frequency measurements on the NASA DC-8 during the PEM-Tropics Mission using actinic flux spectroradiometry: Instrument description and result (1999), *J. Geophys. Res.*, 104, 5647–5661, 1999.
- 5 Shindell, D. T., Faluvegi, G., Bauer, S. E., et al.: Climate response to projected changes in short-lived species under an A1B scenario from 2000–2050 in the GISS climate model, *J. Geophys. Res.*, 112, D20103, doi:10.1029/2007JD008753, 2007.
- Shindell, D. T., Chin, M., Dentener, F., Doherty, R. M., Faluvegi, G., Fiore, A. M., Hess, P., Koch, D. M., MacKenzie, I. A., Sanderson, M. G., Schultz, M. G., Schulz, M., Stevenson, D. S., Teich, H., Textor, C., Wild, O., Bergmann, D. J., Bey, I., Bian, H., Cuvelier, C., Duncan, B. N., Folberth, G., Horowitz, L. W., Jonson, J., Kaminski, J. W., Marmer, E., Park, R., Pringle, K. J., Schroeder, S., Szopa, S., Takemura, T., Zeng, G., Keating, T. J., and Zuber, A.: A multi-model assessment of pollution transport to the Arctic, *Atmos. Chem. Phys.*, 8, 5353–5372, 2008, <http://www.atmos-chem-phys.net/8/5353/2008/>.
- 15 Shindell, D. and Faluvegi, G.: Climate response to regional radiative forcing during the twentieth century, *Nature Geosci.*, 2, 294–300, 2009.
- Simpson, W. R., von Glasow, R., Riedel, K., Anderson, P., Ariya, P., Bottenheim, J., Burrows, J., Carpenter, L. J., Frieß, U., Goodsite, M. E., Heard, D., Hutterli, M., Jacobi, H.-W., Kaleschke, L., Neff, B., Plane, J., Platt, U., Richter, A., Roscoe, H., Sander, R., Shepson, P., Sodeau, J., Steffen, A., Wagner, T., and Wolff, E.: Halogens and their role in polar boundary-layer ozone depletion, *Atmos. Chem. Phys.*, 7, 4375–4418, 2007a, <http://www.atmos-chem-phys.net/7/4375/2007/>.
- 20 Simpson, W. R., Carlson, D., Hönninger, G., Douglas, T. A., Sturm, M., Perovich, D., and Platt, U.: First-year sea-ice contact predicts bromine monoxide (BrO) levels at Barrow, Alaska better than potential frost flower contact, *Atmos. Chem. Phys.*, 7, 621–627, 2007b, <http://www.atmos-chem-phys.net/7/621/2007/>.
- Singh, H. B., Brune, W. H., Crawford, J. H., Flocke, F., and Jacob, D. J.: Chemistry and transport of pollution over the Gulf of Mexico and the Pacific: spring 2006 INTEX-B campaign overview and first results, *Atmos. Chem. Phys.*, 9, 2301–2318, 2009, <http://www.atmos-chem-phys.net/9/2301/2009/>.
- 30 Slusher, D. L., Huey, L. G., Tanner, D. J., Flocke, F. M., and Roberts, J. M.: A thermal

The ARCTAS aircraft mission: design and executionD. J. Jacob et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The ARCTAS aircraft mission: design and executionD. J. Jacob et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

dissociation-chemical ionization mass spectrometry (TD-CIMS) technique for the simultaneous measurement of peroxyacyl nitrates and dinitrogen pentoxide, *J. Geophys. Res.*, 109, D19315, doi:10.1029/2004JD004670, 2004.

5 Soja, A. J., Tchepakova, N. M., French, N. H. F., et al.: Climate-induced boreal forest change: predictions versus current observations, *Global Planet. Change*, 56, 274–296, 2007.

Soja, A. J., Stocks, B., Maczeck, P., Fromm, M., Servranckx, R., and Turetsky, M., and Benscoter, B.: ARCTAS: the perfect smoke, *The Canadian Smoke Newsletter*, Fall 2008, 2–7, 2008.

10 Steffen, A., Douglas, T., Amyot, M., Ariya, P., Aspmo, K., Berg, T., Bottenheim, J., Brooks, S., Cobbett, F., Dastoor, A., Dommergue, A., Ebinghaus, R., Ferrari, C., Gardfeldt, K., Goodsite, M. E., Lean, D., Poulain, A. J., Scherz, C., Skov, H., Sommar, J., and Temme, C.: A synthesis of atmospheric mercury depletion event chemistry in the atmosphere and snow, *Atmos. Chem. Phys.*, 8, 1445–1482, 2008,

<http://www.atmos-chem-phys.net/8/1445/2008/>.

15 Stocks, B. J., Fosberg, M. A., Lynham, T. J., et al.: Climate change and forest fire potential in Russian and Canadian boreal forests, *Climatic Change*, 38, 1–13, 1998.

Stohl, A.: Characteristics of atmospheric transport into the Arctic troposphere, *J. Geophys. Res.*, 111, D11306, doi:10.1029/2005JD006888, 2006.

20 Stone, R. S., Anderson, G. P., Shettle, E. P., et al.: Radiative impact of boreal smoke in the Arctic: observed and modeled, *J. Geophys. Res.*, 113, D14S16, doi:10.1029/2007JD009657, 2008.

Strawa, A. W., Elleman, R., Hallar, A. G., et al.: Comparison of in situ aerosol extinction and scattering coefficient measurements made during the Aerosol Intensive Operating Period, *J. Geophys. Res.*, 111, D05S03, doi:10.1029/2005JD006056, 2006.

25 Streets, D. G., Zhang, Q., Wang, L. T., et al.: Revisiting China's CO emissions after the Transport and Chemical Evolution over the Pacific (TRACE-P) mission: Synthesis of inventories, atmospheric modeling, and observations, *J. Geophys. Res.*, 111, D14306, doi:10.1029/2002JD003093, 2006.

30 Sullivan, A., Peltier, R. E., Brock, C. A., de Gouw, J. A., Holloway, J. S., Warneke, C., Wollny, A. G., and Weber, R. J.: Airborne measurements of carbonaceous aerosol soluble in water over northeastern United States: Method development and an investigation into water-soluble organic carbon sources, *J. Geophys. Res.*, 111, D23S46, doi:10.1029/2006JD007072, 2006.

Talbot, R., Mao, H., Scheuer, E., Dibb, J., Avery, M., Browell, E., Sachse, G., Vay, S., Blake,

D., Huey, G., and Fuelberg, H.: Factors influencing the large-scale distribution of Hg^o in the Mexico City area and over the North Pacific, *Atmos. Chem. Phys.*, 8, 2103–2114, 2008, <http://www.atmos-chem-phys.net/8/2103/2008/>.

5 Tarasick, D. W., Fioletov, V. E., Wardle, D. I., Kerr, J. B., and Davies, J.: Changes in the vertical distribution of ozone over Canada from ozonesondes: 1980–2001, *J. Geophys. Res.*, 110, D02304, doi:10.1029/2004JD004643, 2005.

Tarasick, D. W., Moran, M. D., Thompson, A. M., et al.: Comparison of Canadian air quality forecast models with tropospheric ozone profile measurements above mid-latitude North America during the IONS/ICARTT Campaign: Evidence for stratospheric input, *J. Geophys. Res.*, 112, D12S22, doi:10.1029/2006JD007782, 2007.

10 Thompson, A. M., Stone, J. B., Witte, J. C., et al.: Intercontinental Transport Experiment Ozone Network Study (IONS, 2004): 2. Tropospheric ozone budgets and variability over northeastern North America, *J. Geophys. Res.*, 112, D12S13, doi:10.1029/2006JD007670, 2007.

15 Trenberth, K., Jones, P. D., Ambenje, P., et al.: *Climate Change 2007: The Physical Science Basis*. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Solomon, S., Qin, D., Manning, M., et al., Cambridge University Press, Cambridge, UK and New York, NY, USA, 2007.

20 Turetsky, M. R., Harden, J. W., Friedli, H. R., Flannigan, M., Payne, N., Crock, J., and Radke, L.: Wildfires threaten mercury stocks in northern soils, *Geophys. Res. Lett.*, 33, L16403, doi:10.1029/2005GL025595, 2006.

Turquety, S., Logan, J. A., Jacob, D. J., Hudman, R. C., Leung, F. Y., Heald, C. L., Yantosca, R. M., Wu, S., Emmons, L. K., Edwards, D. P., and Sachse, G. W.: Inventory of boreal fire emissions for North America in 2004: the importance of peat burning and pyro-convective injection, *J. Geophys. Res.*, 112, D12S03, doi:10.1029/2006JD007281, 2007.

25 Vay, S. A., Woo, J.-H., Anderson, B. E., Thornhill, K. L., Blake, D. R., Westberg, D. J., Kiley, C. M., Avery, M. A., Sachse, G. W., Streets, D., Tsutsumi, Y., and Nolf, S.: The influence of regional-scale anthropogenic emissions on CO₂ distributions over the western North Pacific, *J. Geophys. Res.*, 108, 8801, doi:10.1029/2002JD003094, 2003.

30 Wang, H., Jacob, D. J., Le Sager, P., Streets, D. G., Park, R. J., Gilliland, A. B., and van Donkelaar, A.: Surface ozone background in the United States: Canadian and Mexican pollution influences, *Atmos. Environ.*, 43, 1310–1319, 2009.

Waquet, F., Cairns, B., Knobelspiesse, K., Chowdhary, J., Travis, L. D., Schmid, B., and

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The ARCTAS aircraft mission: design and executionD. J. Jacob et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Mishchenko, M. I.: Polarimetric remote sensing of aerosols over land, *J. Geophys. Res.*, 114, D01206, doi:10.1029/2008JD010619, 2009.

Warneke, C., Bahreini, R., Brioude, J., et al: Biomass burning in Siberia and Kazakhstan as an important source for haze over the Alaskan Arctic in April 2008, *Geophys. Res. Lett.*, 36, L02813, doi:10.1029/2008GL036194, 2009.

Weinheimer, A. J., Walega, J. G., Ridley, B. A., Gary, B. L., Blake, D. R., Blake, N. J., Rowland, F. S., Sachse, G. W., Anderson, B. E., and Collins, J. E.: Meridional distributions of NO_x, NO_y, and other species in the lower stratosphere and upper troposphere during AASE II, *Geophys. Res. Lett.*, 21, 2583–2586, 1994.

Westphal, D. L. and Toon, O. B.: Simulations of microphysical, radiative, and dynamical processes in a continental-scale forest fire smoke plume, *J. Geophys. Res.*, 96, 22379–22400, 1991.

Wetzel, M. A., Shaw, G. E., Slusser, J. R., Borys, R. D., and Cahill, C. F.: Physical, chemical, and ultraviolet radiative characteristics of aerosol in central Alaska, *J. Geophys. Res.*, 108, 4418, doi:10.1029/2002JD003208, 2003.

Winker, D.W., W.H. Hunt, and M.J. McGill: Initial performance assessment of CALIOP. *Geophys. Res. Lett.*, 34, L19803, 2007.

Wisthaler, A., Hansel, A., Dickerson, R. R., et al.: Organic trace gas measurements by PTR-MS during INDOEX 1999, *J. Geophys. Res.*, 107, 8024, doi:10.1029/2001JD000576, 2002.

Worden, H. M., Bowman, K. W., Worden, J. R., Eldering, A., and Beer, R.: Satellite measurements of the clear-sky greenhouse effect from tropospheric ozone, *Nature Geosci.*, 1, 305–308, 2008.

Zhang, L., Jacob, D. J., Boersma, K. F., Jaffe, D. A., Olson, J. R., Bowman, K. W., Worden, J. R., Thompson, A. M., Avery, M. A., Cohen, R. C., Dibb, J. E., Flock, F. M., Fuelberg, H. E., Huey, L. G., McMillan, W. W., Singh, H. B., and Weinheimer, A. J.: Transpacific transport of ozone pollution and the effect of recent Asian emission increases on air quality in North America: an integrated analysis using satellite, aircraft, ozonesonde, and surface observations, *Atmos. Chem. Phys.*, 8, 6117–6136, 2008, <http://www.atmos-chem-phys.net/8/6117/2008/>.

Zeng, T., Wang, Y., Chance, K., et al.: Halogen-driven low-altitude O₃ and hydrocarbon losses in spring at northern high latitudes. *J. Geophys. Res.*, 111, D17313, doi:10.1029/2005JD006706, 2006.



The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Satellite measurements of atmospheric composition during ARCTAS¹.

Instrument (platform, launch year)	Spectral range ²	Relevant measurements	Reference
GOME-2 (MetOp, 2006)	UV/Vis	Tropospheric ozone, BrO, NO ₂ , formaldehyde, glyoxal	Munro et al. (2007)
CALIOP (Calipso, 2006)	Vis/NIR	Vertical distribution of aerosol optical depth	Winker et al. (2007)
OMI (Aura, 2004)	UV/Vis	Tropospheric ozone, BrO, NO ₂ , formaldehyde, glyoxal, absorbing aerosol	Levelt et al. (2006)
TES (Aura, 2004)	TIR	Tropospheric ozone, CO, methane	Worden et al. (2008)
MLS (Aura, 2004)	Microwave	Upper tropospheric ozone, CO	Livesey et al. (2008)
MODIS (Aqua, 2002; Terra, 1999)	UV/Vis/NIR	Aerosol optical depth and size classes	Chu et al. (2005)
AIRS (Aqua, 2002)	TIR	CO, methane	McMillan et al. (2008)
SCIAMACHY (Envisat, 2002)	UV/Vis/NIR	Tropospheric BrO, NO ₂ , formaldehyde, glyoxal, CO, methane	Sioris et al. (2006)
MOPITT (Terra, 1999)	NIR/TIR	CO	Deeter et al. (2003, 2004)
MISR (Terra, 1999)	Vis/NIR	Aerosol optical depth, injection heights	Kahn et al. (2005, 2007)

¹ All instruments are on polar-orbiting satellites. The Calipso, Aura, and Aqua satellites are on the A-train orbit with overpass at 13:30 local time (LT); other overpass times are 10:45 LT for Envisat and 10:30 LT for Terra. All instruments observe in the nadir only except MLS (limb) and SCIAMACHY (nadir and limb).

² NIR=near (solar) IR; TIR=thermal IR. All instruments are passive radiometers except CALIOP (active lidar).

The ARCTAS aircraft mission: design and executionD. J. Jacob et al.

Table 2. Operational characteristics of the ARCTAS aircraft.

	DC-8	P-3	B-200
Ceiling, km	12	7	9
Ground speed, knots	240–480	270–330	80–290
Endurance, h	10	8	5
Payload, tons	20	7	0.5

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3a. DC-8 research payload in ARCTAS.

Species, parameters ¹	Method ²	Investigator	Reference
CO, CH ₄ , N ₂ O, H ₂ O	TDLAS	G. Diskin, NASA LaRC	Sachse et al. (1987), Diskin et al. (2002)
CO ₂	Non-dispersive IR analyzer	S. Vay, NASA LaRC	Vay et al. (2003)
Ozone, NO, NO ₂ , NO _y	Chemiluminescence	A. Weinheimer, NCAR	Weinheimer et al. (1994)
Ozone and aerosol vertical distribution (remote)	DIAL	E. Browell, NASA LaRC	Fenn et al. (1999)
NO ₂ , PANs, alkyl nitrates	TD-LIF	R. Cohen, UC Berkeley	Cleary et al. (2002)
HNO ₃ , soluble Br, aerosol ionic composition	mist chamber/IC	J. Dibb, U. New Hampshire	Scheuer et al. (2003), McNaughton et al. (2007)
HNO ₃ , peroxides, HCN, organic acids	CIMS	P. Wennberg, Caltech	Crounse et al. (2006, 2009)
PANs, HNO ₄ , SO ₂ , halogens	CIMS	G. Huey, Georgia Tech	Slusher et al. (2004), Kim et al. (2007)
OH, HO ₂	LIF	W. Brune, Penn State	Brune et al. (1999)
OH, HO ₂ , RO ₂ , H ₂ SO ₄	CIMS	C. Cantrell, NCAR	Cantrell et al. (2003)
HCHO	TDLAS	A. Fried, NCAR	Fried et al. (2008)
NMHCs, halocarbons, alkyl nitrates	WAS-GC	D. Blake, UC Irvine	Blake et al. (2003)
NMHCs, OVOCs, halocarbons, CH ₃ CN	GC-MS	E. Apel, NCAR; D. Riemer, U. Miami	Apel et al. (2003)
Organic gases	PTR-MS	A. Wisthaler, U. Innsbruck	Wisthaler et al. (2002)
Aerosol number, size distribution, volatility, optical properties, CCN, EC/OC, cloud microphysics, LWC	CCN counter, OPC, aethelometer, nephelometer	B. Anderson, NASA LaRC	Anderson et al. (1998)
Size-resolved aerosol composition, volatility	HR-ToF-AMS	J.L. Jimenez, U. Colorado	DeCarlo et al. (2008)
Aerosol inorganic and organic composition	PILS	R. Weber, Georgia Tech	Sullivan et al. (2006)
Black carbon aerosol	SP2	Y. Kondo, U. Tokyo	Moteki and Kondo (2007, 2008)
Elemental, reactive, and particulate mercury	AFS	R. Talbot, U. New Hampshire	Talbot et al. (2008)
Actinic fluxes and photolytic frequencies	spectral radiometry	R. Shetter, NCAR	Shetter and Miller (1999)

¹ Chemical concentrations unless otherwise indicated. NO_y=total reactive nitrogen oxides, including NO_x and its oxidation products; PANs=peroxyacetyl nitrates; NMHCs=nonmethane hydrocarbons; OVOCs=oxygenated volatile organic compounds; CCN=cloud condensation nuclei; EC=elemental carbon; OC=organic carbon; LWC=liquid water content.

² DIAL=Differential Absorption Lidar; TD-LIF=Thermal-Dissociation Laser Induced Fluorescence; CIMS=Chemical Ionization Mass Spectrometry; TD-LAS=Tunable Diode Laser Absorption Spectroscopy; WAS-GC=Whole Air Sampling – Gas Chromatography; GC-MS=Gas Chromatography – Mass Spectrometry; PTR-MS=Proton Transfer Reaction – Mass Spectrometry; IC=Ion Chromatography; CCN=cloud condensation nuclei; OPC=Optical Particle Counter; HR-ToF-AMS=High-Resolution Time-of-Flight Aerosol Mass Spectrometer; PILS=Particle Into Liquid Sampler; SP2=Single Particle Soot Photometer; AFS=Atomic Fluorescence Spectrophotometry.

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Table 3b. P-3 research payload in ARCTAS.

Parameters ¹	Method ²	Investigator	Reference
Aerosol number, size, composition, volatility, optical properties	OPC, PSAP, TDMA, CN counter	A. Clarke, U. Hawaii	Clarke et al. (2004)
Size-resolved aerosol composition and volatility, black carbon aerosol, light scattering	ToF-AMS, SP2, nephelometer	A. Clarke, U. Hawaii	Anderson et al. (1996), Schwarz et al. (2006), DiCarlo et al. (2008)
CCN	CCN counter	A. Nenes, Georgia Tech	Medina et al. (2007)
Aerosol optical properties	Cavity ring-down extinction	A. Strawa, NASA ARC	Strawa et al. (2006)
AOD, H ₂ O column	Tracking Sun photometer	J. Redemann, NASA ARC	Russell et al. (2007)
Solar spectral irradiance	Spectrometer	S. Schmidt, U. Colorado	Kylling et al. (2005)
BRDF	CAR	C. Gatebe, U. Maryland and NASA GSFC	Gatebe et al. (2003)
Radiation fluxes	Broadband radiometers	A. Bucholtz, Naval Res. Lab	Bucholtz et al. (2008)
Carbon monoxide (CO)	TLAS	J. Podolske, NASA ARC	Podolske and Lowenstein (1993)

¹ OD=aerosol optical depth; BRDF=bidirectional reflectance distribution function; CCN=cloud condensation nuclei.

² OPC=Optical Particle Counter; PSAP=Particle Soot Absorption Photometer; CN=condensation nuclei; ToF-AMS=Time of Flight Aerosol Mass Spectrometer; TDMA=Tandem Differential Mobility Analyzer; SP2=Single Particle Soot Photometer; CAR=Cloud Absorption Radiometer; TLAS=Tunable Laser Absorption Spectroscopy.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Table 3c. B-200 research payload in ARCTAS.

Parameters (remote sensing)	Method ¹	Investigator	Reference
Vertical distribution of aerosol properties including extinction at 532 nm, backscatter and depolarization at 532 and 1064 nm	HSRL	R. Ferrare, NASA LaRC	Hair et al. (2008)
Total and linearly polarized reflectance, aerosol and cloud optical properties (optical depth, size parameters, refractive index)	RSP ²	B. Cairns, NASA GISS	Waquet et al. (2009)

¹ HSRL=High Spectral Resolution Lidar; RSP=Research Scanning Polarimeter.

² Summer campaign only.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3d. ARC-IONS¹ ozonesonde sites during ARCTAS.

Site	Spring launches ²	Summer launches ³
Barrow, Alaska (71 N, 157 W)	20	
Boulder, Colorado (40 N, 105 W)		13
Bratt's Lake, Saskatchewan (50 N, 109 W)	15	13
Churchill, Manitoba (59 N, 94 W)	17	9
Egbert, Ontario (44 N, 80 W)		11
Eureka, Nunavut (80 N, 86 W)	19	1
Goose Bay, Newfoundland (53 N, 60 W)		14
Kelowna, British Columbia (49 N, 122 W)	13	13
Narragansett, Rhode Island (41 N, 72 W)	4	3
Resolute, Nunavut (75 N, 95 W)	17	2
Sable Island, Nova Scotia (45 N, 60 W)	12	14
Stony Plain (Edmonton), Alberta (54 N, 114 W)	19	15
Summit, Greenland (73 N, 39 W)	19	18
Trinidad Head, California (40 N, 124 W)	18	16
Whitehorse, Yukon Territory (61 N, 135 W)	12	14
Yarmouth, Nova Scotia (44 N, 66 W)		14
Yellowknife, Northwest Territories (63 N, 115 W)		19

¹ Arctic Intensive Ozonesonde Network Study. Lead investigators were S. Oltmans, NOAA and A. Thompson, Pennsylvania State University.

² Covering the ARCTAS-A period 1–20 April 2008.

³ Covering the ARCTAS-B period 26 June–12 July 2008.

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Table 3e. Ground-based investigations during ARCTAS.

Parameters	Method ¹	Investigator	Reference
Remote and in situ measurements	Instrumented trailer	A. Thompson, Penn State	Thompson et al. (2007)
Aerosol vertical profiles	Lidar	G. Shaw, UA Fairbanks	Wetzel et al. (2003)
Tropospheric BrO	MAX-DOAS	W, Simpson, UA Fairbanks	Simpson et al. (2007b)

¹ MAX-DOAS=Multi-Axis Differential Optical Absorption Spectroscopy.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Table 3f. Supporting investigations during ARCTAS.

Product ¹	Model, data bases	Investigator	Reference
Meteorological forecasts	Various sources, WRF, WRF-Chem	H. Fuelberg, Florida State U.	Fuelberg et al. (2009)
CTM forecasts	FLEXPART	J. Burkhardt, NILU	Hirdman et al. (2009)
	STEM	G. Carmichael, U. Iowa	Adikhary et al. (2008)
	GEOS-5	J. Rodriguez, NASA GSFC; D. Jacob, Harvard	Liang et al. (2009)
	MOZART, CAM-Chem	L. Emmons, NCAR	Arellano et al. (2007); Emmons et al. (2009)
	REAM	Y. Wang, Georgia Tech	Choi et al. (2008)
Near-real-time fire data	Various products	M. Fromm, Naval Res. Lab	Fromm and Servranckx (2003)
Near-real-time satellite data	CALIOP	D. Winker, NASA LaRC	Winker et al. (2007)
	OMI	K. Chance, Harvard-Smithsonian	Levelt et al. (2006)
	TES	J. Worden, NASA/JPL	Worden et al. (2008)
	AIRS	J. Warner, U. Maryland	McMillan et al. (2008)
	MODIS	A. Chu, NASA/GSFC	Chu et al. (2005)
	MISR	R. Kahn, NASA/GFSC	Kahn et al. (2005)
	MOPITT	L. Emmons, NCAR	Deeter et al. (2003)
Emission inventories	Various products	G. Carmichael, U. Iowa	Streets et al. (2006)
Photochemical modeling	LaRC box model	J. Olson, NASA LaRC	Olson et al. (2006)
Near-real-time global CTM analyses	GEOS-Chem	D. Jacob, Harvard	Zhang et al. (2008)

¹CTM = chemical transport model.

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 4a. DC-8 flights during the ARCTAS-A spring deployment¹.

Flight #	Location ²	Date	Special features
3	Palmdale-Fairbanks	1 April	TES and MLS validation, Asian plume
4	Fairbanks-Thule	4 April	OMI BrO validation
5	Thule-Fairbanks	5 April	Summit spiral, OMI BrO validation, North American plume
6	Fairbanks-Iqaluit	8 April	Prudhoe Bay overpass, intercomparison with P-3, spiral over Eureka, OMI BrO validation
7	Iqaluit-Fairbanks	9 April	OMI BrO validation, coordination with B-200, Barrow missed approach ³
8	Fairbanks local 1	12 April	N. Pole flight, Asian and European plumes, Prudhoe Bay missed approach
9	Fairbanks local 2	16 April	Asian plume, coordination with B-200
10	Fairbanks local 3	17 April	N. Pole flight, European plumes, Prudhoe Asian and Bay missed approach
11	Fairbanks-Palmdale	19 April	MLS/CALIPSO underflights, P-3 intercomparison, Asian plume

¹ Flights were typically 8 h in duration.

² Palmdale, California (35 N, 118 W) was the aircraft home base.

³ “Missed approach” refers to an airport runway approach with re-climb just before landing, thus allowing vertical profiling down to the ground.

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 4b. P-3 flights during the ARCTAS-A spring deployment¹.

Flight #	Location ²	Date	Special features
3	Wallops-Yellowknife	31 March	US pollution, Arctic haze including dust
4	Yellowknife-Fairbanks	1 April	Pollution layers, spiral and legs under CALIPSO and OMI.
5	Fairbanks local 1	6 April	Spiral and legs under B-200 and CALIPSO, OMI square spiral and radiation legs, BRDF and albedo over Elson Lagoon ³
6	Fairbanks-Thule	8 April	DC-8 intercomparison, spiral over Eureka during CALIPSO and OMI overpasses, BRDF and albedo
7	Thule-Fairbanks	9 April	Square spiral on CALIPSO and OMI track, BRDF and albedo of ice and cloud, AOD gradient and stack, pollution plume.
8	Fairbanks local 2	13 April	Cloud albedo and BRDF, Asian pollution
9	Fairbanks local 3	15 April	Intercomparison and radiation stack with NOAA P-3 under B-200, BRDF and albedo at Elson Lagoon, radiation leg on CALIPSO and OMI track.
10	Fairbanks-Moffett Field	19 April	DC-8 intercomparison on CALIPSO and OMI track, steps in DC-8 lidar curtain, square spiral during CALIPSO and OMI overpass.

¹ Flights were typically 6 h in duration.

² Wallops, Virginia (38 N, 75 W), and Moffett Field, California (37 N, 122 W) were the aircraft home bases.

³ Sea ice expanse just north of Barrow.

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 4c. B-200 flights during the ARCTAS-A spring deployment¹.

Location	Date	Special features
Barrow local 1	3 April	CALIPSO validation over Arctic Ocean NW of Barrow
Barrow local 2	4 April	Coordinated pattern with DOE Convair-580, multiple passes over Barrow ARM site
Barrow local 3	6 April	CALIPSO validation SW of Barrow, coordination with P-3, dust layers in upper troposphere
Barrow local 4	8 April	Two flights coordinated with DOE Convair-580, extensive low cloud over Barrow, aerosol layers aloft
Barrow local 5	9 April	CALIPSO validation over W Alaska, coordinated flight with DC-8 under CALIPSO track and near Barrow
Barrow local 6	10 April	CALIPSO validation NW of Barrow
Barrow local 7	12 April	Coordinated CALIPSO track with DC-8 and NOAA P-3 SW of Barrow, biomass burning plumes
Barrow-Fairbanks	13 April	Coordinated CALIPSO track with P-3 over W Alaska, coordination with DOE Convair-580 near Barrow
Fairbanks local	15 April	Coordination with P-3 and NOAA P-3 west of Fairbanks and with CALIPSO track over central Alaska, biomass burning plumes
Fairbanks-Nome-Barrow	16 April	Two flights; coordination with DC-8 and CALIPSO track over western Alaska, Asian plumes
Barrow local 8	17 April	CALIPSO validation NW of Barrow, coordination with DC-8 east of Barrow
Barrow local 9	19 April	CALIPSO validation NW of Barrow, coordination with DOE Convair-580 and NOAA P-3 near Barrow, Siberian biomass burning plume sampled with other aircraft over ARM site

¹ Flights were typically 5 h in duration.

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Table 5a. DC-8 flights during the ARCTAS-B summer deployment¹.

Flight #	Location	Date	Special features
12	Palmdale local 1	18 June	Los Angeles Basin and offshore ship emissions
13	Palmdale local 2	20 June	California Central valley, urban and biogenic emissions
14	Palmdale local 3	22 June	California upwind boundary conditions
15	Palmdale local 4	24 June	Los Angeles Basin diurnal evolution and offshore ship emissions
16	Palmdale-Cold Lake	26 June	California fire plumes, OMI validation, HIAPER intercomparison
17	Cold Lake local 1	29 June	Siberian fire plume, pyroconvective outflow, Saskatchewan fires
18	Cold Lake local 2	1 July	Near-field evolution of Saskatchewan fire plumes
19	Cold Lake local 3	4 July	Near-field evolution of Saskatchewan fire plumes, California and
20	Cold Lake local 4	5 July	Saskatchewan fire plumes, biogenic emissions Siberian fire plumes, TES validation
21	Cold Lake-Thule	8 July	Siberian fires and Asian pollution
22	Thule local	9 July	MOPITT validation over Summit, intercomparison with DLR Falcon,
23	Thule-Cold Lake	10 July	P-3 intercomparison, Alberta oil sands emissions Siberian fires and Asian pollution
24	Cold Lake-Los Angeles	14 July	California fire plumes

¹ Flights 12–15 were part of the ARCTAS-CARB deployment. Flight durations were typically 8 h.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 5b. P-3 flights during the ARCTAS-B summer deployment¹.

Flight #	Location	Date	Special features
12	Moffett Field	24 June	19 ship plume crossings, profiles under Terra and Aqua, ocean BRDF (calm & 50-knot winds), fire and pollution plumes.
13/14 ²	Moffett Field-Cold Lake	26 June	Lake Tahoe smoke radiation study in DC-8 lidar curtain and under Terra, lake albedo and BRDF, Central Valley smoke gradient
15	Cold Lake local 1	28 June	Fresh fire plumes (including pyrocumulus), coordination with B-200
16	Cold Lake local 2	29 June	Fresh fire plumes, Fort Mc. Murray plume, BRDF and albedo over lake and land
17	Cold Lake local 3	30 June	CALIPSO and OMI underflight, black and white smoke, in situ and radiation under B-200, BRDF over smoke, Asian pollution with dust
18	Cold Lake local 4	2 July	Radiation work under B-200, aerosol evolution in fire plumes, pyrocumulus
19	Cold Lake local 5	3 July	Ft. McMurray AERONET spiral in regional fire haze, MODIS-MISR-MOPITT spiral, B-200 and CALIPSO-OMI underpass spiral, radiation stack, forest BRDF, fresh fire plumes and downwind evolution, radiation-AOD-in situ ramp.
20	Cold Lake local 6	6 July	Radiation profiles and legs under MISR, BRDF of water surfaces, cloud edges and dense smoke spirals under B-200, pyrocumulus outflow
21	Cold Lake local 7	7 July	Siberian fire plume, aerosol effects on MODIS cloud sensing, radiation stacks
22	Cold Lake local 8	9 July	Fresh fire plumes, spirals and radiation legs under MISR, radiation legs and stacks under B-200 and CALIPSO-OMI
23	Cold Lake local 9	10 July	Fresh fire plumes, Ft. McMurray plume, DC-8 intercomparison under B-200
24	Cold Lake-Wallops	14 July	Transit

¹ Flight 12 was part of the ARCTAS-CARB deployment. Flight durations were typically 6 h.

² Flight 13 was from Moffett Field to Edmonton, Alberta (Canadian customs) and flight 14 was from Edmonton to Cold Lake.

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 5c. B-200 flights during the ARCTAS-B summer deployment¹.

Location	Date	Special features
Yellowknife local 1	28 June	CALIPSO validation track, coordination with P-3 over Saskatchewan fires
Yellowknife local 2	29 June	Coordination with DC-8 over Saskatchewan fires and smoke plumes
Yellowknife local 3	30 June	CALIPSO validation and coordination with P-3, Saskatchewan fire plumes, Asian pollution, dust
Yellowknife local 4	2 July	Two flights; coordination with P-3 over fresh and aged fire plumes; CALIPSO validation on second flight
Yellowknife local 5	3 July	CALIPSO validation and P-3 coordination along CALIPSO track
Yellowknife local 6	6 July	Two flights; second flight was coordination with P-3 for fires and plumes
Yellowknife-Cold Lake-Edmonton	7 July	Two flights: Asian/Siberian plume over Alberta
Edmonton local and Edmonton-Yellowknife	8 July	Two flights: Asian/Siberian plume over western Alberta/eastern British Columbia; CALIPSO validation over northern Alberta
Yellowknife local 7	9 July	CALIPSO validation, coordination with P-3 along CALIPSO track and in fresh fire plumes
Yellowknife local 8	10 July	CALIPSO validation, coordination with P-3 and DC-8 near Yellowknife

¹ Flight durations were typically 5 h.

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.




Fig. 1a. Flight tracks for the ARCTAS-A spring deployment (Table 4a–c). Yellow circles are ozonesonde launch sites (Table 3d).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.




Fig. 1b. Flight tracks for the ARCTAS-B summer deployment (Table 5a–c). Yellow circles are ozonesonde launch sites (Table 3d).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.




Fig. 1c. Flight tracks for the ARCTAS-CARB deployment (Table 5a–b).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The ARCTAS aircraft mission: design and executionD. J. Jacob et al.




Fig. 2. Monthly fire counts for April and July 2008 from the MODIS Aqua satellite instrument (13:30 local overpass time), cloud- and overpass-corrected (V5, available from <http://neespi.gsfc.nasa.gov/myd14cm1.shtml>).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The ARCTAS aircraft mission: design and execution

D. J. Jacob et al.




Fig. 3. Aerosol scattering coefficient ($\lambda=550$ nm) measured at Poker Flat Research Range near Fairbanks, Alaska in March–June 2008. The ARCTAS and ARCPAC intensive flight periods are indicated.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)