

This discussion paper is/has been under review for the journal *Atmospheric Chemistry and Physics (ACP)*. Please refer to the corresponding final paper in *ACP* if available.

**Hydration and
dehydration at the
tropical tropopause**

C. Schiller et al.

Hydration and dehydration at the tropical tropopause

C. Schiller, J.-U. Groß, P. Konopka, F. Plöger, F. H. Silva dos Santos, and N. Spelten

Forschungszentrum Jülich GmbH, ICG-1, 52425 Jülich, Germany

Received: 22 July 2009 – Accepted: 27 July 2009 – Published: 24 August 2009

Correspondence to: C. Schiller (c.schiller@fz-juelich.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

High-resolution water measurements from three tropical airborne missions in Northern Australia, Southern Brazil and West Africa in different seasons are analysed to study the transport and transformation of water in the tropical tropopause layer (TTL) and its impact on the stratosphere. The mean profiles are quite different according to the season and location of the campaigns, with lowest mixing ratios below 2 ppmv at the cold point tropopause during the Australian mission in November/December and high TTL mixing ratios during the African measurements in August. We present backward trajectory calculations considering freeze-drying of the air to the minimum saturation mixing ratio and initialised with climatological satellite data. This trajectory-based reconstruction of water agrees well with the observed H₂O average profiles and therefore demonstrates that the water vapour set point in the TTL is primarily determined by the Lagrangian saturation history. Deep convection was found to moisten the TTL, in several events even above the cold point up to 420 K potential temperatures. However, our study does not provide evidence for a larger impact of these highly-localised events on global scales.

1 Introduction

The entry of water vapour into the stratosphere is controlled by the interplay of transport and freeze-drying primarily in the tropics (e.g. Fueglistaler et al., 2009, and references therein). Several hypotheses have been proposed to explain the dryness of the stratosphere and the variability of the water entry. The average temperature and corresponding saturation ice mixing ratio at the tropical tropopause are higher than the observed average water entry. Therefore, Newell and Gould-Stewart (1981) proposed fountain regions for the entry into the stratosphere, where tropopause temperatures are coldest, as the region of the Western Pacific during the northern hemispheric winter and the Bay of Bengal during northern hemispheric summer. Taking up the idea of dedicated regions of effective freeze-drying, Holton and Gettelman (2001) proposed

ACPD

9, 17495–17529, 2009

Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



that horizontal advection through these cold tropopause regions is causing the dryness of air before entering the stratosphere, rather than enhanced upwelling in these particular regions. Several climatological studies based on a Lagrangian approach (e.g. Jensen and Pfister, 2004; Fueglistaler et al., 2005; Fueglistaler and Haynes, 2005) demonstrated that saturation mixing ratios along trajectories provide a good first-order estimate for stratospheric water vapour mixing ratios and their variability on seasonal and interannual scales. Such studies confirm the important role of the Western Pacific and the Asian monsoon regions to control the moisture flux into the stratosphere.

As an alternative hypothesis to large-scale dehydration, Danielsen (1993) and Sherwood and Dessler (2000) postulated that dehydration occurs primarily in deep overshooting convection. Extremely low temperatures in cumulonimbus turrets may lead to extremely dry air, providing that the air is exposed to lowest temperatures for a sufficiently long time to allow for the ice particles to sediment-out. However, based on part of the data of our study, Corti et al. (2008) presented experimental evidence that overshooting convection has rather a hydrating than dehydrating effect close to the tropical tropopause. Similar conclusions have been drawn from model simulations recently (Chaboureau et al., 2007; Grosvenor et al., 2007; Jensen et al., 2007). Upscaling of overshoot events still reveals a large uncertainty for the estimation of their global impact on the stratospheric water budget.

In this paper, we present airborne high-resolution water measurements from three tropical sites at different seasons, i.e. from Northern Australia and from Southern Brazil during boreal winter and from West Africa during boreal summer. The profiles from the different campaigns are compared in order to determine the characteristic water distribution for each region and season studied. Trajectory studies are used to compare the recent saturation history of the probed air masses with the observed water vapour mixing ratios. We investigate to what extent water vapour can be reconstructed based on trajectories which consider convection only in large scale mean vertical transport rates. The impact of deep convection is studied for individual cases providing independent evidence for persistent moistening of the tropopause region.

2 Experiments

The water measurements presented here were obtained by the Fast In-situ Stratospheric Hygrometer (FISH) which is based on the Lyman- α photofragment fluorescence technique. Instrument and calibration procedure are described in Zöger et al. (1999). The FISH inlet is mounted forward-facing thus measuring total water, i.e. the sum of gas-phase water and ice particles. The procedure to correct the data for oversampling of ice particles is discussed in Schiller et al. (2008). FISH was mounted on the Russian M55 Geophysica high-altitude aircraft with a maximum ceiling of 20 km. Thus measurements in the TTL and lower stratosphere were possible.

We present data from three tropical experiments (Table 1): The Tropical Convection, Cirrus and Nitrogen Oxide experiment (TroCCiNOx) in January and February 2005 from Araçatuba, Brazil, the Stratospheric-Climate Links with Emphasis on the Upper Troposphere and Lower Stratosphere (SCOUT-O3) tropical mission in November and December 2005 from Darwin, Australia, and the African Monsoon Multidisciplinary Analysis (AMMA/SCOUT-O3) experiment in August 2006 from Ouagadougou, Burkina Faso. All three campaigns covered the convective season at the locations in Southern America, West Africa and Northern Australia, with the two southern hemisphere missions taking place during boreal winter and the northern hemisphere mission in Africa during boreal summer.

The SCOUT-O3 flight routes from Darwin are displayed in Fig. 1a. They include local flights in the vicinity of convection as well as long-range flights to probe the inflow and outflow region of convection and the large scale background TTL. Also the measurements during the transfer flights from and to Europe via Cyprus, Arabian Emirates, India, Thailand, and Borneo covered a wide area in the tropics. An overview on the flight patterns, objectives and meteorology of the campaign is given in Vaughan et al. (2008) and Brunner et al. (2009), respectively. Compared to the mesoscale convective system observed over Brazil and West Africa, regularly occurring isolated storms localised over the Tiwi Islands north of Darwin, so-called “Hector” storms (May et al., 2009, and references therein), were investigated during several flights.

Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Similarly, the TroCCiNOx flights include local flights to investigate convective systems (4, 5 and 18 February 2005) and long-range flights to investigate the background TTL and stratosphere (Fig. 1b). Measurements were also made during the transfer route from and to Europe via Spain, Cape Verde Islands, and Recife (Brazil). The operational region in Brazil was located at the edge of the tropics, so also the mixing of tropical with extratropical air was detected during some flights (Konopka et al., 2007). Continental convective systems penetrating the tropopause have been probed by the aircraft on 4 February 2005 (see also Chaboureau et al., 2007) and 5 February 2005.

During the flights over Africa in the frame of AMMA/SCOUT-O3 (overview of flight patterns and meteorology by Cairo et al., 2009), no direct observations of deep convective systems could be made within the range of the aircraft; flight routes are given in Fig. 1c. The multi-cell convective systems of the African monsoon occurred typically upwind, i.e. east of the flight routes as could be identified from satellite IR images. The instrumented transfers from and to Europe covered mainly the transition to extratropical regions.

3 Data/Results

In this paper, the vertical distribution of water vapour in the TTL and lower stratosphere is discussed and compared for the different campaign locations and seasons. The flights listed in Table 1 include measurements of at least two profiles during ascent and descent, most of the local flights include additional profiling information from dives. The vertical information may, according to the flight pattern, be gathered over horizontal distances of several hundred kilometres.

In Fig. 2, the measured water profiles are displayed for the three campaigns, colour-coded with respect to the relative humidity with respect to ice (RH_{ice}). As FISH measures total water, measurements above 100% RH_{ice} in Fig. 2 may include observations in clouds, in particular for RH_{ice} exceeding the threshold of homogeneous ice formation (Koop et al., 2000). The profiles are rather compact around a mean value – except

data obtained in clouds and specific situations to be discussed in Sect. 4.2. They show specific characteristics for each campaign, reflecting the regional and seasonal conditions.

The lowest mixing ratios of less than 2 ppmv are observed during SCOUT-O3 at potential temperatures $\theta=370\text{--}380\text{ K}$, almost coinciding with the local cold point tropopause which was observed around 375 K on average for this period and location (Brunner et al., 2009). The corresponding temperatures were low at these altitudes, i.e. on average -85°C , therefore frequently RH_{ice} close to saturation or clouds were observed in the vicinity of the tropopause. Striking features are layers of high H_2O and cloud occurrence above the tropopause from observation of deep convection above Tiwi Island, the Hector storms (see Sect. 4.2). High RH_{ice} at the cold point tropopause is indicative for effective freeze-drying down to very low mixing ratios in this region and season.

Also during TroCCiNOx, very low mixing ratios were observed, in few cases also close to 2 ppmv. The average minimum is found at slightly higher altitudes, but still close to the average cold point tropopause, which was found between 370 and 380 K, depending on the actual conditions. As the most obvious difference to the SCOUT-O3 observations, mean cold point temperatures were higher, i.e. -79°C ; thus most of the measurements were made at subsaturated conditions without clouds. Only for the flights on 4 and 5 February 2005 when convection reached the tropopause, clouds and high RH_{ice} were observed in that region. Again, enhanced water vapour and ice particles were injected into the stratosphere up to about 415 K during these events.

The AMMA/SCOUT-O3 profiles look very different to those of the two other campaigns, both in absolute values and in the stratospheric gradient. At the cold-point tropopause with an average location at 370 K for this campaign, the water mixing ratio has a mean value of 5 ppmv and shows a pronounced change of the slope, but decreases further in the stratosphere. Balloon measurements in the same region (Khaykin et al., 2009) show the H_2O minimum at approximately 20 km altitude, corresponding to the maximum ceiling of the Geophysica aircraft from which our data

Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



are obtained. Here, the mean mixing ratio obtained from the FLASH-B instrument is 4.4 ppmv and thus consistent with our measurements at these altitudes (4.1 ppmv). Temperatures at the tropopause are significantly higher than during SCOUT-O3 but similar to those from TroCCiNOx. With the higher H₂O values, RH_{ice} frequently is close to 100% in a thin layer around the cold point, sometimes accompanied by cirrus cloud occurrence. Though no Geophysica flights directly above deep convection were made during AMMA/SCOUT-O3, layers of enhanced H₂O above the tropopause are detected in three profiles.

For comparison, our data from February-March 1999 over the Indian Ocean during the APE-THESEO campaign (MacKenzie et al., 2006) show similar to the SCOUT-O3 data extremely low H₂O mixing ratios with saturation at the cold point tropopause. However, during that campaign no convection reaching the tropopause level was probed, but ultrathin cirrus at the tropopause occurred quite frequently (Peter et al., 2003). The water minimum was at slightly higher altitudes than during SCOUT-O3.

Figure 3 summarises the average water profiles obtained during the four tropical Geophysica missions. Here, data in convection-induced ice clouds have not been included as these phenomena lead to a local moistening which might not be representative for the global H₂O distribution (see Sect. 4). Figure 3 also shows frostpoint hygrometer measurements over Indonesia during the SOWER campaign in early January 2006 (Vömel et al., 2007), which are obtained close to our SCOUT-O3 data in a nearby region. The averaged SCOUT-O3 and SOWER profiles agree well above 380 K in terms of absolute values and slope, at lower altitudes the local conditions are likely the reason for slightly higher mixing ratios over Darwin with a strong convective impact. Also the profiles of Kelly et al. (1993) obtained during the STEP campaign January 1987 from Darwin are very close to the SCOUT-O3 profiles in quantity and slope. The TroCCiNOx and APE-THESEO data show slightly lower values above 410 K compared to SCOUT-O3. As these campaigns were carried out three months later in season, the difference would be consistent with an upward propagation of the dehydration signal, the so-called tape recorder (Mote et al., 1996) in that time: using a clear-sky cooling

Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



rate of 0.6 K/d for this region (e.g. Corti et al., 2005), air parcels are lifted from the cold point tropopause to the 410–430 K region in this time. We cannot exclude, however, that a year-to-year variability is overlaid to this phenomenon (Sect. 4.1.3), and the regional influence at these altitudes leads to zonal asymmetries, but to first order the data reflect the upward propagation of the tape recorder signal with season. Close to the tropopause, the H₂O mixing ratios during TroCCiNO_x are higher than the very low values during SOWER, SCOUT-O3 and APE-THESEO and therefore account for moister and warmer conditions at the edge of the tropics over South America. The AMMA/SCOUT-O3 obtained in NH summer are higher by about 2–3 ppmv at 380 K compared to the other campaigns, consistent with the higher water vapour entry in that season and similar to e.g. the historical Panama data in September 1980 by Kley et al. (1982) and sonde data from Costa Rica (Vömel et al., 2007). Above 410 K, the absolute mixing ratios are very similar for all campaigns (including the SOWER, Panama and Cost Rica data), thus the tape recorder amplitude is already smeared-out or masked by regional or year-to year variability. However, the different slopes between the NH summer and winter data are still indicative for the upwelling of the freeze-drying signal from the cold point region with season.

4 Discussion

The water vapour profiles obtained during the three missions TroCCiNO_x, SCOUT-O3 and AMMA/SCOUT-O3 provide a sample of data representing different tropical sites and seasons, each with its specific characteristics. In this Section, we will examine the hydrating and dehydrating processes causing the specific H₂O distributions in particular in the vicinity of the cold point tropopause of the individual missions. Though the coverage of these selected missions covers neither the complete tropical belt nor a full seasonal cycle, the analysis will provide a thorough insight in tropical hydration and dehydration mechanisms and the role of specific tropical regions. Emphasis will be laid on the role of deep convection and the large-scale advection during which the water vapour content of air parcels are changed.

4.1 Dehydration at the cold point tropopause

For the Darwin flights during SCOUT-O3, saturation conditions were observed frequently in a broad altitude band around the cold point, indicative of local dehydration down to the very low mixing ratios of 2 ppmv or less. For AMMA/SCOUT-O3, saturation is reached up to the cold point in part of the profiles, and for TroCCiNO_x, H₂O mixing ratios at the tropopause were usually lower than the local saturation. Therefore, in many cases the corresponding freeze drying events must have occurred earlier and at different locations.

We use backward trajectories ending at the flight paths of the three missions to determine the temperature history of the observed air parcels and to reconstruct its water content. The trajectories are calculated using the Chemical Lagrangian Model of the Stratosphere (CLaMS), which is driven by ECMWF meteorological fields. For the vertical parameterisation of the tropopause region, a hybrid coordinate is chosen as described by Konopka (2007). Above a pressure of 100 hPa, approximately the tropical tropopause pressure, the hybrid vertical coordinate is equal to the potential temperature, with the cross-isentropic vertical velocity deduced from clear-sky radiation (Morcrette scheme). Below, coordinate surfaces smoothly transform into being parallel to pressure surfaces, with the vertical velocity smoothly transforming into the ECMWF vertical wind. Hence, our trajectory calculation is purely diabatic above the tropopause, gradually transforming into kinematic below.

Sensitivity studies with the cross-isentropic velocity deduced from the ERA-Interim reanalysis total diabatic heating rate, including clear-sky and cloud radiative effects and latent heat release (Plöger et al., 2009), show that the change in the resulting water profiles is less than 0.1 ppmv. Therefore, restricting to clear-sky radiative heating rates above 100 hPa is not a shortcoming of our study.

$RH_{ice}=100\%$ is found to be the average and most frequent value in cirrus clouds down to the low temperature conditions in the tropics (Krämer et al., 2009). The nucleation of such clouds occurs however at higher RH_{ice} (Koop, 2000), i.e. at higher

Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



initial water concentrations which are then freeze-dried to a lower value. Jensen et al. (2001) pointed out, that in the tropics low ice crystal numbers and rapid sedimentation might not allow dehydration of air from the nucleation threshold down to saturation ($RH_{ice}=100\%$). Therefore, we tested the sensitivity of saturation mixing ratio along the trajectories considering a relative humidity over ice $RH_{ice}=100\%$ and 130% , respectively. Hereby, the mixing ratio at the end of the trajectory can be changed by up to 0.3 ppmv , providing that a major dehydration had occurred (see Sect. 4.1.3).

4.1.1 Saturation history

Figure 4a–c shows averaged total water data profiles measured by FISH (black dots, including their standard deviation grey-shaded) together with the cold point history for the tropical flights of the three campaigns SCOUT-O3, TroCCiNOx and AMMA/SCOUT-O3, respectively; FISH data include the local flights plus transfers made between 0° and 30° S , between 30° S and 30° N , and between 0° and 30° N , respectively. Figure 4a–c shows profiles of the minimum saturation water vapour mixing ratio calculated along backward trajectories, for 10-day backward trajectories (green crosses) and for 30-day backward trajectories (red crosses). A longer trajectory history results in lower values of the minimum saturation mixing ratios primarily for two reasons: first, the likelihood to encounter very low temperature regions increases with trajectory length. Second, due to the diabatic ascent of air, a larger part of the trajectories ending in the low stratosphere has passed the cold point region when a longer trajectory history is considered. The horizontal dashed lines denote the mean altitude of the cold point along the backward trajectories (green/red for 10/30-day backward trajectories). In the right side of each panel, the fraction of trajectories which originate from the 380 K isentrope or below (black line) is displayed. Further, we show the fraction of trajectories whose minimum saturation mixing ratio is higher by 0.5 ppmv than the corresponding FISH measurement (magenta solid line) or lower by 0.5 ppmv (dashed-dotted magenta line), respectively.

Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hydration and dehydration at the tropical tropopause

C. Schiller et al.

For SCOUT-O3 (Fig. 4a), the average of the FISH data is consistent with the trajectory history from approximately 375 to 405 K. The trajectory length has only a small influence. This implies, that air masses are dehydrated locally or near-by in the same region over this rather large altitude range. A more detailed analysis of the freeze-drying evolution over the SCOUT-O3 period around the cold point and its modulation by Kelvin waves is given by Brunner and Schiller (2009). Further, additional injections of water to these altitudes (see Sect. 4.2) can be freeze-dried back to these low values. Higher up in the stratosphere, the measured H₂O data are much lower than the recent saturation along the trajectories. Thus, their water vapour set point goes back to earlier seasons, and the fraction of trajectories that have passed the cold point tropopause within the last month is close to zero for $\theta > 410$ K.

Below 380 K, measured FISH data are however higher than the minimum saturation mixing ratio along the back trajectories. This implies that the potentially dry air should have been moistened recently – most likely by deep convection which occurred quite frequently in the operational area (Brunner et al., 2009). But also higher up, i.e. up to $\theta = 420$ K, a fraction of 10–50% of the trajectories infers a water mixing ratio lower than the observation by at least 0.5 ppmv (magenta dashed-dotted line), indicative for moistening of the lowermost stratosphere by overshooting convection (see Sect. 4.2), or ECMWF temperatures being too low.

For TroCCiNOx (Fig. 4b), a large change of the saturation mixing ratio along the backward trajectories is evident when increasing their length from 10 to 30 days. The 30-day trajectories are by far closer to the low observed values at 375–400 K than the shorter ones. This implies, that the set point of the low water vapour at these altitudes dates back up to a month and likely from a remote region (see Sect. 4.1.3). The reported frequent subsaturation close to the cold point and above (Fig. 2b) already showed that the operational area at the edge to the subtropics is not an area of active dehydration – at least when air with low mixing ratios is advected. Below 380 K, again a large fraction of trajectories (30 day only) implies a lower H₂O mixing ratio than the observations show. Again, deep convection is one source of recent injections up to

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the cold point, in few cases even above (e.g. Chaboureau et al., 2007). In the vicinity of the subtropical jet, however also mixing with extratropical and tropospheric air is an efficient source of additional water, as demonstrated in case studies for the TroCCiNOx data (Konopka et al., 2007).

For the AMMA/SCOUT-O3 observations (Fig. 4c), saturation along the backward trajectories can explain the H₂O measurements only up to the cold point around 370 K. Even the minimum saturation over 30 days does not approach the measurements much closer in the stratosphere anymore. In the TTL, the much higher temperatures result in much higher saturation mixing ratios than for the other two campaigns. The observations are closer to this value; the slight enhancement is again indicative for moistening by convection. There is however also a small fraction of trajectories, which end at higher observed H₂O mixing ratios than expected from the temperature history (magenta dashed-dotted line). We discuss in Sect. 4.2 that also over Africa convection penetrating the tropopause is moistening the lowest layers of the stratosphere. It is worth to note that the range of trajectories which have passed the cold point (black line at the right) ends below 400 K, consistent with the slower upwelling in this season (e.g. Randel et al., 2007).

4.1.2 Tropical regions of dehydration

The backward trajectory calculations are suited to identify the tropical region where saturation at the lowest mixing ratio preferably occurs. Figure 5 shows the cold points along the 30-day backward trajectories for the three campaigns in the potential temperature range 375–410 K, colour-coded for the corresponding minimum saturation mixing ratio if it agrees with the observation within an interval of ± 0.5 ppmv.

The lowest temperatures for the Darwin measurement (Fig. 5 upper panel) occur close to the aircraft tracks in the equatorial Western Pacific. This region has been identified also in previous studies as the tropical region with the highest dehydration potential in this season (e.g. Fueglistaler et al., 2005). The observed low temperatures around Darwin and the frequent observation of high RH_{ice} at low H₂O mixing ratios

Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



are indicative for active dehydration also in the operational area. Thus, the outflow and overshoots from deep convection with local injection of water from the troposphere (Sect. 4.2) are likely instantaneously freeze-dried back to extremely low mixing ratios.

As discussed above, for TroCCiNO_x only saturation along longer trajectories can explain the observed low mixing ratios over Southern Brazil. The air masses probed by the aircraft have been dried in regions more equatorwards (Fig. 5 central panel), in particular over South America and over the Western and Central Pacific during the preceding 30 days. Also the ECMWF-based climatology compiled by Fueglistaler et al. (2005) identified these two regions to be coldest. Our measurements at the edge to the subtropics show, that the dehydration to lowest mixing ratios at these cold spots is advected also to remote regions with higher temperatures.

For AMMA/SCOUT-O₃, the air masses probed over Africa generally origin from the East passing also the Asian monsoon region (Fig. 5, lower panel). Lowest temperatures corresponding to a saturation mixing ratio of 4–6 ppmv occur over India and the Gulf of Bengal. Thus, the H₂O mixing ratios close to the cold point observed over Africa are broadly consistent with the colder set point in the Southern flank of the Asian Monsoon. So the Asian monsoon is not only a major source of water vapour as concluded from previous studies (e.g. Randel and Park, 2006; James et al., 2008), its cold point characteristics at the same location also regulate the entry of water vapour into the stratosphere and determine the water vapour concentrations also in the downwind regions over Africa.

4.1.3 Trajectory-based reconstruction of H₂O with HALOE climatological data

In contrast to the trajectory calculations in Sect. 4.1.1, we now extend the discussion of the saturation history by initialising the trajectory end points with data from the Halogen Occultation Experiment (HALOE) onboard the Upper Atmosphere Research Satellite (UARS). A climatological data set of H₂O of this experiment is compiled for the years 1991 to 2002 in Grob and Russell (2005). For the initialisation, the zonal average for the respective latitude bin at the backward trajectory end point is chosen. Figure 6a–c

Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



shows in analogy to Fig. 4 the HALOE-initialised water vapour profiles after modifications by 10-day and 30-day backward trajectory calculations (green and red dots, respectively), compared to the actually measured FISH data for the three tropical campaigns. The simulations are done for RH_{ice} of 100% (crosses) and 130% (circles). The black line in the main panels denotes the mean HALOE climatological H_2O profile used for initialisation.

For SCOUT-O3 (Fig. 6a), the change between the HALOE initialisation profile (black line) and the reconstructed mixing ratios at the observation point (green and red dots, for trajectory lengths of 10-day and 30-day, respectively) is largest. The large modifications below 410 K are already expected from the discussion in Sect. 4.1.1, as the cold point is frequently reached along the trajectories, also in the lower stratosphere: The fraction of trajectories along which a cold point below the HALOE initial value is encountered, is 100–10% in the θ -range 380–410 K (right subpanel). Above the cold point, the calculated H_2O profiles (30-day trajectories) match extremely well the mean profiles measured by FISH, and those for the shorter 10-day history are only slightly higher. Below the cold point, recent convection has enhanced the actual H_2O mixing ratio compared to the simulated values, as already discussed above.

The TroCCiNOx comparison provides also a good agreement between the reconstructed and measured H_2O mean profiles above $\theta=380\text{ K}$ (Fig. 6b). Here, the modifications between the climatological HALOE profiles along the trajectories are by far weaker than for the SCOUT-O3 period, i.e. only 50–10% of the trajectories at 380–400 K meet a cold point below the HALOE initial value (right subpanel). Below 380 K, the measured mean H_2O mixing ratio is higher than the reconstructed value, indicative for additional recent moistening (see Sect. 4.1.1) due to vertical mixing along the subtropical jet (Konopka et al., 2007).

For AMMA/SCOUT-O3 (Fig. 6c), virtually no change of the initial HALOE profile occurs along the back trajectories above 375 K. Here, the slope of the profiles is very similar, though the HALOE climatological data are lower by approximately 0.4 ppmv (or 10%) above 390 K. Good agreement is found at 370–390 K, while below the

Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



measurements are again higher than the simulations, as for the other campaigns.

The systematic difference between the reconstructions, based on HALOE data, and the FISH profiles in the stratosphere during AMMA/SCOUT-O3 is likely caused by instrument uncertainties. In a comprehensive comparison of HALOE data with in-situ instruments (Kley et al., 2000), HALOE measurements of H₂O were found at the low end compared to other hygrometers. In particular, FISH and HALOE data differed by about 8%, which is almost the same discrepancy we find in the AMMA/SCOUT-O3 data again. We note that this discrepancy is within the uncertainty of both instruments, and Kley et al. (2000) did not assign one single instrument as the absolute reference for atmospheric water measurements. However, also in other studies HALOE data appeared to be lower than measurements from other reference instruments. If we assume a low bias of the HALOE measurements on the order of 8%, the agreement between the simulations and FISH measurements would improve for the AMMA/SCOUT-O3 case. Initialisation with higher HALOE data would however not change the H₂O reconstruction for TroCCiNOX and SCOUT-O3 to a larger extent, as most of these data are lowered to the saturation mixing ratio at the minimum cold point along the back trajectories anyway. Therefore, our tropical measurements may be a confirmation for the previously found moderate low bias of HALOE data compared to FISH measurements. As noted already in Sect. 3, mean FISH profiles agree also well with those of the AMMA balloon data using the FLASH-B hygrometer (Khaykin et al., 2009), the slight discrepancy at 19–20 km of 0.2 ppmv is even lower than the total accuracy of the instruments implies (approx. 8% for FISH and 10% for FLASH-B).

Beside such potential systematic instrument discrepancies, a deviation of our FISH measurements from simulations using a climatological initialisation may also be caused by atmospheric variability – in particular for the AMMA/SCOUT-O3 case when the H₂O concentration is not altered along the back trajectories for a large altitude range. Randel et al. (2004) discussed, inter alia based on HALOE data, interannual changes of stratospheric water vapour which could be attributed to changes of temperatures at the tropical tropopause associated with the quasi-biennial oscillation (QBO) and El

Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Niño-Southern oscillation. Hereby, H₂O anomalies on the order of a few tenths ppmv are observed, which are of the same order of magnitude as the observed discrepancies during AMMA/SCOUT-O3. After the unexpected drop of water vapour entry around 2001 (e.g. Randel et al., 2006), recent satellite observations (MIPAS/ENVISAT; ODIN/SMR; MLS/AURA; unpublished results) indicate that water vapour in the tropical lower stratosphere has approached values for our observation period in 2005/2006 which are comparable to the mean values of the period considered in the HALOE climatology of Grooß and Russel (2005). In summary, the observed discrepancies between HALOE and FISH data during AMMA/SCOUT-O3 in the stratosphere may be caused by the interannual variability of water vapour entry, though there is no quantitative assessment of such an effect possible for the moment. Support for the attribution of the low SCOUT-O3 measurements in Darwin to low variability is given by the balloon soundings in Indonesia in January 2006 which showed indeed lower water vapour profiles than in the following years (H. Vömel, personal communication).

Zonal asymmetry may also have caused the observed discrepancies during AMMA/SCOUT-O3. In extension of the HALOE climatology by Grooß and Russel (2005), Fig. 7 shows the zonal distribution of H₂O averaged for 0–15° in August derived from HALOE measurements for the years 1992–2005 as well as the deviation from the zonal mean. We note that with the resolution of 24°-latitude bins (i.e. according to the HALOE orbits) only 5–30 data points contribute per bin, despite the large HALOE data set. Therefore, the significance in particular for the lower isentropes is limited. The increased mixing ratios above Africa (peak at 0° longitude) and Asia (peak at 100° E longitude) imply an influence of the monsoon regions on the TTL. They are strongest at 380 K close to the cold point region with up to 0.5 ppmv for the Asian monsoon region. The zonal modulation is weakened with increasing altitude to an amplitude of 0.1 ppmv at 425 K, close to the altitude where our data set still see sporadically enhanced H₂O layers (Sect. 4.2), and almost vanishes above. From the climatological HALOE perspective, the influence of the monsoon regions is apparent but limited to a few tenths of ppmv in the TTL above the cold point, i.e. 3–9% additional water compared to the

Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



zonal mean. Hereby, the direct contribution of the African monsoon might still be over-estimated as this region is located downwind of the Asian monsoon input region.

For comparison, the specific humidity from ECMWF SHECMWF interpolated to the flight route is plotted in Fig. 6a–c (dashed-dotted black lines). For the SCOUT-O3 averages (panel a), SHECMWF and FISH profiles agree well up to 380 K potential temperature. In the stratosphere SHECMWF has a low bias by 1–2 ppmv compared to both the FISH data and the reconstructed H_2O , and the slope is less steep than that of the observations. For the TroCCiNOx campaign (Fig. 6b), the discrepancies between FISH measurements and SHECMWF are qualitatively very similar as during SCOUT-O3, but somewhat lower in magnitude. During AMMA/SCOUT-O3, SHECMWF is again lower than the FISH profile in stratosphere and thus matches the reconstructed water; around the cold point, SHECMWF exceeds the observations. As a general tendency, SHECMWF underestimates the measured H_2O mixing ratios in the stratosphere by 0.5–2 ppmv, with larger discrepancies at higher altitudes. Similar differences between matched HALOE/MLS data and SHECMWF (ERA-40 analysis) are also apparent in a zonal mean climatology (Dethof, 2003; Fig. 3 therein), i.e. a larger bias and less steep gradients of SHECMWF compared to the satellite zonal mean data for the tropical SH in January and a somewhat better agreement for the AMMA/SCOUT-O3 latitudes and season at higher mixing ratios.

4.2 Moistening by deep convection

In Sect. 4.1, we concluded that the major outflow of convection below the cold point results in a moistening of the lower TTL. However, this additional water vapour does not result in an increased transport of H_2O into the stratosphere, as it is trapped at the cold point during the subsequent ascent. As an example, observations in the aged outflow of a strong Hector event on 30 November 2005 showed saturated air masses over a large area at and below the cold point at mixing ratios of 2–4 ppmv with only a few remnants of particle plumes. Therefore, already a few hours after a convective event, the air masses from the outflow might be freeze-dried back to the saturation conditions.

Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Figure 2 however shows layers of significantly enhanced water up to 420 K in several profiles of all three campaigns, which are indicative of convective overshoots. In Schiller et al. (2008), we determined ice water content (IWC) of these ice clouds from the Darwin and Brazilian experiments. IWC in these clouds at the top of deep convection can be up to tens of ppmv and is thus significantly higher than the climatological range of IWC from other observations at these low temperatures (Plate 8 in Schiller et al., 2003) and the ambient mixing ratio. Such high IWC are therefore indicative of rapid transport from the troposphere as in convection. Previously, Kelly et al. (1993) reported observations of particles above 400 K above Hector storms, however with IWC lower than 2 ppmv. Corti et al. (2008) and de Reus et al. (2008) showed that our measurements of enhanced water are correlated with particle measurements, in several cases with diameters of 40 μm and above. In the Corti et al. (2008) study, hypotheses other than convective origin of these particles could be excluded. The observations provide evidence, that these particles injected above the cold point have a moistening effect and that vice versa, there is no indication for a “convective dehydration” as proposed in previous studies (Danielsen, 1993; Sherwood and Dessler, 2000). They confirm model calculations by Jensen et al. (2007) postulating such a moistening effect if particles are injected in subsaturated air above the tropopause.

The measurements over Africa also show layers of enhanced water up to 420 K in single profiles (Fig. 2c). In contrast to the observations during SCOUT-O3 and TroC-CiNOx, these measurements were not obtained in the vicinity of deep convection and no particles are detected in these layers (except a very moderate enhancement of backscatter ratio in the multi-angle spectrometer on 8 August 2006; F. Cairo personal communication). On the other hand, back trajectory calculation show that these observations were made downwind of large mesoscale convective systems detected with good coincidence in time and location by the Meteosat-08 observations (CLaMS trajectories, not shown; for a more detailed study see Fierli et al., 2009). Even if overshoots into the stratosphere may not be fully resolved by cloud top heights derived using brightness temperature differences, they have likely occurred above the identified

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



deep convective systems underlying. These observations, which are similar to those reported from AMMA balloon-borne measurements in the same region (Khaykin et al., 2009), provide evidence, that deep convection injects water irreversibly above the cold point, as these layers with H₂O enhanced by 2–3 ppmv can still be detected about 10–20 h after the convective event in subsaturated air without remnants of particles. Therefore, we can exclude that such injections occur only as a temporary lifting above the tropopause without mixing with the environment as well as a complete sedimentation of particles back to tropospheric altitudes.

The observed layers with enhanced water vapour above the cold point are a highly localised phenomenon on horizontal scales of several 10 km. Probing these events above Tiwi Island during SCOUT-O3, identified an area on the order of 2000 km² (50 km diameter) with particles above the tropopause. Similarly, during AMMA the layers occurred only in single profiles while the next profile a few 100 km apart showed again the undisturbed background profile. The observations during the three campaigns, also taking into account the balloon observations by Khaykin et al. (2009), are quite frequent for such a small-scale phenomenon. On the other hand, its fraction of the total number of measured profile has to be regarded as a biased upper limit, since the flight programme in all missions was focused on the observation of convective systems and their outflow regions.

5 Conclusions

The three tropical airborne campaigns discussed in this study yield different but complementary information on the water vapour distribution and related processes in the TTL. The measurements during SCOUT-O3 from Darwin, Australia in November and December 2005 were characterised by lowest temperatures and mixing ratios close to the cold point. Frequent cloud occurrence, high relative humidity and backward trajectory calculations demonstrated that in the operational area and nearby regions efficient dehydration down to lowest mixing ratios of less than 2 ppmv occurs. During TroCCiNOx over Southern Brazil in February 2005, almost the same low mixing ratios

were observed but at lower RH_{ice} . The freeze-drying events had occurred several weeks before in cold regions equatorwards. The AMMA/SCOUT-O3 mission from West Africa in August 2006 showed a complete different mean water profile with higher mixing ratios in the TTL according to the season. The region is strongly influenced by air masses from the Asian monsoon region, which also appeared to be the region determining the mean water vapour abundance observed. Though the observations of water vapour are highly variable between the three campaigns and even between single flights of one mission, they allow for a consistent interpretation towards a conclusive understanding of water vapour transport and transformation through the TTL into the stratosphere.

From our data, the tropical water vapour entry into the stratosphere is found to be a highly complex interplay between freeze-drying in the coldest regions of the TTL and fresh supply of water by convection or by mixing with tropospheric air. Based on our studies, the 420 K isentrope appears to be the transition between a regionally controlled regime and a zonally well mixed compartment at higher altitudes. This is a quantitative confirmation of the TTL concept discussed in Fueglistaler et al. (2009). It marks the highest altitude of direct impact of overshooting convection. Further, recent condensation and sedimentation of particles may have altered the composition below 420 K leading to regional deviations from the zonal mean. Though an interannual variability of the TTL parameters has been observed to propagate further up and is visible in the different tape recorder signals, the local and regional variability in the TTL becomes smeared out above 420 K.

The main outflow of convection below the cold point as well as frequently observed local overshoots into the stratosphere have been unequivocally identified to moisten their environment. Below the tropopause, our measurements show significantly higher H_2O mixing ratios than inferred from the backtrajectory history which does not consider recent individual convective systems. However, no significant remnants of this high water released in the lower TTL can be found above the cold point anymore as this additional water is removed on its way through the cold point.

Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Hydration and
dehydration at the
tropical tropopause**C. Schiller et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Further, we found layers of enhanced water vapour above the cold point which could be traced to direct injection by overshooting turrets. These irreversible injections of H₂O up to 420 K were observed to be isolated and thus highly localised, i.e. with horizontal extensions on the order of 50 km. Even if an estimate of the amount of water injected in such a single event is possible (Chemel et al., 2009), an upscaling of its global impact on the stratospheric water vapour budget might be possible only with large uncertainties (Corti et al., 2009; Fierli et al., 2009). A global distribution of convection penetrating the tropopause has been derived from satellite data (Gettelman et al., 2002; Liu and Zipser, 2005). Our high resolution measurements confirm the maximum depth of 1.5 km above the cold point as found in these studies. Based on a brightness temperature analysis of satellite cloud data, Gettelman et al. (2002) estimated a zonal mean fraction area of 0.1% of convection penetrating the tropopause, while the TRMM data analysis of Liu and Zipser results in a still lower frequency. The difference, also apparent in the geographical distribution, is likely caused by biases of the different data and their retrieval. However, the regions and seasons we investigated during SCOUT-O3, TroCCiNOx and AMMA/SCOUT-O3 are identified in both studies to be representative for intensive deep convective activity; other key regions are the Western Pacific, the Congo basin, the Indian monsoon region and Central America (Panama region).

The comparison of our data with the HALOE-initialised reconstruction of H₂O could however give a hint on the maximum possible impact of overshoots on the stratospheric water vapour budget. For SCOUT-O3 and to a lesser extent also for TroCCiNOx conditions, additional water injected above the cold point can still be freeze-dried to the low mean values there. The comparison of the AMMA/SCOUT-O3 data with the climatological HALOE profile might suggest enhanced water vapour abundance over Africa (Fig. 6 lower panel). Most likely, the observed discrepancy however reflects the formerly identified low bias of HALOE compared to FISH data, in particular as the offset ranges to higher altitudes than the highest observed overshoots. An effect by inter-annual variability may be on the order of 10% difference (Sect. 4.1.3) and could – depending on its sign – mask a potentially larger discrepancy between our data and

the climatology for this particular year. From the comparison between the HALOE-based water reconstruction and the AMMA/SCOUT-O3 data set we might not be able to quantify precisely the convective impact on the lower stratospheric water content, as there are several remaining uncertainties. On the other hand, based on the magnitude of these uncertainties we do not find evidence for an enhancement of H₂O over Africa of more than 10%. Zonal asymmetries in the HALOE climatological data (Fig. 7) imply such an increase over the African monsoon region and therefore corroborate the finding of a potential moistening of the TTL up to 420 K – but only to a moderate degree.

Trajectory-based reconstruction of water vapour, initialised with climatological HALOE data, yield a remarkable agreement with the in-situ observations presented in this study above the tropopause for the different regions and seasons studied. Therefore the Lagrangian saturation history on time scales of hours to 1 month appears to be appropriate and sufficient to describe the average H₂O distribution at the cold point tropopause and in the first kilometres above. So, our results confirm previous findings from similar Lagrangian studies (e.g. Fueglistaler et al., 2005; Fueglistaler and Haynes, 2005), however now demonstrated with a data set of much finer scales in time and space and thereby covering different tropical regions and seasons. Jensen and Pfister (2004) tested with a more sophisticated Lagrangian ice cloud model the sensitivity of the stratospheric vapour entry on cloud properties and their dependence on wave perturbations and other parameters. While cloud frequency and distribution depended sensitively on these parameters, they also concluded that final water vapour mixing ratios are primarily controlled by the minimum temperatures encountered. Our study confirms these studies and thereby the preceding ideas of Holton and Gettelman (2001) with new experimental evidence, that despite the complex interplay of various processes leading to a highly variable water distribution in the tropics, stratospheric water vapour entry is primarily controlled by the coldest regions in the TTL, in which the air is freeze-dried to its final set point during its way into the stratosphere. The successful trajectory-based reconstruction of H₂O above the tropopause is first of all determined by the reliability of the horizontal wind fields and the temperature distribution, though

Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



convection is only insufficiently parameterized in the underlying ECMWF meteorological data. It is therefore unlikely, that the individually observed deep convective events have a major impact on the global water budget above the tropopause.

Acknowledgements. Discussion of the results and developing ideas of their meaning with many members of the project teams of SCOUT-O3, TroCCiNOx and AMMA/SCOUT-O3 is gratefully acknowledged. We thank MDB, DLR and ERS-Ltd operating and managing the aircraft under the challenging tropical conditions. The study was primarily conducted under the SCOUT-O3 Integrated Project (EC Contract GOCE-CT-2004-505390), with additional campaign funding by the EC projects TroCCiNOx and AMMA as well as by INSU and the Geophysica EEIG.

References

Brunner, D., Siegmund, P., May, P. T., Chappel, L., Schiller, C., Müller, R., Peter, T., Fueglistaler, S., MacKenzie, A. R., Fix, A., Schlager, H., Allen, G., Fjaeraa, A. M., Streibel, M., and Harris, N. R. P.: The SCOUT-O3 Darwin Aircraft Campaign: rationale and meteorology, *Atmos. Chem. Phys.*, 9, 93–117, 2009,

<http://www.atmos-chem-phys.net/9/93/2009/>.

Brunner, D. and Schiller, C.: Large fluctuations of tropopause moisture over the Maritime Continent induced by a Kelvin wave during the SCOUT-O3 campaign in Darwin, Australia, *Atmos. Chem. Phys.*, in preparation, 2009.

Cairo, F., Pommereau, J. P., Law, K. S., et al.: An overview of the SCOUT-O3-AMMA stratospheric aircraft, balloons and sondes campaign in West Africa, August 2006: rationale, roadmap and highlights, *Atmos. Chem. Phys.*, submitted, 2009.

Chaboureaud, J. P., Cammas, J.-P., Duronet, J., et al.: A numerical study of tropical cross-tropopause transport by convective overshoots, *Atmos. Chem. Phys.*, 7, 1731–1740, 2007, <http://www.atmos-chem-phys.net/7/1731/2007/>.

Chemel, C., Russo, M. R., Pyle, J. A., Sokhi, R. J., and Schiller, C.: Quantifying the imprint of a severe Hector thunderstorm during ACTIVE/SCOUT-O3 onto the water content in the upper troposphere/lower stratosphere, *Mon. Weather Rev.*, in press, 2009.

Corti, T., Luo, B. P., Peter, T., Vömel, H., and Fu, Q.: Mean radiative energy balance and vertical mass fluxes in the equatorial upper troposphere and lower stratosphere, *Geophys. Res. Lett.*, 32, L06802, doi:10.1029/2004GL021889, 2005.

Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Hydration and
dehydration at the
tropical tropopause**C. Schiller et al.

Corti, T., Luo, B. P., deReus, M., Brunner, D., Cairo, F., Mahoney, M. J., Matucci, G., Matthey, R., Mitev, V., dos Santos, F. H., Schiller, C., Shur, G., Sitnikov, N. M., Spelten, N., Vössing, H. J., Borrmann, S., and Peter, T.: Unprecedented evidence for overshooting convection hydrating the tropical stratosphere, *Geophys. Res. Lett.*, 35, L10810, doi:10.1029/2008GL033641, 2008.

Danielsen, E. F.: In situ evidence of rapid, vertical, irreversible transport of lower tropospheric air into the lower tropical stratosphere by convective cloud turrets and by larger-scale upwelling in tropical cyclones, *J. Geophys. Res.*, 98, 8665–8681, 1993.

de Reus, M., Borrmann, S., Heymsfield, A. J., Weigel, R., Schiller, C., Mitev, V., Frey, W., Kunkel, D., Kürten, A., Curtius, J., Sitnikov, N. M., Ulanovsky, A., and Ravegnani, F.: Evidence for ice particles in the tropical stratosphere from in-situ measurements, *Atmos. Chem. Phys. Discuss.*, 8, 19313–19355, 2008, <http://www.atmos-chem-phys-discuss.net/8/19313/2008/>.

Dethof, A.: Aspects of modelling and assimilation for the stratosphere at ECMWF, available via: www.atmosp.physics.utoronto.ca/SPARC/News21/21_Dethof.html, SPARC newsletter 21, 2003.

Fierli, F., Orlandi, E., Cairo, F., Law, K. S., Schiller, C., Cagnazzo, C., Viciani, S., Volk, M., Schlager, H., and Ravegnani, F.: Characterization of the convective outflow in the UTLS region during the AMMA campaign Part 1: impact in the upper troposphere and mesoscale simulations, *Atmos. Chem. Phys.*, in preparation, 2009.

Fueglistaler, S., Bonazzola, M., Haynes, P. H., and Peter, T.: Stratospheric water vapour predicted from the Lagrangian temperature history of air entering the stratosphere in the tropics, *J. Geophys. Res.*, 110, D08017, doi:10.1029/2004JD005516, 2005.

Fueglistaler, S. and Haynes, P. H.: Control of interannual and longer-term variability of stratospheric water vapor, *J. Geophys. Res.*, 110, D24108, doi:10.1029/2006JD007273, 2005.

Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkins, I., Fu, Q., and Mote, P. W.: Tropical tropopause layer, *Rev. Geophys.*, 47, RG1004, doi:10.1029/2008RG000267, 2009.

Gettelman, A., Salby, M. L., and Sassi, F.: Distribution and influence of convection in the tropical tropopause region, *J. Geophys. Res.*, 107(D10), 4080, doi:10.1029/2001JD001048, 2002.

Groß, J.-U. and Russell III, J. M.: Technical note: A stratospheric climatology for O₃, H₂O, CH₄, NO_x, HCL and HF derived from HALOE measurements, *Atmos. Chem. Phys.*, 5, 2797–2807, 2005, <http://www.atmos-chem-phys.net/5/2797/2005/>.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Grosvenor, D. P., Choullarton, T. W., Coe, H., and Held, G.: A study of the effect of overshooting deep convection on the water content in the TTL and lower stratosphere from cloud resolving model simulations, *Atmos. Chem. Phys.*, 7, 4977–5002, 2007, <http://www.atmos-chem-phys.net/7/4977/2007/>.

5 Holton, J. R. and Gettelman, A.: Horizontal transport and the dehydration of the stratosphere, *Geophys. Res. Lett.*, 28, 2799–2802, 2001.

James R., Bonazzola, M., Legras, B., Surbled, K., and Fueglistaler, S.: Water vapor transport and dehydration above convective outflow during Asian monsoon, *Geophys. Res. Lett.*, 35, L20810, doi:10.1029/2008GL035441, 2008.

10 Jensen, E. J., Pfister, L., Ackerman, A. S., Tabazadeh, A., and Toon, O. B.: A conceptual model of the dehydration of air due to freeze-drying by optically thin, laminar cirrus rising slowly across the tropical tropopause, *J. Geophys. Res.*, 106, 17237–17252, 2001.

Jensen, E., and Pfister, L.: Transport and freeze-drying in the tropical tropopause layer, *J. Geophys. Res.*, 109, D02207, doi:10.1029/2003JD004022, 2004.

15 Jensen, E. J., Ackerman, A. S., and Smith, J. A.: Can overshooting convection dehydrate the tropical tropopause layer?, *J. Geophys. Res.*, 112, D11209, doi:10.1029/2006JD007943, 2007.

Kelly, K. K., Proffitt, M. H., Chan, K. R., Loewenstein, M., Podolske, J. R., Strahan, S. E., Wilson, J. C., and Kley, D.: Water vapour and cloud water measurements over Darwin during the STEP 1987 tropical mission, *J. Geophys. Res.*, 98, 8713–8723, 1993.

20 Khaykin, S., Pommereau, J.-P., Korshunov, L., Yushkov, V., Nielsen, J., Larsen, N., Christensen, T., Garnier, A., Lukyanov, A., and Williams, E.: Hydration of the lower stratosphere by ice crystal geysers over land convective systems, *Atmos. Chem. Phys.*, 9, 2275–2287, 2009, <http://www.atmos-chem-phys.net/9/2275/2009/>.

25 Kley, D., Schmeltekopf, A. L., Kelly, K., Winkler, R. H., Thomson, T. L., and MacFarland, M.: Transport of water through the tropical tropopause, *Geophys. Res. Lett.*, 9, 617–620, 1982.

Kley, D., Russell III, J., Phillips, C. (Eds.), A. Gettelman, J. Harries, P. Mote, S. Oltmans, E. Remsberg, K. Rosenlof, and C. Schiller: SPARC Assessment of Water Vapour in the Stratosphere and Upper Troposphere, WCRP-113, WMO/TD No. 1043, SPARC Report No. 2, 2000.

30 Konopka, P., Günther, G., Müller, R., Santos, F. H., Schiller, C., Ulanovsky, A., Schlager, H., Volk, C. M., Viciani, S., Pan, L., McKenna, D. S., and Riese, M.: Mixing-driven troposphere to stratosphere transport (TST) across the TTL, *Atmos. Chem. Phys.*, 7, 3285–3308, 2007,

Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Hydration and
dehydration at the
tropical tropopause**

C. Schiller et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

<http://www.atmos-chem-phys.net/7/3285/2007/>.

Koop, T., Luo, B. P., Tsias, A., and Peter, T.: Water activity as the determinant for homogeneous ice nucleation in aqueous solutions, *Nature*, 406, 611–614, 2000.

Krämer, M., Schiller, C., Afchine, A., Bauer, R., Gensch, I., Mangold, A., Schlicht, S., Spelten, N., Sitnikov, N., Borrmann, S., de Reus, M., and Spichtinger, P.: Ice supersaturations and cirrus cloud crystal numbers, *Atmos. Chem. Phys.*, 9, 3505–3522, 2009,
<http://www.atmos-chem-phys.net/9/3505/2009/>.

Liu, C. L. and Zipser, E. J.: Global distribution of convection penetrating the tropical tropopause, *J. Geophys. Res.*, 110, D23104, doi:10.1029/2005JD006063, 2005.

MacKenzie, A. R., Schiller, C., Adriani, A., Beuermann, J., Cairo, F., Corti, T., Gensch, I., Krämer, M., Kröger, C., Lepouchov, B., Lowe, D., Merkulov, S., Mitev, V., Oulanovsky, A., Peter, Th., Ravegnani, F., Rohs, S., Rudakov, V., Salter, P., Stefanutti, L., and Yushkov, V.: The tropopause and hygropause in the equatorial Indian Ocean during February and March 1999, *J. Geophys. Res.*, 111(D18), D18112, doi:10.1029/2005JD006639, 2006.

May, P. T., Allen, G., Vaughan, G., and Connolly, P.: Aerosol and thermodynamic effects on tropical cloud systems during TWICE and ACTIVE, *Atmos. Chem. Phys.*, 9, 15–24, 2009,
<http://www.atmos-chem-phys.net/9/15/2009/>.

Mote, P. W., Rosenlof, K. H., McIntyre, M. E., Carr, E. S., Gille, J. C., Holton, J. R., Kinnersley, J. S., Pumphrey, H. G., Russell III, J. M., and Watery, J. W.: An atmospheric tape recorder: The imprint of tropical tropopause temperatures on stratospheric water vapour, *J. Geophys. Res.*, 101, 3989–4006, 1996.

Newell, R. E. and Gould-Stewart, S.: A stratospheric fountain, *J. Atmos. Sci.*, 38, 2789–2796, 1981.

Peter, Th., Luo, B. P., Wernli, H., Wirth, M., Kiemle, C., Flentje, H., Yushkov, V. A., Khattatov, V., Rudakov, V., Thomas, A., Borrmann, S., Toci, G., Mazzinghi, P., Beuermann, J., Schiller, C., Cairo, F., Di Donfrancesco, G., Adriani, A., Volk, C. M., Strom, J., Noone, K., Mitev, V., MacKenzie, R. A., Carslaw, K. S., Trautmann, T., Santacesaria, V., and Stefanutti, L.: Ultrathin tropical tropopause clouds: I. Cloud morphology and occurrence, *Atmos. Chem. Phys.*, 3, 1083–1091, 2003,
<http://www.atmos-chem-phys.net/3/1083/2003/>.

Plöger, F., Konopka, P., Günther, G., Groöß, J.-U., and Müller, R.: Impact of the vertical velocity scheme on modeling transport in the tropical tropopause layer (TTL), *J. Geophys. Res.*, submitted, 2009.

**Hydration and
dehydration at the
tropical tropopause**C. Schiller et al.

Randel, W. J., Wu, F., Oltmans, S., Rosenlof, K., and Nedoluha, G. E.: Interannual changes of stratospheric water vapor and correlations with tropical tropopause temperatures, *J. Atmos. Sci.*, 61, 2133–2148, 2004.

Randel, W. J., Wu, F., Vömel, H., Nedoluha, G. E., and Forster, P.: Decreases in stratospheric water vapor since 2001: links to changes in the tropical tropopause and the Brewer-Dobson circulation, *J. Geophys. Res.*, 111, D12312, doi:10.1029/2005JD006744, 2006.

Randel, W. J., and M. Park, Deep convective influence on the Asian monsoon anticyclone and associated tracer variability observed with Atmospheric Infrared Sounder (AIRS), *J. Geophys. Res.*, 111, D12314, doi:10.1029/2005JD006490, 2006.

Randel, W. J., Park, M., Wu, F., and Livesey, N.: A large annual cycle in ozone above the tropical tropopause linked to the Brewer-Dobson circulation, *J. Atmos. Sci.*, 64 (12), 4479–4488, 2007.

Schiller, C., M. Krämer, A. Afchine, N. Spelten, N. Sitnikov, The ice water content of Arctic, mid latitude and tropical cirrus, *J. Geophys. Res.*, 113, D24208, doi:10.1029/2008JD010342, 2008.

Sherwood, S. C. and Dessler, A. E.: On the control of stratospheric humidity, *Geophys. Res. Lett.*, 27, 2513–2516, 2000.

Vaughan, G., Schiller, C., MacKenzie, A. R., Bower, K., Peter, T., Schlager, H., Harris, N. R. P., and May, P. T.: SCOUT-O3 /ACTIVE: High-altitude aircraft measurements around deep tropical convection, *B. Am. Meteorol. Soc.*, 89, 647–662, doi:10.1175/BAMS-89-5-647, 2008.

Vömel, H., Barnes, J. E., Forno, R. N., Fujiwara, M., Hasebe, F., Iwasaki, S., Kivi, R., Komala, N., Kyrö, E., Leblanc, T., Morel, B., Ogino, S.-Y., Read, W. G., Ryan, S. C., Saraspriya, S., Selkirk, H., Shiotani, M., Valverde Canossa, J., and Whiteman, D. N.: Validation of Aura Microwave Limb Sounder water vapor by balloonborne Cryogenic Frost point Hygrometer measurements, *J. Geophys. Res.*, 112, D24S37, doi:10.1029/2007JD008698, 2007.

Zöger, M., Afchine, A., Eicke, N., Gerhards, M.-T., Klein, E., McKenna, D. S., Mörschel, U., Schmidt, U., Tan, V., Tuitjer, F., Woyke, T., and Schiller, C.: Fast in situ stratospheric hygrometers: A new family of balloonborne and airborne Lyman α photofragment fluorescence hygrometers, *J. Geophys. Res.*, 104, 1807–1816, 1999.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Hydration and dehydration at the tropical tropopause

C. Schiller et al.

Table 1. Overview of tropical aircraft experiments for this study.

Experiment	Base	Time	No. of local + transfer flights	reference
SCOUT-O3	Darwin, Australia (12° S, 139° W)	31 October– 17 December 2005	8+12	Vaughan et al. (2008) Brunner et al. (2009) http://www.ozone-sec.ch.cam.ac.uk/scout_o3/
TroCCiNOx	Aracatuba, Brazil (21° S, 58° E)	18 January– 18 February 2005	8+6	http://www.pa.op.dlr.de/troccinox
AMMA-SCOUT-O3	Ouagadougou, Burkina Faso (12° N, 1° W)	29 July– 17 August 2006	5+4	Cairo et al. (2009) http://www.isac.cnr.it/~utls/m55amma/

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hydration and dehydration at the tropical tropopause

C. Schiller et al.




Fig. 1. Routes of local flights during **(a)** SCOUT-O3 with operation base in Darwin, Australia **(b)** TroCCiNOx with operation base in Araçatuba, Brazil and **(c)** AMMA/SCOUT-O3 with operation base in Ouagadougou, Burkina Faso. Parts of the transfer flights to and from Europe are included.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hydration and
dehydration at the
tropical tropopause

C. Schiller et al.




Fig. 2. Vertical profiles of total water during the tropical aircraft campaigns **(a)** SCOUT-O3 in Northern Australia (November 2005), **(b)** TroCCiNOx in Brazil (February 2005), and **(c)** AMMA/SCOUT-O3 in West Africa (August 2008). Colour code denotes the relative humidity with respect to ice.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Hydration and dehydration at the tropical tropopause

C. Schiller et al.




Fig. 3. Comparison of mean water vapour profiles of the TroCCiNOx, SCOUT-O3, AMMA/SCOUT-O3 campaigns with those obtained during the APE-THESEO, SOWER, Panama and Costa Rica missions (references see text). FISH data measured in clouds are not considered.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hydration and dehydration at the tropical tropopause

C. Schiller et al.




Fig. 4. Mean vertical profiles of H_2O measured during **(a)** SCOUT-O3 **(b)** TroCCiNOx and **(c)** AMMA/SCOUT-O3, including tropical parts of transfer routes to and from Europe (black dots, binned for θ intervals of 2 K; grey envelope denotes standard deviation). Minimum saturation mixing ratios along back trajectories are given by green (10-day trajectories) and red (30-day trajectories) crosses (freeze-drying along the trajectories to $\text{RH}_{\text{ice}}=100\%$). The green and red dashed lines indicate the average cold point altitude along the back trajectories. The right side of each panel displays the fraction of trajectories, along which the minimum saturation is lower (solid magenta line) or higher (dashed-dotted magenta line) by 0.5 ppmv than the measured mixing ratio. The black solid line represents the fraction of trajectories which have passed the cold point during the last 30 days.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hydration and dehydration at the tropical tropopause

C. Schiller et al.




Fig. 5. Location of coldest temperatures along backward trajectories ($\theta=375\text{--}410\text{ K}$) with end points along the flight tracks during (a) SCOUT-O3, (b) TroCCiNOx and (c) AMMA/SCOUT-O3 (considered flight routes including part of the transfers displayed by grey triangles). The minimum saturation mixing ratio along the backtrajectories if corresponding to the aircraft measurement is marked in colour.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hydration and dehydration at the tropical tropopause

C. Schiller et al.




Fig. 6. Comparison between mean measured H₂O profiles (as in Fig. 4, panels (a)–(c) again display the three campaigns) and simulated ones; freeze-drying along the trajectories to RH_{ice}=100% (crosses) and 130% (circles). Here, backtrajectory calculations of the water vapour mixing ratio at the flight route are initialised with H₂O climatological data from the HALOE satellite experiment (green dots based on 10-day trajectories, red dots on 30-day trajectories). The black profiles show the mean HALOE profiles and their standard deviation used for initialisation of the trajectory calculations. The dashed-dotted profile is the averaged ECMWF specific humidity value interpolated to the flight tracks. The right subpanels show the fraction of trajectories along which a cold point below the HALOE initial value is encountered.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hydration and dehydration at the tropical tropopause

C. Schiller et al.




Fig. 7. HALOE climatology of H₂O at 0–15° N for August of years 1992–2005 as function of longitude constructed similar as in Grooß and Russel (2005) (upper panel) and deviations from the mean at four different isentropes (lower panel).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

